

# SIMPLICIAL VOLUME OF MANIFOLDS FIBERING WITH CONNECTED STRUCTURE GROUP

THORBEN KASTENHOLZ

**ABSTRACT.** In this note we investigate the simplicial volume of fiber bundles with connected structure group. We are able to show that if the structure group is either compact or a Lie group, or if the fiber is aspherical that the simplicial volume of the total space agrees with the simplicial volume of the trivial bundle.

## 1. INTRODUCTION

For a homology class  $\alpha$  of a topological space, one can define its  $\ell_1$ -norm via

$$\|\alpha\| = \inf \left\{ \sum_i |\lambda_i| \mid \alpha \text{ is represented by } \sum_i \lambda_i \sigma_i \right\}$$

For a closed connected and oriented manifold  $M$  the norm of the fundamental class  $\| [M] \|$  is called the simplicial volume of  $M$  and denoted by  $\|M\|$ . This was first investigated by Gromov in [2] and has various relations to topology and geometry. It is a consequence of Gromov's Mapping Theorem [2, The Mapping Theorem, Section 3.1] that the simplicial volume of a manifold only depends on the image of the fundamental class in the group homology of its fundamental group with respect to the map that classifies the universal covering.

Another consequence of Gromov's Mapping Theorem is that the simplicial volume of every manifold  $E$ , which fibers over a manifold  $B$  with a fiber that has amenable fundamental group and has positive dimension, vanishes. Interestingly, if one instead requires that the base manifold has amenable fundamental group, while having no requirements on the fiber, then an analogue of this result can not hold. In fact every mapping torus of a pseudo-Anosov surface automorphism is a hyperbolic 3-manifold and hence has non-zero simplicial volume. Strengthening the restrictions on the base manifold leads to the following question:

**Question 1.1.** *Suppose a closed, connected and oriented manifold  $E$  fibers over a simply-connected manifold  $B$  with positive dimension, does the simplicial volume of  $E$  vanish?*

As a partial answer to [Question 1.1](#) we are able to show the following:

**Theorem A.** *Let  $F \rightarrow E \rightarrow B$  denote a bundle of closed, connected and oriented manifolds with either*

- (a) *compact structure group, or*
- (b) *a connected Lie group as structure group, or*
- (c) *aspherical fiber and connected structure group*

*then the simplicial volume of  $E$  agrees with the simplicial volume of the trivial bundle  $F \times B$ .*

---

Date: April 24, 2024.

Note that if the base manifold is simply-connected, then the structure group of the bundle can always be reduced to a connected group (See [Section 2](#) for an introduction to structure groups). Hence in the context of [Question 1.1](#) it is quite natural to consider bundles with connected structure group and no restriction on the fundamental group of the base manifold as in [Theorem A](#).

As a previous result, it was shown in [\[4\]](#) that if  $B$  is either a simply-connected sphere, or a flexible and 2-connected manifold and the fundamental group of the fiber satisfies the Novikov conjecture, then the simplicial volume of the total space vanishes.

The structure group being compact is equivalent to the existence of a metric on the fiber which is invariant under the structure group. It is further equivalent to the existence of a metric on the total space such that the fibers are totally geodesic submanifolds (See for example [Theorem 1.11](#) and [Theorem 1.12](#) in [\[1\]](#)). Furthermore since isometry groups are Lie groups, this shows that the first case is actually contained in the second case (if one passes to an effective action of the structure group). Additionally every Lie group is homotopy equivalent to its maximally compact structure group, hence the first and second point are actually equivalent.

[Theorem A](#) will be a consequence of the following more general result:

**Theorem B.** *Let  $F \rightarrow E \rightarrow B$  denote a fiber bundle of closed oriented connected manifolds with structure group  $G$ . If  $\pi_0(G, e)$  is finite and a multiple of every rational homology class in the universal covering  $\widetilde{BG}$  can be represented by a disjoint union of spaces that fiber over simply-connected spheres, then the simplicial volume of  $E$  agrees with the simplicial volume of the trivial bundle  $B \times F$ .*

*In particular this holds if  $\widetilde{BG}$  has the rational homotopy type of a product of simply-connected Eilenberg-MacLane spaces,*

The proof of [Theorem B](#) can roughly be divided into two parts. Let  $\pi$  denote the fundamental group of the base manifold. In the first part we show that the simplicial volume of the total space only depends on the image of the fundamental class of the base manifold under the canonical map to  $BG \times B\pi$  (See [Proposition 3.1](#) for the precise statement). This step only requires that  $G$  is connected.

For the second step, we will establish in [Section 4](#) that homology classes in  $BG$  can be represented by a product of simply-connected spheres, provided that  $G$  is connected and  $BG$  has the rational homotopy type of a product of Eilenberg-MacLane spaces and then combine these representatives with [Theorem C](#) in [\[4\]](#) to finish the proof of [Theorem B](#). Finally we will also show in [Section 4](#) that the classifying spaces of the structure groups in [Theorem A](#) have the rational homotopy type of a product of Eilenberg-MacLane spaces.

**Remark 1.2.** The proof presented in [Section 3](#) actually shows that if the answer to [Question 1.1](#) is yes, then the simplicial volume of the total space of any bundle with connected structure group agrees with the simplicial volume of the trivial bundle.

**Acknowledgements.** The author would like to thank Mark Pedron and Manuel Krannich for helpful discussions about rational homotopy theory.

## 2. FIBER BUNDLES AND STRUCTURE GROUPS

This section contains a short introduction to fiber bundles and their structure groups, for a more thorough introduction see [\[3\]](#).

Let  $\pi_E: E \rightarrow B$  denote a smooth fiber bundle with fiber  $F$ , i.e. there exists an open covering  $U_i \subset B$  and fiber preserving diffeomorphisms  $\Phi_i: \pi_E^{-1}(U_i) \rightarrow U_i \times F$  such that

$\Phi_i \circ \Phi_j^{-1}: U_i \cap U_j \times F \rightarrow U_i \cap U_j \times F$  is a fiber preserving diffeomorphism. In particular this yields maps  $\overline{\Phi}_{i,j}: U_i \cap U_j \rightarrow \text{Diff}(F)$  such that

$$(1) \quad \overline{\Phi}_{k,i} \circ \overline{\Phi}_{i,j} = \overline{\Phi}_{k,j}$$

These maps are usually called **cocycles of E** and since these cocycles carry the gluing information of the bundle, one can check that there is a one-to-one correspondence between maps satisfying the cocycle condition (1) and fiber bundles. Notice that the same cocycles can be used to define a  $\text{Diff}(F)$ -principal bundle  $E_{\text{Diff}(F)}$  which comes equipped with a map  $E_{\text{Diff}(F)} \times F \rightarrow E$ . Conversely, using the left action of  $\text{Diff}(F)$  on  $F$ , one can associate to every  $\text{Diff}(F)$ -principal bundle  $P$  a  $F$ -bundle by taking the fiber product  $P \times F/\text{Diff}(F)$ . This yields a one-to-one correspondence between  $\text{Diff}(F)$ -bundles and  $F$ -bundles.

There is the universal  $\text{Diff}(F)$ -principal bundle  $\text{Diff}(F) \rightarrow E\text{Diff}(F) \rightarrow B\text{Diff}(F)$  and for every  $\text{Diff}(F)$ -principal bundle  $P$  over  $B$ , there exists a up to homotopy unique map  $\zeta_P: B \rightarrow B\text{Diff}(F)$  such that the pullback  $\zeta_P^* E\text{Diff}(F)$  is isomorphic to  $P$ . Analogously, for every  $F$ -bundle  $E$  over  $B$ , there exists a up to homotopy unique map  $\zeta_E: B \rightarrow B\text{Diff}(F)$  such that  $E$  is isomorphic to  $\zeta_E^* \gamma_F$ , with  $\gamma_F = E\text{Diff}(F) \times_{\text{Diff}(F)} F$ . For that reason  $B\text{Diff}(F)$  is called the classifying space for  $\text{Diff}(F)$  and we call  $\gamma_F$ , the universal  $F$  bundle.

Let now  $G$  denote a topological group admitting a continuous group action on  $F$ . Such a group action is equivalent to a continuous group homomorphism

$$\phi: G \rightarrow \text{Diff}(F)$$

We say that **the structure group of a bundle  $E \rightarrow B$  can be reduced to  $G$** , if there exists an open covering of  $B$  and trivializations over these open maps, such that the cocycles  $\overline{\Phi}_{i,j}: U_i \cap U_j \rightarrow \text{Diff}(F)$  can be lifted to maps  $\widetilde{\Phi}_{i,j}: U_i \cap U_j \rightarrow G$  such that the cocycle condition  $\overline{\Phi}_{k,i} \circ \overline{\Phi}_{i,j} = \overline{\Phi}_{k,j}$  still holds. An  $F$ -bundle together with a choice of these lifts is called a  **$F$ -bundle with structure group  $G$** .

The group homomorphism  $\phi$  also yields a map on classifying spaces  $B\phi: BG \rightarrow B\text{Diff}(F)$  and the structure group of a bundle can be reduced to  $G$  if and only if the classifying map lifts to  $BG$  and a reduction of the structure group corresponds to such a lift. Similar to the previous considerations, for every  $F$ -bundle  $E \rightarrow B$  with structure group  $G$ , there exists a up to homotopy unique map  $\zeta_E: B \rightarrow BG$  such that  $E$  is isomorphic as a bundle with structure group  $G$  (meaning that the fiberwise isomorphisms are also elements in  $G$ ) to  $\zeta_E^* (B\phi^* \gamma_F)$ . We denote  $B\phi^* \gamma_F$  by  $\gamma_{F,G}$  and call it the universal  $F$ -bundle with structure group  $G$ .

**Example 2.1.** Let  $E \rightarrow B$  denote a smooth  $\mathbf{R}^n$ , bundle, then a reduction of the structure group of  $E$  to  $GL_n(\mathbf{R}) \subset \text{Diff}(\mathbf{R}^n)$  equips  $E$  with a vector bundle structure. Furthermore the structure group of such a bundle can be reduced to  $GL_n^+(\mathbf{R})$  if and only if the bundle is orientable and a choice of orientation corresponds to a reduction of the structure group to  $GL_n^+(\mathbf{R})$ .

One can proceed similarly with spin-structures, or metrics, or complex structures.

Another important example for us is the following:

**Example 2.2.** Let  $G$  denote a topological group and  $G_0$  its identity component, then one easily sees that  $BG_0 \rightarrow BG$  is a universal covering map. As a consequence, one can reduce the structure group of every bundle  $E \rightarrow B$  with structure group  $G$  and simply-connected base  $B$  to  $G_0$ .

Note that the very same considerations will also work for topological bundles i.e. bundles with structure group  $\text{Homeo}(F)$  and even fibrations, where one takes the associative  $H$ -space of self-homotopy equivalences  $\text{Aut}(F)$  as  $G$  (and the cocycle condition only has to hold up to homotopy).

### 3. PROOF OF THEOREM B

The following section contains the proof of Theorem B. The proof consists of two parts, first we show a connection between the simplicial volume of the total space of a bundle and the homology classes in  $BG \times B\pi$ . This step works for every connected structure group. Finally we use the existence of simple representatives of all homology class in  $BG$  to finish the proof.

Let  $F \rightarrow E \rightarrow B$  denote a fiber bundle of compact, connected and orientable manifolds with connected structure group  $G$  and  $n$ -dimensional base manifold.

Let  $\zeta_E: B \rightarrow BG$  denote the classifying map of the bundle, i.e. there exists an isomorphism  $\zeta_E^* \gamma_{F,G} \cong E$ , where  $\gamma_{F,G}$  denotes the tautological  $F$  bundle with structure group  $G$  over  $BG$  as introduced in Section 2.

Let us denote the fundamental group of  $B$  by  $\pi$ , and let  $\zeta_B: B \rightarrow B\pi$  denote a classifying map of the universal covering. Consider the map

$$f = \zeta_E \times \zeta_B: B \rightarrow BG \times B\pi$$

which is covered by a map  $\bar{f}: E \rightarrow \gamma_{F,G} \times B\pi$ . We denote the projection  $\gamma_{F,G} \times B\pi \rightarrow BG \times B\pi$  by  $\pi_{\gamma_{F,G}}$ .

The corresponding long exact sequences of the homotopy groups of a fibration yield the following diagram:

$$\begin{array}{ccccccccccc} \dots & \longrightarrow & \pi_2(B, *) & \longrightarrow & \zeta_B^* \pi_1(F, *) & \longrightarrow & \pi_1(E, *) & \longrightarrow & \pi_1(B, *) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow = & & \downarrow \bar{f}_* & & \downarrow \cong & & \\ \dots & \longrightarrow & \pi_2(BG \times B\pi, *) & \longrightarrow & \pi_1(F, *) & \longrightarrow & \pi_1(\gamma_{F,G} \times B\pi, *) & \longrightarrow & \pi_1(BG \times B\pi, *) & \longrightarrow & 0 \end{array}$$

A simple diagram chase shows that  $\bar{f}_*$  is a surjection with amenable kernel, in particular by Gromov's mapping theorem,  $\bar{f}$  induces an isometry on homology with respect to the  $\ell_1$ -norm. Therefore the simplicial volume of  $E$  agrees with the  $\ell_1$ -norm of  $\bar{f}([E])$ , which equals  $\pi_{\gamma_{F,G}}^! (f_*([B]))$ , where  $\pi_{\gamma_{F,G}}^!$  denotes the fiber transfer. In particular  $\pi_{\gamma_{F,G}}^! (f_*([B]))$  only depends on  $f([B])$ . All in all this yields

**Proposition 3.1.** *Let  $F \rightarrow E \rightarrow B$  and  $F \rightarrow E' \rightarrow B'$  denote two bundles with connected structure group  $G$  and suppose further that the fundamental groups of  $B$  and  $B'$  are isomorphic. Let us denote their fundamental group by  $\pi$ . Assume further that there exists a  $\lambda \in \mathbf{R}$  such that*

$$(\zeta_E \times \zeta_B)_*([B]) = \lambda (\zeta_{E'} \times \zeta_{B'})_*([B']) \in H_*(BG \times B\pi; \mathbf{R})$$

*then  $\|E\| = \lambda \|E'\|$ .*

We will now focus on classifying space admitting nice representatives of their homology classes. This is encapsulated in the following definition:

**Definition 3.2.** Let  $X$  denote a simply-connected topological space, we call its rational homology **generated by fiber bundles over spheres** if there exists maps

$$f_i: E_i \rightarrow X$$

where  $i$  lies in some index set and  $E_i$  is a closed, oriented manifold that fibers over a simply-connected sphere, such that the set  $\{f_{i*}([E_i])\}$  generates the rational homology groups of  $X$ .

Let us from here on forth assume that the rational homology of  $BG$  is generated by fiber bundles over spheres. We will show in [Section 4](#) that this includes the three cases:

- $G$  is a compact group
- $G$  is a Lie group
- $G$  is the identity component of the  $H$ -space of self-homotopy equivalences of an aspherical manifold.

By Künneths theorem, the  $n$ -th rational homology group of  $BG \times B\pi$  is isomorphic to

$$\bigoplus_i H_i(BG; \mathbb{Q}) \otimes H_{n-i}(B\pi; \mathbb{Q})$$

Hence  $f_*([B])$  is the sum of classes  $\alpha_i$  with

$$\alpha_i \in H_i(BG; \mathbb{Q}) \otimes H_{n-i}(B\pi; \mathbb{Q})$$

By assumption any  $\alpha_i$  with  $i \geq 1$  can be represented by a disjoint union of manifolds fibering over simply-connected spheres times some representative of a class in  $H_{n-i}(B\pi; \mathbb{Q})$ . Therefore the norm of  $\pi_{\gamma_{F,G}}^!(\alpha_i)$  which equals the simplicial volume of the pullback of  $\gamma_{F,G}$  to the aforementioned representative vanishes by Theorem C in [\[4\]](#) as the simplicial volume of any space fibering over a simply-connected sphere vanishes. Therefore we conclude that the simplicial volume of  $E$  agrees with the  $\ell_1$ -norm of  $\pi_{\gamma_{F,G}}^!(\alpha_0)$ , but  $\alpha_0$  is an element of  $H_0(BG; \mathbb{Q}) \otimes H_n(B\pi; \mathbb{Q})$  to be more precise it equals  $\zeta_{B*}([B])$ , in particular it is represented by  $* \times \zeta_B = \zeta_{F \times B} \times \zeta_B: B \rightarrow BG \times B\pi$  and therefore  $\pi_{\gamma_{F,G}}^!(\alpha_0)$  is represented by  $F \times B$  which concludes the proof of [Theorem B](#).

#### 4. FINDING SMALL REPRESENTATIVES

The goal of this section is to show that the rational homology of the classifying space  $BG$  is generated by fiber bundles over spheres if:

- $G$  is a compact group
- $G$  is a Lie group
- $G$  is the identity component of the  $H$ -space of self-homotopy equivalences of an aspherical manifold.

In order to accomplish that, we will show that the rational homology of spaces that are rationally homotopy equivalent to a product of Eilenberg-MacLane spaces is generated by fiber bundles over spheres. In fact, we will show something slightly stronger:

**Definition 4.1.** Let  $X$  denote a simply-connected topological space, we call its rational homology **generated by products of spheres** if there exists maps

$$f_i: \prod_{l=1}^{k_i} S^{n_{i,l}} \rightarrow X$$

where  $i$  lies in some index set, and  $n_{i,l} \geq 2$ , such that  $\left\{f_{i*}\left(\left[\prod_{l=1}^{k_i} S^{n_{i,l}}\right]\right)\right\}$  generates the rational homology groups of  $X$ .

The following lemma shows that this property is preserved under rational homotopy equivalences. This follows immediately from the fact that products of spheres are universal (See [\[6\]](#)). Nevertheless we include a proof here.

**Lemma 4.2.** *Suppose  $f: X \rightarrow X'$  is a rational homotopy equivalence and suppose further that the rational homology of  $X'$  is generated by products of spheres, then the same holds for  $X$ .*

*Proof.* Let  $S = \prod_i S^{n_i}$  denote a product of spheres and  $g: S \rightarrow X'$  a map. We will show that there exists a map  $h: S \rightarrow S$  of non-zero degree such that  $g \circ h$  lifts along  $f$ .

Note that  $f$  being a rational homotopy equivalence implies that all homotopy groups of the homotopy fiber  $\text{HoFib}(f)$  are torsion. We will construct a lift of  $g$  via induction over its skeleta  $S^{(d)}$ , precomposing with a map of non-zero degree map where necessary.

Let  $h_k: S \rightarrow S$  denote the map that is the product of degree  $k$  maps on the factors of  $S$ . By the Künneth theorem, this map sends every integral homology class to a  $k$ -fold multiple of another integral homology class, additionally we have  $h_k \circ h_{k'} = h_{kk'}$ .

Since  $f$  induces an isomorphism on rational homotopy groups, there exists a  $k_0$  such that, we can lift  $g \circ h_{k_0}|_{\bigvee_i S^{n_i}}$  to  $X$ . In particular we can lift  $g \circ h_{k_0}$  to the first  $d$ -skeleton which is not a point.

Let us now proceed inductively and assume that a lift on the  $d$  skeleton of  $S$  of  $g \circ h_{k_d}$  is already constructed. The obstruction class  $o_{d+1}$  to extending this lift to the  $d+1$  skeleton is an element of

$$H^{d+1}(S^{(d+1)}, S^{(d)}; \pi_d(\text{HoFib}(f), *))$$

Let us consider the long exact sequence of integral cohomology for this pair:

$$0 \rightarrow H^d(S^{(d+1)}; \mathbb{Z}) \rightarrow H^d(S^{(d)}; \mathbb{Z}) \rightarrow H^{d+1}(S^{(d+1)}, S^{(d)}; \mathbb{Z}) \rightarrow H^{d+1}(S^{(d+1)}; \mathbb{Z}) \rightarrow 0$$

Since the cellular chain complex of a sphere is concentrated in a single degree, one concludes that the map  $H^d(S^{(d+1)}; \mathbb{Z}) \rightarrow H^d(S^{(d)}; \mathbb{Z})$  is an isomorphism, hence  $H^{d+1}(S^{(d+1)}, S^{(d)}; \mathbb{Z})$  is isomorphic to  $H^{d+1}(S^{(d+1)}; \mathbb{Z})$ . Therefore  $h_k$  also sends every class in  $H^{d+1}(S^{(d+1)}; \mathbb{Z})$  to a  $k$ -fold multiple of another class. Hence there exists a  $k_{d+1}$  such that  $h_{k_{d+1}}^*(o_{d+1})$  vanishes. By naturality of the obstruction class we conclude that the lift of  $g \circ h_{k_d} \circ h_{k_{d+1}} = g \circ h_{k_d k_{d+1}}$  can be extended to the  $d+1$ -skeleton  $S^{(d+1)}$ .  $\square$

Let us now restrict to spaces rationally equivalent to products of rational Eilenberg-MacLane spaces for which it is shown quite easily that their rational homology is generated by products of spheres.

**Proposition 4.3.** *Let  $X$  denote a space that has the rational homotopy type of a product of simply-connected rational Eilenberg-MacLane spaces, then its rational homology is generated by products of spheres.*

*Proof.* Note that the rational cohomology ring of  $K(\mathbb{Q}, n)$  is either an exterior algebra with a single generator in degree  $n$  if  $n$  is odd, or a polynomial algebra generated by a single generator in degree  $n$  if  $n$  is even. Evidently, if  $n$  is odd, then every rational homology class is representable by spheres. If  $n$  is even, consider a generator in degree  $kn$  and let  $\iota_k: (S^n)^k \rightarrow K(\mathbb{Q}, n)$ , denote the map that classifies  $\alpha = \sum_{i=1}^k \pi_i^*([S^n]^*)$  where  $\pi_i$  is the projection onto the  $i$ -th factor in  $(S^n)^k$  and  $[S^n]^*$  denotes the dual fundamental class. Since  $\alpha^k$  is a non-zero multiple of the dual fundamental class of  $(S^n)^k$ , it follows that  $\iota_k$  induces an isomorphism in rational cohomology in degree  $kn$ . Hence  $\iota_k$  represents a multiple of the generator in the  $kn$ -th homology of  $K(\mathbb{Q}, n)$ .

Since the rational homology groups of a product are generated by cross products of classes in the factors it follows that any rational homology class in the product of simply-connected rational Eilenberg-MacLane spaces can be represented as a sum of classes representable by products of spheres.

Therefore [Lemma 4.2](#) yields the desired result.  $\square$

The following proposition ties the previous considerations to the main cases in question.

**Proposition 4.4.** *Let  $K$  denote a connected topological group whose underlying topological space is of finite type, then the classifying space  $BK$  has the rational homotopy type of a product of simply-connected rational Eilenberg-MacLane spaces. This class evidently contains compact groups and Lie groups.*

*If  $X$  is an aspherical topological space, then the classifying space of the identity component of its space of self-homotopy equivalences  $B\text{Aut}_0(X)$  is rationally homotopy equivalent to a  $K(V, 2)$ , where  $V = Z(\pi_1(X, *)) \otimes \mathbb{Q}$  and  $Z(\pi_1(X, *))$  denotes the center of the fundamental group of  $X$ .*

*Proof.* For the first part, note that by Borel-Serres theorem (See for example [5, Theorem 6.38]) the cohomology ring of  $BK$  is a polynomial ring with finitely many generators in even degree. Therefore the product of the maps classifying the generators  $BK \rightarrow \prod_i K(\mathbb{Q}, n_i)$  induces an isomorphism in rational cohomology by the Künneth theorem and hence a rational homotopy equivalence.

For the second part note that  $\pi_i(\text{Aut}(X), \text{Id})$  is  $\text{Out}(\pi_1(X, *))$  if  $i = 0$ ,  $Z(\pi_1(X, *))$  if  $i = 1$  and zero else. Therefore  $B\text{Aut}_0(X)$  is already a  $K(Z(\pi_1(X, )), 2)$ .  $\square$

As a consequence, we immediately obtain the following, which combined with [Theorem B](#) yields [Theorem A](#).

**Corollary 4.5.** *Let  $G$  denote either a connected compact group, a Lie group or the identity component of the homotopy automorphisms of an aspherical topological space, then the rational homology of  $BG$  is generated by products of spheres, in particular it is generated by fiber bundles over spheres.*

## REFERENCES

- [1] Richard H. jun. Escobales. Riemannian submersions with totally geodesic fibers. *J. Differ. Geom.*, 10:253–276, 1975.
- [2] M. Gromov. Volume and bounded cohomology. *Publ. Math., Inst. Hautes Étud. Sci.*, 56:5–99, 1982.
- [3] Dale H. Husemoller. *Fibre bundles.*, volume 20 of *Grad. Texts Math.* Berlin: Springer-Verlag, 3rd ed. edition, 1993.
- [4] Thorben Kastenholz and Jens Reinhold. Simplicial volume and essentiality of manifolds fibered over spheres. *J. Topol.*, 16(1):192–206, 2023.
- [5] John McCleary. *A user's guide to spectral sequences.*, volume 58 of *Camb. Stud. Adv. Math.* Cambridge: Cambridge University Press, 2nd ed. edition, 2001.
- [6] Mamoru Mimura, Ronald C. O'Neill, and Hirosi Toda. On p-equivalence in the sense of Serre. *Jpn. J. Math.*, 40:1–10, 1971.

KARLSRUHER INSTITUT FÜR TECHNOLOGIE, ENGLERSTRASSE 2, 76131 KARLSRUHE, GERMANY  
 Email address: thorben.kastenholz@kit.edu