AN AVERAGING FORMULA FOR NIELSEN NUMBERS OF AFFINE *n*-VALUED MAPS ON INFRA-NILMANIFOLDS

KAREL DEKIMPE AND LORE DE WEERDT

ABSTRACT. In [8, 9], the authors developed a nice formula to compute the Nielsen number of a self-map on an infra-nilmanifold. For the case of nilmanifolds this formula was extended to *n*-valued maps in [4]. In this paper, we extend these results further and establish the averaging formula to compute the Nielsen number of any *n*-valued affine map on an infra-nilmanifold.

INTRODUCTION

Let M be a closed manifold (or more generally a compact polyhedron) and let $f: M \to M$ be a self-map of M. In Nielsen-Reidemeister fixed point theory one attaches to each such map f a number N(f), the Nielsen number of f, which is a (sharp) lower bound for the number of fixed points of any map g which is homotopic to f. In this paper we are concerned with infra-nilmanifolds. It is known that any self-map on an infra-nilmanifold is homotopic to a so-called affine map ([10]). These maps are rather algebraic in nature and one has developed an averaging formula for computing their Nielsen number ([8, 9]). Since the Nielsen number of a map is invariant under a homotopy, this formula allows us to compute the Nielsen number of any self-map on an infra-nilmanifold. For *n*-valued maps the situation is more complicated. It is no longer true that any *n*-valued map on an infranilmanifold is homotopic to an affine n-valued map ([6]), but nevertheless, these affine *n*-valued maps still form a large and interesting class of *n*-valued maps on infra-nilmanifolds. In this paper, we obtain a natural generalisation of the formula from [9] for computing the Nielsen number of any affine nvalued map on an infra-nilmanifold.

In the first section we recall the basic Nielsen theory for *n*-valued maps and describe a general averaging formula (1.1) for computing the Nielsen number of an *n*-valued map. In the second section, we recall the necessary concepts concerning infra-nilmanifolds and affine *n*-valued maps. In the third section we establish a decomposition of the fixed point set of an *n*valued map on an infra-nilmanifold, which allows us to rewrite the general averaging formula into a more algebraic one (3.1). In the fourth section we further examine this formula in the case of affine *n*-valued maps, to finally arrive at the main result of this paper (Theorem 4.9), which is an averaging formula generalising in a natural way the formula from [9].

Research supported by Methusalem grant $\rm METH/21/03-long$ term structural funding of the Flemish Government.

1. Nielsen theory of n-valued maps

Let X be a connected compact manifold with fundamental group π and universal cover $p: \tilde{X} \to X$. We will view π as being the group of covering transformations of p.

An *n*-valued map $f : X \multimap X$ is a continuous (i.e. upper and lower semi-continuous) set-valued function, sending each point of X to a set of *n* distinct points in X. Equivalently, by the work of Brown and Gonçalves [2], the map f can be viewed as a continuous single-valued function from X to the unordered configuration space

$$D_n(X) = \{\{x_1, \dots, x_n\} \subseteq X \mid x_i \neq x_j \text{ if } i \neq j\},\$$

i.e. the quotient of the space $F_n(X) = \{(x_1, \ldots, x_n) \in X^n \mid x_i \neq x_j \text{ if } i \neq j\}$ under the action of the permutation group Σ_n . The fixed point set of an *n*-valued map f is $\text{Fix}(f) = \{x \in X \mid x \in f(x)\}.$

In what follows we recall some of the results of Brown et al. [1], where the study of fixed points of f was done by considering fixed points of *lifts* of f to the *orbit configuration space*

$$F_n(\tilde{X},\pi) = \{ (\tilde{x}_1,\ldots,\tilde{x}_n) \in \tilde{X}^n \mid p(\tilde{x}_i) \neq p(\tilde{x}_j) \text{ if } i \neq j \},\$$

which is a covering space of $D_n(X)$ with covering map

$$p^n: F_n(X,\pi) \to D_n(X): (\tilde{x}_1,\ldots,\tilde{x}_n) \mapsto \{p(\tilde{x}_1),\ldots,p(\tilde{x}_n)\}.$$

The corresponding covering group is $\pi^n \rtimes \Sigma_n$, which acts on $F_n(\tilde{X}, \pi)$ by

$$(\gamma_1,\ldots,\gamma_n;\sigma)(\tilde{x}_1,\ldots,\tilde{x}_n)=(\gamma_1\tilde{x}_{\sigma^{-1}(1)},\ldots,\gamma_n\tilde{x}_{\sigma^{-1}(n)})$$

A lift of f can then be defined as a map $\tilde{f}: \tilde{X} \to F_n(\tilde{X}, \pi)$ such that

$$\begin{array}{ccc} \tilde{X} & \stackrel{\tilde{f}}{\longrightarrow} & F_n(\tilde{X}, \pi) \\ p & & & \downarrow^{p^n} \\ X & \stackrel{f}{\longrightarrow} & D_n(X) \end{array}$$

commutes. As $F_n(\tilde{X}, \pi)$ is a subspace of \tilde{X}^n , such a lift $\tilde{f} : \tilde{X} \to F_n(\tilde{X}, \pi)$ splits into n maps $\tilde{f}_1, \ldots, \tilde{f}_n : \tilde{X} \to \tilde{X}$, called the *lift-factors* of \tilde{f} .

A lift $\tilde{f} = (\tilde{f}_1, \ldots, \tilde{f}_n)$ induces a morphism $(\phi_1, \ldots, \phi_n; \sigma) : \pi \to \pi^n \rtimes \Sigma_n$ of the covering groups so that for all $\gamma \in \pi$,

$$(\tilde{f}_1\gamma,\ldots,\tilde{f}_n\gamma)=(\phi_1(\gamma)\tilde{f}_{\sigma_{\gamma}^{-1}(1)},\ldots,\phi_n(\gamma)\tilde{f}_{\sigma_{\gamma}^{-1}(n)}).$$

Using this morphism, one can first subdivide $\{1, \ldots, n\}$ into σ -classes, the equivalence classes for the relation given by

$$i \sim j \iff \exists \gamma \in \pi : \sigma_{\gamma}(i) = j.$$

For all *i*, denote by $S_i = \{\gamma \in \pi \mid \sigma_{\gamma}(i) = i\}$ the stabilizer of *i* for this relation, which is a finite index subgroup of π . The restrictions of the maps ϕ_i to S_i are group morphisms, using which one can define an equivalence relation on π by

$$\alpha \sim \beta \iff \exists \gamma \in S_i : \alpha = \gamma \beta \phi_i(\gamma)^{-1}.$$

The set of equivalence classes for this relation is denoted $\mathcal{R}[\phi_i]$, and its number of elements is the *Reidemeister number* $R(\phi_i)$.

Remark 1.1. In general, given two group morphisms $\varphi, \psi : \Pi_1 \to \Pi_2$, one can consider the equivalence relation on Π_2 given by

$$\alpha \sim \beta \iff \exists \gamma \in \Pi_1 : \alpha = \psi(\gamma) \beta \varphi(\gamma)^{-1}.$$

The equivalence classes are called the *Reidemeister coincidence classes* of φ and ψ , and denoted $\mathcal{R}[\varphi, \psi]$. In case Π_1 is a subgroup of Π_2 and ψ is the inclusion $\iota : \Pi_1 \to \Pi_2$, we abbreviate $\mathcal{R}[\varphi, \iota] = \mathcal{R}[\varphi]$.

The σ -classes and Reidemeister classes of ϕ_i partition the fixed point set of f into a disjoint union of *fixed point classes*: if i_1, \ldots, i_r are representatives of the σ -classes, then

$$\operatorname{Fix}(f) = \bigsqcup_{\ell=1}^{r} \bigsqcup_{[\alpha] \in \mathcal{R}[\phi_{i_{\ell}}]} p(\operatorname{Fix}(\alpha \tilde{f}_{i_{\ell}})).$$

To each fixed point class one can associate an *index*, and the fixed point classes with non-zero index are called *essential*. The number of essential fixed point classes is the *Nielsen number* of f, which is a lower bound for the number of fixed points among all maps homotopic to f:

$$N(f) \le \min\{\#\operatorname{Fix}(g) \mid g \sim f\}.$$

By the above, the Nielsen number can be written as

$$N(f) = \sum_{\ell=1}^{r} \sum_{[\alpha] \in \mathcal{R}[\phi_{i_{\ell}}]} \varepsilon_{\alpha, i_{\ell}}$$

where

$$\varepsilon_{\alpha,i} = \begin{cases} 1 & \text{if } p(\operatorname{Fix}(\alpha \tilde{f}_i)) \text{ is an essential fixed point class of } f, \\ 0 & \text{otherwise.} \end{cases}$$

Since the representatives for the σ -classes were chosen arbitrarily, and choosing another representative i_{ℓ} for some ℓ must give the same result for N(f), the number

$$\sum_{[\alpha]\in\mathcal{R}[\phi_{i_{\ell}}]}\varepsilon_{\alpha,i_{\ell}}$$

does not depend on the chosen representative i_{ℓ} . Therefore the formula above can also be written as

(1.1)
$$N(f) = \sum_{i=1}^{n} \frac{1}{[\pi:S_i]} \sum_{[\alpha] \in \mathcal{R}[\phi_i]} \varepsilon_{\alpha,i}.$$

2. INFRA-NILMANIFOLDS AND AFFINE MAPS

The maps we consider in this paper are affine n-valued maps on infranilmanifolds. An infra-nilmanifold, as defined e.g. in [5], is the quotient of a connected and simply connected nilpotent Lie group G under the action of an almost-Bieberbach group $\pi \subseteq G \rtimes \operatorname{Aut}(G)$, i.e. a torsion free cocompact discrete subgroup of $G \rtimes C$, with C a maximal compact subgroup of $\operatorname{Aut}(G)$. Such an almost-Bieberbach group acts properly discontinuously on G and hence π is (isomorphic to) the fundamental group of the infra-nilmanifold $\pi \backslash G$. As before, we will view π as the covering group of the universal covering $G \to \pi \backslash G$. In the special case where this group π is entirely contained in G, the quotient space is called a nilmanifold. Every infra-nilmanifold $\pi \backslash G$ is finitely covered by a nilmanifold $N \backslash G$: it follows from the generalised Bieberbach Theorems that $\pi \cap G$ is of finite index in π , so one can take $N = \pi \cap G$ or any other subgroup $N \subseteq \pi \cap G$ which is of finite index in π . Such a group N is a finitely generated torsion free nilpotent group and is a lattice of the Lie group G.

Nilmanifolds have been studied in detail by Mal'cev; see [11]. We briefly recall some of his results.

For a nilpotent group G of class c, we denote the lower central series by

$$G = \gamma_1(G) \ge \ldots \ge \gamma_c(G) \ge \gamma_{c+1}(G) = 1.$$

For a finitely generated torsion free nilpotent group N of class c we consider another central series whose terms are the *isolators* of the lower central series,

$$N_i = \sqrt[N]{\gamma_i(N)} = \{g \in N \mid \exists k > 0 : g^k \in \gamma_i(N)\}.$$

This series has the useful property that for all i, the quotient N_i/N_{i+1} is free abelian of finite rank, say $N_i/N_{i+1} \cong \mathbb{Z}^{k_i}$. Moreover, for these numbers k_i , the factors of the lower central series of G satisfy $\gamma_i(G)/\gamma_{i+1}(G) \cong \mathbb{R}^{k_i}$. Morphisms of (lattices in) G induce morphisms on these quotients, which can be represented by $k_i \times k_i$ real (or integer) matrices.

If N, N' are lattices in G, any morphism $\phi : N \to N'$ extends uniquely to a Lie group morphism $G \to G$. The latter induces a morphism on the Lie algebra \mathfrak{g} of G. From now on we will refer to this as the Lie algebra morphism induced by $\phi : N \to N'$, and write $\phi_* : \mathfrak{g} \to \mathfrak{g}$, without explicitly mentioning the extension of ϕ to G.

Another useful property of connected and simply connected nilpotent Lie groups G is that the exponential map $\exp : \mathfrak{g} \to G$ is a diffeomorphism [3, Theorem 1.2.1]. In particular, it is invertible, and we denote the inverse map by $\log : G \to \mathfrak{g}$. This allows us to define the continuous function

$$G \times \mathbb{R} \to G : (g, t) \mapsto g^t = \exp(t \log(g))$$

which we will use later on.

On an infra-nilmanifold we can define the notion of an *affine n*-valued map. A map $f: \pi \setminus G \to D_n(\pi \setminus G)$ is affine if it lifts to a map of the form

$$f: G \to F_n(G, \pi): x \mapsto (g_1\varphi_1(x), \dots, g_n\varphi_n(x))$$

with $g_1, \ldots, g_n \in G$ and $\varphi_1, \ldots, \varphi_n \in \text{End}(G)$. We remark that this condition is independent of the chosen lift \tilde{f} . The corresponding maps ϕ_i and σ from section 1 satisfy

(2.1)
$$g_i\varphi_i(\gamma x) = \phi_i(\gamma)g_{\sigma_\gamma^{-1}(i)}\varphi_{\sigma_\gamma^{-1}(i)}(x)$$

for all $\gamma \in \pi$, $x \in G$ and $i \in \{1, \ldots, n\}$.

Nielsen numbers of affine n-valued maps on nilmanifolds have been studied in [4], where the following formula is proven: **Theorem 2.1** ([4, Theorem 6.13]). Let $N \setminus G$ be a nilmanifold. The Nielsen number of an affine map $f : N \setminus G \to D_n(N \setminus G)$ with lift $\tilde{f} : G \to F_n(G, N) :$ $x \mapsto (g_1\varphi_1(x), \ldots, g_n\varphi_n(x))$ is given by

$$N(f) = \sum_{i=1}^{n} |\det(I - (\varphi_i)_*)|.$$

The goal of this paper is to generalise this formula to infra-nilmanifolds. In the single-valued case, this extension from nilmanifolds to infra-nilmanifolds was done by Kim, Lee and Lee in [8]. They proved that any map f on an infra-nilmanifold $\pi \backslash G$ can be lifted to a nilmanifold $N \backslash G$ that finitely covers $\pi \backslash G$, and they expressed the Nielsen number of f as the average of the Nielsen numbers of these lifts.

We cannot follow the same approach in the *n*-valued case, since *n*-valued maps in general no longer admit lifts to a nilmanifold. However, we can decompose the formula for the Nielsen number in a similar way as in [8], and then plug in the knowledge about affine maps on nilmanifolds established in [4] to obtain the desired averaging formula without involving actual lifts to the nilmanifold.

3. Decomposition of the fixed point set

Let $\pi \backslash G$ be an infra-nilmanifold, finitely covered by a nilmanifold $N \backslash G$. For a general map $f : \pi \backslash G \to D_n(\pi \backslash G)$, we will decompose the sum (1.1) into a sum over Reidemeister classes in N and in π/N , generalising the approach in [8].

For all *i*, consider the group S'_i generated by $\{\gamma^{[\pi:N]} \mid \gamma \in S_i\}$. This is a normal subgroup of S_i , and the quotient group S_i/S'_i is periodic, since the order of every element divides $[\pi:N]$. Also note that π (and therefore also S_i/S'_i) is polycyclic-by-finite, since its finite index normal subgroup N is finitely generated and nilpotent. Since periodic polycyclic-by-finite groups are automatically finite, it follows that $[S_i:S'_i]$ is finite. Consequently, also

$$[N:S'_i] = \frac{[\pi:S_i][S_i:S'_i]}{[\pi:N]}$$

is finite. Thus, S'_i is a finite index normal subgroup of S_i and a finite index subgroup of N.

Furthermore, note that the image of S'_i under any morphism $S_i \to \pi$ is contained in N. Hence the inclusion map $\iota : S_i \to \pi$ and the morphism $\phi_i : S_i \to \pi$ induce morphisms

$$\begin{split} \iota':S'_i \to N:\gamma \mapsto \gamma & \bar{\iota}:S_i/S'_i \to \pi/N:\gamma S'_i \mapsto \gamma N \\ \phi'_i:S'_i \to N:\gamma \mapsto \phi_i(\gamma) & \bar{\phi}_i:S_i/S'_i \to \pi/N:\gamma S'_i \mapsto \phi_i(\gamma)N. \end{split}$$

We will decompose the set of Reidemeister classes of ϕ_i using Reidemeister classes associated to $\bar{\phi}_i$ and ϕ'_i .

First, note that

$$\mathcal{R}[\phi_i] = \bigcup_{\bar{\alpha} \in \pi/N} \bigcup_{g \in N} [g\alpha].$$

If $[g\alpha] = [g'\alpha']$ in $\mathcal{R}[\phi_i]$, then certainly $[\bar{\alpha}] = [\bar{\alpha}']$ in $\mathcal{R}[\bar{\phi}_i, \bar{\iota}]$, i.e.

$$\exists \bar{\gamma} \in S_i / S_i' : \bar{\alpha} = \bar{\iota}(\bar{\gamma}) \bar{\alpha}' \bar{\phi}_i(\bar{\gamma})^{-1}$$

Conversely, note that if $[\bar{\alpha}] = [\bar{\alpha}']$ in $\mathcal{R}[\bar{\phi}_i, \bar{\iota}]$, then there is a $g \in N$ so that $[g\alpha] = [\alpha']$ in $\mathcal{R}[\phi_i]$.

For fixed α , we have $[g\alpha] = [g'\alpha]$ in $\mathcal{R}[\phi_i]$ if and only if [g] = [g'] in $\mathcal{R}[\tau_\alpha \phi_i]$, with $\tau_\alpha : \pi \to \pi : \beta \mapsto \alpha \beta \alpha^{-1}$; i.e.

$$\exists \gamma \in S_i : g = \gamma g' \alpha \phi_i(\gamma)^{-1} \alpha^{-1}.$$

Thus, the above are disjoint unions when taken over these respective Reidemeister classes:

$$\mathcal{R}[\phi_i] = \bigsqcup_{[\bar{\alpha}] \in \mathcal{R}[\bar{\phi}_i, \bar{\iota}]} \bigsqcup_{[g] \in \mathcal{R}[\tau_\alpha \phi_i]} [g\alpha].$$

Rather than in $\mathcal{R}[\tau_{\alpha}\phi_i]$, we would like to view the classes [g] with $g \in N$ as elements of $\mathcal{R}[\tau_{\alpha}\phi'_i]$, where

$$g \sim g' \iff \exists \gamma \in S'_i : g = \gamma g' \alpha \phi'_i(\gamma)^{-1} \alpha^{-1}.$$

Since $S'_i \subseteq S_i$, it is clear that [g] = [g'] in $\mathcal{R}[\tau_\alpha \phi'_i]$ implies [g] = [g'] in $\mathcal{R}[\tau_\alpha \phi_i]$, but conversely there can be distinct elements in $\mathcal{R}[\tau_\alpha \phi'_i]$ that are equivalent in $\mathcal{R}[\tau_\alpha \phi_i]$. The following lemma allows us to describe how many different classes in $\mathcal{R}[\tau_\alpha \phi'_i]$ give rise to the same class in $\mathcal{R}[\tau_\alpha \phi_i]$. In this lemma, we use $\operatorname{coin}(\varphi, \psi) = \{\gamma \in \Pi_1 \mid \varphi(\gamma) = \psi(\gamma)\}$ to denote the group of coincidence points of two morphisms $\varphi, \psi : \Pi_1 \to \Pi_2$.

Lemma 3.1 ([7, Lemma 3.1.4]). Given a commutative diagram of groups with exact rows

$$1 \longrightarrow \Gamma_{1} \longrightarrow \Pi_{1} \longrightarrow \Pi_{1}/\Gamma_{1} \longrightarrow 1$$
$$\varphi' \downarrow \psi' \qquad \varphi \downarrow \psi \qquad \bar{\varphi} \downarrow \bar{\psi}$$
$$1 \longrightarrow \Gamma_{2} \longrightarrow \Pi_{2} \longrightarrow \Pi_{2}/\Gamma_{2} \longrightarrow 1$$

such that the quotient groups Π_i/Γ_i are finite. For all $\alpha \in \Pi_2$ and $g \in \Gamma_2$, the number of Reidemeister coincidence classes $[g'] \in \mathcal{R}[\tau_\alpha \varphi', \psi']$ such that [g] = [g'] in $\mathcal{R}[\tau_\alpha \varphi, \psi]$ is equal to

$$[\operatorname{coin}(\tau_{\bar{\alpha}}\bar{\varphi},\bar{\psi}):\overline{\operatorname{coin}(\tau_{g\alpha}\varphi,\psi)}]$$

where $\overline{\operatorname{coin}(\tau_{g\alpha}\varphi,\psi)}$ denotes the image of $\operatorname{coin}(\tau_{g\alpha}\varphi,\psi)$ in Π_1/Γ_1 .

Applying this to the diagram

$$1 \longrightarrow S'_{i} \longrightarrow S_{i} \longrightarrow S_{i}/S'_{i} \longrightarrow 1$$
$$\begin{array}{c} \phi'_{i} \downarrow^{\iota'} & \phi_{i} \downarrow^{\iota} & \bar{\phi}_{i} \downarrow^{\bar{\iota}} \\ 1 \longrightarrow N \longrightarrow \pi \longrightarrow \pi/N \longrightarrow 1 \end{array}$$

learns that for a given $g \in N$, the number of classes $[g'] \in \mathcal{R}[\tau_{\alpha}\phi'_i]$ such that [g] = [g'] in $\mathcal{R}[\tau_{\alpha}\phi_i]$ is equal to

$$\left[\operatorname{coin}(\tau_{\bar{\alpha}}\bar{\phi}_i,\bar{\iota}):\overline{\operatorname{fix}(\tau_{g\alpha}\phi_i)}\right] = \frac{\left|\operatorname{coin}(\tau_{\bar{\alpha}}\bar{\phi}_i,\bar{\iota})\right|}{\left|\operatorname{fix}(\tau_{g\alpha}\phi_i)\right|}$$

where $\overline{\operatorname{fix}(\tau_{g\alpha}\phi_i)}$ denotes the image of $\operatorname{fix}(\tau_{g\alpha}\phi_i) = \operatorname{coin}(\tau_{g\alpha}\phi_i,\iota)$ in S_i/S'_i . Therefore the sum (1.1) decomposes as

$$N(f) = \sum_{i=1}^{n} \frac{1}{[\pi:S_i]} \sum_{[\bar{\alpha}]\in\mathcal{R}[\bar{\phi}_i,\bar{\iota}]} \sum_{[g]\in\mathcal{R}[\tau_{\alpha}\phi'_i]} \frac{|\overline{\mathrm{fix}(\tau_{g\alpha}\phi_i)}|}{|\mathrm{coin}(\tau_{\bar{\alpha}}\bar{\phi}_i,\bar{\iota})|} \varepsilon_{g\alpha,i}.$$

To further simplify this formula, consider the action of S_i/S'_i on π/N given by

$$\bar{\gamma} \cdot \bar{\alpha} = \bar{\iota}(\bar{\gamma})\bar{\alpha}\bar{\phi}_i(\bar{\gamma})^{-1}$$

The orbit of an element $\bar{\alpha}$ for this action is its Reidemeister class $[\bar{\alpha}] \in \mathcal{R}[\bar{\phi}_i, \bar{\iota}]$. The stabilizer of $\bar{\alpha}$ is the set

$$\{\bar{\gamma} \in S_i / S_i' \mid \bar{\iota}(\bar{\gamma})\bar{\alpha}\bar{\phi}_i(\bar{\gamma})^{-1} = \bar{\alpha}\} = \operatorname{coin}(\tau_{\bar{\alpha}}\bar{\phi}_i, \bar{\iota}).$$

Therefore we have

$$[\bar{\alpha}]| \cdot |\operatorname{coin}(\tau_{\bar{\alpha}}\bar{\phi}_i,\bar{\iota})| = |S_i/S_i'| = [S_i:S_i'].$$

This reduces the expression for N(f) to

$$N(f) = \sum_{i=1}^{n} \frac{1}{[\pi : S_i]} \sum_{\bar{\alpha} \in \pi/N} \frac{1}{|[\bar{\alpha}]|} \sum_{[g] \in \mathcal{R}[\tau_{\alpha}\phi'_i]} \frac{|\operatorname{fix}(\tau_{g\alpha}\phi_i)|}{|\operatorname{coin}(\tau_{\bar{\alpha}}\bar{\phi}_i,\bar{\iota})|} \varepsilon_{g\alpha,i}$$

$$(3.1) \qquad = \sum_{i=1}^{n} \frac{1}{[\pi : S'_i]} \sum_{\bar{\alpha} \in \pi/N} \sum_{[g] \in \mathcal{R}[\tau_{\alpha}\phi'_i]} |\overline{\operatorname{fix}(\tau_{g\alpha}\phi_i)}| \varepsilon_{g\alpha,i}.$$

4. Averaging formula for Affine maps

Now we apply this to the special case where f is an affine map with lift $\tilde{f}: G \to F_n(G, \pi): x \mapsto (g_1\varphi_1(x), \ldots, g_n\varphi_n(x))$. In this case, the formula above can be reduced to an expression purely in terms of the morphisms φ_i .

Recall that, for a morphism $\phi' : S' \to N$ between lattices in a connected and simply connected nilpotent Lie group G, we use $(\phi')_* : \mathfrak{g} \to \mathfrak{g}$ to denote the Lie algebra morphism induced by the unique extension of ϕ' to $G \to G$.

Lemma 4.1. Suppose $\pi \setminus G$ is an infra-nilmanifold finitely covered by a nilmanifold $N \setminus G$, and S is a finite index subgroup of π . Let S' be the subgroup generated by $\{\gamma^{[\pi:N]} \mid \gamma \in S\}$. Let $\phi : S \to \pi$ be a morphism and $\phi' : S' \to N$ its restriction to S'. If det $(I - (\phi')_*) \neq 0$, then fix $(\phi) = 1$.

Proof. We first prove that $fix(\phi') = 1$. To this end, let $exp : \mathfrak{g} \to G$ denote the exponential map of G, and $\log : G \to \mathfrak{g}$ its inverse.

Suppose $\gamma \in S'$ is a fixed point of ϕ' . Then $X = \log(\gamma)$ is a fixed point of $(\phi')_*$, i.e. $(I - (\phi')_*)(X) = 0$. Since $\det(I - (\phi')_*) \neq 0$, the only element X satisfying this equation is 0. It follows that $\gamma = \exp(0) = 1$.

Now, suppose $\gamma \in S$ is a fixed point of ϕ , and denote $k = [\pi : N]$. Then $\phi(\gamma^k) = \phi(\gamma)^k = \gamma^k$, so $\gamma^k \in S'$ is a fixed point of ϕ' . By the above, we have $\gamma^k = 1$. Since S is torsion free, it follows that $\gamma = 1$.

We will now show that for an affine map, the number $\varepsilon_{g\alpha,i}$ in (3.1) is independent of $g \in N$, using the results from [4]. We distinguish two cases. In both cases we will need the following lemma. **Lemma 4.2** ([4, Lemma 6.4]). Let G be a connected and simply connected nilpotent Lie group, and $\varphi \in \text{End}(G)$. If φ_* has no eigenvalue 1, the map $G \to G : x \mapsto \varphi(x)x^{-1}$ is bijective.

The first case is the following.

Notation 4.3. For $\alpha \in \pi \subseteq G \rtimes \operatorname{Aut}(G)$ we will write $\alpha = (a_{\alpha}, A_{\alpha})$ with $a_{\alpha} \in G$ and $A_{\alpha} \in \operatorname{Aut}(G)$.

Corollary 4.4. Let $f : \pi \setminus G \to D_n(\pi \setminus G)$ be an affine *n*-valued map with lift $\tilde{f} : x \mapsto (g_1\varphi_1(x), \ldots, g_n\varphi_n(x))$ and corresponding numbers $\varepsilon_{g\alpha,i}$. Fix *i* and $\alpha \in \pi$. If det $(I - (A_\alpha \varphi_i)_*) \neq 0$, then $\varepsilon_{g\alpha,i} = 1$ for all $g \in N$.

Proof. Pick $g \in N$. Note that for all $x \in G$,

$$g\alpha f_i(x) = g\alpha g_i \varphi_i(x) = ga_\alpha A_\alpha(g_i) A_\alpha \varphi_i(x).$$

So if we denote $ga_{\alpha}A_{\alpha}(g_i) = g'$, then $g\alpha \tilde{f}_i = g'A_{\alpha}\varphi_i$.

By Lemma 4.2, the map $G \to G : x \mapsto A_{\alpha}\varphi_i(x)x^{-1}$ is a bijection. Hence there is a unique $x \in G$ such that $A_{\alpha}\varphi_i(x)x^{-1} = (g')^{-1}$, i.e. such that $g'A_{\alpha}\varphi_i(x) = x$. That is, the map $g\alpha \tilde{f}_i = g'A_{\alpha}\varphi_i$ has a unique fixed point. The index of the fixed point class $p(\operatorname{Fix}(g\alpha \tilde{f}_i))$ is then equal to the index of this unique fixed point of $g'A_{\alpha}\varphi_i$.

In [4, Lemma 6.10] it is proven that, if $\varphi : G \to G$ is a morphism so that $G \to G : x \mapsto \varphi(x)x^{-1}$ is a bijection, then for any $h \in G$, the index of $G \to G : x \mapsto h\varphi(x)$ at its unique fixed point is ± 1 . Applying this to $\varphi = A_{\alpha}\varphi_i$ and h = g' gives the desired result.

For the second case, we can use the following lemma, which is a special case of [4, Lemma 6.6]. We give a more polished version of the proof adapted to our specific needs.

Lemma 4.5. Let G be a connected and simply connected nilpotent Lie group, and $\varphi \in \text{End}(G)$. If φ_* has eigenvalue 1, then for all $g \in G$ there is a $g_0 \in G$ such that $g\varphi(x)g_0 \neq x$ for all $x \in G$.

Proof. We prove the claim by induction on the nilpotency class c of G.

If c = 1, then $G = \mathbb{R}^k$ for some k and $\varphi = \varphi_*$ is a linear map represented by a matrix which has eigenvalue 1. We can choose a basis $\{v_1, \ldots, v_k\}$ of \mathbb{R}^k with respect to which the matrix of φ is of the form

$$\begin{bmatrix} 1 & 0 & \cdots & 0 \\ * & * & \cdots & * \\ \vdots & \vdots & \ddots & \vdots \\ * & * & \cdots & * \end{bmatrix}.$$

That is, if we write $V = \langle v_2, \ldots, v_k \rangle$, then $\varphi(\mu v_1 + v) \equiv \mu v_1 \mod V$ for all $\mu \in \mathbb{R}$ and $v \in V$.

Take $\lambda \in \mathbb{R}$ such that $g \equiv \lambda v_1 \mod V$. We claim that $g_0 = (1 - \lambda)v_1$ does the job. Indeed, suppose $x \in \mathbb{R}^k$ satisfies $g + \varphi(x) + g_0 = x$. If we write $x = \mu v_1 + v$ with $v \in V$, this reduces to

$$\lambda v_1 + \mu v_1 + (1 - \lambda)v_1 = \mu v_1 \mod V,$$

or equivalently, $v_1 \in V$, which is a contradiction.

Now suppose c > 1 and the result holds for all groups of smaller nilpotency class. For all i = 1, ..., c, let M_i be the matrix of the morphism induced by φ on $\gamma_i(G)/\gamma_{i+1}(G)$. We distinguish two cases: either one of the matrices $M_1, ..., M_{c-1}$ has eigenvalue 1, or M_c is the only matrix with eigenvalue 1. In the first case, we can apply the induction hypothesis to the map $G/\gamma_c(G) \to G/\gamma_c(G) : \bar{x} \mapsto \overline{\varphi(x)}$ (and the element \bar{g} for g) to obtain a $g_0 \in G$ such that $g\varphi(x)g_0 \neq \bar{x}$ in $G/\gamma_c(G)$, so in particular $g\varphi(x)g_0 \neq x$ in G, for all $x \in G$.

In the second case, we can apply the induction hypothesis to the restriction of φ to $\gamma_c(G)$ (and the identity element for g) to obtain a $g_0 \in \gamma_c(G)$ such that $\varphi(x)g_0 \neq x$ for all $x \in \gamma_c(G)$. On the other hand, we can apply Lemma 4.2 to the map $G/\gamma_c(G) \rightarrow G/\gamma_c(G) : \bar{x} \mapsto \overline{\varphi(x)}$, which has no eigenvalue 1. It follows that there is a unique $\bar{x}_0 \in G/\gamma_c(G)$ such that $\overline{x_0\varphi(x_0)^{-1}} = \bar{g}$. Choose a representative $x_0 \in G$ for \bar{x}_0 , and take $z \in \gamma_c(G)$ such that $g\varphi(x_0) = x_0 z$. We claim that the element $g_0 z^{-1}$ does the job. Indeed, suppose $x \in G$ satisfies $g\varphi(x)g_0 z^{-1} = x$. By uniqueness of \bar{x}_0 , this element x must be of the form $x_0 z'$ for some $z' \in \gamma_c(G)$. Then we have

$$g\varphi(x_0z')g_0z^{-1} = x_0z' \iff g\varphi(x_0)\varphi(z')g_0z^{-1} = x_0z'$$
$$\Leftrightarrow x_0z\varphi(z')g_0z^{-1} = x_0z'$$
$$\Leftrightarrow \varphi(z')g_0 = z',$$

which is impossible for $z' \in \gamma_c(G)$, by definition of g_0 .

Corollary 4.6. Let $f : \pi \setminus G \to D_n(\pi \setminus G)$ be an affine n-valued map with lift $\tilde{f} : x \mapsto (g_1\varphi_1(x), \ldots, g_n\varphi_n(x))$ and corresponding numbers $\varepsilon_{g\alpha,i}$. Fix i and $\alpha \in \pi$. If det $(I - (A_\alpha \varphi_i)_*) = 0$, then $\varepsilon_{g\alpha,i} = 0$ for all $g \in N$.

Proof. Pick $g \in N$ and define g' as in Corollary 4.4. By Lemma 4.5, there exits a $g_0 \in G$ such that $g'A_{\alpha}\varphi_i(x)g_0 \neq x$ for all $x \in G$. Then we have a homotopy

$$\pi \backslash G \times [0,1] \to D_n(\pi \backslash G) : (p(x),t) \mapsto \{p(\tilde{f}_1(x)g_0^t), \dots, p(\tilde{f}_n(x)g_0^t)\}.$$

Under this homotopy, the fixed point class $p(\text{Fix}(g\alpha \tilde{f}_i))$ of f corresponds to the fixed point class $p(\text{Fix}(g\alpha \tilde{f}_i g_0))$. Since $g\alpha \tilde{f}_i g_0$ has no fixed points by definition of g_0 , the index of this fixed point class is 0. As indices are preserved under homotopies, the index of $p(\text{Fix}(g\alpha \tilde{f}_i))$ is also 0.

On the other hand, the results in [4] give an expression for the number of terms in the second sum in (3.1):

Lemma 4.7. For a morphism $\phi' : S' \to N$, with N finitely generated, torsion free and nilpotent, and S' a finite index subgroup of N, one has

$$R(\phi') = [N : S'] |\det(I - (\phi')_*)|_{\infty},$$

where $|a|_{\infty}$ is defined as |a| if $a \neq 0$, and ∞ if a = 0.

Proof. Let c be the nilpotency class of N. For $i = 1, \ldots, c$, write $N_i = \sqrt[N]{\gamma_i(N)}$ and $S'_i = \sqrt[S']{\gamma_i(S')}$. If M_i denotes the matrix of the morphism $S'_i N_{i+1}/N_{i+1} \to N_i/N_{i+1}$ induced by ϕ' , it is proven in [4, Theorem 4.14] that

$$R(\phi') = [N:S'] |\det(I-M)|_{\infty},$$

where

$$M = \begin{bmatrix} M_1 & 0 & \cdots & 0 \\ 0 & M_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & M_c \end{bmatrix}.$$

On the other hand, it is proven in [4, Theorem 5.4] that the matrix of $(\phi')_*$ with respect to a well-chosen basis for \mathfrak{g} is of the form

$$\begin{bmatrix} M_1 & 0 & \cdots & 0 \\ * & M_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ * & * & \cdots & M_c \end{bmatrix}$$

Thus, we can replace $\det(I - M)$ by $\det(I - (\phi')_*)$, which gives the desired formula.

Lastly, we observe:

Lemma 4.8. Let $f : \pi \backslash G \to D_n(\pi \backslash G)$ be an affine n-valued map with lift $\tilde{f} : x \mapsto (g_1 \varphi_1(x), \ldots, g_n \varphi_n(x))$ and corresponding morphisms $\phi'_i : S'_i \to N$. For all i and $\alpha \in \pi$, for all $g \in N$,

$$\det(I - (\tau_{q\alpha}\phi'_i)_*) = \det(I - (A_\alpha\varphi_i)_*).$$

Proof. We may choose a basis $v_{1,1}, v_{1,2}, \ldots, v_{1,k_1}, v_{2,1}, \ldots, v_{c,k_c}$ for \mathfrak{g} in such a way that the vectors $v_{i,1}, v_{i,2}, \ldots, v_{i,k_i}, v_{i+1,1}, \ldots, v_{c,k_c}$ form a basis of $\gamma_i(\mathfrak{g})$, the *i*-th term in the lower central series of the Lie algebra \mathfrak{g} . With respect to this basis, all morphisms $\mathfrak{g} \to \mathfrak{g}$ are represented by blocked lower triangular matrices (with blocks of size $k_i \times k_i$ on the diagonal).

Now note that for the morphisms $\tau_{g\alpha}$, $\phi'_i : S'_i \to N$, and therefore also for their extensions $G \to G$,

$$\tau_{g\alpha} = c_{ga_{\alpha}} A_{\alpha}$$
 and $\phi'_i = c_{g_i} \varphi_i$

where $c_g: G \to G$ denotes conjugation with $g \in G$. (The equality for ϕ_i follows by evaluating (2.1) at x = 1 and $\gamma \in S'_i$.)

For all $g \in G$, the map $c_g : G \to G$ induces the identity on $\gamma_i(G)/\gamma_{i+1}(G)$, so the maps $(c_g)_*$ induce the identity on $\gamma_i(\mathfrak{g})/\gamma_{i+1}(\mathfrak{g})$ and the matrix representation of $(c_g)_*$ has identity blocks on the diagonal. It follows that the matrices of $(\tau_{g\alpha})_*$ and $(\phi'_i)_*$ have the same blocks on the diagonal as those of $(A_\alpha)_*$ and $(\varphi_i)_*$, respectively; therefore

$$\det(I - (\tau_{q\alpha}\phi'_i)_*) = \det(I - (A_{\alpha}\varphi_i)_*).$$

Putting all these results together gives the averaging formula:

Theorem 4.9. The Nielsen number of an affine map $f : \pi \setminus G \to D_n(\pi \setminus G)$ with lift $\tilde{f} : x \mapsto (g_1\varphi_1(x), \ldots, g_n\varphi_n(x))$ is given by

$$N(f) = \frac{1}{[\pi : N]} \sum_{\bar{\alpha} \in \pi/N} \sum_{i=1}^{n} |\det(I - (A_{\alpha}\varphi_i)_*)|.$$

10

Proof. Recall from section 3 that

$$N(f) = \sum_{i=1}^{n} \frac{1}{[\pi : S'_i]} \sum_{\bar{\alpha} \in \pi/N} \sum_{[g] \in \mathcal{R}[\tau_{\alpha} \phi'_i]} |\overline{\operatorname{fix}(\tau_{g\alpha} \phi_i)}| \varepsilon_{g\alpha,i}.$$

In case det $(I - (A_{\alpha}\varphi_i)_*) \neq 0$, Lemma 4.1 implies fix $(\tau_{g\alpha}\phi_i) = 1$, so in particular $|\overline{\text{fix}(\tau_{g\alpha}\phi_i)}| = 1$; Corollary 4.4 implies $\varepsilon_{g\alpha,i} = 1$ for all g; and Lemmas 4.7 and 4.8 imply

$$R(\tau_{\alpha}\phi'_{i}) = [N:S'_{i}]|\det(I - (A_{\alpha}\varphi_{i})_{*})|.$$

Thus, in this case,

$$\sum_{[g]\in\mathcal{R}[\tau_{\alpha}\phi'_{i}]} |\overline{\operatorname{fix}(\tau_{g\alpha}\phi_{i})}| \varepsilon_{g\alpha,i} = [N:S'_{i}]|\det(I - (A_{\alpha}\varphi_{i})_{*})|.$$

In case det $(I - (A_{\alpha}\varphi_i)_*) = 0$, Corollary 4.6 implies $\varepsilon_{g\alpha,i} = 0$ for all g, so

$$\sum_{[g]\in\mathcal{R}[\tau_{\alpha}\phi_{i}']} |\overline{\operatorname{fix}(\tau_{g\alpha}\phi_{i})}| \varepsilon_{g\alpha,i} = 0 = [N:S_{i}']|\det(I - (A_{\alpha}\varphi_{i})_{*})|.$$

It follows that

$$N(f) = \sum_{i=1}^{n} \frac{1}{[\pi : S'_i]} \sum_{\bar{\alpha} \in \pi/N} [N : S'_i] \det(I - (A_\alpha \varphi_i)_*)|$$
$$= \frac{1}{[\pi : N]} \sum_{\bar{\alpha} \in \pi/N} \sum_{i=1}^{n} |\det(I - (A_\alpha \varphi_i)_*)|.$$

5. Example

To illustrate the formula, consider the following example. Take $G = \mathbb{R}^2$ and let $\pi \subseteq \mathbb{R}^2 \rtimes \operatorname{GL}_2(\mathbb{R})$ be the group generated by

$$a = \left(\begin{bmatrix} 1\\0 \end{bmatrix}, \begin{bmatrix} 1&0\\0&1 \end{bmatrix} \right), \quad b = \left(\begin{bmatrix} 0\\\frac{1}{2} \end{bmatrix}, \begin{bmatrix} -1&0\\0&1 \end{bmatrix} \right)$$

Then $\pi \setminus \mathbb{R}^2$ is an infra-nilmanifold isomorphic to the Klein bottle. Consider the map given by

$$\tilde{f}: \mathbb{R}^2 \to F_2(\mathbb{R}^2, \pi): (t_1, t_2) \mapsto ((0, \frac{t_2}{2}), (0, \frac{t_2}{2} - \frac{1}{4})).$$

This map induces a well-defined 2-valued map $f : \pi \setminus \mathbb{R}^2 \to D_2(\pi \setminus \mathbb{R}^2)$ on the Klein bottle, since for all $(t_1, t_2) \in \mathbb{R}^2$

$$\begin{split} \tilde{f}(a(t_1, t_2)) &= ((0, \frac{t_2}{2}), (0, \frac{t_2}{2} - \frac{1}{4})) = (1, 1; 1)\tilde{f}(t_1, t_2) \\ \tilde{f}(b(t_1, t_2)) &= (b(0, \frac{t_2}{2} - \frac{1}{4}), (0, \frac{t_2}{2})) = (b, 1; \sigma)\tilde{f}(t_1, t_2) \end{split}$$

with $\sigma \in \Sigma_2$ the permutation that interchanges 1 and 2.

Note that f is an affine map, with

$$g_1 = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \quad g_2 = \begin{bmatrix} 0 \\ -\frac{1}{4} \end{bmatrix}, \quad \varphi_1 = \varphi_2 = \begin{bmatrix} 0 & 0 \\ 0 & \frac{1}{2} \end{bmatrix}.$$

To compute the Nielsen number of f, we consider the nilmanifold $Z \setminus \mathbb{R}^2$ where $Z = \pi \cap \mathbb{R}^2 = \langle a, b^2 \rangle \cong \mathbb{Z}^2$, i.e. the 2-torus. The group π/Z has two elements, represented by

$$\left(\begin{bmatrix}0\\0\end{bmatrix}, \begin{bmatrix}1&0\\0&1\end{bmatrix}\right) \quad \text{and} \quad \left(\begin{bmatrix}0\\0\end{bmatrix}, \begin{bmatrix}-1&0\\0&1\end{bmatrix}\right).$$

Thus, by Theorem 4.9

$$\begin{split} N(f) &= \frac{1}{2} \sum_{\bar{\alpha} \in \pi/Z} \sum_{i=1}^{2} \left| \det \left(I - A_{\alpha} \begin{bmatrix} 0 & 0 \\ 0 & \frac{1}{2} \end{bmatrix} \right) \right| \\ &= \sum_{\bar{\alpha} \in \pi/Z} \left| \det \left(I - A_{\alpha} \begin{bmatrix} 0 & 0 \\ 0 & \frac{1}{2} \end{bmatrix} \right) \right| \\ &= \left| \det \left(I - \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & \frac{1}{2} \end{bmatrix} \right) \right| + \left| \det \left(I - \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & \frac{1}{2} \end{bmatrix} \right) \right| \\ &= \left| \det \begin{bmatrix} 1 & 0 \\ 0 & \frac{1}{2} \end{bmatrix} \right| + \left| \det \begin{bmatrix} 1 & 0 \\ 0 & \frac{1}{2} \end{bmatrix} \right| = 1. \end{split}$$

A direct calculation shows that the map f has one isolated fixed point: indeed, for all $k,\ell\in\mathbb{Z}$ we have

$$(t_1, t_2) = a^k b^\ell (0, \frac{t_2}{2}) \quad \Leftrightarrow \quad (t_1, t_2) = (k, \frac{t_2}{2} + \frac{\ell}{2}) \\ \Leftrightarrow \quad (t_1, t_2) = (k, \ell)$$

and

$$(t_1, t_2) = a^k b^\ell (0, \frac{t_2}{2} - \frac{1}{4}) \iff (t_1, t_2) = (k, \frac{t_2}{2} - \frac{1}{4} + \frac{\ell}{2}) \Leftrightarrow (t_1, t_2) = (k, -\frac{1}{2} + \ell);$$

and all these points (k, ℓ) and $(k, -\frac{1}{2} + \ell)$ in \mathbb{R}^2 project to the same point on the Klein bottle. Thus, the minimal number of fixed points is actually attained in this case.

Remark 5.1. This map f is an example of a map on the Klein bottle that does not lift to the torus $Z \setminus \mathbb{R}^2$. In fact, there is no nilmanifold (torus) $N \setminus \mathbb{R}^2$ that finitely covers $\pi \setminus \mathbb{R}^2$ so that f lifts to $N \setminus \mathbb{R}^2$: in order for such a lift to exist, the induced morphism of covering groups $f_* = (\phi_1, \phi_2; \sigma) : \pi \to \pi^2 \rtimes \Sigma_2$ defined in section 1 must satisfy $f_*(N) \subseteq N^2 \rtimes \Sigma_2$. However, note that $f_*(b^2) = (b, b; 1)$. If $N \setminus \mathbb{R}^2$ is a nilmanifold that finitely covers $\pi \setminus \mathbb{R}^2$, then Nmust be a finite index subgroup of Z. Hence there is a smallest $\ell \in \mathbb{Z}_{>0}$ so that $(b^2)^{\ell} \in N$. But then $f_*((b^2)^{\ell}) = (b^{\ell}, b^{\ell}; 1)$ is not an element of $N^2 \rtimes \Sigma_2$, by minimality of ℓ .

This illustrates the fact that a classical averaging formula in terms of lifts as in [8] is not feasible in the *n*-valued case.

References

- R. F. Brown, C. Deconinck, K. Dekimpe, and P. C. Staecker. Lifting Classes for the Fixed Point Theory of n-Valued Maps. *Topology and its Applications*, 274:107–125, 2020.
- [2] R. F. Brown and D. L. Gonçalves. On the topology of n-valued maps. Advanced Fixed Point Theory, 8(2):205-220, 2018.

- [3] L. Corwin and F. P. Greenleaf. Representations of nilpotent Lie groups and their applications: Volume 1, Part 1, Basic theory and examples, volume 18. Cambridge University Press, 1990.
- [4] C. Deconinck and K. Dekimpe. Nielsen numbers of affine n-valued maps on nilmanifolds. Journal of Fixed Point Theory and Applications, 25(4):Paper No. 84, 29, 2023.
- [5] K. Dekimpe. A users' guide to infra-nilmanifolds and almost-Bieberbach groups, volume 40 of Adv. Lect. Math. (ALM), Handbook of group actions. Vol. III, pages 215– 262. Int. Press, Somerville, MA, 2018.
- [6] K. Dekimpe and L. D. Weerdt. Non-affine n-valued maps on tori. arXiv:2401.05053 (to appear in Topology and its Applications), 2024.
- [7] K. Y. Ha, J. B. Lee, and P. Penninckx. Formulas for the Reidemeister, Lefschetz and Nielsen coincidence number of maps between infra-nilmanifolds. *Journal of Fixed Point Theory and Applications*, 2012(39):1–23, 2012.
- [8] S. W. Kim, J. B. Lee, and K. B. Lee. Averaging formula for Nielsen numbers. Nagoya Mathematical Journal, 178:37–53, 2005.
- J. B. Lee and K. B. Lee. Lefschetz numbers for continuous maps, and periods for expanding maps on infra-nilmanifolds. *Journal of Geometry and Physics*, 56(10):2011–2023, 2006.
- [10] K. B. Lee. Maps on infra-nilmanifolds. Pacific Journal of Mathematics, 168(1):157– 166, 1995.
- [11] A. I. Mal'cev. On a class of homogeneous spaces. Izvestiya Akademii Nauk SSSR. Seriya Matematicheskaya, 13:9–32, 1949.

KU LEUVEN CAMPUS KULAK KORTRIJK, 8500 KORTRIJK, BELGIUM Email address: Karel.Dekimpe@kuleuven.be Email address: Lore.DeWeerdt@kuleuven.be