(Non-)Vanishing of high-dimensional group cohomology

Benjamin Brück

Church–Farb–Putman formulated stability and vanishing conjectures for the high-dimensional cohomology of $SL_n(\mathbb{Z})$, surface mapping class groups and automorphism groups of free groups. This is a survey on the current status of these conjectures and their generalisations.

1 Introduction

This article is concerned with the rational cohomology of groups "similar to $SL_n(\mathbb{Z})$ " such as the surface mapping class group $MCG(\Sigma_g)$ and the automorphism group of the free group $Aut(F_n)$. Each such group Γ has finite virtual cohomological dimension $vcd(\Gamma) \in \mathbb{N}$. This implies that its rational cohomology can only be non-trivial in finitely many degrees because it vanishes in degrees above $vcd(\Gamma)$,

$$H^k(\Gamma; \mathbb{Q}) = 0$$
 for $k > \operatorname{vcd}(\Gamma)$.

The low-dimensional cohomology ($k \ll \operatorname{vcd}(\Gamma)$) of these groups is comparably well understood, mainly through homological stability results. However, much less is known about their high-dimensional cohomology ($k \approx \operatorname{vcd}(\Gamma)$). In [23], Church–Farb–Putman conjectured certain patterns in this unstable cohomology.

Duality. The point of departure for these conjectures is that $SL_n(\mathbb{Z})$, $MCG(\Sigma_g)$ and $Aut(F_n)$ are all virtual Bieri–Eckmann duality groups, which means that they satisfy an analogue of Poincaré duality. For $SL_n(\mathbb{Z})$, this follows from work by Borel–Serre [6]: Let R be a number ring, i.e. the ring of integers in a number field \mathbb{K} , and \mathcal{G} a Chevalley–Demazure group scheme (for example $\mathcal{G} = SL_n$, Sp_{2n} , PGL_n , $Spin_{2n+1}$ or SO_{2n}). Then $\Gamma = \mathcal{G}(R)$ is an arithmetic subgroup of $\mathcal{G}(\mathbb{K})$ and Borel–Serre's result implies that for all codimensions $0 \le i \le vcd(\Gamma) < \infty$, there is an isomorphism

$$H^{\operatorname{vcd}(\Gamma)-i}(\Gamma; \mathbb{Q}) \cong H_i(\Gamma; \operatorname{St}(\Gamma) \otimes \mathbb{Q}).$$
 (1)

Here $St(\Gamma)$ is the *Steinberg module*, i.e. the top-degree homology of the Tits building $\Delta(\Gamma)$ associated to $\mathcal{G}(\mathbb{K})$. The duality given by Eq. (1) allows one to study high-dimensional cohomology with rational coefficients by investigating low-dimensional

²⁰²⁰ Mathematics Subject Classification. Primary 57M07; Secondary 11F75, 20F28, 57K20. *Keywords.* group cohomology, duality groups, Steinberg module.

¹This building is a highly symmetric simplicial Γ-complex. For $\Gamma = SL_n(R)$ and $Sp_{2n}(R)$, definitions are given in Section 2.2.2 and Section 6.2.

homology with coefficients in the dualising module $St(\Gamma)$. The trade-off is that the latter requires a good understanding of $St(\Gamma)$, for example by knowing a sufficiently nice generating set, presentation or more generally a partial resolution.

This article. The aim of this article is to give an overview of progress that has been made since Church–Farb–Putman's article [23] appeared. On the one hand, this text collects vanishing and non-vanishing results for the high-dimensional cohomology of various duality groups. On the other hand, it describes partial resolutions of the dualising modules that were used to obtain some of these results.

Section 2, Section 3 and Section 4 contain the original conjectures of Church–Farb–Putman and explain their current status. Section 5, Section 6 and Section 7 give generalisations to different types of Chevalley groups that have recently emerged. Section 8 summarises results for further groups such as congruence subgroups and concludes with an overview.

Related work. This article is intended as a comparably short overview that gives pointers to further literature but does not contain many definitions or proofs. A more accessible account of some of the results presented here is given in lecture notes [49] by Patzt–Wilson. These include background material, exercises and open problems.

$2 \operatorname{SL}_n(\mathbb{Z})$

For the integral special linear group, [23, Conjecture 2] says the following:

Conjecture 1 (Church–Farb–Putman).
$$H^{\binom{n}{2}-i}(\mathrm{SL}_n(\mathbb{Z});\mathbb{Q})=0$$
 for $n\geq i+2$.

Here, $\binom{n}{2} = \operatorname{vcd}(\operatorname{SL}_n(\mathbb{Z}))$, so Conjecture 1 claims that the rational cohomology of $\operatorname{SL}_n(\mathbb{Z})$ vanishes in "low codimension i". This vanishing conjecture is equivalent to a high-dimensional stability statement [23, Conjecture 1]: Church–Farb–Putman described a map $\operatorname{St}(\operatorname{SL}_n(\mathbb{Z})) \to \operatorname{St}(\operatorname{SL}_{n+1}(\mathbb{Z}))$ that induces the homomorphism in the top row of the following commutative diagram.

$$H_{i}(\mathrm{SL}_{n}(\mathbb{Z}); \mathrm{St}(\mathrm{SL}_{n}(\mathbb{Z})) \otimes \mathbb{Q}) \longrightarrow H_{i}(\mathrm{SL}_{n+1}(\mathbb{Z}); \mathrm{St}(\mathrm{SL}_{n+1}(\mathbb{Z})) \otimes \mathbb{Q}) \qquad (2)$$

$$\cong \qquad \qquad \cong$$

$$H^{\binom{n}{2}-i}(\mathrm{SL}_{n}(\mathbb{Z}); \mathbb{Q}) \longrightarrow H^{\binom{n+1}{2}-i}(\mathrm{SL}_{n+1}(\mathbb{Z}); \mathbb{Q})$$

The vertical isomorphisms are a case of Borel–Serre Duality as stated in Eq. (1) and define the homomorphism in the bottom row. Using the definitions of all these maps, one can see that Conjecture 1 holds if the horizontal "stabilisation" maps in Eq. (2) are isomorphisms for $n \ge i + 2$.




Figure 1. $H^k(\mathrm{SL}_n(\mathbb{Z});\mathbb{Q})$. The pink area $\binom{n}{2} - (n-2) \le k \le \binom{n}{2}$ is the range of Conjecture 1. Triangle, square and circle mark existing vanishing results (Eq. (3), Eq. (4) and Eq. (5)); brown the known non-trivial classes of highest degree (Eq. (6) for $n \ge 8$).

2.1 Results

Low rank. For small n, the cohomology groups $H^k(\mathrm{SL}_n(\mathbb{Z});\mathbb{Q})$ can be explicitly calculated using computers. At the moment, this has been done for $n \le 7$ and for some small k when $n \le 10$ [27, 41, 55, 57]. These low-rank computations are all consistent with Conjecture 1.

Low codimension. That Conjecture 1 holds for codimension i = 0 was proven by Lee–Szczarba [40] who more generally showed that if R is a Euclidean number ring, then the top-degree cohomology of $SL_n(R)$ is trivial,

$$H^{\text{vcd}(\mathrm{SL}_n(R))}(\mathrm{SL}_n(R);\mathbb{Q}) = 0 \text{ for } R \text{ Euclidean and } n \geq 2.$$
 (3)

Church–Farb–Putman [24] reproved Lee–Szczarba's result² using a "geometric" argument (see Section 2.2). With similar techniques, Church–Putman [25] subsequently showed that Conjecture 1 also holds for codimension i = 1,

$$H^{\binom{n}{2}-1}(\mathrm{SL}_n(\mathbb{Z});\mathbb{Q}) = 0 \text{ for } n \ge 3.$$

²In fact, they proved a stronger statement, see [24, Theorem C].

The currently strongest known bound is due to Brück–Miller–Patzt–Sroka–Wilson [14] who showed the case i = 2 of Conjecture 1, with a slightly stronger bound on n,

$$H^{\binom{n}{2}-2}(\mathrm{SL}_n(\mathbb{Z});\mathbb{Q}) = 0 \text{ for } n \ge 3.$$
 (5)

Limitations for the vanishing. Combining recent results by Ash [2] and Brown [11], one can see that if Conjecture 1 holds, then it is sharp for $n \equiv 0, 1, 2, 4 \mod 6$ and close to being sharp otherwise. Their results show that

$$H^{\binom{n}{2}-(n-1)}(\mathrm{SL}_n(\mathbb{Z});\mathbb{Q}) \neq 0 \text{ for } n \equiv 0, 1, 2, 4 \mod 6,$$

$$H^{\binom{n}{2}-n}(\mathrm{SL}_n(\mathbb{Z});\mathbb{Q}) \neq 0 \text{ for } n \equiv 3, 5 \mod 6.$$
(6)

Low-rank computations show that $H^{(\frac{3}{2})-2}(SL_3(\mathbb{Z});\mathbb{Q})$ and $H^{(\frac{5}{2})-4}(SL_5(\mathbb{Z});\mathbb{Q})$ are trivial, so the classes from Eq. (6) are in the highest possible degree for $n \le 7$ [55, Table 5], see also Fig. 1.

2.2 Techniques: Partial resolutions of the Steinberg module

The vanishing results in Eq. (3), Eq. (4) and Eq. (5) were all obtained by applying Borel–Serre Duality (Eq. (1)) and then showing that $H_i(SL_n(\mathbb{Z}); St(SL_n(\mathbb{Z})) \otimes \mathbb{Q})$ is trivial for appropriate i and n. This was done using specific partial resolutions of the Steinberg module $St(SL_n(\mathbb{Z}))$.

2.2.1 How a resolution helps to show vanishing. Assume that we have a partial flat resolution of $St(SL_n(\mathbb{Z}))$, i.e. an exact sequence

$$C_k \to C_{k-1} \to \cdots \to C_1 \to C_0 \to \operatorname{St}(\operatorname{SL}_n(\mathbb{Z})) \to 0,$$

where each C_i is a flat $\mathrm{SL}_n(\mathbb{Z})$ -module. Then it follows from standard facts in group homology [25, Lemma 3.1] that $H_*(\mathrm{SL}_n(\mathbb{Z});\mathrm{St}(\mathrm{SL}_n(\mathbb{Z}))\otimes \mathbb{Q})$ is given by the homology of the chain complex

$$C_k \otimes_{\operatorname{SL}_n(\mathbb{Z})} \mathbb{Q} \to C_{k-1} \otimes_{\operatorname{SL}_n(\mathbb{Z})} \mathbb{Q} \to \cdots \to C_1 \otimes_{\operatorname{SL}_n(\mathbb{Z})} \mathbb{Q} \to C_0 \otimes_{\operatorname{SL}_n(\mathbb{Z})} \mathbb{Q} \to 0.$$

In particular, if for some i and n, the coinvariants $C_i \otimes_{\mathrm{SL}_n(\mathbb{Z})} \mathbb{Q}$ vanish, then one gets

$$0 = H_i(\mathrm{SL}_n(\mathbb{Z}); \mathrm{St}(\mathrm{SL}_n(\mathbb{Z})) \otimes \mathbb{Q}) \cong H^{\binom{n}{2} - i}(\mathrm{SL}_n(\mathbb{Z}); \mathbb{Q}).$$

This yields a sufficient criterion for showing Conjecture 1, which was used to prove it for codimensions i = 0, 1, 2.




Figure 2. An apartment in $\Delta(SL_3(R))$, corresponding to a basis $\vec{v}_1, \vec{v}_2, \vec{v}_3$ of \mathbb{K}^3 .

2.2.2 What the resolutions look like. In all the partial resolutions considered here, C_i is a free abelian group that is finitely generated as an $SL_n(\mathbb{Z})$ -module. It has an explicit description of the finitely many orbits of generators and each such generator has finite stabiliser (this is related to flatness [25, Lemma 3.2]). A partial resolution of length 0 is just a surjective equivariant map $C_0 \twoheadrightarrow St(SL_n(\mathbb{Z}))$. This is essentially the same as a generating set of $St(SL_n(\mathbb{Z}))$ that is invariant under the action of $SL_n(\mathbb{Z})$ and has only finitely many orbits of generators. Similarly, a partial resolution of length 1 gives a presentation of $St(SL_n(\mathbb{Z}))$. We will describe the partial resolutions that were used for proving Conjecture 1 in codimensions i = 0, 1 in this form.

Buildings and apartments. For later use, we first consider the more general situation where R is a Dedekind domain with fraction field $\mathbb{K} = \operatorname{frac}(R)$ and \mathcal{G} a Chevalley–Demazure group scheme. We then specialise to $\mathcal{G} = \operatorname{SL}_n$, $R = \mathbb{Z}$, $\mathbb{K} = \mathbb{Q}$. The Steinberg module $\operatorname{St}(\mathcal{G}(R))$ is the top-degree homology of the spherical building $\Delta(\mathcal{G}(R))$ associated to $\mathcal{G}(\mathbb{K})$. This building is a highly symmetric simplicial complex equipped with an action of $\mathcal{G}(\mathbb{K})$. It can be described as a union of certain subcomplexes, its apartments. Each apartment Σ is isomorphic to the Coxeter complex of the (finite) Weyl group of \mathcal{G} and hence its geometric realisation is a sphere of dimension $\operatorname{rk}(\mathcal{G}) - 1$. Its fundamental class gives an element $[\Sigma] \in \operatorname{St}(\mathcal{G}(R)) = \widetilde{H}_{\operatorname{rk}(\mathcal{G})-1}(\Delta(\mathcal{G}(R)))$. By the Solomon–Tits Theorem ([56], [1, Theorem 4.127]), these classes generate $\operatorname{St}(\mathcal{G}(R))$.

 \mathbf{SL}_n . The building $\Delta(\mathrm{SL}_n(R))$ is the order complex of the poset of non-zero proper subspaces of \mathbb{K}^n . An apartment is determined by a tuple³ $[v_1, \ldots, v_n]$, where each $v_i = \langle \vec{v}_i \rangle$ is a line such that $\vec{v}_1, \ldots, \vec{v}_n$ is a basis of \mathbb{K}^n . If R is Euclidean, not all apartments are needed to generate $\mathrm{St}(\mathrm{SL}_n(R))$: Ash–Rudolph [3] showed that already the

³The equivalence class of such a tuple is sometimes called a (higher dimensional) "modular symbol" [3].

⁴The apartment of $[v_1, \ldots, v_n]$ is the full subcomplex on subspaces spanned by proper subsets of $\{\vec{v}_1, \ldots, \vec{v}_n\}$, see Fig. 2. This is isomorphic to the barycentric subdivision of the boundary of an (n-1)-simplex, the Coxeter complex of type A_{n-1} .




Figure 3. A 2-simplex in BA₂ whose boundary leads to Relation 2. in St(SL₂(\mathbb{Z})). Its 1-faces correspond to apartment classes $[v_1, v_2]$, $[v_1, \langle \vec{v}_1 + \vec{v}_2 \rangle]$, $[\langle \vec{v}_1 + \vec{v}_2 \rangle, v_2]$ and are contained in B₂.

classes of *integral apartments*, those $[v_1, \ldots, v_n]$ where $\vec{v}_1, \ldots, \vec{v}_n$ can be chosen as a basis of R^n , form a generating set. Taking C_0 to be the free abelian group on all (oriented) integral apartment classes gives a length-0 partial resolution $C_0 \to \operatorname{St}(\operatorname{SL}_n(R))$. Because $\operatorname{SL}_n(R)$ acts transitively on the set of integral apartments, C_0 is cyclic as an $\operatorname{SL}_n(R)$ -module. This description allows one to show that the coinvariants $C_0 \otimes \mathbb{Q}$ are trivial, which implies vanishing of the top-degree cohomology of $\operatorname{SL}_n(R)$ in Eq. (3).

Church–Farb–Putman [24] reproved this integral generation result in a geometric way, that is via showing that a certain simplicial complex B_n is (n-2)-connected. The (n-1)-simplices of B_n are in 1-to-1 correspondence with the integral apartments of $\Delta(\operatorname{SL}_n(R))$ and C_0 is given by the module of simplicial (n-1)-chains. Using similar techniques, Church–Putman [25, Theorem B] also reproved a result by Bykovskiĭ [19] that gives all relations between integral apartment classes. This yields a presentation, i.e. length-1 partial resolution $C_1 \to C_0 \to \operatorname{St}(\operatorname{SL}_n(\mathbb{Z}))$. It describes $\operatorname{St}(\operatorname{SL}_n(\mathbb{Z}))$ as the free abelian group generated by symbols $[v_1, \ldots, v_n]$ modulo the following relations:

1.
$$[v_1, \dots, v_n] = (-1)^{\operatorname{len}(\pi)} \cdot [v_{\pi(1)}, \dots, v_{\pi(n)}] \, \forall \pi \in \operatorname{Sym}_n;$$

2. $[v_1, \dots, v_n] = [v_1, \langle \vec{v}_1 + \vec{v}_2 \rangle, v_3, \dots, v_n] + [\langle \vec{v}_1 + \vec{v}_2 \rangle, v_2, v_3, \dots, v_n].$ (7)

Both relations in (7) show up as the boundaries of simplices in a simplicial complex $BA_n \supseteq B_n$, see Fig. 3. Showing that BA_n is (n-1)-connected is the main technical difficulty in Church–Putman's geometric proof for the above presentation.

To prove the codimension i = 2 case of Conjecture 1 (see Eq. (5)), Brück–Miller–Patzt–Sroka–Wilson extended this presentation to a length-2 partial resolution of $St(SL_n(\mathbb{Z}))$ [14, Section 1.2]. For this, they showed that a certain simplicial complex $BAA_n \supseteq BA_n$ is n-connected. To handle the combinatorial complexity of BAA_n , their argument involves computer calculations.

3 MCG(Σ_g)

The rational cohomology of the surface mapping class group $MCG(\Sigma_g)$ agrees with that of the moduli space \mathcal{M}_g of genus-g Riemann surfaces. Church–Farb–Putman's analogue of Conjecture 1 in this setup [23, Conjecture 9] asked whether

$$H^{4g-5-i}(\mathrm{MCG}(\Sigma_g);\mathbb{Q}) \cong H^{4g-5-i}(\mathcal{M}_g;\mathbb{Q}) = 0 \text{ for } g \gg i?$$
 (8)

This conjecture, which turned out to be false in general, would have also followed from a conjecture by Kontsevich, see [47, Remark 7.5].

Harer [32] and Ivanov [37] showed that the group $MCG(\Sigma_g)$ is a virtual duality group of dimension $vcd(MCG(\Sigma_g)) = 4g - 5$. More precisely, there are isomorphisms

$$H^{4g-5-i}(MCG(\Sigma_g); \mathbb{Q}) \cong H_i(MCG(\Sigma_g); D_g \otimes \mathbb{Q}), \tag{9}$$

where the dualising module D_g is given by the only non-trivial homology group of the curve complex. Similarly to the case of $\mathrm{SL}_n(\mathbb{Z})$ above, this gives access to the high-dimensional cohomology of $\mathrm{MCG}(\Sigma_g)$ if one can find appropriate partial resolutions of the dualising module D_g . Broaddus [10] found such a length-0 partial resolution by giving a generating set for D_g . Church–Farb–Putman [22] used this generating set to show that Eq. (8) is true in codimension i=0, which was also shown independently by Morita–Sakasai–Suzuki [46],

$$H^{4g-5}(\mathrm{MCG}(\Sigma_g); \mathbb{Q}) \cong H^{4g-5}(\mathcal{M}_g; \mathbb{Q}) = 0 \text{ for } g \ge 2.$$
 (10)

However, Chan–Galatius–Payne [20] and later Payne–Willwacher [50] found many non-trivial classes in the high-dimensional cohomology of \mathcal{M}_g , showing that

$$H^{4g-5-i}(\mathrm{MCG}(\Sigma_g);\mathbb{Q}) \neq 0 \text{ for } i \in \{1,3,4,6,7,9,10,11,13,14\} \,, \, g \gg 0.$$

In particular, Eq. (8) is false for all these codimensions i.

Since then, related techniques also yielded high-dimensional cohomology classes for mapping class groups of marked surfaces [21] and of handlebodies [7,31].

4 Automorphisms of free groups

Among the three families of groups considered in [23], the situation is least clear for the group $Aut(F_n)$ of automorphisms of the free group and the related group $Out(F_n)$ of outer automorphisms. These groups have virtual cohomological dimensions $vcd(Aut(F_n)) = 2n - 2$ and $vcd(Out(F_n)) = 2n - 3$. Morita [45, page 390] conjectured a top-degree vanishing phenomenon for $Out(F_n)$,

$$H^{2n-3}(\text{Out}(F_n); \mathbb{Q}) = 0 \text{ for } n \ge 2?$$
 (11)

Church–Farb–Putman did not conjecture vanishing but rather that the high-dimensional cohomology would only depend on the parity of n [23, Conjecture 12],

$$H^{2n-2-i}(\operatorname{Aut}(F_n);\mathbb{Q}), H^{2n-3-i}(\operatorname{Out}(F_n);\mathbb{Q})$$
 determined by $n+2\mathbb{Z}$ for $n\gg i$? (12)

After these conjectures were made, Bartholdi [4] computed the rational cohomology of $Out(F_7)$ and showed that it is non-trivial in the top degree,

$$H^{11}(\operatorname{Out}(F_7); \mathbb{Q}) \cong \mathbb{Q}.$$

This disproved Morita's conjecture in Eq. (11). While Eq. (12) might still be true, Bartholdi's result makes this seem much less likely: It implies that a high-dimensional stabilisation for $\operatorname{Out}(F_n)$ as claimed by Eq. (12) can only occur for $n \geq 7$. In contrast to that, the analogous stabilisation for $\operatorname{SL}_n(\mathbb{Z})$ and $\operatorname{MCG}(\Sigma_g)$ already holds for $n \geq 2$ (Eq. (3)) and $g \geq 2$ (Eq. (10)).

Bestvina–Feighn [5] showed that $Aut(F_n)$ and $Out(F_n)$ are virtual duality groups. This indicates that it might be possible to study their high-dimensional cohomology with similar techniques as for $SL_n(\mathbb{Z})$ and $MCG(\Sigma_g)$ described above. The problem with this is that until today, no sufficiently explicit descriptions of the dualising modules are known. Hatcher–Vogtmann [33] suggested that the top-degree homology of the free factor complex could be the dualising module of $Aut(F_n)$, but Himes–Miller–Nariman–Putman [34] showed that this is not the case, at least for n = 5. Further candidates for geometric descriptions of the dualising modules are studied in [12] and [18].

$5 \operatorname{SL}_n(R)$

In the past years, vanishing patterns as in Conjecture 1 have been studied for several further families of groups "similar to $SL_n(\mathbb{Z})$ ". One such family is $SL_n(R)$, where R is a number ring different from \mathbb{Z} .

5.1 Vanishing

Lee–Szczarba's vanishing result for the top-degree cohomology of $SL_n(R)$ is already formulated in such a setting and applies to all Euclidean number rings R (see Eq. (3)).

Kupers–Miller–Patzt–Wilson [39] proved that this vanishing extends to codimension i = 1 for two specific such Euclidean number rings, namely the Gaussian integers O_{-1} and the Eisenstein integers O_{-3} :

$$H^{(n^2-n)-1}(\mathrm{SL}_n(R);\mathbb{Q}) = 0 \text{ for } R \in \{O_{-1}, O_{-3}\} \text{ and } n \ge 3,$$
 (13)

where $n^2 - n = \text{vcd}(SL_n(R))$. To show this, they proved that for these rings, the Steinberg module $St(SL_n(R))$ has a "generalised Bykovskii" presentation" that looks like the




Figure 4. Number rings under the assumption of GRH.

presentation of $St(SL_n(\mathbb{Z}))$ in Eq. (7). They also showed that this presentation does not hold if R is a quadratic number ring that is Euclidean but not additively generated by units [39, Theorem C].

5.2 Non-vanishing

If one drops the assumption that R be Euclidean, there is no vanishing in general, not even in top degree. Recall that the class group cl(R) is the abelian group given by isomorphism classes of rank-1 projective R-modules with the tensor product. A ring R is a principal ideal domain (PID) if its class group is trivial, |cl(R)| = 1.

 SL_n over non-PIDs. Church–Farb–Putman [24] showed that $\operatorname{SL}_n(R)$ has non-trivial top-degree cohomology if R is a number ring that is not a PID. More precisely, they proved that

$$\dim_{\mathbb{Q}} H^{\operatorname{vcd}(\operatorname{SL}_{n}(R))}(\operatorname{SL}_{n}(R); \mathbb{Q}) \ge (|\operatorname{cl}(R)| - 1)^{(n-1)} \text{ for } n \ge 2.$$
 (14)

Their proof also shows that if R is a Dedekind domain with $2 \le |\operatorname{cl}(R)| < \infty$, then $\operatorname{St}(\operatorname{SL}_n(R))$ is not generated by integral apartment classes for $n \ge 2$ [24, Theorem B].

 SL_n over non-Euclidean PIDs. Every Euclidean ring is a PID. Hence, combining the vanishing result of Lee–Szczarba (Eq. (3)) with Eq. (14), the only number rings for which it is unknown whether $\operatorname{SL}_n(R)$ has vanishing top-degree cohomology are non-Euclidean PIDs. Assuming the Generalised Riemann Hypothesis (GRH), there are only four such number rings [60], see Fig. 4. These are the rings of integers O_d in $\mathbb{Q}(\sqrt{d})$ for $d \in \{-19, -43, -67, -163\}$. Miller–Patzt–Wilson–Yasaki [44] showed that for the last three, the top-degree cohomology of SL_{2n} does not vanish,

$$H^{\text{vcd}(SL_n(O_d))}(SL_n(O_d); \mathbb{Q}) \neq 0 \text{ for } n \text{ even and } d \in \{-43, -67, -163\}.$$
 (15)

They also refined Church–Farb–Putman's non-integrality result from the previous paragraph in the case of number rings: They showed that if R is a number ring and GRH holds, then $St(SL_n(R))$ is generated by integral apartments if and only if R is Euclidean [44, Corollary 1.2].

6 $\operatorname{Sp}_{2n}(R)$

Another way to find groups similar to $SL_n(\mathbb{Z})$ is to replace SL_n by other group schemes. Here, in particular the family of symplectic groups $Sp_{2n}(R)$ has been studied.

6.1 Vanishing

It follows from work of Gunnells that if R is a Euclidean number ring, then the cohomology of $\operatorname{Sp}_{2n}(R)$ vanishes in its virtual cohomological dimension,

$$H^{\text{vcd}(\operatorname{Sp}_{2n}(R))}(\operatorname{Sp}_{2n}(R);\mathbb{Q}) = 0 \text{ for } R \text{ Euclidean and } n \ge 1.$$
 (16)

This analogue of Lee–Szczarba's SL_n -result (Eq. (3)) first appeared in work of Brück–Santos Rego–Sroka [16], but was supposedly known to experts before. For the case $R = \mathbb{Z}$, Brück–Patzt–Sroka [17,58] gave an alternative geometric proof that is independent of Gunnells' work (see Section 6.2). They later [15] used these geometric techniques to show that cohomology vanishing also occurs in codimension 1,

$$H^{n^2-1}(\operatorname{Sp}_{2n}(\mathbb{Z}); \mathbb{Q}) = 0 \text{ for } n \ge 2.$$
(17)

6.2 Partial resolutions of $St(Sp_{2n}(R))$

Similarly to the setting of SL_n (see Section 2.2), the vanishing results in Eq. (16) and Eq. (17) were obtained using Borel–Serre Duality, which here says that

$$H^{\operatorname{vcd}(\operatorname{Sp}_{2n}(R))-i}(\operatorname{Sp}_{2n}(R);\mathbb{Q}) \cong H_i(\operatorname{Sp}_{2n}(R);\operatorname{St}(\operatorname{Sp}_{2n}(R))\otimes\mathbb{Q}),$$

together with explicit partial resolutions of the Steinberg module $St(Sp_{2n}(R))$.

To describe these partial resolutions, let R again be a Dedekind domain and \mathbb{K} its fraction field. Let ω be a symplectic form on \mathbb{K}^{2n} , i.e. a non-degenerate alternating bilinear form $\omega: \mathbb{K}^{2n} \times \mathbb{K}^{2n} \to \mathbb{K}$. The symplectic group $\operatorname{Sp}_{2n}(R)$ is given by all elements of $\operatorname{GL}_{2n}(R) \subseteq \operatorname{GL}_{2n}(\mathbb{K})$ that preserve ω . A subspace $V \subseteq \mathbb{K}^{2n}$ is called isotropic if $\omega|_{V\times V}$ is trivial. A basis $\vec{v}_1, \vec{v}_{\bar{1}}, \ldots, \vec{v}_n, \vec{v}_{\bar{n}}$ of \mathbb{K}^{2n} is called symplectic if

$$\omega(\vec{v}_i,\vec{v}_j) = \omega(\vec{v}_{\bar{i}},\vec{v}_{\bar{j}}) = 0 \quad \text{and} \quad \omega(\vec{v}_i,\vec{v}_{\bar{j}}) = \delta_{ij},$$

where δ_{ij} is the Kronecker delta.

The Steinberg module $\operatorname{St}(\operatorname{Sp}_{2n}(R))$ is the (n-1)-st homology group of the building $\Delta(\operatorname{Sp}_{2n}(R))$. This building is the poset of non-zero isotropic subspaces of \mathbb{K}^{2n} . An apartment in $\Delta(\operatorname{Sp}_{2n}(R))$ is determined by a tuple $[v_1,\ldots,v_{\bar{n}}]$, where each $v_i=\langle\vec{v}_i\rangle$ is a line such that $\vec{v}_1,\ldots,\vec{v}_{\bar{n}}$ is a symplectic basis of \mathbb{K}^{2n} . It is *integral* if $\vec{v}_1,\ldots,\vec{v}_{\bar{n}}$

⁵The apartment of $[v_1, \ldots, v_{\bar{n}}]$ is the full subcomplex on all isotropic subspaces spanned by subsets of $\vec{v}_1, \ldots, \vec{v}_{\bar{n}}$. This is isomorphic to the barycentric subdivision of the boundary of an n-dimensional cross polytope, the Coxeter complex of type B_n .




Figure 5. A subcomplex of IAA₂ whose boundary leads to Relation 3. in $St(Sp_4(\mathbb{Z}))$.

is a basis of R^{2n} . Gunnells [29] showed that $St(Sp_{2n}(R))$ is generated by integral apartments for R Euclidean. Brück–Sroka [17] gave a new, geometric proof for $R = \mathbb{Z}$. With Patzt [15], they extended the techniques to obtain a presentation of $St(Sp_{2n}(\mathbb{Z}))$. They showed [15, Theorem B] that it is the free abelian group generated by symbols $[v_1, \ldots, v_{\bar{n}}]$ modulo the following relations:

1.
$$[v_{1}, \ldots, v_{\bar{n}}] = (-1)^{\operatorname{len}(\pi)} \cdot [v_{\pi(1)}, \ldots, v_{\pi(\bar{n})}] \, \forall \pi \in \operatorname{Sym}_{n}^{\pm};$$

2. $[v_{1}, \ldots, v_{\bar{n}}] = [v_{1}, \langle \vec{v}_{1} + \vec{v}_{\bar{1}} \rangle, v_{2}, \ldots, v_{\bar{n}}] + [\langle \vec{v}_{1} + \vec{v}_{\bar{1}} \rangle, v_{\bar{1}}, v_{2}, \ldots, v_{\bar{n}}];$
3. $[v_{1}, \ldots, v_{\bar{n}}] = [v_{1}, \langle \vec{v}_{\bar{1}} - \vec{v}_{\bar{2}} \rangle, \langle \vec{v}_{1} + \vec{v}_{2} \rangle, v_{\bar{2}}, v_{3}, \ldots, v_{\bar{n}}]$
 $+ [\langle \vec{v}_{\bar{1}} - \vec{v}_{\bar{2}} \rangle, v_{2}, \langle \vec{v}_{1} + \vec{v}_{2} \rangle, v_{\bar{1}}, v_{3}, \ldots, v_{\bar{n}}].$ (18)

Here, $\operatorname{Sym}_n^{\pm}$ is the signed permutation group, the Coxeter group of type $B_n = C_n$. Similar to the case of $\operatorname{SL}_n(\mathbb{Z})$, these three relations occur as boundaries in an *n*-connected complex IAA_n, see Fig. 5.

6.3 Non-vanishing

Different sources of non-trivial cohomology classes provide limitations for a high-dimensional vanishing of $H^*(\operatorname{Sp}_{2n}(R); \mathbb{Q})$.

 $\operatorname{Sp}_{2n}(\mathbb{Z})$ and the moduli space \mathcal{A}_n . The rational cohomology of $\operatorname{Sp}_{2n}(\mathbb{Z})$ equals that of \mathcal{A}_n , the moduli space of principally polarised abelian varieties of dimension 2n. Brandt–Bruce–Chan–Melo–Moreland–Wolfe [8] computed the top-weight rational cohomology of this moduli space for $n \leq 7$. The classes of highest degree that they found are in codimension n, so their results imply that

$$H^{n^2-n}(\operatorname{Sp}_{2n}(\mathbb{Z});\mathbb{Q}) \cong H^{n^2-n}(\mathcal{A}_n;\mathbb{Q}) \neq 0 \text{ for } n \leq 7.$$
(19)

 \mathbf{Sp}_{2n} over non-PIDs. Brück-Himes [13] gave an analogue of Church-Farb-Putman's non-vanishing result in Eq. (14). They proved that for number rings R, the top-degree

cohomology of $Sp_{2n}(R)$ is non-trivial if R is not a PID,

$$\dim_{\mathbb{Q}} H^{\text{vcd}(\text{Sp}_{2n}(R))}(\text{Sp}_{2n}(R); \mathbb{Q}) \ge (|\operatorname{cl}(R)| - 1)^n \text{ for } n \ge 1.$$
 (20)

They also showed that if R is a Dedekind domain with $2 \le |\operatorname{cl}(R)| < \infty$, then $\operatorname{St}(\operatorname{Sp}_{2n}(R))$ is not generated by integral apartment classes for $n \ge 1$ [13, Theorem 1.2].

7 Chevalley groups

For the top degree, i.e. the cohomology in the virtual cohomological dimension, there is a vanishing result that covers a much bigger class of groups than just SL_n and Sp_{2n} .

Let R be a number ring and G a Chevalley–Demazure group scheme. As mentioned in Section 2.2.2, the Steinberg module St(G(R)) is generated by the fundamental classes of apartments in the building $\Delta(G(R))$. Call such a class *integral* if it is a G(R)-translate of the class of the standard apartment⁶. This notion generalises integrality of apartment classes in the special linear and symplectic settings described above.

Tóth [59] showed that if R is Euclidean, then for almost all Chevalley types, the Steinberg module $St(\mathcal{G}(R))$ is generated by integral apartment classes. Brück–Santos Rego–Sroka [16] deduced from this that the top-degree cohomology of $\mathcal{G}(R)$ is trivial,

$$H^{\text{vcd}(\mathcal{G}(R))}(\mathcal{G}(R);\mathbb{Q}) = 0 \text{ for } R \text{ Euclidean, } \mathcal{G} \text{ type } A_n, B_n, C_n, D_n, E_6, E_7.$$
 (21)

They then stated the following conjecture as [16, Question 1.2].

Conjecture 2. Let R be a Euclidean number ring and G a Chevalley-Demazure group scheme. Then

$$H^{\operatorname{vcd}(\mathcal{G}(R))-i}(\mathcal{G}(R);\mathbb{Q}) = 0 \text{ for } i < \operatorname{rk}(\mathcal{G}).$$

For $G = SL_n$ and $R = \mathbb{Z}$, this specialises to Conjecture 1 by Church–Farb–Putman.

Known cases. For i = 0, Conjecture 2 is known to hold for all irreducible \mathcal{G} except those of type F_4 , G_2 or E_8 (Eq. (21)). For i = 1, it it is presently known to be true for four families: for $SL_n(\mathbb{Z})$ (Eq. (4)); for $SL_n(O_{-1})$ and $SL_n(O_{-3})$ (Eq. (13)); and for $Sp_{2n}(\mathbb{Z})$ (Eq. (17)). For i = 2, the only known instance is $SL_n(\mathbb{Z})$ (Eq. (5)). Conjecture 2 is consistent with low-rank cohomology computations for $SL_n(\mathbb{Z})$, $n \le 10$ [27,41,55,57]; for $Sp_{2n}(\mathbb{Z})$, $n \le 4$ [30,35,36]; and for $SL_n(R)$, $n \le 3$, where R is a Euclidean ring of integers in any imaginary quadratic field [26,54].

⁶This designated apartment in $\Delta(\mathcal{G}(R))$ is uniquely determined by the choice of a BN-pair in $\mathcal{G}(\mathbb{K})$, see [1, Chapter 6].

Limitations. Eq. (14) and Eq. (20) show that the vanishing claimed by Conjecture 2 can only occur for rings R that are PIDs. As mentioned above, R being Euclidean only excludes four further rings under the assumption of the Generalised Riemann Hypothesis. This stronger condition seems natural given the non-vanishing results of Miller–Patzt–Wilson–Yasaki (Eq. (15)).

Known cohomology classes show that one also cannot hope for vanishing results in higher codimensions: The highest degree classes known for $SL_n(\mathbb{Z})$ (Eq. (6)) and for $Sp_{2n}(\mathbb{Z})$ (Eq. (19)) are exactly one below the conjectured vanishing range, in codimensions $(n-1) = \text{rk}(SL_n)$ and $n = \text{rk}(Sp_{2n})$, respectively.

8 Further results and overview

Congruence subgroups. The high-dimensional vanishing results presented above depend on the existence of torsion elements in the groups under questions. These are used to show vanishing of the coinvariants of partial resolutions as described in Section 2.2 (see e.g. [16, Proof of Theorem 3.4] or [25, Section 3.2, p. 1012]). In particular, vanishing is not preserved under passing to finite index subgroups. For example, the top-degree cohomology of congruence subgroups in $SL_n(\mathbb{Z})$ is non-trivial and its rank grows quickly with n [43,48,52,53]. The situation is similar for congruence subgroups in $MCG(\Sigma_g)$ [9,28]. Church–Farb–Putman conjectured that there might be some kind of high-dimensional stability for congruence subgroups in $SL_n(\mathbb{Z})$ nonetheless. In [42], Miller–Nagpal–Patzt stated a precise version of this conjecture [42, Conjecture 6.1] using representation stability and proved it in some instances.

 GL_n and configuration spaces. For vanishing and non-vanishing results in the top-degree cohomology of $GL_n(R)$, see [51]. High-dimensional stability patterns in the context of configuration spaces are studied in [38,61].

Overview. Of the original three conjectures in [23], only the one for $SL_n(\mathbb{Z})$ still has a good chance of being true. However, results for further arithmetic groups indicate that there might be a vanishing phenomenon similar to that of Conjecture 1 that occurs in a larger class of groups. The precise form of this still has to be determined, one candidate is given by Conjecture 2. The high-dimensional (non-)vanishing results mentioned in this article are summarised in Table 1.

Acknowledgements. I would like to thank Jeremy Miller and Jennifer C. H. Wilson for their comments, which helped to improve the content and presentation of this article.

Funding. This work was partially supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 427320536 – SFB 1442, as well

Group Γ	i = 0	i = 1	i = 2
$\mathrm{SL}_n(\mathbb{Z})$	Yes [40]	Yes [25]	Yes [14]
$MCG(\Sigma_g)$	Yes [22]	No [20]	
$\operatorname{Aut}(F_n)$	Probably no [4] (low rank calculations)		
$\overline{\mathrm{SL}_n(R)}$	Yes [40]	Yes for	
R Eucl.		$R \in \{O_{-1}, O_{-3}\}$ [39]	
$\mathrm{SL}_n(R)$	No if $R \neq O_{-19}$ [24,44]		
R not Eucl.	(assuming GRH)		
$\operatorname{Sp}_{2n}(\mathbb{Z})$	Yes [17]	Yes [15]	
$\operatorname{Sp}_{2n}(R)$ $R \text{ not PID}$	No [13]		
$G_n(R)$	Yes for type A_n , B_n ,		
R Eucl.	$C_n, D_n, E_6, E_7 [16]$		
$MCG(V_g)$		No [7]	
congruence	No		
subgroups	[9,28,43,48,52,53]		

Table 1. A table showing whether stably, $H^{\text{vcd}(\Gamma)-i}(\Gamma;\mathbb{Q})=0$ for different groups Γ . R denotes a number ring, Σ_g and V_g a genus-g surface and handlebody, F_n the free group of rank n.

as by Germany's Excellence Strategy EXC 2044 – 390685587, Mathematics Münster: Dynamics–Geometry–Structure.

References

- P. Abramenko and K. S. Brown, *Buildings*. Graduate Texts in Mathematics 248, Springer, New York, 2008
- [2] A. Ash, On the cohomology of $SL_n(\mathbb{Z})$. 2024, arXiv:2402.08840
- [3] A. Ash and L. Rudolph, The modular symbol and continued fractions in higher dimensions. *Inventiones Mathematicae* **55** (1979), no. 3, 241–250
- [4] L. Bartholdi, The rational homology of the outer automorphism group of F_7 . New York Journal of Mathematics **22** (2016), 191–197
- [5] M. Bestvina and M. Feighn, The topology at infinity of $Out(F_n)$. Inventiones Mathematicae **140** (2000), no. 3, 651–692
- [6] A. Borel and J.-P. Serre, Corners and arithmetic groups. Commentarii Mathematici Helvetici 48 (1973), 436–491

- [7] M. Borinsky, B. Brück, and T. Willwacher, Weight 2 cohomology of graph complexes of cyclic operads and the handlebody group. 2023, arXiv:2308.16845
- [8] M. Brandt, J. Bruce, M. Chan, M. Melo, G. Moreland, and C. Wolfe, On the top-weight rational cohomology of \mathcal{A}_g . Geometry & Topology 28 (2024), no. 2, 497–538
- [9] T. Brendle, N. Broaddus, and A. Putman, The high-dimensional cohomology of the moduli space of curves with level structures II: punctures and boundary. *Israel Journal of Mathematics* (2023)
- [10] N. Broaddus, Homology of the curve complex and the Steinberg module of the mapping class group. *Duke Mathematical Journal* **161** (2012), no. 10, 1943–1969
- [11] F. Brown, Bordifications of the moduli spaces of tropical curves and abelian varieties, and unstable cohomology of $GL_{\sigma}(\mathbb{Z})$ and $SL_{\sigma}(\mathbb{Z})$. 2023, arXiv:2309.12753
- [12] B. Brück and R. Gupta, Homotopy type of the complex of free factors of a free group. *Proceedings of the London Mathematical Society. Third Series* **121** (2020), no. 6, 1737–1765
- [13] B. Brück and Z. Himes, Top-degree rational cohomology in the symplectic group of a number ring. 2023, arXiv:2309.05456
- [14] B. Brück, J. Miller, P. Patzt, R. Sroka, and J. C. H. Wilson, On the codimension-two cohomology of $SL_n(\mathbb{Z})$. 2022, arXiv:2204.11967v1
- [15] B. Brück, P. Patzt, and R. J. Sroka, A presentation of symplectic steinberg modules and cohomology of $\operatorname{Sp}_{2n}(\mathbb{Z})$. 2023, arXiv:2306.03180
- [16] B. Brück, Y. Santos Rego, and R. J. Sroka, On the top-dimensional cohomology of arithmetic Chevalley groups. 2022, arXiv:2210.12784v1
- [17] B. Brück and R. J. Sroka, Apartment classes of integral symplectic groups. 2023, arXiv: 2305.17697
- [18] K.-U. Bux, P. Smillie, and K. Vogtmann, On the bordification of outer space. *Journal of the London Mathematical Society. Second Series* **98** (2018), no. 1, 12–34
- [19] V. A. Bykovskiĭ, Generating elements of the annihilating ideal for modular symbols. *Functional Analysis and Its Applications* **37** (2003), no. 4, 263–272
- [20] M. Chan, S. Galatius, and S. Payne, Tropical curves, graph complexes, and top weight cohomology of \mathcal{M}_g . Journal of the American Mathematical Society **34** (2021), no. 2, 565–594
- [21] M. Chan, S. Galatius, and S. Payne, Topology of moduli spaces of tropical curves with marked points. In *Facets of algebraic geometry. a collection in honor of william fulton's* 80th birthday. volume 1, pp. 77–131, Cambridge: Cambridge University Press, 2022
- [22] T. Church, B. Farb, and A. Putman, The rational cohomology of the mapping class group vanishes in its virtual cohomological dimension. *International Mathematics Research Notices. IMRN* (2012), no. 21, 5025–5030
- [23] T. Church, B. Farb, and A. Putman, A stability conjecture for the unstable cohomology of $SL_n\mathbb{Z}$, mapping class groups, and $Aut(F_n)$. In *Algebraic topology: applications and new directions*, pp. 55–70, Contemp. Math. 620, Amer. Math. Soc., Providence, RI, 2014

- [24] T. Church, B. Farb, and A. Putman, Integrality in the Steinberg module and the topdimensional cohomology of SL_nO_K. American Journal of Mathematics **141** (2019), no. 5, 1375–1419
- [25] T. Church and A. Putman, The codimension-one cohomology of $SL_n\mathbb{Z}$. Geometry & Topology 21 (2017), no. 2, 999–1032
- [26] M. Dutour Sikirić, H. Gangl, P. E. Gunnells, J. Hanke, A. Schürmann, and D. Yasaki, On the cohomology of linear groups over imaginary quadratic fields. *Journal of Pure and Applied Algebra* **220** (2016), no. 7, 2564–2589
- [27] P. Elbaz-Vincent, H. Gangl, and C. Soulé, Perfect forms, K-theory and the cohomology of modular groups. Advances in Mathematics 245 (2013), 587–624
- [28] N. J. Fullarton and A. Putman, The high-dimensional cohomology of the moduli space of curves with level structures. *Journal of the European Mathematical Society (JEMS)* **22** (2020), no. 4, 1261–1287
- [29] P. E. Gunnells, Symplectic modular symbols. *Duke Mathematical Journal* **102** (2000), no. 2, 329–350
- [30] R. Hain, The rational cohomology ring of the moduli space of abelian 3-folds. *Mathematical Research Letters* **9** (2002), no. 4, 473–491
- [31] L. Hainaut and D. Petersen, Top weight cohomology of moduli spaces of riemann surfaces and handlebodies, 2023, arXiv:2308.16845
- [32] J. L. Harer, The virtual cohomological dimension of the mapping class group of an orientable surface. *Inventiones Mathematicae* 84 (1986), no. 1, 157–176
- [33] A. Hatcher and K. Vogtmann, The complex of free factors of a free group. *The Quarterly Journal of Mathematics. Oxford. Second Series* **49** (1998), no. 196, 459–468
- [34] Z. Himes, J. Miller, S. Nariman, and A. Putman, The free factor complex and the dualizing module for the automorphism group of a free group. *IMRN. International Mathematics Research Notices* **2023** (2023), no. 22, 19020–19068
- [35] K. Hulek and O. Tommasi, Cohomology of the second Voronoi compactification of A₄. Documenta Mathematica 17 (2012), 195–244
- [36] J.-i. Igusa, On Siegel modular forms of genus two. *American Journal of Mathematics* **84** (1962), 175–200
- [37] N. V. Ivanov, Complexes of curves and Teichmüller modular groups. Akademiya Nauk SSSR i Moskovskoe Matematicheskoe Obshchestvo. Uspekhi Matematicheskikh Nauk 42 (1987), no. 3(255), 49–91, 255
- [38] B. Knudsen, J. Miller, and P. Tosteson, Extremal stability for configuration spaces. Mathematische Annalen 386 (2023), no. 3-4, 1695–1716
- [39] A. Kupers, J. Miller, P. Patzt, and J. C. H. Wilson, On the generalized Bykovskiĭ presentation of Steinberg modules. *International Mathematics Research Notices*. *IMRN* (2022), no. 13, 10347–10401
- [40] R. Lee and R. H. Szczarba, On the homology and cohomology of congruence subgroups. *Inventiones Mathematicae* **33** (1976), no. 1, 15–53

- [41] R. Lee and R. H. Szczarba, On the torsion in $K_4(\mathbb{Z})$ and $K_5(\mathbb{Z})$. Duke Mathematical Journal **45** (1978), 101–129
- [42] J. Miller, R. Nagpal, and P. Patzt, Stability in the high-dimensional cohomology of congruence subgroups. *Compositio Mathematica* **156** (2020), no. 4, 822–861
- [43] J. Miller, P. Patzt, and A. Putman, On the top dimensional cohomology groups of congruence subgroups of $SL_n(\mathbb{Z})$. Geometry & Topology 25 (2021), no. 2, 999–1058
- [44] J. Miller, P. Patzt, J. C. H. Wilson, and D. Yasaki, Non-integrality of some Steinberg modules. *Journal of Topology* 13 (2020), no. 2, 441–459
- [45] S. Morita, Structure of the mapping class groups of surfaces: a survey and a prospect. In *Proceedings of the Kirbyfest (Berkeley, CA, 1998)*, pp. 349–406, Geom. Topol. Monogr. 2, Geom. Topol. Publ., Coventry, 1999
- [46] S. Morita, T. Sakasai, and M. Suzuki, Abelianizations of derivation Lie algebras of the free associative algebra and the free Lie algebra. *Duke Mathematical Journal* 162 (2013), no. 5, 965–1002
- [47] S. Morita, T. Sakasai, and M. Suzuki, Computations in formal symplectic geometry and characteristic classes of moduli spaces, *Quantum Topology* **6** (2015), no. 1, 139–182
- [48] A. Paraschivescu, On a generalization of the double coset formula. *Duke Mathematical Journal* **89** (1997), no. 1, 1–8
- [49] P. Patzt and J. C. H. Wilson, The high-degree rational cohomology of $SL_n(\mathbb{Z})$ and its principal congruent subgroups. 2022, URL https://dept.math.lsa.umich.edu/~jchw/COGENT-School.pdf, lecture notes of a COGENT Summer School 2022
- [50] S. Payne and T. Willwacher, Weight two compactly supported cohomology of moduli spaces of curves. 2021, arXiv:2110.05711v1
- [51] A. Putman and D. Studenmund, The dualizing module and top-dimensional cohomology group of $GL_n(O)$. *Mathematische Zeitschrift* **300** (2022), no. 1, 1–31
- [52] M. Scalamandre, A Solomon-Tits theorem for rings. 2023, arXiv:2310.07175
- [53] J. Schwermer, Eisenstein series and the top degree cohomology of arithmetic subgroups of SL_n/\mathbb{Q} . Journal für die Reine und Angewandte Mathematik 777 (2021), 127–155
- [54] J. Schwermer and K. Vogtmann, The integral homology of SL_2 and PSL_2 of Euclidean imaginary quadratic integers. *Commentarii Mathematici Helvetici* **58** (1983), 573–598
- [55] M. D. Sikirić, P. Elbaz-Vincent, A. Kupers, and J. Martinet, Voronoi complexes in higher dimensions, cohomology of $GL_N(Z)$ for $N \ge 8$ and the triviality of $K_8(Z)$. 2019, arxiv: 1910.11598v1
- [56] L. Solomon, The Steinberg character of a finite group with BN-pair. In Theory of Finite Groups (Symposium, Harvard Univ., Cambridge, Mass., 1968), edited by R. Brauer and C. han Sah, pp. 213–221, Benjamin, New York, 1969
- [57] C. Soulé, The cohomology of SL₃(**Z**). *Topology. An International Journal of Mathematics* **17** (1978), no. 1, 1–22
- [58] R. J. Sroka, *Patterns in the homology of algebras: Vanishing, stability, and higher structures*. Ph.D. thesis, University of Copenhagen, 2021, URL https://noter.math.ku.dk/phd21rs.pdf, ISBN 978-87-7125-043-5

- [59] Á. Tóth, On the Steinberg module of Chevalley groups. *Manuscripta Mathematica* **116** (2005), no. 3, 277–295
- [60] P. J. Weinberger, On Euclidean rings of algebraic integers. Analytic Number Theory, Proc. Sympos. Pure Math. 24, St. Louis Univ. Missouri 1972, 321-332., 1973
- [61] M. Yameen, A remark on extremal stability of configuration spaces. *Journal of Pure and Applied Algebra* **227** (2023), no. 1, 5