# Positive Moments Forever: Undecidable and Decidable Cases

Gemma De les Coves<sup>1</sup>, Joshua Graf<sup>1</sup>, Andreas Klingler<sup>\*1</sup>, and Tim Netzer<sup>2</sup>

<sup>1</sup>Institute for Theoretical Physics, University of Innsbruck, Austria <sup>2</sup>Department of Mathematics, University of Innsbruck, Austria

April 24, 2024

#### Abstract

Is there an algorithm to determine attributes such as positivity or non-zeroness of linear recurrence sequences? This long-standing question is known as Skolem's problem. In this paper, we study the complexity of an equivalent problem, namely the (generalized) moment membership problem for matrices. We show that this problem is decidable for orthogonal, unitary and real eigenvalue matrices, and undecidable for matrices over certain commutative and non-commutative polynomial rings. Our results imply that the positivity problem for simple unitary linear recurrence sequences is decidable, and is undecidable for linear recurrence sequences over the ring of commutative polynomials. As a byproduct, we prove a free version of Pólya's theorem.

# Contents

1	Introduction	<b>2</b>				
2	Problem statement					
	2.1 Relation to the membership problem for linear recurrence sequences $\ldots$ .	5				
3	Decidable cases					
	3.1 Known results: small order	6				
	3.2 Orthogonal and unitary matrices	6				
	3.3 Matrices with a unique dominant eigenvalue or real eigenvalues	9				
	3.4 Further generalizations	10				
4	Undecidable cases	12				
	4.1 Commutative polynomial rings	12				
	4.2 Non-commutative polynomial rings	13				
5	Conclusion	15				
A	A version of the matrix mortality problem	16				
	*Corresponding author. E-mail: andreas.klingler@uibk.ac.at					

## 1 Introduction

Some problems may look innocent yet be formally very difficult — perhaps uncomputable — or even worse, their computability may be unknown. *Skolem's problem* exemplifies this uncertainty, focusing on the behavior of *linear recurrence sequences* (LRS), where each term in the sequence is generated linearly from its predecessors. Examples of LRS include well-known sequences like the Fibonacci sequence or those derived from discretizing differential equations. Despite their simplicity, LRS are fundamental in various mathematical and computer science domains, notably in generating pseudo-random numbers [18], describing the dynamics of cellular automata [11], and many other applications [6].

Specifically, an LRS of order s is given by

$$u_n = a_1 u_{n-1} + a_2 u_{n-2} + \dots + a_s u_{n-s}$$

where  $a_1, \ldots, a_s \in \mathcal{R}$  are fixed elements in a ring  $\mathcal{R}$ , usually commutative. Together with initial values  $u_1, \ldots, u_s \in \mathcal{R}$ , this gives rise to a sequence  $(u_n)_{n \in \mathbb{N}}$  in  $\mathcal{R}$ . While several important examples of LRS are over the ring  $\mathcal{R} = \mathbb{Z}$ , many interesting examples are defined over other rings. For example, the Chebyshev polynomials are defined via the LRS

$$T_n(x) = 2xT_{n-1}(x) - T_{n-2}(x)$$
 with  $T_1(x) \coloneqq x$  and  $T_0(x) \coloneqq 1$ 

over the commutative ring  $\mathbb{Z}[x]$  of univariate polynomials.

Skolem's problem is a long-standing open question concerning LRS over  $\mathbb{Z}$  [13]. It asks whether an algorithm exists that decides if an LRS attains the value 0 for some  $n \in \mathbb{N}$ . While partial solutions to Skolem's problem are known, implying decidability for order  $s \leq 4$  [19, 20], they do not apply to recurrences of order five or more. A modification of Skolem's problem is the *positivity problem* for LRS. Instead of asking whether the LRS is non-zero, it asks whether it stays non-negative. In this case it is also unclear whether an algorithm exists that decides the positivity problem, as decidability is proven only for  $s \leq 5$  [14, 15].

Examples for LRS are *moment sequences*, in which we have

$$u_n = \operatorname{tr}(A^n),$$

or generalized moment sequences, in which

$$u_n = \varphi(A^n)$$

for a given matrix  $A \in \operatorname{Mat}_{s}(\mathcal{R})$  and a linear functional  $\varphi$  on  $\operatorname{Mat}_{s}(\mathcal{R})$ . Over a commutative ring  $\mathcal{R}$ , such generalized moment sequences are as expressive as LRS, i.e. every LRS can be expressed as a moment sequence and vice versa. For this reason, decidability results for generalized moment sequences translate to decidability results for Skolem's problem and the positivity problem.

In this paper, we study the decidability of the moment membership problem. That is, we consider the problem: For an  $s \times s$  matrix A, decide whether

$$\operatorname{tr}(A^n) \in \mathcal{P} \quad \forall n \in \mathbb{N}$$

where  $\mathcal{P}$  is a fixed set. This set usually contains elements that are positive in some sense, so we call the problem also the *moment positivity problem*. Most of our results also hold for generalized moments of the form  $\varphi(A^n)$  as above.

One decisive factor in the complexity of the problem is the instance set  $\mathcal{D}$  of the matrices, which allows us to distinguish between our two main results:

Decidable cases	Undecidable cases
Unitary and Orthogonal matrices	Comm. polynomials $\mathbb{Z}[x_1, \ldots, x_d]$
(Section 3.2)	(Section 4.1)
Dominant or real eigenvalue matrices	Non-comm. polynomials $\mathbb{Z}\langle z_1, \ldots, z_d \rangle$
(Section $3.3$ )	(Section $4.2$ )

**Tab. 1.** For which instance sets is the (generalized) moment membership problem decidable or undecidable? This table summarizes the results of this paper.

- $\triangleright$  We restrict the instance set  $\mathcal{D} \subseteq \operatorname{Mat}_{s}(\mathbb{Z})$  and prove decidability of the problem for a large subclass of integer matrices.
- $\triangleright$  We enlarge the instance set  $\operatorname{Mat}_{s}(\mathbb{Z}) \subseteq \mathcal{D}$  and prove that the problem is undecidable for matrices whose entries are elements of certain unital rings  $\mathcal{R}$ , for certain  $\mathcal{P} \subseteq \mathcal{R}$ .

**Contributions.** Specifically, we determine the complexity of the moment membership problem in the following cases (see Table 1):

- ▷ Decidability. The moment positivity problem is decidable for orthogonal matrices (Theorem 8), unitary matrices (Corollary 11), and matrices with a unique dominant eigenvalue or only real eigenvalues (Theorem 13). It follows that the positivity problem is decidable for simple unitary LRS, i.e. LRS whose characteristic polynomial only has simple roots of modulus 1, as well as for LRS whose characteristic polynomial has a unique dominant root, or only real roots.
- $\triangleright$  Undecidability. The generalized problem is undecidable for the ring of multivariate commutative polynomials (Theorem 18) as well as for non-commutative polynomials, where  $\mathcal{P}$  is the set of polynomials with nonnegative coefficients (Theorem 22). This implies that the corresponding positivity problem for LRS over commutative polynomials is undecidable.
- ▷ Free Pólya's Theorem. As a side result, we prove a free version of Pólya's theorem (Theorem 20). We show that a non-commutative polynomial has nonnegative coefficients if and only if it is entrywise nonnegative on the set of entrywise nonnegative matrices.

This paper is structured as follows. In Section 2 we present the problem statement and show the relation of moment problems to LRS. In Section 3 we present cases in which the moment problem is decidable. This includes a review of known results (Section 3.1), the decidability for orthogonal and unitary matrices (Section 3.2), and the decidability for matrices with unique largest eigenvalue or only real roots (Section 3.3). In Section 4, we prove that the moment problem is undecidable over certain commutative and noncommutative rings, as well as a non-commutative version of Pólya's Theorem. In Section 5 we conclude, and in Appendix A we prove a version of the matrix mortality problem for the undecidability result of commutative polynomials.

### 2 Problem statement

Let  $\mathcal{R}$  be a unital ring (not necessarily commutative), and let  $A \in \operatorname{Mat}_{s}(\mathcal{R})$  be an  $s \times s$  square matrix with entries from  $\mathcal{R}$ . For  $n \ge 0$  the *n*-th moment of A is defined as

$$\mu_n(A) \coloneqq \operatorname{tr}(A^n)$$

where tr denotes the usual trace of a matrix, i.e. the sum of its diagonal entries. The moments of A are clearly elements from  $\mathcal{R}$ , as for  $A = (a_{ij})_{i,j=1,\dots,s}$  we have

$$\mu_0(A) = \underbrace{1_{\mathcal{R}} + \dots + 1_{\mathcal{R}}}_{s}$$
 and  $\mu_1(A) = \sum_{i=1}^s a_{ii}$ 

and for  $n \ge 2$ 

$$\mu_n(A) = \sum_{i_1,\dots,i_n=1}^s a_{i_1i_2} \cdot a_{i_2i_3} \cdots a_{i_{n-1}i_n} \cdot a_{i_ni_1}.$$

Depending on the ring  $\mathcal{R}$ , the moments are studied in different contexts, as the following example shows.

**Example 1.** Let V be a  $\mathbb{C}$ -vector space. Consider the tensor algebra

$$\mathcal{R} \coloneqq T(V) \coloneqq \bigoplus_{m \ge 0} V^{\otimes m} = \mathbb{C} \oplus V \oplus (V \otimes V) \oplus \cdots$$

 $\mathcal{R}$  forms a unital ring with tensor product as multiplication. Actually,  $\mathcal{R}$  is an N-graded unital C-algebra.

For  $A = (a_{ij})_{i,j} \in \operatorname{Mat}_s(V)$  (which embeds into  $\operatorname{Mat}_s(\mathcal{R})$ ), we obtain

$$\mu_n(A) = \sum_{i_1,\dots,i_n=1}^s a_{i_1i_2} \otimes a_{i_2i_3} \otimes \dots \otimes a_{i_{n-1}i_n} \otimes a_{i_ni_1}, \tag{1}$$

where the *n*-th moment is homogeneous of degree *n* in  $\mathcal{R}$ . The expression in (1) is called a *translational invariant matrix product state* in tensor network theory. The number *s* is usually called the *bond dimension* of the tensor  $\mu_n(A)$ .

Now assume that  $\mathcal{R}$  is also equipped with a subset  $\mathcal{P} \subseteq \mathcal{R}$ . In our results and applications, this will always be a set of elements that are *positive* in some sense. Further  $\mathcal{D} \subseteq \operatorname{Mat}_{s}(\mathcal{R})$  will be the set containing all instances of our decision problem. The general decision problem addressed in this paper is the following:

**Problem 2** (Moment positivity problem). Let  $s, \mathcal{P}, \mathcal{D}$  be fixed as above. For  $A \in \mathcal{D}$  decide whether all moments  $\mu_n(A)$  belong to  $\mathcal{P}$ .

Note that  $\mathcal{D}, \mathcal{P}, s$  are fixed in our formulation of the problem. We are thus looking for an algorithm (tailored to  $\mathcal{R}, \mathcal{P}, s$  and  $\mathcal{D}$ ) that upon an input of any instance  $A \in \mathcal{D}$  stops after a finite time, and returns yes if all moments of A belong to  $\mathcal{P}$ , and no if at least one moment of A does not belong to  $\mathcal{P}$ . If such an algorithm exist, we call the moment membership problem *decidable*, otherwise we call it *undecidable*.

Note that if the ring operations are computable and membership of single elements in  $\mathcal{P}$  is decidable, the moments membership problem is clearly semi-decidable in the following sense. Given  $A \in \operatorname{Mat}_{s}(\mathcal{R})$ , we compute higher and higher moments of A, and check membership in  $\mathcal{P}$ . If some moment does *not* belong to  $\mathcal{P}$ , we will know after a finite time. However, this algorithm runs forever in case that all moments do belong to  $\mathcal{P}$ . So the hard part of the problem is certifying membership of all moments in  $\mathcal{P}$ . We will make use of the semi-decidability in Theorem 8.

#### 2.1 Relation to the membership problem for linear recurrence sequences

In the following, we review the relation of the moment problem to the positivity problem for linear recurrence sequences. A linear recurrence sequence (LRS)  $(u_n)_{n \in \mathbb{N}} \in \mathcal{R}^{\mathbb{N}}$  is a sequence whose elements are related to each other linearly, i.e.

$$u_n = a_1 u_{n-1} + a_2 u_{n-2} + \dots + a_s u_{n-s} \tag{2}$$

for all n > s. We call s the *order* of the recurrence relation. The positivity problem for LRS is the following:

**Problem 3.** Given an LRS as in (2) with parameters  $a_1, \ldots, a_s \in \mathcal{R}$  and initial values  $u_1, \ldots, u_s \in \mathcal{R}$ , decide whether  $u_n \in \mathcal{P}$  for all  $n \in \mathbb{N}$ .

We start with the (well-known) observation that every generalized moment sequence is an LRS, if  $\mathcal{R}$  is commutative.

**Lemma 4.** Let  $\mathcal{R}$  be a commutative unital ring, and let  $A \in \operatorname{Mat}_{s}(\mathcal{R})$ . Then  $(\varphi(A^{n}))_{n \in \mathbb{N}}$  is an LRS of order s, for every  $\mathcal{R}$ -linear map  $\varphi \colon \operatorname{Mat}_{s}(\mathcal{R}) \to \mathcal{R}$ .

*Proof.* Let  $p(x) = x^s - a_1 x^{s-1} - \cdots - a_s$  be the characteristic polynomial of the matrix A. By the Cayley–Hamilton theorem for commutative rings (see for example [10, Chapter XIV.3]), we have that

$$A^{s} = a_{1}A^{s-1} + a_{2}A^{s-2} + \dots + a_{s}I$$
(3)

and therefore

$$A^{n} = a_{1}A^{n-1} + a_{2}A^{n-2} + \dots + a_{s}A^{n-s}$$

for all  $n \ge s$ . Applying  $\varphi$  proves the statement.

ι

It is unclear whether a similar statement to Lemma 4 is true for non-commutative rings. While there exist versions of the Cayley–Hamilton theorem for non-commutative rings (see for example [7, 17]), they cannot be applied to obtain an equation similar to (3).

The next observation states that LRS are equivalent to generalized moment sequences. It can be found in [14]:

**Lemma 5.** Let  $(u_n)_{n \in \mathbb{N}}$  be a sequence in a commutative unital ring  $\mathcal{R}$ . The following are equivalent:

- (i)  $(u_n)_{n \in \mathbb{N}}$  is a LRS of order s.
- (ii) There is a matrix  $A \in \operatorname{Mat}_{s}(\mathcal{R})$  and two vectors  $v, w \in \mathcal{R}^{s}$  such that  $u_{n} = v^{t}A^{n-s}w$  for all n > s.

*Proof.* For  $(i) \Rightarrow (ii)$  assume that the recurrence is given by

$$u_n = a_1 u_{n-1} + a_2 u_{n-2} + \dots + a_s u_{n-s}.$$

Using the companion matrix

$$A = \begin{pmatrix} a_1 & 1 & & \\ a_2 & 1 & & \\ \vdots & & \ddots & \\ a_{s-1} & & & 1 \\ a_s & & & \end{pmatrix}$$

we have that  $u_n = v^t A^{n-s} w$  where  $v = (u_s, u_{s-1}, ..., u_1)^t$  and  $w = (1, 0, ..., 0)^t$ .

The proof of  $(ii) \Rightarrow (i)$  is analogous to Lemma 4, by replacing tr by the function  $A \mapsto v^t A w$ . Note that the recurrence starts to hold only for n > 2s, but for our purposes this is irrelevant.

## 3 Decidable cases

In the following we present cases in which the moment membership problem is decidable. This includes known results for small s (Section 3.1), the moment positivity problem for unitary and orthogonal matrices (Section 3.2), and for matrices with a unique largest eigenvalue or only real eigenvalues(Section 3.3). Throughout this section we will consider  $\mathcal{P} = \mathbb{R}_{\geq 0}$ .

#### 3.1 Known results: small order

We first review known results on the decidability of the moment positivity problem. The results are about LRS, but in view of Lemma 4, they immediately transfer to moments.

**Theorem 6.** The moment positivity problem is decidable in the following cases:

(i)  $s \leq 5, \mathcal{D} = \operatorname{Mat}_s(\mathbb{Q}).$ 

(ii)  $s \leq 9, \mathcal{D} \subseteq \operatorname{Mat}_{s}(\mathbb{Q})$  the set of matrices with simple eigenvalues.

The proof of (i) is contained in [14], the proof of (ii) goes back to [15]. Decidability for other values of s is unknown.

The positivity problem of LRS is closely related to *Skolem's Problem* which asks if some sequence element equals 0. The best result in this context is that Skolem's Problem is NP-hard [2]. The decidability of the positivity problem implies decidability of Skolem's Problem. This follows for an integer LRS because  $u_n \neq 0$  if and only if  $u_n^2 - 1 \ge 0$ . If  $(u_n)_{n \in \mathbb{N}}$  is an LRS of order s, then  $u_n^2 - 1$  is an LRS of order  $s^2$ . Moreover, since Skolem's Problem is NP-hard, the positivity problem is NP-hard as well.

#### 3.2 Orthogonal and unitary matrices

We now show that the moment positivity problem for orthogonal (Theorem 8) and unitary matrices (Corollary 11) is decidable. The proof strategy is very similar to [3].

A set  $X \subseteq \mathbb{R}^m$  is algebraic if there are polynomials  $p_1, \ldots, p_n \colon \mathbb{R}^m \to \mathbb{R}$  such that

$$X = \{ x \in \mathbb{R}^m \mid p_1(x) = \dots = p_n(x) = 0 \}.$$

In this case, X is the algebraic variety defined by  $p_1, \ldots, p_n$ , and we write  $X = \mathcal{V}(p_1, \ldots, p_n)$ . Even if the set of defining polynomials is infinite, there always exists a finite choice of polynomials defining the same algebraic variety, by Hilbert's basis theorem. Since we work over  $\mathbb{R}$ , we can even reduce it to a single polynomial by taking the sum of squares of the defining polynomials.

For matrices  $A_1, \ldots, A_d \in \operatorname{Mat}_s(\mathbb{R})$ , let

$$\langle A_1, A_2, \dots, A_d \rangle \coloneqq \{A_{k_1} \cdots A_{k_\ell} \mid \ell \in \mathbb{N}, k_1, \dots, k_\ell = 1, \dots, d\}$$

be the semigroup generated by  $A_1, \ldots, A_d$ . We denote by  $\overline{\langle A_1, \ldots, A_d \rangle}$  the topological closure inside  $Mat_s(\mathbb{R})$  with respect to the Euclidean topology.

**Lemma 7.** Let  $A_1, \ldots, A_d \in O_s(\mathbb{Q})$  be orthogonal  $s \times s$  matrices with rational entries. Then  $\mathcal{G} \coloneqq \overline{\langle A_1, \ldots, A_d \rangle}$  is a compact algebraic group. Moreover there is a recursively enumerable sequence of rational polynomials  $(p_k)_{k \in \mathbb{N}}$  defining  $\mathcal{G}$  inside  $\operatorname{Mat}_s(\mathbb{R})$ .

*Proof.* Compactness of  $\mathcal{G}$  is obvious. To prove that  $\mathcal{G}$  is a group we only have to show that  $A^{-1} \in \mathcal{G}$  for every  $A \in \mathcal{G}$ . Consider the sequence  $(A^k)_{k \in \mathbb{N}}$ . By compactness, there exists a converging subsequence. In other words, for every  $\varepsilon > 0$ , there exists  $n_2 > n_1 + 1$  such that

$$||A^{n_1} - A^{n_2}|| < \epsilon$$

where  $\|\cdot\|$  is the operator norm. Since  $\|A \cdot B\| = \|B\|$  for every matrix B, we obtain

$$||A^{-1} - A^{n_2 - n_1 - 1}|| < \varepsilon.$$

This shows that  $A^{-1} \in \mathcal{G}$ .

Now note that every compact group  $\mathcal{G} \subseteq \operatorname{Mat}_{s}(\mathbb{R})$  is algebraic (see for example [12, Chapter 3, Section 4.4]). In particular, it is shown there that

$$\mathcal{G} = \mathcal{V}(\mathbb{R}[X]^{\mathcal{G}}) := \mathcal{V}(p \in \mathbb{R}[X] \mid p(I_s) = 0, \ p(gX) = p(X) \text{ for all } g \in \mathcal{G}),$$

where  $I_s$  is the identity matrix of size s.

Now note that if  $\mathcal{G}$  is generated by  $A_1, \ldots, A_d$ , then the invariance only needs to be checked with respect to the generators, i.e.

$$\mathcal{G} = \mathcal{V}\Big(p \in \mathbb{R}[X] \mid p(I_s) = 0, \ p(A_i X) = p(X) \ \text{ for } i = 1, \dots, d\Big).$$

Since the conditions  $p(I_s) = 0$  and  $p(A_iX) = p(X)$  are linear in the coefficients of p, there exists a basis  $(p_k)_{k \in \mathbb{N}}$  of the space of solutions of these conditions. Moreover, the coefficients of the basis vectors  $p_k$  can be chosen from  $\mathbb{Q}$ , since all conditions are rational. We now have

$$\mathcal{G} = \mathcal{V}(p_k \mid k \in \mathbb{N}).$$

The polynomials  $p_k$  can be computed recursively by solving the system of linear equations over the space of polynomials with degree d, and by increasing d iteratively.

Note that the statement is not true if  $\mathbb{R}$  is replaced by  $\mathbb{C}$ . For example the group

$$\mathcal{G} \coloneqq \Big\{ e^{i\theta} \mid \theta \in [0, 2\pi) \Big\},\,$$

seen as a subset of  $1 \times 1$  matrices, is not algebraic. Yet, we shall see that the moment problem also generalizes to unitary matrices (see Corollary 11).

Since  $\mathbb{R}[X]$  is a Noetherian ring, there exists  $n \in \mathbb{N}$  such that

$$\mathcal{G} = \mathcal{V}(p_1, \ldots, p_n).$$

This will be an important ingredient to prove the decidability of the moment problem. Note however that n can be arbitrarily large and it is unclear whether n is computable or not.

**Theorem 8.** The moment positivity problem for  $\mathcal{D} = O_s(\mathbb{Q})$  is decidable.

*Proof.* We will present two procedures, each certifying either yes- or no-instances in finite time. Letting these algorithms run in parallel will result in a decision algorithm for the problem.

Certifying no-instances for  $A \in O_s(\mathbb{Q})$  is achieved by iteratively checking whether  $\operatorname{tr}(A^n) \ge 0$  holds for every n. If A is a no-instance, this algorithm will halt when detecting  $\operatorname{tr}(A^n) < 0$  for the first time.

We now present an algorithm to certify yes-instances in finite time. For a given  $A \in O_s(\mathbb{Q})$ , the moment membership problem can be rephrased as

$$\forall B \in \langle A \rangle : \operatorname{tr}(B) \ge 0.$$

By the continuity of the trace, this is equivalent to

$$\forall B \in \overline{\langle A \rangle} : \operatorname{tr}(B) \ge 0. \tag{4}$$

By Lemma 7 there exists a recursively enumerable sequence of polynomials  $(p_k)_{k \in \mathbb{N}}$  and some  $n \in \mathbb{N}$  such that

$$\overline{\langle A \rangle} = \mathcal{V}(p_1, \dots, p_n).$$

Now step k of the algorithm verifies the statement

$$\forall B \in \mathcal{V}(p_1, \dots, p_k) \colon \operatorname{tr}(B) \ge 0 \tag{5}$$

which is decidable by the Tarski–Seidenberg Theorem, since it is a statement in first order logic. As soon as Eq. (5) is true for the first time, the algorithm halts and outputs a correct yes-answer. This will be the case after at most n steps, if A is a yes-instance.  $\Box$ 

**Remark 9.** The previous statement can be generalized in two directions:

(i) By the same argument, the following problem is also decidable: Given  $A_1, \ldots, A_d \in O_s(\mathbb{Q})$  for a fixed matrix size s, decide if:

$$\forall \ell \in \mathbb{N} \; \forall k_1, \dots, k_\ell \in \{1, \dots, d\} \colon \operatorname{tr}(A_{k_1} \cdots A_{k_\ell}) \ge 0.$$

Note that this decision problem for arbitrary matrices is undecidable [4].

(*ii*) The proof remains true if tr is replaced by any other continuous function. This in particular implies that the generalized problem

$$\forall n \in \mathbb{N} \colon \varphi(A^n) \ge 0$$

is decidable.

We now generalize the result to unitary matrices, by embedding them into orthogonal matrices of larger size. We denote by  $\mathbb{Q}[i]$  the field of complex numbers with rational real and imaginary parts, and we denote the set of  $s \times s$  unitary matrices with entries in  $\mathbb{Q}[i]$  by  $U_s(\mathbb{Q}[i])$ .

Lemma 10. The map

$$\Psi \colon \mathbf{U}_s(\mathbb{Q}[i]) \to \mathbf{O}_{2s}(\mathbb{Q})$$
$$U = A + iB \mapsto \begin{pmatrix} A & -B \\ B & A \end{pmatrix}$$

is a group homomorphism. Moreover we have

$$\operatorname{tr}(U) = \frac{1}{2} \operatorname{tr} \left( \Psi(U) \cdot \begin{pmatrix} I_s & iI_s \\ -iI_s & I_s \end{pmatrix} \right).$$

 $\triangle$ 

*Proof.* The map is well defined since  $\Psi(U)$  is orthogonal if and only if U is unitary. The rest is immediate.

The main results of this section are summarized in the following two corollaries.

**Corollary 11.** For each  $s \ge 1$ , the moment positivity problem for matrices from  $U_s(\mathbb{Q}[i])$  is decidable.

*Proof.* It follows immediately from Lemma 10, Theorem 8 and Remark 9 (ii).

**Corollary 12.** The positivity problem is decidable for simple unitary LRS, i.e.

$$u_n = a_1 u_{n-1} + \dots + a_s u_{n-s}$$

with  $a_1, \ldots, a_s \in \mathbb{Q}[i]$ , where the roots of  $p(x) = x^s - a_1 x^{s-1} - a_2 x^{s-2} - \cdots - a_s$  are all simple and of modulus 1.

*Proof.* We choose a unitary matrix  $A \in U_s(\mathbb{C})$  whose eigenvalues are the roots of p, and whose entries are computable numbers. For example, one can take a diagonal matrix with the specified roots on the diagonal. We obtain the recurrence

$$A^n = a_1 A^{n-1} + \dots + a_s A^{n-s}$$

for all  $n \ge s$ , and since the roots are all simple, p is actually the minimal polynomial of A. So  $I_s, A, A^2, \ldots, A^{s-1}$  are linearly independent, and we can thus find a linear functional  $\varphi$  on  $\operatorname{Mat}_s(\mathbb{C})$  with  $\varphi(A^i) = u_i$  for  $i = 0, \ldots, s - 1$ . Now, by Remark 9 and Lemma 10 above, it is decidable whether  $\varphi(A^i) \ge 0$  holds for all i, and since this sequence fulfills the same recurrence and initial conditions as  $(u_i)_{i\ge 1}$ , the two sequences coincide.

#### 3.3 Matrices with a unique dominant eigenvalue or real eigenvalues

In the following, we show that for matrices with a unique dominant eigenvalue, and for matrices with only real eigenvalues, the moment problem is decidable. Note that the idea for the case of a unique dominating eigenvalue is already present in [14], but restricted to multiplicity 1 and matrices of size at most s = 5.

**Theorem 13.** The moment positivity problem is decidable in the following cases:

- (i)  $\mathcal{R} = \mathbb{Q}$ , s arbitrary, and the set of instances restricted to matrices with a unique dominant eigenvalue.
- (ii)  $\mathcal{R} = \mathbb{Q}$ , s arbitrary, and the set of instances restricted to matrices with only real eigenvalues.

*Proof.* We provide algorithms that decide the moments positivity problem for the stated instance sets. We can assume without loss of generality that  $A \in \operatorname{Mat}_{s}(\mathbb{Z})$ , by possibly multiplying the matrix with the largest denominator of its entries.

For (i) let  $A \in \operatorname{Mat}_{s}(\mathbb{Z})$  have a unique dominant eigenvalue. Since A has real entries, the non-real eigenvalues of A come in conjugate pairs. Since there is exactly one eigenvalue  $\lambda_{1}$  of largest absolute value, it must be real. We let k denote its multiplicity and obtain

$$|\mu_n(A) - k \cdot \lambda_1^n| \leq (s-k)|\lambda_2|^n,$$

where  $\lambda_2$  denotes the second largest eigenvalue in absolute value. Thus it suffices to check  $\mu_n(A) \ge 0$  for n up to

$$\frac{\log(s/k-1)}{\log(|\lambda_1|) - \log(|\lambda_2|)}.$$

(*ii*): In this case only odd moments matter, since the even moments are always nonnegative. If the dominant eigenvalues all have the same sign, then we can apply (*i*). Otherwise, since odd powers of eigenvalues with the same absolute values but different signs cancel out, we can reduce the problem to a smaller matrix, where the dominant eigenvalues do have the same sign.  $\Box$ 

#### 3.4 Further generalizations

In the following, we present a generalization of the statements in Section 3.2 and Section 3.3. For a matrix  $A \in Mat_s(\mathbb{R})$ , we denote by spec(A) the multi-set of all eigenvalues of A (where multiple eigenvalues are represented by multiple elements of spec(A)). Express

$$\operatorname{spec}(A) = \operatorname{per}_1(A) \cup \operatorname{per}_2(A) \cup \cdots \cup \operatorname{per}_s(A)$$

as a partition of peripheral spectra, i.e. eigenvalues of the same absolute value, in decreasing order (i.e.  $per_1(A)$  contains the dominant eigenvalues,  $per_2(A)$  the eigenvalues of second largest absolute value...). Note that  $per_i(A)$  can be empty if A has multiple eigenvalues of same absolute value. Moreover, let

$$\mu_n^{(i)}(A)\coloneqq \sum_{\lambda\in \mathsf{per}_i(A)} \left(\frac{\lambda}{|\lambda|}\right)^n.$$

We define

$$\eta_i(A) = \begin{cases} \inf_{n \in \mathbb{N}} \mu_n^{(i)}(A) & : \text{ if } \mathsf{per}_i(A) \neq \emptyset \\ \infty & : \text{ if } \mathsf{per}_i(A) = \emptyset \end{cases}$$

and

$$\gamma_i(A) = \begin{cases} \sup_{n \in \mathbb{N}} \mu_{pn+q}^{(i)}(A) & : \text{ if } \mathsf{per}_i(A) \neq \emptyset \\ -\infty & : \text{ if } \mathsf{per}_i(A) = \emptyset \end{cases}$$

where  $p, q \ge 1$  are arbitrary but fixed integers. So we compute the supremum along an arithmetic progression.

**Lemma 14.** The following two problems are decidable:

- (i) Given<sup>1</sup>  $A \in Mat_s(\mathbb{R}), c \in \mathbb{R}$ , decide whether  $\eta_i(A) \ge c$ .
- (ii) Given  $A \in \operatorname{Mat}_{s}(\mathbb{R}), c \in \mathbb{R}$ , decide whether  $\gamma_{i}(A) \leq c$ .

*Proof.* The decision algorithms are very similar to one from the proof of Theorem 8. To construct an algorithm for (i), let the following two procedures run in parallel:

(a) Evaluate  $\mu_n^{(i)}(A)$  for increasing  $n \in \mathbb{N}$ . Halt if  $\mu_n^{(i)}(A) < c$ .

 $<sup>^{1}</sup>$ To assume that the inputs have a finite description, we restrict to algebraic numbers, i.e. numbers that can be represented as roots of an integer polynomial. This is enough to apply Lemma 14 in the proof of Theorem 15.

(b) Check the statement

$$\forall B \in \mathcal{V}(p_1, \dots, p_k) \colon \operatorname{tr}(B) \ge c$$

for increasing  $k \in \mathbb{N}$ , where  $(p_{\ell})_{\ell \in \mathbb{N}}$  define the variety  $\overline{\langle U \rangle}$ , where U is the diagonal matrix with eigenvalues  $\lambda/|\lambda|$  for  $\lambda \in \mathsf{per}_i(A)$ . Halt if the statement is true.

If A, c is a no-instance of (i), then (a) will eventually halt; if A, c is a yes-instance, (b) will eventually halt, for the same reason as in the proof of Theorem 8.

The algorithm for (ii) is very similar. Let the following two procedures run in parallel:

- (a) Evaluate  $\mu_{pn+q}^{(i)}(A)$  for increasing  $n \in \mathbb{N}$ . Halt if  $\mu_{pn+q}^{(i)}(A) > c$ .
- (b) Check the statement

$$\forall B \in \mathcal{V}(p_1, \dots, p_k) \colon \operatorname{tr}(U^q B) \leqslant c$$

for increasing  $k \in \mathbb{N}$ , where  $(p_{\ell})_{\ell \in \mathbb{N}}$  define the group  $\overline{\langle U^p \rangle}$ , where U is the diagonal matrix with eigenvalues  $\lambda/|\lambda|$  for  $\lambda \in \mathsf{per}_i(A)$ . Halt if the statement is true.

In (b) we only evaluate odd moments; recall Remark 9 (ii).

It is unclear whether  $\eta_i(A) > c$  or  $\eta_i(A) = c$  is decidable. This is due to the fact that we do not know whether  $\mu_n^{(i)}(A)$  attains the infimum/supremum for finite n.

**Theorem 15.** For a fixed parameter  $\varepsilon > 0$ , the moment positivity problem is decidable for all non-zero matrices A satisfying one of the following conditions:

- (i)  $\exists k \in \mathbb{N} : \eta_1(A), \dots, \eta_k(A) \ge 0, \eta_{k+1}(A) \ge \varepsilon.$
- (*ii*)  $\exists k \in \mathbb{N} \colon \gamma_1(A), \dots, \gamma_k(A) \leq 0, \gamma_{k+1}(A) \leq -\varepsilon.$

(*iii*) 
$$\eta_1(A) < 0$$
.

If (ii) or (iii) are satisfied, then A is automatically a no-instance. If (i) is satisfied, then A can be a yes or a no-instance. Moreover, each of the above criteria is decidable.

*Proof.* First, checking whether A satisfies (i), (ii) or (iii) is decidable by Lemma 14, since there are only finitely many of these statements to check.

To prove (i), assume that  $\eta_{k+1}(A) \neq \infty$  (the other case is trivial). Let  $\lambda_i \in \mathsf{per}_i(A)$ . We have that

$$\mu_n(A) = \sum_{i=1}^d |\lambda_i|^n \mu_n^{(i)}(A) \ge |\lambda_{k+1}|^n \left(\varepsilon - s \sum_{i=k+2}^d \left(\frac{|\lambda_i|}{|\lambda_{k+1}|}\right)^n\right)$$

which is positive for

$$n \ge \frac{\log(\varepsilon) - \log(sd)}{\log(|\lambda_{k+2}|) - \log(|\lambda_{k+1}|)}$$

So we only need to check finitely many instances of the problem.

For (ii) we have that

$$\mu_m(A) = \sum_{i=1}^d |\lambda_i|^m \mu_m^{(i)}(A) < |\lambda_{k+1}|^m \left(-\varepsilon + s \sum_{i=k+2}^d \left(\frac{|\lambda_i|}{|\lambda_{k+1}|}\right)^m\right).$$

Now there clearly exists some m of the form pn+q such that the right hand side is negative.

For (*iii*) note that  $\eta_1(A) < 0$  is decidable since  $\eta_1(A) \ge 0$  is decidable by Lemma 14. Let  $0 < \delta < -\eta_1(A)$ . Then there exists an increasing sequence  $(n_\ell)_{\ell \in \mathbb{N}}$  such that  $\mu_{n_\ell}^{(1)}(A) < \eta_1(A) + \delta < 0$  for all  $\ell$ .<sup>2</sup> Therefore we have

$$\mu_{n_{\ell}}(A) = \sum_{i=1}^{d} |\lambda_i|^{n_{\ell}} \mu_{n_{\ell}}^{(i)}(A) < |\lambda_1|^{n_{\ell}} \left( \eta_1(A) + \delta + s \sum_{i=2}^{d} \left( \frac{|\lambda_i|}{|\lambda_1|} \right)^{n_{\ell}} \right).$$

Again there exists  $\ell_0$  such that  $\mu_{n_{\ell_0}}(A) < 0$ .

## 4 Undecidable cases

We now present two finitely generated rings for which the moment membership problem is undecidable. Specifically, in Section 4.1 we prove that the moment membership problem is undecidable for the ring of commutative polynomials  $\mathcal{R} = \mathbb{Z}[x_1, \ldots, x_d]$  if n is sufficiently large. In Section 4.2, we show that the moment membership problem is also undecidable for the space of non-commutative polynomials  $\mathcal{R} = \mathbb{Z}\langle z_1, \ldots, z_d \rangle$ . For a similar undecidable problem including nonnegative and sos polynomials polynomials over infinitely many variables, we refer to [5, Section 6].

#### 4.1 Commutative polynomial rings

In the following, we show that the generalized moment membership problem for  $\mathcal{R} = \mathbb{Z}[x_1, \ldots, x_d]$  and the cone

 $\mathcal{P}_{\text{coeff}} \coloneqq \{ p \in \mathbb{Z}[x_1, \dots, x_d] \mid \text{all coefficients of } p \text{ are nonnegative} \}$ 

is undecidable. The proof relies on the following undecidable problem, for which a proof can be found in Appendix A.

**Lemma 16.** For large enough values of s and d, and a suitable matrix  $N \in Mat_s(\mathbb{Z})$ , the following problem is undecidable: Given  $A_1, \ldots, A_d \in Mat_s(\mathbb{Z})$ , do there exist  $n_1, \ldots, n_d \in \mathbb{N}$  with

$$\operatorname{tr}(A_1^{n_1} \cdot A_2^{n_2} \cdots A_d^{n_d} \cdot N) < 0?$$

**Problem 17.** Let  $M \in Mat_s(\mathcal{R})$  be a fixed matrix. For an input  $A \in Mat_s(\mathcal{R})$ , decide whether

$$\operatorname{tr}(A^n \cdot M) \in \mathcal{P}$$

holds for all  $n \ge 1$ .

**Theorem 18.** If  $s, d \in \mathbb{N}$  are large enough, and M is chosen suitably, then Problem 17 is undecidable for  $\mathcal{R} = \mathbb{Z}[x_1, \ldots, x_d]$  and  $\mathcal{P}_{coeff}$ .

$$\eta_i(U) \leqslant \operatorname{tr}(U^{n_\ell}) \leqslant \eta_i(U) + \frac{1}{\ell}$$

<sup>&</sup>lt;sup>2</sup>This follows from the fact that for a unitary matrix U the group  $\overline{\{U^n \mid n \in \mathbb{N}\}}$  is either finite or contains no isolated points. Because if the set contains an isolated point, then all elements are isolated. But a compact set which contains only isolated points is finite. Hence there exists an increasing sequence  $(n_\ell)_\ell$  such that

*Proof.* Given  $A_1, \ldots, A_d \in \operatorname{Mat}_s(\mathbb{Z})$ , set

$$A = \sum_{i=1}^{d} \left( \sum_{1 \leq j \leq i} e_j e_i^t \right) \otimes A_i \cdot x_i \in \operatorname{Mat}_{ds}(\mathcal{R}).$$

Moreover, define

$$M = mm^t \otimes N$$

with m = (1, ..., 1) and N as in Lemma 16. We have that

$$\operatorname{tr}(A^{n}M) = \sum_{1 \leq i_{1} \leq \dots \leq i_{n} \leq d} i_{1} \cdot \operatorname{tr}(A_{i_{1}} \cdots A_{i_{n}} \cdot N) \cdot x_{i_{1}} \cdots x_{i_{n}}$$
$$= \sum_{n_{1} + \dots + n_{d} = n} \underbrace{c_{n_{1},\dots,n_{d}}}_{\geq 1} \cdot \operatorname{tr}(A_{1}^{n_{1}} \cdots A_{d}^{n_{d}} \cdot N) \cdot x_{1}^{n_{1}} \cdots x_{d}^{n_{d}}$$

where  $c_{n_1,\ldots,n_d} = \min\{i \mid n_i \neq 0\}$ . Thus Problem 17 reduces to the undecidable problem from Lemma 16.

**Remark 19.** Since the sequence  $tr(A^nM)$  is clearly a LRS (see Lemma 4), the previous result shows that positivity of LRS over  $\mathcal{R} = \mathbb{Z}[x_1, \ldots, x_d]$  is undecidable in general.  $\triangle$ 

## 4.2 Non-commutative polynomial rings

We now consider the ring  $\mathcal{R} = \mathbb{Z}\langle z_1, \ldots, z_d \rangle$  of non-commutative polynomials, and show that its moment membership problem is undecidable for the cone of polynomials with positive coefficients. As a Z-module, a basis of  $\mathcal{R}$  consists of all words in the letters  $z_1, \ldots, z_d$  where the order of letters *does* matter. Concatenation of words extends to a multiplication making  $\mathcal{R}$  a unital ring, where 1 corresponds to the empty word. There is a slightly different way to define this object, namely just as the tensor algebra (cf. Example 1)

$$\mathbb{Z}\langle z_1,\ldots,z_d\rangle=T(\mathbb{Z}^d).$$

The equivalence of definitions is apparent when identifying a word  $z_{k_1} \cdots z_{k_m}$  with the element  $e_{k_1} \otimes \cdots \otimes e_{k_m} \in (\mathbb{Z}^d)^{\otimes m}$ , where  $e_r$  denotes the *r*-th standard basis vector in  $\mathbb{Z}^d$ . We equip  $\mathcal{R}$  with two (a priori) different sets of positive elements:

$$\mathcal{P}_{\text{coeff}} \coloneqq \mathbb{Z}_{\geq 0} \langle z_1, \dots, z_d \rangle = \{ p \in \mathbb{Z} \langle z_1, \dots, z_d \rangle \mid \text{all coefficients of } p \text{ are nonnegative} \}$$

$$\mathcal{P}_{\text{eval}} \coloneqq \{ p \in \mathbb{Z} \langle z_1, \dots, z_d \rangle \mid \forall \ell, A_1, \dots, A_d \in \text{Mat}_{\ell}(\mathbb{Z}_{\geq 0}) \colon p(A_1, \dots, A_d) \in \text{Mat}_{\ell}(\mathbb{Z}_{\geq 0}) \}.$$

We now show that both cones coincide, which is a free version of Pólya's Theorem.

**Theorem 20** (Free Pólya's Theorem). Let  $p \in \mathbb{C}\langle z_1, \ldots, z_d \rangle$  with  $m \coloneqq \deg(p)$ . Then the following are equivalent:

- (i) All coefficients of p are nonnegative reals.
- (ii) For all  $A_1, \ldots, A_d \in \operatorname{Mat}_{m+1}(\mathbb{Z}_{\geq 0})$  we have

$$p(A_1,\ldots,A_d) \in \operatorname{Mat}_{m+1}(\mathbb{R}_{\geq 0}).$$

In particular,  $\mathcal{P}_{coeff} = \mathcal{P}_{eval}$ , and in the definition of  $\mathcal{P}_{eval}$  one can restrict  $\ell$  to deg(p) + 1.

*Proof.*  $(i) \Rightarrow (ii)$  is obvious (even without the restriction on the matrix size m + 1). For  $(ii) \Rightarrow (i)$  we construct matrices  $A_1, \ldots, A_d$  that allow us to isolate a single coefficient of p.

Let  $z_{k_1} \cdots z_{k_\ell}$  be a word in the letters  $z_1, \ldots, z_d$ . For  $j = 1, \ldots, d$  define

$$A_j \coloneqq \sum_{i=1,\dots,\ell; \ k_i=j} E_{i,i+1} \in \operatorname{Mat}_{\ell+1}(\mathbb{Z}_{\geq 0}).$$

where  $E_{i,j}$  denotes the matrix (of size  $\ell + 1$ ) with a 1 in position (i, j) and zeros elsewhere. For  $t_1, \ldots, t_r \in \{1, \ldots, d\}$  we have

$$A_{t_{1}} \cdots A_{t_{r}} = \sum_{\substack{i \\ k_{i} = t_{1} \\ k_{i+1} = t_{2} \\ \vdots \\ k_{i+r-1} = t_{r}}} E_{i,i+r} \in \operatorname{Mat}_{\ell+1}(\mathbb{Z}_{\geq 0}).$$

In particular, the  $(1, \ell + 1)$ -entry of a product  $A_{t_1} \cdots A_{t_r}$  is 1 if and only if  $r = \ell$  and  $(k_1, \ldots, k_\ell) = (t_1, \ldots, t_\ell)$ ; in all other cases it is zero. So  $p(A_1, \ldots, A_d)$  contains in its upper right entry precisely the coefficient of p at the word  $z_{k_1} \cdots z_{k_\ell}$ .

Since all words appearing in p are of length at most deg(p) = m, we can do this procedure with matrices  $A_j$  of size at most m + 1, and thus clearly with matrices of size exactly m + 1.

**Remark 21.** Pólya's theorem [16, 8] states that for every homogeneous polynomial  $p \in \mathbb{R}[x_1, \ldots, x_d]$  that is strictly positive on the *d*-simplex

$$\Delta_d \coloneqq \left\{ (a_1, \dots, a_d) \in \mathbb{R}^d \mid a_i \ge 0, \sum_{i=1}^d a_i = 1 \right\},\$$

the polynomial

$$(x_1 + \dots + x_d)^n \cdot p(x_1, \dots, x_d)$$

has positive coefficients, for sufficiently large  $n \in \mathbb{N}$ . In Theorem 20, the space of nonnegative matrices takes the role of the *d*-simplex. While in the commutative case we have to multiply p with an additional polynomial, this is not so in the free version.  $\triangle$ 

We now show that for these cones the moment membership problem is undecidable.

**Theorem 22.** Let  $d, s \ge 7$ . Then the moment membership problem for  $\mathcal{R} = \mathbb{Z}\langle z_1, \ldots, z_d \rangle$ ,  $\mathcal{P}_{\text{coeff}} = \mathcal{P}_{\text{eval}}$  and s is undecidable. This remains true if we restrict the instances to linear matrix polynomials, i.e.  $A \in \text{Mat}_s(\mathbb{Z}\langle z_1, \ldots, z_d \rangle)$  whose entries are linear forms in  $z_1, \ldots, z_d$ .

*Proof.* For  $A = \sum_{k=1}^{d} z_k A_k$  with  $A_k \in \operatorname{Mat}_s(\mathbb{Z})$  we have

$$\mu_n(A) = \sum_{k_1, \dots, k_n = 1}^d \operatorname{tr}(A_{k_1} \cdots A_{k_n}) \cdot z_{k_1} \cdots z_{k_n}.$$

So  $\mu_n(A) \in \mathcal{P}_{\text{coeff}}$  means that  $\operatorname{tr}(A_{k_1} \cdots A_{k_n}) \ge 0$  for all  $k_1, \ldots, k_n = 1, \ldots, d$ . Undecidability of this problem was proven in [4, Lemma 3].

# 5 Conclusion

We have studied the moment membership problem (Problem 2) for matrices over a ring. We have shown that there is a relation to LRS for commutative rings (Lemma 4 and Lemma 5) and that the moments positivity problem is decidable in many cases, including unitary and orthogonal matrices (Theorem 8 and Corollary 11) as well as matrices with a unique dominating eigenvalue or only real eigenvalues (Theorem 13). Finally, we have shown that the generalized moment membership problem is undecidable over the ring of commutative and non-commutative polynomials, where the positivity cone is given by the set of polynomials with non-negative coefficients (Theorem 18 and Theorem 22).

The central open question remains, namely whether the moment membership problem is decidable or undecidable for  $\mathcal{R} = \mathbb{Q}$  and  $\mathcal{P} = [0, \infty)$ . In the context of rings it would be interesting whether it is undecidable for commutative polynomials for the cone of sumof-square polynomials or the non-negative polynomials. This might be the case, as these cones have a richer structure than that of polynomials with nonnegative coefficients.

# References

- P. Bell, V. Halava, T. Harju, J. Karhumaki, and I. Potapov. Matrix Equations and Hilbert's Tenth Problem. Int. J. Algebra Comput., 18:1231–1241, 12 2008. doi:10.1142/S0218196708004925.
- [2] V. D. Blondel and N. Portier. The presence of a zero in an integer linear recurrent sequence is NP-hard to decide. *Linear Algebra Appl.*, 351:91–98, 2002. doi:10.1016/S0024-3795(01)00466-9.
- [3] V. D. Blondel, E. Jeandel, P. Koiran, and N. Portier. Decidable and undecidable problems about quantum automata. SIAM J. Comput., 34:1464–1473, 2005. doi:10.1137/S0097539703425861.
- [4] G. De las Cuevas, T. S. Cubitt, J. I. Cirac, M. M. Wolf, and D. Pérez-García. Fundamental limitations in the purifications of tensor networks. J. Math. Phys., 57(7): 071902, 2016. doi:10.1063/1.4954983.
- [5] G. De las Cuevas, A. Klingler, and T. Netzer. Polynomial decompositions with invariance and positivity inspired by tensors. arXiv:2109.06680, 2021. doi:10.48550/arXiv.2109.06680.
- [6] G. Everest, A. van der Poorten, I. Shparlinski, and T. Ward. *Recurrence Sequences*. American Mathematical Soc., 2003. ISBN 1470423154.
- [7] R. Grosu. The Cayley-Hamilton Theorem for Noncommutative Semirings. In Implementation and Application of Automata, pages 143–153, Berlin, Heidelberg, 2011. Springer Berlin Heidelberg. ISBN 978-3-642-18098-9. doi:10.1007/978-3-642-18098-9\_16.
- [8] G. H. Hardy, J. E. Littlewood, and G. Pólya. *Inequalities*. Cambridge University Press, 1952.
- [9] A. Klingler, M. van der Eyden, S. Stengele, T. Reinhart, and G. De las Cuevas. Many bounded versions of undecidable problems are NP-hard. *SciPost Physics*, 14(6):173, 2023. doi:10.21468/SciPostPhys.14.6.173.

- [10] S. Lang. Algebra, volume 211. Springer Science & Business Media, 2012. ISBN 9781461300410.
- [11] O. Martin, A. M. Odlyzko, and S. Wolfram. Algebraic properties of cellular automata. Comm. Math. Phys., 93:219–258, 1984.
- [12] A. Onishchik and E. Vinberg. Lie Groups and Algebraic Groups. Springer, 1990.
- [13] J. Ouaknine and J. Worrell. Decision problems for linear recurrence sequences. In International Workshop on Reachability Problems, pages 21–28. Springer, 2012. doi:10.1007/978-3-642-33512-9\_3.
- [14] J. Ouaknine and J. Worrell. Positivity problems for low-order linear recurrence sequences. In Proceedings of the Twenty-Fifth Annual ACM-SIAM Symposium on Discrete Algorithms, pages 366–379. ACM, New York, 2014.
- [15] J. Ouaknine and J. Worrell. On the positivity problem for simple linear recurrence sequences. In Automata, languages, and programming. Part II, volume 8573 of Lecture Notes in Comput. Sci., pages 318–329. Springer, Heidelberg, 2014. doi:10.1007/978-3-662-43951-7\_27.
- [16] G. Pólya. Uber positive Darstellung von Polynomen. Vierteljschr. Naturforsch. Ges. Zürich, 73(141-145):2, 1928.
- [17] J. O. Szigeti. Caley-Hamilton Theorem for Matrices over an arbitrary Ring. Serdica Math. J, 32:269–276, 2006.
- [18] R. C. Tausworthe. Random numbers generated by linear recurrence modulo two. Mathematics of Computation, 19(90):201–209, 1965.
- [19] R. Tijdeman, M. Mignotte, and T. Shorey. The distance between terms of an algebraic recurrence sequence. Journal f
  ür die reine und angewandte Mathematik, 349:63–76, 1984.
- [20] N. K. Vereshchagin. The problem of appearance of a zero in a linear recurrence sequence. Mat. Zametki, 38(2):609–615, 1985.

## A A version of the matrix mortality problem

In this appendix we prove Lemma 16 with a similar proof idea as in [4, 9], which is based on a reduction from an undecidable version of the matrix mortality problem.

We start with a version of the matrix mortality problem known to be undecidable.

**Problem 23** (Matrix Mortality version). Let  $A_1, \ldots, A_d \in \text{Mat}_s(\mathbb{Z})$ . Does there exist a choice of  $n_1, \ldots, n_d \in \mathbb{N}$  such that

$$A_1^{n_1} \cdot A_2^{n_2} \cdots A_d^{n_d} = 0?$$

**Theorem 24.** If s and d are large enough, then Problem 23 is undecidable.

For a proof of Theorem 24 we refer to [1]. We are now ready to prove Lemma 16.

Proof of Lemma 16. We provide a reduction from Problem 23. First, fix the matrix

$$N = \begin{pmatrix} \mathbf{0} & 0\\ 0 & 1 \end{pmatrix} + \sum_{i,j=1}^{s} \begin{pmatrix} E_{ij} \otimes E_{ij} & 0\\ 0 & 0 \end{pmatrix} \in \operatorname{Mat}_{s}(\mathbb{Z})^{\otimes 2} \oplus \mathbb{Z} \subseteq \operatorname{Mat}_{s^{2}+1}(\mathbb{Z})$$

where  $E_{ij} = e_i e_j^t$  with  $e_k$  being the kth standard vector. For every matrix in  $\operatorname{Mat}_{s^2+1}(\mathbb{Z})$  of the form

$$Y = \begin{pmatrix} X \otimes X & 0 \\ 0 & a \end{pmatrix}$$

we have

$$\operatorname{tr}(YN) = a + \sum_{i,j=1}^{s} X_{ij}^2.$$

For an instance  $A_1, \ldots, A_d \in Mat_s(\mathbb{Z})$  of Problem 23, define the following d+1 matrices:

$$B_i = \begin{pmatrix} A_i \otimes A_i & 0\\ 0 & 1 \end{pmatrix} \quad \text{for } i = 1, \dots, d$$

and

$$B_{d+1} = \begin{pmatrix} I_s \otimes I_s & 0\\ 0 & -1 \end{pmatrix}$$

where  $I_s$  is the identity matrix of size s.

Let  $n_1, \ldots, n_d \in \mathbb{N}$  be such that

$$A_1^{n_1}\cdots A_d^{n_d}=0$$

Choosing  $n_{d+1} = 1$  we obtain

$$\operatorname{tr}(B_1^{n_1}\cdots B_{d+1}^{n_{d+1}}\cdot N) = -1 + \sum_{i,j=1}^s (A_1^{n_1}\cdots A_d^{n_d})_{ij}^2 = -1 < 0.$$

Conversely, let  $n_1, \ldots, n_{d+1} \in \mathbb{N}$  be such that  $\operatorname{tr}(B_1^{n_1} \cdots B_{d+1}^{n_{d+1}} \cdot N) < 0$ . This is only possible for  $n_{d+1}$  odd and

$$\sum_{i,j=1}^{s} \left( A_1^{n_1} \cdots A_d^{n_d} \right)_{ij}^2 = 0,$$

which implies  $A_1^{n_1} \cdots A_d^{n_d} = 0.$