Convergence of martingales via enriched dagger categories

Paolo Perrone and Ruben Van Belle

University of Oxford

Abstract

We provide a categorical proof of convergence for martingales and backward martingales in mean, using enriched category theory. The enrichment we use is in topological spaces, with their canonical closed monoidal structure, which encodes a version of pointwise convergence.

We work in a topologically enriched dagger category of probability spaces and Markov kernels up to almost sure equality. In this category we can describe conditional expectations exactly as dagger-split idempotent morphisms, and filtrations can be encoded as directed nets of split idempotents, with their canonical partial order structure. As we show, every increasing (or decreasing) net of idempotents tends topologically to its supremum (or infimum).

Random variables on a probability space form contravariant functors into categories of Hilbert and Banach spaces, which we can enrich topologically using the L^p norms. Martingales and backward martingales can be defined in terms of these functors. Since enriched functors preserve convergence of nets, we obtain convergence in the L^p norms. The convergence result for backward martingales indexed by an arbitrary net, in particular, seems to be new.

By changing the functor, one can describe more general notions of conditional expectations and martingales, and if the functor is enriched, one automatically obtains a convergence result. For instance, one can recover the Bochner-based notion of vector-valued conditional expectation, and the convergence of martingales with values in an arbitrary Banach space.

This work seems to be the first application of topologically enriched categories to analysis and probability in the literature. We hope that this enrichment, so often overlooked in the past, will be used in the future to obtain further convergence results categorically.

Contents

1	Intr	roduction	2	
2	Categorical background			
	2.1	Dagger categories	6	
	2.2	Idempotents and their splittings	8	
	2.3	The order of split idempotents	9	
	2.4	Sequential and filtered suprema and infima	12	
3	Ker	nels and random variables, categorically	16	
	3.1	Categories of Markov kernels	16	
	3.2	Functors of random variables	17	
	3.3	Bayesian inversions, conditioning, and dagger structures	21	

4	Idempotent kernels and sub-sigma-algebras	24		
	4.1 Conditional expectation operators	24		
	4.2 The preorder of sub-sigma-algebras	28		
	4.3 Filtrations and martingales	33		
5	Topological enrichment	40		
	5.1 The enrichments of Ban and Hilb	40		
	5.2 The enrichment of Krn and EKrn	42		
	5.3 The subspace of idempotent kernels	45		
6	Martingale convergence	46		
	6.1 Levi properties	47		
	6.2 The idempotent Levi property of Hilb	47		
	6.3 The idempotent Levi property of EKrn and Levy's theorems	50		
7	Vector-valued martingales	52		
	7.1 Bochner integrals and conditional expectations	54		
	7.2 Topological enrichment and convergence	58		
\mathbf{A}	Topologically enriched categories	59		
	A.1 The tensor product	59		
	A.2 The internal hom	61		
	A.3 Categories enriched in Top	62		
R	References			

1 Introduction

This work is part of an effort, started in the last few years, to express and to generalize the main results of probability theory using the language of category theory. Several results of probability theory and related fields have been recovered in this way, and sometimes extended. Among these, the Kolmogorov and Hewitt-Savage zero-one laws [FR20, EP23], de Finetti's theorem [FGP21, MP22], the Carathéodory extension theorem [VB22], the Radon-Nikodym theorem [VB23], and different versions of the ergodic decomposition theorem [MP23, EP23]. The main categorical structures used to achieve these purposes are *probability monads* [Gir82], *Markov categories* [CJ19, Fri20], and particular *dagger categories* [Kar18].

In this work, we turn to another cornerstone result of probability theory, the martingale convergence theorem. It builds on previous work on martingales through the categorical lens of random variable functors [AR18,DDGS18,VB23], and on a categorical treatment of conditional expectations as particular idempotent morphisms [FGL⁺, EP23].

The idea that martingale convergence is a limit or colimit-like property has already appeared in [Koz16], where it is argued that it is like a universal property, but where *certain key properties hold* only up to a null set. In this work we follow that intuition intuition using a category, first defined in [DDGS18], where the morphisms are taken up to almost sure equality.

Probability spaces, kernels, and random variables. Among the most important structures of probability theory are probability spaces and random variables. In this work we give a categorical account of both: probability spaces as a category, and random variables on them as a functor.

In probability theory it is customary to fix a single probability space, the outcome space, and to work only on that space. However, especially in the context of stochastic processes, one considers several sigma-algebras on that space at once, usually forming a filtration. Technically, these sigmaalgebras give rise to different, but related, probability spaces. The categorical description that we use formalizes this intuition: we want probability spaces, with fixed sigma-algebras, to be the objects of a category, and refinements and coarse-grainings of these spaces to be the morphisms. Concretely we work with a category whose objects are probability spaces, and whose morphisms are measure-preserving Markov kernels quotiented under a version of almost sure equality. This category, in the standard Borel case, was first defined in [DDGS18].

Given a probability space (X, \mathcal{A}, p) , one can consider the random variables on it which are measurable for \mathcal{A} and integrable for p. Choosing a different probability space (for example, a different sigma-algebra), one obtains a different set of random variables. Moreover, refinements and coarse-grainings of the probability spaces have an effect on random variables, for example via conditional expectations. This is encoded, category-theoretically, by modelling random variables as a *functor* on the category of probability spaces. This idea was previously developed in [VB23, DDGS18, AR18], and here we give a further generalization of it. For an example of how this works, consider a probability space (X, \mathcal{A}, p) and a sub-sigma-algebra $\mathcal{B} \subseteq \mathcal{A}$. We have arrows between probability spaces as in the diagram below on the left,

$$\begin{array}{ccc} (X, \mathcal{A}, p) & & L^1(X, \mathcal{A}, p) \\ \pi \downarrow \uparrow \pi^+ & & \pi^* \uparrow \downarrow (\pi^+)^* \\ (X, \mathcal{B}, p) & & L^1(X, \mathcal{B}, p) \end{array}$$

where we denote the restriction of p to \mathcal{B} again by p. The map π is the (kernel induced by the) set-theoretical identity $X \to X$, which coarse-grains the sigma-algebra from \mathcal{A} to \mathcal{B} . The kernel π^+ is the Bayesian inverse of π , which is guaranteed to exist, for example, if (X, \mathcal{A}) is a standard Borel space. Random variables are now a contravariant functor. This means that we can assign to each probability space, for example (X, \mathcal{A}, p) , the Banach space $L^1(X, \mathcal{A}, p)$ of integrable random variables on it, again quotiented under almost sure equality. Moreover, we can assign to each morphism, i.e. Markov kernel, a corresponding map between Banach spaces, in the opposite direction, as in the diagram above on the right. For example,

- The map $\pi^* : L^1(X, \mathcal{B}, p) \to L^1(X, \mathcal{A}, p)$, induced by the set-theoretical identity, is basically an inclusion map, saying that all \mathcal{B} -measurable random variables are \mathcal{A} -measurable as well;
- The map $(\pi^+)^* : L^1(X, \mathcal{A}, p) \to L^1(X, \mathcal{B}, p)$, induced by the Bayesian inverse, takes an \mathcal{A} -measurable random variable and gives its conditional expectation given \mathcal{B} .

In other words, the relations between the different probability spaces (for example, in a filtration) and their effects on random variables are captured categorically by means of a category and a functor. For more details, see Section 3.

This approach also lets us consider more general notions of random variables, for example vector-valued ones, integrable in the Bochner sense. These will simply form a different functor on our category. For more details, see Section 7.

Filtrations and martingales. A filtration on a probability space (X, \mathcal{A}, p) is a sequence, or a net, of sub-sigma-algebras $\mathcal{B}_i \subseteq \mathcal{A}$ which are increasingly finer, that is, $\mathcal{B}_i \subseteq \mathcal{B}_j$ whenever $i \leq j$. Usually the filtration is indexed by a total order, which can be interpreted as time, so that as time progresses, our knowledge about the system progresses as well: we are able to make finer and finer distinctions, and to assess the probability of more and more possible events. One can then form the *join* sigma algebra $\mathcal{B}_{\infty} = \bigvee_i \mathcal{B}_i$, which can be interpreted as encoding all the knowledge that one can possibly learn from the process, if one had infinite time. From the point of view of category theory, this sigma-algebra can be obtained as a particular directed limit or colimit of subobjects.

A martingale on the filtration, following this interpretation, is a collection of random variables f_i which, intuitively, follow the refinement of the filtration. More in detail, each f_i needs to be measurable for the sigma-algebra \mathcal{B}_i , i.e. it respects the state of knowledge that we have at "time" i. Moreover, for each $i \leq j$, we have that f_i is the conditional expectation of f_j . In other words, f_i is a "coarse graining" of f_j which "forgets" or "averages over" all the distinctions that we would be able to make at time j, but that we cannot yet make at time i. Dually, we can see f_j as a "refinement" of f_i , which incorporates the additional "knowledge" that \mathcal{B}_j has over \mathcal{B}_i .

The martingale convergence theorem says that, under some conditions,

- The f_i admits a common universal refinement f, measurable for the join sigma algebra \mathcal{B}_{∞} , and moreover,
- The f_i tend to f topologically.

In other words, as our knowledge increases, our refinements become better and better "approximations of the real thing". This "approximation" part is achieved, categorically, by means of a topological enrichment.

Topological enrichment. It is sometimes said that algebra concerns itself with equations and analysis concerns itself with inequalities. While this is of course a simplification, it is true that in ordinary category theory it is not obvious how to express approximations and convergence, and the latter are extremely important in analysis and in probability theory. Luckily, however, category theory has *enriched* versions, enhancements of the notion of category where the sets of arrows are not just sets, but come equipped with additional structures, compatibly with composition. In particular, one can equip these sets with a metric or with a topology. This allows us to talk about convergence of measures and random variables categorically, and in a certain way, it adds approximations and inequalities to an otherwise purely algebraic, equational environment.

In [VB23], a metric enrichment on a category of probability spaces and measure-preserving functions was used to study certain properties of martingales. Also, in [Per24], a metric enrichment on a category of kernels was used to recover classical concepts of information theory, such as entropy and data processing inequalities. In this work we bring the two ideas together, using enriched categories to study martingales, but using categories of kernels. Moreover, instead of metrics we endow our sets of arrows with more general topologies. The reason is that the notion of convergence, and this is too strong for our purposes. We are interested in a version of *uniform* convergence, and this is achieved by using a topological enrichment, where the sets of arrows have a topology which in general not metrizable.¹ In the literature, this enrichment has not been used very often.

¹The topologies we consider, however, can be induced by *families* of metrics, and so we expect our categories to be enriched not only in topological spaces, but also in uniform spaces [Isb64]. We will not pursue this idea in the present work.

In particular, it seems that it was never used to express, in terms of category theory, convergence results in probability or analysis. In that regard, this work might be the first one of its kind, where topological convergence and category theory are used together in this way.

Let's see how this works. First of all, one actually *can* express certain ideas of approximation and convergence in ordinary categories: via particular limits and colimits (hence the name *limit*). For example, the diagram in the category of sets formed by finite sets and inclusion maps

 $\{0\} \longleftrightarrow \{0,1\} \longleftrightarrow \{0,1,2\} \longleftrightarrow \{0,1,2,3\} \longleftrightarrow \ldots$

has as colimit the set of natural numbers \mathbb{N} . Similarly, and dually, one can express an infinite cartesian product as a limit of its finite factors. (This is particularly important in probability theory, as it is related to Kolmogorov's extension theorem, see [FR20].)

The martingale convergence theorem, categorically, can be expressed as the fact that *if we* express an object as a categorical limit (a probability space as a limit of increasingly finer sigmaalgebras), we also have a topological convergence associated with it (the conditional expectations of the random variables converge to a common refinement).

This correspondence between categorical limits and topological ones is a phenomenon that, in different ways, has been noticed before, for example in the category of Hilbert spaces [DMH24], as well as for martingales [Koz16]. In this work, seemingly for the first time in the literature, we make this correspondence mathematically precise, by means of what we call the *idempotent Levi property* of a topologically enriched category. The details are explained in Section 6.

Our results. The main result of this paper is a version of the convergence theorem in mean for martingales and for backward martingales, which we prove for filtrations indexed by directed nets of arbitrary cardinality (Corollaries 6.12 and 6.13), and for random variables taking values in an arbitrary Banach space (Corollary 7.8).

Along the way, there are a number of additional results proven in this paper, which can be of independent interest. First of all, we prove that random variables, including vector-valued ones, are functorial on probability spaces (Propositions 3.6 and 7.5). We also prove that sub-sigma-algebras of a standard Borel space, up to null sets, are in bijection with almost surely idempotent kernels by means of taking conditional expectations and invariant sigma-algebras, and that the order of idempotents coincide with the (almost surely) inclusion order of sub-sigma-algebras (Theorems 4.6 and 4.18). We then give a characterization of martingales in terms of idempotents (Proposition 4.28), and in terms of limits and colimits of Banach and Hilbert spaces (Corollaries 4.29 and 4.30). The last step to prove martingale convergence is what we call the *Levi property* of a topologically enriched category, which says that directed nets of idempotents converge topologically. We prove that such a property holds in the category of Hilbert spaces (Propositions 6.4 and 6.6), and in our category of probability spaces (Theorem 6.9).

As it is well known, there are results on convergence of martingales not only in mean, but also almost surely, at least for the case of sequential filtrations. In this work we focus on convergence in mean, which implies convergence in probability, and leave the case of almost sure convergence to future work. Since almost sure convergence does not arise from a topology, we expect that a categorical treatment of such a convergence will require much more refined enrichments.

Outline.

- In Section 2 we give an overview of the main categorical concepts we need in this paper, dagger categories, and idempotents and their splittings. We show how, in categories of vector spaces, idempotents are projections onto subspaces. We conclude the section by looking at filtrations of subspaces and their limits.
- In Section 3 we turn to the main categories of interest for this work, of probability spaces and Markov kernels between them up to almost sure equality. We also define functors of random variables, and show how conditioning equips our categories with a dagger structure.
- In Section 4 we join the structures introduced in the previous two sections, studying split idempotents in categories of kernels. We show that they are tightly related to sub-sigma-algebras and to conditional expectations of random variables. In particular they allow to give categorical descriptions of filtrations and of martingales.
- In Section 5 we provide our categories of kernels and of vector spaces with a topological enrichment, i.e. a way to talk about convergence of morphisms in a "pointwise-like" sense. We also show that the random variables functors are enriched, meaning that the functorial assignment on morphisms is continuous.
- In Section 6 we finally show the main result of this work, namely convergence in mean martingales and backward martingales, for arbitrary nets, using categorical arguments. The main idea is what we call the *Levi property*, which roughly says that convergence in the order implies topological convergence. The convergence of backward martingales indexed by arbitrary nets seems to be a new result.
- In Section 7 we show that our convergence result can be extended to much more general notions of random variables, provided they form an enriched functor. We give the example of martingales with values in an arbitrary Banach space, using a Bochner-based approach to vector valued conditional expectations.
- Finally, in Appendix A we give some background on topologically enriched categories, and on the closed monoidal structure of the enriching category Top.

Acknowledgements. We want to thank Sam Staton and the members of his group for the enlightening discussions, and for providing a greatly supportive and inspiring research environment. We also want to thank Matthew Di Meglio for the helpful pointers on the category of Hilbert spaces.

Research for both authors is funded by Sam Staton's ERC grant "BLaSt – a Better Language for Statistics".

2 Categorical background

2.1 Dagger categories

A dagger category can be interpreted as an undirected version of a category, similarly to directed and undirected graphs. Just as for undirected graphs, in a dagger category if two objects are connected by a morphism $X \to Y$, then they are also connected by a morphism $Y \to X$, and we can think of these two arrows as a single entity which we can "walk either way". More specifically, to each morphism $f: X \to Y$ we assign a morphism $f^+: Y \to X$, not necessarily invertible, in a way which preserves identity and composition. Here is the precise definition.

Definition 2.1. A dagger category is a category C together with a functor $(-)^+$: $C^{op} \to C$ which is

- The identity on objects, i.e. $X^+ = X$ for all X of C;
- Involutive, i.e. for all morphisms, $(f^+)^+ = f$.

Very often, dagger structures are used to encode "Euclidean" geometric structures, or notions of "orthogonality", as the following examples show.

Example 2.2. The category Euc has as objects Euclidean spaces (finite-dimensional real vector spaces with an inner product), and as morphisms linear maps, or equivalently matrices. The transposition of matrices gives a dagger, as $(AB)^t = A^t B^t$. (Note that the inner product is necessary in order to have a coordinate-independent transposition.)

Example 2.3. The category Hilb of real Hilbert spaces and bounded linear maps can be given a dagger structure via the *adjoint*. Given Hilbert spaces X and Y and a bounded linear map $f: X \to Y$, its adjoint is the unique bounded linear map $f^+: Y \to X$ satisfying

$$\langle f(x), y \rangle = \langle x, f^+(y) \rangle$$

for all $x \in X$ and $y \in Y$. The brackets in the equation above are, in order, the inner product of Y, and the one of X.

Definition 2.4. A morphism $f: X \to Y$ in a dagger category is called

- a dagger monomorphism or isometry if $f^+ \circ f = id_X$;
- a dagger epimorphism or co-isometry if $f \circ f^+ = id_Y$;
- a dagger isomorphism or unitary if it is invertible, and $f^{-1} = f^+$;
- self-dual or self-adjoint if X = Y and $f^+ = f$.

For brevity we will just say "dagger mono" for "dagger monomorphism", as well as "dagger monic", et cetera. Note that a dagger mono is in particular a split mono, a dagger epi is in particular a split epi, and a dagger iso is in particular an isomorphism. Equivalently, a dagger isomorphism is a morphism which is both dagger monic and dagger epic.

Example 2.5. In Euc,

• The dagger monomorphisms are exactly the linear isometries, or linear isometric embeddings, namely the matrices which preserve the Euclidean norms and inner products of vectors:

$$\langle f(x), f(x') \rangle = \langle x, f^+ f(x') \rangle = \langle x, x' \rangle.$$

Dagger epimorphisms are their transposes;

• The dagger isomorphisms are exactly the orthogonal matrices;

• The self-dual morphisms are exactly the symmetric matrices.

Example 2.6. In Hilb,

- The dagger monomorphisms are exactly the linear isometries (and their duals are sometimes called *co-isometries* of Hilbert spaces);
- The dagger isomorphisms are exactly the unitary maps;
- The self-dual morphisms are exactly the self-adjoint operators.

Dagger monos and dagger epis are respectively closed under composition. In what follows we will write

- C_{dm} for the wide subcategory of dagger monos of C;
- C_{de} for the wide subcategory of dagger epis of C.

2.2 Idempotents and their splittings

Let's now look at idempotents. We first start with idempotents in general, and then turn to the dagger setting. An **idempotent** morphism in a category C is an endomorphism $e: X \to X$ such that $e \circ e = e$.

Definition 2.7. A splitting of an idempotent $e : X \to X$ consists of an object A and morphisms $\iota : A \to X$ and $\pi : X \to A$ such that $\pi \circ \iota = id_A$ and $\iota \circ \pi = e$.

Note that this implies that ι is split monic, and π is split epic.

Very often, split idempotents encode the idea of "projections onto a subspace", as in the following examples.

Example 2.8. In Euc, all idempotents split. Given an idempotent linear map e on a Euclidean space X, denote by $A \subseteq X$ its image, and by $\iota : A \to X$ the inclusion. By definition of image, for all $x \in X$ we have that $e(x) \in A$, so e specifies a linear map $\pi : A \to A$ by $x \mapsto e(x)$. This way, $e(x) = \iota(\pi(x))$, and for all $y \in A$, for some x, y = e(x), and so by idempotency, $\pi(\iota(y)) = e(e(x)) = e(x) = y$. Therefore $\pi \circ \iota = \mathrm{id}_A$ and $\iota \circ \pi = e$. In other words, we can see the idempotent e as a projection onto a the subspace A.

Example 2.9. In Ban of Banach spaces and bounded linear maps, all idempotents split as well, and the intuition is similar. Let e be an idempotent bounded linear map on a Banach space X. Let $A \subseteq X$ be its image, and denote by $\iota : A \to X$ be the inclusion map. Note first of all that A is a *closed* subspace: suppose that for a sequence (x_n) in X we have that $e(x_n) \to y$. Then since e is continuous and idempotent we have that

$$e(y) = \lim_{n \to \infty} e(e(x_n)) = \lim_{n \to \infty} e(x_n) = y.$$

So $y \in A$, which means that A is closed, and so it is itself a Banach space. Just as in the example above, e defines a map $\pi : X \to A$, and we have that $\pi \circ \iota = id_A$ and $\iota \circ \pi = e$. Once again, we can view the idempotent e as a projection onto the closed subspace A. Note that not every closed subspace of a Banach space admits a (bounded) idempotent operator projecting onto it, and even when it exists, it may not be unique. In Hilb, the situation is completely analogous, with the additional condition that *every closed* subspace admits an idempotent. Uniqueness still fails, but there is a canonical choice, see Example 2.11.

Equivalently, one can construct A as either the set of fixed points of e, or as the quotient set of fibers of e. Categorically, A can be constructed as either the equalizer or the coequalizer of the parallel pair e, id : $X \to X$, and it is preserved by every functor [BD86, Proposition 1].

Let's now turn to the dagger case.

Definition 2.10. Let C be a dagger category.

- A dagger idempotent or projector is a self-dual idempotent morphism. That is, a morphism e : X → X such that e ∘ e = e = e⁺;
- A dagger splitting of an idempotent $e: X \to X$ is a splitting (A, ι, π) such that $\iota = \pi^+$.

Note that in the definition above, we necessarily have that ι is dagger monic and that π is dagger epic. Note also that if an idempotent has a dagger splitting, then it is a dagger idempotent:

$$e^+ = (\iota \circ \pi)^+ = \pi^+ \circ \iota^+ = \iota \circ \pi = e.$$

We have seen that split idempotents often encode "projections onto subspaces", and that dagger structures often encode notions of "orthogonality". Following this intuition, dagger idempotents often encode *orthogonal projections* onto subspaces:

Example 2.11. In Euc as well as in Hilb, consider a dagger idempotent $e: X \to X$ and its splitting (A, ι, π) as in Examples 2.8 and 2.9. We have that for every $x \in X$ and $y \in A$,

$$egin{aligned} &\langle x-e(x),y
angle &=\langle x,y
angle -\langle e(x),y
angle \ &=\langle x,y
angle -\langle x,e(y)
angle \ &=\langle x,y
angle -\langle x,y
angle \ &=0. \end{aligned}$$

Therefore e is an orthogonal projector onto A. To see that our splitting is dagger, notice that for all $x \in X$ and $y \in A$,

$$\langle \pi(x), y \rangle = \langle e(x), y \rangle = \langle x, e(y) \rangle = \langle x, y \rangle = \langle x, \iota(y) \rangle.$$

Therefore $\iota = \pi^+$. Conversely, by taking the adjoint of the inclusion map, every closed subspace admits a (unique) orthogonal projector.

2.3 The order of split idempotents

Subspaces of a vector space form naturally a partial order under inclusion. The same, dually, can be said about quotients. More generally, subobjects and quotient objects form partial orders. Retracts, or split idempotents, can be seen in this light, and inherit a partial order with interesting properties.

Proposition 2.12. Let e_1 and e_2 be idempotents $X \to X$ in a category, with splittings (A_1, π_1, ι_1) and (A_2, π_2, ι_2) respectively.



The following conditions are equivalent.

- (i) $e_1 \circ e_2 = e_2 \circ e_1 = e_1;$
- (*ii*) $e_2 \circ \iota_1 = \iota_1$ and $\pi_1 \circ e_2 = \pi_1$;
- (iii) There exists $f: A_1 \to A_2$ and $g: A_2 \to A_1$ such that the following diagrams commute.



Moreover, in the situation above, f and g are unique, and $g \circ f = id_{A_1}$.

If in addition we are in a dagger category, and (ι_1, π_1) and (ι_2, π_2) are dagger splittings, then $f = g^+$.

Proof. (i) \Rightarrow (ii):

 $e_2 \circ \iota_1 = e_2 \circ e_1 \circ \iota_1 = e_1 \circ \iota_1 = \iota_1;$ $\pi_1 \circ e_2 = \pi_1 \circ e_1 \circ e_2 = \pi_1 \circ e_1 = \pi_1.$

 $(ii) \Rightarrow (iii)$: Notice that $\iota_2 \circ f = \iota_1$ implies necessarily that $f = \pi_2 \circ \iota_1$, so that f is unique. This way,

$$\iota_2 \circ f = \iota_2 \circ \pi_2 \circ \iota_1 = e_2 \circ \iota_1 = \iota_1.$$

Similarly, if we want that $g \circ \pi_2 = \pi_1$, necessarily $g = \pi_1 \circ \iota_2$. Now

$$g \circ \pi_2 = \pi_1 \circ \iota_2 \circ \pi_2 = \pi_1 \circ e_2 = \pi_1.$$

To show that g is a retraction,

$$g \circ f = \pi_1 \circ \iota_2 \circ \pi_2 \circ \iota_1 = \pi_1 \circ e_2 \circ \iota_1 = \pi_1 \circ \iota_1 = \mathrm{id}_{A_1}.$$

 $\begin{aligned} (iii) \Rightarrow (i): \\ e_1 \circ e_2 \ = \ \iota_1 \circ \pi_1 \circ \iota_2 \circ \pi_2 \ = \ \iota_1 \circ g \circ \pi_2 \ = \ \iota_1 \circ \pi_1 \ = \ e_1. \\ e_2 \circ e_1 \ = \ \iota_2 \circ \pi_2 \circ \iota_1 \circ \pi_1 \ = \ \iota_2 \circ f \circ \pi_1 \ = \ \iota_1 \circ \pi_1 \ = \ e_1. \end{aligned}$

If we are in a dagger category and $\pi_1 = \iota_1^+$ and $\pi_2 = \iota_2^+$, then

$$g^+ = (\pi_1 \circ \iota_2)^+ = \iota_2^+ \circ \pi_1^+ = \pi_2 \circ \iota_1 = f.$$

Definition 2.13. In the hypotheses of the previous proposition, we say that $e_1 \leq e_2$ if any (hence all) the conditions are satisfied.

Remark 2.14. In the dagger case it is even simpler to check that $e_1 \leq e_2$: by taking the dagger,

- $e_1 \circ e_2 = e_1$ if and only if $e_2 \circ e_1 = e_1$;
- $e_2 \circ \iota_1 = \iota_1$ if and only if $\pi_1 \circ e_2 = \pi_1$;
- f as in Proposition 2.12 exists if and only if g exists.

Example 2.15. In Euc, Ban and Hilb, in the hypotheses of the previous proposition, we have that $e_1 \leq e_2$ if and only if, as subspaces, $A_1 \subseteq A_2$. The equivalent conditions read:

- (i) Projecting first onto A_2 and then onto A_1 is the same as just projecting onto A_1 directly;
- (ii) Projecting an element of A_1 onto A_2 leaves it unchanged;
- (iii) The projection $X \to A_1$ and the inclusion $A_1 \to X$ both factor through A_2 .

Proposition 2.16. The relation \leq on the set of split idempotents on X is a partial order.

Proof. For reflexivity, notice that $e \circ e = e$ by idempotency. For transitivity, suppose that $e_1 \circ e_2 = e_1$ and $e_2 \circ e_3 = e_2$. Then

$$e_1 \circ e_3 = e_1 \circ e_2 \circ e_3 = e_1 \circ e_2 = e_1.$$

For antisymmetry, notice that $e_1 = e_1 \circ e_2 = e_2$ implies $e_1 = e_2$.

We can equivalently define the order of idempotents as the following category.

Definition 2.17. Let X be an object of a category C. The category SI(X) has:

- As objects, objects A of C together with a split idempotent $(\iota : A \to X, \pi : X \to A);$
- As morphisms $(A_1, \iota_1, \pi_1) \to (A_2, \iota_2, \pi_2)$, pairs of morphisms $f_{12}: A_1 \to A_2$ and $g_{21}: A_2 \to A_1$ and making the top and bottom triangles in the following diagram commute.



This category is a preorder, equivalent to the poset of split idempotents on X. In the dagger case, we can define things similarly.

Definition 2.18. Let X be an object of a dagger category C. The category DSI(X) has:

• As objects, objects A of C together with a dagger-split idempotent $(\iota : A \to X, \pi : X \to A);$

As morphisms (A₁, ι₁, π₁) → (A₂, ι₂, π₂), pairs of morphisms f₁₂ : A₁ → A₂ and making the top and bottom triangles in the following diagram commute.



In the dagger case, thanks to the self-duality, can have two equivalent descriptions of the DSI(X). The slice category C_{dm}/X has:

- As objects, objects A of C together with a dagger monomorphism $\iota : A \to X$;
- As morphisms, dagger monomorphisms $f_{12}: A_1 \to A_2$ making the following triangle commute.

$$\begin{array}{c} A_1 \xrightarrow{f_{12}} A_2 \\ \downarrow & \swarrow \\ \iota_1 & \swarrow \\ \chi \\ X \end{array}$$

Dually, the coslice category X/C_{de} has:

- As objects, objects Y of C together with a dagger epimorphism $\pi: X \to Y$;
- As morphisms, dagger epimorphisms $g_{21}: Y_2 \to Y_1$ making the following triangle commute.



The category C_{dm}/X is a preorder, equivalent to the poset of dagger-split idempotents on X, and the category X/C_{de} is equivalent to the *opposite* of the poset of dagger-split idempotents on X.

2.4 Sequential and filtered suprema and infima

Consider an ascending chain $e_1 \leq e_2 \leq \ldots$ of idempotents on X. Equivalently it is an inductive sequence in SI(X), as follows:



The sequential colimit A_{∞} of the A_i in SI(X), if it exists, is equivalently the supremum in the order of idempotents. The same thing is true for increasing (directed) nets instead of sequences.

Similarly, consider a descending chain $e_1 \ge e_2 \ge \ldots$ of idempotents. This is equivalently a coinductive sequence in SI(X), as follows:



The sequential limit A_{∞} of the A_i in SI(X), if it exists, is equivalently the infimum in the order of idempotents. The same is true for decreasing nets instead of sequences.

Similar remarks can be made for dagger-split idempotents.

Example 2.19. In Euc and Hilb, increasing and decreasing sequences of dagger idempotents are increasing and decreasing sequences of (closed) subspaces. Their suprema are given by (the closure of) their union, and their infima are given by their intersection. The same is true for nets.

Example 2.20. In Ban, not every closed subspace gives rise to an idempotent, and when it does it may not be unique, so we need a little care. What we can say is that

- (i) Given an increasing net of idempotents e_λ : X → X projecting onto closed subspaces A_λ ⊆ X, denote by A_∞ the closure of their union. Then if the supremum of the e_λ in the idempotent order exists, it projects onto A_∞.
- (ii) Given a decreasing net of idempotents $e_{\lambda} : X \to X$ projecting onto closed subspaces $A_{\lambda} \subseteq X$, denote by A_{∞} their intersection. Then if the infimum of the e_{λ} in the idempotent order exists, it projects onto A_{∞} .

Moreover, in the category $\operatorname{Ban}_{\leq 1}$ of Banach spaces and *1-Lipschitz* linear maps (not just bounded),

- nets of increasing subspaces have the closure of their union as colimit of the inclusion maps;
- nets of decreasing subspaces have their intersection as limit of the inclusion maps.

This is shown in the following propositions. The same statements will also hold in the subcategory $\text{Hilb}_{\leq 1}$ of Hilbert spaces and 1-Lipschitz linear maps, and we can interpret them in terms of the category DSI.

Proposition 2.21. Let X be a Banach space. Let $(A_{\lambda})_{\lambda \in \Lambda}$ be an increasing net of closed subspaces of X, i.e. such that for all all $\lambda \leq \mu$, $A_{\lambda} \subseteq A_{\mu}$. Let

$$A_{\infty} = \operatorname{cl}\left(\bigcup_{\lambda} A_{\lambda}\right).$$

For $\lambda \leq \mu$, denote by $\iota_{\lambda,\mu} : A_{\lambda} \to A_{\mu}$ the inclusion. Then A_{∞} is the (filtered) colimit in $\operatorname{Ban}_{\leq 1}$ of the diagram formed by the $\iota_{\lambda,\mu}$.

Proof. Let B be a Banach space, and consider a cone of 1-Lipschitz linear maps $c_{\lambda} : A_{\lambda} \to B$ making the following diagram commute for each $\mu \geq \lambda$.

 $\begin{array}{c|c}
A_{\lambda} & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\
 & & \\$

We have to show that there is a unique map $c: A_{\infty} \to B$ making the following diagram commute.



Now define first the map

$$c':\bigcup_{\lambda\in\Lambda}A_\lambda\to B$$

by $c'(a_{\lambda}) = c_{\lambda}(a_{\lambda})$ for all $a_{\lambda} \in A_{\lambda}$. Note that since (2.1) commutes, this map does not depend on the choice of λ . Since all the c_{λ} are 1-Lipschitz, so is c', and so by density c' admits a unique (linear) extension to A_{∞} .

Proposition 2.22. Let X be a Banach space. Let $(A_{\lambda})_{\lambda \in \Lambda}$ be a decreasing net of closed subspaces of X, i.e. such that for all all $\lambda \leq \mu$, $A_{\lambda} \supseteq A_{\mu}$. Let

$$A_{\infty} = \bigcap_{\lambda} A_{\lambda}.$$

For $\lambda \leq \mu$, denote by $\iota_{\mu,\lambda} : A_{\mu} \to A_{\lambda}$ the inclusion. Then A_{∞} is the (cofiltered) limit in Ban and $\operatorname{Ban}_{\leq 1}$ of the diagram formed by the $\iota_{\mu,\lambda}$.

Proof. Let B be a Banach space, and consider a cone consisting of bounded linear maps $c_{\lambda} : B \to A_{\lambda}$ (for each λ) making the following diagram commute for each $\lambda \leq \mu$.



The commutativity of the diagram above says that for every $b \in B$, and for every $\mu \geq \lambda$, we have that $c_{\lambda}(b) \in A_{\mu}$. Since any $\lambda, \lambda' \in \Lambda$ admit an upper bound μ , for every $\lambda, \lambda' \in \Lambda, c_{\lambda}(b) \in A_{\mu} \subseteq A_{\lambda'}$.

Therefore, for every λ , $c_{\lambda}(b) \in \bigcap_{\lambda'} A_{\lambda'} = A_{\infty}$. In other words, all the c_{λ} factor uniquely through a map $c: B \to A_{\infty}$ as follows,



where ι_{λ} and ι_{μ} are the inclusions. That is, A_{∞} is the limit of the cofiltered diagram.

Since these statements also hold in $\text{Hilb}_{\leq 1}$, which is a dagger category, by dualizing we get the following statements, involving the projections instead of the inclusions:

Corollary 2.23. Let X be a Hilbert space. Let $(A_{\lambda})_{\lambda \in \Lambda}$ be an increasing net of closed subspaces of X, i.e. such that for all all $\lambda \leq \mu$, $A_{\lambda} \subseteq A_{\mu}$. Let

$$A_{\infty} = \operatorname{cl}\left(\bigcup_{\lambda} A_{\lambda}\right).$$

For $\lambda \leq \mu$, denote by $\pi_{\mu,\lambda} : A_{\mu} \to A_{\lambda}$ the orthogonal projection. Then A_{∞} is the (cofiltered) limit in Hilb_{<1} of the diagram formed by the $\pi_{\mu,\lambda}$.

Corollary 2.24. Let X be a Hilbert space. Let $(A_{\lambda})_{\lambda \in \Lambda}$ be a decreasing net of closed subspaces of X, i.e. such that for all all $\lambda \leq \mu$, $A_{\lambda} \supseteq A_{\mu}$. Let

$$A_{\infty} = \bigcap_{\lambda} A_{\lambda}.$$

For $\lambda \leq \mu$, denote by $\pi_{\lambda,\mu} : A_{\lambda} \to A_{\mu}$ the projection. Then A_{∞} is the (filtered) colimit in $\operatorname{Hilb}_{\leq 1}$ of the diagram formed by the $\pi_{\lambda,\mu}$.

Note that the same statements does not hold more in general for Ban. A counterexample for the analogue of Corollary 2.23 in the Banach case will be given in Example 4.31. A counterexample for the analogue for Corollary 2.24 in the Banach case will be given in Example 6.8.

Let's now interpret this statement, for Hilb, in terms of the category DSI. Note first of all that the dagger monic and the dagger epic morphisms of Hilb are 1-Lipschitz. Now, it is well known that for every category C and object X, the forgetful functor

$$\begin{array}{ccc} X/\mathcal{C} & \longrightarrow & C \\ (Y, X \xrightarrow{f} Y) & \longmapsto & Y \end{array}$$

creates all limits and all connected colimits which exist in C [Rie16, Proposition 3.3.8]. Similarly, the forgetful functor $C/X \rightarrow C$ creates all colimits and all connected limits which exist in C. Setting now $C = Hilb_{dm}$, we have that the forgetful functor $DSI(X) \cong X/Hilb_{dm} \rightarrow Hilb_{dm}$ creates all existing connected limits and colimits, in particular filtered colimits and cofiltered limits. This, together with the previous statements, gives an alternative proof of the fact that intersections and closures of unions are the directed infima and suprema in the order of dagger idempotents of Hilb.

3 Kernels and random variables, categorically

We now turn our attention to probability theory. We define a (dagger) category of probability spaces, with Markov kernels as morphisms, and functors encoding random variables.

3.1 Categories of Markov kernels

We now present categories, Krn, GKrn and EKrn, of probability spaces and Markov kernels up to almost sure equality. Krn was first defined in [DDGS18]. These categories should not be confused with the category, often denoted by Stoch, of *measurable* spaces and Markov kernels, without quotienting under almost sure equality. The two constructions are however related, as shown in the abstract setting of [Fri20, Section 13]. In [EP23] (where GKrn is called PS(Stoch) and GKrn is called PS(BorelStoch)) some of the properties of these categories were studied, which are quite relevant for the present work.

Definition 3.1. Let (X, \mathcal{A}) and (Y, \mathcal{B}) be measurable spaces. A Markov kernel $k : X \to Y$ is an assignment

$$\begin{array}{ccc} X \times \mathcal{B} & \longrightarrow & [0,1] \\ (x,B) & \longmapsto & k(B|x) \end{array}$$

which for every B is measurable as a function of x, and for every x is a probability measure in B. Given Markov kernels $k: X \to Y$ and $\ell: Y \to Z$ we define the Markov kernel $\ell \circ k: X \to Z$ as

follows:

$$(\ell \circ k)(C|x) \coloneqq \int_Y \ell(C|y) \, k(dy|x).$$

Given probability spaces (X, \mathcal{A}, p) and (Y, \mathcal{B}, q) , a Markov kernel $k : X \to Y$ is called **measure**preserving if and only if for every $B \in \mathcal{B}$,

$$\int_X k(B|x) \, p(dx) = q(B).$$

Given probability spaces (X, \mathcal{A}, p) and (Y, \mathcal{B}, q) , two Markov kernels $k, h : X \to Y$ are *p*-almost surely equal if and only if for every measurable $B \in \mathcal{B}$, for *p*-almost all $x \in X$,

$$k(B|x) = h(B|x).$$

Definition 3.2. The category GKrn has

- As objects, probability spaces;
- As morphisms, measure-preserving Markov kernels, quotiented under almost-sure equality.

The "G" in "GKrn" stands for "general", as we also consider the following subcategories:

Definition 3.3. The category Krn is the full subcategory of GKrn whose objects are probability spaces (X, \mathcal{A}, p) , where the measurable space (X, \mathcal{A}) is standard Borel;

Definition 3.4. We call a probability space **essentially standard Borel** if and only if it is isomorphic in GKrn to a standard Borel space. The category EKrn is the full subcategory of GKrn whose objects are essentially standard Borel.

The inclusion functor $\text{Krn} \rightarrow \text{EKrn}$ is an equivalence of categories (but not an isomorphism, since it's only essentially surjective on objects).

In Krn (and EKrn, as we will see), regular conditionals always exist. Therefore, for those categories, we can equivalently take as morphisms $(X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$ probability measures on $X \times Y$ with marginals p and q, i.e. the so-called *couplings* or *transport plans*. Given a coupling c, we obtain a Markov kernel $X \to Y$ (unique up to p-almost-sure equality) by conditioning. Conversely, given a Markov kernel $k : X \to Y$, we get a coupling by defining

$$c(A \times B) = \int_A k(B|x) \ p(dx).$$

It was shown in [EP23, Corollary 3.18] that given a standard Borel probability space (X, \mathcal{A}, p) and any sub-sigma-algebra $\mathcal{B} \subseteq \mathcal{A}$, the probability space (X, \mathcal{B}, p) (where we denote the restriction of p to \mathcal{B} again by p) is essentially standard Borel.

Every measurable function defines a particular "deterministic" kernel as follows. (We will define determinism precisely in Definition 3.16.)

Definition 3.5. Let $f : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, p)$ be a measurable, measure-preserving function (i.e. for all $B \in \mathcal{B}$, $f^{-1}(B) \in \mathcal{A}$ and $p(f^{-1}(B)) = q(B)$). The function f defines a kernel $(X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$ as follows.

$$\delta_f(B|x) = 1_B(x) = \begin{cases} 1 & f(x) \in B; \\ 0 & f(x) \notin B. \end{cases}$$

If we denote by Prob the category of probability spaces and measure-preserving maps, we have an identity-on-objects functor δ : Prob \rightarrow GKrn.

In particular, given a probability space (X, \mathcal{A}, p) and a sub-sigma-algebra $\mathcal{B} \subseteq \mathcal{A}$, the settheoretic identity defines a kernel $\delta_{id} : (X, \mathcal{A}, p) \to (X, \mathcal{B}, p)$ given by

$$\delta_{\rm id}(B|x) = 1_B(x) \tag{3.1}$$

for all $B \in \mathcal{B}$ and $x \in X$. Note that since $\mathcal{B} \subseteq \mathcal{A}$, this assignment is \mathcal{B} -measurable.

3.2 Functors of random variables

Random variables, up to almost sure equality, are functorial on EKrn. Covariant functors of random variables were introduced in [AR18] and independently [DDGS18, Section 4] and [VB23]. Here we use a contravariant version, first defined in [DDGS18, Section 4] in the Borel case, which can be turned covariant by means of the dagger structure of EKrn.

As usual, a **random variable** (**RV**) on a probability space (X, \mathcal{A}, p) is an \mathcal{A} -measurable function $f : X \to \mathbb{R}$. We call f

• integrable if and only if

$$\int_X |f(x)| \, p(dx) < \infty;$$

• *n*-integrable if and only if

$$\int_X |f(x)|^n \, p(dx) < \infty;$$

• almost surely bounded if and only if there exists $A \in \mathcal{A}$ of *p*-measure one such that *f* restricted to *A* is bounded.

Unless otherwise stated, we will not consider non-integrable RVs in the present work.

Again as usual, we say that two RVs f and g on (X, \mathcal{A}, p) are **almost surely equal** if there exists $A \in \mathcal{A}$ of p-measure one such that the restrictions of f and g to A are equal. We denote by

- $L^1(X, \mathcal{A}, p)$ the set of integrable RVs on (X, \mathcal{A}, p) quotiented under almost sure equality;
- $L^n(X, \mathcal{A}, p)$ the set of *n*-integrable RVs on (X, \mathcal{A}, p) quotiented under almost sure equality;²
- $L^{\infty}(X, \mathcal{A}, p)$ the set of almost surely bounded RVs on (X, \mathcal{A}, p) quotiented under almost sure equality.

Since p is a probability measure, every almost surely bounded RV on (X, \mathcal{A}, p) is n-integrable for every n. Also, for $m \ge n \ge 1$, every m-integrable function is n-integrable: for $x \ge 0$, the function $x \mapsto x^{m/n}$ is convex, and so by Jensen's inequality,

$$\left(\int_X |f(x)|^n p(dx)\right)^{m/n} \leq \int_X |f(x)|^m p(dx),$$

so that

$$\|f\|_{L^n} \le \|f\|_{L^m}. \tag{3.2}$$

In particular, if the right-hand side is finite, so is the left-hand side. So, for each probability space, $L^{\infty} \subseteq \cdots \subseteq L^n \subseteq L^{n-1} \subseteq \cdots \subseteq L^1$.

The spaces L^n of random variables form contravariant functors on GKrn. Denote by Ban the category of Banach spaces and bounded linear maps. Equip $L^n(X, \mathcal{A}, p)$ and $L^{\infty}(X, \mathcal{A}, p)$ with their usual norms,

$$||f||_{L^n} \coloneqq \left(\int_X |f(x)|^n p(dx)\right)^{1/n}$$
 and $||f||_{L^\infty} \coloneqq \operatorname{ess\,sup} |f|.$

We have functors $L^n : \operatorname{GKrn}^{\operatorname{op}} \to \operatorname{Ban}$, one for each n, which work as follows.

- On objects, we map a probability space (X, \mathcal{A}, p) to the Banach space $L^n(X, \mathcal{A}, p)$;
- On morphisms, given a measure-preserving kernel $k : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$, we get a bounded linear map $k^* : L^n(Y, \mathcal{B}, q) \to L^n(X, \mathcal{A}, p)$ acting on random variables $g \in L^n(Y, \mathcal{B}, q)$ by

$$k^*g(x) \coloneqq \int_Y g(y) \, k(dy|x). \tag{3.3}$$

Here is the precise statement.

Proposition 3.6. Let $k : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$ be a measure-preserving kernel, and let $g \in L^n(Y, \mathcal{B}, q)$, for n finite or infinite. Then the assignment

$$x \longmapsto k^* g(x) \coloneqq \int_Y g(y) \, k(dy|x)$$

²Here we use L^n rather than L^p in order to avoid confusion with the probability measure p.

is a well-defined element of $L^n(X, \mathcal{A}, p)$.

Moreover, the assignment $k^* : L^n(Y, \mathcal{B}, q) \to L^n(X, \mathcal{A}, p)$ given by $g \mapsto k^*g$ is linear and 1-Lipschitz, so that L^n is a functor $\operatorname{GKrn}^{\operatorname{op}} \to \operatorname{Ban}$.

Proof for finite n. Since k is measure-preserving, by approximating via simple functions,

$$\int_X \int_Y |g(y)|^n \, k(dy|x) \, p(dx) \; = \; \int_Y |g(y)|^n \, q(dy) \; < \; \infty$$

and so the integrals

$$\int_{Y} |g(y)|^n \, k(dy|x)$$

are finite for p-almost all $x \in X$. Even if n > 1, again by Jensen's inequality we have that

$$\left(\int_{Y} |g(y)| \, k(dy|x)\right)^n \leq \int_{Y} |g(y)|^n \, k(dy|x) < \infty.$$

Therefore the integral

$$x\longmapsto \int_Y g(y)\,k(dy|x),$$

is defined for p-almost all $x \in X$, and the assignment specifies a unique element of $L^n(X, \mathcal{A}, p)$ (measurability can be shown by means of the usual approximation via simple functions). We denote this element by k^*g . In other words, given $k : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$, we get a well-defined function $k^* : L^n(Y, \mathcal{B}, q) \to L^n(X, \mathcal{A}, p)$. given by $g \mapsto k^*g$. This function is linear by linearity of integration. To see that it is 1-Lipschitz, once again by Jensen,

$$\begin{aligned} (\|k^*g\|_{L^n})^n &= \int_X \left| \int_Y g(y) \, k(dy|x) \right|^n p(dx) \\ &\leq \int_X \left(\int_Y |g(y)| \, k(dy|x) \right)^n p(dx) \\ &\leq \int_X \int_Y |g(y)|^n \, k(dy|x) \, p(dx) \\ &= \int_Y |g(y)|^n \, q(dy) \\ &= (\|g\|_{L^n})^n. \end{aligned}$$

Before looking at the case of $n = \infty$, let's prove the following auxiliary statement, which can be consider a sort of "unitality" property of the map k^* .

Lemma 3.7. Let $k : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$ be a measure-preserving kernel. Then for every subset $B \in \mathcal{B}$ of q-measure one there exists a subset $A \in \mathcal{A}$ of p-measure one such that for all $x \in A$, k(B|x) = 1.

Proof of Lemma 3.7. Consider $B \in \mathcal{B}$ of measure one, and denote its complement by \overline{B} . Then since k is measure-preserving,

$$0 = q(\bar{B}) = \int_Y k(\bar{B}|x) p(dx),$$

which means that for *p*-almost all $x \in X$, $k(\overline{B}|x) = 0$, i.e. k(B|x) = 1.

Proof of Proposition 3.6 for infinite n. Let $g \in L^{\infty}(Y, \mathcal{B}, q)$. By definition there exists $B \in \mathcal{B}$ of q-measure one such that g restricted to B is bounded. Let K be an upper bound. Since k is measure-preserving, just as for finite n,

$$\int_X \int_Y |g(y)| \, k(dy|x) \, p(dx) \ = \ \int_Y |g(y)| \, q(dy) \ = \ \int_B |g(y)| \, q(dy) \ \le \ K.$$

Therefore the integrals

$$\int_{Y} |g(y)| \, k(dy|x)$$

are finite for *p*-almost all $x \in X$, and so the integral

$$x\longmapsto \int_Y g(y)\,k(dy|x),$$

is defined for p-almost all $x \in X$. Denote this value by $k^*g(x)$.

Moreover, by Lemma 3.7, there exists $A \in \mathcal{A}$ of *p*-measure one such that for all $x \in A$, k(B|x) = 1. Therefore, for all $x \in A$,

$$|k^*g(x)| = \left| \int_Y g(y) \, k(dy|x) \right| \le \int_Y |g(y)| \, k(dy|x) = \int_B |g(y)| \, k(dy|x) \le K.$$

In other words, the restriction of k^*g to A is also bounded by K. So we get a well-defined function $k^* : L^{\infty}(Y, \mathcal{B}, q) \to L^{\infty}(X, \mathcal{A}, p)$. Now since every essential upper bound for |g| is an essential upper bound for $|k^*g|$, we have that

$$\|k^*g\|_{L^{\infty}} = \operatorname{ess\,sup}|k^*g| \le \operatorname{ess\,sup}|g| = \|g\|_{L^{\infty}}.$$

Moreover, the inclusions $L^n(X, \mathcal{A}, p) \subseteq L^m(X, \mathcal{A}, p)$ for $m \leq n$ form a natural transformation $L^n \Rightarrow L^m$.

Let's now see two particular cases of this functorial action.

Example 3.8. A special case of a Markov kernel is a probability measure, which can be seen as a Markov kernel from the one-point space 1. Denote by $(1, \mathcal{N}, \delta_u)$ be the unique probability space on the one-point set 1, where u is the unique element of 1. Given a probability space (X, \mathcal{A}, p) , the only measure-preserving kernel $(1, \mathcal{N}, \delta_u) \to (X, \mathcal{A}, p)$ is given by \tilde{p} , defined in terms of p as follows: for every $A \in \mathcal{A}$,

$$\tilde{p}(A|u) = p(A).$$

Now given any $g \in L^1(X, \mathcal{A}, p)$ (or L^n), we have that $L^1(1, \mathcal{N}, \delta_u) \cong \mathbb{R}$, and

$$ilde{p}^*g(u) \ = \ \int_X g(x)\, ilde{p}(dx|u) \ = \ \int_X g(x)\, p(dx) \ = \ \mathbb{E}[g]$$

In other words, for those kernels which are probability measures, the functorial action of L^1 (and L^n) is exactly giving the expectation values of random variables.

Example 3.9. Let's now consider the case of "deterministic" Markov kernels defined by functions. Let $f : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$ be a measure-preserving function, and consider the induced kernel δ_f as in (3.1). Given a random variable $g \in L^1(Y, \mathcal{B}, q)$ (or L^n), we have for every $x \in X$,

$$(\delta_f)^*g(x) = \int_Y g(y)\,\delta_f(dy|x) = g(f(x)).$$

In other words, $(\delta_f)^*g = g \circ f$. That is, for the kernels in the form δ_f for some measurable function f, the functorial action of L^1 (and L^n) is exactly precomposition of functions. We can view $f : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$ as a "reparametrization" or "refinement" of the outcome space on which our RVs are defined: any RV on Y defines a RV on X, but in general the sigma-algebra \mathcal{A} may be finer than (the one induced by) \mathcal{B} .

In other words, the functorial action of L^1 (and L^n) on kernels is a common generalization of expectations and of refinements of the outcome space.

Proposition 3.10. The functors L^n : GKrn^{op} \rightarrow Ban are faithful. In other words, for measurepreserving kernels $k, h : (X, \mathcal{A}, p) \rightarrow (Y, \mathcal{B}, q)$ we have that k = h p-almost surely if and only if for every random variable $g \in L^n(Y, \mathcal{B}, q)$, $k^*g = h^*g$ p-almost surely.

Proof. It suffices to test the a.s. equality of k and h on the indicator functions of sets of \mathcal{B} , which are in $L^n(Y, \mathcal{B}, q)$ for all n (including $n = \infty$).

Before we leave this section, let's also look at how to talk about *single* random variables categorically, as opposed to spaces of random variables. To this end, notice that a point of a Banach space B is encoded exactly by a (bounded) linear map $f : \mathbb{R} \to B$ (by looking at where f maps $1 \in \mathbb{R}$). Therefore an L^n random variable on (X, \mathcal{A}, p) can be described categorically as a morphism $f : \mathbb{R} \to L^n(X, \mathcal{A}, p)$ of Ban. Its expectation is the composition

$$\mathbb{R} \longrightarrow L^n(X, \mathcal{A}, p) \longrightarrow L^n(1, \mathcal{N}, \delta_u) \cong \mathbb{R}.$$

3.3 Bayesian inversions, conditioning, and dagger structures

The categories Krn and EKrn come equipped with a dagger structure which is very relevant for probability, given by Bayesian inverses. Here is the definition.

Definition 3.11. Let $k : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$ be a measure-preserving kernel. A **Bayesian** inverse of k is a measure-preserving kernel $k^+ : (Y, \mathcal{B}, q) \to (X, \mathcal{B}, p)$ such that for every $A \in \mathcal{A}$ and $B \in \mathcal{B}$,

$$\int_{A} k(B|x) p(dx) = \int_{B} k^{+}(A|y) q(dy).$$
(3.4)

When a Bayesian inverse exists, it is unique almost surely. It is easy to see that the identity kernel is its own Bayesian inverse, and that Bayesian inverses are closed under composition. Moreover, every invertible kernel (invertible in GKrn) has as inverse its Bayesian inverse.

In what follows it is useful to set some notation for conditional expectations. Given a probability space (X, \mathcal{A}, p) , a sub-sigma-algebra $\mathcal{B} \subseteq \mathcal{A}$ and an integrable RV $f \in L^1(X, \mathcal{A}, p)$, recall that a **conditional expectation of** f given \mathcal{B} is a RV $g \in L^1(X, \mathcal{B}, p)$ such that for every $B \in \mathcal{B}$,

$$\int_{B} g(x) \, p(dx) = \int_{B} f(x) \, p(dx).$$

Such a conditional expectation, if it exists, is unique almost surely. We denote it by

$$x \mapsto \mathbb{E}[f|\mathcal{B}](x).$$

We also write, for $A \in \mathcal{A}$, the shorthand

$$\mathbb{P}[A|\mathcal{B}](x) \coloneqq \mathbb{E}[1_A|\mathcal{B}](x),$$

where 1_A is the indicator function.

We will use the following classic measure-theoretic result, which we restate in our notation:

Theorem 3.12 (Rokhlin's disintegration theorem). Let (X, \mathcal{A}, p) be a standard Borel probability space. Consider a sub-sigma algebra $\mathcal{B} \subseteq \mathcal{A}$, and the kernel $\pi := \delta_{id} : (X, \mathcal{A}, p) \to (X, \mathcal{B}, p)$ induced by the set-theoretical identity of X via (3.1). Then π has a Bayesian inverse (or **disintegration**) $\pi^+ : (X, \mathcal{B}, p) \to (X, \mathcal{A}, p)$ such that for all $A \in \mathcal{A}$, for p-almost all $x \in X$,

$$\pi^+(A|x) = \mathbb{P}[A|\mathcal{B}](x).$$

On random variables, the functorial action of kernel π^+ : $(X, \mathcal{B}, p) \to (X, \mathcal{A}, p)$ takes an \mathcal{A} measurable random variable g and returns its conditional expectation with respect to \mathcal{B} :

$$(\pi^{+})^{*}g = ((\delta_{\rm id})^{+})^{*}g = \mathbb{E}[g|\mathcal{B}].$$
(3.5)

Here is a well known, important consequence of the theorem:

Proposition 3.13. Every measure-preserving kernel between standard Borel spaces admits a Bayesian inverse.

Proof. Let (X, \mathcal{A}) and (Y, \mathcal{B}) be standard Borel spaces, and let $k : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$ be a measure-preserving kernel. Form the joint probability measure r on $(X \times Y, \mathcal{A} \otimes \mathcal{B})$ specified on rectangles by

$$r(A \times B) \coloneqq \int_A k(B|x) \, p(dx)$$

Let's show that the projection to the second marginal $\pi_2 : (X \times Y, \mathcal{A} \otimes \mathcal{B}, r) \to (Y, \mathcal{B}, q)$ has a Bayesian inverse. By [EP23, Proposition 2.7], we can rewrite (Y, \mathcal{B}, q) , up to isomorphism of GKrn, as the set $X \times Y$ equipped with the sigma-algebra $\pi_2^{-1}(\mathcal{B})$ induced by the projection π_2 , and the measure r restricted to $\pi_2^{-1}(\mathcal{B})$. Therefore π_2 can be equivalently written as a deterministic kernel $(X \times Y, \mathcal{A} \otimes \mathcal{B}, r) \to (X \times Y, \pi_2^{-1}(\mathcal{B}), r) \cong (Y, \mathcal{B}, q)$. By Theorem 3.12, we then have a Bayesian inverse $\pi_2^+ : (Y, \mathcal{B}, q) \cong (X \times Y, \pi_2^{-1}(\mathcal{B}), r) \to (X \times Y, \mathcal{A} \otimes \mathcal{B}, r)$ as follows,

$$\pi_2^+(A \times B|y) = \mathbb{E}\big(1_A \cdot 1_B | \pi_2^{-1}(\mathcal{B})\big)(x, y)$$

for $A \in \mathcal{A}$ and $B \in \mathcal{B}$, and for $x \in X$ and $y \in Y$ (the quantity, almost surely, does not depend on x). Composing it with the projection $\pi_1 : X \times Y \to X$ we obtain a kernel $h : Y \to X$ given by

$$h(A|y) = \mathbb{E}(1_A | \pi_2^{-1}(\mathcal{B}))(x, y),$$

which again, almost surely, does not depend on x. So now for $A \in \mathcal{A}$ and $B \in \mathcal{B}$,

$$\int_{B} h(A|y) q(dy) = \int_{\pi_{2}^{-1}(B)} \mathbb{E}(1_{A}|\pi_{2}^{-1}(\mathcal{B}))(x, y) r(dx \, dy)$$
$$= \int_{\pi_{2}^{-1}(B)} 1_{A}(x) r(dx \, dy)$$

$$= r(A \times B)$$
$$= \int_{A} k(B|x) p(dx)$$

so that h is a Bayesian inverse of k.

Bayesian inversion is therefore defined on all morphisms, and it makes Krn a dagger category (see also [DDGS18, Theorem 2.10]). Since isomorphisms of GKrn admit Bayesian inverses as well, this also implies that every kernel between *essentially* standard Borel spaces admits a Bayesian inverse as well, and so EKrn is a dagger category as well.

Remark 3.14. Since Krn and EKrn are dagger categories, we can also define a *covariant* version of the random variable functors, $Krn \rightarrow Ban$, by precomposing with the dagger:

$$\operatorname{Krn} \xrightarrow{(-)^+} \operatorname{Krn}^{\operatorname{op}} \xrightarrow{L^n} \operatorname{Ban}$$

We can restrict this functor to the deterministic kernels in the form δ_f . If we denote by BorelMeas the category of standard Borel probability spaces and measure-preserving functions, the resulting functor BorelMeas \rightarrow Ban acts as follows. First of all, given a standard Borel space (X, \mathcal{A}, p) and a sub-sigma-algebra $\mathcal{B} \subseteq \mathcal{A}$, the map $L^n(X, \mathcal{A}, p) \rightarrow L^n(X, \mathcal{B}, A)$ takes an \mathcal{A} -measurable RV and gives its conditional expectation, as in (3.5). More generally, given a measure-preserving function $f : (X, \mathcal{A}, p) \rightarrow (Y, \mathcal{B}, q)$, the map $L^n(X, \mathcal{A}, p) \rightarrow L^n(Y, \mathcal{B}, q)$ takes a RV g on X and gives the following RV on Y:

$$y \mapsto \mathbb{E}[g|f^{-1}(\mathcal{B})](x) \quad \text{for } x \in f^{-1}(y),$$

where $f^{-1}(\mathcal{B}) \subseteq \mathcal{A}$ is the pullback sigma-algebra induced by f, and where the quantity above does not depend on the choice of $x \in f^{-1}(y)$. This functor was defined in [AR18] with the name \mathcal{E} and in [VB23] with the name RV.

Let's now focus on the functor L^2 . Given a probability space (X, \mathcal{A}, p) , the space $L^2(X, p)$ is a Hilbert space, with inner product given by

$$\langle f,g\rangle \;=\; \int_X f(x)\,g(x)\,p(dx).$$

Therefore we can consider L^2 as a functor $\operatorname{GKrn} \to \operatorname{Hilb}$.

Proposition 3.15. The functor $L^2 : \text{EKrn} \to \text{Hilb}$ is a dagger functor, meaning that the following diagram commutes.

$$\begin{array}{ccc} \operatorname{EKrn^{op}} & \xrightarrow{L^2} & \operatorname{Hilb} \\ & & \downarrow (-)^+ & & \downarrow (-)^+ \\ \operatorname{EKrn^{op}} & \xrightarrow{L^2} & \operatorname{Hilb} \end{array}$$

Proof. On objects, the assert is trivial. On morphisms, consider a measure-preserving kernel $k : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$. The commutativity of the diagram says that $(k^*)^+ = (k^+)^*$ In other words, given random variables $f \in L^2(X, \mathcal{A}, p)$ and $g \in L^2(Y, \mathcal{B}, q)$,

$$\langle f, k^*g \rangle = \langle (k^+)^*f, g \rangle.$$

In terms of integrals, equivalently,

$$\int_X \int_Y f(x) \, g(y) \, k(dy|x) \, p(dx) \; = \; \int_X \int_Y f(x) \, g(y) \, k^+(dx|y) \, q(dy).$$

This now follows from (3.4) by approximating f and g via simple functions.

One of the first probabilistic concepts that we can express in terms of the dagger structure is (almost sure) determinism. Let's start with a definition, which follows [Fri20, Section 10 and Section 13].

Definition 3.16. A measure-preserving kernel $(X, \mathcal{A}, p) \rightarrow (Y, \mathcal{B}, q)$ is called **deterministic** if for every $x \in X$ and for every $B \in \mathcal{B}$,

$$k(B|x) = 0 \quad or \quad k(B|x) = 1.$$

The kernel k is called p-almost surely deterministic if for every $B \in \mathcal{B}$, the set of $x \in X$ where $k(B|x) \in \{0,1\}$ has p-measure one.³

Every kernel in the form δ_f for a measurable function f is deterministic. The converse in general is not true, since the sigma-algebra on the codomain may fail to separate points. However, every kernel between standard Borel spaces is deterministic if and only if it is in the form δ_f [Fri20, Example 10.5].

Almost sure determinism, in Krn and EKrn, can be expressed in terms of the dagger structure as follows:

Proposition 3.17 ([EP23, Proposition 2.5]). A Markov kernel k admitting a Bayesian inverse k^+ is almost surely deterministic if and only if $k \circ k^+ = id$. In particular, a morphism of EKrn is almost surely deterministic if and only if it is a dagger epi.

Since every isomorphism of EKrn is almost surely deterministic, it follows that every isomorphism of EKrn is a dagger iso.

4 Idempotent kernels and sub-sigma-algebras

We now study (dagger) idempotents in the category EKrn. We will show that these correspond to sub-sigma-algebras up to almost sure equality, and that their functorial action on random variables will be given by conditional expectation. Similarly, the functorial action of a filtration of sub-sigmaalgebras will give martingales.

4.1 Conditional expectation operators

Let's look at the idempotent morphisms of Krn and EKrn. Since everything is defined up to almost sure equality, from the measure-theoretic point of view the kernels we are interested in are *almost surely* idempotent, in the following sense.

³Note that this may be more general than requiring that there exists a measure-one subset independent of B on which $k(B|x) \in \{0, 1\}$. If \mathcal{B} is countably generated, for example if (Y, \mathcal{B}) is standard Borel, the two notions coincide.

Definition 4.1. A measure-preserving kernel $e : (X, \mathcal{A}, p) \to (X, \mathcal{A}, p)$ is *p*-almost surely idempotent if and only if for every $B \in \mathcal{A}$ and for *p*-almost all $x \in X$, we have

$$\int_X e(B|x') e(dx'|x) = e(B|x)$$

Equivalently, if for all $A, B \in \mathcal{A}$,

$$\int_A \int_X e(B|x') e(dx'|x) p(dx) = \int_A e(B|x) p(dx)$$

The almost sure equivalence classes of these kernels, between standard Borel probability spaces, are precisely the idempotent morphisms of GKrn. In terms of random variables, equivalently, a kernel e is a.s. idempotent if and only if for every RV f, $e^*e^*f = e^*f$ almost surely. That is, if the corresponding operator e^* on $L^n(X, \mathcal{A}, p)$ is idempotent.

Here is our main example of idempotent kernel. As we will show, all idempotent kernels can be written in this form.

Definition 4.2. Let (X, \mathcal{A}, p) be a standard Borel probability space, and consider a sub-sigmaalgebra $\mathcal{B} \subseteq \mathcal{A}$. The conditional expectation operator is the equivalence class of Markov kernels $e_{\mathcal{B}}: (X, \mathcal{A}, p) \to (X, \mathcal{A}, p)$ given by

$$e_{\mathcal{B}}(A|x) \coloneqq \mathbb{P}[A|\mathcal{B}](x) = \mathbb{E}[1_A|\mathcal{B}](x),$$

for each $A \in \mathcal{A}$ and for p-almost all $x \in X$.

Note that by the disintegration theorem (Theorem 3.12) we know that these conditional expectations can indeed be assembled to a kernel $(X, \mathcal{B}, p) \to (X, \mathcal{A}, p)$. Since $\mathcal{B} \subseteq \mathcal{A}$, this assignment is also measurable as a kernel $(X, \mathcal{A}, p) \to (X, \mathcal{A}, p)$.

The action of $e_{\mathcal{B}}$ on random variables, almost surely, is conditional expectation (hence the name):

$$(e_{\mathcal{B}})^* f(x) = \int_X f(x') e_{\mathcal{B}}(dx'|x) = \mathbb{E}[f|\mathcal{B}](x).$$

By idempotency of conditional expectations, we then have that $(e_{\mathcal{B}})^* : L^n(X, \mathcal{A}, p) \to L^n(X, \mathcal{A}, p)$ is idempotent, and hence so is $e_{\mathcal{B}}$.

The idempotent $e_{\mathcal{B}}$ is split by \mathcal{B} :

Lemma 4.3. Let (X, \mathcal{A}, p) be a standard Borel probability space, and let $\mathcal{B} \subseteq \mathcal{A}$ be a sub-sigmaalgebra. A splitting of the idempotent $e_{\mathcal{B}} : (X, \mathcal{A}, p) \to (X, \mathcal{A}, p)$ in GKrn is given by (X, \mathcal{B}, p) , together with the map $\pi = \delta_{id} : (X, \mathcal{A}, p) \to (X, \mathcal{B}, p)$ induced by the set-theoretic identity, and its Bayesian inverse $\pi^+ = (\delta_{id})^+ : (X, \mathcal{B}, p) \to (X, \mathcal{A}, p)$ (which exists by Theorem 3.12).

Proof. First of all, we have that $\pi \circ \pi^+ = \mathrm{id}_{(X,\mathcal{B},p)}$ almost surely, since π is deterministic (using Proposition 3.17). To show that $\pi^+ \circ \pi = e_{\mathcal{B}}$ almost surely, notice that for every $A \in \mathcal{A}$ and for almost all $x \in X$, using Theorem 3.12,

$$(\pi^{+} \circ \pi)(A|x) = \int_{X} \pi^{+}(A|x') \pi(dx'|x) = \int_{X} \mathbb{P}[A|\mathcal{B}](x') \pi(dx'|x) = \mathbb{P}[A|\mathcal{B}](x) = e_{\mathcal{B}}(A|x).$$

In [EP23, Theorem 3.14] it was proven that all idempotents of Krn split, and since Krn is a full subcategory of GKrn, the splitting in GKrn is going to be isomorphic to the one in Krn, i.e., it is in EKrn. In this work we prefer to use sub-sigma-algebras, which are technically not standard Borel (only essentially so), but easier and more explicit to work with. This is why we are interested in the category EKrn.

It turns out that all idempotent morphisms of Krn can be written in the form $e_{\mathcal{B}}$ for some subsigma-algebra \mathcal{B} , and that this gives a splitting of the idempotent (in EKrn). This sub-sigma-algebra is the *invariant sigma-algebra*, defined here:

Definition 4.4. Let $k : (X, \mathcal{A}, p) \to (X, \mathcal{A}, p)$ be a measure-preserving kernel. A measurable subset $B \in \mathcal{A}$ is called *p*-almost surely invariant under k if and only if for *p*-almost all $x \in X$, we have

$$k(B|x) = 1_B(x).$$

Equivalently, if for all $A \in \mathcal{A}$,

$$\int_A k(B|x) p(dx) = \int_A 1_B(x) p(dx) = p(A \cap B).$$

Almost surely invariant sets form a sub-sigma-algebra of \mathcal{A} , which we denote by \mathcal{I}_k , and call the **invariant sigma-algebra**.

Remark 4.5. For every measure-preserving kernel $k : (X, \mathcal{A}, p) \to (X, \mathcal{A}, p)$, the invariant sigmaalgebra \mathcal{I}_k contains all the null sets. Indeed, suppose that for $B \in \mathcal{A}$, p(B) = 0. Then for all $A \in \mathcal{A}$,

$$\int_A k(B|x) p(dx) \le \int_X k(B|x) p(dx) = p(B) = 0 = p(A \cap B).$$

In particular, if (X, \mathcal{A}, p) is a complete measure space, so is (X, \mathcal{I}_k, p) . Similarly, \mathcal{I}_k also contains all the full-measure sets. A converse to this statement, for the standard Borel case, will be given in Corollary 4.17.

Invariant sigma-algebras split idempotents as follows.

Theorem 4.6. Let $e: (X, \mathcal{A}, p) \to (X, \mathcal{A}, p)$ be an idempotent morphism of Krn. Consider the invariant sigma-algebra \mathcal{I}_e . Form the kernel $\pi: (X, \mathcal{A}, p) \to (X, \mathcal{I}_e, p)$ induced by the set-theoretic identity, as in Theorem 3.12, and consider its Bayesian inverse $\pi^+: (X, \mathcal{I}_e, p) \to (X, \mathcal{A}, p)$ (which exists by Theorem 3.12). Then $((X, \mathcal{I}_e, p), \pi^+, \pi)$ is a splitting of e in GKrn.

We will use the following auxiliary statements.

Lemma 4.7 ([EP23, Lemmas 3.7 and 3.9]). Let $k : (X, \mathcal{A}, p) \to (X, \mathcal{A}, p)$ be a measure-preserving kernel. A bounded \mathcal{A} -measurable function $f : X \to \mathbb{R}$ is \mathcal{I}_k -measurable if and only if for p-almost all x,

$$k^*f(x) = f(x).$$

(Such functions are sometimes called harmonic.)

Lemma 4.8. Let $k : (X, \mathcal{A}, p) \to (X, \mathcal{A}, p)$ be a measure-preserving kernel, and let $A \in \mathcal{I}_k$. Then for every $B \in \mathcal{A}$, for p-almost all $x \in X$,

$$k(A|x) k(B|x) = k(A \cap B|x).$$

This lemma can be considered as an instance of *relative positivity in a Markov category*, see [Fri20, Example 13.19] and [FGHL⁺23, Section 2.5].

Proof of Lemma 4.8. Define the sets

$$X_0 \coloneqq \{ x \in X : k(A|x) = 0 \}, \qquad X_1 \coloneqq \{ x \in X : k(A|x) = 1 \}.$$

Since $A \in \mathcal{I}_k$, we have that $k(A|x) = 1_A(x)$ *p*-almost surely, and so $p(X_0 \cup X_1) = 1$. Now let $B \in \mathcal{A}$. For $x \in X_0$,

$$k(A|x) k(B|x) = 0 = k(A \cap B|x).$$

Similarly, for $x \in X_1$,

$$k(A|x) k(B|x) = k(B|x) = k(A \cap B|x)$$

We are now ready to prove the theorem.

Proof of Theorem 4.6. By Lemma 4.3, it suffices to show that $e = e_{\mathcal{I}_e}$ (almost surely). So let $A \in \mathcal{A}$. For p-almost all $x \in X$,

$$e_{\mathcal{I}_e}(A|x) = \mathbb{P}[A|\mathcal{I}_e](x) = \mathbb{E}[1_A|\mathcal{I}_e](x).$$

To show that this conditional expectation is almost surely equal to e(A|x), it suffices to show that e(A|x) is a conditional expectation in the form above. This means that:

- (i) The function $x \mapsto e(A|x)$ is \mathcal{I}_e -measurable;
- (ii) For every $B \in \mathcal{I}_e$,

$$\int_B e(A|x) p(dx) = \int_B 1_A(x) p(dx) = p(A \cap B).$$

To prove (i), notice that since e is idempotent, the function $x \mapsto e(A|x)$ satisfies

$$\int_X e(A|x') e(dx'|x) = e(A|x) \quad \text{for } p\text{-almost all } x$$

Therefore it is harmonic in the sense of Lemma 4.7, and so it is \mathcal{I}_e -measurable. To prove (i) we use, in order, the fact that $B \in \mathcal{I}_e$, Lemma 4.8, and the fact that e is measure-preserving:

$$\int_{B} e(A|x) p(dx) = \int_{X} e(A|x) e(B|x) p(dx)$$
$$= \int_{X} e(A \cap B|x) p(dx)$$
$$= p(A \cap B).$$

Therefore $e(A|x) = e_{\mathcal{I}_e}(A|x)$, and by Lemma 4.3, \mathcal{I}_e splits e.

As we remarked, (X, \mathcal{I}_e, p) is in EKrn, which is a dagger category, and so this splitting, given by π and its Bayesian inverse, is a dagger splitting.

Corollary 4.9 ([EP23, end of Section 3]). Every idempotent morphism of EKrn is a dagger idempotent.

This can be interpreted as a detailed balance condition: every measure-preserving, idempotent Markov kernel induces a reversible process. Also, since we have seen that the functor L^2 : EKrn^{op} \rightarrow Hilb preserves the dagger, every idempotent kernel induces a self-adjoint, idempotent operator $e^* : L^2(X, \mathcal{A}, p) \rightarrow L^2(X, \mathcal{A}, p)$, i.e. an orthogonal projector. (Conversely, a kernel induces an orthogonal projector if and only if it is idempotent.)

Remark 4.10. Since idempotent splittings are preserved by every functor, and so we also have that for every *n*, the idempotent $e^* : L^n(X, \mathcal{A}, p) \to L^n(X, \mathcal{A}, p)$ is split by $L^2(X, \mathcal{I}_e, p)$. Mind that the functors L^n reverse the direction of the arrows, so that the idempotent operator e^* factors as follows,

$$L^{n}(X,\mathcal{A},p) \xrightarrow{(\pi^{+})^{*}} L^{n}(X,\mathcal{I}_{e},p) \xrightarrow{\pi^{*}} L^{n}(X,\mathcal{A},p)$$
$$f \longmapsto \mathbb{E}[f|\mathcal{I}_{e}] \longmapsto \mathbb{E}[f|\mathcal{I}_{e}].$$

The map π^* is the inclusion of \mathcal{I}_e -measurable RVs into all \mathcal{A} -measurable ones, and the map $(\pi^+)^*$ is the projection of \mathcal{A} -measurable RVs onto \mathcal{I}_e -measurable ones given by taking conditional expectations. As it is well known, for the L^2 case this projection is orthogonal.

4.2 The preorder of sub-sigma-algebras

The assignments $e \mapsto \mathcal{I}_e$ and $\mathcal{B} \mapsto e_{\mathcal{B}}$ between idempotents and sub-sigma-algebras are almost inverse to each other. The reason why they are not exactly inverses is that conditional expectations cannot distinguish sigma-algebras when they only differ by null sets. Let's make this precise. First of all, recall that given sub-sigma-algebras $\mathcal{B}, \mathcal{C} \subseteq \mathcal{A}$, we say that \mathcal{B} is coarser than \mathcal{C} if and only if $\mathcal{B} \subseteq \mathcal{C}$, i.e. if every $B \in \mathcal{B}$ is also in \mathcal{C} . We now present an almost-sure version of this idea.

Definition 4.11. Let (X, \mathcal{A}, p) be a probability space, and let $\mathcal{B}, \mathcal{C} \subseteq \mathcal{A}$ be sub-sigma-algebras.

- We say that a subset $B \in \mathcal{B}$ is almost surely in C if there exists some $C \in C$ such that $p(B \triangle C) = 0$. (Here $B \triangle C$ denotes the symmetric difference, i.e. $B \triangle C := (B \setminus C) \cup (C \setminus B)$. Equivalently, 1_B and 1_C are p-almost surely equal.)
- We say that B is almost surely coarser than C (and that C is almost surely finer than B) if every B ∈ B is almost surely in C. In this case we write B ≤ C.
- We say that \mathcal{B} and \mathcal{C} are almost surely equivalent, and we write $\mathcal{B} \simeq \mathcal{C}$, if $\mathcal{B} \lesssim \mathcal{C}$ and $\mathcal{C} \lesssim \mathcal{B}$.

Let's also define this idea from the point of view of random variables.

Definition 4.12. Let (X, \mathcal{A}, p) be a probability space, and let $\mathcal{C} \subseteq \mathcal{A}$ be a sub-sigma-algebra. We say that an \mathcal{A} -measurable random variable (not necessarily integrable) f is almost surely \mathcal{C} -measurable if and only if it is p-almost surely equal to a \mathcal{C} -measurable function. That is, if there exists $A \in \mathcal{A}$ of p-measure one and a \mathcal{C} -measurable RVg such that f and g are equal when restricted to A.

The following statements give further equivalent descriptions of this preorder. We start with equivalent condition to "being almost surely in a sigma-algebra":

Lemma 4.13. Let (X, \mathcal{A}, p) be a standard Borel probability space, and let $\mathcal{B}, \mathcal{C} \subseteq \mathcal{A}$ be a sub-sigmaalgebra. For $B \in \mathcal{B}$, the following conditions are equivalent.

- (i) B is almost surely in C;
- (ii) The indicator function 1_B is almost surely C-measurable;
- (iii) For p-almost all x, $\mathbb{P}[B|\mathcal{C}](x) = 1_B(x)$, i.e. the indicator function 1_B is p-almost surely equal to some (hence every) conditional expectation $\mathbb{E}[1_B|\mathcal{C}]$;
- (iv) $B \in \mathcal{I}_{e_{\mathcal{C}}}$, where $e_{\mathcal{C}}$ is any kernel (in its equivalence class) constructed as in Definition 4.2, and $\mathcal{I}_{e_{\mathcal{C}}}$ is its invariant sigma-algebra as in Definition 4.4.

Proof. For $(i) \Rightarrow (ii)$, notice that B is almost surely in C if and only if there exists $C \in C$ such that $1_B(x) = 1_C(x)$ for p-almost all $x \in X$. Now 1_C is C-measurable.

For $(ii) \Rightarrow (i)$, suppose that 1_B is *p*-almost surely equal to a *C*-measurable function $f: X \to \mathbb{R}$. Then the set $C = f^{-1}(1)$ is in \mathcal{C} , and it differs from B only by a measure zero set.

For $(ii) \Rightarrow (iii)$, suppose that 1_B is *p*-almost surely equal to a *C*-measurable function $f : X \to \mathbb{R}$. Since conditional expectations respect almost-sure equality, we have that for *p*-almost all *x*,

$$\mathbb{E}[1_B|\mathcal{C}](x) = \mathbb{E}[f|\mathcal{C}](x) = f(x) = 1_B(x).$$

For $(iii) \Rightarrow (ii)$, notice that if $\mathbb{E}[1_B|\mathcal{C}](x) = 1_B(x)$ *p*-almost surely, then 1_B is *p*-almost surely equal to a \mathcal{C} -measurable random variable, namely $\mathbb{E}[1_B|\mathcal{C}]$.

For $(iii) \Leftrightarrow (iv)$, notice that the condition that $B \in \mathcal{I}_{e_{\mathcal{C}}}$ means exactly that for *p*-almost all x, $e_{\mathcal{C}}(B|x) = 1_B(x)$. This in turn means exactly that for *p*-almost all $x \in X$, $\mathbb{P}[B|\mathcal{C}](x) = 1_B(x)$. \Box

Here are analogous, equivalent conditions for "almost sure measurability" of functions, which further motivate the name.

Lemma 4.14. Let (X, \mathcal{A}, p) be a probability space, let $\mathcal{C} \subseteq \mathcal{A}$ be a sub-sigma-algebra, and let f be an \mathcal{A} -measurable random variable. The following conditions are equivalent.

- (i) f is almost surely C-measurable;
- (ii) For every Borel set $B \subseteq \mathbb{R}$, $f^{-1}(B)$ is almost surely in \mathcal{C} .
- (iii) (for f integrable) f is almost surely equal to its conditional expectation $\mathbb{E}[f|\mathcal{C}]$;
- (iv) f is $\mathcal{I}_{e_{\mathcal{C}}}$ -measurable.

Proof. For $(i) \Rightarrow (ii)$, suppose that f is almost surely equal to a C-measurable RV g. Then for every Borel set $B \subseteq \mathbb{R}$, $p(f^{-1}(B) \bigtriangleup g^{-1}(B)) = 0$. Therefore $f^{-1}(B)$ is almost surely in C.

For $(ii) \Rightarrow (i)$, let f_+ and f_- be the positive and negative parts of $f = f_+ - f_-$. We can express f_+ (and f_-) as pointwise suprema of a monotone sequence of simple functions,

$$f_+(x) = \sup_n f_n(x)$$
 $f_n(x) = \sum_{i_n=1}^n \alpha_{n,i_n} \mathbf{1}_{A_{n,i_n}}(x),$

for $n \in \mathbb{N}$ and $i_n = 1, ..., n$, where the sets A_{n,i_n} are in the form $f^{-1}(B)$ for some Borel set $B \subseteq \mathbb{R}$. By hypothesis, for each n and i_n we can find a set $C_{n,i_n} \in \mathcal{C}$ such that $p(C_{n,i_n} \triangle A_{n,i_n}) = 0$. Define now

$$g_+(x) = \sup_n g_n(x)$$
 $g_n(x) = \sum_{i_n=1}^n \alpha_{n,i_n} \mathbf{1}_{C_{n,i_n}}(x).$

We have that g_+ is C-measurable, and almost surely equal to f_+ . The same can be done for f_- , and this way we can have C-measurable function g which is almost surely equal to f.

For $(i) \Leftrightarrow (iii)$, notice that $\mathbb{E}[f|\mathcal{C}]$ is \mathcal{C} -measurable.

The equivalence $(iii) \Leftrightarrow (iv)$ follows from the equivalence between the items (i) and (iv) of Lemma 4.13.

The previous two lemmas have the following consequences:

Corollary 4.15. Let (X, \mathcal{A}, p) be a standard Borel probability space, and let $\mathcal{B}, \mathcal{C} \subseteq \mathcal{A}$ be sub-sigmaalgebras. The following conditions are equivalent.

- (i) $\mathcal{B} \lesssim \mathcal{C}$;
- (ii) For every $B \in \mathcal{B}$, 1_B is almost surely \mathcal{C} -measurable;
- (iii) For every $B \in \mathcal{B}$, $\mathbb{P}[B|\mathcal{C}] = 1_B$ p-almost surely;
- (iv) $\mathcal{B} \subseteq \mathcal{I}_{e_{\mathcal{C}}}$.
- (v) Every \mathcal{B} -measurable random variable is almost surely \mathcal{C} -measurable;
- (vi) Every \mathcal{B} -measurable, integrable random variable f is a.s. equal to the conditional expectation $\mathbb{E}[f|\mathcal{C}];$
- (vii) Every \mathcal{B} -measurable random variable f is $\mathcal{I}_{e_{\mathcal{C}}}$ -measurable.
- (viii) For every $A \in \mathcal{A}$, the conditional expectation $\mathbb{P}[A|\mathcal{B}]$ is almost surely \mathcal{C} -measurable;
- (ix) For every A-measurable, integrable random variable f, the conditional expectation $\mathbb{E}[f|\mathcal{B}]$ is almost surely C-measurable.

Corollary 4.16. Let (X, \mathcal{A}, p) be a standard Borel probability space, and let $\mathcal{B}, \mathcal{C} \subseteq \mathcal{A}$ be sub-sigmaalgebras.

- (i) By setting C = B in Corollary 4.15, we see that for every $B \subseteq A$, $B \subseteq I_{e_B}$.
- (ii) By setting $\mathcal{B} = \mathcal{I}_{ec}$ in Corollary 4.15, and relabeling, we see that for every $\mathcal{B} \subseteq \mathcal{A}$, $\mathcal{I}_{e_{\mathcal{B}}} \leq \mathcal{B}$. Using the point above, this gives that $\mathcal{I}_{e_{\mathcal{B}}} \simeq \mathcal{B}$.
- (iii) The two points above show that $\mathcal{I}_{e_{\mathcal{B}}}$ is the finest (in the strict sense) sigma-algebra in its almost sure equivalence class.
- (iv) For every $\mathcal{B}, \mathcal{C} \subseteq \mathcal{A}$, we have that $\mathcal{B} \lesssim \mathcal{C}$ if and only if $\mathcal{I}_{e_{\mathcal{B}}} \subseteq \mathcal{I}_{e_{\mathcal{C}}}$. In particular, $\mathcal{B} \simeq \mathcal{C}$ if and only if $\mathcal{I}_{e_{\mathcal{B}}} = \mathcal{I}_{e_{\mathcal{C}}}$.

Moreover, from Lemma 4.13 we get a converse to Remark 4.5:

Corollary 4.17. Let (X, \mathcal{A}, p) be a standard Borel probability space, and let $\mathcal{B} \subseteq \mathcal{A}$ be a sub-sigmaalgebra. We have that $\mathcal{B} = \mathcal{I}_{e_{\mathcal{B}}}$ if and only if \mathcal{B} contains all the null sets.

Proof. First of all, by Corollary 4.16, $\mathcal{B} \subseteq \mathcal{I}_{e_{\mathcal{B}}}$. Now suppose that \mathcal{B} contains all the null sets, and suppose that $A \in \mathcal{I}_{e_{\mathcal{B}}}$. Then by Lemma 4.13(i), A is almost surely in \mathcal{B} , meaning that there exists $B \in \mathcal{B}$ such that $p(A \bigtriangleup B) = 0$. But we can now write A as

$$A = (B \cup (A \setminus B)) \setminus (B \setminus A).$$

By hypothesis, the set B is in \mathcal{B} , and so are $A \setminus B$ and $B \setminus A$, since they are null. Therefore $A \in \mathcal{B}$. The converse is given by Remark 4.5.

Because of this, one can either work with almost sure equivalence classes of sub-sigma-algebras, or with sub-sigma-algebras which contain all the null sets.

We are now ready for the main statement of this section.

Theorem 4.18. Let (X, \mathcal{A}, p) be a standard Borel probability space. The assignments $e \mapsto \mathcal{I}_e$ and $\mathcal{B} \mapsto e_{\mathcal{B}}$ establish a bijection between

- Almost sure equivalence classes of measure-preserving, a.s. idempotent kernels $e: (X, \mathcal{A}, p) \rightarrow (X, \mathcal{A}, p)$, and
- Almost sure equivalence classes of sub-sigma-algebras $\mathcal{B} \subseteq \mathcal{A}$.

Moreover, this bijection is order-preserving, in the sense that $\mathcal{B} \leq \mathcal{C}$ if and only if $e_{\mathcal{B}} \leq e_{\mathcal{C}}$ in the order of idempotents.

Proof. Note that we already have $\mathcal{I}_{e_{\mathcal{B}}} \simeq \mathcal{B}$ by Corollary 4.16(ii). To prove the bijectivity of the assignment, it suffices to show that the map $e \mapsto \mathcal{I}_e$ is injective (between equivalence classes). This is however guaranteed by Theorem 4.6: each idempotent e can be written (split) as $\pi^+ \circ \pi$, where π is the canonical kernel $\pi : (X, \mathcal{A}, p) \to (X, \mathcal{I}_e, p)$.

Let's now turn to the order. By Corollary 4.16(iv), it suffices to show that for idempotents e, e', we have $e \leq e'$ if and only $\mathcal{I}_e \subseteq \mathcal{I}_{e'}$. So first suppose that $e \leq e'$ in the order of idempotents, which means that $e \circ e' = e' \circ e = e$. Suppose that $B \in \mathcal{I}_e$, which means that for *p*-almost all *x*, $e(B|x) = 1_B(x)$. Equivalently, $e(B|x) = 1_B(x)$ for all $x \in A$ for some set $A \in \mathcal{A}$ with p(A) = 1. Now since e' is measure-preserving,

$$1 = p(A) = \int_X e'(A|x) \, p(dx),$$

which means that for p-almost all $x \in A$, e'(A|x) = 1 as well. Therefore for p-almost all x,

$$e'(B|x) = \int_X 1_B(x') e'(dx'|x)$$
$$= \int_A 1_B(x') e'(dx'|x)$$
$$= \int_A e(B|x) e'(dx'|x)$$
$$= \int_X e(B|x) e'(dx'|x)$$

$$= e \circ e' (B|x)$$
$$= e(B|x)$$
$$= 1_B(x),$$

which means that $B \in \mathcal{I}_{e'}$ as well. Therefore $\mathcal{I}_e \subseteq \mathcal{I}_{e'}$.

Conversely, suppose that $\mathcal{I}_e \subseteq \mathcal{I}_{e'}$. Then the set-theoretic identity on X gives a commutative triangle of measure spaces and (strictly) deterministic, measure-preserving kernels as follows.



where for any sub-sigma-algebra $\mathcal{B} \subseteq \mathcal{C}$, recall that the kernel $\delta_{id} : (X, \mathcal{C}, p) \to (X, \mathcal{B}, p)$ is defined by $\delta_{id}(B|x) \coloneqq 1_B(x)$ for every $B \in \mathcal{B}$ and $x \in X$. Therefore, by Proposition 2.12 (where condition (iii) applies), $e \leq e'$.

Here are further, convenient characterizations.

Corollary 4.19. Let (X, \mathcal{A}, p) be standard Borel, and let $\mathcal{B}, \mathcal{C} \subseteq \mathcal{A}$ be sub-sigma-algebras. We have that $\mathcal{B} \leq \mathcal{C}$ if and only if for every integrable random variable on (X, \mathcal{A}, p) , we have that almost surely,

$$\mathbb{E}\big[\mathbb{E}[f|\mathcal{B}]\big|\mathcal{C}\big] = \mathbb{E}\big[\mathbb{E}[f|\mathcal{C}]\big|\mathcal{B}\big] = \mathbb{E}[f|\mathcal{B}].$$

In particular, $\mathcal{B} \simeq \mathcal{C}$ if and only if for every integrable random variable f, $\mathbb{E}[f|\mathcal{B}]$ and $\mathbb{E}[f|\mathcal{C}]$ are p-almost surely equal.

Corollary 4.20. Let (X, \mathcal{A}, p) be standard Borel, and let $\mathcal{B}, \mathcal{C} \subseteq \mathcal{A}$ be sub-sigma-algebras. Consider the following kernels induced by the set-theoretic identities,

$$(X, \mathcal{A}, p)$$

$$(X, \mathcal{C}, p)$$

$$(X, \mathcal{C}, p)$$

$$(X, \mathcal{B}, p)$$

$$(4.1)$$

and the composite $k := \delta'_{id} \circ (\delta_{id})^+ : (X, \mathcal{C}, p) \to (X, \mathcal{B}, p)$. Then, by Theorem 4.18 and Proposition 2.12, $\mathcal{B} \leq \mathcal{C}$ if and only if k is almost surely deterministic. Moreover, in that case, the map k is the only almost surely deterministic kernel making the diagram (4.1) commute.

In particular, $\mathcal{B} \simeq \mathcal{C}$ if and only if k is an isomorphism.

In other words, \mathcal{B} is almost surely coarser than \mathcal{C} if and only if there is an almost surely deterministic kernel $(X, \mathcal{C}, p) \to (X, \mathcal{B}, p)$ commuting with the maps in (4.1). This generalizes the fact that $\mathcal{B} \subseteq \mathcal{C}$ if and only if the identity (function or kernel) $(X, \mathcal{C}) \to (X, \mathcal{B})$ is \mathcal{C} -measurable.

This also means that almost surely equivalent sub-sigma-algebras of a standard Borel space give rise to isomorphic objects of GKrn (or even of Krn, up to isomorphism). For example, the objects (X, \mathcal{B}, p) and $(X, \mathcal{I}_{e_{\mathcal{B}}}, p)$ are isomorphic. (This last fact can also be seen from the fact that both objects split the same idempotent $e_{\mathcal{B}}$.) **Remark 4.21.** In any category, every idempotent is bounded above by the identity: for every $e: X \to X$, $e \circ id = id \circ e = e$, i.e. $e \leq id$ as idempotents. In a general there may be no lower bound, but there is one in EKrn: consider the "constant random kernel" $k_p: (X, \mathcal{A}, p) \to (X, \mathcal{A}, p)$ given by

$$k_p(A|x) = p(A)$$

for all $x \in X$ and $A \in A$. This corresponds to conditional expectation with respect to the trivial sigma-algebra, i.e. *un* conditional expectation:

$$f \longmapsto \mathbb{E}[f] = \int_X f \, dp.$$

As it is well known, for every random variable f and every sub-sigma-algebra $\mathcal{B} \subseteq \mathcal{A}$,

$$\mathbb{E}\big[\mathbb{E}[f|\mathcal{B}]\big] = \mathbb{E}\big[\mathbb{E}[f]|\mathcal{B}\big] = \mathbb{E}[f|\mathcal{B}].$$

Therefore, for all idempotents $e, e_p \circ e = e \circ e_p = e_p$, i.e. $e_p \leq e$.

4.3 Filtrations and martingales

Let's now turn to martingales. We first need to introduce filtrations, which in our categorical setting correspond to filtered diagrams of dagger-split idempotents.

Definition 4.22. Let (X, \mathcal{A}, p) be a probability space, and let $(\mathcal{B}_{\lambda})_{\lambda \in \Lambda}$ be a net (or sequence) of sub-sigma-algebras of \mathcal{A} . We say that (\mathcal{B}_{λ}) is

- An increasing filtration if for all $\lambda \leq \mu$, $\mathcal{B}_{\lambda} \subseteq \mathcal{B}_{\mu}$;
- A decreasing filtration if for all $\lambda \leq \mu$, $\mathcal{B}_{\lambda} \supseteq \mathcal{B}_{\mu}$;
- An almost surely increasing filtration if for all $\lambda \leq \mu$, $\mathcal{B}_{\lambda} \lesssim \mathcal{B}_{\mu}$;
- An almost surely decreasing filtration if for all $\lambda \leq \mu$, $\mathcal{B}_{\lambda} \gtrsim \mathcal{B}_{\mu}$.

For sub-sigma-algebras of a standard Borel space, whenever $\mathcal{B} \subseteq \mathcal{C}$, by Corollary 4.20 we have a canonical almost surely deterministic morphism $(X, \mathcal{C}, p) \to (X, \mathcal{B}, p)$ (note the direction of the arrow). Therefore every almost surely increasing filtration gives rise to a cofiltered (projective) net of measure spaces and almost surely deterministic kernels, which in the sequential case looks as follows.

$$(X, \mathcal{B}_0, p) \longleftarrow (X, \mathcal{B}_1, p) \longleftarrow (X, \mathcal{B}_2, p) \longleftarrow \dots \longleftarrow (X, \mathcal{B}_n, p) \longleftarrow \dots$$
(4.2)

(We draw the arrow in this way to keep the filtration in the "forward" direction.) Similarly, an almost surely decreasing filtration gives rise to a filtered (inductive) net, which in the sequential case looks as follows.

$$\dots \longleftarrow (X, \mathcal{B}_n, p) \longleftarrow \dots \longleftarrow (X, \mathcal{B}_2, p) \longleftarrow (X, \mathcal{B}_1, p) \longleftarrow (X, \mathcal{B}_0, p) \tag{4.3}$$

If the filtrations are strictly increasing or decreasing, meaning not just almost surely, the morphisms can be taken to be deterministic, not just almost surely. On a standard Borel space there is essentially no difference between "increasing" and "almost surely increasing". On one hand, every increasing filtration is almost surely increasing. A weak converse is given by the following proposition. **Proposition 4.23.** Let (X, \mathcal{A}, p) be standard Borel, and let $(\mathcal{B}_{\lambda})_{\lambda \in \Lambda}$ be an almost surely increasing filtration. Then there exists an increasing (not just almost surely) filtration $(\mathcal{B}'_{\lambda})_{\lambda \in \Lambda}$ such that for all $\lambda, \mathcal{B}'_{\lambda} \simeq \mathcal{B}_{\lambda}$, and moreover for each $\lambda \leq \mu$ each of the following squares commute,

$$(X, \mathcal{B}_{\mu}, p) \longrightarrow (X, \mathcal{B}_{\lambda}, p)$$
$$\downarrow \cong \qquad \qquad \qquad \downarrow \cong$$
$$(X, \mathcal{B}'_{\mu}, p) \longrightarrow (X, \mathcal{B}'_{\lambda}, p)$$

where the maps are the ones canonically induced by Corollary 4.20. The same can be said for the decreasing case.

Proof. Take $\mathcal{B}'_{\lambda} = \mathcal{I}_{e_{\mathcal{B}_{\lambda}}}$.

We know by the results of Section 2.4 that the limit of an increasing filtration (in the subcategory of a.s. deterministic kernels, i.e. the dagger epis) is given by the supremum in the almost-surely-coarse preorder (\leq), and the colimit of a decreasing filtration is given by the infimum. However, relating the suprema and infima for the almost-surely-coarse preorder (\leq) and for the inclusion order (\subset) between sigma-algebras requires a little care.

First of all, call a sub-sigma-algebra **null-set-complete** if it contains all the null sets, or equivalently, by Corollary 4.17, if it is in the form \mathcal{I}_k for some kernel k. It is easy to see that the joins (for the inclusion order) and the meets (i.e. intersection) of null-set-complete sub-sigma-algebras are again null-set complete.

Somewhat conversely, on a standard Borel space we can turn any sub-sigma-algebra into an almost surely equivalent, null-set-complete one via the assignment $\mathcal{B} \mapsto \mathcal{I}_{e_{\mathcal{B}}}$. This assignment preserves joins (for the inclusion order). To see this, denote by \mathcal{N} the sub-sigma-algebra generated by all null sets, i.e. the one of sets $A \in \mathcal{A}$ for which either p(A) = 0 or p(A) = 1. By Corollary 4.17, we have that $\mathcal{I}_{e_{\mathcal{B}}}$ is the join of \mathcal{B} and \mathcal{N} , and so the assignment $\mathcal{B} \mapsto \mathcal{B} \vee \mathcal{N}$ preserves all joins.

However, this assignment in general does not preserve meets (intersections). For an example, consider the set $\{x, y, z\}$ with the discrete sigma-algebra \mathcal{A} and the measure p(x) = p(z) = 1/2, p(y) = 0. Consider now the sub-sigma-algebras \mathcal{B} and \mathcal{C} generated by the partitions $\{\{x, y\}, \{z\}\}$ and $\{\{x\}, \{y, z\}\}$. We have that $\mathcal{B} \cap \mathcal{C}$ is the trivial sigma-algebra, and so $\mathcal{I}_{e_{\mathcal{B}} \cap \mathcal{C}} = \mathcal{N}$. On the other hand $\mathcal{B} \simeq \mathcal{C}$, so that by Corollary 4.16(iv), $\mathcal{I}_{e_{\mathcal{B}}} = \mathcal{I}_{e_{\mathcal{C}}} \cap \mathcal{I}_{e_{\mathcal{C}}} = \mathcal{A}$.

Remark 4.24. For the readers familiar with order theory, we can model the situation as follows. Denote by ΣX the set of sub-sigma algebras of \mathcal{A} , denote by $NC\Sigma X$ the set of null-set-complete sub-sigma-algebras, and denote by I the set of idempotents on (X, \mathcal{A}, p) with the idempotent order. We have respectively a Galois connection, an isomorphism, and an equivalence of preorders as in the following diagram:

$$(\Sigma X, \subseteq) \xleftarrow{\top} (NC\Sigma X, \subseteq) \xleftarrow{\cong} (I, \leq) \xleftarrow{\cong} (\Sigma X, \lesssim)$$
$$\mathcal{B} \xrightarrow{\top} \mathcal{I}_{e_{\mathcal{B}}}$$

Now the assignment $\mathcal{B} \mapsto \mathcal{I}_{e_{\mathcal{B}}}$, being a lower (or left) adjoint, preserves all joins (but not all meets).

For sequential filtrations, in any case, the necessary meets are preserved, as the following proposition shows.

Proposition 4.25. Let (X, \mathcal{A}, p) be standard Borel. The assignment $\mathcal{B} \mapsto \mathcal{I}_{e_{\mathcal{B}}}$ preserves intersections of decreasing filtrations, i.e. sequential infima for the inclusion relation (\subseteq) .

Proof. Let $(B_n)_{n=0}^{\infty}$ be a decreasing filtration. We have to prove that

$$\bigcap_{n} (\mathcal{I}_{e_{\mathcal{B}_{n}}}) = \mathcal{I}_{e_{(\bigcap_{n} \mathcal{B}_{n})}}$$

Since the assignment $\mathcal{B} \mapsto \mathcal{I}_{e_{\mathcal{B}}}$ is monotone, we already have that $\mathcal{I}_{e_{(\bigcap_{n} \mathcal{B}_{n})}} \subseteq \bigcap_{n} (\mathcal{I}_{e_{\mathcal{B}_{n}}})$. We have to prove the other direction.

So let $A \in \bigcap_n (\mathcal{I}_{e_{\mathcal{B}_n}})$. This means that for every $n, A \in \mathcal{I}_{e_{\mathcal{B}_n}}$, which means (using Lemma 4.13) that A is p-almost surely in each \mathcal{B}_n . In other words, for each n there exists $B_n \in \mathcal{B}_n$ such that $p(A \triangle B_n) = 0$. Now since for each n we have that $\mathcal{B}_{n+1} \subseteq \mathcal{B}_n$, we have that for every n and for every $k \ge n$, $B_k \in \mathcal{B}_n$ as well. Define now for each n the set

$$B'_n \coloneqq \bigcup_{k \ge n} B_k$$

For every n we have that

- $B'_n \in \mathcal{B}_n;$
- $B'_n \supseteq B'_{n+1};$
- $p(B'_n \bigtriangleup A) = 0.$

The first item implies, using again that $\mathcal{B}_{n+1} \subseteq \mathcal{B}_n$, that for every n and for every $k \ge n$, $B'_k \in \mathcal{B}_n$ as well. The second item says exactly that the sequence (B'_n) is nonincreasing. With these two remarks in mind, define now the set

$$B \coloneqq \bigcap_{k=0}^{\infty} B'_k.$$

Note that since the sequence (B'_n) is nonincreasing, we could define B equivalently as

$$B \coloneqq \bigcap_{k \ge n} B'_k$$

for any *n*. Therefore $B \in \mathcal{B}_n$ for every *n*, and hence $B \in \bigcap_n \mathcal{B}_n$. Moreover, once again by construction, $p(A \triangle B) = 0$. Therefore *A* is almost surely in $\bigcap_n \mathcal{B}_n$, which means, using again Lemma 4.13, that $A \in \mathcal{I}_{e_{(\bigcap_n \mathcal{B}_n)}}$.

For non-sequential filtrations (nets), instead, one has to be careful, since the infimum in the idempotent order is given by the intersection of the *null-set-completion* $\mathcal{I}_{e(\bigcap_{n} \mathcal{B}_{\lambda})}$ of the \mathcal{B}_{λ} . This difficulty is avoided if one starts with a decreasing net of null-set-complete sub-sigma-algebras. In what follows, to avoid ambiguity, we will denote by $\bigcap_{\lambda} \mathcal{B}_{\lambda}$ the ordinary intersection of sigma-algebras, and by $\bigwedge_{\lambda} \mathcal{B}_{\lambda}$ the intersection of the null-set completions (giving the meet in the idempotent order).

Let's now use the ideas of Section 2.4. Given a standard Borel space (X, \mathcal{A}, p) , the category $DSI(X, \mathcal{A}, p)$ has (equivalently) as objects, sub-sigma-algebras $\mathcal{B} \subseteq \mathcal{A}$ up to almost sure equality, together with the maps

$$(X, \mathcal{A}, p) \xrightarrow[\iota]{\pi} (X, \mathcal{B}, p),$$

where $\pi = \delta_{id}$ and $\iota = \pi^+$ is the disintegration. As morphisms, corresponding to (almost sure) inclusions $\mathcal{B}_1 \leq \mathcal{B}_2$, it has almost surely deterministic kernels $\pi_{2,1} : (X, \mathcal{B}_2, p) \to (X, \mathcal{B}_1, p)$ making the upper and lower triangles in the following diagram commute.



Equivalently, it can be seen as the slice category $\operatorname{Krn}_{\operatorname{dm}}/(X, \mathcal{A}, p)$, or the opposite of $(X, \mathcal{A}, p)/\operatorname{Krn}_{\operatorname{de}}$.

Corollary 4.26. Let (X, \mathcal{A}, p) be standard Borel. In the category $DSI(X, \mathcal{A}, p)$ (equivalently, $Krn_{dm}/(X, \mathcal{A}, p)$ and $(X, (\mathcal{A}, p)/Krn_{de})^{op}$):

- The limit of an increasing filtration (B_λ) on (X, A, p) exists, and it is given by the join sigmaalgebra V_λ B_λ, i.e. the smallest (coarsest) sigma-algebra containing all the B_λ;
- The colimit of a decreasing filtration (\mathcal{B}_{λ}) on (X, \mathcal{A}, p) exists, and is given by the meet (intersection of the null-set completions) sigma-algebra $\bigwedge_{\lambda} \mathcal{B}_{\lambda}$.

Let's now apply the random variable functors to filtrations. We focus on increasing filtrations, decreasing filtrations are analogous. If we apply the L^1 (or L^n) functor to an increasing filtration, which in the sequential case looks as follows,

$$(X, \mathcal{B}_0, p) \xleftarrow{\pi_{1,0}} (X, \mathcal{B}_1, p) \xleftarrow{\pi_{2,1}} (X, \mathcal{B}_2, p) \xleftarrow{\pi_{3,2}} \dots$$

where we denoted by $\pi_{n,n-1}$ the canonical map $(X, \mathcal{B}_n, p) \to (X, \mathcal{B}_{n-1}, p)$, we get a net with the arrows reversed:

$$L^1(X, \mathcal{B}_0, p) \xrightarrow{(\pi_{1,0})^*} L^1(X, \mathcal{B}_1, p) \xrightarrow{(\pi_{2,1})^*} L^1(X, \mathcal{B}_2, p) \xrightarrow{(\pi_{3,2})^*} \dots$$

For example, the map $(\pi_{2,1})^* : L^1(X, \mathcal{B}_1, p) \to L^1(X, \mathcal{B}_2, p)$ includes the \mathcal{B}_1 -measurable random variables into the space of \mathcal{B}_2 -measurable ones, which is possible since \mathcal{B}_2 is finer or equal to \mathcal{B}_1 . (Note that, technically, $L^1(X, \mathcal{B}_1, p)$ is not exactly a subset of $L^1(X, \mathcal{B}_2, p)$, since the latter may have more measure zero sets, but it defines a subset canonically.)

We can also look at the disintegrations $\pi_{n,n-1}^+$: $(X, \mathcal{B}_{n-1}, p) \to (X, \mathcal{B}_n, p)$, fitting into the original diagram as follows.

$$(X, \mathcal{B}_0, p) \xrightarrow[\pi_{1,0}^+]{\pi_{1,0}^+} (X, \mathcal{B}_1, p) \xrightarrow[\pi_{2,1}^+]{\pi_{1,2}^+} (X, \mathcal{B}_2, p) \xrightarrow[\pi_{3,2}^+]{\pi_{3,2}^+} \dots$$

Applying the L^1 (or L^n) functor we get the following maps.

$$L^{1}(X, \mathcal{B}_{0}, p) \xleftarrow{(\pi_{1,0}^{+})^{*}}_{(\pi_{1,0})^{*}} L^{1}(X, \mathcal{B}_{1}, p) \xleftarrow{(\pi_{1,2}^{+})^{*}}_{(\pi_{2,1})^{*}} L^{1}(X, \mathcal{B}_{2}, p) \xleftarrow{(\pi_{3,2}^{+})^{*}}_{(\pi_{3,2})^{*}} \dots$$

The new maps encode conditional expectations: for example, the map $(\pi_{2,1}^+)^* : L^1(X, \mathcal{B}_2, p) \to L^1(X, \mathcal{B}_1, p)$ takes the conditional expectation of \mathcal{B}_2 -measurable random variables given \mathcal{B}_1 (which is possible since \mathcal{B}_2 is finer or equal to \mathcal{B}_1). Focusing on these maps,

$$L^1(X, \mathcal{B}_0, p) \xleftarrow{(\pi_{1,0}^+)^*} L^1(X, \mathcal{B}_1, p) \xleftarrow{(\pi_{1,2}^+)^*} L^1(X, \mathcal{B}_2, p) \xleftarrow{(\pi_{3,2}^+)^*} \dots$$

we can encode a *martingale* as a collection of random variables, one for every space, compatible with the arrows:

Definition 4.27. Let (X, \mathcal{A}, p) be a standard Borel probability space.

A (forward) martingale on (X, \mathcal{A}, p) consists of

- An increasing filtration $(\mathcal{B}_{\lambda})_{\lambda \in \Lambda}$ on (X, \mathcal{A}, p) ;
- For each $\lambda \in \Lambda$, a random variable $f_{\lambda} \in L^1(X, \mathcal{B}_{\lambda}, p)$,

such that for all $\lambda \leq \mu$, f_{λ} is a conditional expectation of f_{μ} given \mathcal{B}_{λ} . A **backward martingale** on (X, \mathcal{A}, p) consists of

- A decreasing filtration $(\mathcal{B}_{\lambda})_{\lambda \in \Lambda}$ on (X, \mathcal{A}, p) ;
- For each $\lambda \in \Lambda$, a random variable $f_{\lambda} \in L^1(X, \mathcal{B}_{\lambda}, p)$,

such that for all $\lambda \leq \mu$, f_{μ} is a conditional expectation of f_{λ} given \mathcal{B}_{μ} .

Recall that we can encode a random variable as a bounded linear map $\mathbb{R} \to (X, \mathcal{A}, p)$. Similarly, we can encode a martingale as a commutative diagram, which in the sequential case looks as follows,

$$L^{1}(X, \mathcal{B}_{0}, p) \xleftarrow{f_{0}} L^{1}(X, \mathcal{B}_{1}, p) \xleftarrow{f_{1}} L^{1}(X, \mathcal{B}_{1}, p) \xleftarrow{f_{2}} L^{1}(X, \mathcal{B}_{2}, p) \xleftarrow{f_{1}} (\pi^{+}_{3,2})^{*} \cdots$$

and a backward martingale is described analogously. Note that the f_n commute with the maps $(\pi_{n,n-1}^+)^*$ (since they are conditional expectations of each other), but *not* with the maps $(\pi_{n,n-1})^*$ in general: that would mean that all the f_n are almost surely equal, and almost surely \mathcal{B}_i -measurable for all i.

Note also that, by the results above, we can equivalently work with *almost surely* increasing (resp. decreasing) filtrations (but in the decreasing, non-sequential case, taking the infimum will require some care).

Let's now consider the elements \mathcal{B}_{λ} of the filtration as sub-sigma-algebras of \mathcal{A} . We have canonical maps as follows.

$$(X, \mathcal{A}, p) \xrightarrow[\pi_{\lambda}]{\pi_{\lambda}} (X, \mathcal{B}_{\lambda}, p)$$

where $\pi_{\lambda} : (X, \mathcal{A}, p) \to (X, \mathcal{B}_{\lambda}, p)$ is the kernel induced by the set-theoretic identity, and π_{λ}^+ is the corresponding disintegration. The pair $(\pi_{\lambda}, \pi_{\lambda}^+)$ forms a (dagger-)split idempotent, and those are preserved by all functors. Therefore we have again a split idempotent

$$L^1(X, \mathcal{A}, p) \xleftarrow{(\pi_\lambda)^*}{(\pi_\lambda^+)^*} L^1(X, \mathcal{B}_\lambda, p),$$

where the map $(\pi_{\lambda})^* : L^1(X, \mathcal{B}_{\lambda}, p) \to L^1(X, \mathcal{A}, p)$ is (almost) an inclusion, and the map $(\pi_{\lambda}^+)^* : L^1(X, \mathcal{A}, p) \to L^1(X, \mathcal{B}_{\lambda}, p)$ forms conditional expectations.

If we consider the functor L^2 : EKrn \rightarrow Hilb, which preserves the dagger structure, we have that the maps $(\pi_{\lambda})^* : L^2(X, \mathcal{B}_{\lambda}, p) \rightarrow L^2(X, \mathcal{A}, p)$ and $(\pi_{\lambda}^+)^* : L^2(X, \mathcal{A}, p) \rightarrow L^2(X, \mathcal{B}_{\lambda}, p)$ form moreover a *dagger*-split idempotent, just like π_{λ} and π_{λ}^+ .

We have seen above that, in the order of split idempotents of (X, \mathcal{A}, p) , the join sigma-algebra is the supremum of an increasing filtration, and the meet (intersection of the null-set completions) sigma-algebra is the infimum of a decreasing filtration. It is natural to ask if the L^n functors preserve these suprema and infima, and the answer is affirmative.

Proposition 4.28. Let (X, \mathcal{A}, p) be a standard Borel probability space, and let $n \in \mathbb{N}$.

- (i) Given an increasing filtration $(\mathcal{B}_{\lambda})_{\lambda \in \Lambda}$ with supremum $\mathcal{B}_{\infty} = \bigvee_{\lambda \in \Lambda} \mathcal{B}_{\lambda}$, the space $L^{n}(X, \mathcal{B}_{\infty}, p)$ is the supremum of the $L^{n}(X, \mathcal{B}_{\lambda}, p)$ in the order of split idempotents on $L^{n}(X, \mathcal{A}, p)$ (equivalently, the colimit in $SI(L^{n}(X, \mathcal{A}, p))$).
- (ii) Given a decreasing filtration $(\mathcal{C}_{\lambda})_{\lambda \in \Lambda}$ with infimum $\mathcal{C}_{\infty} = \bigwedge_{\lambda \in \Lambda} \mathcal{C}_{\lambda}$, the space $L^{n}(X, \mathcal{C}_{\infty}, p)$ is the infimum of the $L^{n}(X, \mathcal{C}_{\lambda}, p)$ in the order of split idempotents on $L^{n}(X, \mathcal{A}, p)$ (equivalently, the limit in $SI(L^{n}(X, \mathcal{A}, p))$).

Proof. First of all, without loss of generality, by possibly replacing \mathcal{B}_{λ} and \mathcal{B}_{∞} with their null-set completions, we can assume that \mathcal{B}_{λ} - and \mathcal{B}_{∞} -measurability and almost sure measurability coincide. (Same for \mathcal{C}_{λ} and \mathcal{C}_{∞} .) Moreover, this way the infimum will simply be given by the intersection ogf sigma-algebras.

(i) Notice that for split idempotents of Banach spaces, in order for $L^n(X, \mathcal{B}_{\infty}, p)$ to be a supremum it suffices to check that it is the closure of the union of the $L^n(X, \mathcal{B}_{\lambda}, p)$, and that it admits a retract. As we know, the retract is given by the conditional expectation map $(\pi^+)^*$: $L^n(X, \mathcal{A}, p) \to L^n(X, \mathcal{B}_{\infty}, p)$, where $\pi : (X, \mathcal{A}, p) \to (X, \mathcal{B}_{\infty}, p)$ is the kernel induced by the set-theoretic identity.

To prove that $L^n(X, \mathcal{B}_{\infty}, p)$ is the closure of the union, let first of all $B \in \mathcal{B}_{\infty}$. By [VB23, Lemma 3.14], there exists a sequence $(B_i)_{i=0}^{\infty}$ of sets from the union $\bigcup_{\lambda \in \Lambda} \mathcal{B}_{\lambda}$ such that $p(B \triangle B_i) \to 0$. Therefore 1_B is in the closure of the union of the $L^n(X, \mathcal{B}_{\lambda}, p)$. The same claim can now be extended to all \mathcal{B}_{∞} -measurable function by approximating them via simple functions. Conversely, suppose f is in the closure of the union of the $L^n(X, \mathcal{B}_{\lambda}, p)$. Then there exists a sequence $(f_i)_{i=0}^{\infty}$ in $\bigcup_{\lambda \in \Lambda} L^n(X, \mathcal{B}_{\lambda}, p)$ tending to f, and all the functions f_i are also in $L^n(X, \mathcal{B}_{\infty}, p)$. But now since $L^n(X, \mathcal{B}_{\infty}, p)$ is complete, $f \in L^n(X, \mathcal{B}_{\infty}, p)$ as well. So $f \in L^n(X, \mathcal{B}_{\infty}, p)$ if and only if it is in the closure of the union of the $L^n(X, \mathcal{B}_{\lambda}, p)$.

(ii) As above, notice that for split idempotents of Banach spaces, in order for $L^n(X, \mathcal{C}_{\infty}, p)$ to be an infimum it suffices to check that it is the intersection of the $L^n(X, \mathcal{C}_{\lambda}, p)$, and that it admits a retract. Again as above, the retract is given by the conditional expectation map $(\pi^+)^* : L^n(X, \mathcal{A}, p) \to L^n(X, \mathcal{C}_{\infty}, p)$, where $\pi : (X, \mathcal{A}, p) \to (X, \mathcal{C}_{\infty}, p)$ is the kernel induced by the set-theoretic identity.

To prove that $L^n(X, \mathcal{C}_{\infty}, p)$ is the intersection, suppose that $f \in L^n(X, \mathcal{A}, p)$ is \mathcal{C}_{λ} -measurable for all λ . Then for every Borel set $B \subseteq \mathbb{R}$, $f^{-1}(B) \in \mathcal{C}_{\lambda}$ for all λ , so that $f^{-1}(C) \in \mathcal{C}_{\infty}$, and hence f is \mathcal{C}_{∞} -measurable. Conversely, if f is \mathcal{C}_{∞} -measurable, then it is \mathcal{C}_{λ} -measurable for all λ . So $f \in L^n(X, \mathcal{C}, p)$ if and only if for all $\lambda, f \in L^n(X, \mathcal{C}_{\lambda}, p)$. Corollary 4.29. By Proposition 2.21 and Proposition 2.22, we have that

- In the increasing case, $L^n(X, \mathcal{B}_{\infty}, p)$ is the colimit in $\operatorname{Ban}_{\leq 1}$ of the $L^n(X, \mathcal{B}_i, p)$ and the inclusion maps $(\pi_{i+1,i})^* : L^n(X, \mathcal{B}_i, p) \to L^n(X, \mathcal{B}_{i+1}, p)$.
- In the decreasing case, $L^n(X, \mathcal{B}_{\infty}, p)$ is the limit in $\operatorname{Ban}_{\leq 1}$ of the $L^n(X, \mathcal{B}_i, p)$ and the inclusion maps $(\pi_{i,i+1})^* : L^n(X, \mathcal{B}_{i+1}, p) \to L^n(X, \mathcal{B}_i, p).$

Moreover, in the case of L^2 , which is a Hilbert space, the dagger structure gives us the following dual statements:

Corollary 4.30. By Corollary 2.23 and Corollary 2.24, we have that

- In the increasing case, $L^2(X, \mathcal{B}_{\infty}, p)$ is the limit in $\operatorname{Hilb}_{\leq 1}$ of the $L^2(X, \mathcal{B}_i, p)$ and the projection maps $(\pi_{i+1,i}^+)^* : L^2(X, \mathcal{B}_{i+1}, p) \to L^2(X, \mathcal{B}_i, p).$
- In the decreasing case, $L^2(X, \mathcal{B}_{\infty}, p)$ is the colimit in $\operatorname{Hilb}_{\leq 1}$ of the $L^2(X, \mathcal{B}_i, p)$ and the inclusion maps $(\pi_{i+1,i}^+)^* : L^2(X, \mathcal{B}_i, p) \to L^2(X, \mathcal{B}_{i+1}, p).$

The latter property says in particular that given an L^2 -bounded martingale (for example, sequential) (f_n) on a filtration (\mathcal{B}_n) , forming a cone as below,

$$L^{2}(X, \mathcal{B}_{0}, p) \xleftarrow[(\pi^{+}_{1,0})^{*}]{\mathbb{K}} L^{2}(X, \mathcal{B}_{1}, p) \xleftarrow[(\pi^{+}_{2,1})^{*}]{\mathbb{K}} \cdots \xleftarrow[(\pi^{+}_{2,1})^{*}]{\mathbb{K}} \cdots$$

there exists a unique random variable f as above making the diagram commute, i.e. a \mathcal{B}_{∞} measurable random variable f such that such that for all n, $f_n = \mathbb{E}[f|\mathcal{B}_n]$. This is in general
not the case for L^1 , in particular if the f_i are not uniformly integrable.

Example 4.31. Let X = [0, 1] with its Borel sigma-algebra. For $n \in \mathbb{N}$, let \mathcal{B}_n be the sigma-algebra generated by the inverse powers of 2, i.e. by the intervals $[0, 1/2^n], (1/n, 2/2^n], \dots, ((2^n - 1)/2^n, 1]$. The functions f_i given by

$$f_n(x) := \begin{cases} 2^n & x \le 1/2^n \\ 0 & x > 1/2^n \end{cases}$$

form a martingale on $(\mathcal{B}_{\backslash})_{n \in \mathbb{N}}$, and we have that for all n,

$$||f_n||_{L^1} = \int_{[0,1]} |f_n(x)| \, dx = \int_{[0,1/2^n]} 2^n \, dx = 1.$$

However there does not exist an f such that for all n, $\mathbb{E}[f|\mathcal{B}_n] = f_n$.

What we always get, by having a colimit in $SI(L^1(X, \mathcal{A}, p))$, is that if we have a random variable $f \in L^1(X, \mathcal{A}, p)$, and its conditional expectations f_n with respects to a filtration $(\mathcal{B}_n)_{n=0}^{\infty}$, then the conditional expectation of f given $\mathcal{B}_{\infty} = \bigvee_n \mathcal{B}_n$ is a \mathcal{B} -measurable random variable f_{∞} such that

for all $n, f_n = \mathbb{E}[f_\infty | \mathcal{B}_n]$. In diagrams, given $f : \mathbb{R} \to L^1(X, \mathcal{A}, p)$,



the conditional expectations f_n can be expressed as $(\pi_n^+)^* f \in L^1(X, \mathcal{B}_n, p)$, and the map f_∞ can be written as $(\pi_\infty^+)^* f \in L^1(X, \mathcal{B}_\infty, p)$. By further postcomposing with the maps $(\pi_n)^*$ and $(\pi_\infty)^*$, one can of course consider all these maps again as random variables on (X, \mathcal{A}, p) . Analogous remarks can be made for decreasing filtrations.

The martingale convergence theorem says that moreover, this map f_{∞} is the limit (in the topological sense) of the f_n . We can state and prove this categorically as well, but in order to talk about topological convergence we need an additional ingredient: a topological enrichment.

5 Topological enrichment

We now equip both our categories of kernels and our categories of vector spaces with a topological enrichment. This amounts to equipping each hom-set C(X, Y) with a topology which makes the composition of morphisms continuous, and it allows us to talk about convergence of morphisms. When morphisms are functions, in our examples this convergence is the pointwise one. More details on this enrichment are given in Appendix A.

5.1 The enrichments of Ban and Hilb

Let X and Y be Banach spaces. We equip the set of bounded linear maps Ban(X, Y) with the topology of pointwise convergence. That is, we say that a net $(f_{\lambda} : X \to Y)_{\lambda \in \Lambda}$ tends to $f : X \to Y$ if and only if for all $x \in X$, $f_{\lambda}(x) \to f(x)$ in Y, or equivalently,

$$||f(x) - f_{\lambda}(x)||_{Y} \to 0$$

Note that we are not requiring that the convergence be uniform in x.

A subbasis of this topology is given by the sets

$$\{f : \|f(x) - y\|_Y < r\}$$

for $x \in X$, $y \in Y$, and r > 0.

In the case of X = Y, this topology is sometimes called the strong operator topology.

Proposition 5.1. This choice of topology for each pair of Banach spaces X and Y makes Ban enriched in Top, in the sense of Appendix A.

The statement follows immediately since we can consider Ban a subcategory of Top (with the corresponding enrichment). We nevertheless provide a direct proof for clarity.

Proof. We have to prove that for all Banach spaces X, Y and Z, the composition map

$$\begin{array}{ccc} \operatorname{Ban}(X,Y) \times \operatorname{Ban}(Y,Z) & \longrightarrow & \operatorname{Ban}(X,Z) \\ & & (f,g) & \longmapsto & g \circ f \end{array}$$

is separately continuous in both variables.

- (i) To prove continuity in f (of the postcomposition with g), suppose that $f_{\lambda} \to f : X \to Y$. This means that for all $x \in X$, $f_{\lambda}(x) \to f(x)$ in Y. Since g is continuous, then $g(f_{\lambda}(x)) \to g(f(x))$. This holds for all x, which means exactly that $g \circ f_{\lambda} \to g \circ f$.
- (ii) To prove continuity in g (of the precomposition with f), suppose that $g_{\lambda} \to g: Y \to Z$. This means that for all $y \in Y$, $g_{\lambda}(y) \to g(y)$ in Z. Setting y = f(x), we see that for all $x \in X$, $g_{\lambda}(f(x)) \to g(f(x))$, i.e. $g_{\lambda} \circ f \to g \circ f$.

Let's now turn to Hilbert spaces. One can consider Hilb as a subcategory of Ban, and inherit the enrichment accordingly. However, this enrichment is not compatible with the dagger structure, i.e. it does not respect the symmetry: if $f_{\lambda} \to f : X \to Y$, it is not necessarily true that $f_{\lambda}^+ \to f^+$: $Y \to X$. Therefore we define the following topology: we say that a net $(f_{\lambda} : X \to Y)_{\lambda=0}^{\infty}$ tends to $f: X \to Y$ if and only if

- (i) for all $x \in X$, $f_{\lambda}(x) \to f(x)$ in Y; and
- (ii) for all $y \in Y$, $f_{\lambda}^+(y) \to f^+(y)$ in X.

Once again, we do not require that the convergence be uniform in x. A subbasis of this topology is given by the sets

$$\{f : \|f(x) - y\|_Y < r, \|f^+(y) - x\|_X < r\}$$

for $x \in X$, $y \in Y$, and r > 0.

In the case of X = Y, this topology is sometimes called the strong-* operator topology. Note that for self-adjoint operators, this topology coincide with the strong operator topology.

Proposition 5.2. This choice of topology for each pair of Banach spaces X and Y makes Hilb enriched in Top, in the sense of Appendix A.

Proof. We have to prove that for all Banach spaces X, Y and Z, the composition map

$$\operatorname{Hilb}(X,Y) \times \operatorname{Hilb}(Y,Z) \longrightarrow \operatorname{Hilb}(X,Z)$$
$$(f,g) \longmapsto g \circ f$$

is separately continuous in both variables.

To prove continuity in f (of the postcomposition with g), suppose that $f_{\lambda} \to f : X \to Y$. This means that for all $x \in X$, $f_{\lambda}(x) \to f(x)$ in Y, and for all $y \in Y$, $f_{\lambda}^+(y) \to f^+(y)$ in X. Now since g is continuous, then $g(f_{\lambda}(x)) \to g(f(x))$ for all x. Moreover, setting $y = g^+(z)$, we have that for all $z \in Z$, $f_{\lambda}^+(g^+(z)) \to f^+(g^+(z))$, i.e. $(g \circ f_{\lambda})^+(z) = (g \circ f)^+(z)$. Therefore $g \circ f_{\lambda} \to g \circ f$.

41

Continuity in q is completely analogous, and dual.

This choice of topology makes the functor $(-)^+$: Hilb \rightarrow Hilb an equivalence of Top-enriched categories. Therefore we have a *topologically enriched dagger category*.

5.2 The enrichment of Krn and EKrn

A Markov kernel $(X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$, intuitively, is at the same time a measurable function on X, and a probability measure on Y. Because of this, we define a notion of convergence which is like the L^1 convergence of measures on X, and like the setwise convergence of probability measure on Y. More in detail, consider probability spaces (X, \mathcal{A}, p) and (Y, \mathcal{B}, q) , and let $(k_{\lambda} : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q))_{\lambda \in \Lambda}$ be a sequence of measure-preserving kernels (up to a.s. equality).

(i) We say that k_λ → k in the **one-sided topology** if and only if for every every measurable set B ∈ B,

$$\int_X \left| k(B|x) - k_\lambda(B|x) \right| p(dx) \to 0;$$

(ii) If (X, \mathcal{A}, p) and (Y, \mathcal{B}, q) are essentially standard Borel, we say that (k_{λ}) tends to k in the **two-sided topology** if and only if both $k_{\lambda} \to k$ and $k_{\lambda}^+ \to k^+$ in the one-sided topology.

Just as for Hilb, we will have that the two-sided topology makes EKrn an enriched *dagger* category. On those kernels $e: X \to X$ which are their own Bayesian inverse (for example, idempotent kernels), the one-sided topology and the two-sided topology coincide.

Theorem 5.3. Let (X, \mathcal{A}, p) and (Y, \mathcal{B}, q) be probability spaces, and let $(k_{\lambda} : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q))_{\lambda \in \Lambda}$ be a sequence of measure-preserving kernels (up to a.s. equality). Then the k_{λ} tend to a kernel k in the one-sided topology if and only if for any (hence all) n (including $n = \infty$), the maps $(k_{\lambda})^* : L^n(Y, \mathcal{B}, q) \to L^n(Y, \mathcal{B}, q)$ tend to k^* .

In other words, the map

$$\operatorname{EKrn}((X, \mathcal{A}, p), (Y, \mathcal{B}, q)) \longrightarrow \operatorname{Ban}(L^{n}(Y, \mathcal{B}, q), L^{n}(X, \mathcal{A}, p))$$
$$k \longmapsto k^{*}$$

induces a homeomorphism onto its image.

Let's prove the theorem using the following auxiliary statement.

Lemma 5.4. Let X and Y be metric spaces, and let $D \subseteq X$ be a dense subset. Let $f : X \to Y$ be 1-Lipschitz, and $(f_{\lambda} : X \to Y)$ be a net of 1-Lipschitz functions. Then $f_{\lambda} \to f$ pointwise if and only if for all $d \in D$, $f_{\lambda}(d) \to f(d)$.

Proof of Lemma 5.4. Clearly, if the f_{λ} tend to f pointwise, so do their restrictions to D.

Conversely, suppose that for all $d \in D$, $f_{\lambda}(d) \to f(d)$. Now let $x \in X$. By density we can find a sequence (x_i) in D tending to x. For every $\varepsilon > 0$ there exists $I_{\varepsilon} \in \mathbb{N}$ such that for all $i \geq I_{\varepsilon}$, $d(x, x_i) < \varepsilon$. Moreover, for all $i \in \mathbb{N}$, since $x_i \in D$ we have that $f_{\lambda}(x_i) \to f(x_i)$. So for all $\varepsilon > 0$ and for all $i \in \mathbb{N}$ we can find $\lambda_{\varepsilon,i}$ such that for all $\mu \geq \lambda_{\varepsilon,i}$, $d(f_{\mu}(x_i), f(x_i)) < \varepsilon$. Given $\varepsilon > 0$, take now $i \geq I_{\varepsilon}$ and $\mu \geq \lambda_{\varepsilon,i}$. Since f and f_{μ} are 1-Lipschitz,

$$d(f_{\mu}(x), f(x)) \leq d(f_{\mu}(x), f_{\mu}(x_{i})) + d(f_{\mu}(x_{i}), f(x_{i})) + d(f(x_{i}), f(x))$$

$$\leq d(x, x_{i}) + d(f_{\mu}(x_{i}), f(x_{i})) + d(x_{i}, x)$$

 $< 3\varepsilon$.

Therefore $f_{\lambda}(x) \to f(x)$.

Proof of Theorem 5.3. We can restate the theorem as the fact that the following conditions are equivalent for all n:

(i) For all $B \in \mathcal{B}$,

$$\int_X \left| k(B|x) - k_\lambda(B|x) \right| p(dx) \to 0;$$

(ii) For all $g \in L^n(Y, \mathcal{B}, q)$,

$$\|k^*g - k^*_\lambda g\|_{L^n} \to 0.$$

Since the maps k^* and k^*_{λ} are 1-Lipschitz (Proposition 3.6), using Lemma 5.4 we can equivalently test condition (ii) on simple functions, i.e. those in the form

$$g(x) = \sum_{i=1}^k \alpha_i \, \mathbf{1}_{B_i}(x),$$

where $\alpha_i \in \mathbb{R}$ and $B_i \in \mathcal{B}$, which are dense in L^n . We can then equivalently rewrite the integral in (ii) as

$$\left\|\sum_{i=1}^{k} \alpha_k \left(k(B_i|x) - k_\lambda(B_i|x)\right)\right\|_{L^n} \leq \sum_{i=1}^{k} |\alpha_k| \left\|\left(k(B_i|x) - k_\lambda(B_i|x)\right)\right\|_{L^n},$$

and so we can rewrite condition (ii) equivalently as

(iii) For all $B \in \mathcal{B}$,

$$\left\| \left(k(B|x) - k_{\lambda}(B|x) \right) \right\|_{L^{n}} \to 0.$$

With this in mind, let's distinguish the few cases.

- For n = 1, conditions (i) and (iii) are exactly the same.
- For $0 < n < \infty$, notice that for all $f \in L^n(X, \mathcal{A}, p)$,

$$(||f||_{L^{n}})^{n} = \int_{X} |f|^{n} dp$$

= $\int_{X} |f| |f|^{n-1} dp$
 $\leq \int_{X} |f| \operatorname{ess\,sup} |f|^{n-1} dp$
= $||f||_{L^{1}} ||f||_{L^{\infty}}^{n-1}.$

Setting now $f = k(B|x) - k_{\lambda}(B|x)$, so that $||f||_{L^{\infty}} \leq 1$, we have that $(||f||_{L^{n}})^{n} \leq (||f||_{L^{1}})$, and so $(i) \Rightarrow (ii)$ The converse statement $(iii) \Rightarrow (i)$ follow by Jensen's inequality.

• The $n = \infty$ case follows by letting $n \to \infty$ in the case above.

Corollary 5.5. Let (X, \mathcal{A}, p) and (Y, \mathcal{B}, q) be probability spaces, and consider a net of measurepreserving kernels $(k_{\lambda} : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q))_{\lambda \in \Lambda}$ (up to a.s. equality). Then the k_{λ} tend to a kernel k in the two-sided topology if and only the maps $(k_{\lambda})^* : L^2(Y, \mathcal{B}, q) \to L^2(X, \mathcal{A}, p)$ tend to k^* in the topology of Hilb.

Proof. By Theorem 5.3, we have that $k_{\lambda} \to k$ in the one-sided topology if and only if $(k_{\lambda})^*$: $L^2(Y, \mathcal{B}, q) \to L^2(X, \mathcal{A}, p)$ tends to k^* pointwise. Similarly, $k_{\lambda}^+ \to k^+$ in the one-sided topology if and only if $(k_{\lambda}^+)^* : L^2(X, \mathcal{A}, p) \to L^2(Y, \mathcal{B}, q)$ tends to $(k^+)^*$ pointwise. Now $k_{\lambda} \to k$ in the two-sided topology if and only if $k_{\lambda} \to k$ and $k_{\lambda}^+ \to k^+$ in the one-sided topology, which happens if and only if $(k_{\lambda})^* \to k^*$ and $(k_{\lambda}^+)^* \to (k^+)^*$ pointwise. The latter condition is precisely convergence in the topology of Hilb.

Proposition 5.6. The one-sided and two-sided topologies enrich GKrn in Top, in the sense of Appendix A.

Proof. Let's first prove the assert for the one-sided topology. We have to prove that for all probability spaces (X, \mathcal{A}, p) , (Y, \mathcal{B}, q) and (Z, \mathcal{C}, r) , the composition map

$$\operatorname{GKrn}((X, \mathcal{A}, p), (Y, \mathcal{B}, q)) \times \operatorname{GKrn}((Y, \mathcal{B}, q), (Z, \mathcal{C}, r)) \longrightarrow \operatorname{GKrn}((X, \mathcal{A}, p), (Z, \mathcal{C}, r))$$
$$(k, h) \longmapsto h \circ k$$

is separately continuous in both variables.

(i) To prove continuity in k (of the postcomposition with h), suppose that $k_{\lambda} \to k$. Then for all $C \in \mathcal{C}$,

$$\int_{X} \left| (h \circ k)(C|x) - (h \circ k_{\lambda})(C|x) \right| p(dx) = \int_{X} \left| \int_{Y} h(C|y) \, k(dy|x) - \int_{Y} h(C|y) \, k_{\lambda}(dy|x) \right| p(dx) \\ = \|k^* h_C - k^*_{\lambda} h_C\|_{L^1}$$

where $h_C(y) = h(C|y)$. By Theorem 5.3, we know that $||k^*h_C - k^*_{\lambda}h_C||_{L^1} \to 0$. This holds for all $C \in \mathcal{C}$, and so $h \circ k_{\lambda} \to h \circ k$.

(ii) To prove continuity in h (of the precomposition with k), suppose that $h_{\lambda} \to h$, meaning that for all $C \in \mathcal{C}$,

$$\int_{Y} \left| h(C|y) - h_{\lambda}(C|y) \right| q(dy) \to 0.$$

Now for every $C \in \mathcal{C}$, by Jensen's inequality and the fact that k is measure-preserving,

$$\begin{split} \int_X \left| (h \circ k)(C|x) - (h_\lambda \circ k)(C|x) \right| p(dx) &= \int_X \left| \int_Y h(C|y) \, k(dy|x) - \int_Y h_\lambda(C|y) \, k(dy|x) \right| p(dx) \\ &\leq \int_X \int_Y \left| h(C|y) - h_\lambda(C|y) \right| k(dy|x) \, p(dx) \\ &= \int_Y \left| h(C|y) - h_\lambda(C|y) \right| q(dy) \to 0. \end{split}$$

Therefore $h_{\lambda} \circ k \to h \circ k$.

Let's now turn to the two-sided topology. Expressing the two-sided topology in terms of the one-sided one, we equivalently have to prove that

(iii) If $k_{\lambda} \to k$ and $k_{\lambda}^+ \to k^+$, then $h \circ k_{\lambda} \to h \circ k$ and $k_{\lambda}^+ \circ h^+ \to k^+ \circ h^+$;

(iv) If $h_{\lambda} \to h$ and $h_{\lambda}^+ \to h^+$, then $h_{\lambda} \circ k \to h \circ k$ and $k^+ \circ h_{\lambda}^+ \to k^+ \circ h^+$.

(All the convergences above are meant in the one-sided topology). To prove (iii), note that if $k_{\lambda} \to k$, then $h \circ k_{\lambda} \to h \circ k$ by (i). Similarly, if $k_{\lambda}^+ \to k^+$, then $k_{\lambda}^+ \circ h^+ \to k^+ \circ h^+$ by (ii) (after renaming). Similarly, to prove (iv), note that if $h_{\lambda} \to h$, then $h_{\lambda} \circ k \to h \circ k$ by (ii), and if $h_{\lambda}^+ \to h^+$, then $k^+ \circ h_{\lambda}^+ \to k^+ \circ h^+$ by (i).

In particular, this way EKrn is an enriched dagger category, like Hilb.

Corollary 5.7. The following functors are enriched:

- $L^n: \operatorname{Krn}^{\operatorname{op}} \to \operatorname{Ban}$, where Krn has either the one-sided or two-sided topology;
- $L^2: \operatorname{Krn}^{\operatorname{op}} \to \operatorname{Hilb}$, where Krn has the two-sided topology;
- Lⁿ (-)⁺: Krn → Ban and L² (-)⁺: Krn → Hilb, where Krn has the two-sided topology. These are the (covariant) functors analogous to the RV functors of [VB23].

Proposition 5.8. The composition of 1-Lipschitz maps between metric spaces is jointly continuous for the pointwise order.

Proof. Let X, Y and Z be metric spaces. Suppose that (f_{λ}) is a net of 1-Lipschitz functions $X \to Y$ tending pointwise to f, and that (g_{λ}) is a net of 1-Lipschitz functions $Y \to Z$ tending pointwise to g. Then for every $x \in X$, for every $\varepsilon > 0$ we can find $\lambda \in \Lambda$ such that for all $\mu \geq \lambda$, both $d(f(x), f_{\mu}(x))$ and $d(g(f(x)), g_{\mu}(f(x)))$ are less than ε . Therefore

$$d(g(f(x)), g_{\mu}(f_{\mu}(x))) \leq d(g(f(x)), g_{\mu}(f(x)) + d(g_{\mu}(f(x)), g_{\mu}(f_{\mu}(x)))$$

$$\leq d(g(f(x)), g_{\mu}(f(x)) + d(f(x), f_{\mu}(x)))$$

$$< 2\varepsilon.$$

Hence $g_{\lambda} \circ f_{\lambda} \to g \circ f$.

Corollary 5.9. The composition in GKrn is jointly continuous, not just separately continuous.

5.3 The subspace of idempotent kernels

Let's now restrict to the case of almost surely idempotent kernels.

Proposition 5.10. Let (X, \mathcal{A}, p) be a probability space, and let (e_{λ}) be a net of a.s. idempotent kernels on (X, \mathcal{A}, p) tending to a kernel e. Then e is a.s. idempotent as well.

In other words, almost surely idempotent kernels form a closed subset of $\operatorname{GKrn}((X, \mathcal{A}, p), (X, \mathcal{A}, p))$.

Proof. Since composition in GKrn is jointly continuous (Corollary 5.9),

$$e \circ e = \lim_{\lambda} (e_{\lambda} \circ e_{\lambda}) = \lim_{\lambda} e_{\lambda} = e.$$

Therefore e is idempotent.

Now let's restrict to the almost surely idempotents on an essentially standard Borel space (X, \mathcal{A}, p) . Notice that since for all a.s. idempotent kernels e we have $e^+ = e$ (almost surely), the one-sided and two-sided topologies coincide.

Proposition 5.11. Let (X, \mathcal{A}, p) be a standard Borel probability space, and let (e_{λ}) and (f_{λ}) be nets of a.s. idempotent kernels on (X, \mathcal{A}, p) , indexed by the same directed set, and tending respectively to kernels e and f. Suppose moreover that in the order of (split) idempotents, $e_{\lambda} \leq f_{\lambda}$ for all i. Then $e \leq f$.

In other words, the order relation of idempotents on (X, \mathcal{A}, p) is closed.

Proof. Since composition in GKrn is jointly continuous (Corollary 5.9),

$$e \circ f = \lim_{\lambda} (e_{\lambda} \circ f_{\lambda}) = \lim_{\lambda} e_{\lambda} = e_{\lambda}$$

The condition $f \circ e = e$ is proven similarly.

We have seen in Section 4 that idempotent kernels on a standard Borel probability space correspond to sub-sigma-algebras (up to almost sure equality). So equivalently, the topology we have on idempotent kernels can be seen as a topology on the set of (equivalence classes of) sub-sigmaalgebras. Concretely, we can say that a net (\mathcal{B}_{λ}) of sub-sigma algebras of (X, \mathcal{A}, p) tends to a sub-sigma-algebra $\mathcal{B} \subseteq \mathcal{A}$ if and only if for the corresponding idempotents we have $e_{\lambda} \to e$, i.e. for all measurable sets $A \in \mathcal{A}$,

$$\int_X \left| \mathbb{P}[A|\mathcal{B}](x) - \mathbb{P}[A|\mathcal{B}_{\lambda}](x) \right| p(dx) \to 0.$$

Equivalently, if for all integrable random variables f,

$$\int_{X} \left| \mathbb{E}[f|\mathcal{B}](x) - \mathbb{E}[f|\mathcal{B}_{\lambda}](x) \right| p(dx) \to 0,$$

i.e. the conditional expectations $\mathbb{E}[f|\mathcal{B}_{\lambda}]$ tend to the conditional expectation $\mathbb{E}[f|\mathcal{B}]$ in L^1 (and also in L^n , when the resulting quantities are finite).

The martingale convergence theorems, which we will prove in the next section, will tell us that

- For an increasing filtration (\mathcal{B}_{λ}) , the \mathcal{B}_{λ} tend topologically to their supremum $\bigvee_{\lambda} \mathcal{B}_{\lambda}$;
- For a decreasing filtration (\mathcal{C}_{λ}) , the \mathcal{C}_{λ} tend topologically to their infimum $\bigwedge_{\lambda} \mathcal{C}_{\lambda}$.

6 Martingale convergence

We now have all the ingredients in place to talk about martingale convergence categorically:

- A way to talk about conditional expectations (Section 4.1);
- A way to talk about filtrations and martingales (Section 4.3);
- A notion of convergence (the topological enrichment from Section 5).

We will show that the topological convergence of martingales can be interpreted as the fact that *convergence in the order implies convergence in the topology*, a phenomenon we call *Levi property*.

6.1 Levi properties

Recall Levi's theorem, which says that every bounded monotone real sequence converges topologically to its supremum. We can define the *Levi property* of a poset analogously:

Definition 6.1. Let X be a topological space equipped with a partial order. We say that

- X has the **upward Levi property** if an upper-bounded upward monotone net (i.e. a sequence (x_{λ}) such that for all $\lambda \leq \mu$, $x_{\lambda} \leq x_{\mu}$, and such that there exists y such that for all λ , $x_{\lambda} \leq y$) has a supremum, and tends to it topologically.
- X has the **downward Levi property** if a lower-bounded downward monotone net has an infimum, and tends to it topologically.

Proposition 6.2. Let X be a topological space equipped with a partial order, and suppose that the order relation is a separately closed (hence jointly closed) subset of $X \times X$. Then if an upward monotone net converges, it converges to it supremum.

Similarly, if a downward monotone net converges, it converges to its infimum.

Proof. We will prove the upward case, the downward case is analogous.

Let (x_{λ}) be an upward monotone net, and let x be its limit. Since the net is increasing, for $\lambda \leq \mu$ we have $x_{\lambda} \leq x_{\mu}$. Since the order is closed, taking the limit in μ we get that $x_{\lambda} \leq x$ for all λ . Therefore x is an upper bound. Now suppose that for some $y, x_{\lambda} \leq y$ for all λ . Then, again since the order is closed, taking the limit in λ gives $x \leq y$. Therefore x is the least upper bound.

Definition 6.3. Let C be a dagger category with a topological enrichment. We say that C has the *idempotent Levi property* if for every object X, the poset of dagger idempotents on X has the upward and downward Levi properties.

Let's spell this out in detail. Let $(e_{\lambda} : X \to X)$ be a net of dagger idempotents on X, with dagger splittings $(A_{\lambda}, \iota_{\lambda}, \pi_{\lambda})$. Suppose first that the net is increasing: for all $\lambda \leq \mu$, $e_{\lambda} \leq e_{\mu}$, i.e. $e_{\lambda} \circ e_{\mu} = e_{\mu} \circ e_{\lambda} = e_{\lambda}$. It is automatically bounded above by the identity (Remark 4.21). The upward Levi property says then that the net e_{λ} has a supremum e, and that $e_{\lambda} \to e$ topologically. Now suppose instead that the net is decreasing: for all $\lambda \leq \mu$, $e_{\lambda} \geq e_{\mu}$ (that is, $e_{\lambda} \circ e_{\mu} = e_{\mu} \circ e_{\lambda} = e_{\mu}$). Suppose moreover it is bounded below. The downward Levi property says then that the net e_{λ} has an infimum e, and that $e_{\lambda} \to e$ topologically. In EKrn, we always have a lower bound, the constant kernel given in of Remark 4.21. In Hilb, we can take the zero subspace.

In both EKrn and Hilb, we know that the suprema and infima exist. We will now show topological convergence.

6.2 The idempotent Levi property of Hilb

In the category of Hilbert spaces, categorical limits often give rise to topological limits of norms (see [DMH24] for more on this). We can formalize this idea by saying that, in Hilb, the idempotent Levi property holds.

Proposition 6.4 (upward Levi property for Hilb). Let X be a Hilbert space. Let $(A_{\lambda})_{\lambda \in \Lambda}$ and A be closed subspaces of X, with corresponding orthogonal projectors $(e_{\lambda})_{\lambda \in \Lambda}$ and e. Suppose moreover

that for all $\lambda \leq \mu$, $A_{\lambda} \subseteq A_{\mu}$, and that

$$A = \operatorname{cl}\left(\bigcup_{\lambda} A_{\lambda}\right).$$

Then $e_{\lambda} \rightarrow e$ pointwise.

Lemma 6.5. Let X be Hilbert space, let $B \subseteq X$ be a closed subset, and let $e_B : X \to X$ be the orthogonal projection onto B. Then for every $x \in X$, the projection $e_B(x)$ is the closest point on B to x, i.e.

$$d(x, e_B(x)) = \inf_{b \in B} d(x, b).$$

Proof of Lemma 6.5. Since e_B is an orthogonal projection (see Example 2.11), we have that for all $b \in B$, the vectors $x - e_B(x)$ and $b - e_B(x)$ are orthogonal. Therefore, for all $b \in B$,

$$||x - b||^{2} = ||(x - e_{B}(x)) - (b - e_{B}(x))||^{2}$$

= $||x - e_{B}(x)||^{2} + ||b - e_{B}(x)||^{2}$
 $\geq ||x - e_{B}(x)||^{2}.$

So, for all $b \in B$, $d(x, e_B(x)) = d(x, b)$.

Proof of Proposition 6.4. Given $x \in A$, note that $e(x) \in A$. Now for all λ , by Lemma 6.5 (setting $B = A_{\lambda}$ and replacing x by e(x)),

$$d(e(x), e_{\lambda}(x)) = d(e(x), e_{\lambda}(e(x)))$$
$$= \inf_{a_{\lambda} \in A_{\lambda}} d(e(x), a_{\lambda}),$$

which tends to zero since e(x) is in the closure of the A_{λ} .

Proposition 6.6 (downward Levi property for Hilb). Let X be a Hilbert space. Let $(A_{\lambda})_{\lambda \in \Lambda}$ and A be closed subspaces of X, with corresponding orthogonal projectors $(e_{\lambda})_{\lambda \in \Lambda}$ and e. Suppose moreover that for all $\lambda \leq \mu$, $A_{\lambda} \supseteq A_{\mu}$, and that

$$A = \bigcap_{\lambda} A_{\lambda}.$$

Then $e_{\lambda} \rightarrow e$ pointwise.

We prove the statement by means of the following lemma, which we learned from Matthew Di Meglio [DMH24].

Lemma 6.7. Let $(A_{\lambda})_{\lambda \in \Lambda}$ be a filtered diagram of Banach spaces and surjective, linear 1-Lipschitz maps $f_{\lambda,\mu} : A_{\lambda} \to A_{\mu}$. Then the colimit in $\operatorname{Ban}_{\leq 1}$ is the space A given by forming the quotient of any of the A_{λ} , under the seminorm

$$a \longmapsto \lim_{\mu \ge \lambda} \|\pi_{\lambda,\mu}(a)\|_{A_{\mu}},$$

with colimiting cocone formed by the quotient maps $q_{\lambda} : A_{\lambda} \to A$.

Proof of Lemma 6.7. First of all, the limit in the definition of the norm converges, since it is a nonincreasing sequence (the $\pi_{\lambda,\mu}$ are 1-Lipschitz) bounded below by zero. Being a limit of seminorms, it is itself a seminorm. Moreover, by filteredness, this seminorm does not depend on the choice of λ we start with.

Consider now a Banach space B and 1-Lipschitz linear maps $c_{\lambda} : A_{\lambda} \to B$ for all λ forming a cocone, i.e. such that for all $\mu \geq \lambda$, the outer triangle in the diagram below commutes.



Define now a map $c: A \to B$ as follows. If we construct A as a quotient of A_{λ} , its elements are equivalence classes $[a_{\lambda}]$, with $a_{\lambda} \in A_{\lambda}$, quotiented by the seminorm. Define now $c([a_{\lambda}]) = c_{\lambda}(a_{\lambda})$.

To show that this map is well defined on equivalence classes, suppose that a_{λ} and b_{λ} are in the same equivalence class, which means that

$$\lim_{\mu > \lambda} \|\pi_{\lambda,\mu}(a_{\lambda} - b_{\lambda})\|_{A_{\mu}} = 0.$$

Then for every $\mu \geq \lambda$,

$$\begin{aligned} \|c_{\lambda}(a_{\lambda}) - c_{\lambda}(b_{\lambda})\|_{B} &= \|c_{\lambda}(a_{\lambda} - b_{\lambda})\|_{B} \\ &= \|c_{\mu}(\pi_{\lambda,\mu}(a_{\lambda} - b_{\lambda}))\|_{B} \\ &\leq \|\pi_{\lambda,\mu}(a_{\lambda} - b_{\lambda})\|_{A_{\mu}}, \end{aligned}$$

and so

$$\|c_{\lambda}(a_{\lambda}) - c_{\lambda}(b_{\lambda})\|_{B} \leq \lim_{\mu \geq \lambda} \|\pi_{\lambda,\mu}(a_{\lambda} - b_{\lambda})\|_{A_{\mu}} = 0,$$

which means that $c_{\lambda}(a_{\lambda}) = c_{\lambda}(b_{\lambda})$, i.e. the map c is well defined on equivalence classes. A similar calculation shows that it is 1-Lipschitz, and linearity is easily checked.

By construction of the map c, we have $c \circ q_{\lambda} = c_{\lambda}$. Uniqueness of the map, and hence the universal property, follows from the fact that the quotient map q_{λ} is epimorphic.

To show that this quotient construction does not depend on the choice of λ , notice first of all that for all $\mu \geq \lambda$, defining c in terms of μ instead of λ gives the same quotient (since the $\mu' \geq \mu$ form a subnet which converges to the same limit). Now by directedness, given $\lambda, \lambda' \in \Lambda$ we can find $\mu \in \Lambda$ such that $\mu \geq \lambda, \lambda'$.

Proof of Proposition 6.6. First of all, by Corollary 2.24 we know that A is the colimit in $\operatorname{Hilb}_{\leq 1}$ of the spaces A_{λ} and their projections. Let $x \in X$. Notice that if $\pi : X \to A$ is the projection, we have that $\pi(e(x)) = \pi(x)$, and so, by Lemma 6.7 (taking $A_{\lambda} = X$), we have that x and e(x) are in the same equivalence class induced by the limiting norm:

$$\lim_{\mu} \|\pi_{\mu}(x - e(x))\| = 0.$$

Now since $A \subseteq A_{\mu}$, $e_{\mu} \circ e = e$, and using that the inclusions $\iota_{\mu} : A_{\mu} \to X$ are isometries,

$$\begin{split} \lim_{\mu} \|e_{\mu}(x) - e(x)\| &= \lim_{\mu} \|e_{\mu}(x) - e_{\mu}(e(x))\| \\ &= \lim_{\mu} \|e_{\mu}(x - e(x))\| \\ &= \lim_{\mu} \|\iota_{\mu}(\pi_{\mu}(x - e(x)))\| \\ &= \lim_{\mu} \|\pi_{\mu}(x - e(x))\| \\ &= 0. \end{split}$$

Note that the proof uses Corollary 2.24, which holds for Hilbert spaces, but not for general Banach spaces. Indeed, for general Banach spaces the statement fails, as the following example shows.

Example 6.8. Let $X = \ell^{\infty}(\mathbb{N})$. Consider the decreasing sequence of closed subsets,

$$A_i = \{ f \in \ell^{\infty}(\mathbb{N}) : \forall x \le i, f(x) = 0 \}.$$

These subspaces admit projectors, given by

$$e_i(f)(x) = \begin{cases} 0 & x \le i; \\ f(x) & x > i. \end{cases}$$

We have that $\bigcap_i A_i = \{0\}$, However, $e_i(f)$ does not tend to zero in general. Indeed, for f = 1 identically,

$$||e_i(f)|| = \sup_x |e_i(f)(x)| = 1$$

for all $i \in \mathbb{N}$.

Indeed, while $\bigcap_i A_i = \{0\}$, it is not true that $\{0\}$ is the colimit of the projections: by Lemma 6.7, the colimit would be given by the quotient under the seminorm $f \mapsto \lim_i ||e_i(f)||$. By the calculation above, however, for f = 1 the resulting seminorm gives 1, not zero.

6.3 The idempotent Levi property of EKrn and Levy's theorems

The idempotent Levi property holds in EKrn with our topological enrichment, and this can be considered a categorical formalization of Levy's (not Levi's) upward and downward theorems in mean.

Theorem 6.9. The category EKrn has the idempotent Levi property.

Proof. Let (X, \mathcal{A}, p) be essentially standard Borel, and let $(e_{\lambda})_{\lambda \in \Lambda}$ be a net of almost surely idempotent kernels. We know that this net is bounded above above and below by Remark 4.21. Also, by Corollary 5.5, we have that e_{λ} tends (either in the one-sided or two-sided topology) to an idempotent e if and only if the operators $e_{\lambda}^* : L^2(X, \mathcal{A}, p) \to L^2(X, \mathcal{A}, p)$ tend to e^* in Hilb.

• If the net is increasing, we know it has a supremum in EKrn, given by the join sigma-algebra. Call this supremum e_{∞} . By Proposition 4.28(i), $e_{\infty}^* : (X, \mathcal{A}, p) \to L^2(X, \mathcal{A}, p)$ is the supremum of the e_{λ}^* in Hilb as well. Now by the upward Levi property for Hilb (Proposition 6.4), e_{λ}^* tends to e_{∞}^* . Therefore $e_{\lambda} \to e_{\infty}$. • Similarly, if the net is decreasing, we know it has an infimum in EKrn, given by the meet sigma-algebra (intersection of the null-set completions). Call this infimum e_{∞} . By Proposition 4.28(ii), $e_{\infty}^* : (X, \mathcal{A}, p) \to L^2(X, \mathcal{A}, p)$ is the infimum of the e_{λ}^* in Hilb as well. Now by the downward Levi property for Hilb (Proposition 6.6), e_{λ}^* tends to e_{∞}^* . Therefore $e_{\lambda} \to e_{\infty}$.

As corollaries, by looking at the actions of idempotents on random variables, we get Levy's upward and downward theorems, for convergence in mean:

Corollary 6.10 (Levy's upward theorem). Let (X, \mathcal{A}, p) be essentially standard Borel, let $(\mathcal{B}_{\lambda})_{\lambda \in \Lambda}$ be an increasing filtration, and let $f \in L^n(X, \mathcal{A}, p)$. Denote the join $\bigvee_{\lambda \in \Lambda} \mathcal{B}_{\lambda}$ by \mathcal{B}_{∞} . Then $\mathbb{E}[f|\mathcal{B}_{\lambda}]$ converges to $\mathbb{E}[f|\mathcal{B}_{\infty}]$ in L^n .

Proof. By Theorem 4.18, the increasing filtration $(\mathcal{B}_{\lambda})_{\lambda \in \Lambda}$ induces an increasing net of dagger idempotents $(e_{\lambda} : X \to X)_{\lambda \in \Lambda}$. Since the join of the filtration is given by \mathcal{B}_{∞} , the join of the dagger idempotents is given by the dagger idempotent $e_{\infty} : X \to X$ induced by \mathcal{B}_{∞} .

Because EKrn satisfies the idempotent Levi property (Theorem 6.9), it follows that the net $(e_{\lambda})_{\lambda \in \Lambda}$ converges topologically to e_{∞} in EKrn $((X, \mathcal{A}, p), (X, \mathcal{A}, p))$, using either the one-sided or two-sided topology (they agree on idempotents). Since L^n is an enriched functor (Corollary 5.7), we see that $(e^*_{\lambda})_{\lambda \in \Lambda}$ converges to e^*_{∞} in $\text{Ban}(L^n(X, \mathcal{A}, p), (X, \mathcal{A}, p))$. This exactly means that

$$\mathbb{E}[f \mid \mathcal{B}_{\lambda}] = e_{\lambda}^* f \to e_{\infty}^* f = \mathbb{E}[f \mid B_{\infty}]$$

in L^n for every $f \in L^n(X, \mathcal{A}, p)$.

Corollary 6.11 (Levy's downward theorem). Let (X, \mathcal{A}, p) be essentially standard Borel, let $(\mathcal{B}_{\lambda})_{\lambda \in \Lambda}$ be a decreasing filtration, and let $f \in L^{n}(X, \mathcal{A}, p)$. Denote the meet $\bigwedge_{\lambda \in \Lambda} \mathcal{B}_{\lambda}$ (intersection of the null-set completions) by \mathcal{B}_{∞} . Then $\mathbb{E}[f|\mathcal{B}_{\lambda}]$ converges to $\mathbb{E}[f|\mathcal{B}_{\infty}]$ in L^{n} .

The proof is analogous to the one of the previous corollary.

We can now combine these theorems with the results at the end of Section 4.3, to obtain the following *martingale convergence theorems* in mean.

Corollary 6.12 (Martingale convergence theorem in mean). Let (X, \mathcal{A}, p) be essentially standard Borel, let $(\mathcal{B}_{\lambda})_{\lambda \in \Lambda}$ be an increasing filtration, and let $(f_{\lambda})_{\lambda \in \Lambda}$ be a martingale where either

- the f_{λ} admit a uniform L^n bound for $n \geq 2$; or
- there exists $f \in L^1(X, \mathcal{A}, p)$ such that for eventually all λ , $f_{\lambda} = \mathbb{E}[f|\mathcal{B}_{\lambda}]$.

Denote the join $\bigvee_{\lambda \in \Lambda} \mathcal{B}_{\lambda}$ by \mathcal{B}_{∞} . Then there exists $f_{\infty} \in L^{n}(X, \mathcal{B}_{\infty}, p)$ such that

- $f_{\lambda} = \mathbb{E}[f_{\infty}|\mathcal{B}_{\lambda}]$ for all λ ;
- f_{λ} converges to f_{∞} in L^n .

Proof. Let us first consider the first case. The increasing filtration induces a cofiltered diagram $\mathcal{B} : \Lambda \to \text{EKrn.}$ By applying the dagger functor and then the L^2 functor we obtain a filtered diagram D:

$$\Lambda \xrightarrow{\mathcal{B}} \operatorname{EKrn} \xrightarrow{\dagger} \operatorname{EKrn}^{\operatorname{op}} \xrightarrow{L^2} \operatorname{Hilb}_{<1}$$

This means that $D(\lambda) = L^2(X, \mathcal{B}_{\lambda}, p)$ and $D(\lambda < \mu) = (\pi_{\lambda,\mu}^{\dagger})^*$, where $\pi_{\lambda,\mu} : (X, \mathcal{B}_{\mu}, p) \to (X, \mathcal{B}_{\lambda}, p)$ is the identity on sets. By Corollary 6.10 the limit of D is given by $L^2(X, \mathcal{B}_{\infty}, p)$ with projection maps $(\pi_{\lambda}^{\dagger})^*$ for every $\lambda \in \Lambda$.

Now for every $\lambda \in \Lambda$ and for $n \geq 2$, $||f_{\lambda}||_{L^2} \leq ||f_{\lambda}||_{L^n}$, and so the uniform L^n bound (that we have by hypothesis) also gives us a uniform L^2 bound. In other words, we can define a real number $r := \sup_{\lambda} \mathbb{E}[f_{\lambda}^2]$. For $\lambda \in \Lambda$, the assignment $r \mapsto f_{\lambda}$ defines a map $\varphi_{\lambda} : \mathbb{R} \to L^2(X, \mathcal{B}_{\lambda}, p)$ in Hilb \leq_1 . Because $(f_{\lambda})_{\lambda \in \Lambda}$ is a martingale we have that \mathbb{R} together with the maps $(\varphi_{\lambda})_{\lambda \in \Lambda}$ form a cone over the diagram L^2 . Therefore there exists a unique map $\varphi_{\infty} : \mathbb{R}$ in Hilb \leq_1 such that $(\pi_{\lambda}^{\dagger})^* \varphi_{\infty} = \varphi_{\lambda}$ for every $\lambda \in \Lambda$. If we define $f_{\infty} := \varphi_{\infty}(r)$, then

$$\mathbb{E}[f_{\infty} \mid \mathcal{B}_{\lambda}] = (\pi_{\lambda}^{\dagger})^* \varphi_{\infty}(r) = \varphi_{\lambda}(r) = f_{\lambda}.$$

By Corollary 6.10, it now follows that $f_{\lambda} \to f_{\infty}$ in L^n .

The second case follows from Corollary 4.30, from the tower property of conditional expectation (taking $f_{\infty} = \mathbb{E}[f \mid \mathcal{B}_{\infty}]$), and again by Corollary 6.10.

Corollary 6.13 (Backward martingale convergence theorem in mean). Let (X, \mathcal{A}, p) be essentially standard Borel, let $(\mathcal{B}_{\lambda})_{\lambda \in \Lambda}$ be a decreasing filtration, and let $(f_{\lambda})_{\lambda \in \Lambda}$ be a martingale with $f_{\lambda} \in$ $L^n(X, \mathcal{B}_{\lambda}, p)$ for $n \geq 1$. Denote the meet $\bigwedge_{\lambda \in \Lambda} \mathcal{B}_{\lambda}$ (intersection of the null-set completions) by \mathcal{B}_{∞} . Then there exists $f_{\infty} \in L^n(X, \mathcal{B}_{\infty}, p)$ such that

- $f_{\infty} = \mathbb{E}[f_{\lambda}|\mathcal{B}_{\infty}]$ for all λ ;
- f_{λ} converges to f_{∞} in L^n .

The proof is analogous to the one of the second case of Corollary 6.12.

This version of the backward martingale convergence theorem, for decreasing filtrations indexed by arbitrary nets, seems to be new.

7 Vector-valued martingales

Our abstract formalism allows to talk about martingales in a very general way, including vectorvalues ones, and gives us a convergence result for all such classes of martingales. Here is the definition in general.

Definition 7.1. Let $F : \text{EKrn}^{\text{op}} \to \text{Ban be any functor.}$

- An *F*-valued random variable on (X, \mathcal{A}, p) is an element of the Banach space $F(X, \mathcal{A}, p)$. (Equivalently, a linear map $\mathbb{R} \to F(X, \mathcal{A}, p)$).
- An *F*-valued martingale on (X, \mathcal{A}, p) consists of an increasing filtration $(\mathcal{B}_{\lambda})_{\lambda \in \Lambda}$ on (X, \mathcal{A}, p) , together with a family of random variables $(f_{\lambda})_{\lambda \in \Lambda}$ such that for all $\lambda \leq \mu$, $f_{\lambda} = F(\pi_{\mu,\lambda}^+)(f_{\mu})$. (Equivalently, with a cone in Ban on the $F(X, \mathcal{B}_{\lambda}, p)$ with tip \mathbb{R} .)
- An F-valued backward martingale is defined analogously, but with a decreasing filtration.

Beside the ordinary RV functors of Proposition 3.6, we will also give the example of vectorvalued random variables in the Bochner (strong) sense. One could generalize this even further, replacing the category Ban with any other category, but we will not do that here.

Thanks to how we have set up the formalism, whenever the functor F is enriched we automatically have a version of the Levy theorems:

Theorem 7.2. Let $F : \text{EKrn}^{\text{op}} \to \text{Ban}$ be an enriched functor, let $(\mathcal{B}_{\lambda})_{\lambda \in \Lambda}$ be an increasing filtration on a standard Borel space (X, \mathcal{A}, p) , and let $f \in F(X, \mathcal{A}, p)$. We have that

(i) The F-images of the retracts

$$(X, \mathcal{A}, p) \xleftarrow{\pi_{\lambda}}{} (X, \mathcal{B}_{\lambda}, p)$$

 $are\ retracts$

$$F(X, \mathcal{A}, p) \xrightarrow{F(\pi_{\lambda})} F(X, \mathcal{B}_{\lambda}, p)$$

in Ban, where $F(\pi_{\lambda})$ is a projection and $F(\pi^+\lambda)$ an inclusion, and the order of split idempotents is preserved;

- (ii) If we denote by f_{λ} the element $F(\pi_{\lambda}^+)(f)$, we have that $(f_{\lambda})_{\lambda \in \Lambda}$ is an *F*-valued martingale;
- (iii) The F-image of the supremum

$$(X, \mathcal{A}, p) \xrightarrow[\pi_{\infty}]{\pi_{\infty}^+} (X, \mathcal{B}_{\infty}, p)$$

is the supremum in the split idempotent order on $F(X, \mathcal{A}, p)$ in Ban;

(iv) If we denote by f_{∞} the element $F(\pi_{\infty}^+)(f)$, we have that f_{λ} tends to f_{∞} topologically (in the norm of $F(X, \mathcal{A}, p)$).

The analogous statement is true for decreasing filtrations and backward martingales.

Proof. We will prove the statement for increasing filtrations and forward martingales, the backward case is analogous.

(i) By (contravariant) functoriality,

$$F(\pi_{\lambda}^{+}) \circ F(\pi_{\lambda}) = F(\pi_{\lambda} \circ \pi_{\lambda}^{+}) = F(\operatorname{id}_{(X,\mathcal{B}_{\lambda},p)}) = \operatorname{id}_{F(X,\mathcal{B}_{\lambda},p)},$$

so that we have a retract. To see that the order is preserved, denote by e_{λ} the idempotent induced by \mathcal{B}_{λ} . If $e_{\lambda} \leq e_{\mu}$, i.e. $e_{\lambda} \circ e_{\mu} = e_{\mu} \circ e_{\lambda} = e_{\lambda}$, again by functoriality we have that

$$F(e_{\lambda}) \circ F(e_{\mu}) = F(e_{\mu} \circ e_{\lambda}) = F(e_{\lambda}),$$

and analogously $F(e_{\mu}) \circ F(e_{\lambda}) = F(e_{\lambda})$, so that $F(e_{\lambda}) \leq F(e_{\mu})$.

(ii) Once again by functoriality, for every $\lambda \leq \mu$,

$$F(\pi_{\mu,\lambda}^{+})(f_{\mu}) = F(\pi_{\mu,\lambda}^{+}) \circ F(\pi_{\mu}^{+})(f) = F(\pi_{\mu}^{+} \circ \pi_{\mu,\lambda}^{+}) = F(\pi_{\lambda}^{+})(f) = f_{\lambda}.$$

- (iii) Denote by e_{∞} the idempotent induced by \mathcal{B}_{∞} . By Theorem 6.9, $e_{\lambda} \to e_{\infty}$ topologically in $\mathrm{EKrn}((X, \mathcal{A}, p), (X, \mathcal{A}, p))$. Since the functor F is enriched, then $F(e_{\lambda}) \to F(e_{\infty})$ topologically in $\mathrm{Ban}(F(X, \mathcal{A}, p), F(X, \mathcal{A}, p))$. Now by Proposition 5.11 and Proposition 6.2 it follows that $F(e_{\infty})$ is a supremum of the $F(e_{\lambda})$.
- (iv) From the point above, we have that $f_{\lambda} = F(e_{\lambda})(f) \to F(e_{\infty})(f) = f_{\infty}$ topologically. \Box Let's now apply this to vector-valued martingales.

7.1 Bochner integrals and conditional expectations

In what follows we will consider random variables with values in a Banach space V. The case where V is separable is particularly simple, but we will not assume it here. (Note that a separable Banach space is in particular a complete separable metric space, so that with its Borel sigma-algebra it is a standard Borel space.)

Where possible, we will follow the terminology and conventions of [Coh13, Appendix E].

Definition 7.3. Let (X, \mathcal{A}) be a measurable space, and let V be a Banach space.

- A V-valued random variable is an A-measurable function $f: X \to V$, where V is taken with the Borel sigma-algebra.
- A V-valued RV f on X is called **strongly measurable** if it is measurable and its image $f(X) \subseteq V$ is separable, and almost surely strongly measurable if it is almost surely equal to a strongly measurable RV.
- A V-valued RV f on X is called simple if it is in the form

$$f(x) = \sum_{i=1}^{n} 1_{A_i}(x) \cdot v_i$$

where $v_i \in V$ and $A_i \in \mathcal{A}$ for all i = 1, ..., n.

Every simple RV is strongly measurable. Also, if V is separable, measurability and strong measurability coincide. Just as ordinarily measurable RVs, strongly measurable RVs are closed under linear combinations [Coh13, Proposition E.3] and under pointwise limits [Coh13, Proposition E.1].

Definition 7.4. In analogy with the real case, we call an a.s. strongly measurable V-valued random variable f **Bochner-integrable** if and only if

$$\int_X \|f(x)\|_V \, p(dx) < \infty,$$

and **Bochner-***n***-***integrable*, for $n \ge 1$, if and only if

$$\int_X (\|f(x)\|_V)^n \, p(dx) \ < \ \infty.$$

We denote the set of a.s. equivalence classes of (a.s. strongly measurable) Bochner-*n*-integrable random variables on (X, \mathcal{A}, p) by $L^n(X, \mathcal{A}, p; V)$. These spaces are known in the literature under the name of *Bochner spaces*. They are Banach spaces with norm

$$\|f\|_{L^{n}(X,\mathcal{A},p;V)} = \left(\int_{X} (\|f(x)\|_{V})^{n} p(dx)\right)^{1/n},$$

analogously to usual L^n spaces. Similarly, we denote by $L^{\infty}(X, \mathcal{A}, p; V)$ the set of equivalence classes of almost surely bounded V-valued random variables, with the essential supremum norm. Just as in the real-valued case, we have that for $m \geq n$, $L^m(X, \mathcal{A}, p; V) \subseteq L^n(X, \mathcal{A}, p; V)$. It can be shown that if a RV f is Bochner-integrable, then there exists a sequence (f_n) of simple V-valued RVs

$$f_n(x) \coloneqq \sum_{i=1}^{\kappa_n} 1_{A_{n,i}}(x) \cdot v_{n,i}$$

tending pointwise to f and such that for all x and n, $||f_n(x)||_V \leq ||f(x)||_V$ [Coh13, Proposition E.2]. In that case, the **Bochner integral** of f is defined by

$$\int_X f(x) \, p(dx) \ \coloneqq \ \lim_{n \to \infty} \int_X f_n(x) \, p(dx) \ \coloneqq \ \lim_{n \to \infty} \sum_{i=1}^{k_n} p(A_{n,i}) \cdot v_{n,i},$$

is finite, and is independent of the choice of (f_n) [Coh13, p. 399]. One can show that Bochner integration is linear in f, and a version of the dominated convergence theorem holds.

We now want to construct functors $L^n(-; V) : \operatorname{GKrn}^{\operatorname{op}} \to \operatorname{Ban}$ analogous to the L^n functors defined in Section 3.2:

- On objects, we map a probability space (X, \mathcal{A}, p) to the Banach space $L^n(X, \mathcal{A}, p; V)$;
- On morphisms, given a measure-preserving kernel $k : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$, we get a bounded linear map $k^* : L^n(Y, \mathcal{B}, q; V) \to L^n(X, \mathcal{A}, p; V)$ acting on random variables $g \in L^n(Y, \mathcal{B}, q; V)$ by

$$k^*g(x) \coloneqq \int_Y g(y) \, k(dy|x). \tag{7.1}$$

Here is the precise statement, the analogous of Proposition 3.6.

Proposition 7.5. Let $k : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$ be a measure-preserving kernel, and let $g \in L^n(Y, \mathcal{B}, q; V)$, for n finite or infinite. Then the assignment

$$x \longmapsto k^* g(x) \coloneqq \int_Y g(y) \, k(dy|x)$$

is a well-defined element of $L^n(X, \mathcal{A}, p; V)$.

Moreover, the assignment $k^* : L^n(Y, \mathcal{B}, q; V) \to L^n(X, \mathcal{A}, p; V)$ given by $g \mapsto k^*g$ is linear and 1-Lipschitz, so that L^n is a functor GKrn^{op} \to Ban.

Lemma 7.6. Let $h: (X, \mathcal{A}) \to \mathbb{R}$ be a measurable function, and let $v \in V$. Then the function

$$(X, \mathcal{A}) \xrightarrow{h \cdot v} V$$
$$x \longmapsto h(x) \cdot v$$

is strongly measurable.

Proof of Lemma 7.6. Let (h_n) be a sequence of simple (real-valued) functions

$$h_n(x) \coloneqq \sum_{i=1}^{k_n} \lambda_{n,i} \, \mathbf{1}_{A_{n,i}}(x)$$

tending pointwise to h. Since the scalar multiplication

$$\begin{aligned} \mathbb{R} \times V & \longrightarrow V \\ (\lambda, v) & \longmapsto \lambda \cdot v \end{aligned}$$

is continuous, we have that the functions

$$x \longmapsto h_n(x) \cdot v = \sum_{i=1}^{k_n} \lambda_{n,i} \, \mathbf{1}_{A_{n,i}}(x) \cdot v$$

tend pointwise to $h \cdot v$. Since strongly measurable functions are closed under linear combinations and pointwise limits, we only have to show that for each n and i, the function $x \mapsto 1_{A_{n,i}}(x) \cdot v$ is strongly measurable. But this is the case: the function is measurable (since the set $A_{n,i}$ is), and it assumes at most only two values (0 and v).

Proof of Proposition 7.5 for finite n. Since k is measure-preserving, by approximation via simple (real-valued) functions,

$$\int_{X} \int_{Y} (\|g(y)\|_{V})^{n} k(dy|x) p(dx) = \int_{Y} (\|g(y)\|_{V})^{n} q(dy) < \infty,$$
(7.2)

and so the integrals

$$\int_{Y} (\|g(y)\|_V)^n \, k(dy|x)$$

are finite for p-almost all $x \in X$. Even if n > 1, again by Jensen's inequality we have that

$$\left(\int_{Y} \|g(y)\|_{V} \, k(dy|x)\right)^{n} \leq \int_{Y} (\|g(y)\|_{V})^{n} \, k(dy|x) < \infty.$$

Therefore the Bochner integral

$$x \longmapsto \int_{Y} g(y) \, k(dy|x), \tag{7.3}$$

is defined for *p*-almost all $x \in X$, and the assignment specifies a unique function $X \to V$ up to a null set. To see that this function is a.s. strongly measurable, let $(g_n)_{n=1}^{\infty}$ be a sequence of simple *V*-valued RVs

$$g_n(y) \coloneqq \sum_{i=1}^{k_n} \mathbb{1}_{B_{n,i}}(y) \cdot v_{n,i}$$

tending pointwise to g and such that $||g_n(y)||_V \leq ||g(y)||_V$. (The sequence exists since g is Bochnerintegrable.) Hence for almost all x,

$$\int_{Y} g(y) \, k(dy|x) \; = \; \lim_{n \to \infty} \int_{Y} g_n(y) \, k(dy|x) \; = \; \lim_{n \to \infty} \sum_{i=1}^{k_n} k(B_{n,i}|x) \cdot v_{n,i}$$

and so by Lemma 7.6 the assignment (7.3) is a pointwise limit of strongly measurable functions. Therefore the assignment (7.3) gives a well-defined element of $L^n(X, \mathcal{A}, p; V)$. We denote this element by k^*g . In other words, given $k : (X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$, we get a well-defined function $k^* : L^n(Y, \mathcal{B}, q; V) \to L^n(X, \mathcal{A}, p; V)$ given by $g \mapsto k^*g$. This function is linear by linearity of integration. To see that it is 1-Lipschitz, once again by Jensen,

$$(\|k^*g\|_{L^n(X,\mathcal{A},p;V)})^n = \int_X \left(\left\| \int_Y g(y) \, k(dy|x) \right\|_V \right)^n p(dx)$$

$$\leq \int_X \left(\int_Y \|g(y)\|_V \, k(dy|x) \right)^n p(dx)$$

$$\leq \int_X \int_Y (\|g(y)\|_V)^n \, k(dy|x) \, p(dx)$$

$$= \int_Y (\|g(y)\|_V)^n \, q(dy)$$

$$= (\|g\|_{L^n(Y,\mathcal{B},q;V)})^n.$$

Proof of Proposition 7.5 for infinite n. Let $g \in L^{\infty}(Y, \mathcal{B}, q; V)$. By definition there exists $B \in \mathcal{B}$ of q-measure one such that g restricted to B is bounded. Let K be an upper bound for the norm $\|g\|_B\|_V$. Since k is measure-preserving,

$$\int_X \int_Y \|g(y)\|_V \, k(dy|x) \, p(dx) \ = \ \int_Y \|g(y)\|_V \, q(dy) \ = \ \int_B \|g(y)\|_V \, q(dy) \ \le \ K.$$

Therefore the integrals

$$\int_Y \|g(y)\|_V \, k(dy|x)$$

are finite for *p*-almost all $x \in X$, and so the Bochner integral

$$x\longmapsto \int_Y g(y)\,k(dy|x),$$

is defined for p-almost all $x \in X$. Denote this value by $k^*g(x)$. As in the case of finite n, this is strongly measurable in X.

Now by Lemma 3.7, there exists $A \in \mathcal{A}$ of *p*-measure one such that for all $x \in A$, k(B|x) = 1. Therefore, for all $x \in A$,

$$\|k^*g(x)\|_V = \left\|\int_Y g(y) \, k(dy|x)\right\|_V \le \int_Y \|g(y)\|_V \, k(dy|x) = \int_B \|g(y)\|_V \, k(dy|x) \le K.$$

In other words, the restriction of k^*g to A is also bounded by K. So we get a well-defined function $k^* : L^{\infty}(Y, \mathcal{B}, q; V) \to L^{\infty}(X, \mathcal{A}, p; V)$. Moreover, since every essential upper bound for $||g||_V$ is an essential upper bound for $||k^*g||_V$, we have that

$$\|k^*g\|_{L^{\infty}} = \operatorname{ess\,sup} \|k^*g\|_V \le \operatorname{ess\,sup} \|g\|_V = \|g\|_{L^{\infty}}.$$

Just as in the real-valued case, the inclusions $L^n(X, \mathcal{A}, p; V) \subseteq L^m(X, \mathcal{A}, p; V)$ for $m \leq n$ form natural transformations.

As an example of the functorial action of $L^n(-; V)$, we can look at *V*-valued conditional expectations. Indeed, let (X, \mathcal{A}, p) be a standard Borel space, and let $\mathcal{B} \subseteq \mathcal{A}$ be a sub-sigma-algebra. The disintegration kernel $\pi^+ : (X, \mathcal{B}, p) \to (X, \mathcal{A}, p)$ given by the conditional expectations

$$\pi^+(A|x) = \mathbb{P}[A|\mathcal{B}](x)$$

induces a linear, 1-Lipschitz map $(\pi^+)^* : L^n(X, \mathcal{A}, p; V) \to L^n(X, \mathcal{B}, p; V)$ which can be considered a generalization of conditional expectation to vector-valued random variables. Because of that, let's use the same notation as for real-valued random-variables: we write

$$L^{n}(X, \mathcal{A}, p; V) \xrightarrow{(\pi^{+})^{*}} L^{n}(X, \mathcal{B}, p; V)$$
$$f \longmapsto \mathbb{E}[f|\mathcal{B}].$$

Note that, just as in the real-valued case:

- $\mathbb{E}[f|\mathcal{B}]$ is (strongly) \mathcal{B} -measurable;
- For every $B \in \mathcal{B}$ we have that

$$\int_{B} \mathbb{E}[f|\mathcal{B}](x) \, p(dx) = \int_{B} f(x) \, p(dx).$$

and $\mathbb{E}[f|\mathcal{B}]$ is determined almost surely uniquely by these properties.

Also, in this notation, a V-valued martingale looks just like a real-valued one: a V-valued martingale on (X, \mathcal{A}, p) with filtration (B_{λ}) is a family of V-valued, strongly \mathcal{B}_{λ} -measurable RVs (f_{λ}) such that for all $\lambda \leq \mu$, $f_{\lambda} = \mathbb{E}[f_{\mu}|\mathcal{B}_{\lambda}]$.

7.2 Topological enrichment and convergence

Let's now turn to topological enrichment and convergence.

Proposition 7.7. For every Banach space V and for each n, the functor $L^n(-; V)$: GKrn^{op} \rightarrow Ban is topologically enriched.

Proof. Let $(k_{\lambda})_{\lambda \in \Lambda}$ be a net of measure-preserving kernels $(X, \mathcal{A}, p) \to (Y, \mathcal{B}, q)$ tending to k. That is, for every $B \in \mathcal{B}$,

$$\int_X |k(B|x) - k_\lambda(B|x)| \, p(dx) \to 0.$$

We have to show that the induced maps $k_{\lambda}^* : L^n(Y, \mathcal{B}, q; V) \to L^n(X, \mathcal{A}, p; V)$ tend to k^* , meaning that for all $g \in L^n(Y, \mathcal{B}, p; V)$,

$$||k^*g - k^*_{\lambda}g||_{L^n(X,\mathcal{A},q;V)} \to 0,$$

i.e.

$$\int_X \|k^*g(x) - k^*_\lambda g(x)\|_V \, p(dx) \to 0.$$

By Lemma 5.4 it suffices to test this convergence on a dense subset of $L^n(Y, \mathcal{B}, q; V)$, and we take the set simple V-valued RVs, i.e. those in the following form,

$$g(y) = \sum_{i=1}^{k} \mathbb{1}_{B_i}(x) \cdot v_i$$

for $v_i \in V$ and $B_i \in \mathcal{B}$. Now

$$k_{\lambda}^*g = \int_Y g(y) \, k(dy|x) = \sum_{i=1}^k k_{\lambda}(B_i|x) \cdot v_i,$$

and the same is true for k, so that

$$\begin{split} \int_{X} \|k^{*}g(x) - k_{\lambda}^{*}g(x)\|_{V} p(dx) &= \int_{X} \left\| \sum_{i=1}^{k} \left(k(B_{i}|x) - k_{\lambda}(B_{i}|x) \right) \cdot v_{i} \right\|_{V} p(dx) \\ &\leq \int_{X} \left| \sum_{i=1}^{k} k(B_{i}|x) - k_{\lambda}(B_{i}|x) \right| \|v_{i}\|_{V} p(dx) \\ &\leq \sum_{i=1}^{k} \left(\int_{X} |k(B_{i}|x) - k_{\lambda}(B_{i}|x)| p(dx) \right) \|v_{i}\|_{V}, \end{split}$$

but the term in the brackets tends to zero by assumption. Hence $k_{\lambda}^* \to k^* g$.

Because of this, Theorem 7.2 can be applied, and it gives us the following result.

Corollary 7.8. Let (X, \mathcal{A}, p) be a standard Borel probability space, and let V be a Banach space.

(i) Let $(\mathcal{B}_{\lambda})_{\lambda \in \Lambda}$ be an increasing filtration and let by \mathcal{B}_{∞} its supremum (join of sigma-algebras). Given $f \in L^{n}(X, \mathcal{A}, p, V)$, denote by f_{λ} and f_{∞} the (V-valued) conditional expectations $\mathbb{E}[f|\mathcal{B}_{\lambda}]$ and $\mathbb{E}[f|\mathcal{B}_{\infty}]$. Then (f_{λ}) is a V-valued martingale, and f_{λ} tends to f_{∞} in mean, meaning that

$$\int_X (\|f(x) - f_\lambda(x)\|_V)^n \, p(dx) \to 0.$$

(ii) Let $(\mathcal{B}_{\lambda})_{\lambda \in \Lambda}$ be a decreasing filtration and let by \mathcal{B}_{∞} its infimum (intersection of the nullset-completions). Given $f \in L^{n}(X, \mathcal{A}, p, V)$, denote by f_{λ} and f_{∞} the (V-valued) conditional expectations $\mathbb{E}[f|\mathcal{B}_{\lambda}]$ and $\mathbb{E}[f|\mathcal{B}_{\infty}]$. Then (f_{λ}) is a V-valued backward martingale, and f_{λ} tends to f_{∞} in mean, as above.

A Topologically enriched categories

The category Top is not cartesian closed. It is however still *monoidal* closed, where as tensor product we take the topology of *separate* continuity, instead of joint continuity.

A.1 The tensor product

Definition A.1. Let X and Y be topological spaces. We call a subset subset $U \subseteq X \times Y$ separately open if and only if

• For each $x \in X$, the set $U_x \subseteq Y$ given by

$$U_x \coloneqq \{y \in Y : (x, y) \in U\}$$

is an open subset of Y;

• For each $y \in Y$, the set $U_y \subseteq X$ given by

$$U_y \coloneqq \{x \in X : (x, y) \in U\}$$

is an open subset of X.

Proposition A.2. Let X and Y be topological spaces. The separately open subsets of $X \times Y$ form a topology.

Proof. First of all, the empty set and the set $X \times Y$, seen as subsets of $X \times Y$, are separately open. Let now U and V be separately open. For each $x \in X$ have that

$$\begin{split} (U \cap V)_x &= \{ y \in Y : (x, y) \in U \cap V \} \\ &= \{ y \in Y : (x, y) \in U, (x, y) \in V \} \\ &= \{ y \in Y : y \in U_x, y \in V_x \} \\ &= U_x \cap V_x, \end{split}$$

which is open. The same is true for $(U \cap V)_y$ for each $y \in Y$. Given a family $(U_i)_{i \in I}$ of separately open subsets of $X \times Y$, for $x \in X$,

$$\left(\bigcup_{i} U_{i}\right)_{x} = \left\{ y \in Y : (x, y) \in \bigcup_{i} U_{i} \right\}$$
$$= \left\{ y \in Y : (x, y) \in U_{i} \text{ for some } i \in I \right\}$$
$$= \bigcup_{i} U_{i} \{ y \in Y : (x, y) \in U_{i} \}$$
$$= \bigcup_{i} (U_{i})_{x},$$

which is open. Once again, the same is true for $(\bigcup_i U_i)_{y}$.

Definition A.3. We call the topology on $X \times Y$ given by the separately open subsets the **topology** of separate continuity or the tensor product topology. We denote the resulting topological space by $X \otimes Y$.

Note that this makes Top a monoidal category, with the one-point space as monoidal unit (so it's semicartesian monoidal).

Proposition A.4. Let X, Y and Z be topological spaces, and let $f : X \times Y \to Z$ be a function. We have that f is separately continuous in X and Y if and only if it is continuous for the topology of separate continuity.

For this proof, and for later convenience, given a function $f: X \times Y \to Z$, define

- For each $x \in X$, the function $f_x : Y \to Z$ given by $f_x(y) = f(x, y)$;
- For each $y \in Y$, the function $f_y : X \to Z$ given by $f_y(x) = f(x, y)$.

Proof. The function f is continuous as a function $f : X \otimes Y \to Z$ if and only if for every open $V \subseteq Z$, the preimage

$$f^{-1}(V) = \{(x, y) \in X \times Y : f(x, y) \in V\}$$

is separately open. This means that

• For every $x \in X$, the set

$$(f^{-1}(V))_x = \{y \in Y : (x, y) \in f^{-1}(V)\} \\= \{y \in Y : f(x, y) \in V\} \\= \{y \in Y : f_x(y) \in V\} \\= f_x^{-1}(V)$$

is open. This means exactly that $f_x: Y \to Z$ is continuous;

• For every $x \in X$, the set

$$(f^{-1}(V))_y = f_y^{-1}(V)$$

is open. This means exactly that $f_y: X \to Z$ is continuous.

Now f is separately continuous if and only if for all x and y, the functions $f_x : Y \to Z$ and $f_y : X \to Z$ are continuous.

A.2 The internal hom

Definition A.5. Let X and Y be topological spaces. Denote by [X, Y], and call it internal hom, the set of continuous functions $X \to Y$, equipped with the topology of pointwise convergence, generated by the (sub-basic) sets

$$S(x,V) \coloneqq \{f: X \to Y : f(x) \in V\}$$

for each $x \in X$ and each open $V \subseteq Y$.

Proposition A.6. Let X and Y be topological spaces. Let $(f_{\alpha})_{\alpha \in \Lambda}$ be a net (or a sequence) in [X, Y]. We have that $f_{\alpha} \to f$ if and only if for every $x \in X$, $f_{\alpha}(x) \to f(x)$.

Proof. We have that $f_{\alpha} \to f$ if and only if every sub-basic open neighborhood S(x, V) of f eventually contains f_{α} . That is, if for each $x \in X$ and for each open $V \subseteq Y$ with $f(x) \in V$, there exists a $\beta \in \Lambda$ such that for all $\alpha \geq \beta$, $f_{\alpha}(x) \in V$. This is exactly saying that for each $x \in X$, the net $(f_{\alpha}(x))_{\alpha \in \Lambda}$ converges to f(x).

Proposition A.7. Let X, Y and Z be topological spaces. We have a natural bijection

$$\operatorname{Top}(X \otimes Y, Z) \xrightarrow{\sharp} \operatorname{Top}(X, [Y, Z])$$
$$f \longmapsto (x \stackrel{f^{\sharp}}{\longmapsto} f_x)$$

induced by the corresponding one of Set.

Therefore Top with this tensor product and internal hom is closed monoidal.

Proof. Note that the hom-tensor adjunction mapping $f \mapsto f^{\sharp}$ for Set is a natural bijection. Therefore it suffices to prove that $f: X \times Y \to Z$ is separately continuous if and only if

• for every $x \in X$, $f^{\sharp}(x) = f_x$ is a continuous function $Y \to Z$;

• the map $f^{\sharp}: X \to [Y, Z]$ is continuous.

Now for each $x \in X$, the function f_x is continuous if and only if for each open $V \subseteq Z$, the set $f_x^{-1}(V) = (f^{-1}(V))_x$ is open. Also, $f^{\sharp} : X \to [Y, Z]$ is continuous if and only if for each $y \in Y$ and each open $V \subseteq Z$, the set

$$(f^{\sharp})^{-1}(S(y,V)) = \{x \in X : f_x \in S(y,V)\} \\ = \{x \in X : f_x(y) \in V\} \\ = \{x \in X : f(x,y) \in V\} \\ = (f^{-1}(V))_y$$

is open. The two conditions together are therefore equivalent to f being separately continuous. \Box

A.3 Categories enriched in Top

Consider Top with the closed monoidal structure of the previous section. Instantiating the usual definition of enriched category, we get:

Definition A.8. A category enriched in Top, or topologically enriched category, amounts to

- An ordinary category C;
- On each hom-set C(X,Y), a topology, in such a way that each composition assignment

$$\begin{array}{ccc} \mathcal{C}(X,Y) \times \mathcal{C}(Y,Z) & \longrightarrow & \mathcal{C}(X,Z) \\ & & (f,g) & \longmapsto & g \circ f \end{array}$$

is separately continuous.

The second condition is equivalent to the following two conditions:

- Given a net (or sequence) $(f_{\alpha} : X \to Y)_{\alpha \in \Lambda}$ tending to $f : X \to Y$, and a morphism $g: Y \to Z$, we have that $g \circ f_{\alpha} \to g \circ f$, i.e. postcomposition is a continuous operation;
- Given a morphism $f: X \to Y$ and a net (or sequence) $(g_{\alpha}: Y \to Z)_{\alpha \in \Lambda}$ tending to $g: Y \to Z$, we have that $g_{\alpha} \circ f \to g \circ f$, i.e. precomposition is a continuous operation.

References

- [AR18] Takanori Adachi and Yoshihiro Ryu. A category of probability spaces, 2018. arXiv:1611.03630. ↑2, 3, 17, 23
- [BD86] Francis Borceux and Dominique Dejean. Cauchy completion in category theory. Cahiers de Topologie et Géométrie Différentielle Catégoriques, 27(2):133–146, 1986. ↑9
- [CJ19] Kenta Cho and Bart Jacobs. Disintegration and Bayesian inversion via string diagrams. Math. Structures Comput. Sci., 29:938–971, 2019. arXiv:1709.00322. ↑2
- [Coh13] Donald L. Cohn. Measure theory. Birkhäuser, 2013. ↑54, 55
- [DDGS18] Fredrik Dahlqvist, Vincent Danos, Ilias Garnier, and Alexandra Silva. Borel kernels and their approximation, categorically. In MFPS 34: Proceedings of the Thirty-Fourth Conference on the Mathematical Foundations of Programming Semantics, volume 341, pages 91–119, 2018. ↑2, 3, 16, 17, 23
- [DMH24] Matthew Di Meglio and Chris Heunen. Dagger categories and the complex numbers: Axioms for the category of finite-dimensional Hilbert spaces and linear contractions, 2024. arXiv:2401.06584. ↑5, 47, 48
- [EP23] Noé Ensarguet and Paolo Perrone. Categorical probability spaces, ergodic decompositions, and transitions to equilibrium, 2023. arXiv:2310.04267. ↑2, 16, 17, 22, 24, 26, 28
- [FGHL⁺23] Tobias Fritz, Tomáš Gonda, Nicholas Gauguin Houghton-Larsen, Antonio Lorenzin, Paolo Perrone, and Dario Stein. Dilations and information flow axioms in categorical probability. *Mathematical Structures in Computer Science*, 33(10), 2023. arXiv:2211.02507. ↑27
- [FGL⁺] Tobias Fritz, Tomáš Gonda, Antonio Lorenzin, Paolo Perrone, and Dario Stein. Absolute continuity, supports and idempotent splitting in categorical probability. arXiv:2308.00651. ↑2
- [FGP21] Tobias Fritz, Tomáš Gonda, and Paolo Perrone. de Finetti's theorem in categorical probability. J. Stoch. Anal., 2(4), 2021. arXiv:2105.02639. ↑2
- [FR20] Tobias Fritz and Eigil Fjeldgren Rischel. Infinite products and zero-one laws in categorical probability. Compositionality, 2:3, 2020. arXiv:1912.02769. ↑2, 5
- [Fri20] Tobias Fritz. A synthetic approach to Markov kernels, conditional independence and theorems on sufficient statistics. Adv. Math., 370:107239, 2020. arXiv:1908.07021. ↑2, 16, 24, 27
- [Gir82] Michèle Giry. A categorical approach to probability theory. In Categorical aspects of topology and analysis, volume 915 of Lecture Notes in Mathematics. Springer, 1982. ↑2
- [Isb64] J. R. Isbell. Uniform spaces. American Mathematical Society, 1964. ↑4

- [Kar18] Martti Karvonen. The Way of the Dagger. PhD thesis, University of Edinburgh, 2018. $\uparrow 2$
- [Koz16] Dexter Kozen. Kolmogorov extension, martingale convergence, and compositionality of processes. In 2016 31st Annual ACM/IEEE Symposium on Logic in Computer Science (LICS), page 692–699. IEEE, 2016. ↑2, 5
- [MP22] Sean Moss and Paolo Perrone. Probability monads with submonads of deterministic states. In Proceedings of the 37th Annual ACM/IEEE Symposium on Logic in Computer Science, pages 1–13, 2022. ↑2
- [MP23] Sean Moss and Paolo Perrone. A category-theoretic proof of the ergodic decomposition theorem. Ergodic Theory Dynam. Systems, pages 1–27, 2023. arXiv:2207.07353. ↑2
- [Per24] Paolo Perrone. Markov Categories and Entropy. IEEE Transactions of Information Theory, 70(3), 2024. arXiv:2212.11719. ↑4
- [Rie16] Emily Riehl. Category theory in context. Mineola, NY: Dover Publications, 2016. https://math.jhu.edu/~eriehl/context.pdf. ↑15
- [VB22] Ruben Van Belle. A categorical proof of the Carathéodory extension theorem, 2022. arXiv:2210.01720. ↑2
- [VB23] Ruben Van Belle. A categorical treatment of the Radon-Nikodym theorem and martingales, 2023. arXiv:2305.03421. ↑2, 3, 4, 17, 23, 38, 45