_

_

Algebras of Interaction and Cooperation

Ulrich Faigle*

April 25, 2024

Abstract: Systems of cooperation and interaction are usually studied in the context of real or complex vector spaces. Additional insight, however, is gained when such systems are represented in vector spaces with multiplicative structures, *i.e.*, in algebras. Algebras, on the other hand, are conveniently viewed as polynomial algebras. In particular, basic interpretations of natural numbers yield natural polynomial algebras and offer a new unifying view on cooperation and interaction. For example, the concept of *Galois transforms* and *zero-dividends* of cooperative games is introduced as a nonlinear analogue of the classical Harsanyi dividends. Moreover, the polynomial model unifies various versions of Fourier transforms. Tensor products of polynomial spaces establish a unifying model with quantum theory and allow to study classical cooperative games as interaction activities in a quantum-theoretic context.

Keywords: Activity system, cooperative game, decision system, evolution, Fourier transform, Galois transform, interaction system, measurement, quantum game.

1 Introduction

What is a *game*? While "playing" is often understood as an activity involving one or more human players, we take here the general game-theoretic approach of [9] and think of an underlying *system* that is characterized by the *states* it can assume. The states typically depend on certain *actions* that *players* may or may not take in order to achieve other states. We do not necessarily require players to be human beings with human interests and feelings *etc.* and thus might refer to the players

^{*}Mathematisches Institut, Universität zu Köln, Weyertal 80, 50931 Köln, Germany eMail: faigle@zpr.uni-koeln.de

also simply as *agents*. The agents my act, interact or cooperate according to the rules within specified contexts.

A mathematical analysis of games, of course, requires mathematical models for the underlying systems and their states. States have to be described as mathematical objects. Similarly, observations on systems should be modeled accordingly as mathematical functions on the collection of states. The present paper concentrates on these aspects. In particular, we refer to *quantum games* if the underlying systems fit or extend standard mathematical models of physical quantum systems.

Questions about optimal strategies for the achievement of certain goals or about the existence of Nash equilibria *etc*. are disregarded. Given appropriate models, such questions lead to mathematical optimization problems that can be studied in their own right¹.

In all of mathematical application analysis, and in game theory in particular, linear models have been proven to be of utmost importance. These models could be formulated and studied as abstract structures. The key in our analysis is the representation of relevant system parameters by polynomials as this point of view ties together many otherwise seemingly different models. Our polynomials are not polynomial functions in variables x_i at the outset, but formal polynomials in indeterminates x_i . Appropriate interpretations of and substitutions into the indeterminates then reveal fundamental aspects of various game-theoretic models.

1.1 Outline of the presentation

The algebraic structures in our analysis are suggested naturally when one represents system states as polynomials rather than vectors. The idea of expressing mathematical system characteristics via polynomials has a long tradition in general algebra (see, e.g., [32]). In fact, classical algebra arose from the wish to solve polynomial equations. The study of cooperative games in terms of an associated polynomial function in n variables was initiated in [27]. Polynomial functions, however, typically imply just one particular algebraic structure: the addition and multiplication rules of scalar-valued functions.

A substantially improved modeling flexibility is gained with *formal* polynomials rather than polynomial functions. In this case, one deals with *indetermi*-

¹See, e.g., [21, 14, 9].

nates instead of *variables*. Depending on the interpretation of the indeterminates, one is then led to various natural algebraic structures.

The present approach is based on the three fundamental aspects of natural numbers: cardinalities of finite sets, binary representation of finite sets and representation of information in terms of (0,1)-sequences (Section 4). Each of these aspects implies its own multiplication rule for formal polynomials.

The polynomial model is then applied to cooperative games and activity systems. Section 6 links polynomials to linear transforms and, in particular, to Fourier transforms. Moreover, the new concept of *Galois transforms* is shown to appear naturally and yields *nonlinear* transforms of cooperative games. Interaction games are treated in Section 7 with the emphasis on the role of a *valuation* on a game-theoretic system as a particular state-dependent Heisenberg type measure on the system. A discussion with a perspective on future work concludes the presentation.

1.2 Relation to earlier work

Often, game theory is regarded as a scientific discipline in its own right². Moreover, cooperative games and strategic games with non-cooperative players are treated separately. However, as questions about the computation of strategies became more and more of interest, the many connections of game theory with other mathematical fields (*e.g.*, mathematical optimization) became prominent. Computational questions have furthermore led to the emergence of game-theoretic research in computer science³. Moreover, a seemingly new area of game theory has been initiated where the games are supposed to be played according to the physical laws of quantum mechanics⁴.

As it turns out, a comprehensive approach to mathematical game theory is possible which ties together various areas of applied mathematics and includes the different aspects above as special cases⁵. In particular, the relationship between game-theoretic cooperation and quantum mechanics has been recognized. In this sense, the present work is a continuation of the mathematical game theory research program begun in [11, 12]. As in the classical foundations of mathe-

²Also the fundamental mathematical theory [23] proceeds this way.

³cf. [24].

⁴See, *e.g.*, [28]

⁵See [9].

matical optimization⁶, our mathematical model is essentially linear. However, it is observed that quadratic (and thus geometric, but nonlinear) measurement models arise naturally from the projection of linear operators onto lower dimensional spaces (*cf.* Section 7.2).

While [13] explores some first aspects, the present analysis pursues more generally the realization that the employment of polynomial algebras instead of pure abstract vector spaces as modeling tools offers a distinctly more refined mathematical analysis and, furthermore, relates game theory to classical algebra. No previous mathematical game-theoretical model seems to have taken this route before.

2 Mathematical preliminaries

Let $\mathbb{N} = \{0, 1, 2, \dots, n, \dots\}$ denote the set of natural numbers⁷. \mathbb{R} is the field of real numbers and \mathbb{C} the field of complex numbers, *i.e.*, numbers of the form

$$z = a + ib$$
 where $a, b \in \mathbb{R}$ and $i^2 = -1$.

The complex number z = a + ib admits a representation in the form of de Moivre:

$$z = re^{it} = r(\cos t + i\sin t)$$
 with real numbers $r, t \ge 0$. (1)

The number $\overline{z}=a-\mathrm{i}b=re^{-\mathrm{i}t}$ is the *conjugate* of $z=a+\mathrm{i}b$ and has the property

$$z\overline{z} = a^2 + b^2 = r^2 = |z|^2$$
.

For arbitrary sets X and Y, $X \times Y$ is the set of all pairs (x, y) of elements $x \in X, y \in Y$. Y^X is the collection of all functions $f: X \to Y$, which may be thought of as *valuations* of the elements of X with values from Y.

The set \mathbb{C}^X of all complex valuations of X is a vector space under the usual addition and scalar multiplication of complex-valued functions. The *support* of $f \in \mathbb{C}^X$ is the set

$$\mathrm{supp}(f)=\{x\in X|f(x)\neq 0\}.$$

If $X = \{x_1, \dots, x_n\}$ is a finite set, \mathbb{C}^X may conveniently be identified with the n-dimensional coordinate space \mathbb{C}^n and, similarly, $\mathbb{C}^{X \times X}$ with the space of all complex $n \times n$ matrices.

⁶See, e.g., [21, 14].

⁷"Die ganzen Zahlen hat der liebe Gott gemacht, alles andere ist Menschenwerk." (L. Kronecker) ("The integers were created by the good Lord, everything else is the work of man.")

2.1 Polynomials

A (formal) polynomial relative to the set X is an element $f \in \mathbb{C}^X$ with finite support. In order to emphasize the polynomial aspect, we write elements $f \in \mathbb{C}^X$ as formal sums

$$f = \sum_{x \in X} f_x x$$
 with the coefficients $f_x = f(x)$.

In the case $supp(f) \subseteq \{x_0, x_1, \dots, x_n\}$, we write the polynomial f also in the form

$$f = \sum_{k=0}^{n} f_k x_k$$
 or, using superscripts $x^k \doteq x_k$, $f = \sum_{k=0}^{n} f_k x^k$.

We denote by $\mathbb{C}(X)$ the complex vector space of all polynomials in \mathbb{C}^X . The elements $x \in X$ are formally just *indeterminates* without a numerical meaning in their own right. However, they can be given particular functional meaning in applications of the polynomial model.

Remark 1 For notational convenience, we will often identify the indeterminate $x \in X$ with the polynomial $1x \in \mathbb{C}(X)$.

2.2 Polynomial functions

If $X = \{x_0(t), x_1(t), x_2(t), \ldots\}$ is a family of complex-valued functions $x_k(t)$ in the variable t, one may think of a polynomial $p \in \mathbb{C}(X)$ as a complex-valued function in the variable t,

$$p(t) = \sum_{k>0} c_k x_k(t),$$

which arises as the corresponding linear combination of the functions in X. In the special case of the functions $x_k(t) = t^k$, p(t) is a *standard polynomial function*:

$$p(t) = c_0 + c_1 x + c_2 t^2 + \ldots + c_n t^n.$$

Remark 2 The representation of polynomials as functions allows the application of the methods of differentiation and intergration in their analysis. For modeling purposes, however, it is important to retain the flexibility of formal polynomials.

Ex. 1 (Wave functions) The choice $x_k(t) = (e^{2\pi i t})^k = e^{2\pi i kt}$ exhibits the wave functions $v : \mathbb{R} \to \mathbb{C}$ of type

$$v(t) = \sum_{k>0} c_k [\cos(2\pi kt)] + i \sin(2\pi kt)]$$

as polynomial functions.

2.3 Power series and generating functions

In the case $X = \{x_0, x_1, x_2, \dots, \}$ of indeterminates that are indexed by the natural numbers, we think of $f \in \mathbb{C}^X$ as a *(formal power series)* with the notational representation

$$f = \sum_{k>0} f_k x^k.$$

Assuming $X=\{1,t,t^2,\ldots\}$ as a set of polynomial functions $x(t)=t^k$, on the other hand, the vector $f\in\mathbb{C}^X$ may define a complex function

$$f(t) = \sum_{k \ge 0} f_k t^k$$

if the sum has a well-defined region of convergence in \mathbb{C} . In this case, the function f(t) is interpreted as *generating* the numerical parametes $f_k \in \mathbb{C}$ via evaluations of derivatives:

$$f_k = \frac{f^{(k)}(0)}{k!}$$
 $(k = 0, 1, 2, ...).$

Generating functions have proven useful in probability theory or in the asymptotic analysis of combinatorial parameter sequences⁸. If X is finite, a generating function is just a polynomial function.

3 Incidence algebra

The function product of two functions $f,g:X\to\mathbb{C}$ is the function $f\cdot g$ obtained from the usual product of the components

$$(f \cdot g)(x) = f(x)g(x) \quad (x \in X)$$
 (2)

 $⁽f \cdot g)(x)$ 8 See, e.g., [1, 2, 15]

and implies the corresponding multiplication not only of polynomials but of general power series. Note that the multiplication

$$(f,g) \mapsto f \cdot g$$

is commutative and *bilinear*, *i.e.*, linear in each of the two components. All the multiplication structures we consider subsequently are derived from the function product, although not all are commutative.

Remark 3 (Hermitian products) *Geometric considerations associate with a polynomial* $f \in \mathbb{C}(X)$ *a measure of its length as its* norm, *namely the the real number*

$$||f|| = \sqrt{\sum_{x \in X} |f_x|^2} = \sqrt{\sum_{x \in X} f_x \cdot \overline{f_x}},$$

which leads to the consideration of the hermitian function product

$$(f,g) \mapsto f \cdot \overline{g},$$

which is not linear but conjugated linear in its second component. The component sum of $f \cdot \overline{q}$ is the (hermitian) inner product

$$(f|g) = \sum_{x} (f \cdot \overline{g})_x = \sum_{x} f_x \overline{g_x} \quad with \quad ||f|| = \sqrt{(f|f)}.$$
 (3)

3.1 Incidence functions

For our puroses, an *incidence function* on the set X is a function

$$m: X \times X \times X \rightarrow \{0, 1\}.$$

Given m, we define the product $p=f\bullet g$ of the polynomials $f,g\in\mathbb{C}(X)$ as the polynomial with the coefficients

$$p_x = \sum_{y,z \in X} m(x,y,z) f_y g_z. \tag{4}$$

The multiplication $(f,g) \mapsto f \bullet g$ on $\mathbb{C}(X)$ is no longer necessarily commutative, but still *bilinear*. We refer to $(\mathbb{C}(X), \bullet)$ as a *(polynomial) incidence algebra*. This notion generalizes Rota's [29] fundamental algebraic model for the combinatorial analysis of discrete functions.

Ex. 2 (Möbius transform [29]) Let (X, \leq) be a partially ordered set and consider the incidence function

$$\zeta(x, y, z) = 1 \iff x \ge y = z.$$

Define the product $h = f \bullet g$ of the polynomials $f, g \in \mathbb{C}(X)$ as the polynomial with the coefficients

$$h_x = \sum_{y \le x} f_y g_y = \sum_{y,z} \zeta(x,y,z) f_y g_z.$$

The choice of unit coefficients $g_y = 1$, yields a linear operator $f \mapsto \zeta(f)$, where

$$\zeta(f)_x = \sum_{y \le x} f_y.$$

It is not difficult to see (and well-known) that the inverse operator $\mu = \zeta^{-1}$ exists. $w = \mu(f)$ is the Möbius transform of f (relative to the given partial order on X) and has the property

$$f_x = \sum_{y \le x} w_y \quad (x \in X).$$

Remark 4 In game theory, the coefficients w_y of the Möbius transform $w = \mu(f)$ are known as the Harsanyi [17] dividends of the valuation f.

If the incidence function m is *finitary*, i.e., if the set

$$m(x) = \{(y, z) \in X \times X \mid m(x, y, z) = 1\}$$
 is finite for all $x \in X$,

the product rule of the algebra $(\mathbb{C}(X), \bullet)$, is easily extended to a product rule in the vector space $\mathbb{C}^*(X)$ of all linear functionals $L: \mathbb{C}(X) \to \mathbb{C}$. The product $F \bullet G$ is defined for $F, G \in \mathbb{C}^*(X)$ as the linear functional that acts on the basis polynomials $x \in X$ of $\mathbb{C}(X)$ as follows:

$$(F \bullet G)(x) = \sum_{(y,z) \in m(x)} F_y G_z. \tag{5}$$

Remark 5 (Umbral calculus) Let $X = \{x^0, x^1, \dots, \}$ be indexed by the natural numbers and consider the indicence function

$$m(x^k, x^i, x^j) = 1 \iff k = i + j.$$

Umbral calculus is a technique to derive combinatorial identities by moving formally from superscripts to subscripts. It can be justified via the application of linear operators and functionals on $\mathbb{C}(X)$ under the general multiplication rule

$$(f \bullet g)_k = \sum_{i \le k} \beta_{ik} f_j g_{k-i}. \tag{6}$$

with suitable connection coefficients β_{ik} . See [30] for more details and many examples.

3.2 Monoidal algebras

The polynomial algebra $(\mathbb{C}(X), \bullet)$ is *monoidal* if the multiplication turns X into a *monoid*, *i.e*, if

- 1. $1 \bullet x = x \bullet 1 = x$ holds for some neutral element $1 \in X$ and all $x \in X$;
- 2. $x \bullet (y \bullet z) = (x \bullet y) \bullet z \in X$ holds for all $x, y, z \in X$.

Conversely, if (X, \bullet) is a monoid, the multiplication extends to a multiplication of arbitrary polynomials in $\mathbb{C}(X)$ *via* the incidence function

$$m(x, y, z) = 1 \iff x = y \bullet z.$$

Ex. 3 (Lattices) The partial order $L = (X, \leq)$ is a semilattice if for all $x, y \in X$, there exists a unique largest element $x \wedge y$ such that

$$x \wedge y \leq x$$
 and $x \wedge y \leq y$.

If also a unique maximal element $x_{max} \in L$ exists, then (X, \wedge) is a monoid with neutral element x_{max} . The semilattice L is a lattice if furthermore, any $x, y \in L$ admit a unique minimal element $x \vee y$ such that

$$x \lor y \ge x$$
 and $x \lor y \ge y$,

i.e, if also the inverted partial order $L^* = (X, \ge)$ is a semilattice. If L has a minimal element x_{min} , then (X, \vee) is a monoid with neutral element x_{min} .

4 Algebra of natural numbers

We denote the usual standard sum of natural numbers $i, j \in \mathbb{N}$ by i+j and their product by $i \cdot j$ or simply ij. However, other important algebraic structures on \mathbb{N} also offer themselves, depending on the representation of natural numbers.

4.1 Binary representation and algebra

Every natural number $k < 2^n$ has a unique representation with n binary digits k_i :

$$k = \sum_{i=0}^{n-1} k_i 2^k \quad (k_i \in \{0, 1\}). \tag{7}$$

So every (0,1)-string $\alpha \in \{0,1\}^{\mathbb{N}}$ with finite support corresponds to a unique natural number

$$a = \sum_{i=0}^{\infty} \alpha_i 2^i.$$

Endowing the set $\{0,1\}$ with the binary addition rule

$$1 \oplus 1 = 0$$
,

one obtains a binary addition rule for all natural numbers:

$$\left(\sum_{i=0}^{\infty} \alpha_i 2^i\right) \oplus \left(\sum_{i=0}^{\infty} \beta_i 2^i\right) = \sum_{i=0}^{\infty} (\alpha_i \oplus \beta_i) 2^i \quad (\alpha_i, \beta_i \in \{0, 1\}), \tag{8}$$

which is commutative and associative. Moreover, 0 is the neutral element (as under the standard addition rule).

Consider now a set $X = \{x^0, x^1, \dots, x^n, \dots\}$ of indeterminates x^n with indices $n \in \mathbb{N}$. The binary addition suggests a commutative and associative multiplication on X:

$$x^i \odot x^j = x^{i \oplus j}$$

with the neutral element $\mathbf{1}=x^0$. The monoid (X,\odot) implies the polynomial algebra $(\mathbb{C}(X),\odot)$ where the (binary) product of the polynomials $f,g\in\mathbb{C}(X)$ is the polynomial $f\odot g$ with the coefficients

$$(f \odot g)_n = \sum_{i \oplus j = n} f_i g_i.$$

IMPARTIAL GAMES. We illustrate binary algebra with the example of the well-known 2-person game Nim, which involves two players and a finite set N that is partitioned into k blocks N_i with $n_i = |N_i|$ objects each. A move of a player consists in the choice of a non-empty block N_i and the subsequent removal of at least one object from N_i . The players move alternatingly. A player loses if

he cannot move on his turn. We associate with the Nim game its *characteristic* polynomial as the binary monomial

$$p(N) = x^{n_1} \odot x^{n_2} \odot \cdots \odot x^{n_k} = x^{n_1 \oplus n_2 \oplus \ldots + \oplus n_k}.$$

By the Sprague-Grundy theory of combinatorial 2-person games one then finds:

- The second player has a winning strategy if p(N) = 1 (= x^0).
- If $p(N) \neq 1$, the first player has a winning strategy.

Remark 6 An actual winning strategy for the Nim game N is easily computed: The current player moves, if possible, the game into a Nim situation N' with characteristic polynomial p(N') = 1.

Nim games are prototypical impartial 2-person games as each impartial game is strategically equivalent to a Nim game.

4.2 Set representations and Boolean algebra

Each (0,1)-string $\alpha \in \{0,1\}^{\mathbb{N}}$ with components α_i describes a unique subset A of \mathbb{N} via

$$A = \operatorname{supp}(\alpha) = \{ i \in \mathbb{N} \mid \alpha_i = 1 \}. \tag{9}$$

In fact, (9) establishes a one-to-one correspondence between $\{0,1\}^{\mathbb{N}}$ and the subsets of \mathbb{N} . Moreover, strings with finite support correspond to finite subsets and, simultaneously, to natural numbers *via* their binary representation (7).

Consider now an arbitrary n-element set $E = \{e_1, \dots, e_n\}$. Each subset $A \subseteq E$ corresponds to a unique (0, 1)-string α of length n,

$$\alpha = \alpha_1 \dots \alpha_n$$
 with $\alpha_i = 1 \longleftrightarrow e_i \in A$,

and also to a natural number

$$a = \sum_{i=1}^{n} \alpha_i 2^{i-1} < 2^n.$$

Consequently, the family \mathcal{E} of all subsets $A \subseteq E$ corresponds to the set

$$\mathbb{N}_n = \{ k \in \mathbb{N} \mid k < 2^n \}$$

⁹See, e.g., [9] for details

of natural numbers or to the family $\mathbb{B}_n = \{0,1\}^n$ of (0,1)-strings of length n. Hence, if $m = 2^n$, each polynomial of the form

$$p = p_0 x_0 + p_1 x_2 + \ldots + p_{m-1} x_{m-1}$$

may equally well be indexed by the subsets of the n-element set \mathcal{E} or the binary n-strings:

$$p = \sum_{k=0}^{m-1} p_k x_k \quad \longleftrightarrow \quad \sum_{A \in \mathcal{E}} p_A x_A \quad \longleftrightarrow \quad \sum_{\alpha \in \mathbb{B}_n} p_\alpha x_\alpha.$$

The set-theoretic interpretation suggests a polynomial algebra based on Boolean lattice operations:

$$x_A \lor x_B = x_{A \cup B}$$
 and $x_A \land x_B = x_{A \cap B}$.

 (X, \vee) is a monoid with neutral element x_{\emptyset} . (X, \wedge) is a monoid with neutral element x_E if attention is restricted to polynomials indexed by the subsets of E.

Ex. 4 (Möbius algebra) Let
$$X = \{x_A \mid A \subseteq E\}$$
 and set $q = \sum_{A \subseteq E} x_A$. Then

$$\sum_{T \subseteq S} p_T = (p \vee q)_S \quad \textit{and} \quad \sum_{T \supseteq S} p_T = (p \wedge q)_S \quad \textit{for any } p \in \mathbb{C}(X) \textit{ and } S \subseteq E.$$

The example generalizes to lattices (see Ex. 3) in a straightforward manner.

4.3 Concatenation and tensor products

The concatenation $\gamma = \alpha \star \beta$ of a string $\alpha \in \mathbb{B}_n$ with a string $\beta \in \mathbb{B}_m$ is the string

$$\gamma = \gamma_1, \dots, \gamma_n \gamma_{n+1} \dots \gamma_{n+m} \in \mathbb{B}_{n+m}$$

where $\gamma_1 \dots \gamma_n = \alpha$ and $\gamma_{n+1} \dots \gamma_{n+m} = \beta$. Its set-theoretic interpretation is straightforward:

• Let N and M be disjoint sets with n=|N| and m=|M| elements. Then $\gamma=\alpha\star\beta$ represents the set

$$A \cup B \subseteq N \cup M$$
,

if α represents the subset $A \subseteq N$ and β the subset $B \subseteq M$.

Conversely, every string $\gamma \in \mathbb{B}_{n+m}$ decomposes into the product $\gamma = \alpha \star \beta$ of unique strings $\alpha \in \mathbb{B}_n$ and $\beta \in \mathbb{B}_m$.

The polynomial perspective on concatenation leads to the *tensor product* of polynomials. Let X and Y be two sets of indeterminates and denote by $X \otimes Y$ the collection of all pairs $x \otimes y$ of indeterminates $x \in X$ and $y \in Y$.

We define the *tensor product* of $f \in \mathbb{C}(X)$ and $g \in \mathbb{C}(Y)$, in analogy with (2), as the formal product

$$\left(\sum_{x \in X} c_x x\right) \otimes \left(\sum_{y \in Y} d_y y\right) = \sum_{(x,y) \in X \times Y} c_x d_y (x \otimes y). \tag{10}$$

The tensor product is bilinear but, in contrast to the function product, not necessarily commutative. The collection of all tensor products is not closed under taking sums and hence not a vector space in its own right. The tensor products generate the vector space

$$\mathbb{C}(X) \otimes \mathbb{C}(Y) = \mathbb{C}(X \otimes Y) \ (\simeq \mathbb{C}(X \times Y)).$$

4.4 Entanglement

If $f \in \mathbb{C}(X)$ and $g \in \mathbb{C}(Y)$ are polynomials of norm ||f|| = 1 = ||g||, then the polynomial $f \otimes g \in \mathbb{C}(X \otimes Y)$ has norm $||f \otimes g|| = 1$. Intuitively, $f \otimes g$ arises as the *independent* join of f and g. Notice however:

• There may be polynomials $h \in \mathbb{C}(X \otimes Y)$ that cannot be *disentangled*, *i.e.*,

$$h \neq f \otimes g$$
 holds for all $f \in \mathbb{C}(X)$ and $g \in \mathbb{C}(Y)$.

This fact has an important consequence in mathematical modeling. In models where systems states are represented by polynomials, states of the tensor product may possibly not be split into independent components.

Remark 7 States that cannot be disentangled may appear counterintuitive in physical system models and have received considerable attention particularly in quantum theory¹¹.

¹⁰However: if one does not refer explicitly to \mathbb{B}_n and \mathbb{B}_m , the decomposition is *not* unique.

¹¹See, *e.g*, [5].

5 Cooperative games

The model of a cooperative TU-game $\Gamma = (N, v)$ involves a set $N = \{e_1, \dots, e_n\}$ of n players. A subset $S \subseteq N$ is a coalition. Letting \mathcal{N} be the family of all coalitions (subsets) of N, a valuation $v : \mathcal{N} \to \mathbb{R}$ is assumed to be given with v(S) as the value generated by the coalition S. We define the characteristic polynomial of Γ in its set-theoretic notation with indeterminates x_S as

$$p^v = \sum_{S \in \mathcal{N}} v_S x_S$$
 where $v_S = v(S)$.

For its numerical interpretation, recall that the members of the set \mathcal{N} correspond to the $m=2^n$ natural numbers $0 \leq s < m$. So the characteristic polynomial takes the form

$$p^v \longleftrightarrow \sum_{s=0}^{m-1} v_s x^s$$

with the index numbers s corresponding to the coalitions $S \in \mathcal{N}$.

Finally, denoting by $|\sigma\rangle$ the indeterminate x_S corresponding to the coalition $S \in \mathcal{N}$ in its representation as a (0,1)-string $\sigma \in \mathbb{B}_n$, we have

$$p^v \longleftrightarrow \sum_{\sigma \in \mathbb{B}_n} v_{\sigma} | \sigma \rangle.$$

Remark 8 Interpreting the valuation v of the cooperative game $\Gamma = (N, v)$ as a function on $\mathbb{B}_n = \{0, 1\}^n$, v is also known as a pseudo-boolean function 12 . Moreover, v is usually called the characteristic function of Γ . In the present context, however, it is useful to allow the latter terminology with a greater flexibility (see the next Section 5.1).

Remark 9 A more general model of cooperative games with restricted cooperation 13 considers only valuations $v: \mathcal{F} \to \mathbb{R}$ on a given subfamily $\mathcal{F} \subseteq \mathcal{N}$ of coalitions. The "augmenting systems" of [6] are special cases of this model, for example. We leave it to the interested reader to extend the present theory accordingly.

¹²See, *e.g.*, [7].

¹³See [8].

5.1 Characteristic functions

There are various functions associated with cooperative games (relative to the set N of players) which result from appropriate substitutions into the characteristic polynomial p^v . Examples from Section 2.2 are the polynomial function $p^v(t)$ and the wave function v(t), both with one real variable t.

If a complex variable t_S is substituted for each indeterminate x_S , the characteristic polynomial p^v yields a linear functional $\hat{v}: \mathbb{C}^{\mathcal{N}} \to \mathbb{C}$ with values

$$\hat{v}(t) = \sum_{S \in \mathcal{N}} v_S t_S \quad (t \in \mathbb{C}^{\mathcal{N}}).$$

Ex. 5 (Multinomial extensions) Owen's [27] multinomial extension of the valuation $v : \mathcal{N} \to \mathbb{R}$ is obtained if n real variables t_1, \ldots, t_n are associated with the n players e_1, \ldots, e_n and the polynomial functions

$$t_S = \prod_{e_i \in S} t_i \prod_{e_j \notin S} (1 - t_j)$$

are substituted into the indeterminates x_S of the characteristic polynomial:

$$p^{v}(t_1,\ldots,t_n) = \sum_{S \in \mathcal{N}} v_S \prod_{e_i \in S} t_i \prod_{e_j \notin S} (1 - t_j).$$

Similar polynomials are obtained if one considers the set \mathbb{C} of complex numbers (or the set \mathbb{R} of real numbers) to be endowed with a lattice structure $L=(\mathbb{C},\vee,\wedge)$ and replaces the usual scalar addition and multiplication by the lattice operations:

$$c+d\mapsto c\vee d$$
 and $c\cdot d\mapsto c\wedge d$.

Ex. 6 (Sugeno integrals) Assume $L = (\mathbb{R}, \wedge, \vee)$ to be a lattice and define for a valuation $v : \mathcal{N} \to \mathbb{R}$ its lattice polynomial as

$$p^v = \bigvee_{S \in \mathcal{N}} v_S \wedge x_S.$$

Associating real variables t_i with the elements e_i of N as in Ex. 5, the substitution of

$$t_S = \bigwedge_{e_i \in S} t_i$$

then yields the lattice polynomial function

$$p^{v}(t) = \bigvee_{S \in \mathcal{N}} v_S \wedge t_S \quad (t \in \mathbb{R}^{\mathcal{N}}).$$

The Sugeno integral¹⁴ relative to a valuation $v : \mathcal{N} \to \mathbb{R}$ is the lattice polynomial with the lattice operations

$$c \lor d = \max\{c, d\}$$
 and $c \land d = \min\{c, d\}$.

5.2 Activity systems

Suppose that the players $i \in N$ in the cooperative game $\Gamma = (N, v)$ decide to become *active* in Γ with respective probabilities π_i . Then the evaluation

$$p^{v}(\pi_1, \dots, \pi_n) = \sum_{S \in \mathcal{N}} v_S \pi_S$$
 with $\pi_S = \prod_{i \in S} \pi_S$

of the multinomial extension yields the expected value of the ensuing cooperation – provided the players take their individual decisions *independently* from each other¹⁵. In general, *i.e.*, without the independence assumption, the expected value results from the substitution of some probability distribution π on the coalition family \mathcal{N} into the indeterminates of the characteristic polynomial:

$$E^{v}(\pi) = \sum_{S \in \mathcal{N}} v(S)\pi(S). \tag{11}$$

So π describes probabilistically a *state* the cooperative Γ is in and the value it is expected to generate in this state.

As a probabilistic augmentation of the cooperation model, let us think of the members of N simply as agents (players, or coalitions of players, or physical entities etc.) and define an activity $system^{16}$ \mathcal{A}_N relative to a finite set $N=\{1,\ldots,n\}$ of agents with collection $\mathcal N$ of subsets as a system whose states are described by complex vectors $u\in\mathbb C^{\mathcal N}$ of unit length $\|u\|=1$ or, equivalenty, (squared) norm

$$||u||^2 = \sum_{S \in \mathcal{N}} |u_S|^2 = 1.$$

¹⁴See [16, 22] for details.

¹⁵The *cooperative fuzzy games* of [3] make a similar independence assumption on the players.

¹⁶According to the standard interpretation of quantum mechanics, finite-dimensional quantum systems are activity systems in the present sense. (See also Section 7.2).

The (real) parameters $|u_S|^2$ specify a probability distribution on \mathcal{N} . Hence, if $v: \mathcal{N} \to \mathbb{R}$ is a measuring function on \mathcal{N} , its expected observation value will be

$$E^{v}(u) = \sum_{S \in \mathcal{N}} v(S)|u_{S}|^{2}$$
(12)

if A_N is in the activity state u. Conversely, every probability distribution π on \mathcal{N} arises from some unit length vector u^{π} . For example:

$$u^{\pi} = (\sqrt{\pi_S} : S \in \mathcal{N}) \in \mathbb{C}^{\mathcal{N}}.$$

Remark 10 (Hermitian multiplication) The states of A_N correspond to polynomials $s \in \mathbb{C}(X)$ with norm ||s|| = 1. The hermitian product $(s, s) = s \cdot \overline{s}$ has the components $(s, s)_T = |s_T|^2$. So the expected value results from the evaluation of the substitution of v into (s, s).

6 Transforms

A transform on $\mathbb{C}(X)$ is an operator $T:\mathbb{C}(X)\to\mathbb{C}(X)$. For example, the Möbius transform (Example 2) μ is a linear transform. In game theory, the underlying partial order is usually assumed to be set-theoretic:

$$x_A < x_B \iff A \subseteq B$$

and yields a Haranyi-type valuation $w=\mu(v)\in\mathbb{R}^{\mathcal{N}}$ for each $v:\mathcal{N}\to\mathbb{R}$ such that

$$v(B) = \sum_{A \subseteq S} w(A)$$
 for all $B \in \mathcal{N}$.

The idea of the characteristic polynomial suggests also structurally important algebraic transforms on $\mathbb{C}(X)$ that are nonlinear and have not received attention in game theory so far.

Throughout this section, we assume an underlying family \mathcal{N} of $m=2^n$ coalitions S of an n-element player set N, which are represented by the natural numbers $0 \leq s < m$.

6.1 Fourier transforms

For a given scalar $\omega \in \mathbb{C}$, the basic ω -transform of an indeterminate x is its scalar adjustment

$$x \mapsto \omega x$$
.

6.2 The discrete Fourier transform

In the case of the standard polynomial algebra $(\mathbb{C}(X), \cdot)$, the indeterminates have the product decomposition

$$x^s = x^1 \cdot x^1 \cdot \dots \cdot x^1$$
 (s times).

So x^1 is the basic indeterminate. The ω -transform $x^1\mapsto \omega x^1$ implies

$$(\omega x^1)^s = \omega^s x^s$$
 for all s

and is represented by its characteristic polynomial

$$f^{(\omega)} = \sum_{k=0}^{m-1} \omega^k x^k.$$

The associated ω -transform F^{ω} on the whole space $\mathbb{C}(X)$ is the linear transform

$$p \mapsto F^{\omega}(p) = f^{\omega} \cdot p.$$

Since F^{ω} is linear, it suffices to keep track of its effect on the basic polynomials x^s :

$$F^{\omega}(x^s) = f^{\omega} \cdot x^s = \sum_{k=0}^{m-1} \omega^{sk} x^k$$
 for all $s = 0, 1, \dots, m-1$.

For the choice $\omega=e^{2\pi\mathrm{i}/m}$, F^{ω} is known as the discrete Fourier transform on $\mathbb{C}(X)$ and yields for $v\in\mathbb{C}^{\mathcal{N}}$,

$$F^{\omega}(v_0x^0 + v_1x + \ldots + v_{m-1}x^{m-1}) = \hat{v}_0x^0 + \hat{v}_1x + \ldots + \hat{v}_{m-1}x^{m-1}$$

with the Fourier coefficients

$$\hat{v}_j = \sum_{k=0}^{m-1} v_k e^{2\pi i jk/m} \quad (j = 0, 1, \dots, m-1).$$

It is easily verified that the discrete Fourier transform has a Fourier type inverse:

$$v_k = \frac{1}{m} \sum_{j=0}^{m-1} \hat{v}_j e^{-2\pi i jk/m} \quad (k = 0, 1, \dots, m-1).$$

Remark 11 Viewed alternatively, the Fourier coefficients \hat{v}_j are obtained by either substituting the interpolation parameters $t_j = e^{2\pi i j/m}$ into the corresponding polynomial function or by the substitution of the parameters j into the corresponding wave function.

6.3 The quantum Fourier transform

The so-called quantum Fourier transform QFT_n is the discrete Fourier transform in disguise when one replaces the indeterminates x^s by the corresponding binary bit strings $|s\rangle \in \mathbb{B}_n$. The effect on the string $|s\rangle \in \mathbb{B}_n$ is:

$$QFT_n|s\rangle = \sum_{k=0}^{m-1} \omega^{sk}|k\rangle = \sum_{k=0}^{m-1} e^{2\pi i sk/m}|k\rangle.$$

Remark 12 The quantum Fourier transform is a major tool in the theory of quantum computation¹⁷.

6.4 The parity transform

In the polynomial algebra $(\mathbb{C}(X), \odot)$ with the binary multiplication

$$x^s \odot x^k = x^{s \oplus k}$$
 for all $s, k \in \mathbb{B}_n$,

the *n* indeterminates x_1, \ldots, x_n associated with the *n* unit elements in \mathbb{B}_n are basic in the sense

$$x_s = \prod_{i:s_i=1} x_i$$
 and hence $\prod_{i:s_i=1} (\omega x_i) = \omega^{|s|} x_s$,

where $|s| = s_1 + \ldots + s_n$ is the usual sum of the components of the vector $s \in \{0,1\}^n$. In set-theoretic notation, the analogue of the polynomial f^ω is the polynomial

$$h^{\omega} = \sum_{S \in \mathcal{N}} \omega^{|S|} x_S$$

which implies the transform

$$p \mapsto h^{\omega} \odot p$$
.

The choice $\omega = -1$ yields the parity transform

$$p \mapsto \hat{p} = h^{-1} \odot p.$$

with the binary Fourier coefficients

$$\hat{v}_S = \sum_{T \subseteq N} (-1)^{|S \cap T|} v_T \quad \text{for any } v = \sum_{S \subseteq N} v_S x_S \in \mathbb{C}(X).$$

Remark 13 The parity transform is also called binary Fourier transform¹⁸. The Hadamard transform H is the normalized parity transform:

$$H(v) = \frac{1}{\sqrt{m}}\hat{v}$$
 with the self-inverse property $H(H(v)) = v$.

6.5 Galois transforms and zero-dividends

Recall from the Fundamental Theorem of Algebra¹⁹ that the polynomial function

$$v(t) = v_0 + v_1 t + v_2 t^2 + \ldots + v_{k-1} t^{k-1} + t^k$$

with $k \geq 1$, admits a parameter vector $g = (g_1, \dots, g_k) \in \mathbb{C}^k$ such that

$$v(t) = (t + g_1)(t + g_2) \cdots (t + g_k). \tag{13}$$

One thus has

$$v_{k-1} = g_1 + g_2 + \ldots + g_k$$

$$v_{k-2} = g_1 g_2 + \ldots + g_2 g_3 + \ldots + g_{k-1} g_n k$$

$$v_{k-3} = g_1 g_2 g_3 + g_1 g_2 g_4 + \ldots$$

$$\vdots$$

$$v_0 = g_1 g_2 \cdots g_k.$$
(14)

Remark 14 The numbers $z_i = -g_i$ are exactly the zeroes of the polynomial function v(t). So the parameter vector (g_1, \ldots, g_k) is unique up to permutations of coordinates.

If v(t) is the characteristic polynomial function of a valuation $v \in \mathbb{C}^m$, we augment g to a vector in \mathbb{C}^m :

$$g(v) = (g_1, \dots, g_k, \dots, g_m)$$
 with $g_\ell = 1$ for $\ell \ge k$.

We call the valuation g(v) a Galois transform of v.

If $\Gamma = (N, v)$ is a cooperative game with characteristic polynomial function

$$v(t) = v_0 + v_1 t + v_2 t^2 + \ldots + v_k t^k$$

¹⁸For interesting applications of the binary transform in game theory, see, e.g., [20, 26].

¹⁹See, e.g., [32]

and $v_k \neq 0$ we normalize v(t) to the polynomial $\tilde{v}(t) = v(t)/v_k$ with the leading coefficient $\tilde{v}_k = 1$ and call the parameters $z_i = -g_i$ relative to the Galois transform $g(\tilde{v})$ of the normalized game $\tilde{\Gamma} = (N, \tilde{v})$ the zero-dividends of Γ .

Notice that the normalized game $\tilde{\Gamma}$ can be reconstructed from its Galois transform.

Ex. 7 (Zero-normalized games) Cooperative game theory often makes the assumption that a cooperative game (N, v) is zero-normalized, i.e., has the property $v(\emptyset) = 0$. Since $\tilde{v}(\emptyset)$ is the product of the zero-dividends g_i of v, one has

$$v(\emptyset) = 0 \iff g_1 g_2 \cdots g_m = 0$$

Hence v is zero-normalized if and only if $g_i = 0$ holds for some zero-dividend g_i of v.

7 Interaction games

Consider a set $N = \{1, ..., n\}$ of general entities i and specify an *interaction game* on N as a valuation

$$V: N \times N \to \mathbb{R}$$
,

where $V_{ij} = V(i, j)$ is the interaction worth²⁰ of $i, j \in N$. Where X is a set of n indeterminates x_i , the associated characteristic polynomial is

$$\chi^V = \sum_{i=1}^n \sum_{j=1}^n V_{ij}(x_i \otimes x_j),$$

which means that interaction games refer to the tensor product space $\mathbb{C}(X \otimes X)$ (and not to the vector space $\mathbb{C}(X)$) or, more precisely, to the real tensor space $\mathbb{R}(X \otimes X)$. Abstractly, V is given as a real $n \times n$ matrix with the coefficients V_{ij} .

Let $A \in \mathbb{R}^{N \times N}$ be an arbitrary real matrix whose coefficients A_{ij} have the interpretation of enhancing the interaction worth relative to V:

• If i and j interact at level A_{ij} , their interaction produces the value $V_{ij}A_{ij}$.

²⁰See, for example, the *interaction indices* in various application modeling contexts [4, 31].

Hence the interaction instance A produces the game's overall value as

$$\chi^{V}(A) = \sum_{i,j \in N} V_{ij} A_{ij} = \operatorname{tr} (A^{T} V)$$
(15)

where the trace tr (C) of a matrix C is the sum of its diagonal coefficients. In other words, $\chi^V(A)$ equals the usual euclidean inner product (V|A) of the two matrices V,A, considered as n^2 -dimensional parameter vectors. This fact yields the dual interpretations:

- 1. An interaction game V is a linear functional on the space of all interaction instances A.
- 2. An interaction instance A is a linear functional on the space of all interaction games V.

7.1 The hermitian perspective

Setting $A^+ = (A + A^T)/2$, one finds that a matrix $A \in \mathbb{R}^{n \times n}$ decomposes into a symmetric matrix A^+ and a skew-symmetric matrix A^- :

$$A = A^{+} + A^{-}$$
 where $(A^{+})^{T} = A^{+}, (A^{-})^{T} = -A^{-}.$ (16)

Moreover, is it straightforward to check that the symmetry decomposition (16) of A is unique. Associate now with $A \in \mathbb{R}^{n \times n}$ the well-defined *hermitian* matrix

$$\hat{A} = A^+ + iA^- \in \mathbb{C}^{n \times n}$$

and let $\mathbb{H}_n \subseteq \mathbb{C}^{n \times n}$ be the family of all hermitian $n \times n$ matrices.

Clearly, \mathbb{H}_n is isomorphic to $\mathbb{R}^{n\times n}$ with respect to the field \mathbb{R} of real scalars. (\mathbb{H}_n is not a complex vectors space, however.) Recall that the *adjoint* C^* of the complex matrix $C\in\mathbb{C}^{n\times n}$ is the transpose \overline{C}^T of the conjugated matrix \overline{C} and note

Lemma 1 For any matrices $C \in \mathbb{C}^{n \times n}$ and $A, B \in \mathbb{R}^{n \times n}$, one has

- 1. $C \in \mathbb{H}_n$ if and only if $C^* = C$ (i.e., C is self-adjoint).
- 2. $(A|B) = \operatorname{tr}(B^T A) = \operatorname{tr}(\hat{B}\hat{A}) = (\hat{A}|\hat{B}).$

Hence:

• Interaction games can equally well be studied within the context of the hermitian matrix space \mathbb{H}_n .

The adjoint u^* of the (column) vector $u \in \mathbb{C}^n$ is the row vector of conjugated components of u. So uu^* is a self-adjoint $n \times n$ matrix. Indeed, an important general characterization of self-adjoint matrices is:

Lemma 2 The matrix $C \in \mathbb{C}^{n \times n}$ is self-adjoint if and only if there are real scalars λ_k and vectors $u_k \in \mathbb{C}^n$ such that

$$C = \sum_{k} \lambda_k u_k u_k^*. \tag{17}$$

Proof. Because the matrices $u_k u_k^*$ are self-adjoint, a matrix C of the form (17) is self-adjoint. To see the converse, recall that $\mathbb{R}^{n\times n}$ and \mathbb{H}_n are isomorphic. So it suffices to consider (0,1)-matrices $A \in \mathbb{R}^{n \times n}$ with exactly one non-zero entry $A_{ij} = 1$. Moreover, we may assume n = 2.

If A is diagonal, one has $\hat{A} = A$, for which the claim is obviously true. So assume, for example,

$$A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$
 and hence $C = \hat{A} = \frac{1}{2} \begin{pmatrix} 0 & 1+\mathrm{i} \\ 1-\mathrm{i} & 0 \end{pmatrix}$

with the two real eigenvalues $\lambda_1 = +\sqrt{2}$ and $\lambda_2 = -\sqrt{2}$. Consequently, the 2dimensional space \mathbb{C}^2 admits a basis of eigenvectors of C, which imply the claim.

Remark 15 The spectral decomposition of self-adjoint matrices²¹ shows that the vectors u_k in (17) can be chosen as eigenvectors of C with real eigenvalues λ_k also in the case n > 2.

Consider, for example, an activity system A_N as in Section 5.2 relative to the valuation $v: \mathcal{N} \to \mathbb{R}$. Let $u \in \mathbb{C}^{\mathcal{N}}$ be a state vector of norm ||u|| = 1 with the associated self-adjoint $m \times m$ matrix $U = uu^*$ of complex coefficients

$$U_{ST} = u_S \overline{u_T}.$$

²¹See, *e.g.*, [33].

Where V = diag(v) is the diagonal interaction matrix with diagonal coefficients $V_{SS} = v(S)$, one now observes:

$$(V|U) = \sum_{S \subseteq N} v(S)\overline{u_S}u_S = \sum_{S \subseteq N} v(S)|u_S|^2 = E^v(u).$$

So \mathcal{A}_N is recognized as the restriction of an interaction game on \mathcal{N} with characteristic function $V = \operatorname{diag}(v)$ to interaction instances of the form uu^* .

7.2 Heisenberg measurements

A measurement operator on the interaction system N is a functional

$$\mu: \mathbb{R}^{N \times N} \to \mathbb{R}$$

which produces the measurement result $\mu(A)$ if the system is in a state that corresponds to the activity instance A. If the functional μ is linear, the measurement acutally represents an interaction game. So there exists a matrix $M \in \mathbb{R}^{n \times n}$ such that

$$\mu(A) = (M|A) = (\hat{M}|\hat{A}),$$

which means that μ can be understood as a linear functional on the (real) vector space \mathbb{H}_n of all hermitian matrices. Assuming

$$\hat{A} = \sum_{k} \lambda_k u_k u_k^*$$

for suitable real parameters λ_k and (complex) vectors $u_k \in \mathbb{C}^n$, one has

$$\mu(A) = \sum_{k} \lambda_k \mu(u_k u_k^*) = \sum_{k} \lambda_k (\hat{M} | u_k u_k^*). \tag{18}$$

A Heisenberg measurement operator²² on the (complex) vector space \mathbb{C}^n is a real-valued functional $\gamma: \mathbb{C}^n \to \mathbb{R}$ of the form

$$\gamma(u)=u^*Gu=(G|uu^*)\quad\text{with }G\in\mathbb{C}^{n\times n}.$$

Since γ is real-valued, one may assume that G is self-adjoint. Notice that a Heisenberg measurement operator is not linear. On the other hand, (18) shows:

²²"Heisenberg" refers to the fact that this measurement model is standard in quantum theory.

• A Heisenberg measurement operator on \mathbb{C}^n arises from the restriction of a linear measurement operator on the interaction system N to interaction instances of the hermitian form uu^* .

Similarly, the linear measurement matrix \hat{M} of the operator μ in (18) is of the form

$$\hat{M} = \sum_{j} \delta_{j} w_{j} w_{j}^{*}$$

for suitable real numbers δ_i and vectors $w_i \in \mathbb{C}^n$, which implies

$$\mu(A) = \sum_{i} \sum_{j} \lambda_{i} \mu_{j}(w_{j} w_{j}^{*} | u_{i} u_{i}^{*}).$$
(19)

Consequently, the fundamental linear interaction measures are recognized to be of the form

$$\mu_w(u) = (ww^*|uu^*) = |w^*u|^2 \quad \text{with} \quad w, u \in \mathbb{C}^n.$$
 (20)

8 Discussion

Linearity plays an important role in mathematical application models. The mathematical analysis, however, will reveal more characteristic features of the underlying systems when the model is not just considered to be a vector space with scalar multiplication, but an algebra, *i.e.*, additionally equipped with a multiplication operation for vectors. Suitable multiplication operations are naturally associated with multiplication rules for polynomials, which renders polynomial models powerful and flexible.

The present approach shows that mathematical models for cooperation and interaction connect with important aspects of classical algebra and combinatorics. For example, the representation of coalitions by natural numbers embeds the representation of cooperative games into the context of Galois theory, *i.e.*, the theory of solving algebraic equations. Future work of exploring this area of mathematical system analysis in more detail appears to be promising.

The polynomial model also underlines the aspect of quantum-theoretic models as interaction systems and, conversely, embeds cooperation and interaction into the setting of physical quantum systems. The evolution of such systems can be mathematically understood in a far broader context (see, *e.g.*[10]). Moreover,

the apparatus of theoretical physics can be brought to bear on general systems of cooperation and interaction. In particular, Hamiltonians of cooperative games can be expected to provide considerable insight into fundamental laws according to which such systems behave.

References

- [1] M. AIGNER, Combinatorial Theory, Springer, 1997.
- [2] M. AIGNER, *A Course in Enumeration*, Springer, 2007, https://doi.org/10.1007/978-3-540-39035-0.
- [3] J.P. AUBIN: *Cooperative fuzzy games*, Math. of Operations Research **6** (1981), 1–13.
- [4] G. BELIAKOV, S. JAMES AND J.-Z. WU: Value and Interaction Indices. In: Discrete Fuzzy Measures. Studies in Fuzziness and Soft Computing **382**, Springer, 2019.
- [5] J. S. Bell: On the Einstein-Podolsky-Rosen paradox, Physics 1 (1964), 195-200.
- [6] J.M. BILBAO: *Cooperative games under augmenting systems*, SIAM J. Discrete Math. **17** (2003), 122–133.
- [7] E. BOROS, E AND P.L.HAMMER, *Pseudo-boolean optimization*, Discrete Appl. Math. **123** (2002), 155–225. doi:10.1016/S0166-218X(01)00341-9
- [8] U. FAIGLE: *Cores of games with restricted cooperation*, ZOR-Methods and Models of Operations Research **33** (1989), 405–422.
- [9] U. FAIGLE, *Mathematical Game Theory*, World Scientific, New Jersey, 2022, ISBN 9789811246692.
- [10] U. FAIGLE AND G. GIERZ: Markovian statistics on evolving systems, Evolving Systems 99 (2018), 213-–225. https://doi.org/10.1007/s12530-017-9186-8
- [11] U. FAIGLE AND M. GRABISCH: Bases and linear transforms of TU-games and cooperation systems, Int. J. of Game Theory **45** (2016), 875–892.

- [12] U. FAIGLE AND M. GRABISCH: *Game-theoretic interaction and decision:* A quantum analysis, Games **8** (2017), DOI:10.3390/g8040048.
- [13] U. FAIGLE AND M. GRABISCH: *Polynomial representation of TU-games*, arXiv (2024), https://arxiv.org/abs/2401.12741
- [14] U. FAIGLE, W. KERN AND G. STILL, Algorithmic Principles of Mathematical Programming. Springer, 2002, DOI 10.1007/978-94-015-9896-5.
- [15] W. FELLER, An Introduction to Probability Theory and Its Applications I, Wiley, New York, 1968.
- [16] M. GRABISCH, Set Functions, Games and Capacities in Decision Making, Springer 2016.
- [17] J.C. HARSANYI: A simplified bargaining model for the n-person cooperative game, International Economic Review 4 (1967), 194–220.
- [18] P. L. HAMMER AND S. RUDEANU, Boolean Methods in Operations Research and Related Areas. Springer, 1968.
- [19] K. IKEDA AND S. AOKI: *Infinitely repeated quantum games and strategic efficiency*, Quantum Inf. Process **20**, 387 (2021). https://doi.org/10.1007/s11128-021-03295-7
- [20] G.KALAI: A Fourier-theoretic perspective on the Condorcet paradox and Arrow's theorem, Advances in Applied Mathematics 29 (2002), 412–426.
- [21] D.G. LUENBERGER, *Optimization by Vector Space Methods*. Wiley Professional Paperback Series, 1998.
- [22] J.-L. MARICHAL: Weighted lattice polynomials, Discrete Mathematics 309 (2009), 814–820.
- [23] J. VON NEUMANN AND O. MORGENSTERN, *Theory of Games and Economic Behavior*, Princeton University Press, 1944.
- [24] N. NISAN, T. ROUGHGARDEN, É. TARDOS AND V.V. VAZIRANI, (EDS.) *Algorithmic Game Theory*, Cambridge University Press, Cambridge, 2007.

- [25] NIELSEN AND I. CHUANG, *Quantum Computation*. Cambrigde University Press, 2000.
- [26] R. O'DONNELL, *Analysis of Boolean Functions*. Cambridge University Press, 2014.
- [27] G. OWEN: Multilinear extension of games, Management Sci. 18 (1972), 64–79.
- [28] E.W. PIOTROWSKI AND J. SŁADKOWSKI: (2003). An invitation to quantum game theory,. International Journal of Theoretical Physics. Springer Nature {bf 42} (2003) 1089-–1099. doi:10.1023/a:1025443111388. ISSN 0020-7748. S2CID 13630647.
- [29] G. C. Rota. On the foundations of combinatorial theory I. Theory of Möbius functions. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete* 2 (1964), 340–368.
- [30] G.-C. ROTA, D. KAHANER AND A. ODLYZKO: *Finite operator calculus*, Journal of Mathematical Analysis and its Applications **42**, June 1973. Reprinted in the book with the same title, Academic Press, New York, 1975.
- [31] *Interaction index an overview*, https://www.sciencedirect.com/topics/mathematics/interaction-index.
- [32] B.L. VAN DER WAERDEN, *Algebra I*, Springer Heidelberg, DOI https://doi.org/10.1007/978-3-642-85527-6
- [33] J. WEIDMANN, *Linear Operators in Hilbert Space*, Graduate Texts in Mathematics **68**, Springer, 2012.