

Strategy Complexity of Büchi Objectives in Concurrent Stochastic Games

Stefan Kiefer
University of Oxford
Oxford, United Kingdom

Mahsa Shirmohammadi
IRIF & CNRS, Université Paris cité
Paris, France

Richard Mayr
University of Edinburgh
Edinburgh, United Kingdom

Patrick Totzke
University of Liverpool
Liverpool, United Kingdom

ABSTRACT

We study 2-player concurrent stochastic Büchi games on countable graphs. Two players, Max and Min, seek respectively to maximize and minimize the probability of visiting a set of target states infinitely often. We show that there always exist ε -optimal Max strategies that use just a step counter plus 1 bit of public memory. This upper bound holds for all countable graphs, but it is a new result even for the special case of finite graphs. The upper bound is tight in the sense that Max strategies that use just a step counter, or just finite memory, are *not* sufficient even on finite game graphs.

The upper bound is a consequence of a slightly stronger new result: ε -optimal Max strategies for the combined Büchi and Transience objective require just 1 bit of public memory (but cannot be memoryless). Our proof techniques also yield a closely related result, that ε -optimal Max strategies for the Transience objective alone (which is only meaningful in infinite graphs) can be memoryless.

CCS CONCEPTS

- **Theory of computation** → Random walks and Markov chains;
- **Mathematics of computing** → Probability and statistics.

KEYWORDS

Stochastic Games, Strategy Complexity, Büchi and Transience Objectives, ε -Optimal Strategies

1 INTRODUCTION

Background. We study 2-player zero-sum concurrent stochastic games on countable (finite or countably infinite) graphs.¹ Introduced by Shapley in his seminal 1953 work [28], and generalized in [12] and [18] to allow infinite state and action sets and non-termination, such games play a central role in the solution of many problems in economics, see [3, 13, 24, 29, 30], evolutionary biology, e.g., [26], and computer science, see [1, 2, 5–7, 9, 23, 29, 31] among others.

Here we consider event-based objectives (rather than general measurable reward functions). The Büchi objective is to visit some

¹Our proofs do not carry over to uncountable state spaces. Apart from the usual issues with measurability in uncountable systems, we also, e.g., partition events into as many parts as there are states (in our case only countably many) and then rely on sigma-additivity.

given subset of the states infinitely often. The Transience objective is to visit no state infinitely often, i.e., every single state is visited at most finitely often (if at all). The players, Max and Min, seek respectively to maximize and minimize the probability of the objective.

A central result in zero-sum 2-player stochastic games with finite action sets is the existence of a *value* for the large class of Borel measurable objectives [19, 21]. That is, for Borel measurable functions f on plays,

$$\sup_{Max} \inf_{Min} f = \text{value} = \inf_{Min} \sup_{Max} f$$

over Max and Min strategies. This holds even under the weaker assumption that one player has countable action sets and the other plays has finite action sets [21][11, Theorem 11]. Throughout this paper we assume that Max has countable action sets and Min has finite action sets. (While the reverse case also has value, our results do not hold for it; see below.)

The amount/type of memory and randomization required for a good (ε -optimal, resp. optimal) strategy for a given player and objective is also called the *strategy complexity*.

Optimal Max strategies for Büchi objectives do not exist in general, not even for the simpler reachability objective in finite-state concurrent games [10]. However, in finite-state concurrent Büchi games, if every state admits an optimal Max strategy then Max also has a memoryless strategy that is optimal from all states [5, Proposition 17]. Optimal Max strategies for Transience do not exist even in countable Markov decision processes (MDPs) [16], and Transience trivially never holds in finite-state MDPs/games. Thus we consider the strategy complexity of ε -optimal strategies.

Our contribution. We show (Theorem 3.1) that ε -optimal Max strategies for the Büchi objective in countable concurrent stochastic games require just a step counter plus 1 bit of public memory. This is a new result even for the subclass of *finite-state* concurrent stochastic games with finitely many actions, where the strategy complexity of Büchi objectives had also been open. (About the strategy complexity for the opposing Min player see Section 7.)

The step counter plus 1 bit upper bound on the strategy complexity of ε -optimal Max strategies is tight, even for finite-state concurrent Büchi games. In the “Bad Match” game [20, Example 13.7], neither finite-memory strategies nor Markov strategies (i.e., strategies that use just a step counter) can be ε -optimal for Max for any $\varepsilon < 1$; see Section 4. (The fact that finite-memory Max strategies are insufficient in finite-state concurrent Büchi games was mentioned (without proof) in [8], who considered a different



counterexample, namely a repeated version of the classic “Snowball game” (aka Hide-or-Run game) of Kumar and Shiau [18].) In the special case of finite-state *turn-based* Büchi games, the strategy complexity is lower, because both players have optimal memoryless deterministic strategies. For countable MDPs, different counterexamples show that neither finite-memory strategies nor Markov strategies can be ε -optimal for Max for the Büchi objective [14].

Our upper bound (Theorem 3.1) is a consequence of a stronger new result (Theorem 3.2) on the strategy complexity of the *combined* Büchi and Transience objective in countable concurrent stochastic games with finite Min action sets: ε -optimal Max strategies require just 1 bit of (public) memory. This upper bound is tight, even for acyclic countable MDPs, in the sense that memoryless Max strategies do not suffice [14]. (Note that plays can only satisfy this combined objective if the target set of the Büchi objective is infinite). The stronger result on the combined Büchi and Transience objective implies the result on the Büchi objective above as follows. We can apply the stronger result to a new derived acyclic game where the step counter is encoded into the states (and thus Transience always holds), and carry the obtained 1-bit Max strategy back to the original game as a step counter plus 1 bit strategy; see Appendix B.

Our upper bound on the strategy complexity of the combined Büchi and Transience objective in countable concurrent stochastic games generalizes a corresponding result for countable MDPs [16, Lemma 4]. Similarly to [16], 1 bit of public memory is used to switch between two modes. Roughly speaking, one mode aims to visit the target set in the short term while the other mode records that this short-term goal has been attained. However, the proof techniques for stochastic games are very different from those used for MDPs. Unlike in MDPs, Max cannot statically plan ahead which states in the Büchi target set will be visited within a given time horizon with high probability, because of the influence of Min’s decisions. Similarly, for transience, Max cannot statically plan that certain states are very unlikely to be visited again after a given time horizon. Again it may depend on Min’s decisions whether certain states are (re-)visited arbitrarily late in the game. Therefore, our construction of the Max strategy is based on a different principle. It proceeds state by state (based on an enumeration of the states), rather than phase by phase as was done for MDPs in [16].

Our third result (Theorem 3.3) concerns the strategy complexity of the Transience objective alone (which is only meaningful in infinite-state games). There always exist memoryless ε -optimal Max strategies for Transience in countable concurrent stochastic games. This generalizes a corresponding result for countable MDPs [16, Theorem 8]. However, unlike for MDPs, these strategies cannot be deterministic in concurrent games, i.e., they need to randomize.

Finally, our upper bounds do not carry over to countable stochastic games with countable Min action sets. Even for the simpler reachability objective, there is a counterexample where Max strategies that just use a step counter plus finite (private) memory are not ε -optimal for any $\varepsilon < 1$. In fact, the counterexample is a countably infinite-state *turn-based* stochastic reachability game with countably infinite Min branching and finite Max branching [17].

2 PRELIMINARIES

A *probability distribution* over a countable set S is a function $\alpha : S \rightarrow [0, 1]$ with $\sum_{s \in S} \alpha(s) = 1$. The *support* of α is the set $\{s \in S \mid \alpha(s) > 0\}$. We write $\mathcal{D}(S)$ for the set of all probability distributions over S .

Stochastic Games. We study stochastic games between two players, Max and Min. A (*concurrent*) *game* \mathcal{G} is played on a countable set of states S . For each state $s \in S$ there are non-empty countable *action* sets $A(s)$ and $B(s)$ for Max and Min, respectively. A *mixed action* for Max (resp. Min) in state s is a distribution over $A(s)$ (resp. $B(s)$). Let $A \stackrel{\text{def}}{=} \bigcup_{s \in S} A(s)$ and $B \stackrel{\text{def}}{=} \bigcup_{s \in S} B(s)$.

Let $Z \stackrel{\text{def}}{=} \{(s, a, b) \mid s \in S, a \in A(s), b \in B(s)\}$. For every triple $(s, a, b) \in Z$ there is a distribution $p(s, a, b) \in \mathcal{D}(S)$ over successor states. We call a state $s \in S$ a *sink* state if $p(s, a, b)(s) = 1$ for all $a \in A(s)$ and $b \in B(s)$. We extend the *transition function* p to mixed actions $\alpha \in \mathcal{D}(A(s))$ and $\beta \in \mathcal{D}(B(s))$ by letting

$$p(s, \alpha, \beta) \stackrel{\text{def}}{=} \sum_{a \in A(s)} \sum_{b \in B(s)} \alpha(a) \beta(b) p(s, a, b),$$

which is a distribution over S . Let $s_0 \in S$ be an initial state; a *play* from s_0 is an infinite sequence in Z^ω where the first triple contains s_0 . Starting from s_0 , the game is played in steps $\{0, 1, 2, \dots\}$, where at every step $n \in \mathbb{N}$, the play is in some state s_n . At step n , Max and Min independently and simultaneously choose mixed actions $\alpha_n \in \mathcal{D}(A(s_n))$ and $\beta_n \in \mathcal{D}(B(s_n))$. Then two actions $a_n \in A$ and $b_n \in B$ are sampled according to α_n and β_n , respectively, a state $s_{n+1} \in S$ is chosen according to $p(s_n, a_n, b_n)$, before both players are informed of the outcomes a_n, b_n, s_{n+1} and the next step starts from s_{n+1} . The game is called *turn-based* if every $p(s, a, b)$ either solely depends on (s, a) or solely on (s, b) and thus, for every state s , only the action of one player matters.

Strategies. We write H_n for the set of *histories* at step $n \in \mathbb{N}$, which is defined as $H_0 \stackrel{\text{def}}{=} S$ and $H_n \stackrel{\text{def}}{=} Z^n \times S$ for all $n > 0$. The set of all histories is $H \stackrel{\text{def}}{=} \bigcup_{n \in \mathbb{N}} H_n$. For each history $h = (s_0, a_0, b_0) \cdots (s_{n-1}, a_{n-1}, b_{n-1}) s_n \in H_n$, let $s_h \stackrel{\text{def}}{=} s_n$ denote its final state. Define the *length* of h by $|h| \stackrel{\text{def}}{=} n$. For any history h' starting in state s_h , we write hh' for the concatenation of the histories $(s_0, a_0, b_0) \cdots (s_{n-1}, a_{n-1}, b_{n-1}) h'$.

A *strategy* for Max is a function σ that to each history $h \in H$ assigns a mixed action $\sigma(h) \in \mathcal{D}(A(s_h))$. It is called *deterministic* if $\sigma(h)$ is a Dirac distribution for every history h . Mixed actions and strategies for Min are defined analogously. We write Σ and Π for the sets of strategies for Max and Min, respectively.

Probability Measures and Optimality. An initial state s_0 and a pair of strategies σ, π for Max and Min induce a probability measure where the events, also called *objectives* here, are measurable sets of plays (see e.g. [11]). We write $\mathcal{P}_{\mathcal{G}, s_0, \sigma, \pi}(E)$ for the probability of the event E starting from s_0 . It is initially defined for the cylinder sets generated by the histories and then extended to the sigma-algebra by Carathéodory’s unique extension theorem [4]. Given some Borel measurable reward function v on plays, we will write $\mathcal{E}_{\mathcal{G}, s_0, \sigma, \pi}$ for the expectation w.r.t. $\mathcal{P}_{\mathcal{G}, s_0, \sigma, \pi}$ and v . We may drop \mathcal{G} from the subscript when it is understood. For a game \mathcal{G} and objective E fixed,

the *lower value* and *upper value* of a state s_0 are defined as

$$\begin{aligned} \text{val}_{\mathcal{G},E}^{\downarrow}(s_0) &\stackrel{\text{def}}{=} \sup_{\sigma \in \Sigma} \inf_{\pi \in \Pi} \mathcal{P}_{\mathcal{G},s_0,\sigma,\pi}(E) \quad \text{and} \\ \text{val}_{\mathcal{G},E}^{\uparrow}(s_0) &\stackrel{\text{def}}{=} \inf_{\pi \in \Pi} \sup_{\sigma \in \Sigma} \mathcal{P}_{\mathcal{G},s_0,\sigma,\pi}(E). \end{aligned}$$

The inequality $\text{val}_{\mathcal{G},E}^{\downarrow}(s_0) \leq \text{val}_{\mathcal{G},E}^{\uparrow}(s_0)$ holds by definition. If $\text{val}_{\mathcal{G},E}^{\downarrow}(s_0) = \text{val}_{\mathcal{G},E}^{\uparrow}(s_0)$, then this quantity is called the *value* of s_0 , denoted by $\text{val}_{\mathcal{G},E}(s_0)$. For all Borel objectives, the value exists if all action sets are finite, and even if, for all states s , we have that $A(s)$ is finite or $B(s)$ is finite [21][11, Theorem 11]. We always assume the latter, so that $\text{val}_{\mathcal{G},E}(s_0)$ exists.

For $\varepsilon \geq 0$, a Max strategy σ is called ε -*optimal* from s_0 if for all Min strategies π we have $\mathcal{P}_{\mathcal{G},s_0,\sigma,\pi}(E) \geq \text{val}_{\mathcal{G},E}(s_0) - \varepsilon$. It is called *multiplicatively ε -optimal* if it satisfies $\mathcal{P}_{\mathcal{G},s_0,\sigma,\pi}(E) \geq \text{val}_{\mathcal{G},E}(s_0)(1 - \varepsilon)$. Clearly, σ is ε -optimal if it is multiplicatively ε -optimal. A 0-optimal strategy is also called *optimal*.

Objectives. Given a set $T \subseteq S$ of target states, the *reachability* objective $\text{Reach}(T)$ is the set of plays that visit T at least once, i.e., $s_h \in T$ holds for some history h that is a prefix of the play. The objective $\text{Reach}_L(T)$ additionally requires that all states in h before the visit to T are in L . The dual *safety* objective $\text{Avoid}(T) \stackrel{\text{def}}{=} Z^\omega \setminus \text{Reach}(T)$ consists of the plays that never visit T . We are mainly interested in *Büchi* objectives $\text{Bu}(T)$ that consist of all those plays that visit the set of target states $T \subseteq S$ infinitely often. (If the set T is infinite then some plays that satisfy $\text{Bu}(T)$ might not visit any particular state in T infinitely often.) Whenever possible, we will assume a fixed target set T and simply write Reach , Avoid and Bu for these objectives. The *transience* objective Tr contains those plays in which every state of the game is eventually avoided. That is, a play $(s_0, a_0, b_0)(s_1, a_1, b_1) \cdots \in Z^\omega$ is in Tr if for every $s \in S$ there exist n_s so that $s \neq s_i$ for $i \geq n_s$. The *transient Büchi* objective is $\text{TB} \stackrel{\text{def}}{=} \text{Tr} \cap \text{Bu}$. Clearly, TB can only be met if T is infinite.

Strategy Classes. We formalize the amount of *memory* needed to implement strategies. A Max (analogously Min) strategy is *based on* a set M of *memory modes* if it can be described by an initial mode $m_0 \in M$ and a pair of functions

$$\begin{aligned} \sigma_{act} : M \times S &\rightarrow \mathcal{D}(A) \quad \text{and} \\ \sigma_{up} : S \times M \times A \times B \times S &\rightarrow \mathcal{D}(M), \end{aligned}$$

selecting mixed actions and updating the memory mode, respectively. For any history $h = (s_0, a_0, b_0) \cdots (s_{n-1}, a_{n-1}, b_{n-1})s_n \in H_n$, $\sigma(h) = \sigma_{act}(m_n, s_n)$ and m_{n+1} is drawn randomly according to the distribution $\sigma_{up}(s_n, m_n, a_n, b_n, s_{n+1})$. For M -based strategy σ and $m \in M$ we will write $\sigma[m]$ to denote the strategy with σ_{act} and σ_{up} as in σ but with initial mode $m_0 = m$.

A memory-based strategy is *public* if at each step of the game, its current memory mode is known to the opponent, and *private* otherwise. Note that if all memory updates $\sigma_{up}(\cdot)$ are Dirac then σ is public, since the opponent can keep track of its memory.

A strategy σ is *memoryless* if it is based on a singleton set M , i.e., $\sigma(h) = \sigma(h')$ for any two histories h, h' with $s_h = s_{h'}$. A strategy is *1-bit* if it is based on $M = \{0, 1\}$. A *step counter* is a special case of infinite memory in the form of a discrete clock (by default starting as 0) that gets incremented by 1 at every step, independently of

the actions of the players. Strategies that use just a step counter are also called *Markov* strategies. A strategy σ is *1-bit Markov* if it is based on a step counter and two extra memory modes. That is, $M = \mathbb{N} \times \{0, 1\}$ and at any step n the update function σ_{up} guarantees that the mode m_n is in $\{n\} \times \{0, 1\}$.

3 THE MAIN RESULTS

Our results concern the memory requirement of Max strategies for the Büchi (Bu), Transience (Tr) and Transient Büchi (TB) objectives in concurrent games.

Throughout the paper, \mathcal{G} refers to a concurrent 2-player game with a countable state space S , where Max has countable action sets and Min has finite action sets.

THEOREM 3.1. *Let \mathcal{G} be a game with Büchi objective and $\varepsilon > 0$. Max has a 1-bit Markov strategy σ such that*

- (1) $\sigma[0]$ is multiplicatively ε -optimal from every state,² and
- (2) all memory updates $\sigma_{up}(\cdot)$ are Dirac (hence the memory is public)

Moreover, if \mathcal{G} is turn-based then there exists such a strategy that is deterministic.

The 1-bit Markov upper bound for the Büchi objective of Theorem 3.1 is a new result even for the special case of *finite-state* concurrent Büchi games. Moreover, this upper bound is tight in the sense that Markov strategies or finite-memory strategies are not sufficient, neither for finite-state concurrent games [20, Example 13.7], [8] nor for countable MDPs [14]. We re-visit some of these lower bounds in Section 4 where we discuss in detail the example of the “Bad Match”, in a slightly adapted formulation as a finite-state concurrent Büchi game.

Our strongest result is that for transient Büchi objectives: Max has public 1-bit strategies that are uniformly ε -optimal (i.e., ε -optimal from every state).

THEOREM 3.2. *Let \mathcal{G} be a game with TB objective and $\varepsilon > 0$. Max has a 1-bit strategy σ so that*

- (1) $\sigma[0]$ is multiplicatively ε -optimal from every state, and
- (2) all memory updates $\sigma_{up}(\cdot)$ are Dirac (hence the memory is public).

Moreover, if \mathcal{G} is turn-based then there exists such a strategy that is deterministic.

This implies Theorem 3.1. Indeed, following [16], for any game \mathcal{G} on S we can consider the game on the product $S \times \mathbb{N}$ in which the second component acts as the step counter. The resulting game \mathcal{G}' is acyclic and therefore all plays are transient. By applying Theorem 3.2 to \mathcal{G}' , we obtain an ε -optimal 1-bit Max strategy, which can be carried back to \mathcal{G} as a 1-bit Markov strategy. See Lemma B.1 in Appendix B for more details.

Our third main contribution is that in games with Transience objectives, Max has memoryless strategies that are uniformly ε -optimal.

²Recall that $\sigma[0]$ means that the 1-bit memory is initially set to 0. In contrast, the initial value of the step counter does not matter. The strategy is multiplicatively ε -optimal from every state, regardless of the initial value of the step counter, and thus in particular in the default case where the step counter is initially 0.




Figure 1: The Bad Match. Max and Min actions, where they influence the motion, are depicted in blue and red, respectively. The objective is $\text{Bu}(\{w, s\})$. No Markov or finite-memory strategy for Max is ε -optimal from d , for any $\varepsilon < 1$.

THEOREM 3.3. *Let \mathcal{G} be a game with Tr objective and $\varepsilon > 0$. Max has a memoryless strategy that is multiplicatively ε -optimal from every state.*

Moreover, if \mathcal{G} is turn-based then there exists such a strategy that is deterministic.

The technique we use to prove this result is similar to, but simpler than the one used for Theorem 3.2. We therefore first focus on transience games in Section 5.

4 LOWER BOUNDS

The game depicted in Figure 1 is called the “Bad Match”. It was originally discussed with an expected lim sup objective with state-based rewards in $\{-1, 0, 1\}$ [20, 32]. Here we slightly adapt its presentation as a concurrent Büchi game. There are action sets $A = B = \{0, 1\}$ and the objective is Bu with target states $\{s, w\}$. The initial state is d . The transition distributions are Dirac and satisfy $p(d, 1, 1)(w) = p(d, 1, 0)(l) = p(d, 0, 1)(t) = p(d, 0, 0)(s) = 1$; the motion from states w, l, s, t follows the unique depicted edge under all actions. Intuitively, w and l are winning and losing sinks respectively, whereas s and t represent a temporary win and loss for Max, respectively.

In this game, all states except l have value 1 by [20, Lemma 13.8]; see also Proposition 4.4 below. In the following we consider the strategy complexity of the Bad Match from Max’s point of view. First, Max does not have any optimal strategy from d , but only ε -optimal strategies.

PROPOSITION 4.1. *In the Bad Match, Max does not have any optimal strategy from state d .*

PROOF. Either Max always plays action 0, in which case Min can win by always playing action 1. Or, otherwise, there exists some finite history h in which Max always plays action 0, after which Max plays action 1 with positive probability $\delta > 0$ for the first time. Then Min can make Max’s attainment $\leq 1 - \delta < 1$ by first playing according to h , and then playing action 0. \square

The following two propositions state lower bounds for the complexity of Max strategies: neither finite-memory strategies nor Markov strategies for Max are ε -optimal.

PROPOSITION 4.2. *Every finite-memory Max strategy σ in the Bad Match is worthless: $\inf_{\pi \in \Pi} \mathcal{P}_{\mathcal{G}, d, \sigma, \pi}(\text{Bu}) = 0$.*

PROOF. Let σ be a Max strategy based on some finite memory set M , and fix an $\varepsilon > 0$. We show that Min can win with probability $1 - \varepsilon$, i.e., that there is a Min strategy π so that $\mathcal{P}_{\mathcal{G}, d, \sigma, \pi}(\text{Bu}) \leq \varepsilon$.

Choose $\varepsilon_1, \varepsilon_2 > 0$ with $\varepsilon = \varepsilon_1 + \varepsilon_2$ and first consider the Min strategy, π_1 , which in each step plays action 1 with probability ε_1 and action 0 with probability $1 - \varepsilon_1$. Then σ and π_1 induce a Markov chain with finite state space $S \times M$. Since π_1 plays both actions with constant nonzero probabilities, all states (x, m) in a bottom strongly-connected component (BSCC) of this Markov chain must satisfy one of the following properties. Either $x \in \{w, l\}$ or m is a Max memory mode from which Max can never play action 1 in the future. By standard properties of finite-state Markov chains, there is a number K such that, independently of the start state, the probability of having entered a bottom strongly-connected component (BSCC) in the first K steps is at least $1 - \varepsilon_2$. In particular, K depends only on the induced Markov chain, and thus only on \mathcal{G} and the strategies σ and π_1 . Now we define Min’s counter-strategy π , that depends on K and thus on σ . The Min strategy π plays like π_1 in the first K steps and action 1 always thereafter.

Consider the strategies σ and π . The probability that within K steps the permanently winning state w is reached is at most ε_1 ; this follows from the definition of π_1 . The probability that within K steps the finite-state Markov chain has not yet reached a BSCC is at most ε_2 . It follows that the probability that within K steps Max has permanently lost or reaches a memory mode from where he will never play action 1 is at least $1 - \varepsilon_1 - \varepsilon_2 = 1 - \varepsilon$. Thus, Min wins with probability at least $1 - \varepsilon$. \square

Notice that Proposition 4.2 even holds for all private finite-memory Max strategies.

We repeat an argument of [20, Lemma 13.9], that similarly, all Markov strategies are worthless for Max.

PROPOSITION 4.3. *Every Markov strategy σ for Max in the Bad Match is worthless: $\inf_{\pi \in \Pi} \mathcal{P}_{\mathcal{G}, d, \sigma, \pi}(\text{Bu}) = 0$.*

PROOF. Let σ be a Markov strategy for Max and for any step $n \geq 0$, let r_n be the probability with which σ chooses action 1 at step $2n$ (in state d). There are two cases, whether or not the sum of r_n converges or diverges as n goes to ∞ .

Case $\sum_{n=0}^{\infty} r_n = \infty$. Then almost surely, there are infinitely many steps in which Max chooses action 1. If Min’s strategy π always chooses action 0, she guarantees that the game almost surely reaches the losing sink l . Thus, $\mathcal{P}_{\mathcal{G}, d, \sigma, \pi}(\text{Bu}) = 0$.

Case $\sum_{n=0}^{\infty} r_n < \infty$. Let $\varepsilon > 0$. Then we can choose K so that $\sum_{n=K}^{\infty} r_n \leq \varepsilon$. In particular, ε bounds the probability with which Max ever plays action 1 after step K . Let π be the Min strategy that chooses 0 until step K and 1 thereafter. This guarantees that until step K , the game will surely avoid the winning sink w . If in the first K steps, Max plays action 1 then the game moves to the losing sink l and is won by Min. Otherwise, if the game does not reach l within the first K steps, then the probability that Max always plays

0, and the play therefore alternates between non-target states d and t indefinitely, is at least $1 - \varepsilon$. Hence, $\mathcal{P}_{\mathcal{G},d,\sigma,\pi}(\text{Bu}) \leq \varepsilon$. \square

We now show that there exists an ε -optimal Max strategy in the Bad Match that uses just a step counter plus one extra bit of public memory.

PROPOSITION 4.4. *The value of state d in the Bad Match is 1 and Max has ε -optimal 1-bit Markov strategies.*

PROOF. Maitra and Sudderth [20, Lemma 13.8] prove that state d has value 1 by showing that no Min strategy can enforce a value strictly below 1 (i.e., the upper value of d is 1) and then appealing to the determinacy of the game.

We now give an alternative proof by explicitly constructing a Max strategy σ that is ε -optimal from d . Afterwards we argue how σ can be modified to give a 1-bit Markov strategy.

First, recall that finite-state concurrent games with *reachability* objectives always have memoryless randomized ε -optimal strategies [20, Section 7.7]. In particular, in the Bad Match as depicted in Figure 1, Max has such a strategy to reach the target set $\{s, w\}$ from d with probability $\geq 1 - \varepsilon$. He can do so by playing action 1 with probability ε and action 0 with probability $1 - \varepsilon$. Indeed, consider this Max strategy and an arbitrary Min strategy. Let E be the event that Min eventually plays action 0. Conditioned under E , there is a first time that Min plays action 0. Then either Max has already played action 1 before, in which case he has already won via state w , or not, in which case Max plays action 0 with probability $1 - \varepsilon$, so he wins via state s with probability $1 - \varepsilon$. Therefore, conditioned under E , Max wins with probability $\geq 1 - \varepsilon$. On the other hand, conditioned under not- E , Min always plays action 1 and thus Max wins almost surely by reaching state w . It follows that Max wins (reaches a state in $\{s, w\}$) with probability at least $1 - \varepsilon$.

Now consider the following Max strategy σ from d for the Büchi objective. It proceeds in phases 1, 2, 3, \dots , separated by visits to state s (unless state w has been reached). In every phase i , Max plays an ε_i -optimal strategy towards reaching the set $\{s, w\}$. If w is reached then the play is winning for Max anyway, otherwise the play returns from s to d and the next phase $i + 1$ begins. By choosing $\varepsilon_i \stackrel{\text{def}}{=} \varepsilon 2^{-i}$, we obtain that the Büchi objective is satisfied with probability $\geq 1 - \varepsilon$. This strategy σ uses infinite memory, since it needs to keep track of the current phase i (and this cannot be done with just a step counter by Proposition 4.3).

We construct a Max strategy σ' that is similar to σ . It also works in phases, but each phase i has a pre-determined length l_i . In each phase i , Max plays a memoryless ε_i -optimal strategy towards reaching the target set $\{s, w\}$, and the length l_i is chosen sufficiently large such that the probability of reaching the target before the end of the phase is $\geq 1 - \varepsilon_i - \varepsilon'_i$ for some sufficiently small $\varepsilon'_i > 0$. If w is reached then the play is winning for Max anyway. Otherwise, for the rest of the current phase, Max plays conservatively by always choosing action 0. Then the next phase $i + 1$ begins, etc. By choosing $\varepsilon_i \stackrel{\text{def}}{=} \varepsilon'_i \stackrel{\text{def}}{=} \varepsilon 2^{-(i+1)}$, we obtain that the Büchi objective is satisfied with probability $\geq 1 - \varepsilon$. The strategy σ' is a 1-bit Markov strategy. It uses the step counter to determine the current phase, since the lengths l_i are pre-determined, and it uses 1 bit of public memory to remember whether it has already visited state s in the current phase and thus needs to play conservatively until the end of this

phase. The 1 bit ensures that σ' loses $\leq \varepsilon_i + \varepsilon'_i = \varepsilon 2^{-i}$ only *once* in each phase i , and not multiple times per phase. \square

We show, as Theorem 3.1, that such an ε -optimal 1-bit Markov strategy for Max always exists, even in countably infinite-state stochastic Büchi games; see Section 6.

5 TRANSIENCE GAMES

In this section we prove that Max has memoryless strategies that are uniformly ε -optimal for Transience.

THEOREM 3.3. *Let \mathcal{G} be a game with Tr objective and $\varepsilon > 0$. Max has a memoryless strategy that is multiplicatively ε -optimal from every state.*

Moreover, if \mathcal{G} is turn-based then there exists such a strategy that is deterministic.

We now fix a game \mathcal{G} with the set S of states and with the transience objective Tr . From \mathcal{G} , we construct a *leaky* game \mathcal{G}_\perp , described below, where $\perp \notin S$ is a fresh losing sink state. Denote by $\text{Avoid}(\perp)$ the event of not visiting \perp . The goal of the construction is that, for every memoryless strategy of Max together with any (general) strategy of Min, the events Tr and $\text{Avoid}(\perp)$ are equal up to measure zero. The intuition is that it then suffices to construct strategies in \mathcal{G}_\perp that focus on avoiding \perp , which is conceptually easier.

Towards this goal, for every state $s \in S$ and every Max action $a \in A(s)$, we replace this Max action a by a countably infinite collection of Max actions a_η for every *rational* number $\eta \in (0, 1)$. The Max action a_η leads with probability η (regardless of Min's action) to \perp , and with probability $1 - \eta$ behaves like the original Max action a ; see Figure 2 for an illustration of the construction. Hence the set of Max actions in \mathcal{G}_\perp is countably infinite. Below, we show that for all events $E \subseteq S^\omega$ (note that $\perp \notin S$) and states s , the values of s for E in games \mathcal{G} and \mathcal{G}_\perp are equal. In particular, the equality holds for the transience objective Tr .

LEMMA 5.1. *For all events $E \subseteq S^\omega$ and all states $s \in S$:*

$$\text{val}_{\mathcal{G},E}(s) = \text{val}_{\mathcal{G}_\perp,E}(s).$$

The lemma follows by showing that, for every $\varepsilon > 0$, every Max strategy in \mathcal{G} can be converted to a strategy in \mathcal{G}_\perp with only an additional ε error. Indeed, consider any Max strategy σ in \mathcal{G} . We define σ_\perp to play at the history h exactly the same mixed action as σ plays, but with the leakage $\eta = \varepsilon \cdot 2^{-(|h|+1)}$. See Appendix C for a detailed proof.

As mentioned above, the leaky game \mathcal{G}_\perp has the following important property.

LEMMA 5.2. *In \mathcal{G}_\perp , for all memoryless Max strategies σ , all Min strategies π and states s , the events Tr and $\text{Avoid}(\perp)$ differ only by a nullset. Formally,*

$$\mathcal{P}_{\mathcal{G}_\perp,s,\sigma,\pi}((\text{Tr} \setminus \text{Avoid}(\perp)) \cup (\text{Avoid}(\perp) \setminus \text{Tr})) = 0.$$

The lemma follows by the strong fairness of probabilistic choices. Informally speaking, if not Tr then some state is visited infinitely often, but for every visit to such a state there will be a fixed positive probability leaking to \perp (depending only on the state), and thus a transition to \perp is eventually taken. See Appendix C for a detailed proof.




Figure 2: In the game \mathcal{G} , the distribution $p(s, a, b)$ is such that $\sum_{i \in \mathbb{N}} p(s, a, b)(s_i) = 1$. In the leaky game \mathcal{G}_\perp , for every rational number $\eta \in (0, 1)$, the probability distribution $p_\perp(s, a_\eta, b)$ is defined such that the support (s, a, b) also includes \perp . More precisely, we have $p_\perp(s, a_\eta, b)(s_i) = (1 - \eta)p(s, a, b)(s_i)$ and $p_\perp(s, a_\eta, b)(\perp) = \eta$.

5.1 Key Technical Lemmas

The following three lemmas, Lemma 5.3–Lemma 5.5, are the main technical tools, that in Section 5.2 combined with an argument on a martingale defined on the transience values in \mathcal{G}_\perp , prove Theorem 3.3.

By Lemma 5.1, all states have the same value in transience games \mathcal{G} and \mathcal{G}_\perp . By Lemma C.1 in Appendix C, we can carry back ε -optimal memoryless strategies in \mathcal{G}_\perp to \mathcal{G} . Subsequently, it suffices to prove that Max has memoryless ε -optimal strategies in the game \mathcal{G}_\perp . Below, since we mostly focus on the game \mathcal{G}_\perp we may drop it from the subscripts in values, probabilities and expectations, if understood from the context.

Fix some start state $s_0 \neq \perp$ in \mathcal{G}_\perp . Without restriction, we assume that \perp is the only state with value 0 in \mathcal{G}_\perp and thus $\text{val}_{\text{Tr}}(s_0) > 0$. Fix u such that

$$0 \leq u < \text{val}_{\text{Tr}}(s_0). \quad (1)$$

For all $n \geq 1$, let \mathcal{F}_n be the sigma-algebra generated by the cylinder sets corresponding to the histories $hs \in Z^* S$ where h starts in s_0 and visits s_0 at most $n - 1$ times; note that hs visits s_0 n th time if $s = s_0$ and h visits s_0 exactly $n - 1$ times.

Let

$$\tau : Z^\omega \rightarrow \{1, 2, \dots\} \cup \{\infty\} \quad (2)$$

denote the random variable that counts the number of visits to s_0 . Define a sequence $(X_n)_{n=1}^\infty$ of random variables by

$$X_n \stackrel{\text{def}}{=} \begin{cases} u & \text{if } \tau \geq n \\ 1 & \text{if } \tau < n \text{ and } \text{Tr} \\ 0 & \text{if } \tau < n \text{ and not } \text{Tr}. \end{cases} \quad (3)$$

Intuitively, the X_n encode increasingly strict conditions about the number of visits to s_0 . Since $\tau \geq n$ and $(\tau < n) \cap \text{Tr}$ are \mathcal{F}_n -measurable, the X_n are \mathcal{F}_n -measurable.

Denote by Ret the event to return to the state s_0 , that is, the set of all plays that start in s_0 and revisit it at some later step.

LEMMA 5.3. *Max has a strategy σ such that, for all $\pi \in \Pi$,*

$$\mathcal{P}_{s_0, \sigma, \pi}(\text{Tr} \setminus \text{Ret}) + \mathcal{P}_{s_0, \sigma, \pi}(\text{Ret})u > u.$$

Moreover, if \mathcal{G}_\perp is turn-based then there exists such a strategy that is deterministic.

PROOF. Towards a contradiction suppose that for all Max strategies $\tilde{\sigma}$ there is a Min strategy π with

$$\mathcal{P}_{s_0, \tilde{\sigma}, \pi}(\text{Tr} \setminus \text{Ret}) + \mathcal{P}_{s_0, \tilde{\sigma}, \pi}(\text{Ret})u \leq u. \quad (4)$$

Let σ be an arbitrary Max strategy. For any history $\mathbf{h} \in H$ that ends with a visit to s_0 , there is a unique strategy, denoted by $\sigma_{\mathbf{h}}$, whose behavior is exactly like the behavior of σ after \mathbf{h} ; formally, $\sigma_{\mathbf{h}}(h) = \sigma(\mathbf{h}h)$ for all histories $h \in H$.

By (4), for any history \mathbf{h} with $s_{\mathbf{h}} = s_0$ and the unique $\sigma_{\mathbf{h}}$ just defined, there is $\pi_{\mathbf{h}}$ with

$$\mathcal{P}_{s_0, \sigma_{\mathbf{h}}, \pi_{\mathbf{h}}}(\text{Tr} \setminus \text{Ret}) + \mathcal{P}_{s_0, \sigma_{\mathbf{h}}, \pi_{\mathbf{h}}}(\text{Ret})u \leq u. \quad (5)$$

Let π be the Min strategy that upon every visit to state s_0 via a history \mathbf{h} plays $\pi_{\mathbf{h}}$ (until, if it exists, the next visit of s_0). Formally, we define $\pi(\mathbf{h}h) = \pi_{\mathbf{h}}(h)$ if the history \mathbf{h} ends with a visit to s_0 and h contains s_0 only at the start.

Consider the probability measure induced by \mathcal{G}_\perp under σ and π ; and recall that X_n is \mathcal{F}_n -measurable. We show the following:

CLAIM 5.3.1. *The sequence $(X_n)_{n=1}^\infty$ is a supermartingale with respect to the filtration $(\mathcal{F}_n)_{n=1}^\infty$:*

$$\mathcal{E}_{s_0, \sigma, \pi}(X_{n+1} \mid \mathcal{F}_n) \leq X_n.$$

The proof of the claim rest on (5). The details are technical and can be found in Appendix C.

For all $n \geq 1$, the events $\tau + 1 = n$ and $\tau + 1 \leq n$ are \mathcal{F}_n -measurable. This implies that $\tau + 1$ is a stopping time with respect to the filtration $(\mathcal{F}_n)_{n=1}^\infty$, given the convention that $\infty + 1$ is equal ∞ . Since $|X_n| \leq 1$ for all n , using the optional stopping theorem (OST)³, we obtain the following

$$\begin{aligned} u &= \mathcal{E}_{s_0, \sigma, \pi}(X_1) \\ &\geq \mathcal{E}_{s_0, \sigma, \pi}(X_{\tau+1}) && \text{by OST and Claim 5.3.1} \\ &\geq \mathcal{E}_{s_0, \sigma, \pi}(X_{\tau+1} \mid \text{Tr}) \mathcal{P}_{s_0, \sigma, \pi}(\text{Tr}) \end{aligned}$$

Since on Tr the random variable $\tau + 1 < \infty$ and $X_{\tau+1} = 1$, we obtain $\mathcal{E}_{s_0, \sigma, \pi}(X_{\tau+1} \mid \text{Tr}) = 1$. By the latter and the argument above, we get

$$u \geq \mathcal{P}_{s_0, \sigma, \pi}(\text{Tr}).$$

Since the strategy σ was chosen arbitrarily, it follows that

$$u \geq \text{val}_{\text{Tr}}(s_0),$$

³Throughout this paper we use a general version of the optional stopping theorem (OST) from [22, Proposition IV-5-24, Corollary IV-2-25].

a contradiction with (1). The following claim concludes the proof of Lemma 5.3. A proof can be found in Appendix C.

CLAIM 5.3.2. *If \mathcal{G}_\perp is turn-based, there is a strategy σ , as described in Lemma 5.3, that is deterministic in the first step.*

□

Let σ be as described in Lemma 5.3. Denote by σ^* the strategy obtained from σ by restarting every time s_0 is revisited. Formally, we define

$$\sigma^*(h_1 h_2) = \sigma(h_2) \quad (6)$$

if the history h_1 ends with a visit to s_0 and h_2 has no visits to s_0 (except at the start).

The purpose of the next lemma is to establish the existence of a good strategy that always takes the *same* mixed action whenever the play returns to s_0 . Note that a strategy that is merely ε -optimal in every subgame is not sufficient here, and hence we need the stronger property of Lemma 5.3 to get the strategy σ^* above.

LEMMA 5.4. $\mathcal{P}_{s_0, \sigma^*, \pi}(\text{Tr}) \geq u$ for all $\pi \in \Pi$.

PROOF. Let π be an arbitrary Min strategy. Consider the probability measure induced by \mathcal{G}_\perp under σ^* and π .

CLAIM 5.4.1. *The sequence $(X_n)_{n=1}^\infty$ is a submartingale with respect to the filtration $(\mathcal{F}_n)_{n=1}^\infty$:*

$$\mathcal{E}_{s_0, \sigma^*, \pi}(X_{n+1} \mid \mathcal{F}_n) \geq X_n.$$

Similarly to the proof of Lemma 5.3, in order to use the optional stopping theorem, observe that $\tau + 1$ is a stopping time with respect to the filtration $(\mathcal{F}_n)_{n=1}^\infty$. This together with the optional stopping theorem imply that

$$\begin{aligned} u &= \mathcal{E}_{s_0, \sigma^*, \pi}(X_1) \\ &\leq \mathcal{E}_{s_0, \sigma^*, \pi}(X_{\tau+1}) && \text{by OST and Claim 5.4.1} \\ &\leq \mathcal{P}_{s_0, \sigma^*, \pi}(X_{\tau+1} \neq 0) && \text{since } 0 \leq X_{\tau+1} \leq 1. \end{aligned}$$

By definition of the X_n , see (3), if $\tau < \infty$ and not Tr then $X_{\tau+1} = 0$ holds. By the above, we note that

$$\begin{aligned} u &\leq \mathcal{P}_{s_0, \sigma^*, \pi}(X_{\tau+1} \neq 0) \\ &\leq \mathcal{P}_{s_0, \sigma^*, \pi}(\tau = \infty \text{ or Tr}) \\ &\leq \mathcal{P}_{s_0, \sigma^*, \pi}(\tau = \infty) + \mathcal{P}_{s_0, \sigma^*, \pi}(\text{Tr}). \end{aligned}$$

By definition of the leaky game \mathcal{G}_\perp and the strategy σ^* , there is $\delta > 0$ such that upon every visit of s_0 the probability of falling into \perp in the very next step is at least δ . As a result,

$$\mathcal{P}_{s_0, \sigma^*, \pi}(\tau = \infty) = 0$$

holds, implying that $u \leq \mathcal{P}_{s_0, \sigma^*, \pi}(\text{Tr})$, as required. □

Let α be a Max mixed action at s_0 . Define \mathcal{G}_α to be a game similar to \mathcal{G}_\perp where Max is only able to play α at s_0 . We formalize the definition of \mathcal{G}_α by introducing the notion of *fixing a mixed action α in \mathcal{G}* , that is, to obtain a new game identical to \mathcal{G}_\perp except that

- (1) the set of Max actions at s_0 is a singleton set $\{\alpha\}$; and
- (2) for all Min actions $b \in B(s_0)$, the transition function of \mathcal{G}_α for (s_0, α, b) is exactly the probability distribution defined by α and b in \mathcal{G}_\perp .

The following lemma shows that one can fix a mixed action at s_0 so that the value of no state drops significantly.

LEMMA 5.5. *Let $s_0 \in S$ be a state in \mathcal{G}_\perp (hence $s_0 \neq \perp$) and let $r \in (0, 1)$. There is a Max mixed action α at s_0 such that*

$$\text{val}_{\mathcal{G}_\alpha, \text{Tr}}(s) \geq r \text{val}_{\mathcal{G}_\perp, \text{Tr}}(s)$$

holds for all states $s \in S$. If \mathcal{G}_\perp is turn-based, there is such an α that is Dirac.

PROOF. Let $s \in S$ be chosen arbitrary. Let

$$u \stackrel{\text{def}}{=} r \text{val}_{\mathcal{G}_\perp, \text{Tr}}(s_0) \quad (7)$$

ensuring that $u < \text{val}_{\mathcal{G}_\perp, \text{Tr}}(s_0)$ as required in (1). Let σ be a strategy satisfying the condition in Lemma 5.3 for this u . From σ , we obtain σ^* as described in (6). Let the mixed action α , a probability distribution over $A(s_0)$, be the action of σ^* at history s_0 . If \mathcal{G}_\perp is turn-based, α is Dirac.

Let $x \in (0, 1)$ be chosen arbitrary. We finally define $\tilde{\sigma}$ from σ^* : Starting in s , this strategy $\tilde{\sigma}$ first plays a strategy σ_s with

$$\inf_{\pi} \mathcal{P}_{\mathcal{G}_\perp, s, \sigma_s, \pi}(\text{Tr}) \geq x \text{val}_{\mathcal{G}_\perp, \text{Tr}}(s), \quad (8)$$

and whenever (if at all) the play visits s_0 , it switches to σ^* .

Let $\tilde{\pi}$ be an arbitrary Min strategy. We redefine the random variable τ as given in (2), allowing for the possibility that $\tau = 0$ to accommodate plays that never visit s_0 . For the u in (7), define the sequence $(X_n)_{n=1}^\infty$ of random variables as in (3). Define

$$X_0 \stackrel{\text{def}}{=} r x \text{val}_{\mathcal{G}_\perp, \text{Tr}}(s), \quad (9)$$

and let \mathcal{F}_0 be the sigma-algebra generated by the cylinder set corresponding to the history that consists only of the initial state s . Similar to the X_n being \mathcal{F}_n -measurable, the random variable X_0 is also \mathcal{F}_0 -measurable.

CLAIM 5.5.1. *The sequence $(X_n)_{n=0}^\infty$ is a submartingale with respect to the filtration $(\mathcal{F}_n)_{n=0}^\infty$:*

$$\mathcal{E}_{\mathcal{G}_\perp, s_0, \tilde{\sigma}, \tilde{\pi}}(X_{n+1} \mid \mathcal{F}_n) \geq X_n.$$

The proof of this Claim is a combination of reasoning for $\mathcal{E}_{\mathcal{G}_\perp, s, \tilde{\sigma}, \tilde{\pi}}(X_1 \mid \mathcal{F}_0) \geq X_0$ and Claim 5.4.1; we provide a detailed proof in Appendix C.

Proceeding similarly as in the proof of Lemma 5.4, we can get that

$$\begin{aligned} r x \text{val}_{\mathcal{G}_\perp, \text{Tr}}(s) &= \mathcal{E}_{\mathcal{G}_\perp, s, \tilde{\sigma}, \tilde{\pi}}(X_0) && \text{by definition in (9)} \\ &\leq \mathcal{E}_{\mathcal{G}_\perp, s, \tilde{\sigma}, \tilde{\pi}}(X_{\tau+1}) && \text{by OST and Claim 5.5.1} \\ &\leq \mathcal{P}_{\mathcal{G}_\perp, s, \tilde{\sigma}, \tilde{\pi}}(\text{Tr}) && \text{similarly to Lemma 5.4.} \end{aligned}$$

By its construction, $\tilde{\sigma}$ chooses α each time the play visits s_0 . Recall that \mathcal{G}_α is the game obtained from \mathcal{G}_\perp by fixing α at s_0 , ensuring that Max is only able to play α at s_0 . Hence, the strategy $\tilde{\sigma}$ is uniquely translated to a corresponding strategy $\tilde{\sigma}_\alpha$ in \mathcal{G}_α in a natural way. Following $\tilde{\sigma}_\alpha$ in \mathcal{G}_α , we obtain

$$r x \text{val}_{\mathcal{G}_\perp, \text{Tr}}(s) \leq \mathcal{P}_{\mathcal{G}_\alpha, s, \tilde{\sigma}_\alpha, \tilde{\pi}}(\text{Tr}).$$

Since x and $\tilde{\pi}$ were arbitrary, it follows that

$$r \text{val}_{\mathcal{G}_\perp, \text{Tr}}(s) \leq \text{val}_{\mathcal{G}_\alpha, \text{Tr}}(s),$$

as required. □

5.2 Proof of Theorem 3.3

We are now in a position to prove Theorem 3.3 which states that Max has memoryless strategies that are ε -optimal from every state in a transience game.

In the sequel, for clarity, we assume that the set of states in \mathcal{G}_\perp , excluding \perp , is $\{1, 2, \dots\}$. Let $r \in (0, 1)$ be arbitrary. Define $\mathcal{G}_0 \stackrel{\text{def}}{=} \mathcal{G}_\perp$. We inductively define a sequence $(\mathcal{G}_i)_{i \geq 1}$ of games and a sequence $(\alpha_i)_{i \geq 1}$ of Max mixed actions. For all $i \geq 1$, invoking Lemma 5.5, there exists a probability distribution α_i on $A(i)$ such that for all $s \in S$

$$\text{val}_{\mathcal{G}_i, \text{Tr}}(s) \geq r^{2^{-i}} \text{val}_{\mathcal{G}_{i-1}, \text{Tr}}(s) \quad (10)$$

where the game \mathcal{G}_i is obtained from \mathcal{G}_{i-1} by fixing the mixed actions α_i at i .

The collection of the mixed actions α_i at states i defines a memoryless strategy, denoted by σ , for Max in \mathcal{G}_\perp . By Lemma 5.5, if \mathcal{G}_\perp is turn-based, σ can be made deterministic. Below, we will prove that, for all states $s \in S$ and for all Min strategies π ,

$$\mathcal{P}_{\mathcal{G}_\perp, s, \sigma, \pi}(\text{Tr}) \geq r \text{val}_{\mathcal{G}_\perp, \text{Tr}}(s), \quad (11)$$

proving Theorem 3.3.

For all $i \geq 0$, define $r_i \stackrel{\text{def}}{=} \prod_{j>i} r^{2^{-j}}$, that is intuitively the proportion of the value in the game \mathcal{G}_i that will be maintained by all *future fixings* $j > i$. For brevity, write $v_i(s)$ for $\text{val}_{\mathcal{G}_i, \text{Tr}}(s)$. It follows from the definition of the games \mathcal{G}_i in (10) that for all $i \geq 1$ and all states s ,

$$r_i v_i(s) \geq r_{i-1} v_{i-1}(s). \quad (12)$$

Recall that H_n is the set of histories at step n . These histories visit $n+1$ states (though not necessarily $n+1$ different states). For all $n \geq 1$, let \mathcal{F}_n be the sigma-algebra generated by the cylinder sets corresponding to H_{n-1} .

Let π be an arbitrary Min strategy. Given a play visiting the sequence s_1, s_2, \dots of states, for all $n \geq 1$, we define the random variable

$$m(n) \stackrel{\text{def}}{=} \max\{s_1, s_2, \dots, s_n\}.$$

We further define the random variable Y_n , taking values in $[0, 1]$, by

$$Y_n \stackrel{\text{def}}{=} r_{m(n)} v_{m(n)}(s_n). \quad (13)$$

Intuitively, the variable Y_n is the value of s_n after having *fixed* an action in the states $\{1, \dots, m(n)\}$, scaled down by $r_{m(n)}$, that is the proportion of the value in the game \mathcal{G}_i that will be maintained by all *future fixings* $i > m(n)$. Both random variables $m(n)$ and Y_n are \mathcal{F}_n -measurable.

Let s_1 be an arbitrary state, excluding \perp . We aim to prove the inequality in (11). Consider a random play visiting the sequence s_1, s_2, \dots of states. For all $n \geq 1$, we have

$$\begin{aligned} & \mathcal{E}_{s_1, \sigma, \pi}(Y_{n+1} \mid \mathcal{F}_n) \\ &= \mathcal{E}_{s_1, \sigma, \pi}(r_{m(n+1)} v_{m(n+1)}(s_{n+1}) \mid \mathcal{F}_n) \quad \text{by (13)} \\ &\geq \mathcal{E}_{s_1, \sigma, \pi}(r_{m(n)} v_{m(n)}(s_{n+1}) \mid \mathcal{F}_n) \quad \text{by (12)} \end{aligned}$$

But then recall that in state s_n , strategy σ plays the mixed action fixed in $\mathcal{G}_{m(n)}$, and, therefore, the expected value of the values in $\mathcal{G}_{m(n)}$ does not drop in one step. Hence, we get

$$\begin{aligned} & \geq \mathcal{E}_{s_1, \sigma, \pi}(r_{m(n)} v_{m(n)}(s_n) \mid \mathcal{F}_n) \\ &= Y_n. \end{aligned}$$

Consequently, the sequence $(Y_n)_{n=1}^\infty$ is a submartingale with respect to the filtration $(\mathcal{F}_n)_{n=1}^\infty$. This implies that, in game \mathcal{G}_\perp , for all $n \geq 1$,

$$\mathcal{E}_{s_1, \sigma, \pi}(Y_n) \geq \mathcal{E}_{s_1, \sigma, \pi}(Y_1). \quad (14)$$

We can now conclude the proof as follows. In game \mathcal{G}_\perp , by Lemma 5.2 and σ being memoryless, we get that

$$\mathcal{P}_{s_1, \sigma, \pi}(\text{Tr}) = \mathcal{P}_{s_1, \sigma, \pi}(\text{Avoid}(\perp))$$

Since \perp is a sink state and has value 0 for transience, we have

$$\geq \mathcal{P}_{s_1, \sigma, \pi}\left(\limsup_{n \rightarrow \infty} Y_n > 0\right)$$

Furthermore, since $Y_n \leq 1$, it follows that

$$\begin{aligned} & \geq \mathcal{E}_{s_1, \sigma, \pi}\left(\limsup_{n \rightarrow \infty} Y_n\right) \\ & \geq \limsup_{n \rightarrow \infty} \mathcal{E}_{s_1, \sigma, \pi}(Y_n) \quad \text{reverse Fatou lemma} \\ & \geq \mathcal{E}_{s_1, \sigma, \pi}(Y_1) \quad \text{by (14)} \\ & = r_1 v_1(s_1) \quad \text{by (13)} \\ & \geq r_0 v_0(s_1) \quad \text{by (12)} \end{aligned}$$

By (10) and definition of r_0 , we finally can conclude that

$$\mathcal{P}_{s_1, \sigma, \pi}(\text{Tr}) \geq r \text{val}_{\mathcal{G}_\perp, \text{Tr}}(s_1),$$

as required in (11).

6 BÜCHI GAMES

In a game \mathcal{G} with state space S and $T \subseteq S$ denote by Bu the Büchi objective, i.e., the objective to visit the set T infinitely often. (If the set T is infinite then some plays that satisfy Bu might not visit any particular state in T infinitely often.) Define also the transient Büchi objective, $\text{TB} \stackrel{\text{def}}{=} \text{Tr} \cap \text{Bu}$. Clearly, TB can only be met if T is infinite.

We show the following main result, which, as argued in Section 3, implies Theorem 3.1.

THEOREM 3.2. *Let \mathcal{G} be a game with TB objective and $\varepsilon > 0$. Max has a 1-bit strategy σ so that*

- (1) $\sigma[0]$ is multiplicatively ε -optimal from every state, and
- (2) all memory updates $\sigma_{\text{up}}(\cdot)$ are Dirac (hence the memory is public).

Moreover, if \mathcal{G} is turn-based then there exists such a strategy that is deterministic.

Fix a game \mathcal{G} , and consider the derived leaky game \mathcal{G}_\perp as defined in Section 5. By Lemma 5.1, all states have the same value in \mathcal{G} and \mathcal{G}_\perp wrt. the TB objective. By Lemma C.1 in Appendix C, we can carry back ε -optimal 1-bit strategies in \mathcal{G}_\perp to \mathcal{G} . Thus, it suffices to prove that Max has ε -optimal 1-bit strategies in the game \mathcal{G}_\perp .

We will assume without loss of generality that $\perp \notin S$ is the only state with $\text{val}_{\text{TB}}(\perp) = 0$, as we can redirect all transitions that lead to a value-0 state to \perp , without changing the value of any state. Moreover, by Lemma C.2 in Appendix C, in \mathcal{G}_\perp , under every finite-memory Max strategy and arbitrary Min strategy, the events Tr and $\text{Avoid}(\perp)$ are equal up to measure zero.

Definition 6.1. A bridge in \mathcal{G}_\perp is a triple $\Lambda = (I, L, \sigma_0)$ where $I, L \subseteq S$ (thus $\perp \notin I$) are disjoint finite sets of states and σ_0 is a mapping that maps every $s \in I \cup L$ to a mixed Max action $\sigma_0(s) \in \mathcal{D}(A(s))$. We write $\langle \Lambda \rangle$ for the event that after each visit of a state in I , eventually $I \cup L$ will be left, at which point immediately a state in T is visited.

A Max strategy σ is said to be *consistent* with Λ if, in any play against any Min strategy π , after every visit of a state in I , strategy σ plays as specified by σ_0 until $I \cup L$ is left. A bridge is called *deterministic* if, for every $s \in I \cup L$, $\sigma_0(s)$ is Dirac, i.e., it chooses a pure action.

To prove Theorem 3.2, we need the following key lemma, which is shown in Appendix D.1. Its role is similar to the role of Lemma 5.5 in Section 5.

LEMMA 6.2. *Let $I \subseteq S$ be a finite set of states in \mathcal{G}_\perp (thus $\perp \notin I$). Let $r \in (0, 1)$. There are a bridge $\Lambda = (I, L, \sigma_0)$ and a Max strategy σ consistent with Λ so that*

$$\inf_{\pi} \mathcal{P}_{s, \sigma, \pi}(\langle \Lambda \rangle \cap \text{TB}) \geq r \text{val}_{\text{TB}}(s)$$

holds for all states $s \in S$. If \mathcal{G}_\perp is turn-based then there is a deterministic such bridge.

Now we prove the main theorem.

PROOF OF THEOREM 3.2. Let $r \in (0, 1)$ be arbitrary. It is convenient to assume that $S = \{2, 4, 6, \dots\}$. We define a sequence $(\mathcal{G}_i)_{i \geq 1}$ of games inductively. Each game \mathcal{G}_i will have a state space

$$S_i \stackrel{\text{def}}{=} S \cup L'_1 \cup \dots \cup L'_i,$$

where S, L'_1, L'_2, \dots are pairwise disjoint and $L'_i \subseteq \{3, 5, 7, \dots\}$. We will later associate the even and odd state numbers in these derived games with different memory modes 0 and 1 of a 1-bit strategy in the original game \mathcal{G}_\perp , respectively.

Define $\mathcal{G}_0 \stackrel{\text{def}}{=} \mathcal{G}_\perp$ and

$$I_0 \stackrel{\text{def}}{=} L_0 \stackrel{\text{def}}{=} L'_0 \stackrel{\text{def}}{=} \emptyset.$$

For all $i \geq 1$, define

$$I_i \stackrel{\text{def}}{=} I_{i-1} \cup L_{i-1} \cup L'_{i-1} \cup \{2i\}.$$

By Lemma 6.2, in \mathcal{G}_{i-1} there are a bridge $\Lambda_i = (I_i, L_i, \sigma_i)$ and a Max strategy σ consistent with Λ_i so that for every $s \in S_{i-1}$ we have

$$\inf_{\pi} \mathcal{P}_{\mathcal{G}_{i-1}, s, \sigma, \pi}(\langle \Lambda_i \rangle \cap \text{TB}) \geq r^{2^{-i}} \text{val}_{\mathcal{G}_{i-1}, \text{TB}}(s).$$

If the game is turn-based then the bridge Λ_i can be made deterministic. Since $I_i \cap L_i = \emptyset$ and

$$I_i \supseteq L'_1 \cup \dots \cup L'_{i-1},$$

all states in L_i are in S , i.e., are “even”. See the top of Figure 3.

We construct \mathcal{G}_i from \mathcal{G}_{i-1} by adding fresh, “odd”, copies of the states in L_i , namely

$$L'_i \stackrel{\text{def}}{=} \{2j + 1 \mid 2j \in L_i\}.$$

In \mathcal{G}_i we “plaster” the bridge Λ_i , but using L'_i instead of L_i . (In MDPs/games, “plastering” means fixing the choices of a player in a subset of the states in a certain defined way; here according to the bridge. See [25] for an early use of this technique.) This is to enforce that upon every entry into I_i , a version of σ_i is played by

Max until $I_i \cup L'_i$ is left, at which point either a state in T is visited or the play falls into \perp . More precisely, we replace the Max actions of states $s \in I_i$ by the single action $\sigma_i(s)$, and states $2j + 1 \in L'_i$ get the single action $\sigma_i(2j)$. The successor state distributions of those actions are modified so that transitions to states $2j \in L_i$ are redirected to the “odd sibling” $2j + 1 \in L'_i$, transitions to $I_i \cup (T \setminus L_i)$ are kept, and all other transitions are redirected to \perp . The states in L_i remain untouched for now (but will be plastered in \mathcal{G}_{i+1}). See the bottom of Figure 3.

From the properties of the bridge Λ_i , for all $s \in S_{i-1}$, we have

$$\begin{aligned} \text{val}_{\mathcal{G}_i, \text{TB}}(s) &\geq \inf_{\pi} \mathcal{P}_{\mathcal{G}_{i-1}, s, \sigma, \pi}(\langle \Lambda_i \rangle \cap \text{TB}) \\ &\geq r^{2^{-i}} \text{val}_{\mathcal{G}_{i-1}, \text{TB}}(s). \end{aligned} \quad (15)$$

There is a natural limit of the games \mathcal{G}_i , say \mathcal{G}_∞ , with state space

$$S_\infty \stackrel{\text{def}}{=} S \cup \bigcup_{i \geq 1} L'_i,$$

where all choices of Max have been fixed. (One does not need to fix actions in \perp , since its value is zero anyway.)

The game \mathcal{G}_∞ can be seen as an MDP where only Min is still active. From the point of view of Max, the fixed choices of Max in \mathcal{G}_∞ define a two-mode (public memory with deterministic updates) Max strategy, say σ , for \mathcal{G}_\perp , where

- (1) each (even) state $s \in S$ in \mathcal{G}_∞ corresponds to state s in \mathcal{G}_\perp and σ being in memory mode 0,
- (2) and each (odd) state $s \in \bigcup_{i \geq 1} L'_i$ in \mathcal{G}_∞ corresponds to state $s - 1$ in \mathcal{G}_\perp and σ being in memory mode 1.
- (3) The start mode is 0.
- (4) The mode update corresponding to transitions that were redirected to \perp in the construction of \mathcal{G}_∞ can be defined arbitrarily, say to mode 0.

Note that, while the actions of σ can be mixed in general, its memory updates are always deterministic, switching the memory mode after visiting I_i and T , respectively. However, if the game is turn-based then all the bridges can be made deterministic, and thus σ is completely deterministic.

Recall that $\sigma[0]$ refers to σ starting in memory mode 0. It remains to show that $\sigma[0]$ is nearly optimal from every state, that is,

$$\inf_{\pi} \mathcal{P}_{\mathcal{G}_\perp, s, \sigma[0], \pi}(\text{TB}) \geq r \text{val}_{\mathcal{G}_\perp}(s)$$

holds for all $s \in S$.

For all $s \in S$, the following holds:

$$\inf_{\pi} \mathcal{P}_{\mathcal{G}_\perp, s, \sigma[0], \pi}(\text{TB}) \geq \inf_{\pi} \mathcal{P}_{\mathcal{G}_\infty, s, \pi}(\text{TB}).$$

We note in passing that the inequality above might be strict due to the transitions redirected to \perp in \mathcal{G}_∞ . The game \mathcal{G}_∞ has been obtained from \mathcal{G}_\perp by fixing a finite-memory (actually 1-bit) Max strategy. Thus, by our construction of \mathcal{G}_\perp , the events Tr and $\text{Avoid}(\perp)$ are equal up to measure zero in \mathcal{G}_∞ , under any Min strategy (by Lemma C.2). In \mathcal{G}_∞ , for any state $i \in S_\infty$, the only way out of the finite set $I_i \cup L'_i$ is via a state in T or falling into \perp . Thus, in \mathcal{G}_∞ the events Bu and Tr are also equal up to measure zero under any Min




Figure 3: Illustration of the construction in Theorem 3.2. The top picture indicates a bridge $\Lambda_i = (I_i, L_i, \sigma_i)$ in game \mathcal{G}_{i-1} . The bottom picture indicates how in game \mathcal{G}_i the bridge Λ_i is “plastered” using L'_i instead of L_i .

strategy. Hence, for all $s \in S$ we have

$$\begin{aligned}
 \inf_{\pi} \mathcal{P}_{\mathcal{G}_{\perp}, s, \sigma[0], \pi}(\text{TB}) &\geq \inf_{\pi} \mathcal{P}_{\mathcal{G}_{\infty}, s, \pi}(\text{TB}) \\
 &= \inf_{\pi} \mathcal{P}_{\mathcal{G}_{\infty}, s, \pi}(\text{Tr}) \\
 &= \inf_{\pi} \mathcal{P}_{\mathcal{G}_{\infty}, s, \pi}(\text{Avoid}(\perp)).
 \end{aligned} \tag{16}$$

The rest of the proof proceeds similarly to Section 5. Let us write $v_i(s) \stackrel{\text{def}}{=} \text{val}_{\mathcal{G}_i, \text{TB}}(s)$ in the following to avoid clutter. For $i \geq 0$ define $r_i \stackrel{\text{def}}{=} \prod_{j>i} r^{2^{-j}}$. It follows from (15) that for all $i \geq 1$ and all

$s \in S_{i-1}$ we have

$$r_i v_i(s) \geq r_{i-1} v_{i-1}(s). \tag{17}$$

Recall that H_{n-1} is the set of histories with the first n states. For all $n \geq 1$, let \mathcal{F}_n be the sigma-algebra generated by the cylinder sets corresponding to H_{n-1} . For a play visiting states s_1, s_2, \dots and $n \geq 1$ define

$$m(n) \stackrel{\text{def}}{=} \max\{s_1, \dots, s_n\}$$

and the random variable Y_n , taking values in $[0, 1]$, by

$$Y_n \stackrel{\text{def}}{=} r_{m(n)} v_{m(n)}(s_n).$$

Both $m(n)$ and Y_n are \mathcal{F}_n -measurable. Intuitively, Y_n is obtained by scaling down $v_{m(n)} = \text{val}_{\mathcal{G}_{m(n)}}(s_n)$, which is the value in $\mathcal{G}_{m(n)}$ where (at least) the states s_1, \dots, s_n have been plastered, by a lower bound, $r_{m(n)}$, on the proportion of the value that will be maintained by “future” plasterings.

Let $s_1 \in S$ be an arbitrary state. It suffices to show that

$$\inf_{\pi} \mathcal{P}_{\mathcal{G}_{\perp, s_1, \sigma[0], \pi}}(\text{TB}) \geq \text{rval}_{\mathcal{G}_{\perp}}(s_1).$$

The following claim follows from the fact that $(Y_n)_{n=1}^{\infty}$ is a submartingale with respect to the filtration $(\mathcal{F}_n)_{n=1}^{\infty}$; see Appendix D.

CLAIM 6.2.1. *For any Min strategy π ,*

$$\mathcal{E}_{\mathcal{G}_{\infty, s_1, \pi}} Y_n \geq \mathcal{E}_{\mathcal{G}_{\infty, s_1, \pi}} Y_1 \quad \text{for all } n \geq 1.$$

Hence,

$$\begin{aligned} & \inf_{\pi} \mathcal{P}_{\mathcal{G}_{\perp, s_1, \sigma[0], \pi}}(\text{TB}) \\ & \geq \inf_{\pi} \mathcal{P}_{\mathcal{G}_{\infty, s_1, \pi}}(\text{Avoid}(\perp)) \quad \text{by (16)} \end{aligned}$$

Since \perp is a sink state and has value 0, we have

$$\geq \inf_{\pi} \mathcal{P}_{\mathcal{G}_{\infty, s_1, \pi}} \left(\limsup_{n \rightarrow \infty} Y_n > 0 \right)$$

Furthermore, since $Y_n \leq 1$, it follows that

$$\begin{aligned} & \geq \inf_{\pi} \mathcal{E}_{\mathcal{G}_{\infty, s_1, \pi}} \left(\limsup_{n \rightarrow \infty} Y_n \right) \\ & \geq \inf_{\pi} \limsup_{n \rightarrow \infty} \mathcal{E}_{\mathcal{G}_{\infty, s_1, \pi}} Y_n \quad \text{reverse Fatou lemma} \\ & \geq \inf_{\pi} \mathcal{E}_{\mathcal{G}_{\infty, s_1, \pi}} Y_1 \quad \text{by Claim 6.2.1} \\ & = r_{s_1} v_{s_1}(s_1) \quad \text{definitions of } Y_1, m(1) \\ & \geq r_0 v_0(s_1) \quad \text{by (17)} \\ & = \text{rval}_{\mathcal{G}_{\perp}}(s_1) \quad \text{definitions of } r_0, \mathcal{G}_{\perp}, \end{aligned}$$

as required. \square

7 STRATEGY COMPLEXITY FOR MINIMIZER

We have shown that ε -optimal Max strategies in countable concurrent Büchi games with finite Min action sets require just a step counter plus 1 bit of public memory (but cannot use less memory in general, even if the game is finite). This upper bound does not hold for the strategy complexity of the opposing Min player. From player Min’s point of view, the objective is to maximize the probability of satisfying the dual *co-Büchi* objective of visiting the set of target states only *finitely* often. I.e., Min in a Büchi game is the same as Max in a *co-Büchi* game. However, a step counter plus arbitrary finite (private) memory are *not* sufficient for ε -optimal Max strategies for the *co-Büchi* objective in countable concurrent games. In fact, this lower bounds holds even in countable finitely-branching turn-based *co-Büchi* games [17]. On the other hand, in *finite-state* concurrent games, ε -optimal Max strategies for the *co-Büchi* objective can be memoryless randomized [8], and this holds even for the more general \liminf objective [27]. *Optimal* Max strategies for the *co-Büchi* objective need not exist, but might require memory if they do. Bordais et al.[5, Section 6] give an example of a finite-state *co-Büchi* game where optimal Max strategies exist but require infinite memory.

In countably infinite *co-Büchi* games with an infinite set of target states T there can exist plays that visit T infinitely often, but visit each state in T only finitely often. By the standard definition, such plays do not satisfy the *co-Büchi* objective. One could define a weaker objective, *weak-co-Büchi*, that only requires that each state in T is visited only finitely often. *Weak-co-Büchi* coincides with *co-Büchi* if T is finite, and thus in particular in all finite-state games. However, in infinite-state games, *weak-co-Büchi* could be very different. In fact, it can be seen as a generalized transience objective, where one demands transience only wrt. the states in the set T instead of all states. We conjecture that ε -optimal Max strategies for this generalized transience objective could be memoryless, along similar lines as our proofs in Section 5.

REFERENCES

- [1] E. Altman, K. Avrachenkov, N. Bonneau, Merouane M. Debbah, R. El-Azouzi, and D. Menasche. 2007. Constrained stochastic games in wireless networks. In *IEEE global communications conference (GLOBECOM)*, IEEE (Ed.), Washington.
- [2] E. Altman, K. Avrachenkov, R. Marquez, and G. B. Miller. 2005. Zero-sum constrained stochastic games with independent state processes. *Math. Methods Oper. Res.* 62, 3 (2005), 375–386. <https://doi.org/10.1007/s00186-005-0034-4>
- [3] Michael Bacharach. 2019. *Economics and the Theory of Games*. CRC Press.
- [4] Patrick Billingsley. 1995. *Probability and Measure*. Wiley, New York, NY. Third edition.
- [5] Benjamin Bordais, Patricia Bouyer, and Stéphane Le Roux. 2022. Playing (Almost-)Optimally in Concurrent Büchi and Co-Büchi Games. In *42nd IARCS Annual Conference on Foundations of Software Technology and Theoretical Computer Science (FSTTCS 2022) (Leibniz International Proceedings in Informatics (LIPIcs), Vol. 250)*, Anuj Dawar and Venkatesan Guruswami (Eds.). Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl, Germany, 33:1–33:18. <https://doi.org/10.4230/LIPIcs.FSTTCS.2022.33>
- [6] Benjamin Bordais, Patricia Bouyer, and Stéphane Le Roux. 2022. Optimal Strategies in Concurrent Reachability Games. In *30th EACSL Annual Conference on Computer Science Logic, CSL 2022, February 14-19, 2022, Göttingen, Germany (Virtual Conference) (LIPIcs, Vol. 216)*, Florin Manea and Alex Simpson (Eds.). Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 7:1–7:17. <https://doi.org/10.4230/LIPIcs.CSL.2022.7>
- [7] P. Bouyer, N. Markey, M. Randour, A. Sangnier, and D. Stan. 2016. Reachability in Networks of Register Protocols under Stochastic Schedulers. In *International Colloquium on Automata, Languages and Programming (ICALP) (Leibniz International Proceedings in Informatics (LIPIcs), Vol. 55)*. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, Dagstuhl, Germany, 106:1–106:14.
- [8] Luca de Alfaro and Thomas Henzinger. 2000. Concurrent Omega-Regular Games. In *Proceedings of the 15th Annual IEEE Symposium on Logic in Computer Science (LICS 2000)*. IEEE.
- [9] L. de Alfaro and T. A. Henzinger. 2001. Interface automata. In *International Symposium on Foundations of Software Engineering*. ACM, 109–120. <https://doi.org/10.1145/503209>
- [10] L. de Alfaro, T. A. Henzinger, and O. Kupferman. 1998. Concurrent Reachability Games. In *39th Annual Symposium on Foundations of Computer Science, FOCS '98, November 8-11, 1998, Palo Alto, California, USA*. IEEE Computer Society, 564–575. <https://doi.org/10.1109/SFCS.1998.743507>
- [11] J Flesch, A. Predtetchinski, and W. Sudderth. 2020. Positive zero-sum stochastic games with countable state and action spaces. *Applied Mathematics and Optimization* 82 (2020), 499–516. Issue 2.
- [12] D. Gillette. 1958. Stochastic Games with Zero Stop Probabilities. In *Contributions to the Theory of Games (AM-39), Volume III*. Princeton University Press, 179–188. <https://doi.org/10.1515/9781400882151-011>
- [13] A. Jaskiewicz and A. S. Nowak. 2011. Stochastic Games with Unbounded Payoffs: Applications to Robust Control in Economics. *Dynamic Games and Applications* 1, 2 (2011), 253–279. <https://doi.org/10.1007/s13235-011-0013-8>
- [14] Stefan Kiefer, Richard Mayr, Mahsa Shirmohammadi, and Patrick Totzke. 2019. Büchi Objectives in Countable MDPs. In *International Colloquium on Automata, Languages and Programming (ICALP) (LIPIcs, Vol. 132)*. 119:1–119:14. <https://doi.org/10.4230/LIPIcs.ICALP.2019.119> Full version at <https://arxiv.org/abs/1904.11573>.
- [15] Stefan Kiefer, Richard Mayr, Mahsa Shirmohammadi, and Patrick Totzke. 2020. Strategy Complexity of Parity Objectives in Countable MDPs. In *International Conference on Concurrency Theory (CONCUR)*. 7:1–17. <https://doi.org/10.4230/LIPIcs.CONCUR.2020.39>

- [16] Stefan Kiefer, Richard Mayr, Mahsa Shirmohammadi, and Patrick Totzke. 2021. Transience in Countable MDPs. In *International Conference on Concurrency Theory (CONCUR) (LIPIcs, Vol. 203)*. Full version at <https://arxiv.org/abs/2012.13739>.
- [17] Stefan Kiefer, Richard Mayr, Mahsa Shirmohammadi, and Patrick Totzke. 2024. Strategy Complexity of Reachability in Countable Stochastic 2-Player Games. *Dynamic Games and Applications* (2024). To appear, <https://arxiv.org/abs/2203.12024>.
- [18] P. R. Kumar and T. H. Shiao. 1981. Existence of Value and Randomized Strategies in Zero-Sum Discrete-Time Stochastic Dynamic Games. *SIAM Journal on Control and Optimization* 19, 5 (1981), 617–634. <https://doi.org/10.1137/0319039>
- [19] A. Maitra and W. Sudderth. 1998. Finitely Additive Stochastic Games with Borel Measurable Payoffs. *International Journal of Game Theory* 27, 2 (July 1998), 257–267. <https://doi.org/10.1007/s001820050071>
- [20] Ashok P. Maitra and William D. Sudderth. 1996. *Discrete Gambling and Stochastic Games*. Springer-Verlag.
- [21] Donald A. Martin. 1998. The determinacy of Blackwell games. *Journal of Symbolic Logic* 63, 4 (1998), 1565–1581. <https://doi.org/10.2307/2586667>
- [22] J. Neveu. 1975. *Discrete-Parameter Martingales*. North-Holland Publishing Company.
- [23] Abraham Neyman and Sylvain Sorin. 2003. *Stochastic Games and Applications*. Springer, Dordrecht. <https://doi.org/10.1007/978-94-010-0189-2>
- [24] A. Nowak and Krzysztof Szajowski. 2005. *Advances in Dynamic Games: Applications to Economics, Finance, Optimization, and Stochastic Control*. Birkhaeuser, Boston. <https://doi.org/10.1007/b138903>
- [25] Donald Ornstein. 1969. On the Existence of Stationary Optimal Strategies. *Proc. Amer. Math. Soc.* 20, 2 (1969), 563–569. <https://doi.org/10.2307/2035700>
- [26] T. E. S. Raghavan, T.S. Ferguson, T. Parthasarathy, and O. Vrieze. 2012. *Stochastic Games and Related Topics: in honor of Professor LS Shapley*. Vol. 7. Springer Science & Business Media. <https://doi.org/10.1007/978-94-011-3760-7>
- [27] Piercesare Secchi. 1998. On the existence of good stationary strategies for nonleavable stochastic games. *International Journal of Game Theory* 27 (1998), 61–81. Issue 1. <https://doi.org/10.1007/BF01243195>
- [28] Lloyd S Shapley. 1953. Stochastic games. *Proceedings of the National Academy of Sciences* 39, 10 (1953), 1095–1100. <https://doi.org/10.1073/pnas.39.10.1095>
- [29] Eilon Solan and Nicolas Vieille. 2015. Stochastic games. *Proceedings of the National Academy of Sciences* 112, 45 (2015), 13743–13746. <https://doi.org/10.1073/pnas.1513508112> arXiv:<https://www.pnas.org/content/112/45/13743.full.pdf>
- [30] Sylvain Sorin. 1992. Repeated games with complete information. In *Handbook of Game Theory with Economic Applications*, R.J. Aumann and S. Hart (Eds.). Handbook of Game Theory with Economic Applications, Vol. 1. Elsevier, Chapter 4, 71–107. <https://ideas.repec.org/h/eee/gamchp/1-04.html>
- [31] Mária Svorenová and Marta Kwiatkowska. 2016. Quantitative verification and strategy synthesis for stochastic games. *European Journal of Control* 30 (2016), 15–30.
- [32] F. Thuijsman. 1992. *Optimality and Equilibria in Stochastic Games*. Number no. 82 in CWI Tract - Centrum voor Wiskunde en Informatica. Centrum voor Wiskunde en Informatica. <https://books.google.co.uk/books?id=sfzuAAAAAAJ>

A MATHEMATICAL BACKGROUND

LEMMA A.1. *Given an infinite sequence $(a_k)_{k \in \mathbb{N}}$ of real numbers with $0 \leq a_k \leq 1$. Then*

$$1 - \sum_{k=1}^n a_k \leq \prod_{k=1}^n (1 - a_k).$$

PROOF. The proof is by an induction on n . The base case vacuously holds. Assume that $1 - \sum_{k=1}^{n-1} a_k \leq \prod_{k=1}^{n-1} (1 - a_k)$ holds, then

$$\begin{aligned} \prod_{k=1}^n (1 - a_k) &= (1 - a_n) \prod_{k=1}^{n-1} (1 - a_k) \\ &\geq (1 - a_n) \left(1 - \sum_{k=1}^{n-1} a_k\right) \\ &\geq 1 - a_n + a_n \sum_{k=1}^{n-1} a_k - \sum_{k=1}^{n-1} a_k \\ &\geq 1 - \sum_{k=1}^n a_k \end{aligned}$$

□

B REDUCTION TO ACYCLIC GAMES IN SECTION 3

Following [15, Lemma 4], to show the existence of 1-bit Markov strategies in Büchi games, it suffices to show that 1-bit strategies suffice in a Büchi game that is made acyclic by encoding a step counter in the state space.

LEMMA B.1. *Suppose that for every acyclic Büchi game \mathcal{G}' and $\varepsilon > 0$ Max has a 1-bit strategy σ' such that*

- (1) $\sigma'[0]$ is multiplicatively ε -optimal from every state,
- (2) all memory updates $\sigma'_{up}(\cdot)$ are Dirac, and
- (3) σ' is deterministic if \mathcal{G}' is turn-based.

Then for every Büchi game \mathcal{G} and $\varepsilon > 0$ Max has a 1-bit Markov strategy σ such that

- (1) $\sigma[0]$ is multiplicatively ε -optimal from every state,
- (2) all memory updates $\sigma_{up}(\cdot)$ are Dirac, and
- (3) σ is deterministic if \mathcal{G} is turn-based.

PROOF. Let \mathcal{G} be a Büchi game with set S of states, where at each state $s \in S$ Max and Min have action sets $A(s)$ and $B(s)$, respectively. Let p be the transition function of \mathcal{G} . Let $\varepsilon > 0$.

We transform \mathcal{G} into an acyclic Büchi game \mathcal{G}' by encoding a step counter into the states. That is, the set of states in \mathcal{G}' is $S \times \mathbb{N}$. For each state (s, k) , with $k \in \mathbb{N}$, Max and Min have action $A(s)$ and $B(s)$, respectively. The transition function p' of \mathcal{G}' is defined by

$$p'((s, k), a, b)(s', j) \begin{cases} p(s, a, b)(s') & \text{if } j = k + 1 \\ 0 & \text{otherwise.} \end{cases}$$

Note that for each state $s \in S$ and each $k \in \mathbb{N}$ the value of s in \mathcal{G} equals the value of (s, k) in \mathcal{G}' .

By the assumption of the lemma statement, in \mathcal{G}' Max has a 1-bit strategy σ' with the properties (1)–(3) from the assumption. Obtain from σ' the 1-bit Markov strategy σ , which, at state s with memory mode $b \in \{0, 1\}$ and step counter n , plays like σ' at state (s, n) with mode b . Then σ satisfies the properties (1)–(3) from the conclusion of the lemma statement. In particular, since $\sigma'[0]$ is ε -optimal from every state in \mathcal{G}' , it follows that $\sigma[0]$ is ε -optimal from every state in \mathcal{G} . Indeed, $\sigma[0]$ is ε -optimal from every state in \mathcal{G} and for every value of the step counter, thus specifically also for the initial step counter value of 0. □

C MISSING PROOFS IN SECTION 5

LEMMA 5.1. *For all events $E \subseteq S^\omega$ and all states $s \in S$:*

$$\text{val}_{\mathcal{G}, E}(s) = \text{val}_{\mathcal{G}_\perp, E}(s).$$

PROOF. By the construction of \mathcal{G}_\perp , due to the fact that there is leakage in every step of the game, the value of every state s in \mathcal{G}_\perp is not more than the value of s in \mathcal{G} . Indeed, in every step there is some positive leakage to \perp and that is an error for the objective E . Below, we argue that despite the leakage the value of states remains equal in \mathcal{G} and \mathcal{G}_\perp .

Given $\varepsilon > 0$ and any Max strategy σ in \mathcal{G} we derive a Max strategy σ_\perp in \mathcal{G}_\perp , with only an additional ε error for E . Let σ_\perp be such that $\sigma_\perp(h_\perp) = a_\eta$ where $\eta = \varepsilon \cdot 2^{-(|h_\perp|+1)}$ and $\sigma(h) = a$, where h is the history in \mathcal{G} corresponding to h_\perp in \mathcal{G}_\perp ; in particular, every Max action a_q , with $q \in (0, 1)$, in h_\perp is replaced with the action a in h . Since $E \subseteq \text{Avoid}(\perp)$, for all Min strategies π , we have

$$\begin{aligned} \mathcal{P}_{\mathcal{G}_\perp, s, \sigma_\perp, \pi}(E) &= \mathcal{P}_{\mathcal{G}_\perp, s, \sigma_\perp, \pi}(E \mid \text{Avoid}(\perp)) \mathcal{P}_{\mathcal{G}_\perp, s, \sigma_\perp, \pi}(\text{Avoid}(\perp)) \\ &= \mathcal{P}_{\mathcal{G}, s, \sigma, \pi}(E) \mathcal{P}_{\mathcal{G}_\perp, s, \sigma_\perp, \pi}(\text{Avoid}(\perp)) && \text{by definition of } \sigma_\perp \\ &\geq \mathcal{P}_{\mathcal{G}, s, \sigma, \pi}(E) \left(1 - \sum_{i=1}^{\infty} \frac{\varepsilon}{2^i}\right) && \text{by Lemma A.1} \\ &\geq \mathcal{P}_{\mathcal{G}, s, \sigma, \pi}(E)(1 - \varepsilon). \end{aligned}$$

For all $i \geq 3$, every (multiplicatively) 2^{-i} -optimal strategy σ in \mathcal{G} transferred to \mathcal{G}_\perp in the described way above yields a (multiplicatively) 2^{-i+1} -optimal strategy σ_\perp in \mathcal{G}_\perp , concluding the proof. \square

The next lemma shows how to carry Max strategies in \mathcal{G}_\perp back to \mathcal{G} .

LEMMA C.1. *Let $E \subseteq S^\omega$ be an event. Let σ_\perp be an M -based Max strategy in \mathcal{G}_\perp , that is ε -optimal from every state (with a fixed initial mode). There exists an M -based Max strategy σ in \mathcal{G} , that is ε -optimal from every state (with a fixed initial mode).*

PROOF. Define the Max strategy σ in \mathcal{G} as follows. For each history h in \mathcal{G} , there exists a corresponding history h_\perp in \mathcal{G}_\perp that does not visit \perp , and every Max action a_q , with $q \in (0, 1)$, in h_\perp is replaced with the action a in h . Define $\sigma(h) = a$ if there exists $\eta \in (0, 1)$ such that $\sigma_\perp(h_\perp) = a_\eta$.

Clearly, σ simulates exactly σ_\perp except the leakage to \perp , which is an error for E . This implies that σ is ε -optimal from every state (with the given initial mode). Moreover, by definition, σ can be implemented with the exact set M of memory modes as σ_\perp does. \square

In order to prove Lemma 5.2, we prove a stronger result that implies this Lemma and is also used in Section 6.

LEMMA C.2. *In \mathcal{G}_\perp , for all finite-memory Max strategies σ , all Min strategies π and states s , the events Tr and $\text{Avoid}(\perp)$ differ only by a nullset. Formally,*

$$\mathcal{P}_{\mathcal{G}_\perp, s, \sigma, \pi}((\text{Tr} \setminus \text{Avoid}(\perp)) \cup (\text{Avoid}(\perp) \setminus \text{Tr})) = 0.$$

PROOF. Consider the set $\{1, 2, \dots\}$ of states in \mathcal{G}_\perp , excluding \perp . Write $E^n(i)$ for the event that the state i is visited at least n times. The event $E^\infty(i)$, defined by

$$E^\infty(i) \stackrel{\text{def}}{=} \bigcap_{n \geq i} E^n(i),$$

is the event that state i is visited infinitely often. Let \uplus denote the disjoint union. Since \perp is absorbing, we have

$$\text{Avoid}(\perp) = \text{Tr} \uplus \bigcup_{i \in \mathbb{N}} E^\infty(i).$$

It suffices to show that, for all $i \in \mathbb{N}$,

$$\mathcal{P}_{\mathcal{G}_\perp, s, \sigma, \pi}(E^\infty(i)) = 0.$$

Let M be the finite set of memory modes of the Max strategy σ . By definition, $E^{n+1}(s_i) \subseteq E^n(s_i)$ holds for all $n \in \mathbb{N}$. Then it follows that

$$\begin{aligned} \mathcal{P}_{\mathcal{G}_\perp, s, \sigma, \pi}(E^\infty(i)) &= \lim_{n \rightarrow \infty} \mathcal{P}_{\mathcal{G}_\perp, s, \sigma, \pi}(E^n(i)) && \text{by continuity of measures} \\ &\leq \lim_{n \rightarrow \infty} (1 - \eta_i)^{n-1} = 0 && \text{since no transition to } \perp \text{ is taken so far} \end{aligned}$$

where $\eta_i \stackrel{\text{def}}{=} \min_{m \in M} \sum_{a \in A(i)} \sum_{\eta \in (0, 1) \cap \mathbb{Q}} \eta \sigma_{act}(m, i)(a_\eta)$ is a lower bound for the probability of going to \perp in that step. By the construction of \mathcal{G}_\perp , at every state i and memory mode m , Max chooses among actions a_η with $a \in A(i)$ and $\eta \in (0, 1) \cap \mathbb{Q}$, and thus $\eta_i > 0$. \square

CLAIM 5.3.1. *The sequence $(X_n)_{n=1}^\infty$ is a supermartingale with respect to the filtration $(\mathcal{F}_n)_{n=1}^\infty$:*

$$\mathcal{E}_{s_0, \sigma, \pi}(X_{n+1} \mid \mathcal{F}_n) \leq X_n.$$

PROOF. For all $n \geq 1$, write $C_n \subseteq Z^* \times \{s_0\}$ for the set of histories that end in s_0 and visit s_0 exactly n times. For histories $h \in C_n$, write $\text{Cyl}(h) \subseteq Z^\omega$ for the cylinder set corresponding to h .

Let $n \geq 1$ and $h \in C_n$. We first show that

$$\mathcal{P}_{s_0, \sigma, \pi}(\text{Tr and } \tau = n \mid \text{Cyl}(h)) + \mathcal{P}_{s_0, \sigma, \pi}(\tau > n \mid \text{Cyl}(h)) \cdot u \leq u \quad (18)$$

holds. Indeed, by the definition of τ , the plays in $(\text{Tr and } \tau = n)$ are transient plays that visit s_0 exactly n times, so when starting with h , they have already visited s_0 for n times and will never return to s_0 . Recall the relation between strategies σ and σ_h and π_h and π :

$$\begin{array}{ccc}
 \sigma & \xrightarrow{\text{conditioned to } h} & \sigma_h \\
 & & \downarrow \\
 \pi & \xleftarrow{\text{reset after each visit to } s_0} & \pi_h
 \end{array}
 \quad \text{chosen to satisfy (5)}$$

Similarly, the plays in $\tau > n$ visit s_0 at least $n + 1$ times, so when starting with h , they must return to s_0 . Combining these two observations together with (5), we have

$$\begin{aligned}
 & \mathcal{P}_{s_0, \sigma, \pi}(\text{Tr and } \tau = n \mid \text{Cyl}(h)) + \mathcal{P}_{s_0, \sigma, \pi}(\tau > n \mid \text{Cyl}(h)) \cdot u \\
 &= \mathcal{P}_{s_0, \sigma_h, \pi_h}(\text{Tr} \setminus \text{Ret}) + \mathcal{P}_{s_0, \sigma_h, \pi_h}(\text{Ret}) \cdot u && \text{by def. of } \pi \\
 &\leq u && \text{by (5)}
 \end{aligned}$$

which completes the proof of (18).

Given an event E denote by $\mathbb{1}_E$ the indicator function of E . The next step in the proof is to show that, for all $n \geq 1$,

$$\mathcal{E}_{s_0, \sigma, \pi}(\mathbb{1}_{\tau \geq n} X_n \mid \mathcal{F}_n) = \sum_{h \in C_n} \mathbb{1}_{\text{Cyl}(h)} \mathcal{E}_{s_0, \sigma, \pi}(X_n \mid \text{Cyl}(h)). \quad (19)$$

For all $n \geq 1$, the event $\{\tau \geq n\}$ is naturally partitioned as $\{\tau \geq n\} = \uplus_{h \in C_n} \text{Cyl}(h)$. Since $\mathbb{1}_{\tau \geq n}$ is measurable, and then using the above-mentioned partition of $\{\tau \geq n\}$, we have

$$\begin{aligned}
 \mathcal{E}_{s_0, \sigma, \pi}(\mathbb{1}_{\tau \geq n} X_n \mid \mathcal{F}_n) &= \mathbb{1}_{\tau \geq n} \mathcal{E}_{s_0, \sigma, \pi}(X_n \mid \mathcal{F}_n) \\
 &= \sum_{h \in C_n} \mathbb{1}_{\text{Cyl}(h)} \mathcal{E}_{s_0, \sigma, \pi}(X_n \mid \mathcal{F}_n)
 \end{aligned}$$

Let $h \in C_n$, implying that h visits s_0 for n times. Recall that \mathcal{F}_n is the sigma-algebra generated by the cylinder sets corresponding to the histories $hs \in Z^* \times S$ such that h visit s_0 less than n times. Hence, each generating cylinder set of \mathcal{F}_n either is included in $\text{Cyl}(h)$ or has empty intersection with $\text{Cyl}(h)$. Due to this fact, the equality

$$\mathbb{1}_{\text{Cyl}(h)} \mathcal{E}_{s_0, \sigma, \pi}(X_n \mid \mathcal{F}_n) = \mathbb{1}_{\text{Cyl}(h)} \mathcal{E}_{s_0, \sigma, \pi}(X_n \mid \text{Cyl}(h))$$

holds, which concludes the proof of (19).

Finally to prove the statement of the claim, we will show that $\mathcal{E}_{s_0, \sigma, \pi}(X_{n+1} \mid \mathcal{F}_n) \leq X_n$ as follows. We can rewrite $\mathcal{E}_{s_0, \sigma, \pi}(X_{n+1} \mid \mathcal{F}_n)$ as

$$\mathcal{E}_{s_0, \sigma, \pi}(\mathbb{1}_{\tau < n} X_{n+1} + \mathbb{1}_{\tau \geq n} X_{n+1} \mid \mathcal{F}_n).$$

But since $X_{n+1} = X_n$ when $\tau < n$ the above sum in turn can be written as

$$\mathcal{E}_{s_0, \sigma, \pi}(\mathbb{1}_{\tau < n} X_n) + \mathcal{E}_{s_0, \sigma, \pi}(\mathbb{1}_{\tau \geq n} X_{n+1} \mid \mathcal{F}_n).$$

Recall that $\mathbb{1}_{\tau \geq n}$ is \mathcal{F}_n -measurable, then

$$\begin{aligned}
 \mathcal{E}_{s_0, \sigma, \pi}(X_{n+1} \mid \mathcal{F}_n) &= \mathbb{1}_{\tau < n} X_n + \mathcal{E}_{s_0, \sigma, \pi}(\mathbb{1}_{\tau \geq n} X_{n+1} \mid \mathcal{F}_n) \\
 &\stackrel{\text{by (19)}}{=} \mathbb{1}_{\tau < n} X_n + \sum_{h \in C_n} \mathbb{1}_{\text{Cyl}(h)} \mathcal{E}_{s_0, \sigma, \pi}(X_{n+1} \mid \text{Cyl}(h)) \\
 &= \mathbb{1}_{\tau < n} X_n + \sum_{h \in C_n} \mathbb{1}_{\text{Cyl}(h)} (\mathcal{P}_{s_0, \sigma, \pi}(\text{Tr and } \tau = n \mid \text{Cyl}(h)) + \mathcal{P}_{s_0, \sigma, \pi}(\tau > n \mid \text{Cyl}(h)) \cdot u) \\
 &\stackrel{\text{by (18)}}{\leq} \mathbb{1}_{\tau < n} X_n + \sum_{h \in C_n} \mathbb{1}_{\text{Cyl}(h)} u \\
 &= \mathbb{1}_{\tau < n} X_n + \mathbb{1}_{\tau \geq n} u \\
 &= \mathbb{1}_{\tau < n} X_n + \mathbb{1}_{\tau \geq n} X_n \\
 &= X_n.
 \end{aligned}$$

That is, the sequence $(X_n)_{n=1}^{\infty}$ is a supermartingale with respect to the filtration $(\mathcal{F}_n)_{n=1}^{\infty}$. \square

CLAIM 5.3.2. *If \mathcal{G}_{\perp} is turn-based, there is a strategy σ , as described in Lemma 5.3, that is deterministic in the first step.*

PROOF. Let $|A(s_0)| > 1$. Towards a contradiction, suppose that for all $\tilde{\sigma}$ that are deterministic in the first step there is π such that $\mathcal{P}_{s_0, \tilde{\sigma}, \pi}(\text{Tr} \setminus \text{Ret}) + \mathcal{P}_{s_0, \tilde{\sigma}, \pi}(\text{Ret})u \leq u$. Let σ be an arbitrary Max strategy. It can be decomposed, in a natural way, into a distribution $\alpha \in \mathcal{D}(A(s_0))$ and strategies $(\sigma_a)_{a \in A(s_0)}$, where σ_a deterministically plays a in the first step. By the assumption above, for each $a \in A(s_0)$ there is π_a with $\mathcal{P}_{s_0, \sigma_a, \pi_a}(\text{Tr} \setminus \text{Ret}) + \mathcal{P}_{s_0, \sigma_a, \pi_a}(\text{Ret})u \leq u$. Since \mathcal{G}_\perp is turn-based, we have $|B(s_0)| = 1$, and so there is a Min strategy π that plays π_a for the $a \in A(s_0)$ that Max is observed to choose in the first step. Then we have

$$\begin{aligned} \mathcal{P}_{s_0, \sigma, \pi}(\text{Tr} \setminus \text{Ret}) + \mathcal{P}_{s_0, \sigma, \pi}(\text{Ret})u &= \sum_{a \in A(s_0)} \alpha(a) (\mathcal{P}_{s_0, \sigma_a, \pi_a}(\text{Tr} \setminus \text{Ret}) + \mathcal{P}_{s_0, \sigma_a, \pi_a}(\text{Ret})u) \\ &\leq \sum_{a \in A(s_0)} \alpha(a)u = u. \end{aligned}$$

Since σ was arbitrary, we have derived (4). From there the proof by contradiction proceeds identically to the concurrent case. \square

CLAIM 5.4.1. *The sequence $(X_n)_{n=1}^\infty$ is a submartingale with respect to the filtration $(\mathcal{F}_n)_{n=1}^\infty$:*

$$\mathcal{E}_{s_0, \sigma^*, \pi}(X_{n+1} \mid \mathcal{F}_n) \geq X_n.$$

PROOF. This proof is analogous to the proof of Claim 5.3.1.

As in Claim 5.3.1, we write $C_n \subseteq Z^* \times \{s_0\}$, with $n \geq 1$, for the set of histories that end in s_0 and visit s_0 exactly n times. For histories $h \in C_n$, write $\text{Cyl}(h) \subseteq Z^\omega$ for the cylinder set corresponding to h .

Given a history $\mathbf{h} \in H$ we define $\pi_{\mathbf{h}}$ for the strategy whose behavior is exactly like the behavior of π after \mathbf{h} . Formally, for all histories $h \in H$ define $\pi_{\mathbf{h}}(h) = \pi(\mathbf{h}h)$.

Let $n \geq 1$ and $h \in C_n$. Similar to (18) in Claim 5.3.1, the following can be shown:

$$\begin{aligned} &\mathcal{P}_{s_0, \sigma^*, \pi}(\text{Tr and } \tau = n \mid \text{Cyl}(h)) + \mathcal{P}_{s_0, \sigma^*, \pi}(\tau > n \mid \text{Cyl}(h)) \cdot u \\ &= \mathcal{P}_{s_0, \sigma, \pi_{\mathbf{h}}}(\text{Tr} \setminus \text{Ret}) + \mathcal{P}_{s_0, \sigma, \pi_{\mathbf{h}}}(\text{Ret}) \cdot u && \text{by definition of } \sigma^* \text{ and } \pi_{\mathbf{h}} \\ &> u && \text{by Lemma 5.3.} \end{aligned} \tag{20}$$

Next, identical to (19) in the proof of Claim 5.3.1 we show that, for all $n \geq 1$,

$$\mathcal{E}_{s_0, \sigma^*, \pi}(\mathbb{1}_{\tau \geq n} X_n \mid \mathcal{F}_n) = \sum_{h \in C_n} \mathbb{1}_{\text{Cyl}(h)} \mathcal{E}_{s_0, \sigma^*, \pi}(X_n \mid \text{Cyl}(h)), \tag{21}$$

where $\mathbb{1}_E$ is the indicator function of E .

Finally, analogously, we conclude that

$$\begin{aligned} \mathcal{E}_{s_0, \sigma^*, \pi}(X_{n+1} \mid \mathcal{F}_n) &= \mathbb{1}_{\tau < n} X_n + \mathcal{E}_{s_0, \sigma^*, \pi}(\mathbb{1}_{\tau \geq n} X_{n+1} \mid \mathcal{F}_n) \\ &\stackrel{\text{by (21)}}{=} \mathbb{1}_{\tau < n} X_n + \sum_{h \in C_n} \mathbb{1}_{\text{Cyl}(h)} \mathcal{E}_{s_0, \sigma^*, \pi}(X_{n+1} \mid \text{Cyl}(h)) \\ &= \mathbb{1}_{\tau < n} X_n + \sum_{h \in C_n} \mathbb{1}_{\text{Cyl}(h)} (\mathcal{P}_{s_0, \sigma^*, \pi}(\text{Tr and } \tau = n \mid \text{Cyl}(h)) + \mathcal{P}_{s_0, \sigma^*, \pi}(\tau > n \mid \text{Cyl}(h)) \cdot u) \\ &\stackrel{\text{by (20)}}{\geq} \mathbb{1}_{\tau < n} X_n + \sum_{h \in C_n} \mathbb{1}_{\text{Cyl}(h)} u \\ &= \mathbb{1}_{\tau < n} X_n + \mathbb{1}_{\tau \geq n} u \\ &= \mathbb{1}_{\tau < n} X_n + \mathbb{1}_{\tau \geq n} X_n \\ &= X_n. \end{aligned}$$

That is, the sequence $(X_n)_{n=1}^\infty$ is a submartingale with respect to the filtration $(\mathcal{F}_n)_{n=1}^\infty$. \square

CLAIM 5.5.1. *The sequence $(X_n)_{n=0}^\infty$ is a submartingale with respect to the filtration $(\mathcal{F}_n)_{n=0}^\infty$:*

$$\mathcal{E}_{\mathcal{G}_\perp, s_0, \tilde{\sigma}, \tilde{\pi}}(X_{n+1} \mid \mathcal{F}_n) \geq X_n.$$

PROOF. In this proof all probabilities and expectations refer to the game \mathcal{G}_\perp . Denote by Vis the event that s_0 is visited. Following the definition of X_1 in (3):

$$\begin{aligned} &\mathcal{E}_{s, \tilde{\sigma}, \tilde{\pi}}(X_1 \mid \mathcal{F}_0) \\ &= \mathcal{P}_{s, \sigma_s, \tilde{\pi}}(\text{Vis}) \cdot u + \mathcal{P}_{s, \sigma_s, \tilde{\pi}}(\text{Tr and not Vis}) \cdot 1 \\ &\quad + \mathcal{P}_{s, \sigma_s, \tilde{\pi}}(\text{not Tr and not Vis}) \cdot 0 \end{aligned}$$

Substituting u from (1) we obtain

$$\begin{aligned}
 &= r \mathcal{P}_{s, \sigma_s, \tilde{\pi}}(\text{Vis}) \text{val}_{\text{Tr}}(s_0) + \mathcal{P}_{s, \sigma_s, \tilde{\pi}}(\text{Tr and not Vis}) \\
 &\geq r (\mathcal{P}_{s, \sigma_s, \tilde{\pi}}(\text{Vis}) \text{val}_{\text{Tr}}(s_0) + \mathcal{P}_{s, \sigma_s, \tilde{\pi}}(\text{Tr and not Vis})) \\
 &\geq r \inf_{\pi} (\mathcal{P}_{s, \sigma_s, \pi}(\text{Vis}) \text{val}_{\text{Tr}}(s_0) + \mathcal{P}_{s, \sigma_s, \pi}(\text{Tr and not Vis})) \\
 &\geq r \inf_{\pi} \mathcal{P}_{s, \sigma_s, \pi}(\text{Tr})
 \end{aligned}$$

By (8), we further have that

$$\begin{aligned}
 &\geq r x \text{val}_{\text{Tr}}(s) \\
 &= X_0.
 \end{aligned}$$

Now the proof given for the Claim 5.4.1 showing that $(X_n)_{n=1}^{\infty}$ is a submartingale with respect to the filtration $(\mathcal{F}_n)_{n=1}^{\infty}$, combined with the calculation above concludes the statement of the Claim. \square

D MISSING PROOFS FROM SECTION 6

CLAIM 6.2.1. *For any Min strategy π ,*

$$\mathcal{E}_{\mathcal{G}_{\infty}, s_1, \pi} Y_n \geq \mathcal{E}_{\mathcal{G}_{\infty}, s_1, \pi} Y_1 \quad \text{for all } n \geq 1.$$

PROOF. For all $n \geq 1$ and any Min strategy π we have

$$\begin{aligned}
 &\mathcal{E}_{\mathcal{G}_{\infty}, s_1, \pi}(Y_{n+1} \mid \mathcal{F}_n) \\
 &= \mathcal{E}_{\mathcal{G}_{\infty}, s_1, \pi}(r_{m(n+1)} v_{m(n+1)}(s_{n+1}) \mid \mathcal{F}_n) \\
 &\geq \mathcal{E}_{\mathcal{G}_{\infty}, s_1, \pi}(r_{m(n)} v_{m(n)}(s_{n+1}) \mid \mathcal{F}_n) && \text{by (17)} \\
 &\geq \mathcal{E}_{\mathcal{G}_{\infty}, s_1, \pi}(r_{m(n)} v_{m(n)}(s_n) \mid \mathcal{F}_n) && \text{see below} \\
 &= Y_n,
 \end{aligned}$$

where the inequality marked with “see below” follows from the observation that the successor state distribution in state s_n of \mathcal{G}_{∞} has been fixed already in $\mathcal{G}_{m(n)}$, and, therefore, the expected value, in terms of \mathcal{G}_{∞} , cannot drop in one step (though it might increase if π plays poorly). Thus, Y_1, Y_2, \dots is a submartingale w.r.t. $\mathcal{E}_{\mathcal{G}_{\infty}, s_1, \pi}$ for any Min strategy π and thus the claim follows. \square

D.1 Proof of Lemma 6.2

It remains to show Lemma 6.2. In the rest of this section, all measures and expectations are wrt. the leaky game \mathcal{G}_{\perp} . First we will need the following general technical lemma.

LEMMA D.1. *Let $f : (0, 1]^n \rightarrow (0, 1]^n$ be a continuous function. For any $x \in (0, 1)^n$ there is $y \in (0, 1)^n$ such that for all $i \in \{1, \dots, n\}$*

$$f(y)_i > y_i \implies y_i > x_i \quad \text{and} \quad f(y)_i \leq y_i \implies y_i = x_i.$$

PROOF. Define $y^{(1)} \stackrel{\text{def}}{=} x$. For any $m \geq 1$ for which $y^{(m)}$ has been defined, let

$$E^{(m)} \stackrel{\text{def}}{=} \{i \in \{1, \dots, n\} \mid f(y^{(m)})_i > y_i^{(m)} = x_i\}.$$

If $E^{(m)} = \emptyset$, define $y \stackrel{\text{def}}{=} y^{(m)}$. Otherwise, since f is continuous, we can define a vector $y^{(m+1)}$ with $y^{(m+1)} \geq y^{(m)}$ (in all components) such that $1 \geq f(y^{(m+1)})_i > y_i^{(m+1)} > y_i^{(m)}$ holds for all $i \in \{1, \dots, n\}$ with $f(y^{(m)})_i > y_i^{(m)}$. Since the $E^{(m)}$ are pairwise disjoint, this defines the required vector y . \square

Fix a finite set of states $I \subseteq S$ with $\perp \notin I$ for the rest of the section. For $s \in I$, define Ret_s as the following event: visit a state in $T \setminus I$; then (necessarily at least one step later) visit a state in I and the first such visited state in I is s . Events $(\text{Ret}_s)_{s \in I}$ are pairwise disjoint. Define $\text{Ret} \stackrel{\text{def}}{=} \bigcup_{s \in I} \text{Ret}_s$. Towards a proof of Lemma 6.2 we first prove the following lemma. (Here and in similar cases, we often write u_s for $u(s)$, where $u \in (0, 1]^I$ and $s \in I$.)

LEMMA D.2. *Let $q \in (0, 1)$. There are $u \in (0, 1)^I$ with $u_s \geq q \text{val}_{\text{TB}}(s)$ for all $s \in I$ and for all $s_0 \in I$ there is a Max strategy σ such that*

$$\inf_{\pi} \left(\mathcal{P}_{s_0, \sigma, \pi}(\text{TB} \setminus \text{Ret}) + \sum_{s \in I} \mathcal{P}_{s_0, \sigma, \pi}(\text{Ret}_s) u_s \right) > u_{s_0}$$

PROOF. Recall that we assume without loss of generality that \perp is the only state with $\text{val}_{\text{TB}}(\perp) = 0$ and $\perp \notin I$. Consider the function $f : \mathbb{R}^I \rightarrow \mathbb{R}^I$ defined by, for all $s_0 \in I$,

$$f(x)_{s_0} \stackrel{\text{def}}{=} \sup_{\sigma} \inf_{\pi} \left(\mathcal{P}_{s_0, \sigma, \pi}(\text{TB} \setminus \text{Ret}) + \sum_{s \in I} \mathcal{P}_{s_0, \sigma, \pi}(\text{Ret}_s) x_s \right)$$

From the assumptions on I and the disjointedness of the events $(\text{Ret}_s)_{s \in I}$, it follows that f is also $f : (0, 1]^I \rightarrow (0, 1]^I$. Since f is continuous, we can use Lemma D.1, instantiated with $n = |I|$ and $x_{s_0} = q\text{val}_{\text{TB}}(s_0) \in (0, 1)$, and obtain that there is $u \in (0, 1)^I$ such that for all $s_0 \in I$ we have $u_{s_0} \geq q\text{val}_{\text{TB}}(s_0)$, and we have $f(u)_{s_0} > u_{s_0}$ if and only if $u_{s_0} > q\text{val}_{\text{TB}}(s_0)$. Let $J \stackrel{\text{def}}{=} \{s_0 \in I \mid u_{s_0} = q\text{val}_{\text{TB}}(s_0)\}$. It suffices to show that $J = \emptyset$.

Choose $q' \in (q, 1)$ such that $u_s > q'\text{val}_{\text{TB}}(s)$ for all $s \in I \setminus J$. For $s_0 \in J$ and a strategy σ write

$$g(s_0, \sigma) \stackrel{\text{def}}{=} \inf_{\pi} \left(\mathcal{P}_{s_0, \sigma, \pi}(\text{TB} \setminus \text{Ret}) + \sum_{s \in I} \mathcal{P}_{s_0, \sigma, \pi}(\text{Ret}_s) u_s \right).$$

For all $s_0 \in J$ and all σ we have $g(s_0, \sigma) \leq u_{s_0}$. It follows from the construction of \mathcal{G}_{\perp} that there is always a *strictly* “better” strategy σ' , i.e., $g(s_0, \sigma') > g(s_0, \sigma)$. Thus, for all $s_0 \in J$ and all σ we have $g(s_0, \sigma) < u_{s_0}$. Hence,

$$\forall s_0 \in J \forall \sigma \exists \pi : \mathcal{P}_{s_0, \sigma, \pi}(\text{TB} \setminus \text{Ret}) + \sum_{s \in I} \mathcal{P}_{s_0, \sigma, \pi}(\text{Ret}_s) u_s \leq u_{s_0}. \quad (22)$$

(We could even have $< u_{s_0}$ above, but that is not needed.) For a play starting from a state in I , we define in the following a sequence of states, namely $s^{(1)}, s^{(2)}, \dots \in I$. This sequence may be finite, in which case $s^{(n)}$ is no longer defined from some n on. Each defined $s^{(n)}$ naturally corresponds to a visit of I . Specifically, denote by $s^{(1)} \in I$ the initial state, and, for $n \geq 1$, denote by $s^{(n+1)} \in I$ the state for which the play suffix started from $s^{(n)}$ satisfies $\text{Ret}_{s^{(n+1)}}$. In other words, after the visit of $s^{(n)}$ a state in $T \setminus I$ is visited and then a state in I is visited and the first such visited state is $s^{(n+1)}$. Denote by τ the random variable, taking values in $\{1, 2, \dots\} \cup \{\infty\}$, such that for all $n \in \{1, 2, \dots\}$ we have that $s^{(n)}$ is defined if and only if $\tau \geq n$. Intuitively, τ is the number of visits of I with visits of $T \setminus I$ in between. For all $n \in \{1, 2, \dots\}$, let \mathcal{F}_n be the sigma-algebra generated by the cylinder sets corresponding to the histories $w \in Z^* \times S$ such that w includes at most the visits $s^{(1)}, \dots, s^{(n-1)}$ (i.e., the state s at the end of history w may be the visit $s^{(n)}$). Note that the event $\tau \geq n$ and the random states $s^{(1)}, \dots, s^{(\min\{\tau, n\})}$ are \mathcal{F}_n -measurable.

Let τ' be the random variable, taking values in $\{1, 2, \dots\} \cup \{\infty\}$, such that if there is $n \in \{1, 2, \dots\}$ with $s^{(1)}, \dots, s^{(n-1)} \in J$ and $s^{(n)} \in I \setminus J$ then $\tau' \stackrel{\text{def}}{=} n$, and $\tau' \stackrel{\text{def}}{=} \tau + 1$ if such n does not exist. Note that *either* $\infty > \tau' \leq \tau$ and $s^{(\tau')} \in I \setminus J$ is defined or $\tau' = \tau + 1$ and $s^{(\tau')}$ is not defined. Random variable τ' is a stopping time with respect to the filtration $\mathcal{F}_1, \mathcal{F}_2, \dots$. Define random variables X_1, X_2, \dots by

$$X_n \stackrel{\text{def}}{=} \begin{cases} u_{s^{(n)}} & \text{if } \tau' > n \\ u_{s^{(\tau')}} & \text{if } \tau' \leq n \text{ and } \tau' \leq \tau \\ 1 & \text{if } \tau' = \tau + 1 \leq n \text{ and TB} \\ 0 & \text{if } \tau' = \tau + 1 \leq n \text{ and not TB.} \end{cases}$$

Each X_n is \mathcal{F}_n -measurable.

Let σ be an arbitrary strategy. The behavior of σ after any visit of I is determined by the history leading up to the visit. Thus, for every history h that ends with a visit of I there is a strategy, say σ_h , whose behavior is exactly like the behavior of σ after h . By (22) and the choice of q' , for every $s_0 \in I$ and every h that ends with a visit of s_0 there exists a (sufficiently good) Min strategy π_h such that if $s_0 \in J$ then

$$\mathcal{P}_{s_0, \sigma_h, \pi_h}(\text{TB} \setminus \text{Ret}) + \sum_{s \in I} \mathcal{P}_{s_0, \sigma_h, \pi_h}(\text{Ret}_s) u_s \leq u_{s_0} \quad (23)$$

and if $s_0 \in I \setminus J$ then (since $u_{s_0} > q'\text{val}_{\text{TB}}(s_0)$)

$$u_{s_0} \geq q' \mathcal{P}_{s_0, \sigma_h, \pi_h}(\text{TB}). \quad (24)$$

Let π be the Min strategy that upon every visit $s^{(n)}$ with $n \leq \min\{\tau, \tau'\}$ plays π_h (until the visit $s^{(n+1)}$ with $n+1 \leq \min\{\tau, \tau'\}$ if this visit exists, and forever otherwise).

For all $n \geq 1$ and $s_0 \in J$, write $C_{n, s_0} \subseteq Z^* \times \{s_0\}$ for the set of histories that include the visits $s^{(1)}, \dots, s^{(n)} \in J$ and end with $s^{(n)} = s_0$. For any $h \in C_{n, s_0}$, write $\text{Cyl}(h) \subseteq Z^{\omega}$ for the cylinder set corresponding to h . For all $n \geq 1$ we have $\{\tau' > n\} = \bigcup_{s_0 \in J} \bigcup_{h \in C_{n, s_0}} \text{Cyl}(h)$, where the union is disjoint. From the definitions of π and π_h (Equations (23) and (24)) it follows that for all $\tilde{s} \in J$ and $n \geq 1$ and $s_0 \in J$ and $h \in C_{n, s_0}$

$$\begin{aligned} & \mathcal{E}_{\tilde{s}, \sigma, \pi}(X_{n+1} \mid \text{Cyl}(h)) \\ &= \mathcal{P}_{\tilde{s}, \sigma, \pi}(\tau' - 1 = \tau = n \text{ and TB} \mid \text{Cyl}(h)) \\ & \quad + \sum_{s \in I} \mathcal{P}_{\tilde{s}, \sigma, \pi}(\tau \geq n+1 \text{ and } s^{(n+1)} = s \mid \text{Cyl}(h)) u_s \\ &= \mathcal{P}_{s_0, \sigma_h, \pi_h}(\text{TB} \setminus \text{Ret}) + \sum_{s \in I} \mathcal{P}_{s_0, \sigma_h, \pi_h}(\text{Ret}_s) u_s \\ &\leq u_{s_0}. \end{aligned} \quad (25)$$

Towards a contradiction, suppose that $J \neq \emptyset$, i.e., there is $\tilde{s} \in J$. To avoid clutter, let us drop the subscripts from $\mathcal{P}_{\tilde{s},\sigma,\pi}$ and $\mathcal{E}_{\tilde{s},\sigma,\pi}$ in the following. For all $n \geq 1$

$$\begin{aligned}
 \mathcal{E}(X_{n+1} \mid \mathcal{F}_n) &= \mathbb{1}_{\tau' \leq n} X_n + \sum_{s_0 \in J} \sum_{h \in \mathcal{C}_{n,s_0}} \mathbb{1}_{\text{Cyl}(h)} \mathcal{E}(X_{n+1} \mid \text{Cyl}(h)) \\
 &\stackrel{(25)}{\leq} \mathbb{1}_{\tau' \leq n} X_n + \sum_{s_0 \in J} \sum_{h \in \mathcal{C}_{n,s_0}} \mathbb{1}_{\text{Cyl}(h)} u_{s_0} \\
 &= \mathbb{1}_{\tau' \leq n} X_n + \sum_{s_0 \in J} \mathbb{1}_{\tau' > n, s^{(n)}=s_0} u_{s_0} \\
 &= \mathbb{1}_{\tau' \leq n} X_n + \mathbb{1}_{\tau' > n} u_{s^{(n)}} \\
 &= \mathbb{1}_{\tau' \leq n} X_n + \mathbb{1}_{\tau' > n} X_n \\
 &= X_n,
 \end{aligned}$$

where $\mathbb{1}_E$ is the indicator function of E . That is, X_1, X_2, \dots is a supermartingale with respect to the filtration $\mathcal{F}_1, \mathcal{F}_2, \dots$. Using the optional stopping theorem (noting that $|X_n| \leq \max\{1, \max_{s \in I} u_s\} \leq 1$ for all n), we obtain

$$\begin{aligned}
 q \text{val}_{\text{TB}}(\tilde{s}) &= u_{\tilde{s}} = \mathcal{E}X_1 \geq \mathcal{E}X_{\tau'} \\
 &\geq \mathcal{P}(\tau' = \tau + 1 \wedge \text{TB}) \\
 &\quad + \mathcal{P}(\infty > \tau' \leq \tau) \mathcal{E}(u_{s^{(\tau')}} \mid \infty > \tau' \leq \tau).
 \end{aligned} \tag{26}$$

Considering the second summand of the sum above, we have

$$\begin{aligned}
 &\mathcal{P}(\infty > \tau' \leq \tau) \mathcal{E}(u_{s^{(\tau')}} \mid \infty > \tau' \leq \tau) \\
 &= \sum_{s \in I \cup J} \mathcal{P}(\infty > \tau' \leq \tau \wedge s^{(\tau')} = s) u_s \\
 &\geq q' \sum_{s \in I \cup J} \mathcal{P}(\infty > \tau' \leq \tau \wedge s^{(\tau')} = s) \mathcal{P}(\text{TB} \mid \infty > \tau' \leq \tau \wedge s^{(\tau')} = s) \\
 &= q' \sum_{s \in I \cup J} \mathcal{P}(\infty > \tau' \leq \tau \wedge s^{(\tau')} = s \wedge \text{TB}) \\
 &= q' \mathcal{P}(\infty > \tau' \leq \tau \wedge \text{TB}),
 \end{aligned}$$

where the inequality follows from the definition of π . By combining this with (26) we obtain

$$\begin{aligned}
 q \text{val}_{\text{TB}}(\tilde{s}) &\geq q' (\mathcal{P}(\tau' = \tau + 1 \text{ and TB}) + \mathcal{P}(\infty > \tau' \leq \tau \text{ and TB})) \\
 &= q' \mathcal{P}_{\tilde{s},\sigma,\pi}(\text{TB}).
 \end{aligned}$$

Since the strategy σ was chosen arbitrarily, it follows that $q \text{val}_{\text{TB}}(\tilde{s}) \geq q' \text{val}_{\text{TB}}(\tilde{s})$. Since $\text{val}_{\text{TB}}(\tilde{s}) > 0$, we obtain $q \geq q'$, a contradiction. \square

For $s \in I$, define Vis_s as the event that a state in I is visited and the first such visited state is s . Events $(\text{Vis}_s)_{s \in I}$ are pairwise disjoint. Define $\text{Vis} \stackrel{\text{def}}{=} \bigcup_{s \in I} \text{Vis}_s$.

Let $u \in (0, 1)^I$ be as in Lemma D.2. Extend u so that $u \in (0, 1]^{I \cup T}$ by defining, for all $f \in T$,

$$u_f \stackrel{\text{def}}{=} \sup_{\sigma} \inf_{\pi} \left(\mathcal{P}_{f,\sigma,\pi}(\text{TB} \setminus \text{Vis}) + \sum_{s \in I} \mathcal{P}_{f,\sigma,\pi}(\text{Vis}_s) u_s \right). \tag{27}$$

We may have $u_f = 1$ for $f \in T \setminus I$. Since $I \cap T$ may be non-empty, the definition of u_f above for $u_f \in I \cap T$ might conflict with the construction of u from Lemma D.2. However, for $u_f \in I \cap T$ the definition above reduces to $u_f = u_f$, since f is the first state in I that is visited (immediately), i.e., $\mathcal{P}_{f,\sigma,\pi}(\text{Vis}_f) = 1$ holds for all $f \in I \cap T$ and all σ, π . Thus the extended u is well-defined.

Let U denote the random variable such that $U = u_f$ where f is the first visited state in $T \setminus I$, and $U = 0$ if $T \setminus I$ is not visited. Next we prove the following lemma.

LEMMA D.3. *Let σ be the strategy from Lemma D.2. We have*

$$\inf_{\pi} \mathcal{E}_{s_0,\sigma,\pi} U > u_{s_0} \quad \text{for all } s_0 \in I.$$

PROOF. Let $s_0 \in I$. For a play starting from s_0 denote by $f^{(1)} \in T$ the first visited state in $T \setminus I$. State $f^{(1)}$ is undefined if $T \setminus I$ is not visited. We have

$$\begin{aligned}
 & \inf_{\pi} \mathcal{E}_{s_0, \sigma, \pi} U \\
 &= \inf_{\pi} \sum_{f \in T \setminus I} \mathcal{P}_{s_0, \sigma, \pi}(f^{(1)} = f) u_f \\
 &\geq \inf_{\pi} \sum_{f \in T \setminus I} \left(\mathcal{P}_{s_0, \sigma, \pi}(f^{(1)} = f \text{ and } \text{TB} \setminus \text{Ret}) \right. \\
 &\quad \left. + \sum_{s \in I} \mathcal{P}_{s_0, \sigma, \pi}(f^{(1)} = f \text{ and } \text{Ret}_s) u_s \right) \\
 &= \inf_{\pi} \mathcal{P}_{s_0, \sigma, \pi}(\text{TB} \setminus \text{Ret}) + \sum_{s \in I} \mathcal{P}_{s_0, \sigma, \pi}(\text{Ret}_s) u_s \\
 &> u_{s_0}.
 \end{aligned}$$

The equality on the second line follows from the definitions of U and $f^{(1)}$. The inequality on the third line follows from a case split whether the play ever returns to I and the definitions of u_f , Vis , Vis_s , Ret and Ret_s ; cf. (27). The equality on the fourth line follows from the fact that TB and all Ret_s require a visit of $T \setminus I$, and the inequality on the last line follows from Lemma D.2. \square

We will need the following lemma on the existence of ε -optimal memoryless Max strategies for reachability objectives. It is a slight extension of previous results; see [20, Section 7.7], and [11, Theorem 12.1]. Recall that the objective $\text{Reach}_L(\top)$ is satisfied by plays that stay inside the set of states L until they reach the target state \top .

LEMMA D.4 ([17, LEMMA 5]). *Consider a concurrent game with state space S , where Max may have infinite action sets but Min has only finite action sets. Let $S_0 \subseteq S$ be a finite set of states, and $\top \in S$ a target state. For all $\varepsilon > 0$ there are a memoryless Max strategy σ_0 and a finite set $L \subseteq S$ such that $\inf_{\pi} \mathcal{P}_{s, \sigma_0, \pi}(\text{Reach}_L(\top)) \geq \text{val}_{\text{Reach}\top}(s) - \varepsilon$ holds for all $s \in S_0$. If the game is turn-based, there is a deterministic such σ_0 .*

Lemma D.4 is used in the proof of the following lemma, to “replace” the general Max strategy σ from Lemmas D.2 and D.3 by a memoryless one. For a bridge $\Lambda = (I, L, \sigma_0)$ (where I is as before) and plays starting from I , let U_{Λ} denote the random variable such that $U_{\Lambda} = u_f$ if $f \in T$ is the state that is first visited upon leaving $I \cup L$, and $U_{\Lambda} = 0$ if $I \cup L$ is not left or the state first visited upon leaving $I \cup L$ is not in T .

LEMMA D.5. *There is a bridge $\Lambda = (I, L, \sigma_0)$ (where I is as before) such that $L \cap T = \emptyset$ and for every Max strategy σ consistent with Λ and for all $s_0 \in I$ we have $\inf_{\pi} \mathcal{E}_{s_0, \sigma, \pi} U_{\Lambda} > u_{s_0}$. If \mathcal{G}_{\perp} is turn-based then σ_0 is deterministic.*

PROOF. We transform the “weighted reachability” objective defined by the random variable U to a reachability objective by constructing a new game, $\hat{\mathcal{G}}$, from \mathcal{G}_{\perp} as follows. Add a fresh state, say \top , and for each $f \in T \setminus I$ redirect each incoming transition so that with probability u_f it goes to \top and with probability $1 - u_f$ to the losing sink \perp . (This is possible, since $u_f \in (0, 1]$.)

Let σ_1 be the strategy from Lemma D.3; i.e., we have that $\inf_{\pi} \mathcal{E}_{\mathcal{G}_{\perp}, s, \sigma_1, \pi} U > u_s$ for all $s \in I$. Let $\varepsilon > 0$ be such that $\inf_{\pi} \mathcal{E}_{\mathcal{G}_{\perp}, s, \sigma_1, \pi} U - \varepsilon > u_s$ holds for all $s \in I$. By Lemma D.4 there are a memoryless (if \mathcal{G}_{\perp} is turn-based: memoryless deterministic) strategy σ_0 and a finite set $L \subseteq (S \setminus T) \cup I$ such that $\inf_{\pi} \mathcal{P}_{\hat{\mathcal{G}}, s, \sigma_0, \pi}(\text{Reach}_L(\top)) \geq \text{val}_{\hat{\mathcal{G}}, \text{Reach}\top}(s) - \varepsilon$ holds for all $s \in I$. Let $s \in I$ be arbitrary, and let σ be an arbitrary strategy consistent with $(I, L \setminus I, \sigma_0)$. It suffices to show that $\inf_{\pi} \mathcal{E}_{\mathcal{G}_{\perp}, s, \sigma, \pi} U_{\Lambda} > u_s$. We have

$$\begin{aligned}
 \inf_{\pi} \mathcal{E}_{\mathcal{G}_{\perp}, s, \sigma, \pi} U_{\Lambda} &= \inf_{\pi} \mathcal{P}_{\hat{\mathcal{G}}, s, \sigma, \pi}(\text{Reach}_L(\top)) \\
 &= \inf_{\pi} \mathcal{P}_{\hat{\mathcal{G}}, s, \sigma_0, \pi}(\text{Reach}_L(\top)) \\
 &\geq \text{val}_{\hat{\mathcal{G}}, \text{Reach}\top}(s) - \varepsilon \\
 &\geq \inf_{\pi} \mathcal{P}_{\hat{\mathcal{G}}, s, \sigma_1, \pi}(\text{Reach}\top) - \varepsilon \\
 &= \inf_{\pi} \mathcal{E}_{\mathcal{G}_{\perp}, s, \sigma_1, \pi} U - \varepsilon \\
 &> u_s,
 \end{aligned}$$

as required. \square

Let $\Lambda = (I, L, \sigma_0)$ be the bridge from Lemma D.5. For plays starting from I , define $\langle \Lambda \rangle_1$ as the following event: leave $I \cup L$; upon leaving $I \cup L$ immediately visit a state in T . Plays from I that are in $\langle \Lambda \rangle$ are also in $\langle \Lambda \rangle_1$ (but not vice-versa, since $\langle \Lambda \rangle$ requires this to happen after each visit to I , and not just once as for $\langle \Lambda \rangle_1$).

The following lemma strengthens Lemma D.2.

LEMMA D.6. *There is a Max strategy σ consistent with Λ such that, for all $s_0 \in I$,*

$$\inf_{\pi} \left(\mathcal{P}_{s_0, \sigma, \pi}(\langle \Lambda \rangle \cap \text{TB} \setminus \text{Ret}) + \sum_{s \in I} \mathcal{P}_{s_0, \sigma, \pi}(\langle \Lambda \rangle_1 \cap \text{Ret}_s) u_s \right) > u_{s_0}$$

PROOF. By Lemma D.5 there is $x \in (0, 1)$ such that for all σ consistent with Λ and all $s_0 \in I$ we have $x \inf_{\pi} \mathcal{E}_{s_0, \sigma, \pi} U_{\Lambda} > u_{s_0}$. By the definition of u_f for $f \in T$ (Equation (27)), there is a Max strategy σ_2 such that, for all $f \in T \setminus I$,

$$\inf_{\pi} \left(\mathcal{P}_{f, \sigma_2, \pi}(\text{TB} \setminus \text{Vis}) + \sum_{s \in I} \mathcal{P}_{f, \sigma_2, \pi}(\text{Vis}_s) u_s \right) \geq x u_f$$

We can choose σ_2 to be consistent with Λ . Let $s_0 \in I$ be arbitrary. For a play starting from s_0 denote by $f_{\Lambda} \in T$ the state that is first visited upon leaving $I \cup L$. State f_{Λ} is undefined if $I \cup L$ is not left or the state first visited upon leaving $I \cup L$ is not in T . Let σ be a strategy consistent with Λ which switches to σ_2 as soon as f_{Λ} is visited. We have

$$\begin{aligned} & \inf_{\pi} \left(\mathcal{P}_{s_0, \sigma, \pi}(\langle \Lambda \rangle \cap \text{TB} \setminus \text{Ret}) + \sum_{s \in I} \mathcal{P}_{s_0, \sigma, \pi}(\langle \Lambda \rangle_1 \cap \text{Ret}_s) u_s \right) \\ &= \inf_{\pi} \left(\mathcal{P}_{s_0, \sigma, \pi}(\langle \Lambda \rangle_1 \cap \text{TB} \setminus \text{Ret}) + \sum_{s \in I} \mathcal{P}_{s_0, \sigma, \pi}(\langle \Lambda \rangle_1 \cap \text{Ret}_s) u_s \right) \\ &= \inf_{\pi} \sum_{f \in T \setminus I} \left(\mathcal{P}_{s_0, \sigma, \pi}(f_{\Lambda} = f \text{ and } \text{TB} \setminus \text{Ret}) \right. \\ & \quad \left. + \sum_{s \in I} \mathcal{P}_{s_0, \sigma, \pi}(f_{\Lambda} = f \text{ and } \text{Ret}_s) u_s \right) \\ &= \inf_{\pi} \sum_{f \in T \setminus I} \left(\mathcal{P}_{s_0, \sigma, \pi}(f_{\Lambda} = f) \cdot \left(\mathcal{P}_{f, \sigma_2, \pi}(\text{TB} \setminus \text{Vis}) \right. \right. \\ & \quad \left. \left. + \sum_{s \in I} \mathcal{P}_{f, \sigma_2, \pi}(\text{Vis}_s) u_s \right) \right) \\ &\geq x \inf_{\pi} \sum_{f \in T \setminus I} \mathcal{P}_{s_0, \sigma, \pi}(f_{\Lambda} = f) u_f \\ &= x \inf_{\pi} \mathcal{E}_{s_0, \sigma, \pi} U_{\Lambda} \\ &> u_{s_0}, \end{aligned}$$

as required. \square

Now we can prove Lemma 6.2.

LEMMA 6.2. *Let $I \subseteq S$ be a finite set of states in \mathcal{G}_{\perp} (thus $\perp \notin I$). Let $r \in (0, 1)$. There are a bridge $\Lambda = (I, L, \sigma_0)$ and a Max strategy σ consistent with Λ so that*

$$\inf_{\pi} \mathcal{P}_{s, \sigma, \pi}(\langle \Lambda \rangle \cap \text{TB}) \geq r \text{val}_{\text{TB}}(s)$$

holds for all states $s \in S$. If \mathcal{G}_{\perp} is turn-based then there is a deterministic such bridge.

PROOF OF LEMMA 6.2. Choose $q, x \in (0, 1)$ such that $qx \geq r$. Let $u \in (0, 1]^{I \cup T}$ and the bridge $\Lambda = (I, L, \sigma_0)$ be as defined before. Let σ be the strategy from Lemma D.6. Define a strategy σ^* as follows. For any initial state \tilde{s} , strategy σ^* first plays a strategy $\sigma_{\tilde{s}}$ with $\inf_{\pi} \mathcal{P}_{\tilde{s}, \sigma_{\tilde{s}}, \pi}(\text{TB}) \geq x \text{val}_{\text{TB}}(\tilde{s})$; if I is entered (which is at once if $\tilde{s} \in I$), strategy σ^* switches to σ ; from then on, whenever $\langle \Lambda \rangle_1$ occurs and then the play returns to I , strategy σ^* restarts σ upon that return. Since σ is consistent with Λ , so is σ^* . Let $\tilde{s} \in S$ be an arbitrary initial state, and let π be an arbitrary Min strategy. It suffices to show that $\mathcal{P}_{\tilde{s}, \sigma^*, \pi}(\langle \Lambda \rangle \cap \text{TB}) \geq r \text{val}_{\text{TB}}(\tilde{s})$.

For a play starting from \tilde{s} , we define in the following a sequence of states, namely $s^{(1)}, s^{(2)}, \dots \in I$. This sequence may be finite, in which case $s^{(n)}$ is no longer defined from some n on. Each defined $s^{(n)}$ naturally corresponds to a visit of I . Specifically, denote by $s^{(1)} \in I$ the first visited state in I , and, for $n \geq 1$, denote by $s^{(n+1)} \in I$ the state for which the play suffix started from $s^{(n)}$ satisfies $\langle \Lambda \rangle_1 \cap \text{Ret}_{s^{(n+1)}}$. In other words, after the visit of $s^{(n)}$ the event $\langle \Lambda \rangle_1$ occurs and then a state in I is visited and the first such visited state is $s^{(n+1)}$. Note that σ^* restarts σ exactly upon each visit $s^{(n)}$. Denote by τ the random variable, taking values in $\{0, 1, 2, \dots\} \cup \{\infty\}$, such that for all $n \in \{0, 1, 2, \dots\}$ we have that $s^{(n)}$ is defined if and only if $\tau \geq n$. Let \mathcal{F}_0 be the sigma-algebra generated by the cylinder set corresponding to the history that consists only of the initial state \tilde{s} . For all $n \in \{1, 2, \dots\}$, let \mathcal{F}_n be the sigma-algebra generated by the cylinder sets corresponding to the histories $w \in Z^* \times S$ such that w includes at most the visits $s^{(1)}, \dots, s^{(n-1)}$ (i.e., the state s at the end of history w may be the visit $s^{(n)}$). For all $n \geq 0$, the event $\tau \geq n$ and the random states $s^{(1)}, \dots, s^{(\min\{\tau, n\})}$ are \mathcal{F}_n -measurable.

Define random variables $X_0 \stackrel{\text{def}}{=} r \text{val}_{\text{TB}}(\tilde{s})$ and X_1, X_2, \dots by

$$X_n \stackrel{\text{def}}{=} \begin{cases} u_{s^{(n)}} & \text{if } \tau \geq n \\ 1 & \text{if } \tau < n \text{ and } \langle \Lambda \rangle \cap \text{TB} \\ 0 & \text{if } \tau < n \text{ and not } (\langle \Lambda \rangle \cap \text{TB}). \end{cases}$$

Each X_n is \mathcal{F}_n -measurable. We have

$$\begin{aligned}
\mathcal{E}_{\tilde{s}, \sigma^*, \pi}(X_1 | \mathcal{F}_0) &= \sum_{s \in I} \mathcal{P}_{\tilde{s}, \sigma_s, \pi}(\text{Vis}_s) \cdot u_s + \mathcal{P}_{\tilde{s}, \sigma_s, \pi}(\langle \Lambda \rangle \cap \text{TB} \setminus \text{Vis}) \cdot 1 \\
&= \sum_{s \in I} \mathcal{P}_{\tilde{s}, \sigma_s, \pi}(\text{Vis}_s) u_s + \mathcal{P}_{\tilde{s}, \sigma_s, \pi}(\text{TB} \setminus \text{Vis}) \\
&\geq q \sum_{s \in I} \mathcal{P}_{\tilde{s}, \sigma_s, \pi}(\text{Vis}_s) \text{val}_{\text{TB}}(s) + \mathcal{P}_{\tilde{s}, \sigma_s, \pi}(\text{TB} \setminus \text{Vis}) \\
&\geq q \left(\sum_{s \in I} \mathcal{P}_{\tilde{s}, \sigma_s, \pi}(\text{Vis}_s) \text{val}_{\text{TB}}(s) + \mathcal{P}_{\tilde{s}, \sigma_s, \pi}(\text{TB} \setminus \text{Vis}) \right) \\
&\geq q \inf_{\pi'} \left(\sum_{s \in I} \mathcal{P}_{\tilde{s}, \sigma_s, \pi'}(\text{Vis}_s) \text{val}_{\text{TB}}(s) + \mathcal{P}_{\tilde{s}, \sigma_s, \pi'}(\text{TB} \setminus \text{Vis}) \right) \\
&\geq q \inf_{\pi'} \mathcal{P}_{\tilde{s}, \sigma_s, \pi'}(\text{TB}) \\
&\geq q x \text{val}_{\text{TB}}(\tilde{s}) \\
&\geq r \text{val}_{\text{TB}}(\tilde{s}) \\
&= X_0.
\end{aligned}$$

For all $n \geq 1$ and $s_0 \in I$, write $C_{n, s_0} \subseteq Z^* \times \{s_0\}$ for the set of histories that include the visits $s^{(1)}, \dots, s^{(n)} \in I$ and end with $s^{(n)} = s_0$. For any $h \in C_{n, s_0}$, write $\text{Cyl}(h) \subseteq Z^\omega$ for the cylinder set corresponding to h . For all $n \geq 1$ we have $\{\tau \geq n\} = \bigcup_{s_0 \in I} \bigcup_{h \in C_{n, s_0}} \text{Cyl}(h)$, where the union is disjoint. For any history h write π_h for the Min strategy whose behavior is exactly like the behavior of π after h . By the definition of σ^* and by Lemma D.6 we have for all $n \geq 1$ and $s_0 \in I$ and $h \in C_{n, s_0}$

$$\begin{aligned}
\mathcal{E}_{\tilde{s}, \sigma^*, \pi}(X_{n+1} | \text{Cyl}(h)) &= \mathcal{P}_{\tilde{s}, \sigma^*, \pi}(\tau = n \text{ and } \langle \Lambda \rangle \cap \text{TB} | \text{Cyl}(h)) \\
&\quad + \sum_{s \in I} \mathcal{P}_{\tilde{s}, \sigma^*, \pi}(\tau \geq n+1 \text{ and } s^{(n+1)} = s | \text{Cyl}(h)) u_s \\
&= \mathcal{P}_{s_0, \sigma, \pi_h}(\langle \Lambda \rangle \cap \text{TB} \setminus \text{Ret}) + \sum_{s \in I} \mathcal{P}_{s_0, \sigma, \pi_h}(\langle \Lambda \rangle_1 \cap \text{Ret}_s) u_s \\
&> u_{s_0}.
\end{aligned} \tag{28}$$

To avoid clutter, let us drop the subscripts from $\mathcal{P}_{\tilde{s}, \sigma^*, \pi}$ and $\mathcal{E}_{\tilde{s}, \sigma^*, \pi}$ in the following. For all $n \geq 1$

$$\begin{aligned}
\mathcal{E}(X_{n+1} | \mathcal{F}_n) &= \mathbb{1}_{\tau < n} X_n + \sum_{s_0 \in I} \sum_{h \in C_{n, s_0}} \mathbb{1}_{\text{Cyl}(h)} \mathcal{E}(X_{n+1} | \text{Cyl}(h)) \\
&\stackrel{(28)}{\geq} \mathbb{1}_{\tau < n} X_n + \sum_{s_0 \in I} \sum_{h \in C_{n, s_0}} \mathbb{1}_{\text{Cyl}(h)} u_{s_0} \\
&= \mathbb{1}_{\tau < n} X_n + \sum_{s_0 \in I} \mathbb{1}_{\tau \geq n, s^{(n)} = s_0} u_{s_0} \\
&= \mathbb{1}_{\tau < n} X_n + \mathbb{1}_{\tau \geq n} u_{s^{(n)}} \\
&= \mathbb{1}_{\tau < n} X_n + \mathbb{1}_{\tau \geq n} X_n \\
&= X_n,
\end{aligned}$$

where $\mathbb{1}_E$ is the indicator function of E . That is, X_0, X_1, X_2, \dots is a submartingale with respect to the filtration $\mathcal{F}_0, \mathcal{F}_1, \mathcal{F}_2, \dots$. Random variable $\tau + 1$ is a stopping time with respect to the same filtration. Using the optional stopping theorem (noting that $|X_n| \leq \max\{1, \max_{s \in I} u_s\} \leq 1$ for all n), we obtain

$$r \text{val}_{\text{TB}}(\tilde{s}) = \mathcal{E}X_0 \leq \mathcal{E}X_{\tau+1} \leq \mathcal{P}(X_{\tau+1} \neq 0),$$

as $0 \leq X_{\tau+1} \leq 1$. It follows from the definition of the random variables X_n that, if $\tau < \infty$ and not $(\langle \Lambda \rangle \cap \text{TB})$, then $X_{\tau+1} = 0$. Thus,

$$\begin{aligned}
r \text{val}_{\text{TB}}(\tilde{s}) &\leq \mathcal{P}(X_{\tau+1} \neq 0) \leq \mathcal{P}(\tau = \infty \text{ or } (\langle \Lambda \rangle \cap \text{TB})) \\
&\leq \mathcal{P}(\tau = \infty) + \mathcal{P}(\langle \Lambda \rangle \cap \text{TB}).
\end{aligned}$$

By the construction of \mathcal{G}_\perp and the strategy σ^* , there is $\delta > 0$ such that upon every visit of I corresponding to $s^{(1)}, s^{(2)}, \dots \in I$ the probability of falling into \perp in the very next step is at least δ . (This is because σ^* plays a fixed memoryless strategy upon entering I .) Thus, $\mathcal{P}(\tau = \infty) = 0$. Hence, $r \text{val}_{\text{TB}}(\tilde{s}) \leq \mathcal{P}(\langle \Lambda \rangle \cap \text{TB})$, as required. \square