# MATRIX ANALOGUES OF SOME ASYMPTOTIC METHODS FOR LAPLACE INTEGRALS

PENG-CHENG HANG

ABSTRACT. The matrix analogues of Laplace's method and Watson's lemma are derived via the approach described by Williams and Wong [J. Approx. Theory 24 (4) (1974), 378-384]. Some examples are also given.

#### 1. Introduction

Williams and Wong [14] studied the operator analogue of Watson's lemma. In their work, a key assumption for the matrix form of Watson's lemma is that the matrix is normal. Luke and Barker [7] tried to weaken the "normal" condition, but the constant  $M_1$  in [7, Eq. (40)] is not absolute. which motivates us to find a correct answer.

This paper is organised as follows. In Section 2, we deduce an operator version of Laplace's method. In Section 3, we make a revision of Luke and Barker's result [7, Theorem 3]. This paper concludes with some examples in Section 4.

#### 2. Operator analogue of Laplace's method

In this section, we adopt the definition of asymptotic expansions of operator-valued functions introduced in [14, Sect. 3] and establish the operator analogue of Laplace's method.

Let X be a Banach space over the complex field  $\mathbb{C}$  and Y be a dense subspace of X. Let A be a closed linear operator from Y into X. The spectrum and resolvent of A are denoted by  $\sigma(A)$  and  $R_{\lambda}(A) = (\lambda I - A)^{-1}$ , respectively. According to [14, Sect. 2], if, in addition, A satisfies the following conditions:

• (C<sub>1</sub>) There exists a positive  $\Delta$  such that

$$\sigma(A) \subset \left\{ \lambda \in \mathbb{C} : \lambda \neq 0 \text{ and } |\arg \lambda| \leqslant \frac{\pi}{2} - \Delta \right\},$$

• (C<sub>2</sub>) Let  $\omega(A) := \inf \{ \Re(\lambda) : \lambda \in \sigma(A) \}$ . There exists M > 0 and  $0 < \omega_1 \leq \omega(A)$  such that for any positive integer n,

$$\|R_{\lambda}(A)^{n}\| \leq M(\omega_{1}-\lambda)^{-n} \quad (\lambda < \omega_{1}),$$
(2.1)

then  $e^{tA}$  is well-defined for real t. Then we can define for  $0 < \alpha < \infty$  that

$$A^{-\alpha} = \frac{1}{\Gamma(\alpha)} \int_0^\infty t^{\alpha-1} \mathrm{e}^{-tA} \mathrm{d}t.$$
 (2.2)

Furthermore, there exists a strongly continuous semigroup,  $\{e^{-tA}: 0 \leq t < \infty\}$ , of bounded linear operators on X such that

$$\left\| \mathbf{e}^{-tA} \right\| \leqslant M \mathbf{e}^{-t\omega_1}, \quad \text{for all } 0 < t < \infty.$$
(2.3)

Let us state our main results. Our proof is based on the classical procedure in [16, p. 57-60].

**Theorem 2.1.** Let  $\{A_{\alpha}\}$  be a net of closed linear operators from Y into X, each satisfying conditions  $(C_1)$  and  $(C_2)$  with the same  $\Delta$  and M and such that there is some positive  $\eta$  with  $\omega_1(A_\alpha) \ge \eta \, \omega(A_\alpha)$  for each  $A_\alpha$ . Define the integral

$$I(z) := \int_{a}^{b} \varphi(x) \mathrm{e}^{-zh(x)} \mathrm{d}x,$$

where h(x) is real,  $a \in \mathbb{R}$  is finite and b(>a) is finite or infinite.

<sup>2020</sup> Mathematics Subject Classification. 33C47, 41A60, 44A10, 47A56.

Key words and phrases. Asymptotics, Watson's lemma, Laplace's method, special matrix functions.

#### P.-C. HANG

Assume that (i) h(x) > h(a) for all  $x \in (a, b)$ , and for every  $\delta > 0$  the infimum of h(x) - h(a) in  $[a + \delta, b)$  is positive; (ii) h'(x) and  $\varphi(x)$  are continuous in a neighborhood of x = a, except possibly at a; (iii) as  $x \to a^+$ , the following expansions hold

$$h(x) \sim h(a) + \sum_{s=0}^{\infty} a_s (x-a)^{s+\mu}, \quad \varphi(x) \sim \sum_{s=0}^{\infty} b_s (x-a)^{s+\lambda-1},$$

where the expansion for h(x) is differentiable,  $\lambda$  and  $\mu$  are positive constants, and  $a_0 > 0$ ; and (iv) the integral I(z) converges absolutely for all sufficiently large z.

Then the bounded linear operator  $I(A_{\alpha})$  has the asymptotic expansion

$$I(A_{\alpha}) \sim e^{-h(a)A_{\alpha}} \sum_{s=0}^{\infty} \Gamma\left(\frac{s+\lambda}{\mu}\right) c_s A_{\alpha}^{-\frac{s+\lambda}{\mu}}, \quad as \ \left\|A_{\alpha}^{-1}\right\| \to 0,$$
(2.4)

where the coefficients  $c_s$  are defined by the expansion

$$\frac{\varphi(x)}{h'(x)} \sim \sum_{s=0}^{\infty} c_s t^{(s+\lambda-\mu)/\mu}, \quad as \ t = h(x) - h(a) \to 0^+.$$
(2.5)

In particular,  $c_0 = b_0 \mu^{-1} a_0^{-\lambda/\mu}$ .

*Proof.* Let  $\omega_{1,\alpha} = \omega_1(A_\alpha)$  and  $\omega_\alpha = \omega(A_\alpha)$ . Then  $\omega_{1,\alpha}^{-1} \leq ||A_\alpha^{-1}|| (\eta \sin \Delta)^{-1}$  (see [14, Eq. (3)]). Thus  $\omega_{1,\alpha} \to +\infty$ , as  $||A_\alpha^{-1}|| \to 0$ . In view of this, using (2.3) and the condition (iv) gives

$$\|I(A_{\alpha})\| \leq \left\| e^{-h(a)A_{\alpha}} \right\| \int_{a}^{b} |\varphi(x)| \left\| e^{-(h(x)-h(a))A_{\alpha}} \right\| dx$$
$$\leq M \left\| e^{-h(a)A_{\alpha}} \right\| \int_{a}^{b} |\varphi(x)| \cdot e^{-\omega_{1,\alpha}(h(x)-h(a))} dx < \infty,$$

which confirms the boundness of  $I(A_{\alpha})$  for suitable  $\alpha$ .

Conditions (ii) and (iii) imply the existence of a number c in (a, b) such that h'(x) and  $\varphi(x)$  are continuous in (a, c], and h'(x) is positive there. Put T = h(c) - h(a) > 0 and introduce the new variable t = h(x) - h(a). Since h(x) is increasing in (a, c), we may write

$$e^{h(a)A_{\alpha}} \int_{a}^{c} \varphi(x) e^{-h(x)A_{\alpha}} dx = \int_{0}^{T} f(t) e^{-tA_{\alpha}} dt$$
(2.6)

with f(t) beging the continuous function in (0, T] given by

$$f(t) = \varphi(x) \frac{\mathrm{d}x}{\mathrm{d}t} = \frac{\varphi(x)}{h'(x)}$$

Now f(t) has the expansion (2.5) as  $t \to 0^+$ . Applying the operator analogue of Watson's lemma [14, Theorem 1] to the integral on the right-hand side of (2.6), we have

$$\int_{a}^{c} \varphi(x) \mathrm{e}^{-h(x)A_{\alpha}} \mathrm{d}x \sim \mathrm{e}^{-h(a)A_{\alpha}} \sum_{s=0}^{\infty} \Gamma\left(\frac{s+\lambda}{\mu}\right) c_{s} A_{\alpha}^{-\frac{s+\lambda}{\mu}}, \quad \text{as } \left\|A_{\alpha}^{-1}\right\| \to 0.$$
(2.7)

Now estimate the integral on the remaining interval (c, b). Let  $\omega_0$  be a value of z > 0 for which I(z) is absolutely convergent, ant set

$$\varepsilon = \inf_{c \leqslant x < b} (h(x) - h(a)).$$

By condition (i),  $\varepsilon$  is positive. Now for  $\omega_{1,\alpha} \ge \omega_0$ ,

$$\omega_{1,\alpha}[h(x) - h(a)] = (\omega_{1,\alpha} - \omega_0) [h(x) - h(a)] + \omega_0[h(x) - h(a)] \ge (\omega_{1,\alpha} - \omega_0) \varepsilon + \omega_0[h(x) - h(a)].$$
(2.8)

Let  $I_1(A_\alpha) := \int_c^b \varphi(x) e^{-h(x)A_\alpha} dx$ . Applying (2.3), (2.8) and the condition (iv), we get

$$\begin{aligned} \left\| \mathrm{e}^{h(a)A_{\alpha}} I_{1}(A_{\alpha}) \right\| &\leq M \int_{c}^{b} |\varphi(x)| \mathrm{e}^{-\omega_{1,\alpha}(h(x)-h(a))} \mathrm{d}x \\ &\leq M \mathrm{e}^{(\omega_{0}-\omega_{1,\alpha})\varepsilon} \int_{c}^{b} |\varphi(x)| \mathrm{e}^{-\omega_{0}(h(x)-h(a))} \mathrm{d}x = K \mathrm{e}^{-\omega_{1,\alpha}\varepsilon}. \end{aligned}$$

Recall that  $\omega_{1,\alpha}^{-1} \leq ||A_{\alpha}^{-1}|| (\eta \sin \Delta)^{-1}$ . Then

$$\|e^{h(a)A_{\alpha}}I_{1}(A_{\alpha})\| \leq Ke^{-C/\|A_{\alpha}^{-1}\|},$$
(2.9)

where  $C = \varepsilon \eta \sin \Delta$ . The result (2.4) follows from (2.7) and (2.9).

A similar discussion of [14, Corollary] gives the following result.

**Corollary 2.2.** Let  $\{A_{\alpha}\}$  be a net of bounded linear operators on a Hilbert space  $\mathcal{H}$ . If each  $A_{\alpha}$  is normal and satisfies the condition  $(C_1)$  with the same  $\Delta$ , and if  $\varphi(x)$  and h(x) satisfy conditions (i), (ii), (iii) and (iv), then the expansion (2.4) holds.

### 3. The work of Luke and Barker revisited

Luke and Barker [7, Sect. II] showed that their result [7, Theorem 3] satisfies the hypotheses of [14, Theorem 1]. However, the constant  $M_1$  in [7, Eq. (40)] depends on the index m. More explicitly, we have the following fact

$$M_{1} = \sum_{r=0}^{p} (m)_{r} |\lambda_{1} - \lambda|^{-r} > 1 + m |\lambda_{1} - \lambda|^{-1},$$

which implies that under their restrictions, the inequality (2.1) is not valid.

In this section, we shall make a revision of their result. Before this, let us do some preparations. For  $A \in \mathbb{C}^{m \times m}$ , let  $A^+ := \frac{1}{2} (A + A^H)$  and define its 2-norm by

$$|A|| = \sup_{\|x\|_2 = 1} \|Ax\|_2,$$

where for any vector  $x \in \mathbb{C}^{m \times 1}$ ,  $||x||_2 = \sqrt{x^H x}$ . Denote

$$\mu(A) = \min\left\{z \in \mathbb{C} : z \in \sigma\left(A^{+}\right)\right\}, \ \omega(A) = \min\{\Re(\lambda) : \lambda \in \sigma(A)\}.$$

By [3, p. 647], it follows that

$$\left\|\mathbf{e}^{-tA}\right\| \leqslant \mathbf{e}^{-t\mu(A)} \quad (t \ge 0). \tag{3.1}$$

Next we derive the matrix form of Watson's lemma.

**Theorem 3.1.** Let A be a matrix such that the condition  $(C_1)$  (see Sect. 2) is satisfied and there is a positive  $\eta$  with  $\mu(A) \ge \eta \omega(A)$ . Assume that: (a) the function f(t) is a locally integrable function on  $[0, \infty)$  and admits the expansion

$$f(t) = \sum_{n=1}^{\infty} a_n t^{n/r-1}, \quad |t| \le c + \delta,$$

where r, c and  $\delta$  are positive; and (b) there exist positive constants  $M_0$  and b independent of t such that  $|f(t)| < M_0 e^{bt}$  holds for  $t \ge c$ . Then

$$\int_{0}^{\infty} f(t) \mathrm{e}^{-tA} \mathrm{d}t \sim \sum_{n=1}^{\infty} a_n \Gamma(n/r) A^{-n/r}, \quad as \ \left\| A^{-1} \right\| \to 0.$$
(3.2)

*Proof.* Fix an integer  $N \ge 2$ . Clearly, there is a constant C such that for all t > 0,

$$\left| f(t) - \sum_{n=1}^{N-1} a_n t^{n/r-1} \right| \leqslant C t^{N/r-1} e^{bt}.$$
(3.3)

A similar derivation of [14, Eq. (3)] gives

$$\mu(A) \ge \left\| A^{-1} \right\|^{-1} \eta \sin \Delta. \tag{3.4}$$

This together with (3.1) and (2.2) shows that  $A^{-n/r}$  is well-defined. Using (2.2), we can infer that

$$R_N(A) := \int_0^\infty f(t) e^{-tA} dt - \sum_{n=1}^{N-1} a_n \Gamma(n/r) A^{-n/r}$$

satisfies

$$R_N(A) = \int_0^\infty \left( f(t) - \sum_{n=1}^{N-1} a_n t^{n/r-1} \right) e^{-tA} dt.$$

If  $||A^{-1}|| \leq \frac{\eta \sin \Delta}{2b}$ , then  $\mu(A) \geq 2b$  by (3.4) and hence

$$||R_N(A)|| \leqslant CM \int_0^\infty t^{N/r-1} \mathrm{e}^{(b-\mu(A))t} \mathrm{d}t$$

by virtue of (3.1) and (3.3). Since  $\mu(A) \to \infty$ , it follows from (3.4) that

$$||R_N(A)|| = \mathcal{O}\left(\mu(A)^{-N/r}\right) = \mathcal{O}\left(||A^{-1}||^{-N/r}\right)$$

Recall the inequality [14, p. 382]

$$||A^{-1}||^{-N/r} \leq C_1 ||A^{-(N-1)/r}|| ||A^{-1}||^{1/r}$$

where  $C_1 = C_1(N, r)$ . Thus  $||R_N(A)|| = o(||A^{-(N-1)/r}||)$ , which completes the proof.

**Corollary 3.2.** Let A be a normal matrix satisfying the condition  $(C_1)$  and f(t) be a function satisfying the conditions (a) and (b). Then the expansion (3.2) holds.

*Proof.* It suffices to show that  $\mu(A) \ge \eta \,\omega(A)$ . Indeed, the further fact that  $\mu(A) = \omega(A)$  follows from the spectral theorem for normal matrices.

### Remark 3.1.

- (1) Karamata's Tauberian theorem is a converse of Watson's lemma (see [2, Sect. XIII.5]) and its matrix analogue was given in [8].
- (2) New approaches are needed to derive the matrix analogues of asymptotic methods for Mellin convolution integrals developed in [6, 13, 15].

## 4. Examples

In this section, we deal with asymptotics of the gamma matrix function, Bessel matrix functions and matrix-valued Kummer function. Here we assume that A is a matrix in  $\mathbb{C}^{m \times m}$  and the norm ||A|| is a least upper bound on the matrices of complex numbers.

4.1. Gamma matrix function. If A is a matrix such that  $\Re(z) > 0$  for any eigenvalue z of A, then the gamma matrix function  $\Gamma(A)$  is defined as [4, Eq. (1)]

$$\Gamma(A) = \int_0^\infty t^{A-I} \mathrm{e}^{-t} \mathrm{d}t, \quad t^{A-I} \equiv \exp\left((A-I)\log t\right).$$

Moreover, if A + nI is invertible for all integer  $n \ge 0$ , then  $\Gamma(A)$  is invertible, and the reciprocal gamma function is defined as

$$\Gamma^{-1}(A) = A(A+I)\cdots(A+(n-1)I)\Gamma(A+nI)^{-1}, \quad n \ge 1.$$

Clearly,  $\Gamma^{-1}(A) = \Gamma(A)^{-1}$  if  $\sigma(A) \subset \mathbb{C} \setminus \mathbb{Z}_{\leq 0}$ .

Now we can establish asymptotics of the gamma matrix function.

**Theorem 4.1.** Let A be a matrix satisfying: (c)  $\sigma(A) \subset \{\lambda \in \mathbb{C} : \lambda \neq 0, |\arg(\lambda - \theta)| < \Delta\}$  with  $0 < \Delta < \frac{\pi}{2}$  and  $0 < \theta < 2\pi$ ; and (d) for an invertible matrix P such that  $P^{-1}AP = J$  is the Jordan normal form of A, the number  $n(P) := \|P\| \|P^{-1}\|$  is bounded as  $\|A^{-1}\| \to 0$ . Then

$$\Gamma(A) \sim \sqrt{2\pi} e^{-A} A^{A-I/2} \sum_{k=0}^{\infty} g_k A^{-k}, \quad as \ \left\|A^{-1}\right\| \to 0,$$
(4.1)

where  $g_k$  has explicit expressions. In particular,  $g_0 = 1$  and  $g_1 = \frac{1}{12}$ .

*Proof.* Note that the scaled gamma function

$$\Gamma^*(z) := \frac{1}{\sqrt{2\pi}} \mathrm{e}^z z^{1/2-z} \Gamma(z)$$

is analytic in the domain described in the condition (c). Since (4.1) holds for  $\Gamma^*(z)$  as  $|z| \to \infty$ , we can make use of [7, Theorem 1] and obtain that (4.1) holds for  $\Gamma^*(A)$  as  $||A^{-1}|| \to 0$ .

**Theorem 4.2.** If A is Hermitian and postive definite, then for any integer  $N \ge 1$ ,

$$\Gamma(A) = \sqrt{2\pi} e^{-A} A^{A-I/2} \left( \sum_{k=0}^{N-1} g_k A^{-k} + R_N(A) \right),$$

and the reminder  $R_N(A)$  has the estimate

$$\|R_N(A)\|_2 \leqslant \frac{1+\zeta(N)}{(2\pi)^{N+1}} \Gamma(N) \|A^{-1}\|_2^{-N},$$
(4.2)

where  $\zeta$  denotes the Riemann zeta function.

*Proof.* Now take the spectral norm for matrices:  $||A|| \equiv ||A||_2$ . According to [7, p. 100] (see also [11, p. 271]), it holds for  $||B||_2$  that if B is Hermitian and positive definite, then any inequality satisfied by f(z) must also be satisfied by f(B). Thus, (4.2) follows from the error bound  $R_N(z)$  for asymptotics of the scalar gamma function(see [1]) and the fact that A is positive definite.  $\Box$ 

**Remark 4.1.** It is of interest to find the matrix analogues of asymptotic methods for contour integrals, since these methods may be helpful to derive asymptotics of special matrix functions, such as the asymptotics of  $\Gamma(A)$  in the general case  $\sigma(A) \subset \{\lambda \in \mathbb{C} : \lambda \neq 0, |\arg \lambda| < \pi - \delta\}$ .

4.2. Bessel matrix functions. Integral expressions of Bessel matrix functions are given by

$$J_A(z) = \left(\frac{z}{2}\right)^A \frac{\Gamma^{-1}(A+I/2)}{\sqrt{\pi}} \int_{-1}^{1} \left(1-t^2\right)^{A-I/2} \cos(zt) dt, \quad |\arg z| < \pi;$$
  
$$I_A(z) = \left(\frac{z}{2}\right)^A \frac{\Gamma^{-1}(A+I/2)}{\sqrt{\pi}} \int_{-1}^{1} \left(1-t^2\right)^{A-I/2} \cosh(zt) dt, \quad |\arg z| < \frac{\pi}{2}$$

where  $z \neq 0$  and  $\mu(A) > -\frac{1}{2}$ ; see [12, Sect. 3] for details.

Let A be a normal matrix satisfying the condition  $(C_1)$  (see Section 2) and set

$$h(t) = -\log(1-t^2), \ L(A) = \left(\frac{z}{2}\right)^A \Gamma^{-1} (A+I/2) (A-I/2)^{-1/2}.$$

Then Theorem 2.1 confirms that as  $||A^{-1}|| \to 0$ ,

$$J_A(z) \sim L(A), \ z \neq 0, |\arg z| < \pi; \quad I_A(z) \sim L(A), \ z \neq 0, |\arg z| < \frac{\pi}{2}.$$
 (4.3)

Since A is normal, the fact that n(P) = 1 verifies for the spectral norm and thus n(P) is bounded for any matrix norm. Hence, applying Theorem 2.1 and (4.3), we get that as  $||A^{-1}|| \to 0$ ,

$$J_A(z) \sim \frac{1}{\sqrt{2\pi}} \left(\frac{ez}{2}\right)^A A^{-A-I/2}, \quad z \neq 0, |\arg z| < \pi;$$
  
$$I_A(z) \sim \frac{1}{\sqrt{2\pi}} \left(\frac{ez}{2}\right)^A A^{-A-I/2}, \quad z \neq 0, |\arg z| < \frac{\pi}{2}.$$

4.3. Matrix-valued Kummer function. Similar to the scalar case, the matrix-valued confluent hypergoemetric function  $_1F_1$  is defined as

$$_{1}F_{1}(a;b;A) := \sum_{n=0}^{\infty} \frac{(a)_{n}}{(b)_{n}} \frac{A^{n}}{n!}, \quad A \in \mathbb{C}^{m \times m}.$$

Moreover, we have the matrix analogue of Kummer's first formula [9, p. 125, Eq. (2)]

$${}_{1}F_{1}(a;b;A) = e^{A} {}_{1}F_{1}(b-a;b;-A).$$
(4.4)

Let A be a normal matrix satisfying the condition (C<sub>1</sub>). Then  $\mu(A) = \omega(A)$  and further,

 $\lambda_i(A^+) = \Re(\lambda_i(A)), \quad 1 \le i \le n, \tag{4.5}$ 

where  $A^+ = \frac{1}{2} (A + A^H)$  and  $\lambda_i(A)$  is the *i*-th eigenvalue of A. Write  $B = -\mathbf{i} \cdot \log A$  and then  $A^{\mathbf{i}t} = e^{-tB}$ . Now combining (4.5) with (3.1) and the condition (C<sub>1</sub>) gives the following estimate

$$\left\|A^{\mathrm{i}t}\right\| \leqslant \mathrm{e}^{\left(\frac{\pi}{2} - \Delta\right)|t|}, \quad t \in \mathbb{R}.$$
(4.6)

An immediate result of the moment inequality [5, p. 115] is

$$||A^{-r}|| \leq C_r ||A^{-1}||^r, \quad r > 0.$$

P.-C. HANG

Then for any positive integer N,

$$\|A^{-N-1/2}\| \leq \|A^{-N}\| \|A^{-1/2}\| \leq C \|A^{-N}\| \|A^{-1}\|^{1/2}.$$
(4.7)

Applying the estimates (4.6) and (4.7) to the classical procedures in [10, p. 107-108 and 193-195], we may readily obtain the Mellin-Barnes integral representation

$$\frac{\Gamma(a)}{\Gamma(b)} {}_1F_1(a;b;-A) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\Gamma(-s)\Gamma(s+a)}{\Gamma(s+b)} A^s ds$$

and the asymptotic expansion

$$_{1}F_{1}(a;b;-A) \sim \frac{\Gamma(b)}{\Gamma(b-a)} A^{-a} \sum_{n=0}^{\infty} \frac{(a)_{n} (a-b+1)_{n}}{n!} A^{-n}, \text{ as } ||A^{-1}|| \to 0$$

Due to (4.4), we also have the following expansion

$$_{1}F_{1}(a;b;A) \sim \frac{\Gamma(b)}{\Gamma(a)} e^{A} A^{a-b} \sum_{n=0}^{\infty} \frac{(c-a)_{n} (1-a)_{n}}{n!} A^{-n}, \text{ as } ||A^{-1}|| \to 0.$$

#### References

- W. G. C. Boyd, Gamma function asymptotics by an extension of the method of steepest descents, Proc. R. Soc. A: Math. Phys. 447 (1931) (1994), 609–630.
- [2] W. Feller, An introduction to probability theory and its applications, Volume 2. Vol. 81. John Wiley & Sons, 1991.
- [3] I. Higueras, B. Garcia-Celayeta, Logarithmic norms for matrix pencils, SIAM J. Matrix Anal. Appl. 20 (3) (1999), 646–666.
- [4] L. Jódar, J. C. Cortés, Some properties of Gamma and Beta matrix functions, Appl. Math. Lett. 11 (1) (1998), 89–93.
- [5] S. G. Krein, Linear differential equations in Banach space, Amer. Math Soc., 1971.
- [6] J. L. López, Asymptotic expansions of Mellin convolution integrals, SIAM Rev. 50 (2) (2008), 275–293.
- [7] Y. L. Luke, G. P. Barker, On asymptotic expansions for functions of matrix argument, J. Approx. Theory 25 (2) (1979), 93–104.
- [8] M. Meerschaert, H. P. Scheffler, Tauberian theorems for matrix regular variation, Trans. Am. Math. Soc. 365 (4) (2013), 2207–2221.
- [9] E. D. Rainville, Special functions, (1960).
- [10] R. B. Paris, D. Kaminski, Asymptotics and Mellin-Barnes integrals, Vol. 85. Cambridge University Press, 2001.
- [11] F. Riesz, B. S. Nagy, Functional analysis, Courier Corporation, 2012.
- [12] J. Sastre, L. Jódar, Asymptotics of the modified bessel and the incomplete gamma matrix functions, Appl. Math. Lett. 16 (6) (2003), 815–820.
- [13] A. Sidi, Asymptotic expansions of Mellin transforms and analogues of Watson's lemma, SIAM J. Math. Anal. 16 (4) (1985), 896–906.
- [14] J. J. Williams, R. Wong, Asymptotic expansion of operator-valued Laplace transform, J. Approx. Theory 12 (4) (1974), 378–384.
- [15] R. Wong, Explicit error terms for asymptotic expansions of Mellin convolutions, J. Math. Anal. Appl. 72 (2) (1979), 740–756.
- [16] R. Wong, Asymptotic approximations of integrals, Society for Industrial and Applied Mathematics, 2001.

PENG-CHENG HANG, SCHOOL OF MATHEMATICS AND STATISTICS, DONGHUA UNIVERSITY, SHANGHAI 201620, PEOPLE'S REPUBLIC OF CHINA.

Email address: mathroc618@outlook.com