

THE INTEGRAL CHOW RING OF THE STACK OF POINTED HYPERELLIPTIC CURVES

ALBERTO LANDI

ABSTRACT. We study the integral Chow ring of the stack $\mathcal{H}_{g,n}$ parametrizing n -pointed smooth hyperelliptic curves of genus g . We compute the integral Chow ring of $\mathcal{H}_{g,n}$ for $n = 1, 2$ completely, while for $3 \leq n \leq 2g + 2$ we compute it up to the additive order of a single class in degree 2. We obtain partial results also for $n = 2g + 3$. In particular, taking $g = 2$ and recalling that $\mathcal{H}_{2,n} = \mathcal{M}_{2,n}$, our results hold for $\mathrm{CH}^*(\mathcal{M}_{2,n})$ for $1 \leq n \leq 7$.

INTRODUCTION

Since the seminal work of David Mumford [Mum77], several researchers have put their efforts in computing (both integral and rational) Chow rings of natural stacks of curves, as the stack $\overline{\mathcal{M}}_{g,n}$ of stable n -pointed curves of genus g , and its smooth counterpart $\mathcal{M}_{g,n}$.

In this paper we are interested in the closed substack $\mathcal{H}_{g,n}$ of $\mathcal{M}_{g,n}$ classifying n -pointed smooth hyperelliptic curves of genus g . Notice that for $g = 2$ we have $\mathcal{M}_{2,n} = \mathcal{H}_{g,2}$.

Since a considerable number of papers have been written on the intersection theory of stacks, we recall only a few known results about $\mathcal{H}_{g,n}$ and related stacks. In [Lar21], Eric Larson has computed the Chow ring of $\overline{\mathcal{M}}_2$, while the Chow ring of \mathcal{M}_2 was previously computed by Angelo Vistoli in [Vis98]. This last result has been generalized to \mathcal{H}_g by Dan Edidin and Damiano Fulghesu in [EF09] for the even g case, and by Andrea Di Lorenzo in [DL18] for the odd g case. The pointed case was first addressed by Michele Pernice in [Per22], where he studied the $n = 1$ case. However, the result is not completely correct, due to a mistake at the end of the proof; in this article we supplement his ideas to correct it. Finally, Dan Edidin and Zhengning Hu studied the locus of $\mathcal{H}_{g,n}$ where all the sections have image in the Weierstrass divisor (in particular, $1 \leq n \leq 2g + 2$), and they have computed the Chow rings of those loci. We will use their results in this paper. Samir Canning and Hannah Larson ([CL22]) have computed the rational Chow ring of $\mathcal{H}_{g,n}$ for $n \leq 2g + 6$, using a different description of the stack.

We compute the integral Chow ring of $\mathcal{H}_{g,n}$ for $n = 1, 2$ completely, while for $3 \leq n \leq 2g + 3$ we find the generators and almost every relation. As a consequence, we recover [CL22, Corollary 1.5] when $n \leq 2g + 3$.

All our results are based on the techniques of [Lan23], where a new description of $\mathcal{H}_{g,n}$ is found and used to compute its Picard groups for all g and n .

Statement of the results and strategy. We work over a field of characteristic strictly greater than $2g$, where $g \geq 2$ is the genus. The main result of this paper is

Key words and phrases. Chow ring, intersection theory, hyperelliptic curve, moduli.

the (almost) computation of the integral Chow ring of $\mathcal{H}_{g,n}$ for $1 \leq n \leq 2g+3$. This ring has already been computed in the case $n = 1$ in [Per22] by Pernice, but the result is not correct as stated; we correct it by extending his ideas. The presentation of $\text{CH}^*(\mathcal{H}_{g,1})$ with the most geometric meaning that we get is described in the following Theorem; see the first Section for the definitions of the various classes.

Remark 0.1. We use the following notation. If we have a ring A and elements $\beta_1, \dots, \beta_s \in A$, and $p_1, \dots, p_r \in \mathbb{Z}[x_1, \dots, x_s]$ are polynomials in s variables, we write

$$A \simeq \frac{\mathbb{Z}[\beta_1, \dots, \beta_s]}{(p_1(\beta_1, \dots, \beta_s), \dots, p_r(\beta_1, \dots, \beta_s))}$$

to mean that there is a surjective homomorphism

$$\mathbb{Z}[x_1, \dots, x_s] \twoheadrightarrow A$$

sending x_i to α_i and with kernel generated by p_1, \dots, p_r .

Moreover, we use the square brackets to denote the Chow class associated to the closed subscheme written between them.

Theorem 0.2. *Let $\text{char } k > 2g$. Then,*

$$\text{CH}^*(\mathcal{H}_{g,1}) \simeq \frac{\mathbb{Z}[\psi_1, [\mathcal{W}_{g,n}^1]]}{((4g+2)((g+1)\psi_1 - (g-1)[\mathcal{W}_{g,1}^1]), [\mathcal{W}_{g,1}^1]^2 + \psi_1 \cdot [\mathcal{W}_{g,1}^1])}$$

regardless of the parity of g .

In general, we show that $\text{CH}^*(\mathcal{H}_{g,n})$ is generated in degree 1 if $1 \leq n \leq 2g+3$.

Proposition 0.3. *For all $1 \leq n \leq 2g+3$, $\text{CH}^*(\mathcal{H}_{g,n})$ is generated in degree 1.*

The main idea is to use an exact sequence

$$\bigoplus_{i < j} \text{CH}^*(\mathcal{H}_{g,n-1}^{i,j}) \longrightarrow \text{CH}^*(\mathcal{H}_{g,n}) \longrightarrow \text{CH}^*(\mathcal{H}_{g,n}^{\text{far}}) \longrightarrow 0$$

where $\mathcal{H}_{g,n-1}^{i,j}$ is an open substack of $\mathcal{H}_{g,n-1}$, and $\mathcal{H}_{g,n}^{\text{far}}$ is an open substack of $\mathcal{H}_{g,n}$, whose Chow ring is computable for $2 \leq n \leq 2g+3$ (not completely for $n = 2g+3$). Using induction on n and the result for $n = 1$, one gets that each piece on the left and right is generated in degree 1, and concludes that the same holds for the piece in the middle.

Moreover, one can find numerous relations between Chow classes using the results in [Lan23] on the Picard group of $\mathcal{H}_{g,n}$. All this can be used to compute the Chow ring in the case $n = 2$.

Theorem 0.4. *Let $\text{char } g > 2g$. Then, $\text{CH}^*(\mathcal{H}_{g,2})$ is generated by the classes of ψ_1 , $\mathcal{W}_{g,2}^1$ and $\mathcal{Z}_{g,2}^{1,2}$, and the subgroup of relations is generated by*

$$\begin{aligned} & (4g+2)((g+1)\psi_1 - (g-1)[\mathcal{W}_{g,2}^1]), \\ & [\mathcal{Z}_{g,2}^{1,2}]^2 + \psi_1 \cdot [\mathcal{Z}_{g,2}^{1,2}], \quad [\mathcal{W}_{g,2}^1]^2 + \psi_1 \cdot [\mathcal{W}_{g,2}^1], \quad [\mathcal{Z}_{g,2}^{1,2}] \cdot [\mathcal{W}_{g,2}^1], \\ & (2g+1)(g+1)\psi_1^2 + (2g+1)(3g-1)[\mathcal{W}_{g,2}^1]^2 - (2g+1)(g+1)[\mathcal{Z}_{g,2}^{1,2}]^2. \end{aligned}$$

Unfortunately, the relations found are not enough to determine the integral Chow ring in the case of $3 \leq n \leq 2g+2$. The only piece missing is the additive order of the degree 2 class $[\mathcal{Z}_{g,n}^{1,2}]^2$. We summarize the results here.

Theorem 0.5. *Let $\text{char } k > 2g$. Then, $\text{CH}^*(\mathcal{H}_{g,3})$ is generated by the classes of $\mathcal{W}_{g,3}^1$, $\mathcal{Z}_{g,3}^{1,2}$, $\mathcal{Z}_{g,3}^{1,3}$ and $\mathcal{Z}_{g,3}^{2,3}$, and the ideal of relations is generated by*

$$\begin{aligned} & (8g+4)[\mathcal{W}_{g,3}^1] - (4g+2)(g+1)([\mathcal{Z}_{g,3}^{1,2}] + [\mathcal{Z}_{g,3}^{1,3}] - [\mathcal{Z}_{g,3}^{2,3}]), \\ & [\mathcal{Z}_{g,3}^{1,2}] \cdot [\mathcal{Z}_{g,3}^{1,3}], \quad [\mathcal{Z}_{g,3}^{1,2}] \cdot [\mathcal{Z}_{g,3}^{2,3}], \quad [\mathcal{Z}_{g,3}^{1,3}] \cdot [\mathcal{Z}_{g,3}^{2,3}], \quad [\mathcal{Z}_{g,3}^{1,j}] \cdot [\mathcal{W}_{g,3}^1] \text{ for } j = 2, 3, \\ & [\mathcal{W}_{g,3}^1] \cdot [\mathcal{Z}_{g,3}^{2,3}] + (g+1)[\mathcal{Z}_{g,3}^{2,3}]^2, \quad 2[\mathcal{W}_{g,3}^1]^2 - (g+1)[\mathcal{Z}_{g,3}^{2,3}]^2, \\ & (2g+1)(g+1)([\mathcal{Z}_{g,3}^{i,j}]^2 + [\mathcal{Z}_{g,3}^{i',j'}]^2) \text{ for } i \neq j, i' \neq j', \quad b_g[\mathcal{Z}_{g,3}^{i,j}]^2 \text{ for } i \neq j. \end{aligned}$$

where b_g is either $(2g+1)(g+1)$ or $(4g+2)(g+1)$.

Theorem 0.6. *Let $\text{char } k > 2g$. Then, $\text{CH}^*(\mathcal{H}_{g,4})$ is generated by the classes of $\mathcal{W}_{g,4}^1$, $\mathcal{Z}_{g,4}^{1,2}$, $\mathcal{Z}_{g,4}^{1,3}$, $\mathcal{Z}_{g,4}^{1,4}$ and $\mathcal{Z}_{g,4}^{2,3}$, and the ideal of relations is generated by*

$$\begin{aligned} & (8g+4)[\mathcal{W}_{g,4}^1] - (4g+2)(g+1)([\mathcal{Z}_{g,4}^{1,2}] + [\mathcal{Z}_{g,4}^{1,3}] - [\mathcal{Z}_{g,4}^{2,3}]), \\ & [\mathcal{Z}_{g,4}^{1,i}] \cdot [\mathcal{Z}_{g,4}^{1,j}] \text{ for } 1 < i < j, \quad [\mathcal{Z}_{g,4}^{1,j}] \cdot [\mathcal{Z}_{g,4}^{2,3}] \text{ for } j = 2, 3, \quad [\mathcal{Z}_{g,4}^{1,j}] \cdot [\mathcal{W}_{g,3}^1] \text{ for } j > 1, \\ & [\mathcal{Z}_{g,4}^{2,3}] \cdot [\mathcal{W}_{g,3}^1] + (g+1)[\mathcal{Z}_{g,4}^{2,3}]^2, \quad [\mathcal{Z}_{g,4}^{2,3}]^2 = -[\mathcal{Z}_{g,4}^{2,3}] \cdot [\mathcal{Z}_{g,4}^{1,4}] = [\mathcal{Z}_{g,4}^{1,4}]^2, \\ & 2[\mathcal{W}_{g,4}^1]^2 - (g+1)[\mathcal{Z}_{g,4}^{2,3}]^2, \quad (g+1)[\mathcal{Z}_{g,4}^{i,j}]^3 \text{ for all } i \neq j, \\ & (g+1)([\mathcal{Z}_{g,4}^{i,j}]^2 - [\mathcal{Z}_{g,4}^{i',j'}]^2) \text{ for all } i \neq j, i' \neq j', \quad b_g[\mathcal{Z}_{g,4}^{i,j}]^2 \text{ for all } i \neq j \end{aligned}$$

for some integer $b_g = b'_g(g+1)$ such that b'_g divides $4g+2$.

Theorem 0.7. *Let $\text{char } k > 2g$ and $5 \leq n \leq 2g+2$. Then, $\text{CH}^*(\mathcal{H}_{g,n})$ is generated by the classes of $\mathcal{W}_{g,n}^1$, $\mathcal{Z}_{g,n}^{1,i}$ for $1 < i \leq n$ and $\mathcal{Z}_{g,n}^{2,3}$, and the ideal of relations is generated by*

$$\begin{aligned} & (8g+4)[\mathcal{W}_{g,n}^1] - (4g+2)(g+1)([\mathcal{Z}_{g,n}^{1,2}] + [\mathcal{Z}_{g,n}^{1,3}] - [\mathcal{Z}_{g,n}^{2,3}]), \\ & [\mathcal{Z}_{g,n}^{1,i}] \cdot [\mathcal{Z}_{g,n}^{1,j}] \text{ for } 1 < i < j, \quad [\mathcal{Z}_{g,n}^{1,j}] \cdot [\mathcal{Z}_{g,n}^{2,3}] \text{ for } j = 2, 3, \quad [\mathcal{Z}_{g,n}^{1,j}] \cdot [\mathcal{W}_{g,n}^1] \text{ for } j \neq 1, \\ & [\mathcal{Z}_{g,n}^{2,3}] \cdot [\mathcal{W}_{g,n}^1] + (g+1)[\mathcal{Z}_{g,n}^{2,3}]^2, \quad 2[\mathcal{W}_{g,n}^1]^2 - (g+1)[\mathcal{Z}_{g,n}^{2,3}]^2, \\ & [\mathcal{Z}_{g,n}^{2,3}]^2 = -[\mathcal{Z}_{g,n}^{2,3}] \cdot [\mathcal{Z}_{g,n}^{1,i}] = [\mathcal{Z}_{g,n}^{1,j}]^2 \text{ for all } i > 3 \text{ and } j \neq 1, \\ & b_g[\mathcal{Z}_{g,n}^{i,j}]^2 \text{ for all } i \neq j \end{aligned}$$

where b_g is a positive integer dividing $(4g+2)(g+1)$.

Notice that we have some bounds on the order of $[\mathcal{Z}_{g,n}^{2,3}]^2$, but we do not manage to find the order exactly. Notice that $[\mathcal{Z}_{g,n}^{2,3}]^2 = -[\mathcal{Z}_{g,n}^{1,4}] \cdot [\mathcal{Z}_{g,n}^{2,3}]$, for $n \geq 4$, so in that case we are dealing with a proper intersection. It is possible that the implementation of some techniques such as higher Chow groups can be used to compute it, but it does not seem immediate. This technique is complicated to use as the first higher Chow group of $\mathcal{H}_{g,n}^{\text{far}}$ is non-zero.

The case of $2g+3$ is less well behaved. The Chow ring of $\mathcal{H}_{g,n}$ is very different from those for $n \leq 2g+2$, since this stack is now a scheme, thus the Chow ring has to be 0 in degree greater than its dimension.

Theorem 0.8. *Let $\text{char } k > 2g$ and $n = 2g+3$. Then, $\text{CH}^*(\mathcal{H}_{g,2g+3})$ is generated by the classes of $\mathcal{W}_{g,2g+3}^1$, $\mathcal{Z}_{g,2g+3}^{1,i}$ for $1 < i \leq 2g+3$ and $\mathcal{Z}_{g,2g+3}^{2,3}$, and the ideal of*

relations is generated by

$$\begin{aligned}
& (8g+4)[W_{g,2g+3}^1] - (4g+2)(g+1)([Z_{g,2g+3}^{1,2}] + [Z_{g,2g+3}^{1,3}] - [Z_{g,2g+3}^{2,3}]), \\
& [Z_{g,2g+3}^{1,i}] \cdot [Z_{g,2g+3}^{1,j}] \text{ for } 1 < i < j, \quad [Z_{g,2g+3}^{1,j}] \cdot [Z_{g,2g+3}^{2,3}] \text{ for } j = 2, 3, \\
& [Z_{g,2g+3}^{1,j}] \cdot [W_{g,2g+3}^1] \text{ for } j \neq 1, \quad [Z_{g,2g+3}^{2,3}] \cdot [W_{g,2g+3}^1] + (g+1)[Z_{g,2g+3}^{2,3}]^2, \\
& 2[W_{g,2g+3}^1]^2 - (g+1)[Z_{g,2g+3}^{2,3}]^2, \\
& [Z_{g,2g+3}^{2,3}]^2 = -[Z_{g,2g+3}^{2,3}] \cdot [Z_{g,2g+3}^{1,i}] = [Z_{g,2g+3}^{1,j}]^2 \text{ for all } i > 3 \text{ and } j \neq 1, \\
& b_g [Z_{g,n}^{i,j}]^2 \text{ for all } i \neq j, \quad [W_{g,2g+3}^1]^{c_g}
\end{aligned}$$

where b_g is a positive integer dividing $(4g+2)(g+1)$, and $1 < c_g \leq 2g+2$.

The only other thing not known in this case is the multiplicative order of $[W_{g,2g+2}^1]$. This should be easier to be found than the additive order of $[Z_{g,2g+3}^{1,2}]^2$.

Structure of the paper. In the first Section we recall some results obtained in [Lan23], which are essential for the rest of the paper. In the second and third Section we compute the Chow rings of $\mathcal{H}_{g,1}$ and $\mathcal{H}_{g,2}^{\text{far}}$, respectively. Next, in Section 4, we show that for $1 \leq n \leq 2g+3$ the Chow ring of $\mathcal{H}_{g,n}$ is generated in degree 1 and we compute some relations. These are enough to compute the Chow ring of $\mathcal{H}_{g,2}$, as shown in Section 5. In Section 6, we compute $\mathcal{H}_{g,n}^{\text{far}}$ for $3 \leq n \leq 2g+2$, and we conclude the computation of $\mathcal{H}_{g,n}$ for $n \leq 2g+3$, except for the additive order of the degree 2 torsion class $[Z_{g,n}^{1,2}]^2$, and the multiplicative order of $[W_{g,2g+3}^1]$ in the case $n = 2g+3$.

Acknowledgements. I would like to thank Angelo Vistoli for the valuable time spent discussing ideas on this and related subjects. I am also grateful to Dan Edidin and Zhengning Hu for the useful discussions we had. I wish to warmly thank Michele Pernice for his support and taking the time to talk about this work.

1. PRELIMINARIES AND THE GEOMETRY OF $\mathcal{H}_{g,n}$

We will use the description of $\mathcal{H}_{g,n}$ obtained in [Lan23], where the author computed the Picard group of $\mathcal{H}_{g,n}$ for all g and n . We recall the description and results here. We will always assume that the characteristic of the base field does not divide $2g+2$.

1.1. Description of $\mathcal{H}_{g,n}$.

Construction. Let

$$\mathcal{C}_g \xrightarrow{f} \mathcal{P}_g \xrightarrow{\pi} \mathcal{H}_g$$

be the universal family of n -hyperelliptic curves, ramifying over the ‘universal’ Brauer-Severi stack $\mathcal{P}_g \rightarrow \mathcal{H}_g$, and let α be the universal hyperelliptic involution on \mathcal{C}_g . Let \mathcal{C}_g^n be the fiber product of n copies of \mathcal{C}_g over \mathcal{H}_g , and $\Delta_{i,j}$ be the i, j -th diagonal. Notice that $\mathcal{H}_{g,n}$ is the open substack of \mathcal{C}_g^n given by the complement of the extended diagonal. For all $i \neq j$, define $\mathcal{Z}_{g,n}^{i,j} \subset \mathcal{H}_{g,n}$ as the restriction of $\alpha_i^{-1}(\Delta_{i,j})$ to $\mathcal{H}_{g,n}$, which is a closed substack of $\mathcal{H}_{g,n}$. Here, $\alpha_i : \mathcal{C}_g^n \rightarrow \mathcal{C}_g^n$ is the involution which acts as α on the i -th component, and as the identity on the others. Finally, define $\mathcal{H}_{g,n}^{\text{far}}$ as the complement of the $\mathcal{Z}_{g,n}^{i,j}$ ’s in $\mathcal{H}_{g,n}$.

Denote with

$$\mathcal{C}_{g,n} \xrightarrow{f} \mathcal{P}_{g,n} \xrightarrow{\pi} \mathcal{H}_{g,n}$$

the universal family of pointed hyperelliptic curves over $\mathcal{H}_{g,n}$, which ramifies over $\mathcal{P}_{g,n}$. Consider the universal i -th section $\tilde{\sigma}_i : \mathcal{H}_{g,n} \rightarrow \mathcal{C}_{g,n}$, which induces a section $\sigma_i = f \circ \tilde{\sigma}_i$ of $\mathcal{P}_{g,n} \rightarrow \mathcal{H}_{g,n}$. Then, $\mathcal{Z}_{g,n}^{i,j}$ is exactly the locus where $\sigma_i = \sigma_j$, i.e. $\tilde{\sigma}_i$ and $\tilde{\sigma}_j$ differ by the hyperelliptic involution.

Remark 1.1. Recall that $\mathcal{H}_{g,n}$ is integral and smooth for all $g \geq 2$ and all n . From the construction it follows that $\mathcal{Z}_{g,n}^{i,j}$ is integral and smooth for all $i \neq j$, and that $\mathcal{Z}_{g,n}^{i,j} = \mathcal{Z}_{g,n}^{j,i}$. For this reason, we usually assume $i < j$. Moreover, the intersection of $\mathcal{Z}_{g,n}^{i,j}$ and $\mathcal{Z}_{g,n}^{l,m}$ is empty if and only if $\{i, j\}$ and $\{l, m\}$ intersect properly. See [Lan23, Subsection 2.1] for the details in a more general setting.

The $\mathcal{Z}_{g,n}^{i,j}$ define Cartier divisors on $\mathcal{H}_{g,n}$, whose classes are subject to the following relations.

Proposition 1.2. *Let $1 \neq i < j$. Then,*

$$[\mathcal{Z}_{g,n}^{i,j}] = [\mathcal{Z}_{g,n}^{1,i}] + [\mathcal{Z}_{g,n}^{1,j}] + [\mathcal{Z}_{g,n}^{2,3}] - [\mathcal{Z}_{g,n}^{1,2}] - [\mathcal{Z}_{g,n}^{1,3}].$$

Moreover, there are no other relations between those divisors.

Proof. See [Lan23, Corollary 2.7 and Proposition 5.2]. □

Now, we study the open substack $\mathcal{H}_{g,n}^{\text{far}} \subset \mathcal{H}_{g,n}$. For $n \leq 2g+3$, this is isomorphic to the quotient of an open subscheme of a representation of a linear algebraic group by the same group. We report here the precise statements. See [Lan23, Subsections 2.2, 2.3 and 2.4] for details.

Proposition 1.3 ([AV04, Theorem 4.1]). *It holds that*

$$\mathcal{H}_g \simeq \left[\frac{\mathbb{A}^{2g+3} \setminus \Delta_g}{\text{GL}_2/\mu_{g+1}} \right]$$

where Δ_g is the locus of singular forms and a matrix A acts on a form f of degree $2g+2$ as

$$A \cdot f(x, y) = f(A^{-1}(x, y)).$$

Remark 1.4. We use the following notation. Let $3 \leq n \leq 2g+3$ and take a point

$$(a_0, \dots, a_{2g+3-n}, s_1, s_2, s_3, t_1, \dots, t_{n-3}, p_1, \dots, p_{n-3}) \in \mathbb{A}^{2g+3} \times (\mathbb{A}^{n-3} \setminus \tilde{\Delta})$$

where $\tilde{\Delta}$ is the locus where p_1, \dots, p_{n-3} are not mutually distinct and different from 0 and 1. Then, we think at the point as the data of a polynomial

$$f(x, y) = a_0 x^{2g+2} + a_1 x^{2g+1} y + \dots + a_{2g+2} y^{2g+2},$$

where there is no abuse of notation regarding the a_i 's, such that

$$f(0, 1) = s_1^2, \quad f(1, 0) = s_2^2, \quad f(1, 1) = s_3^2, \quad f(1, p_i) = t_i^2 \text{ for all } i.$$

The polynomial f is clearly uniquely determined, see also [Lan23, Lemma 2.20]. We use a similar convention for $n \leq 3$.

Proposition 1.5 ([Per22, Proposition 1.6], [Lan23, Corollary 2.17]). *It holds that*

$$\mathcal{H}_{g,1} \simeq \left[\frac{\mathbb{A}^{2g+3} \setminus \Delta_{g,1}}{\mathbb{B}_2/\mu_{g+1}} \right]$$

where $\Delta_{g,1}$ is the locus corresponding to singular forms, and a lower triangular matrix $A = \begin{bmatrix} a & 0 \\ b & c \end{bmatrix}$ acts on (f, s_1) as

$$A \cdot (f, s_1) = (f(A^{-1}), c^{-(g+1)} s_1).$$

Similarly,

$$\mathcal{H}_{g,2}^{far} \simeq \left[\frac{\mathbb{A}^{2g+3} \setminus \Delta_{g,2}}{(\mathbb{G}_m \times \mathbb{G}_m)/\mu_{g+1}} \right]$$

where where $\Delta_{g,2}$ is the locus corresponding to singular forms, and $A = (a, c)$ acts on (f, s_1, s_2) as

$$A \cdot (f, s_1, s_2) = (f(A^{-1}), c^{-(g+1)} s_1, a^{-(g+1)} s_2).$$

Proposition 1.6 ([Lan23, Proposition 2.21]). *Let $3 \leq n \leq 2g + 3$. Then*

$$\mathcal{H}_{g,n}^{far} \simeq (\mathcal{P}(2^{2g+3-n}, 1^n) \times (\mathbb{A}^{n-3} \setminus \tilde{\Delta})) \setminus \overline{\Delta}_{g,n}$$

where $\mathcal{P}(2^{2g+3-n}, 1^n)$ is the weighted projective stack where the first $2g + 3 - n$ variables have degree 2, while the last n have degree 1, and $\overline{\Delta}_{g,n}$ is the locus corresponding to singular forms. In particular, for $n = 2g + 3$,

$$\mathcal{H}_{g,2g+3}^{far} \simeq (\mathbb{P}^{2g+2} \times (\mathbb{A}^{2g} \setminus \tilde{\Delta})) \setminus \overline{\Delta}_{g,2g+3}$$

and it is a scheme.

We will use these presentations to compute their Chow ring. We conclude the subsection defining one last important divisor of $\mathcal{H}_{g,n}$.

Construction. Let $n \geq 1$. For all $1 \leq i \leq n$, define $\mathcal{W}_{g,n}^i$ to be the divisor of $\mathcal{H}_{g,n}$ given by the points whose images under $\tilde{\sigma}_i$ are in the ramification locus of the uniform cyclic covers parametrized by those points. Rigorously, it is defined as follows. Recall that \mathcal{C}_g^n is the n -fold fibred product of \mathcal{C}_g over \mathcal{H}_g . The ramification locus of the map $\mathcal{C}_g \rightarrow \mathcal{P}_g$ is a closed substack

$$\mathcal{W}_{g,1}^1 \subset \mathcal{C}_g = \mathcal{H}_{g,1}$$

which is étale of degree $2g + 2$ over \mathcal{H}_g . Then, for $n \geq 2$, define

$$\mathcal{W}_{g,n}^1 := (\mathcal{W}_{g,1}^1 \times_{\mathcal{H}_g} \mathcal{C}_g^{n-1}) \cap \mathcal{H}_{g,n} \subset \mathcal{H}_{g,n} \subset \mathcal{C}_g^n.$$

For every other i the definition of $\mathcal{W}_{g,n}^i$ is analogous; alternatively, one can consider the automorphism of $\mathcal{H}_{g,n}$ which exchanges the i -th section with the first.

1.2. The Picard group and classes of divisors. For our computation it will be crucial to know the Picard group of $\mathcal{H}_{g,n}$, which has been computed in [Lan23] in a more general setting. We report the results here.

Theorem 1.7 ([Lan23, Corollary 5.14]). *Suppose that the ground field k is of characteristic not dividing $2g + 2$.*

- *If g is odd, then $\text{Pic}(\mathcal{H}_{g,n}) \simeq \mathbb{Z}^n \oplus \mathbb{Z}/(8g + 4)\mathbb{Z}$.*
- *If g is even, then $\text{Pic}(\mathcal{H}_{g,n}) \simeq \mathbb{Z}^n \oplus \mathbb{Z}/(4g + 2)\mathbb{Z}$.*

Moreover, the torsion part always comes from \mathcal{H}_g .

For $n \geq 3$, it is easy to describe the generators of the Picard group.

Proposition 1.8 ([Lan23, Proposition 5.16]). *Let $n \geq 3$ and assume that the characteristic of k does not divide $2g + 2$. Then*

$$\{[\mathcal{Z}_{g,n}^{1,j}]\}_{2 \leq j \leq n} \cup \{[\mathcal{Z}_{g,n}^{2,3}]\} \cup \{[\mathcal{W}_{g,n}^1]\}$$

is a minimal set of generators of $\text{Pic}(\mathcal{H}_{g,n})$. Moreover,

$$(1) \quad (8g + 4)[\mathcal{W}_{g,n}^1] = (4g + 2)(g + 1)([\mathcal{Z}_{g,n}^{1,2}] + [\mathcal{Z}_{g,n}^{1,3}] - [\mathcal{Z}_{g,n}^{2,3}])$$

is the only relation, regardless of the parity of g .

To address the case of $n \leq 2$ it is important to compute the classes of $\mathcal{W}_{g,n}^1$ and ψ_1 in terms of a fixed base of $\text{Pic}(\mathcal{H}_{g,1})$. Recall that the psi class ψ_i is defined as the pullback of the relative differentials of the universal family under the i -th section. We use the conventions of Theorem 3.2 of [Lan23].

Lemma 1.9 ([Lan23, Lemma 5.7]). *Using the conventions of [Lan23, Theorem 3.2], if g is odd then $[\mathcal{W}_{g,1}^1]$ in $\text{Pic}(\mathcal{H}_{g,1})$ corresponds to*

$$(-(g + 1)/2, 1) \in \mathbb{Z} \oplus \mathbb{Z}/(8g + 4)\mathbb{Z}$$

while if g is even then it corresponds to

$$(-(g + 1), g/2 + 1) \in \mathbb{Z} \oplus \mathbb{Z}/(4g + 2)\mathbb{Z}.$$

The same holds for $[\mathcal{W}_{g,2}^1]$ in $\text{Pic}(\mathcal{H}_{g,2}^{far})$.

Similarly, if g is odd then $[\mathcal{W}_{g,2}^2]$ in $\text{Pic}(\mathcal{H}_{g,2}^{far})$ corresponds to

$$((g + 1)/2, 1) \in \mathbb{Z} \oplus \mathbb{Z}/(8g + 4)\mathbb{Z}$$

while if g is even then it corresponds to

$$(g + 1, -g/2) \in \mathbb{Z} \oplus \mathbb{Z}/(4g + 2)\mathbb{Z}.$$

Proposition 1.10. *The first psi class ψ_1 in $\text{Pic}(\mathcal{H}_{g,1})$ corresponds to*

$$(-(g - 1)/2, 1) \in \mathbb{Z} \oplus \mathbb{Z}/(8g + 4)\mathbb{Z}$$

if g is odd, while if g is even it corresponds to

$$(-(g - 1), g/2) \in \mathbb{Z} \oplus \mathbb{Z}/(4g + 2)\mathbb{Z}.$$

Proof. Consider the linear group $B_2 \times \mathbb{G}_m$ and its action on $\mathbb{A}^{2g+2} \times \mathbb{A}^1 \times \mathbb{A}^3$ described as follows. Let $B \in B_2$ and $\lambda \in \mathbb{G}_m$, and consider $(a_0, \dots, a_{2g+1}, s_1, x_0, x_1, y)$ an element of $\mathbb{A}^{2g+2} \times \mathbb{A}^1 \times \mathbb{A}^3$, which we also denote by (f, s_1, x_0, x_1, y) with the usual convention. Then the action is defined by

$$(B, \lambda) \cdot (f, s_1, x_0, x_1, y) = (\lambda^{-2} f \circ B^{-1}, \lambda^{-1} c^{-(g+1)} s_1, B(x_0, x_1), \lambda^{-1} y)$$

where

$$B = \begin{bmatrix} a & 0 \\ b & c \end{bmatrix}$$

and $B(x_0, x_1) = (ax_0, bx_0 + cx_1)$. Notice that we have homomorphisms

$$B_2/\mu_{g+1} \longrightarrow B_2 \times \mathbb{G}_m \longrightarrow B_2/\mu_{g+1}$$

sending $[A] = \begin{bmatrix} a & 0 \\ b & c \end{bmatrix}$ to $(c^{-1}A, c^{g+1})$, which is easily seen to be well defined, and sending (B, λ) to the class of $\lambda^{1/(g+1)}B$. Notice that the composite is just the identity. Moreover, we can consider the morphism

$$\mathbb{A}^{2g+2} \times \mathbb{A}^1 \times \mathbb{A}^3 \longrightarrow \mathbb{A}^{2g+2} \times \mathbb{A}^1$$

simply given by the projection, which admits the obvious section

$$\mathbb{A}^{2g+2} \times \mathbb{A}^1 \longrightarrow \mathbb{A}^{2g+2} \times \mathbb{A}^1 \times \mathbb{A}^3$$

sending (f, s_1) to $(f, s_1, 0, 1, s_1)$. Notice that both maps are equivariant with respect to the action of the two linear groups and the homomorphisms between them. Moreover, this last section has image in the closed subscheme X' of $\mathbb{A}^{2g+2} \times \mathbb{A}^1 \times \mathbb{A}^3$ defined by the equation $f(x_0, x_1) = y^2$. Notice that this locus is $(\mathbb{B}_2 \times \mathbb{G}_m)$ -invariant. Excising the locus of singular polynomials, we get an open invariant subscheme X of X' . It is not hard to show that the induced morphism

$$\varphi : \left[\frac{X}{\mathbb{B}_2 \times \mathbb{G}_m} \right] \longrightarrow \mathcal{H}_{g,1}$$

is the universal family over $\mathcal{H}_{g,1}$. We can use this construction to compute the class of ψ_1 in $\text{Pic}(\mathcal{H}_{g,1})$.

Notice that, using the exact sequence of differentials, the module of relative differentials of the morphism φ seen at the level of schemes, i.e. without passing to the quotient, is given by

$$\frac{dx_0 \oplus dx_1 \oplus dy}{\langle f_{x_0} dx_0 + f_{x_1} dx_1 - 2y dy \rangle}.$$

Restricting to the section, we get

$$\frac{dx_0 \oplus dx_1 \oplus dy}{\langle a_{2g+1} dx_0 + (2g+2)s_1^2 dx_1 - 2s_1 dy \rangle}$$

which is a rank 2 vector bundle. Take the determinant, which is generated by $dx_0 \wedge dy$ over $s_1 \neq 0$, and by $dx_1 \wedge dy$ over $a_{2g+1} \neq 0$. Notice that, when both s_1 and a_{2g+1} are non-zero,

$$\frac{1}{s_1} dx_0 \wedge dy = \frac{2g+2}{a_{2g+1}} dx_1 \wedge dy.$$

This defines a non-vanishing regular section (recall that the characteristic of the base field does not divide $2g+2$), hence a trivialization of the pullback of the dual sheaf. Now, $(dx_0 \wedge dy)/s_1^2$ is an invariant section generating the line bundle. Notice that the action of \mathbb{B}_2/μ_{g+1} on it is via $a^{-1}c^{-g}$. We conclude by applying the linear base changes as in [Lan23, Theorem 3.2]. \square

Together, the two results above yield the following Proposition.

Proposition 1.11. *The classes ψ_1 and $[\mathcal{W}_{g,1}^1]$ form a minimal set of generators of $\text{Pic}(\mathcal{H}_{g,1})$, and*

$$(4g+2)(g+1)\psi_1 = (4g+2)(g-1)[\mathcal{W}_{g,1}^1]$$

is the only relation. Adjoining $[\mathcal{Z}_{g,2}^{1,2}]$ one gets a minimal set of generators of $\text{Pic}(\mathcal{H}_{g,2})$, with the same relation.

2. THE CHOW RING OF $\mathcal{H}_{g,1}$

In this Section we extend and correct the ideas of Pernice in [Per22] to compute the Chow ring of $\mathcal{H}_{g,1}$. The only difference in the result is the relation in degree 2, which essentially amounts to ‘half’ of the one of Pernice in the even genus case, and to ‘one fourth’ in the odd genus case.

We begin by recalling some facts about Chow envelopes. For the proofs and more details see [EF09] and [Per22]; for the general theory on equivariant Chow rings, see [EG98].

Definition 2.1. Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a proper, representable morphism of quotient stacks. We say that f is Chow-surjective if the morphism of groups

$$f_* : \mathrm{CH}_*(X) \longrightarrow \mathrm{CH}_*(Y)$$

is surjective.

We say that a morphism of algebraic stacks $f : \mathcal{X} \rightarrow \mathcal{Y}$ is a Chow envelope if $f(K) : \mathcal{X}(K) \rightarrow \mathcal{Y}(K)$ is essentially surjective for every extension of fields K/k .

Lemma 2.2 ([Per22, Lemma 3.3]). *Let G be a group scheme over k and suppose we have an action on two algebraic spaces X and Y and a map $f : X \rightarrow Y$ which is G -equivariant. We denote by f^G the induced morphism of quotient stacks and we assume it is proper and representable. If f^G is a Chow envelope, then f^G is Chow-surjective.*

Remark 2.3. If T is a special group acting on an algebraic space X , then the T -torsor $X \rightarrow [X/T]$ is a Chow envelope.

Proposition 2.4 ([Per22, Corollary 3.6]). *Let G, T be two groups schemes over k and suppose we have an action of $G \times T$ on two algebraic spaces X and Y and a map $f : X \rightarrow Y$ which is $G \times T$ -equivariant. Assume that T is a special group. Suppose that f^G is a Chow envelope and $f^{G \times T}$ is a proper representable morphism. Then $f^{G \times T} : [X/(G \times T)] \rightarrow [Y/(G \times T)]$ is Chow-surjective.*

We set $N = 2g + 2$. Thanks to B_2/μ_{g+1} being an extension of the maximal torus and a unipotent subgroup, using Proposition 1.5 we reduce to the computation of the $(\mathbb{G}_m \times \mathbb{G}_m)/\mu_{g+1}$ -equivariant Chow ring of $(\mathbb{A}^{2g+2} \times \mathbb{A}^1) \setminus \Delta_{g,1}$.

Notice that there are isomorphisms

$$(2) \quad \begin{array}{ccc} (\mathbb{G}_m \times \mathbb{G}_m)/\mu_{g+1} & \xrightarrow{\varphi} & \mathbb{G}_m \times \mathbb{G}_m \xrightarrow{\varphi_{\text{even}}} \mathbb{G}_m \times \mathbb{G}_m \\ & & \downarrow \varphi_{\text{odd}} \\ & & \mathbb{G}_m \times \mathbb{G}_m \end{array}$$

where

$$\varphi(a, c) = (ac^{-1}, c^{g+1}), \quad \varphi_{\text{even}}(\lambda, \mu) = (\lambda^{g/2+1}\mu, \lambda^{g/2}\mu), \quad \varphi_{\text{odd}}(\lambda, \mu) = (\lambda^{(g+1)/2}\mu, \lambda).$$

The two composites are the isomorphisms induced by those in [AV04, Proposition 4.4] and [Per22, Proposition 1.7], depending on the parity of the genus. Using these last identifications, for both B_2/μ_{g+1} and GL_2/μ_{g+1} , the map

$$\mathbb{A}^{2g+3} \setminus \Delta_{g,1} \ni (f, s_1) \longrightarrow f \in \mathbb{A}^{2g+3} \setminus \Delta_g$$

inducing

$$\mathcal{H}_{g,1} \longrightarrow \mathcal{H}_g$$

obtained by forgetting the section is equivariant in a natural way. We tacitly consider these identifications, depending on the parity of the genus.

Since we want to use the theory of Chow envelopes, we pass to the projective setting. To do so, we consider the quotient $(\mathbb{A}^{2g+2} \times \mathbb{A}^1) \setminus 0$ by the action of \mathbb{G}_m with weights $(2, \dots, 2, 1)$, and let $\overline{\Delta}_{g,1}$ be the image of $\Delta_{g,1}$ under the quotient. Notice that using the isomorphisms in (2), we have to compute the quotient of the $(\mathbb{G}_m \times \mathbb{G}_m)$ -equivariant Chow ring of $\mathcal{P}(2^N, 1)$ by the image of the $(\mathbb{G}_m \times \mathbb{G}_m)$ -equivariant Chow ring of $\overline{\Delta}_{g,1}$ under the inclusion. Then, one recovers the Chow ring of $\mathcal{H}_{g,1}$ simply by substituting the first Chern class ξ of the tautological bundle of $\mathcal{P}(2^N, 1)$ with an expression in the obvious characters t_1 and t_2 of $\mathbb{G}_m \times \mathbb{G}_m$ (the expression depends on the parity of g).

Remark 2.5. Using the isomorphism φ in (2), the action of $\mathbb{G}_m \times \mathbb{G}_m$ on the space $(\mathbb{A}^{2g+2} \times \mathbb{A}^1) \setminus \Delta_{g,1}$ is with weights $(-(2g+2), \dots, -1, 0)$ and $(-2, \dots, -2, -1)$, respectively. Therefore, one could simply study the quotient of the \mathbb{G}_m -equivariant Chow ring of $\mathcal{P}(2^N, 1)$ by the image of the \mathbb{G}_m -equivariant Chow ring of $\overline{\Delta}_{g,1}$ under the inclusion, where \mathbb{G}_m is the first component the product group scheme. Then, the first Chern class ξ of the tautological bundle of $\mathcal{P}(2^N, 1)$ would correspond to the character $-t_2$ of the second factor in $\mathbb{G}_m \times \mathbb{G}_m$. However, in order to simplify the comparison between the Chow rings of $\mathcal{H}_{g,1}$ and \mathcal{H}_g , it is easier to consider $(\mathbb{G}_m \times \mathbb{G}_m)$ -equivariant Chow rings as explained above.

The dependence on the parity of the genus is very annoying, but we can ignore it in some sense, thanks to the following Remark.

Remark 2.6. We will follow and correct the strategy used by Pernice in [Per22], which is based on comparing the Chow ring of $\mathcal{H}_{g,1}$ with the one of \mathcal{H}_g . Recall that, when g is even, GL_2/μ_{g+1} is isomorphic to GL_2 , while in the odd case it is isomorphic to $\mathbb{G}_m \times \mathrm{PGL}_2$. The first is special, hence when computing $\mathrm{CH}^*(\mathcal{H}_g)$ one can reduce to the maximal torus T , while this cannot be done in the odd case, as PGL_2 is not special. Hence the strategy for the computation of the Chow ring of \mathcal{H}_g depends on the parity of g , see [EF09] and [DL18]. However, using the isomorphism φ in (2), we get that the presentation of the T -equivariant Chow ring of $\mathbb{A}^{2g+3} \setminus \Delta_g$ (see Proposition 1.3) does not depend on the parity of g . Indeed, one can notice that ‘half’ of the generating degree 2 relation in [DL18] is a relation in the T -equivariant Chow ring. Thus, the map obtained by reducing to the maximal torus is not injective when g is odd, as expected. This observation allow us to ignore the parity of the genus. For $\mathrm{CH}^*(\mathcal{H}_{g,1})$, we will show that in degree 2 the generating relation is ‘half’ of the one in the even case, hence a ‘fourth’ of the one in the odd case.

We will use a slightly different notation from the one in [Per22]. Recall that Pernice has constructed maps

$$a_r : \mathbb{P}^r \times \mathcal{P}(2^{N-2r}, 1) \longrightarrow \mathcal{P}(2^N, 1)$$

$$b_r : \mathbb{P}^{r-1} \times \mathcal{P}(2^{N-2r}) \longrightarrow \mathcal{P}(2^N, 1)$$

that factor through $\Delta_{g,1}$, and he proved that a_r and b_r together give a Chow envelope of $\overline{\Delta}_{g,1}$; see [Per22, Proposition 3.12]. The first map sends $(f, (g, s_1))$ to $(f^2g, f(0, 1)s_1)$, where f is a polynomial of degree $r+1$ and g a polynomial of

degree $N - 2r + 1$ such that $g(0, 1) = s_1^2$, while the second sends (f, g) to $(x^2 f^2 g, 0)$, with a similar notation.

Moreover, he has shown that the image of the pushforward under each of those maps is generated by $\alpha_{r,i} := (a_r)_* \xi_r^i$ and $\beta_{r,i} := (b_r)_* \xi_{r-1}^i$, respectively, with i varying. In the first case, ξ_r is the pullback to $\mathbb{P}^r \times \mathcal{P}(2^{N-2r}, 1)$ of the first Chern class of $\mathcal{O}_{\mathbb{P}^r}(1)$, while in the second case ξ_{r-1} is the pullback to $\mathbb{P}^{r-1} \times \mathcal{P}(2^{N-2r+1})$ of the first Chern class of $\mathcal{O}_{\mathbb{P}^{r-1}}(1)$.

2.1. Problems with the previous proof. The main problem of the computation in [Per22] is the following. The idea in that article is to consider cartesian diagrams

$$\begin{array}{ccc} X & \longrightarrow & \mathcal{P}(2^N, 1) \\ \downarrow \psi & & \downarrow \phi \\ \mathbb{P}^r \times \mathbb{P}^{N-2r} & \xrightarrow{\pi_r} & \mathbb{P}^N \end{array}$$

and

$$\begin{array}{ccc} Y & \longrightarrow & \mathcal{P}(2^N, 1) \\ \downarrow \psi' & & \downarrow \phi \\ \mathbb{P}^{r-1} \times \mathbb{P}^{N-2r} & \xrightarrow{\pi'_r} & \mathbb{P}^N \end{array}$$

where the first bottom horizontal map is part of a $(\mathbb{G}_m \times \mathbb{G}_m)$ -Chow envelope of Δ_g , see [EF09, Proposition 4.1], sending (f, g) to $f^2 g$, while the second bottom horizontal map sends (f, g) to $x_0^2 f^2 g$. Then one considers the induced map from $\mathbb{P}^r \times \mathcal{P}(2^{N-2r}, 1)$ to X , which factors a_r , and the one from $\mathbb{P}^{r-1} \times \mathcal{P}(2^{N-2r})$ to Y , which factors b_r . These maps are again proper and representable. One then shows that the pushforward of 1 under those maps is again 1, and uses compatibility of proper pushforward and flat pullback of [Vis98] to conclude that everything comes from ϕ^* , if 1 is in the image of the pullback under ψ' . While it is true that ϕ is flat, see Proposition 2.7, the pullback of 1 under ψ' is not necessarily 1. Indeed, for $r = 1$, we will see that Y is irreducible but not smooth, not even generically reduced. Precisely, the fundamental class $[Y]$ in $\mathrm{CH}_*(Y)$ corresponds to 2. Moreover, notice that the behaviour of ϕ is very similar to the root stack $\sqrt[r]{(\mathcal{O}(1), a_{2g+2})/\mathbb{P}^N}$ over \mathbb{P}^N . Indeed, over the locus where $a_{2g+2} \neq 0$, the map ϕ is an isomorphism, while over the zero locus it is ramified of order 2, and the automorphism groups are the same to those of the root stack. We show that they are indeed equal; this proves also the above facts. The result is slightly more general.

Proposition 2.7. *Let N and r be positive integers. The morphism*

$$\phi : \mathcal{P}(r^N, 1) \longrightarrow \mathbb{P}^N$$

identifies $\mathcal{P}(r^N, 1)$ with the r -th root stack $\sqrt[r]{(\mathcal{O}(1), a_{2g+2})/\mathbb{P}^N}$ over \mathbb{P}^N . In particular, ϕ is flat.

Proof. The above discussion shows that the map $\phi : \mathcal{P}(r^N, 1) \rightarrow \mathbb{P}^N$ factors as

$$\mathcal{P}(r^N, 1) \xrightarrow{\psi} \sqrt[r]{(\mathcal{O}(1), a_{2g+2})/\mathbb{P}^N} \longrightarrow \mathbb{P}^N$$

and ψ is an isomorphism over the locus where $a_{2g+2} \neq 0$. Moreover, ψ induces an isomorphism between the automorphism groups of geometric points, hence it

is representable. Being also proper and birational between smooth stacks, ψ is an isomorphism. The flatness of ϕ follows from [Cad04, Corollary 2.3.6]. \square

Corollary 2.8. *There is a commutative diagram with cartesian squares*

$$\begin{array}{ccccc}
\mathcal{P}(2^{N-1}) & \xrightarrow{i} & \mathcal{P}(2^{N-1})_\epsilon & \xrightarrow{b'_1} & \mathcal{P}(2^N)_\epsilon & \xrightarrow{b''_1} & \mathcal{P}(2^N, 1) \\
& & \downarrow & & \downarrow & & \downarrow \\
& & \mathcal{P}(2^{N-1}) & \longrightarrow & \mathcal{P}(2^N) & \longrightarrow & \mathcal{P}(2^{N+1}) \\
& & \downarrow & & \downarrow & & \downarrow \\
& & \mathbb{P}^{N-2} & \longrightarrow & \mathbb{P}^{N-1} & \longrightarrow & \mathbb{P}^N.
\end{array}$$

where $b''_1 \circ b'_1 \circ i = b_1$, and $\mathcal{P}(2^N)_\epsilon$ (respectively, $\mathcal{P}(2^{N-1})_\epsilon$) is a first order thickening of $\mathcal{P}(2^N)$ (respectively, of $\mathcal{P}(2^{N-1})$).

In particular,

$$[\mathcal{P}(2^{N-1})_\epsilon] = 2 \in \text{CH}_*(\mathcal{P}(2^{N-1})_\epsilon).$$

Proof. The only thing to be proved is that $\mathcal{P}(2^N)_\epsilon$ is a first order thickening of $\mathcal{P}(2^N)$. Notice that $\mathcal{P}(2^N)_\epsilon \rightarrow \mathbb{P}^{N-1}$ is the locus of the root stack $\mathcal{P}(2^N, 1)$ where the defining section a_{2g+2} vanishes. Recall that when a n -root stack over an algebraic stack \mathcal{X} is defined by $(\mathcal{O}_{\mathcal{X}}, f)$, where f is a global section of \mathcal{X} , the root stack is isomorphic to the quotient stack of

$$\text{Spec}_{\mathcal{X}}(\mathcal{O}_{\mathcal{X}}[T]/(T^n - f))$$

by the action of μ_n given by $\zeta \cdot T = \zeta T$. In particular, when $f = 0$ and $n = 2$, we get

$$[(\text{Spec}_{\mathcal{X}}(\mathcal{O}_{\mathcal{X}}[T]/(T^2)))/\mu_2]$$

which is different from the root gerbe $\sqrt[2]{\mathcal{O}_{\mathcal{X}}/\mathcal{X}}$; see [Ols16, Chapter 10.3] or [Cad04, Section 2] for more on root stacks. In particular, $\mathcal{P}(2^N)_\epsilon$ is Zariski-locally on \mathbb{P}^{N-1} of the form described above. \square

Corollary 2.9. *Consider the commutative diagram with cartesian square*

$$\begin{array}{ccccc}
& & \xrightarrow{a_r} & & \\
& & \searrow & & \nearrow \\
\mathbb{P}^r \times \mathcal{P}(2^{N-2r}, 1) & \xrightarrow{a'_r} & X & \longrightarrow & \mathcal{P}(2^N, 1) \\
& \searrow \theta & \downarrow \psi & & \downarrow \phi \\
& & \mathbb{P}^r \times \mathbb{P}^{N-2r} & \xrightarrow{\pi_r} & \mathbb{P}^N.
\end{array}$$

Then ϕ , θ , a'_r and ψ are isomorphisms over the locus where $a_{2g+2} \neq 0$ (hence $s_1 \neq 0$), with all spaces isomorphic to $\mathbb{A}^r \times \mathbb{A}^{N-2r}$.

2.2. Computation of $\alpha_{r,i}$. First we focus on $\alpha_{r,0}$.

Lemma 2.10. *The classes $\alpha_{r,0}$ are contained in the image of ϕ^* .*

Proof. It follows from Corollary 2.9 that $(a'_r)_* 1 = \psi^* 1 = 1$, and we conclude by compatibility of representable proper pushforward and flat pullback. \square

Now we consider $\alpha_{r,i}$ for $0 < i \leq r$. We have a commutative diagram

$$\begin{array}{ccccc} \mathbb{P}^{r-1} \times \mathcal{P}(2^{N-2r}, 1) & \xrightarrow{i} & \mathbb{P}^r \times \mathcal{P}(2^{N-2r}, 1) & \xrightarrow{a_r} & \mathcal{P}(2^N, 1) \\ & \searrow \theta & & \nearrow b_r & \\ & & \mathbb{P}^{r-1} \times \mathcal{P}(2^{N-2r+1}) & & \end{array}$$

where i is the inclusion corresponding to the hyperplane of points in \mathbb{P}^r whose first coordinate is 0, and θ sends $(f, (g, s_1))$ to (f, g) , with the obvious notation. Notice that the T -equivariant class of that hyperplane is the sum of ξ_r and the first Chern class of a character of T , thanks to Lemma 2.4 of [EF09]. This shows that $\alpha_{r,1}$ is generated by $\alpha_{r,0}$ and the image of $(b_{r-1})_*$. One can argue inductively to obtain the same result for all $\alpha_{r,i}$, proving the following Lemma.

Lemma 2.11. *For $0 < i \leq r$, $\alpha_{r,i}$ is generated by $\alpha_{r,0}$ and the image of $(b_{r-1})_*$.*

This reduces us to the computation of the $\beta_{r,i}$'s.

2.3. Computation of $\beta_{r,i}$. For all $1 < r \leq N/2$, consider the commutative diagram

$$\begin{array}{ccc} \mathbb{P}^{r-1} \times \mathcal{P}(2^{N-2r+1}) & \xrightarrow{b_r} & \mathcal{P}(2^N, 1) \\ b'_r \downarrow & \nearrow b_1 & \\ \mathcal{P}(2^{N-1}) & & \end{array}$$

where b'_r sends (f, g) in f^2g . This shows that it is enough to study $(b_1)_*$.

Now, notice that

$$b_1^* : \frac{\mathbb{Z}[h]}{(2^N h^{N+1})} \simeq \text{CH}^*(\mathcal{P}(2^N, 1)) \longrightarrow \text{CH}^*(\mathcal{P}(2^{N-1})) \simeq \frac{\mathbb{Z}[h]}{(2^{N-1} h^{N-1})}$$

satisfies $b_1^* h = h$, hence it is surjective, and the same is true at the level of $(\mathbb{G}_m \times \mathbb{G}_m)$ -equivariant Chow rings. By the projection formula it follows that the image of $(b_1)_*$ is the ideal generated by $(b_1)_* 1 = \beta_{1,0}$. Therefore we are reduced to study $\beta_{1,0}$.

Lemma 2.12. *It holds that*

$$\beta_{1,0} = (s_1 = 0) \cdot (a_{2g+1} = 0) \in \text{CH}_{\mathbb{G}_m \times \mathbb{G}_m}^2(\mathcal{P}(2^N, 1)).$$

Proof. It holds essentially by definition. \square

Remark 2.13. In particular, this shows that in the setting of Subsection 2.1 indeed $(\psi')^* 1 \neq 1$. Moreover, the defining relation of $\text{CH}_T^*(\mathcal{P}(2^N, 1))$ is contained in the ideal generated by $\beta_{1,0}$. Indeed, the relation is given by the product of the T -equivariant classes of the $N+1$ coordinate hyperplanes, and $\beta_{1,0}$ is the product of two of those.

2.4. Final computation. Putting everything together, we have proved that the relations are generated by those coming from \mathcal{H}_g and

$$\beta_{1,0} = (s_1 = 0) \cdot (a_{2g+1} = 0) \in \text{CH}_{\mathbb{G}_m \times \mathbb{G}_m}^2(\mathcal{P}(2^N, 1)).$$

Computing this last class in terms of the characters of $\mathbb{G}_m \times \mathbb{G}_m$ gives the following Theorem, where we use the notation of Pernice.

Theorem 2.14. *Let $\text{char } k > 2g$. If g is even, then*

$$\text{CH}^*(\mathcal{H}_{g,1}) \simeq \frac{\mathbb{Z}[t_0, t_1]}{((4g+2)(t_0+t_1), \frac{g(g-1)}{2}(t_0^2+t_1^2) - g(g+3)t_0t_1 + (2g+1)(t_0+t_1)t_1)}.$$

If g is odd, then

$$\text{CH}^*(\mathcal{H}_{g,1}) \simeq \frac{\mathbb{Z}[\tau, \rho]}{((8g+4)\tau, 2\tau^2 + \frac{g(g+1)}{2}\rho^2 - (2g+1)\tau\rho)}.$$

Remark 2.15. Notice that, for g even, multiplying the degree 2 relation and using the degree 1 relation we get back the degree 2 relation described in [Per22]. For the g odd case, we have to multiply by 4.

2.5. Base changes. We know that $[\mathcal{W}_{g,1}^1]$ and ψ_1 generate the Chow ring, so we write those classes in terms of the generators t_0 and t_1 , so that we can rewrite the above isomorphisms in terms of the classes we are interested in.

Lemma 2.16. *With the notation of Theorem 2.14, we have the following. If g is even, then*

$$[\mathcal{W}_{g,1}^1] = -\frac{g}{2}t_0 + \left(\frac{g}{2} + 1\right)t_1, \quad \psi_1 = -\left(\frac{g}{2} - 1\right)t_0 + \frac{g}{2}t_1.$$

If g is odd, then

$$[\mathcal{W}_{g,1}^1] = \tau - \frac{g+1}{2}\rho, \quad \psi_1 = \tau - \frac{g-1}{2}\rho.$$

Proof. It is a straightforward computation using our previous results 1.9 and 1.10 on the classes $[\mathcal{W}_{g,n}^1]$ and ψ_1 , respectively, together with [Per22, Proposition 1.7]. \square

Making this change of variables, we get the following result.

Corollary 2.17. *Let $g \geq 2$ and $\text{char } k > 2g$. Then,*

$$\text{CH}^*(\mathcal{H}_{g,1}) \simeq \frac{\mathbb{Z}[\psi_1, [\mathcal{W}_{g,n}^1]]}{((4g+2)((g+1)\psi_1 - (g-1)[\mathcal{W}_{g,1}^1]), [\mathcal{W}_{g,1}^1]^2 + \psi_1 \cdot [\mathcal{W}_{g,1}^1])}$$

regardless of the parity of g .

There is another interesting basis that we can consider. Recall that we had isomorphisms

$$\begin{array}{ccc} (\mathbb{G}_m \times \mathbb{G}_m) / \mu_{g+1} & \xrightarrow{\varphi} & \mathbb{G}_m \times \mathbb{G}_m \xrightarrow{\varphi_{\text{even}}} \mathbb{G}_m \times \mathbb{G}_m \\ & & \downarrow \varphi_{\text{odd}} \\ & & \mathbb{G}_m \times \mathbb{G}_m \end{array}$$

where

$$\varphi(a, c) = (ac^{-1}, c^{g+1}), \quad \varphi_{\text{even}}(\lambda, \mu) = (\lambda^{g/2+1}\mu, \lambda^{g/2}\mu), \quad \varphi_{\text{odd}}(\lambda, \mu) = (\lambda^{(g+1)/2}\mu, \lambda).$$

Let l_1 and l_2 be the obvious characters of the copy of $\mathbb{G}_m \times \mathbb{G}_m$ in the middle. Then, the above diagram induces the following base changes

$$t_0 = \left(\frac{g}{2} + 1\right)l_1 + l_2, \quad t_1 = \frac{g}{2}l_1 + l_2$$

and

$$\tau = \frac{g+1}{2}l_1 + l_2, \quad \rho = l_1$$

and the equalities

$$[\mathcal{W}_{g,1}^1] = l_2, \quad \psi_1 = l_1 + l_2.$$

This allows us to rewrite everything in another symmetric way, which highlights again the fact that the presentation does not depend on the genus, even though when seen abstractly the two rings appear slightly different, in particular the Picard group.

Corollary 2.18. *Using the base changes above, we have*

$$\mathrm{CH}^*(\mathcal{H}_{g,1}) \simeq \frac{\mathbb{Z}[l_1, l_2]}{((4g+2)((g+1)l_1 + 2l_2), 2l_2^2 + l_1l_2)}.$$

3. THE CHOW RING OF $\mathcal{H}_{g,2}^{\mathrm{far}}$

In this section, we compute the Chow ring of $\mathcal{H}_{g,2}^{\mathrm{far}}$. To do so, we follow the strategy used for the computation of the Chow ring of $\mathcal{H}_{g,1}$.

First, recall that

$$\mathcal{H}_{g,2}^{\mathrm{far}} \simeq \left[\frac{\mathbb{A}^{2g+3} \setminus \Delta_{g,2}}{(\mathbb{G}_m \times \mathbb{G}_m) / \mu_{g+1}} \right]$$

where the action has weights

$$\begin{pmatrix} -(2g+1) & \dots & -1 & 0 & -(g+1) \\ -1 & \dots & -(2g+1) & -(g+1) & 0 \end{pmatrix}.$$

Again, we pass to the projective setting by considering the quotient of \mathbb{A}^{2g+3} by the action of \mathbb{G}_m with weights $(2, \dots, 2, 1, 1)$, hence obtaining $\mathcal{P}(2^{N-1}, 1, 1)$ and its closed subscheme $\overline{\Delta}_{g,2}$, which is the image of $\Delta_{g,2}$.

Now, we construct morphisms that together will form a Chow envelope of $\overline{\Delta}_{g,2}$. We use the usual convention where, given an homogeneous form f of degree $2g+2$ in two variables x_0, x_1 , and such that $f(0,1) = s_1^2$ and $f(1,0) = s_2^2$, we denote by $(f, s_1, s_2) \in \mathbb{A}^{2g+3}$ the element $(a_1, \dots, a_{2g+1}, s_1, s_2)$, where a_0, \dots, a_{2g+2} are the coefficients of f . We use a similar convention for couples (f, s_1) satisfying $f(0,1) = s_1^2$ and couples (f, s_2) satisfying $f(1,0) = s_2^2$.

First, we define

$$\tilde{a}_r : \mathbb{A}^{r+1} \times \mathbb{A}^{2g+3-2r} \longrightarrow \mathbb{A}^{2g+3}$$

as $(f, (g, s_1, s_2)) \mapsto (f^2g, f(0,1)s_1, f(1,0)s_2)$, which is clearly well defined. In a similar way, we define

$$\tilde{b}_r : \mathbb{A}^r \times \mathbb{A}^{2g+3-2r} \longrightarrow \mathbb{A}^{2g+3}$$

as $(f, (g, s_2)) \mapsto (x_0^2 f^2 g, 0, f(1,0)s_2)$,

$$\tilde{c}_r : \mathbb{A}^r \times \mathbb{A}^{2g+3-2r} \longrightarrow \mathbb{A}^{2g+3}$$

as $(f, (g, s_1)) \mapsto (x_1^2 f^2 g, f(0,1)s_1, 0)$, and finally

$$\tilde{d}_r : \mathbb{A}^{r-1} \times \mathbb{A}^{2g+3-2r} \longrightarrow \mathbb{A}^{2g+3}$$

as $(f, g) \mapsto (x_0^2 x_1^2 f^2 g, 0, 0)$.

Remark 3.1. It is necessary to consider all these maps in order to have a Chow envelope, see [Per22, Remark 3.8].

Now, we define an action of $G_{g,2} := (\mathbb{G}_m \times \mathbb{G}_m)/\mu_{g+1}$ on the source of the maps just defined so that they become $G_{g,2}$ -equivariant. Then, we pass to the projective setting. For this, we identify $G_{g,2}$ with $\mathbb{G}_m \times \mathbb{G}_m$ via the isomorphism φ of (2), so that we do not have to worry about the parity of g .

Given elements $(a, c) \in \mathbb{G}_m \times \mathbb{G}_m$, the action on the source $\mathbb{A}^{r+1} \times \mathbb{A}^{2g+3-2r}$ of \tilde{a}_r is defined as

$$(a, c) \cdot (f(x, y), (g(x, y), s_1, s_2)) = (f(ax, y), (c^2 g(ax, y), cs_1, a^{(g+1)} cs_2)).$$

The definition of the other actions is analogous. For the sake of completeness, we describe them for \tilde{b}_r , \tilde{c}_r and \tilde{d}_r in this order:

$$\begin{aligned} (a, c) \cdot (f(x, y), (g(x, y), s_2)) &= (f(ax, y), (c^2 g(ax, y), a^{(g+1)} cs_2)) \\ (a, c) \cdot (f(x, y), (g(x, y), s_1)) &= (f(ax, y), (c^2 g(ax, y), cs_1)) \\ (a, c) \cdot (f(x, y), g(x, y)) &= (f(ax, y), c^2 g(ax, y)). \end{aligned}$$

It is a straightforward computation to check that the maps just defined are equivariant with respect to these actions. Notice that the first component of $\mathbb{G}_m \times \mathbb{G}_m$ acts trivially on the first factor of each source.

Now, we pass to the projective setting. To do so, we consider also the standard action of \mathbb{G}_m on the first factor of each source, with weight one, and the morphism of group schemes

$$\mathbb{G}_m \times G_{g,2} \longrightarrow G_{g,2}$$

given by $(\lambda, (a, c)) \mapsto (a, \lambda \cdot c)$.

It is again a straightforward calculation to show that the maps defined induce $(\mathbb{G}_m \times \mathbb{G}_m)$ -equivariant maps

$$\begin{aligned} a_r : \mathbb{P}^r \times \mathbb{P}(2^{2g+1-2r}, 1, 1) &\longrightarrow \mathbb{P}(2^{2g+1}, 1, 1) \\ b_r : \mathbb{P}^{r-1} \times \mathbb{P}(2^{2g+2-2r}, 1) &\longrightarrow \mathbb{P}(2^{2g+1}, 1, 1) \\ c_r : \mathbb{P}^{r-1} \times \mathbb{P}(2^{2g+2-2r}, 1) &\longrightarrow \mathbb{P}(2^{2g+1}, 1, 1) \\ d_r : \mathbb{P}^{r-2} \times \mathbb{P}(2^{2g+3-2r}) &\longrightarrow \mathbb{P}(2^{2g+1}, 1, 1). \end{aligned}$$

Lemma 3.2. *The maps a_r , b_r , c_r and d_r are proper and representable morphisms of quotient stacks.*

Proof. This is an analogue of [Per22, Lemma 3.9], and it can be proved in the same way. \square

It is important to notice that for every r the morphisms a_r , b_r , c_r and d_r factor through $\overline{\Delta}_{g,2}$. To show it, one can construct for each map a commutative diagram as in [Per22, Remark 3.10].

If f is a morphism of algebraic stacks, we denote with $X(f)$ the source of f .

Proposition 3.3. *Suppose $\text{char } k > 2g$. Let ω_r be the co-product morphism of a_r , b_r , c_r and d_r , and define $\omega = \bigsqcup_{r=1}^{g+1} \omega_r$. Then ω is surjective at the level of $G_{g,2}$ -equivariant Chow ring, i.e. the map ω_* from*

$$\bigoplus_{r=1}^{g+1} \left[\text{CH}_{G_{g,2}}^*(X(a_r)) \oplus \text{CH}_{G_{g,2}}^*(X(b_r)) \oplus \text{CH}_{G_{g,2}}^*(X(c_r)) \oplus \text{CH}_{G_{g,2}}^*(X(d_r)) \right]$$

to $\mathrm{CH}^*(\overline{\Delta}_{g,2})$ is surjective.

Proof. It follows by construction, and it is analogous to [Per22, Proposition 3.12]. One has to subdivide the analysis in cases depending on whether an element (h, s_1, s_2) with $h = f^2g$ is such that $f(0,1)$ or $f(1,0)$ are 0 or not. We then have four cases, one for each map. This shows that ω is a Chow envelope of $\overline{\Delta}_{g,2}$, hence the statement follows from Lemma 2.2.

The proof of the following Lemma is standard. \square

Lemma 3.4. *The following holds:*

$$\begin{aligned} \mathrm{CH}_{G_{g,2}}^*(\mathbb{P}^r \times \mathbb{P}(2^{N-1-2r}, 1, 1)) &\simeq \frac{\mathbb{Z}[t_0, t_1, \xi_{r,1}, \xi_{r,2}]}{(p(t_0, t_1, \xi_{r,1}), q(t_0, t_1, \xi_{r,2}))} \\ \mathrm{CH}_{G_{g,2}}^*(\mathbb{P}^{r-1} \times \mathbb{P}(2^{N-2r}, 1)) &\simeq \frac{\mathbb{Z}[t_0, t_1, \xi_{r-1,1}, \xi_{r-1,2}]}{(p'(t_0, t_1, \xi_{r-1,1}), q'(t_0, t_1, \xi_{r-1,2}))} \\ \mathrm{CH}_{G_{g,2}}^*(\mathbb{P}^{r-2} \times \mathbb{P}(2^{N-2r+1})) &\simeq \frac{\mathbb{Z}[t_0, t_1, \xi_{r-2,1}, \xi_{r-2,2}]}{(p''(t_0, t_1, \xi_{r-2,1}), q''(\xi_{r-2,2}))}. \end{aligned}$$

Here, $\xi_{r,1}$ and $\xi_{r,2}$ are the pullback under the first and second projection, respectively, of the first Chern class of the tautological bundle $\mathcal{O}(1)$ of the respective projective stack, while p (respectively p' , p'') is a monic polynomial in the variable $\xi_{r,1}$ (respectively $\xi_{r-1,1}$, $\xi_{r-2,1}$) of degree $r+1$ (respectively r , $r-1$), and similarly for q (respectively q' , q''), which is of degree $N+1-2r$.

Recall also that

$$\mathrm{CH}_{G_{g,2}}^*(\mathbb{P}(2^{N-1}, 1, 1)) \simeq \frac{\mathbb{Z}[t_0, t_1, h]}{(h + p_N(t_0, t_1))(h + p_0(t_0, t_1)) \prod_{i=1}^N (2h + p_i(t_0, t_1))}$$

for some polynomials $p_i(t_0, t_1)$ and $h = c_1(\mathcal{O}(1))$.

Lemma 3.5. *Using the notation above,*

$$\begin{aligned} (a_r)^*(h) &= \xi_{r,1} + \xi_{r,2}, & (b_r)^*(h) &= \xi_{r-1,1} + \xi_{r-1,2}, \\ (c_r)^*(h) &= \xi_{r-1,1} + \xi_{r-1,2}, & (d_r)^*(h) &= \xi_{r-2,1} + \xi_{r-2,2}. \end{aligned}$$

Proof. The proof is the same as the one of [Per22, Lemma 4.2]. \square

Therefore, we are left with computing the classes

$$\begin{aligned} \alpha_{r,i} &:= (a_r)_* \xi_{r,1}^i, & \beta_{r,i} &:= (b_r)_* \xi_{r-1,1}^i, \\ \gamma_{r,i} &:= (c_r)_* \xi_{r-1,1}^i, & \delta_{r,i} &:= (d_r)_* \xi_{r-2,1}^i \end{aligned}$$

in $\mathrm{CH}_{G_{g,2}}^*(\mathbb{P}(2^{N-1}, 1, 1))$.

Lemma 3.6. *The image of $\mathrm{CH}_{G_{g,2}}^*(\overline{\Delta}_{g,2})$ in $\mathrm{CH}_{G_{g,2}}^*(\mathbb{P}(2^{N-1}, 1, 1))$ is generated by*

$$\alpha_{r,0} \text{ for } 1 \leq r \leq N/2, \quad \beta_{1,0}, \quad \gamma_{1,0}, \quad \delta_{2,0}.$$

Proof. Notice that we have a commutative diagram of proper morphisms

$$\begin{array}{ccccc} \mathbb{P}^{r-1} \times \mathcal{P}(2^{N-2r-1}, 1, 1) & \xleftarrow{i} & \mathbb{P}^r \times \mathcal{P}(2^{N-2r-1}, 1, 1) & \xrightarrow{a_r} & \mathcal{P}(2^{N-1}, 1, 1) \\ & \searrow \theta & & \nearrow b_r & \\ & & \mathbb{P}^{r-1} \times \mathcal{P}(2^{N-2r}, 1) & & \end{array}$$

where i is the inclusion of the hyperplane where the first coordinate of \mathbb{P}^r is 0. Reasoning in the same way as in Lemma 2.11, we get that $\alpha_{r,i}$ is generated by $\alpha_{r,0}$ and the image of $(b_{r-1})_*$, for every $1 \leq i \leq r$.

Now, consider the commutative diagram

$$\begin{array}{ccc} \mathbb{P}^{r-1} \times \mathcal{P}(2^{N-2r}, 1) & \xrightarrow{b_r} & \mathcal{P}(2^{N-1}, 1, 1) \\ b'_r \downarrow & \nearrow b_1 & \\ \mathcal{P}(2^{N-2}, 1) & & \end{array}$$

where b'_r sends $(f, (g, s_2))$ to $(f^2g, f(1, 0)s_2)$. This shows that b_r factors through b_1 . Moreover, the pullback homomorphism

$$b_1^* : \frac{\mathbb{Z}[h]}{(2^{N-1}h^{N+1})} \simeq \mathrm{CH}^*(\mathcal{P}(2^{N-1}, 1, 1)) \longrightarrow \mathrm{CH}^*(\mathcal{P}(2^{N-2}, 1)) \simeq \frac{\mathbb{Z}[h]}{(2^{N-2}h^{N-1})}$$

is surjective, and so is the pullback at the level of $(\mathbb{G}_m \times \mathbb{G}_m)$ -equivariant Chow rings. By the projection formula it follows that the image of $(b_1)_*$ is generated by $(b_1)_*1 = \beta_{1,0}$.

A completely analogous argument works for $\gamma_{r,0}$ and $\delta_{r,0}$. The statement follows. \square

Let $\phi : \mathcal{P}(2^{N-1}, 1, 1) \rightarrow \mathcal{P}(2^N, 1)$ be the morphism induced by $s_2 \mapsto s_2^2$. Notice that it is representable and flat. The proof of the following Lemma is straightforward.

Lemma 3.7. *Let $\mathbb{A}^N \subset \mathcal{P}(2^N, 1)$ be the open subscheme where s_1 does not vanish. Then, there is a cartesian diagram*

$$\begin{array}{ccc} \mathbb{A}^N & \longrightarrow & \mathcal{P}(2^{N-1}, 1, 1) \\ \downarrow u & & \downarrow \phi \\ \mathbb{A}^N & \longrightarrow & \mathcal{P}(2^N, 1) \end{array}$$

where $u(a_1, \dots, a_{N-1}, s_2) = (a_1, \dots, a_{N-1}, s_2^2)$.

Lemma 3.8. *The classes $\alpha_{r,0}$ are in the image of ϕ^* for all r .*

Proof. Consider the commutative diagram with cartesian square

$$\begin{array}{ccccc} & & a_r & & \\ & & \curvearrowright & & \\ \mathbb{P}^r \times \mathcal{P}(2^{N-2r-1}, 1, 1) & \xrightarrow{a'_r} & X & \longrightarrow & \mathcal{P}(2^{N-1}, 1, 1) \\ & \searrow & \downarrow \psi & & \downarrow \phi \\ & & \mathbb{P}^r \times \mathcal{P}(2^{N-2r}, 1) & \longrightarrow & \mathcal{P}(2^N, 1). \end{array}$$

Now, the set-theoretic image of a_r is the preimage under ϕ of the image of the bottom morphism. We prove that $(a'_r)_*1 = \psi_*1$. It is enough to show it after base changing the above diagram to an open substack of $\mathcal{P}(2^N, 1)$ intersecting the image of a_r . Choose the complement of $(s_1 = 0) \cup (a_{2g+1} = 0)$. Over that substack

(actually, scheme), we have that a'_r is an isomorphism, thanks to the previous Lemma. Indeed, over $s_1 \neq 0$ we have a diagram with cartesian square

$$\begin{array}{ccccc} \mathbb{A}^r \times \mathbb{A}^{N-2r} & \xrightarrow{a'_r} & X^\circ & \longrightarrow & \mathbb{A}^N \\ & & \downarrow & & \downarrow u \\ & & \mathbb{A}^r \times \mathbb{A}^{N-2r} & \longrightarrow & \mathbb{A}^N \end{array}$$

inducing an isomorphism

$$X^\circ \simeq \text{Spec } k[x_1, \dots, x_r, y_1, \dots, y_{N-2r}, t_N] / (t_N^2 - x_r^2 y_{N-2r}).$$

Where $x_r \neq 0$, i.e. over $a_{2g+1} \neq 0$, a'_r is an isomorphism, and the statement follows. \square

Let $\phi' : \mathcal{P}(2^{N-1}, 1, 1) \rightarrow \mathcal{P}(2^N, 1)$ the arrow induced by $s_1 \mapsto s_1^2$.

Lemma 3.9. *The class $\beta_{1,0}$ is the intersection of the divisors $(a_{2g+1} = 0)$ and $(s_1 = 0)$. Similarly, $\gamma_{1,0}$ is the intersection of the divisors $(s_2 = 0)$ and $(a_1 = 0)$. Moreover, $\delta_{2,0} = \beta_{1,0} \cdot \gamma_{1,0}$. In particular, all these classes are contained in the images of ϕ^* and ϕ'^* .*

Proof. It follows immediately from the definition of the maps. \square

Remark 3.10. As in Remark 2.13, the above Lemma shows also that the defining relation of $\text{CH}_T^*(\mathcal{P}(2^{N-1}, 1, 1))$ is contained in the ideal generated by $\beta_{1,0}$.

Notice that, using the notation of Corollary 2.18, it holds

$$[\mathcal{W}_{g,2}^2] = (g+1)l_1 + l_2, \quad \psi_2 = gl_1 + l_2$$

so that

$$\begin{aligned} [\mathcal{W}_{g,2}^2] \cdot ([\mathcal{W}_{g,2}^2] + \psi_2) &= ((g+1)l_1 + l_2)((2g+1)l_1 + 2l_2) \\ &= (g+1)(2g+1)l_1^2 + 2l_2^2 + (4g+3)l_1l_2 \end{aligned}$$

which is the new relation.

Putting everything together we get the following Proposition.

Proposition 3.11. *Let $\text{char } k > 2g$. Then,*

$$\text{CH}^*(\mathcal{H}_{g,2}^{\text{far}}) \simeq \frac{\mathbb{Z}[l_1, l_2]}{((4g+2)((g+1)l_1 + 2l_2), 2l_2^2 + l_1l_2, (g+1)(2g+1)l_1^2 + (4g+2)l_1l_2)}.$$

It is useful to write it also in terms of $[\mathcal{W}_{g,2}^1]$ and ψ_1 .

Corollary 3.12. *Let $\text{char } k > 2g$. Then, $\text{CH}^*(\mathcal{H}_{g,2}^{\text{far}})$ is generated by ψ_1 and $[\mathcal{W}_{g,2}^1]$, and the ideal of relations is generated by*

$$\begin{aligned} &(4g+2)((g+1)\psi_1 - (g-1)[\mathcal{W}_{g,2}^1]), \\ &[\mathcal{W}_{g,2}^1]^2 + \psi_1 \cdot [\mathcal{W}_{g,2}^1], \quad (2g+1)(g+1)\psi_1^2 + (2g+1)(3g-1)[\mathcal{W}_{g,2}^1]^2. \end{aligned}$$

Proof. Using the computations done for Corollary 2.18, we get that

$$[\mathcal{W}_{g,2}^2] = (g+1)\psi_1 - g[\mathcal{W}_{g,2}^1], \quad \psi_2 = g\psi_1 - (g-1)[\mathcal{W}_{g,2}^1]$$

in $\text{Pic}(\mathcal{H}_{g,2}^{\text{far}})$. The statement follows. \square

4. GENERATORS AND SOME RELATIONS OF THE CHOW RING OF $\mathcal{H}_{g,n}$

We have very explicit generators of the Picard group of $\mathcal{H}_{g,n}$, with evident geometric meaning. Together with our concrete description of $\mathcal{H}_{g,n}$, this should decisively help in the computation of the integral Chow ring $\mathcal{H}_{g,n}$, at least when this is generated in degree 1. We know that this does not always happen, since the (rational) Chow ring of the stack of n -pointed hyperelliptic curves is not always tautological. An example is given in [PG01, Theorem 3], where the authors construct a non-tautological class in $\mathcal{M}_{2,20}$. However, we shall prove that for $1 \leq n \leq 2g + 3$ the Chow ring is generated in degree 1. Therefore, Proposition 1.8 gives us also the generators of the Chow ring, and we are left with computing the intersection of those divisors. Thanks to their geometric nature, this is doable.

4.1. Generators. We show that the generators of the Picard group described in Proposition 1.8 actually generate the Chow ring for $1 \leq n \leq 2g + 3$. We already know this for $n = 1$, see Section 2. We want to use induction on n . To do so, we use the fact that $\mathcal{Z}_{g,n}^{i,j}$ is isomorphic to an open substack of $\mathcal{H}_{g,n-1}$.

Definition 4.1. Let $1 \leq i < j \leq n$ and define $\mathcal{H}_{g,n-1}^{i,j}$ to be the open substack of $\mathcal{H}_{g,n-1}$ where the map

$$\mathcal{H}_{g,n-1} \dashrightarrow \mathcal{H}_{g,n}$$

induced by

$$(C \rightarrow S, \sigma_1, \dots, \sigma_{n-1}) \longmapsto (C \rightarrow S, \sigma_1, \dots, \sigma_{j-1}, \alpha \circ \sigma_i, \sigma_j, \dots, \sigma_{n-1})$$

is defined, where α is the hyperelliptic involution. This is exactly the complement of $\mathcal{W}_{g,n-1}^i \cup \bigcup_{l \neq i} \mathcal{Z}_{g,n-1}^{l,i}$, in $\mathcal{H}_{g,n-1}$.

Remark 4.2. By construction, the above map defines a closed immersion

$$u_{i,j} : \mathcal{H}_{g,n-1}^{i,j} \hookrightarrow \mathcal{H}_{g,n}$$

identifying $\mathcal{H}_{g,n-1}^{i,j}$ with $\mathcal{Z}_{g,n}^{i,j}$. See also [Lan23, Remark 2.1].

Lemma 4.3. *Let $1 \leq n \leq 2g + 3$. Then the Chow ring of $\mathcal{H}_{g,n}^{\text{far}}$ is generated in degree 1. In particular, for $n = 1, 2$ the Chow ring is generated by $[\mathcal{W}_{g,1}^1]$ and ψ_1 , while for $n \geq 3$ it is generated by $[\mathcal{W}_{g,n}^1]$.*

Proof. We know that this is true for $n = 1, 2$, see Theorem 2.14 and Proposition 3.11. For $3 \leq n \leq 2g + 3$, thanks to Proposition 1.6, we know that $\mathcal{H}_{g,n}^{\text{far}}$ is an open substack of a weighted projective space, and the localization sequence concludes. The last part follows from the first and Subsection 1.2. \square

Proposition 4.4. *For all $1 \leq n \leq 2g + 3$, $\text{CH}^*(\mathcal{H}_{g,n})$ is generated in degree 1.*

Proof. We already know that the statement holds for $n = 1$ and for $\mathcal{H}_{g,n}^{\text{far}}$. We prove the Proposition by induction on n . Suppose that the statement is true for $n - 1$, and consider the following exact sequence

$$(3) \quad \bigoplus_{i,j} \text{CH}^*(\mathcal{H}_{g,n-1}^{i,j}) \xrightarrow{\sum (u_{i,j})^*} \text{CH}^*(\mathcal{H}_{g,n}) \xrightarrow{\iota^*} \text{CH}^*(\mathcal{H}_{g,n}^{\text{far}}) \longrightarrow 0$$

for all $i < j$. Since $\text{CH}^*(\mathcal{H}_{g,n-1})$ is generated in degree 1, the same holds for the Chow ring of $\mathcal{H}_{g,n-1}^{i,j}$, hence the restriction $(u_{i,j})^*$ is surjective. It follows from the projection formula that the image of $\sum (u_{i,j})_*$ is the ideal generated by the $\mathcal{Z}_{g,n}^{i,j}$.

Now, let β be an element of $\text{CH}^*(\mathcal{H}_{g,n})$. Since $[\mathcal{W}_{g,n}^1]$ and ψ_1 generate $\text{CH}^*(\mathcal{H}_{g,n}^{\text{far}})$, we can modify β with an element generated in degree 1 so that its restriction to $\mathcal{H}_{g,n}^{\text{far}}$ is 0. Therefore, we may assume from now on that $\iota^*(\beta) = 0$. Thanks to the exactness of the sequence (3), α is in the image of $\sum_{i,j} (u_{i,j})_*$, which we have shown to be the ideal generated by the $[\mathcal{Z}_{g,n}^{i,j}]$. Actually, more is true. Indeed, by the inductive hypothesis we know that every element of the Chow ring of $\mathcal{H}_{g,n-1}^{i,j}$ is a product of classes of degree 1, and so it can be written as the pullback via $(u_{i,j})^*$ of a product of classes of degree 1 in the Chow ring of $\mathcal{H}_{g,n}$. It follows from the projection formula that every element in the image of $\sum_{i,j} (u_{i,j})_*$ is a product of classes of degree 1; in particular, the same holds for β . This concludes. \square

Corollary 4.5. *Assume that $\text{char } k > 2g$. Then the classes $[\mathcal{W}_{g,n}^1]$, ψ_1 , $[\mathcal{Z}_{g,2}^{1,2}]$ form a minimal set of generators of $\text{CH}^*(\mathcal{H}_{g,2})$, and*

$$(4) \quad (4g+2)(g+1)\psi_1 = (4g+2)(g-1)[\mathcal{W}_{g,n}^1]$$

is the only relation in degree 1.

Let $n \geq 3$. Then

$$\{[\mathcal{Z}_{g,n}^{1,j}]\}_{2 \leq j \leq n} \cup \{[\mathcal{Z}_{g,n}^{2,3}]\} \cup \{[\mathcal{W}_{g,n}^1]\}$$

is a minimal set of generators of $\text{CH}^*(\mathcal{H}_{g,n})$, and

$$(5) \quad (8g+4)[\mathcal{W}_{g,n}^1] = (4g+2)(g+1)([\mathcal{Z}_{g,n}^{1,2}] + [\mathcal{Z}_{g,n}^{1,3}] - [\mathcal{Z}_{g,n}^{2,3}])$$

is the only relation in degree 1. \square

Remark 4.6. Notice that Proposition 4.4 works for all n such that $\text{CH}^*(\mathcal{H}_{g,n}^{\text{far}})$ and $\text{CH}^*(\mathcal{H}_{g,n-1})$ are generated in degree 1. Thus, the previous observation tells us that the first examples of non-tautological classes are to be looked for in $\text{CH}^*(\mathcal{H}_{g,n}^{\text{far}})$.

4.2. Intersection of Chow classes. We are left with computing the intersections between the divisors in Corollary 4.5.

Lemma 4.7. *Let $n \geq 2$. In $\text{CH}^*(\mathcal{H}_{g,n})$, the following holds.*

$$(6) \quad [\mathcal{Z}_{g,n}^{1,i}] \cdot [\mathcal{W}_{g,n}^1] = 0 \quad \text{for all } i \neq 1$$

$$(7) \quad [\mathcal{Z}_{g,n}^{1,i}] \cdot [\mathcal{Z}_{g,n}^{1,j}] = 0 \quad \text{for all } 1 < i < j$$

$$(8) \quad [\mathcal{Z}_{g,n}^{1,i}] \cdot [\mathcal{Z}_{g,n}^{2,3}] = 0 \quad \text{for } i = 2, 3.$$

Proof. It follows immediately from the fact that the intersection of those divisors is empty. \square

Lemma 4.8. *Let $n \geq i > 3$. Then,*

$$(9) \quad [\mathcal{Z}_{g,n}^{1,i}]^2 = -[\mathcal{Z}_{g,n}^{1,i}] \cdot [\mathcal{Z}_{g,n}^{2,3}] = [\mathcal{Z}_{g,n}^{2,3}]^2.$$

Proof. Notice that, by Proposition 1.2,

$$[\mathcal{Z}_{g,n}^{1,i}] = -[\mathcal{Z}_{g,n}^{2,3}] + [\mathcal{Z}_{g,n}^{1,2}] + [\mathcal{Z}_{g,n}^{3,i}].$$

Multiplying by $[\mathcal{Z}_{g,n}^{1,i}]$ and $[\mathcal{Z}_{g,n}^{2,3}]$ we get the two equalities, respectively. \square

Lemma 4.9. *Let $n \geq 5$ and $i > 3$, then*

$$(10) \quad [\mathcal{Z}_{g,n}^{1,2}]^2 = [\mathcal{Z}_{g,n}^{1,3}]^2 = [\mathcal{Z}_{g,n}^{1,i}]^2 = -[\mathcal{Z}_{g,n}^{1,i}] \cdot [\mathcal{Z}_{g,n}^{2,3}] = [\mathcal{Z}_{g,n}^{2,3}]^2.$$

Proof. It follows from the previous Lemma and the fact that the relation

$$[\mathcal{Z}_{g,n}^{1,4}]^2 = [\mathcal{Z}_{g,n}^{1,5}]^2$$

has to hold even after permuting the indices. \square

In the case $n \geq 5$, thanks to the above Lemmas, we are left with computing the relations $[\mathcal{Z}_{g,n}^{2,3}]^2$, $[\mathcal{Z}_{g,n}^{2,3}] \cdot [\mathcal{W}_{g,n}^1]$ and $[\mathcal{W}_{g,n}^1]^2$. Of course, this takes care only of the degree 2 component of $\text{CH}^*(\mathcal{H}_{g,n})$; however, we will see that, together with knowing $\text{CH}^*(\mathcal{H}_{g,n}^{\text{far}})$, this is enough.

Lemma 4.10. *Let $n \geq 2$. Then, for all $i \neq j$,*

$$[\mathcal{Z}_{g,n}^{i,j}]|_{\mathcal{Z}_{g,n}^{i,j}} = -\psi_i|_{\mathcal{Z}_{g,n}^{i,j}} = -\psi_j|_{\mathcal{Z}_{g,n}^{i,j}}$$

in $\text{Pic}(\mathcal{Z}_{g,n}^{i,j})$. In particular,

$$[\mathcal{Z}_{g,n}^{i,j}]^2 = -[\mathcal{Z}_{g,n}^{i,j}] \cdot \psi_i = -[\mathcal{Z}_{g,n}^{i,j}] \cdot \psi_j$$

in $\text{CH}^*(\mathcal{H}_{g,n})$.

Proof. It can be proved in different ways. For instance, if g is even then it follows from the last two relations of Lemma 4.17, while for $n \geq 3$ it is a consequence of Proposition 4.19. In general, we can argue as follows. Clearly, it is enough to show it for $n = 2$. Let

$$\mathcal{C}_{g,2} \xrightarrow{f} \mathcal{P}_{g,2} \xrightarrow{\pi} \mathcal{H}_{g,2}$$

be the universal family, with universal sections $\tilde{\sigma}_1, \tilde{\sigma}_2$, and universal ‘intermediate’ sections $\sigma_1 := f \circ \tilde{\sigma}_1$ and $\sigma_2 := f \circ \tilde{\sigma}_2$. Let L_i^\vee be the ideal sheaf associated to $\sigma_i(\mathcal{H}_{g,2}) := \mathcal{D}_{g,2}^i \subset \mathcal{P}_{g,2}$. Thanks to [Lan23, Lemma 2.3], we have $\sigma_2^* \mathcal{O}_{\mathcal{D}_{g,2}^1} = \mathcal{O}_{\mathcal{Z}_{g,2}^{1,2}}$. Applying σ_2^* to the exact sequence

$$0 \longrightarrow L_1^\vee \longrightarrow \mathcal{O}_{\mathcal{P}_{g,2}} \longrightarrow \mathcal{O}_{\mathcal{D}_{g,2}^1} \longrightarrow 0$$

we get a surjective map $\sigma_2^*(L_1^\vee) \rightarrow \mathcal{O}_{\mathcal{H}_{g,2}}(-\mathcal{Z}_{g,2}^{1,2})$ between invertible sheaves, hence an isomorphism. It follows that

$$\mathcal{O}_{\mathcal{H}_{g,2}}(\mathcal{Z}_{g,2}^{1,2}) \simeq \sigma_2^* L_1 \simeq \sigma_1^* L_2.$$

Let $\mathcal{C} := \mathcal{C}_{g,2}$, $\mathcal{P} := \mathcal{P}_{g,2}$ and $\mathcal{H} := \mathcal{H}_{g,2}$. Recall that we have an exact sequence of sheaves

$$0 \longrightarrow f^* \omega_{\mathcal{P}/\mathcal{H}} \longrightarrow \omega_{\mathcal{C}/\mathcal{H}} \longrightarrow \Omega_{\mathcal{C}/\mathcal{P}} \longrightarrow 0$$

where $\Omega_{\mathcal{C}/\mathcal{P}}$ is supported on the Weierstrass divisor. Let $u_{1,2} : \mathcal{Z}_{g,2}^{1,2} \hookrightarrow \mathcal{H}_{g,2}$ be the inclusion. Applying $u_{1,2}^* \circ \tilde{\sigma}_1^*$ to that exact sequence, we get

$$u_{1,2}^* \tilde{\sigma}_1^*(\omega_{\mathcal{P}/\mathcal{H}}) \longrightarrow u_{1,2}^* \tilde{\sigma}_1^*(\omega_{\mathcal{C}/\mathcal{H}}) \longrightarrow u_{1,2}^* \tilde{\sigma}_1^*(\Omega_{\mathcal{C}/\mathcal{P}}) \longrightarrow 0.$$

Notice that the middle piece corresponds to the restriction of ψ_1 to $\mathcal{Z}_{g,2}^{1,2}$. Since the image of $\tilde{\sigma}_1 \circ u_{1,2}$ does not intersect the Weierstrass divisor, $u_{1,2}^* \tilde{\sigma}_1^* \Omega_{\mathcal{C}/\mathcal{P}} = 0$, and we get an isomorphism

$$u_{1,2}^* \tilde{\sigma}_1^* \omega_{\mathcal{P}/\mathcal{H}} \xrightarrow{\cong} u_{1,2}^* \tilde{\sigma}_1^* \omega_{\mathcal{C}/\mathcal{H}}.$$

Now, $\sigma_1^* \omega_{\mathcal{P}/\mathcal{H}} \simeq \sigma_1^*(L_1^\vee)$, and clearly $u_{1,2}^* \sigma_1^* \simeq u_{1,2}^* \sigma_2^*$. It follows that

$$u_{1,2}^* \mathcal{O}_{\mathcal{H}_{g,2}}(-\mathcal{Z}_{g,2}^{1,2}) \simeq u_{1,2}^* \sigma_2^* L_1^\vee \simeq u_{1,2}^* \sigma_1^* L_1^\vee \simeq u_{1,2}^* \sigma_1^* \omega_{\mathcal{P}/\mathcal{H}} \simeq u_{1,2}^* \tilde{\sigma}_1^*(\omega_{\mathcal{C}/\mathcal{H}})$$

which proves the Lemma. \square

The following relation is very important, because it lives in degree 1.

Proposition 4.11. *Let $n \geq 3$. Then,*

$$(11) \quad [\mathcal{W}_{g,n}^1] - [\mathcal{W}_{g,n}^2] = (g+1)([\mathcal{Z}_{g,n}^{1,3}] - [\mathcal{Z}_{g,n}^{2,3}]).$$

Proof. The fact that it holds rationally follows from Proposition 1.8 and the rational relation

$$[\mathcal{W}_{g,n}^1] + [\mathcal{W}_{g,n}^2] = (g+1)[\mathcal{Z}_{g,n}^{1,2}]$$

proved in [EH21, Theorem 1.1, Theorem 1.2], or in [Lan23, Lemma 5.10]. Therefore, it is enough to show that the image of $[\mathcal{W}_{g,n}^1] - [\mathcal{W}_{g,n}^2]$ in $\text{Pic}(\mathcal{H}_{g,n}^{\text{far}})$ is 0. This follows immediately from Lemma 1.9 and the definition of the map $\text{Pic}(\mathcal{H}_{g,2}^{\text{far}}) \rightarrow \text{Pic}(\mathcal{H}_{g,3}^{\text{far}})$. \square

Corollary 4.12. *Let $n \geq 3$. Then,*

$$(12) \quad [\mathcal{W}_{g,n}^1] \cdot [\mathcal{Z}_{g,n}^{2,3}] = -(g+1)[\mathcal{Z}_{g,n}^{2,3}]^2.$$

Proof. It follows from the previous Proposition intersecting the relation (11) with $[\mathcal{Z}_{g,n}^{2,3}]$. \square

Corollary 4.13. *Let $n \geq 3$. Then, for all $i \neq j$,*

$$(13) \quad (4g+2)(g+1)[\mathcal{Z}_{g,n}^{i,j}]^2 = 0$$

$$(14) \quad (4g+2)[\mathcal{Z}_{g,n}^{2,3}] \cdot [\mathcal{W}_{g,n}^1] = 0$$

$$(15) \quad (8g+4)[\mathcal{W}_{g,n}^1]^2 = 0$$

Proof. For the first part, it is enough to consider the case of $\mathcal{Z}_{g,n}^{1,2}$. Then it is enough to intersect

$$(8g+4)[\mathcal{W}_{g,n}^1] = (4g+2)(g+1)([\mathcal{Z}_{g,n}^{1,2}] + [\mathcal{Z}_{g,n}^{1,3}] - [\mathcal{Z}_{g,n}^{2,3}])$$

with $[\mathcal{Z}_{g,n}^{1,2}]$. The second part follows from the first and the previous Corollary. The third relation follows from the second, multiplying the relation above by $[\mathcal{W}_{g,n}^1]$. \square

Corollary 4.14. *Let $n \geq 5$. Then, $\text{CH}^2(\mathcal{H}_{g,n})$ is generated by the classes $[\mathcal{W}_{g,n}^1]^2$ and $[\mathcal{Z}_{g,n}^{2,3}]^2$.*

Proof. It follows from Lemma 4.7, Lemma 4.9 and Corollary 4.12. \square

Corollary 4.15. *Let $n \geq 5$. For every $q \geq 3$ the Chow group $\text{CH}^q(\mathcal{H}_{g,n})$ is generated by $[\mathcal{W}_{g,n}^1]^q$. More precisely, the restriction homomorphism*

$$\text{CH}^{q \geq 3}(\mathcal{H}_{g,n}) \longrightarrow \text{CH}^{q \geq 3}(\mathcal{H}_{g,n}^{\text{far}})$$

is an isomorphism of $\text{CH}^(\mathcal{H}_{g,n})$ -modules.*

Proof. We know that the only (possibly) non trivial classes in degree 2 are generated by $[\mathcal{W}_{g,n}^1]^2$ and $[\mathcal{Z}_{g,n}^{2,3}]^2$, see Corollary 4.14. Moreover,

$$[\mathcal{Z}_{g,n}^{2,3}]^3 = [\mathcal{Z}_{g,n}^{1,2}]^2 \cdot [\mathcal{Z}_{g,n}^{2,3}] = 0$$

and similarly for $[\mathcal{W}_{g,n}^1] \cdot [\mathcal{Z}_{g,n}^{2,3}]^2$. Thanks to Corollary 4.12, we have

$$[\mathcal{W}_{g,n}^1]^2 \cdot [\mathcal{Z}_{g,n}^{2,3}] = -(g+1)[\mathcal{W}_{g,n}^1] \cdot [\mathcal{Z}_{g,n}^{2,3}]^2 = 0$$

and the first part of the statement is proved. The second part follows immediately from the first and the exact sequence

$$\bigoplus \mathrm{CH}^*(\mathcal{H}_{g,n}^{i,j}) \xrightarrow{\sum (u_{a,b})^*} \mathrm{CH}^*(\mathcal{H}_{g,n}) \xrightarrow{\iota^*} \mathrm{CH}^*(\mathcal{H}_{g,n}^{\mathrm{far}}) \longrightarrow 0.$$

Indeed, the image of $(u_{i,j})_*$ is the ideal generated by $[\mathcal{Z}_{g,n}^{i,j}]$, hence the map is 0 in degree $q \geq 3$. \square

We are left with computing the relations between $[\mathcal{Z}_{g,n}^{2,3}]^2$ and $[\mathcal{W}_{g,n}^1]^2$. We have already obtained partial information in Corollary 4.13.

The following are other interesting relations, particularly important for low n .

Lemma 4.16. *Let $n \geq 2$. Then,*

$$g[\mathcal{W}_{g,n}^1] + [\mathcal{W}_{g,n}^2] = (g+1)\psi_1 + (g+1)[\mathcal{Z}_{g,n}^{1,2}].$$

Proof. It follows from the computation of the classes ψ_1 , $[\mathcal{W}_{g,n}^1]$ and $[\mathcal{W}_{g,n}^2]$ in $\mathrm{Pic}(\mathcal{H}_{g,n}^{\mathrm{far}})$, and from the rational relation $[\mathcal{W}_{g,n}^1] + [\mathcal{W}_{g,n}^2] = (g+1)[\mathcal{Z}_{g,n}^{1,2}]$. \square

Lemma 4.17. *The first psi class ψ_2 in $\mathrm{Pic}(\mathcal{H}_{g,2}^{\mathrm{far}})$ corresponds to*

$$((g-1)/2, 1) \in \mathbb{Z} \oplus \mathbb{Z}/(8g+4)\mathbb{Z}$$

if g is odd, while if g is even it corresponds to

$$(g-1, -g/2+1) \in \mathbb{Z} \oplus \mathbb{Z}/(4g+2)\mathbb{Z}.$$

Moreover, in $\mathrm{Pic}(\mathcal{H}_{g,2})$ it holds

$$\begin{aligned} (g-1)([\mathcal{W}_{g,n}^1] - [\mathcal{W}_{g,n}^2]) &= (g+1)(\psi_1 - \psi_2) \\ [\mathcal{W}_{g,n}^1] + [\mathcal{W}_{g,n}^2] &= \psi_1 + \psi_2 + 2[\mathcal{Z}_{g,2}^{1,2}] \\ (g-1)[\mathcal{W}_{g,2}^1] + \psi_2 &= g\psi_1 + (g-1)[\mathcal{Z}_{g,2}^{1,2}]. \end{aligned}$$

Proof. The first part follows from the computation in Proposition 1.10, while the second follows from the first and the fact that the relations hold rationally. \square

Lemma 4.18. *For all $n \geq 2$, it holds*

$$(2g+1)(g+1)\psi_1^2 + (2g+1)(3g-1)[\mathcal{W}_{g,n}^1]^2 - (2g+1)(g+1)[\mathcal{Z}_{g,n}^{1,2}]^2.$$

For all $n \geq 3$, $i \neq j$ and $i' \neq j'$, it holds

$$(2g+1)(g+1)[\mathcal{Z}_{g,n}^{i,j}]^2 = (2g+1)(g+1)[\mathcal{Z}_{g,n}^{i',j'}]^2 = 0.$$

Proof. Applying the two above Lemmas to the relation $[\mathcal{W}_{g,2}^2]^2 + \psi_2[\mathcal{W}_{g,2}^2] = 0$ we get the first part. The second follows from the first and Proposition 4.19. \square

The following identity is very important.

Proposition 4.19. *Let $n \geq 3$. Then,*

$$(16) \quad [\mathcal{W}_{g,n}^1] = \psi_1 + [\mathcal{Z}_{g,n}^{1,2}] + [\mathcal{Z}_{g,n}^{1,3}] - [\mathcal{Z}_{g,n}^{2,3}].$$

Proof. From the definition of the map $\text{Pic}(\mathcal{H}_{g,2}^{\text{far}}) \rightarrow \text{Pic}(\mathcal{H}_{g,3}^{\text{far}})$, we know that the image of ψ_1 in $\text{Pic}(\mathcal{H}_{g,n}^{\text{far}})$ is the same as $[\mathcal{W}_{g,n}^1]$, for $n \geq 3$. It follows that their difference is expressible as a sum of the $[\mathcal{Z}_{g,n}^{1,j}]$'s and $[\mathcal{Z}_{g,n}^{2,3}]$ in a unique way. Moreover, putting together Proposition 4.11 and Lemma 4.16, we get that

$$(g+1)[\mathcal{W}_{g,n}^1] = (g+1)\psi_1 + (g+1)([\mathcal{Z}_{g,n}^{1,2}] + [\mathcal{Z}_{g,n}^{1,3}] - [\mathcal{Z}_{g,n}^{2,3}]).$$

Since there are no relations between the $[\mathcal{Z}_{g,n}^{1,j}]$'s and $[\mathcal{Z}_{g,n}^{2,3}]$, the statement follows. \square

Recall that in Corollary 2.17 we have proved that $[\mathcal{W}_{g,1}^1]^2 = -[\mathcal{W}_{g,1}] \cdot \psi_1$. It is worth mentioning that it was possible to prove it using some results in [EH22].

Theorem 4.20 ([EH22, Theorem 1.3]). *The Chow ring of $\mathcal{W}_{g,1}^1$ is*

$$\text{CH}^*(\mathcal{W}_{g,1}^1) = \frac{\mathbb{Z}[\psi_1]}{(4g(2g+1)\psi_1)}.$$

Consider the restriction of $[\mathcal{W}_{g,1}^1]$ to $\mathcal{W}_{g,1}^1$; we want to express it in terms of ψ_1 . This is done in the proof of the above Theorem, see the proofs of [EH22, Corollary 5.4] and [EH22, Corollary 5.7].

Lemma 4.21 ([EH22]). *The restriction of $[\mathcal{W}_{g,1}^1]$ to $\mathcal{W}_{g,1}^1$ is*

$$[\mathcal{W}_{g,1}^1]|_{\mathcal{W}_{g,1}^1} = -\psi_1|_{\mathcal{W}_{g,1}^1}.$$

In particular, in $\text{CH}^*(\mathcal{H}_{g,1})$ we have

$$[\mathcal{W}_{g,1}^1]^2 = -[\mathcal{W}_{g,1}^1] \cdot \psi_1.$$

Proposition 4.22. *Let $n \geq 3$. Then,*

$$(17) \quad 2[\mathcal{W}_{g,n}^1]^2 = -[\mathcal{W}_{g,n}^1] \cdot [\mathcal{Z}_{g,n}^{2,3}] = (g+1)[\mathcal{Z}_{g,n}^{2,3}]^2.$$

Proof. From Proposition 4.19 and Corollary 4.12, we know that

$$[\mathcal{W}_{g,n}^1]^2 = [\mathcal{W}_{g,n}^1] \cdot \psi_1 - [\mathcal{W}_{g,n}^1] \cdot [\mathcal{Z}_{g,n}^{2,3}] = [\mathcal{W}_{g,n}^1] \cdot \psi_1 + (g+1)[\mathcal{Z}_{g,n}^{2,3}]^2$$

and one concludes using the previous Lemma. \square

Since we will see that $[\mathcal{W}_{g,n}^1]^2 \neq 0$ in $\text{CH}^*(\mathcal{H}_{g,n}^{\text{far}})$, we are left with computing the order of $[\mathcal{Z}_{g,n}^{2,3}]^2$. We only know that it divides $(4g+2)(g+1)$ when $n \geq 3$. We will manage to compute it completely only for $n = 2$, and to find lower bounds greater than 1 when $n = 3, 4$.

5. COMPUTATION OF $\text{CH}^*(\mathcal{H}_{g,2})$

We have computed the Chow ring of $\mathcal{H}_{g,2}^{\text{far}}$ in Section 3, Corollary 3.12. Recall that we have an exact sequence

$$\text{CH}^*(\mathcal{H}_{g,1}^{1,2}) \xrightarrow{(u_{1,2})^*} \text{CH}^*(\mathcal{H}_{g,2}) \longrightarrow \text{CH}^*(\mathcal{H}_{g,2}^{\text{far}}) \longrightarrow 0$$

where $\mathcal{H}_{g,1}^{1,2} = \mathcal{H}_{g,1} \setminus \mathcal{W}_{g,1}^1$, see Definition 4.1. Then we can complete the above exact sequence to a commutative diagram

$$(18) \quad \begin{array}{ccccccc} \mathrm{CH}^*(\mathcal{H}_{g,1}^{1,2}) & \xrightarrow{(u_{1,2})^*} & \mathrm{CH}^*(\mathcal{H}_{g,2}) & \longrightarrow & \mathrm{CH}^*(\mathcal{H}_{g,2}^{\mathrm{far}}) & \longrightarrow & 0 \\ & \searrow (u_{1,2})^* & \downarrow & & & & \\ & & \mathrm{CH}^*(\mathcal{H}_{g,2} \setminus \mathcal{W}_{g,n}^1) & & & & \\ & \nearrow q_2^* & & & & & \\ & & & & & & \\ & \nearrow (u_{1,2})^* & & & & & \end{array}$$

where $(u_{1,2})^* q_2^*$ is the identity of $\mathrm{CH}^*(\mathcal{H}_{g,1}^{1,2})$.

This allows us to compute the Chow ring of $\mathcal{H}_{g,2} \setminus \mathcal{W}_{g,2}^1$, which in turn can be used to compute $\mathrm{CH}^*(\mathcal{H}_{g,2})$.

Lemma 5.1. *Suppose $\mathrm{char} k > 2g$. Then,*

$$\mathrm{CH}^*(\mathcal{Z}_{g,2}^{1,2}) \simeq \mathrm{CH}^*(\mathcal{H}_{g,1}^{1,2}) \simeq \frac{\mathbb{Z}[\psi_1]}{((4g+2)(g+1)\psi_1)}.$$

Proposition 5.2. *Suppose $\mathrm{char} k > 2g$. Then, $\mathrm{CH}^*(\mathcal{H}_{g,2} \setminus \mathcal{W}_{g,2}^1)$ is isomorphic to*

$$\frac{\mathbb{Z}[\psi_1, [\mathcal{Z}_{g,2}^{1,2}]]}{((4g+2)(g+1)\psi_1, [\mathcal{Z}_{g,2}^{1,2}]^2 + \psi_1 \cdot [\mathcal{Z}_{g,2}^{1,2}], (2g+1)(g+1)(\psi_1^2 - [\mathcal{Z}_{g,2}^{1,2}]^2))}.$$

Proof. We already know that $\mathrm{CH}^*(\mathcal{H}_{g,2} \setminus \mathcal{W}_{g,2}^1)$ is generated by ψ_1 and $[\mathcal{Z}_{g,2}^{1,2}]$, and that the only relation in degree 1 is $(4g+2)(g+1)\psi_1 = 0$.

Now, we focus on degree 2 relations. We already know that the two degree 2 relations hold, see Lemma 4.10 and Lemma 4.18. Suppose that there is an other general relation

$$d\psi_1^2 + e[\mathcal{Z}_{g,2}^{1,2}]^2 = 0.$$

Restricting $d\psi_1^2 + e[\mathcal{Z}_{g,2}^{1,2}]^2$ to $\mathrm{CH}^*(\mathcal{H}_{g,2}^{\mathrm{far}})$ one gets that d is divisible by $(2g+1)(g+1)$, hence using the last relation we reduce to compute the order of $[\mathcal{Z}_{g,2}^{1,2}]^2$. Now, we know that

$$u_{1,2}^*[\mathcal{Z}_{g,2}^{1,2}] = -u_{1,2}^*\psi_1 = -\psi_1.$$

Therefore, $u_{1,2}^*[\mathcal{Z}_{g,2}^{1,2}]^2$ has order $(4g+2)(g+1)$, thanks to the previous Lemma, and the relation is generated by those in the statement. A similar argument works in degrees greater than 2. \square

Theorem 5.3. *Let $\mathrm{char} g > 2g$. Then, $\mathrm{CH}^*(\mathcal{H}_{g,2})$ is generated by the classes of ψ_1 , $\mathcal{W}_{g,2}^1$ and $\mathcal{Z}_{g,2}^{1,2}$, and the subgroup of relations is generated by*

$$\begin{aligned} & (4g+2)((g+1)\psi_1 - (g-1)[\mathcal{W}_{g,2}^1]), \\ & [\mathcal{Z}_{g,2}^{1,2}]^2 + \psi_1 \cdot [\mathcal{Z}_{g,2}^{1,2}], \quad [\mathcal{W}_{g,2}^1]^2 + \psi_1 \cdot [\mathcal{W}_{g,2}^1], \quad [\mathcal{Z}_{g,2}^{1,2}] \cdot [\mathcal{W}_{g,2}^1], \\ & (2g+1)(g+1)\psi_1^2 + (2g+1)(3g-1)[\mathcal{W}_{g,2}^1]^2 - (2g+1)(g+1)[\mathcal{Z}_{g,2}^{1,2}]^2. \end{aligned}$$

Proof. We already know that those are the generators, and that the relations in degree 1 are multiples of the first relation in the statement. Moreover, we already know that those are indeed relations.

Now, suppose we have a relation in degree $q \geq 2$. Using the middle row, we get a linear relation between ψ_1^q , $[\mathcal{W}_{g,2}^1]^q$ and $[\mathcal{Z}_{g,2}^{1,2}]^q$. Modulo $\mathcal{Z}_{g,2}^{1,2}$ we get a relation in

$\mathrm{CH}^*(\mathcal{H}_{g,2}^{\mathrm{far}})$. Since the ideal of relations in $\mathcal{H}_{g,2}$ surjects to the one in $\mathcal{H}_{g,2}^{\mathrm{far}}$, we reduce to the computation of the order of $(\mathcal{Z}_{g,2}^{1,2})^q$. This is divisible by $(4g+2)(g+1)$, thanks to Proposition 5.2, and it is exactly $(4g+2)(g+1)$ as one can see by multiplying the relation in degree 1 with $[\mathcal{Z}_{g,2}^{1,2}]$ and using Lemma 4.10. \square

6. THE ALMOST COMPUTATION OF $\mathrm{CH}^*(\mathcal{H}_{g,n})$ FOR $3 \leq n \leq 2g+3$

6.1. The Chow ring of $\mathcal{H}_{g,n}^{\mathrm{far}}$ for $3 \leq n \leq 2g+3$. In this Subsection we compute the Chow ring of $\mathcal{H}_{g,n}^{\mathrm{far}}$ for $3 \leq n \leq 2g+2$, and obtain the generator of the ring in the case $n = 2g+3$. In this last case, the only thing we do not manage to compute is the multiplicative order of the generator.

Recall that

$$\mathcal{H}_{g,n}^{\mathrm{far}} \simeq (\mathbb{P}(2^{2g+3-n}, 1^n) \times (\mathbb{A}^{n-3} \setminus \tilde{\Delta})) \setminus \overline{\Delta}_{g,n}.$$

Moreover,

$$\mathrm{CH}^*(\mathbb{P}(2^{2g+3-n}, 1^n) \times (\mathbb{A}^{n-3} \setminus \tilde{\Delta})) \simeq \frac{\mathbb{Z}[h]}{(2^{2g+3-n} h^{2g+3})}.$$

Proposition 6.1. *Let $\mathrm{char} k > 2g$. If $3 \leq n \leq 2g+2$, then*

$$\mathrm{CH}^*(\mathcal{H}_{g,n}^{\mathrm{far}}) \simeq \frac{\mathbb{Z}[[\mathcal{W}_{g,n}^1]]}{((8g+4)[\mathcal{W}_{g,n}^1], 2[\mathcal{W}_{g,n}^1]^2)}.$$

Moreover, the map induced by forgetting the n -th section, i.e. the last one, is an isomorphism for $4 \leq n \leq 2g+2$.

Proof. First, we know that the above relations hold; see Proposition 4.22 for the relation in degree 2. On the other hand, in [EH22] it has been shown that, for $n \leq 2g+2$, the Chow ring of the intersection of all $\mathcal{W}_{g,n}^i$ has Chow ring generated by the restriction of ψ_1 , with ideal of relations generated by $2\psi_1$. Since for $3 \leq n \leq 2g+3$ the restrictions to $\mathcal{H}_{g,n}^{\mathrm{far}}$ of the classes ψ_1 and $[\mathcal{W}_{g,n}^i]$ agree for every i , this concludes. \square

Remark 6.2. The case $n = 2g+3$ is harder, and we do not compute $\mathrm{CH}^*(\mathcal{H}_{g,2g+3}^{\mathrm{far}})$. The difference with the case $n \leq 2g+2$ is that $\mathcal{H}_{g,n}$ is now a scheme, hence the integral Chow ring has to be 0 in degrees above the dimension. However, some things follow easily from what we have done. Recall that

$$\mathcal{H}_{g,2g+3} \simeq (\mathbb{P}^{2g+2} \times (\mathbb{A}^{2g} \setminus \tilde{\Delta})) \setminus \overline{\Delta}_{g,2g+3}$$

and so the Chow ring is generated by $[\mathcal{W}_{g,2g+3}^1]$, the usual relations hold, and $[\mathcal{W}_{g,2g+3}^1]^{2g+2} = 0$. It is not clear what is the exact multiplicative order of $[\mathcal{W}_{g,2g+3}^1]$. Moreover, it is easy to convince oneself that it is not possible to use similar Chow envelopes as in the cases $n = 0, 1, 2$.

6.2. The Chow ring of $\mathcal{H}_{g,3}$. We apply the same method as for $n = 2$. Recall that we have a commutative diagram

$$(19) \quad \begin{array}{ccccccc} \mathrm{CH}^*(\mathcal{H}_{g,2}^{1,3}) & \xrightarrow{(u_{1,3})^*} & \mathrm{CH}^*(\mathcal{H}_{g,3}) & \longrightarrow & \mathrm{CH}^*(\mathcal{H}_{g,3}^{\mathrm{far}}) & \longrightarrow & 0 \\ & \searrow (u_{1,3})^* & \downarrow & & & & \\ & & \mathrm{CH}^*(\mathcal{H}_{g,3} \setminus \mathcal{W}_{g,n}^1 \cup \mathcal{Z}_{g,3}^{1,2}) & & & & \\ & \nearrow q_3^* & & & & & \\ & & \swarrow (u_{1,3})^* & & & & \end{array}$$

where $(u_{1,3})^* q_3^*$ is the identity of $\text{CH}^*(\mathcal{H}_{g,2}^{1,3})$, and the first row is a complex.

This allows us to compute the Chow ring of $\mathcal{H}_{g,3} \setminus \mathcal{W}_{g,3}^1$, which in turn can be used to compute $\text{CH}^*(\mathcal{H}_{g,3})$.

Lemma 6.3. *Let $\text{char } k > 2g$. Then,*

$$\text{CH}^*(\mathcal{Z}_{g,3}^{1,3}) \simeq \text{CH}^*(\mathcal{H}_{g,2}^{1,3}) \simeq \frac{\mathbb{Z}[\psi_1]}{((4g+2)(g+1)\psi_1, (2g+1)(g+1)\psi_1^2)}.$$

Proof. It follows immediately from Theorem 5.3. \square

Proposition 6.4. *Let $\text{char } k > 2g$. Then, $\text{CH}^*(\mathcal{H}_{g,3} \setminus (\mathcal{W}_{g,3}^1 \cup \mathcal{Z}_{g,3}^{1,2}))$ is isomorphic to*

$$\frac{\mathbb{Z}[\psi_1, [\mathcal{Z}_{g,3}^{1,3}]]}{((4g+2)(g+1)\psi_1, [\mathcal{Z}_{g,3}^{1,3}]^2 + \psi_1 \cdot [\mathcal{Z}_{g,3}^{1,3}], (2g+1)(g+1)\psi_1^2, (2g+1)(g+1)[\mathcal{Z}_{g,3}^{1,3}]^2)}.$$

Proof. We already know that $\text{CH}^*(\mathcal{H}_{g,3} \setminus (\mathcal{W}_{g,3}^1 \cup \mathcal{Z}_{g,3}^{1,2}))$ is generated by ψ_1 and $[\mathcal{Z}_{g,3}^{1,3}]$, and that the only relation in degree 1 is $(4g+2)(g+1)\psi_1 = 0$.

We now focus on degree 2 relations; we already know that those in the statement hold (see Lemma 4.18 for the last one). Suppose there is another general relation

$$d\psi_1^2 + e[\mathcal{Z}_{g,3}^{1,3}]^2 = 0.$$

Now, recall that $[\mathcal{W}_{g,3}^1] = \psi_1 + [\mathcal{Z}_{g,3}^{1,2}] + [\mathcal{Z}_{g,3}^{1,3}] - [\mathcal{Z}_{g,3}^{2,3}]$ in $\text{CH}^*(\mathcal{H}_{g,3})$, so using $[\mathcal{Z}_{g,3}^{2,3}]$ instead of ψ_1 we get the relation $d[\mathcal{Z}_{g,3}^{2,3}]^2 + (d+e)[\mathcal{Z}_{g,3}^{1,3}]^2 = 0$. Restricting to $\mathcal{Z}_{g,3}^{1,3}$, we get that $(2g+1)(g+1)$ divides $d+e$. Restricting to $\mathcal{Z}_{g,3}^{2,3}$ one gets that $(2g+1)(g+1)$ divides d , and thus e too. The above follows from the fact that $\mathcal{Z}_{g,n}^{i,j}$ are all isomorphic, thus they have the same Chow ring, which is described in the above Lemma. Therefore, using $(2g+1)(g+1)[\mathcal{Z}_{g,3}^{1,3}]^2 = 0$ we reduce to computing the order of ψ_1^2 , which is $(2g+1)(g+1)$ thanks to the injectivity of q_3^* .

A similar argument works in degrees greater than 2. \square

Theorem 6.5. *Let $\text{char } k > 2g$. Then, $\text{CH}^*(\mathcal{H}_{g,3})$ is generated by the classes of $\mathcal{W}_{g,3}^1$, $\mathcal{Z}_{g,3}^{1,2}$, $\mathcal{Z}_{g,3}^{1,3}$ and $\mathcal{Z}_{g,3}^{2,3}$, and the ideal of relations is generated by*

$$\begin{aligned} & (8g+4)[\mathcal{W}_{g,3}^1] - (4g+2)(g+1)([\mathcal{Z}_{g,3}^{1,2}] + [\mathcal{Z}_{g,3}^{1,3}] - [\mathcal{Z}_{g,3}^{2,3}]), \\ & [\mathcal{Z}_{g,3}^{1,2}] \cdot [\mathcal{Z}_{g,3}^{1,3}], \quad [\mathcal{Z}_{g,3}^{1,2}] \cdot [\mathcal{Z}_{g,3}^{2,3}], \quad [\mathcal{Z}_{g,3}^{1,3}] \cdot [\mathcal{Z}_{g,3}^{2,3}], \quad [\mathcal{Z}_{g,3}^{1,j}] \cdot [\mathcal{W}_{g,3}^1] \text{ for } j = 2, 3, \\ & [\mathcal{W}_{g,3}^1] \cdot [\mathcal{Z}_{g,3}^{2,3}] + (g+1)[\mathcal{Z}_{g,3}^{2,3}]^2, \quad 2[\mathcal{W}_{g,3}^1]^2 - (g+1)[\mathcal{Z}_{g,3}^{2,3}]^2, \\ & (2g+1)(g+1)([\mathcal{Z}_{g,3}^{i,j}]^2 + [\mathcal{Z}_{g,3}^{i',j'}]^2) \text{ for } i \neq j, i' \neq j', \quad b_g[\mathcal{Z}_{g,3}^{i,j}]^2 \text{ for } i \neq j. \end{aligned}$$

where b_g is either $(2g+1)(g+1)$ or $(4g+2)(g+1)$.

Proof. We already know that those are the generators, and that the relations in degree 1 are multiples of the first relation in the statement. Moreover, we already know that those in the statement are relations, and that they generate those in degree ≥ 2 involving $[\mathcal{W}_{g,3}^1]$. We are left with computing the orders of $[\mathcal{Z}_{g,3}^{i,j}]^q$ for $q \geq 2$. Restricting to $\mathcal{Z}_{g,3}^{i,j}$ we get that their order is divisible by $(2g+1)(g+1)$, and we are done (notice that $(2g+1)(g+1)[\mathcal{Z}_{g,3}]^q = 0$ for all $q \geq 3$). \square

6.3. **The Chow ring of $\mathcal{H}_{g,4}$.** We start in the same way as before.

Lemma 6.6. *Suppose $\text{char } k > 2g$. Then,*

$$\text{CH}^*(\mathcal{H}_{g,3}^{1,4}) \simeq \frac{\mathbb{Z}[\psi_1]}{((4g+2)(g+1)\psi_1, (g+1)\psi_1^2)} \simeq \frac{\mathbb{Z}[[\mathcal{Z}_{g,3}^{2,3}]]}{((4g+2)(g+1)[\mathcal{Z}_{g,3}^{2,3}], (g+1)[\mathcal{Z}_{g,3}^{2,3}]^2)}.$$

Proof. It follows immediately from Theorem 6.5. \square

Proposition 6.7. *Suppose $\text{char } k > 2g$. Then,*

$$\begin{aligned} \text{CH}^*(\mathcal{H}_{g,4} \setminus (\mathcal{W}_{g,4}^1 \cup \mathcal{Z}_{g,4}^{1,2} \cup \mathcal{Z}_{g,4}^{1,3})) \\ \simeq \frac{\mathbb{Z}[\psi_1, [\mathcal{Z}_{g,4}^{1,4}]]}{((4g+2)(g+1)\psi_1, [\mathcal{Z}_{g,4}^{1,4}]^2 + \psi_1 \cdot [\mathcal{Z}_{g,4}^{1,4}], (g+1)\psi_1^2, [\mathcal{Z}_{g,4}^{1,4}]^2 - \psi_1^2)} \\ \simeq \frac{\mathbb{Z}[[\mathcal{Z}_{g,4}^{2,3}], [\mathcal{Z}_{g,4}^{1,4}]]}{((4g+2)(g+1)[\mathcal{Z}_{g,4}^{2,3}], [\mathcal{Z}_{g,4}^{1,4}]^2 + [\mathcal{Z}_{g,4}^{2,3}] \cdot [\mathcal{Z}_{g,4}^{1,4}], [\mathcal{Z}_{g,4}^{2,3}]^2 + [\mathcal{Z}_{g,4}^{2,3}] \cdot [\mathcal{Z}_{g,4}^{1,4}], (g+1)[\mathcal{Z}_{g,4}^{2,3}]^2)}. \end{aligned}$$

Proof. It is proved in the same way as Proposition 6.4. \square

Theorem 6.8. *Let $\text{char } k > 2g$. Then, $\text{CH}^*(\mathcal{H}_{g,4})$ is generated by the classes of $\mathcal{W}_{g,4}^1$, $\mathcal{Z}_{g,4}^{1,2}$, $\mathcal{Z}_{g,4}^{1,3}$, $\mathcal{Z}_{g,4}^{1,4}$ and $\mathcal{Z}_{g,4}^{2,3}$, and the ideal of relations is generated by*

$$\begin{aligned} (8g+4)[\mathcal{W}_{g,4}^1] - (4g+2)(g+1)([\mathcal{Z}_{g,4}^{1,2}] + [\mathcal{Z}_{g,4}^{1,3}] - [\mathcal{Z}_{g,4}^{2,3}]), \\ [\mathcal{Z}_{g,4}^{1,i}] \cdot [\mathcal{Z}_{g,4}^{1,j}] \text{ for } 1 < i < j, \quad [\mathcal{Z}_{g,4}^{1,j}] \cdot [\mathcal{Z}_{g,4}^{2,3}] \text{ for } j = 2, 3, \quad [\mathcal{Z}_{g,4}^{1,j}] \cdot [\mathcal{W}_{g,3}^1] \text{ for } j > 1, \\ [\mathcal{Z}_{g,4}^{2,3}] \cdot [\mathcal{W}_{g,3}^1] + (g+1)[\mathcal{Z}_{g,4}^{2,3}]^2, \quad [\mathcal{Z}_{g,4}^{2,3}]^2 = -[\mathcal{Z}_{g,4}^{2,3}] \cdot [\mathcal{Z}_{g,4}^{1,4}] = [\mathcal{Z}_{g,4}^{1,4}]^2, \\ 2[\mathcal{W}_{g,4}^1]^2 - (g+1)[\mathcal{Z}_{g,4}^{2,3}]^2, \quad (g+1)[\mathcal{Z}_{g,4}^{i,j}]^3 \text{ for all } i \neq j, \\ (g+1)([\mathcal{Z}_{g,4}^{i,j}]^2 - [\mathcal{Z}_{g,4}^{i',j'}]^2) \text{ for all } i \neq j, i' \neq j', \quad b_g[\mathcal{Z}_{g,4}^{i,j}]^2 \text{ for all } i \neq j \end{aligned}$$

for some integer $b_g = b'_g(g+1)$ such that b'_g divides $4g+2$.

Proof. We already know that those are the generators and that the relations hold. The only relation which is less clear is the bottom left. However, notice that

$$2[\mathcal{W}_{g,4}^1]^2 = (g+1)[\mathcal{Z}_{g,4}^{2,3}]^2$$

and applying the automorphism exchanging the fourth and third section we get the desired relation. Finally, the same arguments as in Theorem 6.5 apply to show that the relations in the statement generated the ideal of relations. \square

6.4. **The Chow ring of $\mathcal{H}_{g,n}$ for $5 \leq n \leq 2g+3$.**

Definition 6.9. Consider the map $q_n : \mathcal{H}_{g,n} \rightarrow \mathcal{H}_{g,n-1}$ forgetting the last section. Define $\mathcal{H}_{g,n}^o$ to be preimage of $\mathcal{H}_{g,n-1}^{1,n}$ under q_n , which is an open substack of $\mathcal{H}_{g,n}$.

Remark 6.10. We have already used this stack in the previous sections. For instance, $\mathcal{H}_{g,2}^o = \mathcal{H}_{g,2} \setminus \mathcal{W}_{g,2}^1$, while for $n=3$ we have $\mathcal{H}_{g,3}^o = \mathcal{H}_{g,3} \setminus (\mathcal{W}_{g,3}^1 \cup \mathcal{Z}_{g,3}^{1,2})$.

Proposition 6.11. *Let $\text{char } k > 2g$, and let $5 \leq n \leq 2g+3$. Then,*

$$\text{CH}^*(\mathcal{H}_{g,n-1}^{1,n}) \simeq \frac{\mathbb{Z}[\psi_1]}{((4g+2)(g+1)\psi_1, \psi_1^2)}$$

and

$$\begin{aligned} \mathrm{CH}^*(\mathcal{H}_{g,n}^o) &\simeq \frac{\mathbb{Z}[\psi_1, [\mathcal{Z}_{g,n}^{1,n}]]}{((4g+2)(g+1)\psi_1, \psi_1 \cdot [\mathcal{Z}_{g,n}^{1,n}], \psi_1^2, [\mathcal{Z}_{g,n}^{1,n}]^2)} \\ &\simeq \frac{\mathbb{Z}[[\mathcal{Z}_{g,n}^{2,3}], [\mathcal{Z}_{g,n}^{1,n}]]}{((4g+2)(g+1)[\mathcal{Z}_{g,n}^{2,3}], [\mathcal{Z}_{g,n}^{2,3}] \cdot [\mathcal{Z}_{g,n}^{1,n}], [\mathcal{Z}_{g,n}^{2,3}]^2, [\mathcal{Z}_{g,n}^{1,n}]^2)}. \end{aligned}$$

Proof. Everything follows from what we know and the fact that we are killing $[\mathcal{Z}_{g,n}^{1,4}]$ too, so that all the squares are 0. \square

Theorem 6.12. *Let char $k > 2g$ and $5 \leq n \leq 2g+2$. Then, $\mathrm{CH}^*(\mathcal{H}_{g,n})$ is generated by the classes of $\mathcal{W}_{g,n}^1$, $\mathcal{Z}_{g,n}^{1,i}$ for $1 < i \leq n$ and $\mathcal{Z}_{g,n}^{2,3}$, and the ideal of relations is generated by*

$$\begin{aligned} &(8g+4)[\mathcal{W}_{g,n}^1] - (4g+2)(g+1)([\mathcal{Z}_{g,n}^{1,2}] + [\mathcal{Z}_{g,n}^{1,3}] - [\mathcal{Z}_{g,n}^{2,3}]), \\ &[\mathcal{Z}_{g,n}^{1,i}] \cdot [\mathcal{Z}_{g,n}^{1,j}] \text{ for } 1 < i < j, \quad [\mathcal{Z}_{g,n}^{1,j}] \cdot [\mathcal{Z}_{g,n}^{2,3}] \text{ for } j = 2, 3, \quad [\mathcal{Z}_{g,n}^{1,j}] \cdot [\mathcal{W}_{g,n}^1] \text{ for } j \neq 1, \\ &[\mathcal{Z}_{g,n}^{2,3}] \cdot [\mathcal{W}_{g,n}^1] + (g+1)[\mathcal{Z}_{g,n}^{2,3}]^2, \quad 2[\mathcal{W}_{g,n}^1]^2 - (g+1)[\mathcal{Z}_{g,n}^{2,3}]^2, \\ &[\mathcal{Z}_{g,n}^{2,3}]^2 = -[\mathcal{Z}_{g,n}^{2,3}] \cdot [\mathcal{Z}_{g,n}^{1,i}] = [\mathcal{Z}_{g,n}^{1,j}]^2 \text{ for all } i > 3 \text{ and } j \neq 1, \\ &b_g[\mathcal{Z}_{g,n}^{i,j}]^2 \text{ for all } i \neq j \end{aligned}$$

where b_g is a positive integer dividing $(4g+2)(g+1)$.

Proof. It can be proved in the same way as before. \square

Remark 6.13. Notice that the major changes that happen varying n from 2 to 5 are due to the variations of the order of ψ_1^2 in $\mathrm{CH}(\mathcal{H}_{g,n-1}^{1,n})$. This essentially measures how much movable the divisors $[\mathcal{Z}_{g,n}^{i,j}]$ are.

In the same way one can show the following result.

Theorem 6.14. *Let char $k > 2g$ and $n = 2g+3$. Then, $\mathrm{CH}^*(\mathcal{H}_{g,2g+3})$ is generated by the classes of $\mathcal{W}_{g,2g+3}^1$, $\mathcal{Z}_{g,2g+3}^{1,i}$ for $1 < i \leq 2g+3$ and $\mathcal{Z}_{g,2g+3}^{2,3}$, and the ideal of relations is generated by*

$$\begin{aligned} &(8g+4)[\mathcal{W}_{g,2g+3}^1] - (4g+2)(g+1)([\mathcal{Z}_{g,2g+3}^{1,2}] + [\mathcal{Z}_{g,2g+3}^{1,3}] - [\mathcal{Z}_{g,2g+3}^{2,3}]), \\ &[\mathcal{Z}_{g,2g+3}^{1,i}] \cdot [\mathcal{Z}_{g,2g+3}^{1,j}] \text{ for } 1 < i < j, \quad [\mathcal{Z}_{g,2g+3}^{1,j}] \cdot [\mathcal{Z}_{g,2g+3}^{2,3}] \text{ for } j = 2, 3, \\ &[\mathcal{Z}_{g,2g+3}^{1,j}] \cdot [\mathcal{W}_{g,2g+3}^1] \text{ for } j \neq 1, \quad [\mathcal{Z}_{g,2g+3}^{2,3}] \cdot [\mathcal{W}_{g,2g+3}^1] + (g+1)[\mathcal{Z}_{g,2g+3}^{2,3}]^2, \\ &2[\mathcal{W}_{g,2g+3}^1]^2 - (g+1)[\mathcal{Z}_{g,2g+3}^{2,3}]^2, \\ &[\mathcal{Z}_{g,2g+3}^{2,3}]^2 = -[\mathcal{Z}_{g,2g+3}^{2,3}] \cdot [\mathcal{Z}_{g,2g+3}^{1,i}] = [\mathcal{Z}_{g,2g+3}^{1,j}]^2 \text{ for all } i > 3 \text{ and } j \neq 1, \\ &b_g[\mathcal{Z}_{g,n}^{i,j}]^2 \text{ for all } i \neq j, \quad [\mathcal{W}_{g,2g+3}^1]^{c_g} \end{aligned}$$

where b_g is a positive integer dividing $(4g+2)(g+1)$, and $1 < c_g \leq 2g+2$.

Remark 6.15. For $3 \leq n \leq 2g+2$, we are left with computing b_g , which we do not do in this article. There are some difficulties in using higher Chow rings (with \mathbb{Z}_l -coefficients) in this case, since it can be shown that the first higher Chow ring of $\mathcal{H}_{g,n}^{\mathrm{far}}$ is non-zero in the cases we are interested in. Notice that b_g is also the order of $[\mathcal{Z}_{g,n}^{2,3}] \cdot [\mathcal{Z}_{g,n}^{1,4}]$ for $n \geq 4$, so that at least we are dealing with a proper intersection.

REFERENCES

- [AV04] Alessandro Arsie and Angelo Vistoli, *Stacks of cyclic covers of projective spaces*, Compos. Math. **140** (2004), no. 3, 647–666. MR 2041774
- [Cad04] Charles Cadman, *Using stacks to impose tangency conditions on curves*, American Journal of Mathematics **129** (2004).
- [CL22] Samir Canning and Hannah Larson, *The rational Chow rings of moduli spaces of hyperelliptic curves with marked points*, Annali della Scuola Normale Superiore (2022).
- [DL18] Andrea Di Lorenzo, *The Chow ring of the stack of hyperelliptic curves of odd genus*, International Mathematics Research Notices (2018).
- [EF09] Dan Edidin and Damiano Fulghesu, *The integral Chow ring of the stack of hyperelliptic curves of even genus*, Math. Res. Lett. **16** (2009), no. 1, 27–40. MR 2480558
- [EG98] Dan Edidin and William Graham, *Equivariant intersection theory*, Invent. Math. **131** (1998), no. 3, 595–634. MR 1614555
- [EH21] Dan Edidin and Zhengning Hu, *Chow classes of divisors on stacks of pointed hyperelliptic curves*, Annali della Scuola Normale Superiore (2021).
- [EH22] ———, *The integral Chow rings of the stacks of hyperelliptic Weierstrass points*, arXiv preprint arXiv:2208.00556 (2022).
- [Lan23] Alberto Landi, *The Picard group of the stack of pointed smooth cyclic covers of the projective line*, 2023.
- [Lar21] Eric Larson, *The integral Chow ring of \overline{M}_2* , Algebr. Geom. **8** (2021), no. 3, 286–318. MR 4206438
- [Mum77] David Mumford, *Stability of projective varieties*, Monographies de L’Enseignement Mathématique [Monographs of L’Enseignement Mathématique], vol. No. 24, L’Enseignement Mathématique, Geneva, 1977, Lectures given at the “Institut des Hautes Études Scientifiques”, Bures-sur-Yvette, March-April 1976. MR 450273
- [Ols16] Martin Olsson, *Algebraic spaces and stacks*, 2016.
- [Per22] Michele Pernice, *The integral Chow ring of the stack of 1-pointed hyperelliptic curves*, Int. Math. Res. Not. IMRN (2022), no. 15, 11539–11574. MR 4458558
- [PG01] Rahul Pandharipande and Tom Graber, *Constructions of nontautological classes of moduli spaces of curves*, Michigan Mathematical Journal **51** (2001), 93–110.
- [Vis98] Angelo Vistoli, *The Chow ring of \mathcal{M}_2 . Appendix to “Equivariant intersection theory” [Invent. Math. **131** (1998), no. 3, 595–634; MR1614555 (99j:14003a)] by D. Edidin and W. Graham*, Invent. Math. **131** (1998), no. 3, 635–644. MR 1614559

SCUOLA NORMALE SUPERIORE, PIAZZA CAVALIERI 7, 56126 PISA, ITALY
 Email address: alberto.landi@brown.edu