Nonlocal order parameter of pair superfluids

Nitya Cuzzuol 1* , Arianna Montorsi 1† and Luca Barbiero 1

1 Institute for Condensed Matter Physics and Complex Systems, DISAT, Politecnico di Torino, I-10129 Torino, Italy

⋆ nitya.cuzzuol@polito.it , † arianna.montorsi@polito.it

Abstract

Order parameters represent a fundamental resource to characterize quantum matter. We show that pair superfluids can be rigorously defined in terms of a nonlocal order parameter, named odd parity, which derivation is experimentally accessible by local density measurements. As a case of study, we first investigate a constrained Bose-Hubbard model at different densities, both in one and two spatial dimensions. Here, our analysis finds pair superfluidity for relatively strong attractive interactions. The odd parity operator acts as the unique order parameter for such phase irrespectively to the density of the system and its dimensionality. In order to enforce our finding, we confirm the generality of our approach also on a two-component Bose-Hubbard Hamiltonian, which experimental realization represents a timely topic in ultracold atomic systems. Our results shed new light on the role of correlated density fluctuations in pair superfluids. In addition, they provide a powerful tool for the experimental detection of such exotic phases and the characterization of their transition to the normal superfluid phase.

Copyright attribution to authors.

This work is a submission to SciPost Physics.

License information to appear upon publication.

Publication information to appear upon publication.

Received Date Accepted Date Published Date

Contents

1	Introduction	2
2	The model	3
	2.1 Bosonization analysis	4
3	Numerical results	5
	3.1 Results in 1D	6
	3.2 Results in 2D	6
	3.3 Arbitrary Filling	7
	3.4 Two-species system	8
4	Conclusions	9
References		10

1 Introduction

Quantumness is fundamentally rooted in delocalization and entanglement, the latter allowing quantum matter to organize in low temperature phases which escape Landau classification of spontaneously symmetry breaking (SSB) phases [1,2]. Notable examples are topological phases [3–7], as well as insulators with continuum symmetries, i.e. low dimensional Mott insulators [8]. Crucially, while SSB phases are endowed with a local order, the aforementioned examples are characterized by nonlocal orderings which do not violate the Mermin-Wagner theorem [9]. Such intrinsic nonlocality allows for insulators with continuous symmetries both in 1D and 2D to be accurately captured by nonlocal parity operators [10–14], while nonlocal string order parameters [15–17] result the fundamental quantity to capture one dimensional topological insulators. Relevantly, from SSB [18–24] to low dimensional Mott [25–28] and topological [29,30] insulators, experimental setups working with ultracold atoms [31,32] have allowed for an in depth description of such states of quantum matter. Besides the impressive versatility encoded in such quantum simulators, a game changing aspect is represented by highly accurate probing schemes. By indeed making use of either noise correlators [33,34] or quantum gas microscopy [35] both local and nonlocal orderings have been efficiently probed.

Although the above discussion focused on insulating states of matter, the notion of nonlocal orders can be applied also to characterize conducting phases of low-dimensional interacting fermions [12,36–38]. In this regard, nonlocal parity [11,12] and string [17] operators relative to the spin degrees of freedom have been efficiently employed to characterize Luther Emery liquid [11] and superconducting phases [36–38] as well as topological conducting regimes [36,38] respectively. At the same time, the order parameters of gapped conducting phases in low dimensional bosonic matter are still unexplored. In this paper we tackle this important topic.

In particular, we explore conducting phases characterized by bosonic pairing, i. e. pair superfluids (PSFs) whose existence has been predicted in large variety of physical systems [39–52]. Importantly, PSFs are neither associated to a symmetry breaking nor to the appearance of topological order, thus representing an example of a gapped superfluid with continuous symmetry. To perform our analysis, we consider a paradigmatic model of strongly interacting bosons, namely the Bose-Hubbard model with a constraint of a maximum of two bosons per site. It was initially introduced to describe regimes characterized by a large rate of three-body losses in ultracold atomic gases [39, 40]. Previous theoretical works [39, 41] have established that for weakly attractive interaction a normal superfluid (SF) phase occurs while, by increasing the attraction between bosons, the system undergoes a phase transition. Here, a different SF phase, i.e. the PSF where pairs of single bosons become the effective elementary constituents, appears. As we will discuss in details, these different superfluids have been characterized by analyzing the different long distance decay of the single particle and pair-pair correlation function [39]. Although certainly useful, this approach poses significant challenges. On one hand it makes difficult to properly locate the transition point. On the other hand, experimental schemes to probe the long distance behavior of correlators are still characterized by a limited efficiency. Here, we propose an alternative approach based on local density measurements, which can be performed with high accuracy through quantum gas microscopy.

Specifically, our method introduces a new nonlocal parameter, named **odd parity**. The prediction is first confirmed by a bosonization and renormalization group analysis. We then prove also numerically that the odd parity is the unique order parameter of the PSF phase. As PSF can occur also in 2D [41, 43, 44, 46], we extend the definition of odd parity by means of the brane generalization [37,53] and we verify numerically that our approach holds also beyond the one dimensional limit. In addition, since sample imperfections might make it challenging

to accurately control the particle density, we show that our method remains highly robust in a large range of bosonic fillings. Finally, we test the validity of our predictions on an experimentally feasible two component Bose-Hubbard Hamiltonian where the SF-PSF phase transition was recently predicted [54–59].

2 The model




Figure 1: Cartoon of a phase with finite odd parity order (see (8) and (10)) for a 1D (a) and 2D (b) lattice respectively . It amounts to a background of holons (empty sites) and doublons (doubly occupied sites) in which density fluctuations (single bosons) occur only in pairs with finite correlation length, highlighted by red shaded regions. Panels (c) and (d) present the resulting phase diagrams obtained through iDMRG simulations for the 1D system (with $\chi_{max} = 200$) and a lattice system with M = 6 (with $\chi_{max} = 1500$), respectively. The SF region (depicted in blue) corresponds to zero expectation value of odd parity, while the PSF region (depicted in red) is characterized by a non-zero value.

We analyse the Bose-Hubbard model corresponding to the following Hamiltonian

$$H = -t \sum_{\langle R, R' \rangle} (b_R^{\dagger} b_{R'} + \text{h.c.}) + \frac{U}{2} \sum_{R} n_R (n_R - 1) + \mu \sum_{R} n_R$$
 (1)

where R = (x, y) is a generic site of a lattice composed by L rungs and M legs $(1 \le x \le L; 1 \le y \le M, M = 1 \text{ in 1D})$ schematized in figure (1)(a,b). Here, $\langle R, R' \rangle$ indicates nearest-neighbor sites and b_R , b_R^{\dagger} are the bosonic creation and annihilation operators on the site R. In addition, μ is the system chemical potential and the hopping matrix element t = 1 fixes the energy scale. In order to favor the bosonic pairing we focus on regimes with on-site attractive interaction U < 0 and we impose the three body constraint $(b_R^{\dagger})^3 = 0$ which allows to prevent the system collapse. As shown both in the 1D and 2D case in figure (1)(c,d), bosons remain in the SF phase up to a critical value of the attractive interaction, where PSF emerges. To locate the SF-PSF transition one can investigate the long distance behavior of the single particle and pair-pair correlation functions defined as

$$S(r) = \langle b_{x,y}^{\dagger} b_{x+r,y} \rangle \tag{2}$$

$$D(r) = \langle (b^{\dagger})_{x,y}^2 (b)_{x+r,y}^2 \rangle, \tag{3}$$

respectively ¹. In fact, one expects that in the SF phase both correlators exhibit algebraic decay, signaling the presence of quasi-long-range order. In contrast, in the PSF phase, one can argue

¹In (2) and (3), the y-th leg in the lattice is constant but can assume any value. However, since the model

that while S(r) exhibits exponential decay, indicating the suppression of single bosons motion, D(r) should maintain its polynomial decay [39].

These intuitive aspects reflect the specific behavior of two different energy gaps defined as:

$$\Delta_{\alpha=1,2}(L\times M) = E(N+\alpha;L\times M) + E(N-\alpha;L\times M) - 2E(N;L\times M)$$
 (4)

where $E(N; L \times M)$ is the ground state energy of a system of N particles on $L \times M$ sites. Specifically, one can find that, for a finite size system, $\Delta_2 > \Delta_1$ in SF while $\Delta_2 < \Delta_1$ in PSF [42]. On the other hand, in the thermodynamic limit one finds that $\Delta_2 = 0$ for any value of U, while $\Delta_1 > 0$ when entering the PSF regime and thus explaining the exponential decay of S(r).

2.1 Bosonization analysis

The 3-body constraint Bose-Hubbard model at filling one is directly connected to spin-1 XXZ chain with single ion anisotropy [10, 42, 60–62]. This aspect is at the core of the analytical treatment of (1) in 1D by means of a bosonization approach [63]. Specifically, by following [10, 64], the Hamiltonian (1) is first mapped into a spin-1 model, then each spin-1 variable is written as a sum of two spin-1/2, which in turn are mapped onto two spinless fermions via a Jordan Wigner transformation. The two corresponding fermionic ladder operators are bosonized in the standard way, introducing two bosonic fields $\phi_{1,2}(x)$, and their dual fields $\theta_{1,2}(x)$ such that $[\phi_{\alpha}(x), \theta_{\beta}(y)] = i\pi \delta_{\alpha\beta}\Theta(x-y)$, with $\alpha, \beta = 1, 2$ and $\Theta(x-y)$ being the Heaviside step function. Apart from a coupling term negligible in the low energy limit, the resulting Hamiltonian decouples when re-written in terms of the symmetric an anti-symmetric combinations of $\phi_{1,2}(x)$, namely $\phi_{\pm}(x) = \phi_1(x) \pm \phi_2(x)$ and $\theta_{\pm}(x) = (\theta_1(x) \pm \theta_2(x))/2$. Explicitly

$$H \to H_+ + H_- \quad , \tag{5}$$

with

$$H_{+} \equiv \frac{u_{+}}{2\pi} \int dx \left[K_{+} (\partial_{x} \theta_{+})^{2} + \frac{1}{K_{+}} (\partial_{x} \phi_{+})^{2} \right] + \frac{g}{(\pi a)^{2}} \int dx \cos(2\phi_{+})$$
 (6)

$$H_{-} \equiv \frac{u_{-}}{2\pi} \int dx \left[K_{-} (\partial_{x} \theta_{-})^{2} + \frac{1}{K_{-}} (\partial_{x} \phi_{-})^{2} \right] - \frac{g}{(\pi a)^{2}} \int dx \cos(2\phi_{-}) - \frac{t}{\pi a} \int dx \cos(2\theta_{-}).$$
 (7)

Here a is the lattice spacing, g=-aU/2 is the coupling constant, and $K_{\pm}=2(\sqrt{1\pm U/t\pi})^{-1}$ are the Luttinger parameters, while $u_{\pm}=2ta/K_{\pm}$ are the velocities in the \pm -channels. We recognize in H_{\pm} two sine-Gordon models where the condition U<0 implies $K_{+}>2$ and $K_{-}<2$. A renormalization group analysis [10] suggests that in this regime the field ϕ_{+} never pins, whereas ϕ_{-} (for $U<-2t\pi$) can pin to the value 0. The corresponding phase can be identified with the PSF phase.

On general grounds, the bosonization approach can also be exploited to associate to each pinning value of the bosonic fields an appropriate nonlocal operator whose expectation value becomes different from zero in the corresponding phase [11, 12, 17]. In particular, it was shown that the MI phase of the 1D (1) can be characterized by the non vanishing of the expectation value of the parity operator [10], namely $O_P(j) \equiv \prod_{x=1}^{j-1} \exp[i\pi(n_x - 1)]$, which in the continuous limit becomes $O_P(x) \sim \cos(2\phi_+(x))$. In the thermodynamic limit, its expectation value will remain finite when most sites are singly occupied, and holon (empty sites) doublon

is isotropic in both hopping and interaction, any y-th leg leads to the same effects. In one dimension, obviously y = 1.

(doubly occupies sites) fluctuations occur in correlated pairs.

Here we observe that when instead the field ϕ_- pins to the value 0 a different parity operator $O_p^{(o)}$ should become finite, defined as

$$O_{\mathbf{p}}^{(0)}(j) = \prod_{x=1}^{j-1} \exp[i\pi n_x]$$
 (8)

We name it *odd parity*, since its expectation value will be finite when fluctuations occur in the form of correlated pairs of single bosons. In fact, in the continuum limit it turns out to be connected to the bosonic field ϕ_{-} , i.e. the antisymmetric combination of fermionic fields $\phi_{1,2}$. Explicitly

$$O_{\rm p}^{(\rm o)}(x) \sim \cos(2\phi_{-}(x))$$
 , (9)

which expectation value is finite when $\phi_{-}(x) = 0$. Thus the latter should be finite in the whole PSF phase, and vanishing as soon as ϕ_{-} unpins. Figure (1)(a) shows a simple cartoon of the case when this quantity is nonvanishing in 1D.

When extending this analysis to the 2D limit, upon exploiting [13, 14, 53] one may introduce a brane odd parity to capture the PSF phase. This is the product of M odd parities for each x-value in the lattice. Explicitly

$$O_{\mathbf{p}}^{(0)}(j) = \prod_{x=1}^{j-1} \exp\left[i\pi \sum_{y=1}^{M} n_{x,y}\right]$$
 (10)

which is nonvanishing when in the disordered background of holons and doublons density fluctuations occur only in correlated pairs, as shown schematically in figure (1)(b).

3 Numerical results




Figure 2: Analysis of a 1D system (M=1) at $\bar{n}=1$ by performing iDMRG simulations with a maximum bond dimension of $\chi_{\rm max}=200$. The panels (a) and (b) present the analysis of the correlators in (2),(3) respectively. In the panel (c), the odd parity $O_{\rm p}^{\rm (o)}$ is evaluated.

We present in this section the results we obtained by performing iDMRG [65–67] simulations, which allows to accurately take into account the quantumness of a given system. We analyse first the system of a single chain (M=1) and then we extend our investigation to larger systems aiming to reach the 2D limit. Within this algorithm, we can either fix a priori

the particle density, working in the canonical ensemble, or tune the chemical potential, in a grand canonical framework.

Firstly, for both the 1D and 2D cases, we present a validation at fixed filling $\bar{n}=1$ of the effectiveness of exploiting the odd parity in estimating the SF-PSF transition, in comparison to the use of correlators S(r), D(r) as defined in (2, 3). Then we present our resulting phase diagrams in (μ, U) plane, shown in figure (1)(c,d). Next, for both the 1D and 2D cases, we further investigate the filling dependence of the SF-PSF phase transition by imposing $\bar{n} \neq 1$ throughout the transition.

3.1 Results in 1D




Figure 3: Analysis of the gaps defined in (4) by increasing the size of the chain L=60,80,120,200,300. One can see in (a) the single particle gap while in (b) the double particle gap. Only the first one remains finite by extrapolating the thermodynamic limit (TDL) by considering a second order polynomial in 1/L. We keep as maximum bond dimension $\chi_{\rm max}=200$.

Our investigation starts with the study of local correlators in (2) and (3) by fixing $\bar{n}=1$ and enforcing particle number conservation. As illustrated in figure (2)(a) and (b) and mentioned before, we find that strong attractions generate an exponential decay of S(r) while D(r) maintains its quasi-long range order for any U. This behavior can be compared with that of the odd parity, as shown in figure (2)(c). Here, $O_p^{(o)}$ shows a sharp and well defined transition point. Specifically, the odd parity becomes finite at a specific U_c which clearly does correspond to the one where the long-range behavior of S(r) starts to deviate from an algebraic decay. Our numerical approach makes it also possible to accurately complement this analysis by incorporating a study of the gaps $\Delta_{1,2}$. The thermodynamic extrapolation in figure (3) shows indeed that in the SF $\Delta_1 = \Delta_2 = 0$ while, as long as PSF takes place, $\Delta_2 = 0$ but Δ_1 becomes finite. Notably, the condition $\Delta_1 > 0$ turns out to be fulfilled exactly when $O_p^{(o)} > 0$ therefore unambiguously proving the validity and accurateness of our approach.

In order to further enforce our results, restricted so far to the canonical ensemble, we also inspect the odd parity in the grand canonical ensemble by tuning both U and μ . The resulting phase diagram is reported in figure (1)(c), based on the analysis of $O_p^{(o)}$. It confirms the presence of the PSF phase (in red), in which the odd parity order remains robust irrespectively to the particle density.

3.2 Results in 2D

The simulation of 2D systems using MPS algorithms can be challenging due to the efficiency of this method relying on covering a finite ratio of the system's entanglement entropy. How-




Figure 4: Analysis for a 2D system. The first two panels (a) and (b) present the the correlators in (2),(3) respectively, by increasing the onsite attractive interaction U < 0 for M = 6. Panel (c) presents the plot of the SF-PSF transition for 2D systems as a function of the number of legs M. The inset shows the extrapolation for a square 2D lattice of the critical interaction $U_c(2D) = -17.65 \pm 0.05$ imposing $\chi_{max} = 1500$. Similar to the 1D case, a comparison of the three plots reveals that, coinciding with the change in decay observed in S(r) the odd parity becomes finite.

ever, the entanglement entropy grows extensively according to the area law, diminishing the algorithm's efficiency [67,68]. In our iDMRG simulations, we address this challenge by considering an infinite size in the x-direction, achieved through the repetition of a finite unit cell in the y-direction. The use of periodic boundary conditions accelerates convergence to a 2D configuration. The resulting infinite cylinder system can be effectively mapped to a 1D system by following a 'snake-like' path, initially traversing in the y-direction.

We start again from the canonical ensemble by considering the different behaviours of the correlators (2) and (3) at $\bar{n}=1$. As in the previous case, figure (4)(a) and (b) shows that for a sufficiently negative U there is a change in the decay law for S(r) while D(r) conserves his long distance behavior. As before, this point signals the emergence of the PSF phase. Again, we show in figure (4)(c) this transition to coincide with the critical U_c where the odd parity becomes finite.

Then, by increasing M from 3 to 8, we evaluated the odd parity to extrapolate the critical interaction for SF-PSF transition in the C_4 symmetric 2D limit 2 , as illustrated in figure (4)(c). For M=6 we further release the particle number conservation and evaluate again the odd parity by varying the onsite attractive interaction U in the (μ, U) plane. The resulting phase diagram, which should be representative of the 2D limit, is shown in figure (1)(d): also in this case the presence of the PSF phase (in red) is in one-to-one correspondence with the region of finite odd parity. This is confined between the SF phase (in blue), and the two trivial phases (in white) corresponding to the $\bar{n}=0$ and $\bar{n}=2$ cases.

3.3 Arbitrary Filling

The previously discussed results for 1D and 2D systems focused first on the canonical ensemble, where we enforced a filling of $\bar{n}=1$. Subsequently, such condition was released upon exploring the behavior of the system in the grand canonical ensemble, where the number of bosons is not necessarily constant throughout the SF-PSF phase transitions, and we obtained the (μ, U) 1D and 2D phase diagrams. To confirm the picture, we now inspect the SF-PSF transition again in the canonical ensemble, for fixed fillings $\bar{n} \neq 1$.

²We consider the case M = 6 as a reliable approximation of a 2D system, given that the difference between this case and our resulting limit is less than 3%.




Figure 5: Evaluation of the odd parity $O_{\rm p}^{\rm (o)}$ for a fixed $\bar{n} \neq 1$ for the 1D system in (a) and 2D in (b), both by performing iDMRG simulations for $\chi_{\rm max} = 200,1500$ respectively. The $\bar{n}=1$ case is included for comparison with previous results. As evident from figures, for the considered densities there is no change in the nature of the transition when moving away from filling one.

The results are shown in figure (5) for a 1D (a) and a 2D (b) system. There the odd parity (8) was evaluated for fillings greater and smaller than one. The SF and PSF phases are again distinguished through the behavior of odd parity: in the first, the odd parity is zero, whereas in the other the emergence of a nonlocal order is confirmed by its non zero value. A careful analysis reveals how at all fillings the transition is second order, in both 1D and 2D cases³. Moreover, in this way the actual dependence of the critical value of interaction U_c on filling is obtained.

3.4 Two-species system

The presence of the SF-PSF phase transition we discussed in previous sections is not limited to the 3-body constrained BHM. For instance, it is observed also in case of two components hard-core bosons in optical lattice [54–59]. Although, the regime of PSF has not yet been observed, Bose mixtures in optical lattice have been the subject of intense experimental activity [69–76]. The Hamiltonian we are interested in reads

$$H = -t \sum_{i,\sigma=A,B} (b_{i,\sigma}^{\dagger} b_{i+1,\sigma} + \text{h.c.}) + U \sum_{i} n_{i,A} n_{i,B}$$
 (11)

where i is a generic site of the chain for boson species $\sigma = A, B$. For attractive interchain on-site interaction, the system should show a tendency to create pairs of A-B bosons.

As before we can measure the emergence of the odd parity order. The previous definition (8) in this case becomes

$$O_{\rm p}^{(\rm o)}(j) = \prod_{i=1}^{j-1} \exp\left[\sum_{\sigma={\rm A},{\rm B}} i\pi n_{i,\sigma}\right].$$
 (12)

Now the same site on the two chains is either empty or occupied in both chains, whereas the fluctuations occur in the form of correlated pairs of bosons of the different species as depicted in figure (6)(a).

In agreement with results in [59,77], and with bosonization analysis in the fermionic case [11], $O_p^{(o)}$ becomes finite as soon as U < 0, as shown in figure (6)(b).

³This is at variance with the case of fixed chemical potential, in which traversing a transition in the (μ, U) plane may result in the total number of bosons not being conserved, leading to abrupt filling changes. This discontinuity can also manifest in odd parity, transitioning from zero to a finite value.




Figure 6: (a) Schematic representation of the two hard-core bosons chains model. One can see the direct connection to the previous representation in figure (1)(a,b). Now the couple of bosons are composed by A (in blue) and B (in green) bosons. (b) Evaluation of the odd parity in (12) for SF-PSF phase transition. Differently from previous case, the transition occurs for small negative interaction.

4 Conclusions

We unveiled the order parameter of the exotic pair superfluid phase. The latter takes the form of a nonlocal parity operator named odd parity which can be accurately detected through local density measurements. Such quantity is finite whenever in a disordered background of holons and doublons single bosons occur in correlated pairs. The prediction is confirmed by a bosonization analysis and extensive numerical simulations.

Relevantly, we have shown our results to hold both in one and two spatial dimensions, at any density supporting a pair superfluid phase, and in two different models where such state of matter takes place. This aspect confirms the generality and the large range of applicability of our findings. Moreover, we proved the aforementioned points through extensive matrix-product-state simulations where quantum fluctuation are accurately taken into account. This makes our results highly robust and reliable in deep quantum regimes where mean-field approaches are not reliable.

We further stress that a major advantage of nonlocal operators is that they do not break continuous Hamiltonian symmetries, thus describing orders that can persist in low dimensional systems even at non vanishing temperatures. Hence the measurement of odd parity order is even more at hand in present experimental setups, especially because it can be obtained by means of just local density probes. In conclusion, we believe that our results represent an important step towards a more complete and accurate characterization of strongly correlated many-body quantum systems.

Acknowledgments

Funding information N.C. and A.M. thank for the financial support from the ICSC – Centro Nazionale di Ricerca in High Performance Computing, Big Data and Quantum Computing, funded by European Union – NextGenerationEU (Grant number CN00000013). L. B. acknowledges financial support within the DiQut Grant No.2022523NA7 funded by European Union – Next Generation EU, PRIN 2022 program (D.D. 104 - 02/02/2022 Ministero dell'Università e della Ricerca).

Computational resources were provided by HPC@POLITO (http://www.hpc.polito.it).

Calculations were performed using the TeNPy Library (version 0.10.0) [66].

References

- [1] L. D. Landau, E. M. Lifshitz and M. Pitaevskii, *Statistical Physics*, Butterworth-Heinemann, New York (1999).
- [2] K. G. Wilson and J. Kogut, *The renormalization group and the \epsilon expansion*, Phys. Rep. **12**(2), 75 (1974), doi:https://doi.org/10.1016/0370-1573(74)90023-4.
- [3] A. Altland and M. R. Zirnbauer, *Nonstandard symmetry classes in mesoscopic normal-superconducting hybrid structures*, Phys. Rev. B **55**, 1142 (1997), doi:10.1103/PhysRevB.55.1142.
- [4] A. Kitaev, *Periodic table for topological insulators and superconductors*, AIP Conference Proceedings **1134**(1), 22 (2009), doi:10.1063/1.3149495, https://pubs.aip.org/aip/acp/article-pdf/1134/1/22/11584243/22_1_online.pdf.
- [5] X.-G. Wen, *Symmetry-protected topological phases in noninteracting fermion systems*, Phys. Rev. B **85**, 085103 (2012), doi:10.1103/PhysRevB.85.085103.
- [6] F. Pollmann, E. Berg, A. M. Turner and M. Oshikawa, Symmetry protection of topological phases in one-dimensional quantum spin systems, Phys. Rev. B 85, 075125 (2012), doi:10.1103/PhysRevB.85.075125.
- [7] X. Chen, Z.-C. Gu, Z.-X. Liu and X.-G. Wen, Symmetry protected topological orders and the group cohomology of their symmetry group, Phys. Rev. B 87, 155114 (2013), doi:10.1103/PhysRevB.87.155114.
- [8] T. Giamarchi, Quantum Physics in One Dimension, Oxford University Press, ISBN 0198525001 (2004).
- [9] N. D. Mermin and H. Wagner, Absence of ferromagnetism or antiferromagnetism in one- or two-dimensional isotropic heisenberg models, Phys. Rev. Lett. 17, 1133 (1966), doi:10.1103/PhysRevLett.17.1133.
- [10] E. Berg, E. G. Dalla Torre, T. Giamarchi and E. Altman, *Rise and fall of hidden string order of lattice bosons*, Phys. Rev. B **77**, 245119 (2008), doi:10.1103/PhysRevB.77.245119.
- [11] A. Montorsi and M. Roncaglia, *Nonlocal order parameters for the 1d hubbard model*, Phys. Rev. Lett. **109**, 236404 (2012), doi:10.1103/PhysRevLett.109.236404.
- [12] L. Barbiero, A. Montorsi and M. Roncaglia, *How hidden orders generate gaps in one-dimensional fermionic systems*, Phys. Rev. B **88**, 035109 (2013), doi:10.1103/PhysRevB.88.035109.
- [13] S. P. Rath, W. Simeth, M. Endres and W. Zwerger, *Non-local order in mott insulators, duality and wilson loops*, Annals of Physics **334**, 256 (2013), doi:https://doi.org/10.1016/j.aop.2013.04.006.
- [14] S. Fazzini, F. Becca and A. Montorsi, *Nonlocal parity order in the two-dimensional mott insulator*, Phys. Rev. Lett. **118**, 157602 (2017), doi:10.1103/PhysRevLett.118.157602.

[15] M. den Nijs and K. Rommelse, Preroughening transitions in crystal surfaces and valence-bond phases in quantum spin chains, Phys. Rev. B **40**, 4709 (1989), doi:10.1103/PhysRevB.40.4709.

- [16] S. R. Manmana, A. M. Essin, R. M. Noack and V. Gurarie, *Topological invariants and interacting one-dimensional fermionic systems*, Phys. Rev. B **86**, 205119 (2012), doi:10.1103/PhysRevB.86.205119.
- [17] A. Montorsi, F. Dolcini, R. C. Iotti and F. Rossi, *Symmetry-protected topological phases of one-dimensional interacting fermions with spin-charge separation*, Phys. Rev. B **95**, 245108 (2017), doi:10.1103/PhysRevB.95.245108.
- [18] A. Mazurenko, C. S. Chiu, G. Ji, M. F. Parsons, M. Kanász-Nagy, R. Schmidt, F. Grusdt, E. Demler, D. Greif and M. Greiner, *A cold-atom fermi–hubbard antiferromagnet*, Nature **545**(7655), 462 (2017), doi:10.1038/nature22362.
- [19] L. Tanzi, S. M. Roccuzzo, E. Lucioni, F. Famà, A. Fioretti, C. Gabbanini, G. Modugno, A. Recati and S. Stringari, *Supersolid symmetry breaking from compressional oscillations in a dipolar quantum gas*, Nature **574**(7778), 382 (2019), doi:https://doi.org/10.1038/s41586-019-1568-6.
- [20] M. Guo, F. Böttcher, J. Hertkorn, J.-N. Schmidt, M. Wenzel, H. P. Büchler, T. Langen and T. Pfau, *The low-energy goldstone mode in a trapped dipolar supersolid*, Nature **574**(7778), 386 (2019), doi:https://doi.org/10.1038/s41586-019-1569-5.
- [21] L. Chomaz, D. Petter, P. Ilzhöfer, G. Natale, A. Trautmann, C. Politi, G. Durastante, R. M. W. van Bijnen, A. Patscheider, M. Sohmen, M. J. Mark and F. Ferlaino, Long-lived and transient supersolid behaviors in dipolar quantum gases, Phys. Rev. X 9, 021012 (2019), doi:10.1103/PhysRevX.9.021012.
- [22] H. P. Zahn, V. P. Singh, M. N. Kosch, L. Asteria, L. Freystatzky, K. Sengstock, L. Mathey and C. Weitenberg, *Formation of spontaneous density-wave patterns in dc driven lattices*, Phys. Rev. X **12**, 021014 (2022), doi:10.1103/PhysRevX.12.021014.
- [23] L. Su, A. Douglas, M. Szurek, R. Groth, S. F. Ozturk, A. Krahn, A. H. Hébert, G. A. Phelps, S. Ebadi, S. Dickerson, F. Ferlaino, O. Marković et al., Dipolar quantum solids emerging in a hubbard quantum simulator, Nature 622(7984), 724 (2023), doi:10.1038/s41586-023-06614-3.
- [24] H.-J. Shao, Y.-X. Wang, D.-Z. Zhu, Y.-S. Zhu, H.-N. Sun, S.-Y. Chen, C. Zhang, Z.-J. Fan, Y. Deng, X.-C. Yao, Y.-A. Chen and J.-W. Pan, *Observation of the antiferromagnetic phase transition in the fermionic hubbard model* (2024), 2402.14605.
- [25] E. Haller, R. Hart, M. J. Mark, J. G. Danzl, L. Reichsöllner, M. Gustavsson, M. Dalmonte, G. Pupillo and H.-C. Nägerl, *Pinning quantum phase transition for a luttinger liquid of strongly interacting bosons*, Nature **466**(7306), 597 (2010), doi:10.1038/nature09259.
- [26] M. Endres, M. Cheneau, T. Fukuhara, C. Weitenberg, P. Schauß, C. Gross, L. Mazza, M. C. Bañuls, L. Pollet, I. Bloch and S. Kuhr, Observation of correlated particle-hole pairs and string order in low-dimensional mott insulators, Science 334(6053), 200 (2011), doi:10.1126/science.1209284, https://www.science.org/doi/pdf/10.1126/science.1209284.
- [27] D. Wei, D. Adler, K. Srakaew, S. Agrawal, P. Weckesser, I. Bloch and J. Zeiher, *Observation of brane parity order in programmable optical lattices*, Phys. Rev. X **13**, 021042 (2023), doi:10.1103/PhysRevX.13.021042.

[28] J. Hur, W. Lee, K. Kwon, S. Huh, G. Y. Cho and J.-y. Choi, *Measuring nonlocal brane order with error-corrected quantum gas microscopes*, Phys. Rev. X **14**, 011003 (2024), doi:10.1103/PhysRevX.14.011003.

- [29] S. de Léséleuc, V. Lienhard, P. Scholl, D. Barredo, S. Weber, N. Lang, H. P. Büchler, T. Lahaye and A. Browaeys, *Observation of a symmetry-protected topological phase of interacting bosons with rydberg atoms*, Science **365**(6455), 775 (2019), doi:10.1126/science.aav9105, https://www.science.org/doi/pdf/10.1126/science.aav9105.
- [30] P. Sompet, S. Hirthe, D. Bourgund, T. Chalopin, J. Bibo, J. Koepsell, P. Bojović, R. Verresen, F. Pollmann, G. Salomon, C. Gross, T. A. Hilker et al., Realizing the symmetry-protected haldane phase in fermi-hubbard ladders, Nature 606(7914), 484 (2022), doi:10.1038/s41586-022-04688-z.
- [31] M. Lewenstein, A. Sanpera, V. Ahufinger, B. Damski, A. Sen(De) and U. Sen, *Ultracold atomic gases in optical lattices: mimicking condensed matter physics and beyond*, Adv. Phys. **56**(2), 243 (2007), doi:10.1080/00018730701223200.
- [32] C. Gross and I. Bloch, *Quantum simulations with ultracold atoms in optical lattices*, Science **357**(6355), 995 (2017), doi:10.1126/science.aal3837.
- [33] E. Altman, E. Demler and M. D. Lukin, *Probing many-body states of ultracold atoms via noise correlations*, Phys. Rev. A **70**, 013603 (2004), doi:10.1103/PhysRevA.70.013603.
- [34] J. Argüello-Luengo, S. Julià-Farré, M. Lewenstein, C. Weitenberg and L. Barbiero, *Probing spontaneously symmetry-broken phases with spin-charge separation through noise correlation measurements* (2024), 2404.08374.
- [35] C. Gross and W. S. Bakr, *Quantum gas microscopy for single atom and spin detection*, Nat. Phys. **17**(12), 1316 (2021), doi:10.1038/s41567-021-01370-5.
- [36] S. Fazzini, L. Barbiero and A. Montorsi, *Interaction-induced fractionalization and topological superconductivity in the polar molecules anisotropic t j model*, Phys. Rev. Lett. **122**, 106402 (2019), doi:10.1103/PhysRevLett.122.106402.
- [37] L. F. Tocchio, F. Becca and A. Montorsi, *Superconductivity in the Hubbard model: a hidden-order diagnostics from the Luther-Emery phase on ladders*, SciPost Phys. **6**, 018 (2019), doi:10.21468/SciPostPhys.6.2.018.
- [38] A. Montorsi, S. Fazzini and L. Barbiero, *Homogeneous and domain-wall topological haldane conductors with dressed rydberg atoms*, Phys. Rev. A **101**, 043618 (2020), doi:10.1103/PhysRevA.101.043618.
- [39] A. J. Daley, J. M. Taylor, S. Diehl, M. Baranov and P. Zoller, *Atomic three-body loss as a dynamical three-body interaction*, Phys. Rev. Lett. **102**, 040402 (2009), doi:10.1103/PhysRevLett.102.040402.
- [40] L. Mazza, M. Rizzi, M. Lewenstein and J. I. Cirac, *Emerging bosons with three-body interactions from spin-1 atoms in optical lattices*, Phys. Rev. A **82**, 043629 (2010), doi:10.1103/PhysRevA.82.043629.
- [41] S. Diehl, M. Baranov, A. J. Daley and P. Zoller, *Observability of quantum criticality and a continuous supersolid in atomic gases*, Phys. Rev. Lett. **104**, 165301 (2010), doi:10.1103/PhysRevLett.104.165301.

[42] M. Dalmonte, M. Di Dio, L. Barbiero and F. Ortolani, *Homogeneous and inhomogeneous magnetic phases of constrained dipolar bosons*, Phys. Rev. B **83**, 155110 (2011), doi:10.1103/PhysRevB.83.155110.

- [43] Y.-W. Lee and M.-F. Yang, Superfluid-insulator transitions in attractive bose-hubbard model with three-body constraint, Phys. Rev. A 81, 061604 (2010), doi:10.1103/PhysRevA.81.061604.
- [44] L. Bonnes and S. Wessel, *Pair superfluidity of three-body constrained bosons in two dimensions*, Phys. Rev. Lett. **106**, 185302 (2011), doi:10.1103/PhysRevLett.106.185302.
- [45] A. Perali, D. Neilson and A. R. Hamilton, *High-temperature superfluidity in double-bilayer graphene*, Phys. Rev. Lett. **110**, 146803 (2013), doi:10.1103/PhysRevLett.110.146803.
- [46] M. Singh, T. Mishra, R. V. Pai and B. P. Das, *Quantum phases of attractive bosons on a bose-hubbard ladder with three-body constraint*, Phys. Rev. A **90**, 013625 (2014), doi:10.1103/PhysRevA.90.013625.
- [47] A. Macia, G. E. Astrakharchik, F. Mazzanti, S. Giorgini and J. Boronat, *Single-particle versus pair superfluidity in a bilayer system of dipolar bosons*, Phys. Rev. A **90**, 043623 (2014), doi:10.1103/PhysRevA.90.043623.
- [48] G. E. Astrakharchik, R. E. Zillich, F. Mazzanti and J. Boronat, *Gapped spectrum in pair-superfluid bosons*, Phys. Rev. A **94**, 063630 (2016), doi:10.1103/PhysRevA.94.063630.
- [49] T. Bilitewski and N. R. Cooper, *Synthetic dimensions in the strong-coupling limit: Supersolids and pair superfluids*, Phys. Rev. A **94**, 023630 (2016), doi:10.1103/PhysRevA.94.023630.
- [50] C. Hubert, Y. Baruchi, Y. Mazuz-Harpaz, K. Cohen, K. Biermann, M. Lemeshko, K. West, L. Pfeiffer, R. Rapaport and P. Santos, *Attractive dipolar coupling between stacked exciton fluids*, Phys. Rev. X **9**, 021026 (2019), doi:10.1103/PhysRevX.9.021026.
- [51] M. Zimmerman, R. Rapaport and S. Gazit, *Collective interlayer pairing and pair superfluidity in vertically stacked layers of dipolar excitons*, Proceedings of the National Academy of Sciences **119**(30), e2205845119 (2022), doi:10.1073/pnas.2205845119, https://www.pnas.org/doi/pdf/10.1073/pnas.2205845119.
- [52] A. Padhan, R. Parida, S. Lahiri, M. K. Giri and T. Mishra, *Quantum phases of constrained bosons on a two-leg bose-hubbard ladder*, Phys. Rev. A **108**, 013316 (2023), doi:10.1103/PhysRevA.108.013316.
- [53] C. Degli Esposti Boschi, A. Montorsi and M. Roncaglia, *Brane parity orders in the insulating state of hubbard ladders*, Phys. Rev. B **94**, 085119 (2016), doi:10.1103/PhysRevB.94.085119.
- [54] B. Paredes and J. I. Cirac, From cooper pairs to luttinger liquids with bosonic atoms in optical lattices, Phys. Rev. Lett. **90**, 150402 (2003), doi:10.1103/PhysRevLett.90.150402.
- [55] A. Kuklov, N. Prokof'ev and B. Svistunov, Superfluid-superfluid phase transitions in a two-component bose-einstein condensate, doi:10.1103/PhysRevLett.92.030403.
- [56] A. Kuklov, N. Prokof'ev and B. Svistunov, *Commensurate two-component bosons in an optical lattice: Ground state phase diagram*, Phys. Rev. Lett. **92**, 050402 (2004), doi:10.1103/PhysRevLett.92.050402.

[57] A. Hu, L. Mathey, I. Danshita, E. Tiesinga, C. J. Williams and C. W. Clark, *Counterflow and paired superfluidity in one-dimensional bose mixtures in optical lattices*, Phys. Rev. A **80**, 023619 (2009), doi:10.1103/PhysRevA.80.023619.

- [58] A. Hu, L. Mathey, C. J. Williams and C. W. Clark, Noise correlations of onedimensional bose mixtures in optical lattices, Phys. Rev. A 81, 063602 (2010), doi:10.1103/PhysRevA.81.063602.
- [59] B. Grémaud and G. G. Batrouni, *Pairing and pair superfluid density in one-dimensional two-species fermionic and bosonic hubbard models*, Phys. Rev. Lett. **127**, 025301 (2021), doi:10.1103/PhysRevLett.127.025301.
- [60] W. Chen, K. Hida and B. C. Sanctuary, *Ground-state phase diagram of s* = 1 XXZ *chains with uniaxial single-ion-type anisotropy*, Phys. Rev. B **67**, 104401 (2003), doi:10.1103/PhysRevB.67.104401.
- [61] E. G. Dalla Torre, E. Berg and E. Altman, *Hidden order in 1d bose insulators*, Phys. Rev. Lett. **97**, 260401 (2006), doi:10.1103/PhysRevLett.97.260401.
- [62] S. Ejima and H. Fehske, Comparative density-matrix renormalization group study of symmetry-protected topological phases in spin-1 chain and bose-hubbard models, Phys. Rev. B **91**, 045121 (2015), doi:10.1103/PhysRevB.91.045121.
- [63] T. Giamarchi, *Quantum Physics in One Dimension*, Oxford University Press, ISBN 9780198525004, doi:10.1093/acprof:oso/9780198525004.001.0001 (2003).
- [64] H. J. Schulz, *Phase diagrams and correlation exponents for quantum spin chains of arbitrary spin quantum number*, Phys. Rev. B **34**, 6372 (1986), doi:10.1103/PhysRevB.34.6372.
- [65] I. P. McCulloch, *Infinite size density matrix renormalization group, revisited* (2008), 0804. 2509.
- [66] J. Hauschild and F. Pollmann, *Efficient numerical simulations with Tensor Networks: Tensor Network Python (TeNPy)*, SciPost Phys. Lect. Notes p. 5 (2018), doi:10.21468/SciPostPhysLectNotes.5, 1805.00055.
- [67] U. Schollwöck, *The density-matrix renormalization group in the age of matrix product states*, Annals of Physics **326**(1), 96 (2011), doi:https://doi.org/10.1016/j.aop.2010.09.012, January 2011 Special Issue.
- [68] J. Eisert, M. Cramer and M. B. Plenio, *Colloquium: Area laws for the entanglement entropy*, Rev. Mod. Phys. **82**, 277 (2010), doi:10.1103/RevModPhys.82.277.
- [69] J. Catani, L. De Sarlo, G. Barontini, F. Minardi and M. Inguscio, Degenerate bose-bose mixture in a three-dimensional optical lattice, Phys. Rev. A 77, 011603 (2008), doi:10.1103/PhysRevA.77.011603.
- [70] S. Trotzky, P. Cheinet, S. Fölling, M. Feld, U. Schnorrberger, A. M. Rey, A. Polkovnikov, E. A. Demler, M. D. Lukin and I. Bloch, *Time-resolved observation and control of su-perexchange interactions with ultracold atoms in optical lattices*, Science 319(5861), 295 (2008), doi:10.1126/science.1150841, https://www.science.org/doi/pdf/10.1126/science.1150841.

[71] T. Best, S. Will, U. Schneider, L. Hackermüller, D. van Oosten, I. Bloch and D.-S. Lühmann, *Role of interactions in* ⁸⁷Rb—⁴⁰**K** *bose-fermi mixtures in a 3d optical lattice*, Phys. Rev. Lett. **102**, 030408 (2009), doi:10.1103/PhysRevLett.102.030408.

- [72] D. M. Weld, H. Miyake, P. Medley, D. E. Pritchard and W. Ketterle, *Thermometry and refrigeration in a two-component mott insulator of ultracold atoms*, Phys. Rev. A **82**, 051603 (2010), doi:10.1103/PhysRevA.82.051603.
- [73] J. Amato-Grill, N. Jepsen, I. Dimitrova, W. Lunden and W. Ketterle, *Interaction spectroscopy of a two-component mott insulator*, Phys. Rev. A **99**, 033612 (2019), doi:10.1103/PhysRevA.99.033612.
- [74] P. N. Jepsen, J. Amato-Grill, I. Dimitrova, W. W. Ho, E. Demler and W. Ketterle, *Spin transport in a tunable heisenberg model realized with ultracold atoms*, Nature **588**(7838), 403 (2020), doi:10.1038/s41586-020-3033-y.
- [75] W. C. Chung, J. de Hond, J. Xiang, E. Cruz-Colón and W. Ketterle, *Tunable single-ion anisotropy in spin-1 models realized with ultracold atoms*, Phys. Rev. Lett. **126**, 163203 (2021), doi:10.1103/PhysRevLett.126.163203.
- [76] P. N. Jepsen, W. W. Ho, J. Amato-Grill, I. Dimitrova, E. Demler and W. Ketterle, *Transverse spin dynamics in the anisotropic heisenberg model realized with ultracold atoms*, Phys. Rev. X 11, 041054 (2021), doi:10.1103/PhysRevX.11.041054.
- [77] A. Safavi-Naini, B. Capogrosso-Sansone and A. Kuklov, *Quantum phases of hard-core dipolar bosons in coupled one-dimensional optical lattices*, Phys. Rev. A **90**, 043604 (2014), doi:10.1103/PhysRevA.90.043604.