

EXPONENTIAL DECAY FOR FRACTIONAL SCHRÖDINGER PARABOLIC PROBLEMS

JAN W. CHOLEWA¹ AND ANIBAL RODRIGUEZ-BERNAL²

¹Institute of Mathematics
University of Silesia in Katowice
40-007 Katowice, Poland
E-mail: jan.cholewa@us.edu.pl

²Departamento de Análisis Matemático y Matemática Aplicada
Universidad Complutense de Madrid
28040 Madrid, Spain
and
Instituto de Ciencias Matemáticas
CSIC-UAM-UC3M-UCM*, Spain
E-mail: arober@ucm.es

ABSTRACT. We discuss exponential decay in $L^p(\mathbb{R}^N)$, $1 \leq p \leq \infty$, of solutions of a fractional Schrödinger parabolic equation with a locally uniformly integrable potential. The exponential type of the semigroup of solutions is considered and its dependence in $1 \leq p \leq \infty$ is addressed. We characterise a large class of potentials for which solutions decay exponentially.

1. INTRODUCTION

In this paper we discuss exponential decay of solutions of fractional Schrödinger semigroups

$$\begin{cases} u_t + (-\Delta)^\mu u + V(x)u = 0, & x \in \mathbb{R}^N, t > 0 \\ u(x, 0) = u_0(x), & x \in \mathbb{R}^N \end{cases} \quad (1.1)$$

with $0 < \mu \leq 1$, $u_0 \in L^p(\mathbb{R}^N)$ with $1 \leq p \leq \infty$.

The nonnegative potential $V \geq 0$ belongs to the locally uniform space $L_U^{p_0}(\mathbb{R}^N)$, which, for $1 \leq p_0 < \infty$, is composed of the functions $V \in L_{loc}^{p_0}(\mathbb{R}^N)$ such that there exists $C > 0$ such that for all $x_0 \in \mathbb{R}^N$

$$\int_{B(x_0, 1)} |V|^{p_0} \leq C$$

endowed with the norm

$$\|V\|_{L_U^{p_0}(\mathbb{R}^N)} = \sup_{x_0 \in \mathbb{R}^N} \|V\|_{L^{p_0}(B(x_0, 1))}.$$

For $p_0 = \infty$ we define $L_U^\infty(\mathbb{R}^N) = L^\infty(\mathbb{R}^N)$.

For $\mu = 1$ and potentials $0 \leq V \in L^1(\mathbb{R}^N) + L^\infty(\mathbb{R}^N)$ it was proved in [1] that the exponential decay in $L^p(\mathbb{R}^N)$ of solutions of (1.1) holds iff V is sufficiently positive at infinity in the sense that

$$\int_G V(x) dx = \infty \quad (1.2)$$

for any open set $G \subset \mathbb{R}^N$ containing arbitrarily large balls, that is, such that for any $r > 0$ there exists $x_0 \in \mathbb{R}^N$ such that the ball of radius r around x_0 is included in G .

Date: May 15, 2024.

Partially supported by Projects PID2019-103860GB-I00 and PID2022-137074NB-I00, MINECO, Spain.

Key words and phrases: fractional Schrödinger operator, linear evolution problem, stability.

Mathematical Subject Classification 2020: 35J10, 35R11, 47D06, 35B35.

*Partially supported by Severo Ochoa Grant CEX2019-000904-S funded by MCIN/AEI/10.13039/ 501100011033.

To extend this result to solutions of (1.1) with $0 < \mu \leq 1$ and $0 \leq V \in L_U^{p_0}(\mathbb{R}^N)$ we first prove in Section 3 that if $p_0 > \max\{\frac{N}{2\mu}, 1\}$ then for any $1 \leq p \leq \infty$, (1.1) is well posed in $L^p(\mathbb{R}^N)$ and that it defines a contraction semigroup of solutions $u(t, u_0) = S_{\mu, V}(t)u_0$. This semigroup is order preserving, strongly continuous if $1 \leq p < \infty$ and analytic if $1 < p < \infty$, see Proposition 3.1. Moreover, for smooth initial data, solutions can be represented in terms of the fractional unperturbed semigroup (that is, when $V = 0$) and the variation of constants formula. Also, for nonnegative initial data, the larger the potential, the smaller the solution is, see Proposition 3.3. If $V \in L^\infty(\mathbb{R}^N)$ then the variation of constants formula applies to all initial data in $L^p(\mathbb{R}^N)$, see Proposition 3.4, and therefore it is natural to consider potentials $0 \leq V \in L_U^{p_0}(\mathbb{R}^N)$ that can be approximated in $L_U^{p_0}(\mathbb{R}^N)$ by bounded ones and to analyze the convergence of the corresponding semigroups, see Proposition 3.5 and Theorem 3.7.

With these tools, in Section 4 we address the exponential decay of solutions of (1.1) in $L^p(\mathbb{R}^N)$ with $1 \leq p \leq \infty$ for potentials $0 \leq V \in L_U^{p_0}(\mathbb{R}^N)$ satisfying (1.2). For this, we first analyze how the exponential decay, actually, the exponential type of the semigroup $\{S_{\mu, V}(t)\}_{t \geq 0}$, depends on $1 \leq p \leq \infty$. We prove that if $0 < \mu \leq 1$ then the exponential type is either zero or it is negative for all $1 \leq p \leq \infty$ simultaneously. If $\mu = 1$ we recover the result that the exponential type is independent of $1 \leq p \leq \infty$, see Theorem 4.1. Then we prove that if solutions decay exponentially then (1.2) holds. Conversely, if the potential can be approximated by bounded ones, then (1.2) actually characterises the exponential decay, see Theorem 4.9.

Finally, in Section 2 we have included several known results for the unperturbed problem (1.1) with $V = 0$.

In this paper we denote by c or C generic constants that may change from line to line, whose value is not important for the results.

Also we will denote $A \sim B$ to denote quantities (norms or functions, for example) such that there exist positive constants c_1, c_2 such that $c_1 A \leq B \leq c_2 A$. Similarly we use $A \lesssim B$ to indicate $A \leq cB$.

2. BASIC RESULTS ON FRACTIONAL OPERATORS AND SEMIGROUPS

In this section we review several known results for the case of the unperturbed equation, that is, (1.1) with $V = 0$.

First, for the standard heat equation, $\mu = 1$, we have that for a wide class of initial data, including, $L^p(\mathbb{R}^N)$ for $1 \leq p \leq \infty$, the solution of (1.1) is given by

$$S(t)u_0(x) = \int_{\mathbb{R}^N} k(t, x, y)u_0(y) dy \quad t > 0, x \in \mathbb{R}^N, \quad (2.1)$$

for the selfsimilar convolution kernel

$$0 \leq k(t, x, y) = \frac{1}{t^{\frac{N}{2}}} k_0\left(\frac{x-y}{t^{\frac{1}{2}}}\right) = \frac{1}{(4\pi t)^{\frac{N}{2}}} e^{-\frac{|x-y|^2}{4t}}. \quad (2.2)$$

In particular, (2.1) defines an order preserving semigroup of contractions in $L^p(\mathbb{R}^N)$, $1 \leq p \leq \infty$ and

$$\|S(t)\|_{\mathcal{L}(L^p(\mathbb{R}^N))} = 1, \quad t > 0.$$

This semigroup is strongly continuous if $1 \leq p < \infty$ and analytic if $1 < p < \infty$. In particular, if $1 \leq p < \infty$ then for $u_0 \in L^p(\mathbb{R}^N)$ we have

$$S(t)u_0 \rightarrow u_0 \quad \text{as } t \rightarrow 0^+ \quad \text{in } L^p(\mathbb{R}^N) \text{ for any } 1 \leq p < \infty.$$

If $p = \infty$ then for $u_0 \in L^\infty(\mathbb{R}^N)$

$$S(t)u_0 \rightarrow u_0 \quad \text{as } t \rightarrow 0^+ \quad \text{in } L_{loc}^p(\mathbb{R}^N) \text{ for any } 1 \leq p < \infty. \quad (2.3)$$

For all these classical results see e.g. [7].

Finally, because of the regularity of the kernel, we have that for $1 \leq p \leq \infty$ and $u_0 \in L^p(\mathbb{R}^N)$, $u(t) = S(t)u_0$ in (2.1) is of class $C^\infty((0, \infty) \times \mathbb{R}^N)$ and satisfies the heat equation pointwise in $(0, \infty) \times \mathbb{R}^N$.

The semigroup of solutions possesses also several *smoothing* estimates among which we recall that for $1 \leq p \leq q \leq \infty$ there exists a constant $c_{p,q} > 0$ such that

$$\|S(t)\|_{\mathcal{L}(L^p(\mathbb{R}^N), L^q(\mathbb{R}^N))} = \frac{c_{p,q}}{t^{\frac{N}{2}(\frac{1}{p}-\frac{1}{q})}}, \quad t > 0.$$

The fractional powers of $-\Delta$ can be defined in many equivalent ways, see [18]. From the heat semigroup (2.1) it can be defined as

$$(-\Delta)^\mu \phi(x) = \frac{1}{\Gamma(-\mu)} \int_0^\infty (S(t)\phi(x) - \phi(x)) \frac{dt}{t^{1+\mu}}$$

for $0 < \mu < 1$, which coincides with the nonlocal operator defined by

$$(-\Delta)^\mu \phi(x) = C_{N,\mu} \text{ P.V. } \int_{\mathbb{R}^N} \frac{\phi(x) - \phi(z)}{|x - z|^{N+2\mu}} dz \quad (2.4)$$

with $C_{N,\mu} = \frac{2^{2\mu} \mu \Gamma(\frac{N}{2} + \mu)}{\pi^{N/2} \Gamma(1 - \mu)}$, see [13]. When considered in $L^p(\mathbb{R}^N)$ for $1 \leq p \leq \infty$, the domain of $(-\Delta)^\mu$ is the set of functions $\phi \in L^p(\mathbb{R}^N)$ such that $(-\Delta)^\mu \phi \in L^p(\mathbb{R}^N)$ with the natural norm

$$\|\phi\|_{L^p} + \|(-\Delta)^\mu \phi\|_{L^p},$$

see [8, Section A.2] when $1 \leq p < \infty$ and [9, Appendix A] when $p = \infty$. For $1 < p < \infty$ this domain coincides with the Bessel space $H_p^{2\mu}(\mathbb{R}^N)$. When $p = 2$ these spaces will be denoted by $H^{2\mu}(\mathbb{R}^N)$.

Using [8, Lemma 6.2] we have in particular, that for $1 < p < \infty$,

$$\|(-\Delta)^{\frac{1}{2}} \phi\|_{L^p(\mathbb{R}^N)} \sim \sum_{j=1}^N \|\partial_j \phi\|_{L^p(\mathbb{R}^N)} \quad \text{in } H_p^1(\mathbb{R}^N),$$

and for $\mu = \frac{1}{2} + \mu' \in (0, 1)$

$$\|(-\Delta)^\mu \phi\|_{L^p(\mathbb{R}^N)} \sim \sum_{j=1}^N \|(-\Delta)^{\mu'} \partial_j \phi\|_{L^p(\mathbb{R}^N)}.$$

In particular, for $p = 2$ and for $\mu = \frac{1}{2} + \mu' \in (0, 1)$

$$\|(-\Delta)^\mu \phi\|_{L^2(\mathbb{R}^N)} \sim \sum_{j=1}^N [\partial_j \phi]_{W^{2\mu', 2}(\mathbb{R}^N)}$$

where in the right hand side we have the Gagliardo seminorms

$$[\phi]_{W^{\mu,p}(\mathbb{R}^N)}^p = \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|\phi(x) - \phi(y)|^p}{|x - y|^{N+\mu p}} dy dx.$$

The fractional semigroup, that is, the semigroup defined by the fractional Laplacian, can be constructed from the heat semigroup for $0 < \mu < 1$ and $1 \leq p < \infty$ as

$$S_\mu(t)\phi = \int_0^\infty f_{t,\mu}(s) S(s)\phi ds = \int_0^\infty f_{1,\mu}(s) S(st^{\frac{1}{\mu}})\phi ds, \quad t > 0 \quad (2.5)$$

and $S_\mu(0)\phi = \phi$ for $\phi \in L^p(\mathbb{R}^N)$, where $f_{t,\mu}$ is given by

$$0 \leq f_{t,\mu}(\lambda) = \begin{cases} \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} e^{z\lambda - tz^\mu} dz & \lambda \geq 0, \\ 0 & \lambda < 0, \end{cases}$$

$\sigma > 0$, and the branch for z^μ is chosen such that $\text{Re}(z^\mu) > 0$ if $\text{Re}(z) > 0$, and we have $\int_0^\infty f_{t,\mu}(s) ds = 1$, see [8, Appendix A.3], [26, p. 259] and [26, (20'), p. 264]. This semigroup is bounded, strongly continuous and analytic, see [8, Corollary 4.1] and [8, Appendix A.3]. For $p = \infty$ the same construction is possible, although the semigroup is bounded and analytic but not strongly continuous, see [9,

Proposition B.1]. In particular, for $1 \leq p \leq \infty$ and $\mu \in (0, 1)$, as the semigroup is analytic, see [19, Proposition 2.1.1], we have that $u(t) = S_\mu(t)u_0$ with $u_0 \in L^p(\mathbb{R}^N)$ solves

$$u_t + (-\Delta)^\mu u = 0, \quad x \in \mathbb{R}^N, \quad t > 0, \quad u(0) = u_0$$

and for $1 \leq p < \infty$ we have

$$S_\mu(t)u_0 \rightarrow u_0 \quad \text{as } t \rightarrow 0^+ \quad \text{in } L^p(\mathbb{R}^N).$$

When $p = \infty$ the following holds.

Proposition 2.1. *For $u_0 \in L^\infty(\mathbb{R}^N)$ and $0 < \mu < 1$*

$$S_\mu(t)u_0 \rightarrow u_0 \quad \text{as } t \rightarrow 0^+ \quad \text{in } L_{loc}^p(\mathbb{R}^N) \text{ for any } 1 \leq p < \infty.$$

Proof. Using the second expression in (2.5) and [26, formula (14), p. 262], if $u_0 \in L^\infty(\mathbb{R}^N)$ and we take a ball $B \subset \mathbb{R}^N$, using (2.3) and Lebesgue's dominated convergence theorem, we get

$$\|S_\mu(t)u_0 - u_0\|_{L^p(B)} \leq \int_0^\infty f_{1,\mu}(s) \|S(st^\frac{1}{\mu})u_0 - u_0\|_{L^p(B)} \rightarrow 0 \quad \text{as } t \rightarrow 0^+.$$

■

Also, from (2.5) and the properties of the heat semigroup, it is easy to obtain that the fractional semigroup is of contractions and order preserving in $L^p(\mathbb{R}^N)$ for $1 \leq p \leq \infty$. Actually $\|S_\mu(t)\|_{\mathcal{L}(L^p(\mathbb{R}^N))} = 1$ for all $t \geq 0$, see e.g. [8, Corollary 3.7]. Moreover it satisfies, among other, the smoothing estimates

$$\|S_\mu(t)\|_{\mathcal{L}(L^p(\mathbb{R}^N), L^q(\mathbb{R}^N))} = \frac{c_{p,q,\mu}}{t^{\frac{N}{2\mu}(\frac{1}{p}-\frac{1}{q})}}, \quad t > 0, \quad (2.6)$$

for $1 \leq p \leq q \leq \infty$, see [6, Theorem 6.2] and [8, Proposition 6.5].

We also have for $0 \leq \gamma \leq \gamma' \leq 1$, $1 < p \leq q < \infty$

$$\|S_\mu(t)\|_{\mathcal{L}(H_p^{2\gamma}(\mathbb{R}^N), H_q^{2\gamma'}(\mathbb{R}^N))} \leq \frac{1}{t^{\frac{N}{2m\mu}(\frac{1}{p}-\frac{1}{q})}} \max\{c_{p,q,\mu}, \frac{c_{\gamma,\gamma',p,q,\mu}}{t^{\frac{\gamma'-\gamma}{\mu}}}\}, \quad t > 0. \quad (2.7)$$

The fractional semigroup has a C^∞ , positive convolution self similar kernel such that

$$S_\mu(t)u_0(x) = \int_{\mathbb{R}^N} k_\mu(t, x, y) u_0(y) dy \quad t > 0, \quad x \in \mathbb{R}^N, \quad (2.8)$$

$$0 \leq k_\mu(t, x, y) = \frac{1}{t^{\frac{N}{2\mu}}} k_{0,\mu} \left(\frac{x-y}{t^{\frac{1}{2\mu}}} \right) \sim \frac{1}{t^{\frac{N}{2\mu}}} H_\mu \left(\frac{x-y}{t^{\frac{1}{2\mu}}} \right), \quad (2.9)$$

where

$$H_\mu(z) = \min \left\{ 1, \frac{1}{|z|^{N+2\mu}} \right\} \sim I_\mu(z) = \frac{1}{(1 + |z|^2)^{\frac{N+2\mu}{2}}}, \quad z \in \mathbb{R}^N. \quad (2.10)$$

The profile $k_{0,\mu}$ is even and satisfies

$$(-\Delta)^\mu k_{0,\mu} = \frac{x}{2\mu} \nabla k_{0,\mu} + \frac{N}{2\mu} k_{0,\mu},$$

see [10, Section 6], [5, 6]. Since the kernel is C^∞ so the solution of the fractional heat equation in (2.8) is $C^\infty((0, \infty) \times \mathbb{R}^N)$, see [6].

3. THE FRACTIONAL SCHRÖDINGER SEMIGROUP

In this section we study the perturbed equations

$$\begin{cases} u_t + (-\Delta)^\mu u + V(x)u = 0, & x \in \mathbb{R}^N, \quad t > 0 \\ u(x, 0) = u_0(x), & x \in \mathbb{R}^N \end{cases}$$

for potentials in the uniform space $L_U^{p_0}(\mathbb{R}^N)$, with $p_0 \geq 1$, that is, satisfying

$$\|V\|_{L_U^{p_0}(\mathbb{R}^N)} = \sup_{x_0 \in \mathbb{R}^N} \|V\|_{L^{p_0}(B(x_0, 1))} < \infty.$$

Proposition 3.1. Assume $0 \leq V \in L_U^{p_0}(\mathbb{R}^N)$, for $p_0 > \frac{N}{2\mu}$.

Then $-(-\Delta)^\mu - V(x)$ defines an order preserving semigroup of contractions in $L^p(\mathbb{R}^N)$ for any $1 \leq p \leq \infty$ which is strongly continuous if $1 \leq p < \infty$ and analytic for $1 < p < \infty$ and that we denote $\{S_{\mu,V}(t)\}_{t \geq 0}$.

These semigroups are consistent in the sense that the semigroup in $L^p(\mathbb{R}^N)$ and in $L^q(\mathbb{R}^N)$ give the same result for any $t > 0$ if $u_0 \in L^p(\mathbb{R}^N) \cap L^q(\mathbb{R}^N)$.

Also the semigroup in $L^{p'}(\mathbb{R}^N)$ is the adjoint of the semigroup in $L^p(\mathbb{R}^N)$ for $1 \leq p < \infty$ and $\frac{1}{p} + \frac{1}{p'} = 1$.

Proof. Consider the bilinear symmetric, nonnegative definite form

$$a(\phi, \psi) = \int_{\mathbb{R}^N} (-\Delta)^{\frac{\mu}{2}} \phi (-\Delta)^{\frac{\mu}{2}} \psi + \int_{\mathbb{R}^N} V(x) \phi \psi.$$

We first prove that is well defined in the Bessel space $H^\mu(\mathbb{R}^N)$. For this, denote by $\{Q_i\}$ the family of cubes centered at points of integer coordinates in \mathbb{R}^N and with edges of length 1 parallel to the axes. Thus if $V \in L_U^{p_0}(\mathbb{R}^N)$, for any $\phi, \psi \in H^\mu(\mathbb{R}^N)$ we have, using Hölder's inequality,

$$\int_{\mathbb{R}^N} |V(x) \phi \psi| = \sum_i \int_{Q_i} |V(x)| |\phi| |\psi| \leq \sum_i \|V\|_{L^{p_0}(Q_i)} \|\phi\|_{L^r(Q_i)} \|\psi\|_{L^r(Q_i)}$$

with $\frac{1}{p_0} + \frac{2}{r} = 1$ and $H^\mu(\mathbb{R}^N) \subset L^r(\mathbb{R}^N)$, that is $\mu - \frac{N}{2} \geq -\frac{N}{r}$. This implies in turn that we must have $p_0 \geq \frac{N}{2\mu}$ (with strict inequality if $p_0 = 1$ which implies $r = \infty$). Hence, using the embedding $H^\mu(Q_i) \subset L^r(Q_i)$, with constants independent of i , we get

$$\begin{aligned} \int_{\mathbb{R}^N} |V(x)| |\phi| |\psi| &\leq C \|V\|_{L_U^{p_0}(\mathbb{R}^N)} \sum_i \|\phi\|_{L^r(Q_i)} \|\psi\|_{L^r(Q_i)} \\ &\leq C \|V\|_{L_U^{p_0}(\mathbb{R}^N)} \left(\sum_i \|\phi\|_{H^\mu(Q_i)}^2 \right)^{\frac{1}{2}} \left(\sum_i \|\psi\|_{H^\mu(Q_i)}^2 \right)^{\frac{1}{2}} \leq C \|V\|_{L_U^{p_0}(\mathbb{R}^N)} \|\phi\|_{H^\mu(\mathbb{R}^N)} \|\psi\|_{H^\mu(\mathbb{R}^N)}, \end{aligned}$$

by Lemma 3.2 below.

Now if $\phi \in H^\mu(\mathbb{R}^N)$ then $|\phi| \in H^\mu(\mathbb{R}^N)$. To see this we will use the generalised Strook-Varopoulos inequality, see [12, Lemma 5.2], so for $g(\phi), (-\Delta)^\mu \phi \in L^2(\mathbb{R}^N)$

$$\int_{\mathbb{R}^N} g(\phi) (-\Delta)^\mu \phi \geq \int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} G(\phi)|^2 \geq 0 \quad (3.1)$$

with $g' = (G')^2$. In particular, with $g(s) = s$ and $G(s) = |s|$ we get for $\phi, (-\Delta)^\mu \phi \in L^2(\mathbb{R}^N)$, that is, for $\phi \in H^{2\mu}(\mathbb{R}^N)$

$$\int_{\mathbb{R}^N} \phi (-\Delta)^\mu \phi \geq \int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} \phi|^2 \geq 0$$

and we get, for $\phi \in H^{2\mu}(\mathbb{R}^N)$

$$\int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} \phi|^2 \leq \int_{\mathbb{R}^N} \phi (-\Delta)^\mu \phi = \int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} \phi|^2.$$

For $\phi \in H^\mu(\mathbb{R}^N)$, by density, if $\phi_n \in H^{2\mu}(\mathbb{R}^N)$ converges in $H^\mu(\mathbb{R}^N)$ to ϕ , the inequality above and $\phi_n \rightarrow \phi$ in $L^2(\mathbb{R}^N)$ we can assume that $|\phi_n| \rightarrow |\phi|$ weakly in $H^\mu(\mathbb{R}^N)$ and we get for $\phi \in H^\mu(\mathbb{R}^N)$

$$\int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} \phi|^2 \leq \int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} \phi|^2.$$

Thus $|\phi| \in H^\mu(\mathbb{R}^N)$.

In particular, if $V \geq 0$ we have

$$a(|\phi|, |\phi|) \leq a(\phi, \phi), \quad \phi \in H^\mu(\mathbb{R}^N). \quad (3.2)$$

On the other hand, if $0 \leq \phi \in H^\mu(\mathbb{R}^N)$ then $v = \min\{\phi, 1\} \in H^\mu(\mathbb{R}^N)$. To see this observe that $v = g(\phi)$ with g Lipschitz and $g'(s) = \mathcal{X}_{[0,1]}(s)$. Then in (3.1) we have $G(s) = g(s)$ and for smooth ϕ

$$\int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} g(\phi)|^2 \leq \int_{\mathbb{R}^N} g(\phi) (-\Delta)^\mu \phi = \int_{\mathbb{R}^N} (-\Delta)^{\frac{\mu}{2}} g(\phi) (-\Delta)^{\frac{\mu}{2}} \phi$$

and Hölder's inequality gives

$$\int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} g(\phi)|^2 \leq \int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} \phi|^2, \quad \phi \in H^{2\mu}(\mathbb{R}^N).$$

Thus $g(\phi) \in H^\mu(\mathbb{R}^N)$. By density, if $\phi_n \in H^{2\mu}(\mathbb{R}^N)$ converges in $H^\mu(\mathbb{R}^N)$ to ϕ , the inequality above and $g(\phi_n) \rightarrow g(\phi)$ in $L^2(\mathbb{R}^N)$ we can assume that $g(\phi_n) \rightarrow g(\phi)$ weakly in $H^\mu(\mathbb{R}^N)$ and we extend the inequality to $\phi \in H^\mu(\mathbb{R}^N)$.

In particular, if $V \geq 0$ we have

$$a(v, v) \leq a(\phi, \phi), \quad \phi \in H^\mu(\mathbb{R}^N). \quad (3.3)$$

Then from (3.2) and (3.3), [11, Theorem 1.3.2, pag 12] and [11, Theorem 1.3.3, pag 14], we have an order preserving semigroup of contractions in $L^p(\mathbb{R}^N)$ for $1 \leq p \leq \infty$. Moreover, from [11, Theorem 1.4.1, pag 22] the semigroup in $L^p(\mathbb{R}^N)$ is the extension of the one in $L^2(\mathbb{R}^N)$, the semigroup in $L^p(\mathbb{R}^N)$ is the adjoint of the semigroup in $L^{p'}(\mathbb{R}^N)$ for $1 < p \leq \infty$, where $\frac{1}{p} + \frac{1}{p'} = 1$. Also, the semigroup is analytic for $1 < p < \infty$ and strongly continuous for $1 \leq p < \infty$. ■

The following result, used above, was proved in Lemma 2.4 in [4].

Lemma 3.2. *Let $\{Q_i\}$ be the family of cubes centered at points of integer coordinates in \mathbb{R}^N and with edges of length 1 parallel to the axes.*

Then for any $0 \leq s \leq 2$ and $1 < p < \infty$

$$\sum_i \|\phi\|_{H_p^s(Q_i)}^p \leq C \|\phi\|_{H_p^s(\mathbb{R}^N)}^p \quad \text{for all } \phi \in H_p^s(\mathbb{R}^N).$$

The next result shows in particular that, for suitable p , the semigroup $\{S_{\mu,V}(t)\}_{t \geq 0}$ above in $L^p(\mathbb{R}^N)$ can be represented in terms of the fractional semigroup $\{S_\mu(t)\}_{t \geq 0}$ and the variation of constants formula (a.k.a. Duhamel's principle), at least for smooth initial data.

Proposition 3.3. *For $0 \leq V \in L_U^{p_0}(\mathbb{R}^N)$ with $p_0 > \max\{\frac{N}{2\mu}, 1\}$, the semigroup $u(t) = S_{\mu,V}(t)u_0$ satisfies the following.*

- (i) *For $1 < p < \infty$ and $p \leq p_0$, the operator $-(-\Delta)^\mu - V$ with domain $D((-\Delta)^\mu) = H_p^{2\mu}(\mathbb{R}^N)$ is a sectorial operator and it is the generator of the analytic semigroup $S_{\mu,V}(t)$. Moreover, for $u_0 \in H_p^{2\mu}(\mathbb{R}^N)$, we have $u \in C([0, \infty), H_p^{2\mu}(\mathbb{R}^N)) \cap C^1((0, \infty), L^p(\mathbb{R}^N))$ and satisfies the variations of constants formula*

$$u(t) = S_\mu(t)u_0 - \int_0^t S_\mu(t-s)Vu(s)ds, \quad t > 0.$$

- (ii) *For $1 \leq p \leq \infty$ and $u_0 \in L^p(\mathbb{R}^N)$,*

$$|S_{\mu,V}(t)u_0| \leq S_{\mu,V}(t)|u_0| \leq S_\mu(t)|u_0|.$$

In particular, the semigroup $S_{\mu,V}(t)$ satisfies the smoothing estimates (2.6).

- (iii) *For $1 \leq p \leq \infty$ and $0 \leq u_0 \in L^p(\mathbb{R}^N)$, if $V_1 \geq V_2$ then for $u_0 \geq 0$ we have $0 \leq S_{\mu,V_1}(t)u_0 \leq S_{\mu,V_2}(t)u_0$.*

Proof. (i) We show that the domain of $(-\Delta)^\mu + V$ coincides with the domain $D((-\Delta)^\mu) = H_p^{2\mu}(\mathbb{R}^N)$ and is a sectorial operator. Actually, denote by $\{Q_i\}$ the family of cubes centered at points of integer coordinates in \mathbb{R}^N and with edges of length 1 parallel to the axes. Thus if $V \in L_U^{p_0}(\mathbb{R}^N)$, for any $\phi \in H_p^{2\mu}(\mathbb{R}^N)$ we have, using Hölder's inequality,

$$\|V\phi\|_{L^p(\mathbb{R}^N)}^p = \int_{\mathbb{R}^N} |V|^p |\phi|^p = \sum_i \int_{Q_i} |V|^p |\phi|^p \leq \sum_i \|V\|_{L^{p_0}(Q_i)}^p \|\phi\|_{L^p(Q_i)}^p$$

provided we chose r such that $\frac{1}{p} = \frac{1}{p_0} + \frac{1}{r}$, which is possible since $p_0 \geq p$. Now we use the Sobolev embedding

$$\|\phi\|_{L^r(Q_i)} \leq C\|\phi\|_{H_p^{2\mu s}(Q_i)}$$

with constants independent of i , for $-\frac{N}{r} \leq 2\mu s - \frac{N}{p}$, which requires $\frac{N}{p_0} \leq 2\mu s$. Hence we get, using Lemma 3.2,

$$\|V\phi\|_{L^p(\mathbb{R}^N)}^p \leq C\|V\|_{L_U^{p_0}(\mathbb{R}^N)}^p \sum_i \|\phi\|_{H_p^{2\mu s}(Q_i)}^p \leq C\|V\|_{L_U^{p_0}(\mathbb{R}^N)}^p \|\phi\|_{H_p^{2\mu s}(\mathbb{R}^N)}^p. \quad (3.4)$$

Then, since $p_0 > \frac{N}{2\mu}$ we can take $0 < s < 1$ above and then by interpolation

$$\|V\phi\|_{L^p(\mathbb{R}^N)} \leq C\|V\|_{L_U^{p_0}(\mathbb{R}^N)} \|\phi\|_{H_p^{2\mu s}(\mathbb{R}^N)} \leq C\|V\|_{L_U^{p_0}(\mathbb{R}^N)} \|\phi\|_{H_p^{2\mu}(\mathbb{R}^N)}^s \|\phi\|_{L^p(\mathbb{R}^N)}^{1-s}$$

and then, for any $\varepsilon > 0$,

$$\|V\phi\|_{L^p(\mathbb{R}^N)} \leq \varepsilon \|(-\Delta)^\mu \phi\|_{L^p(\mathbb{R}^N)} + C_\varepsilon \|\phi\|_{L^p(\mathbb{R}^N)}, \quad \phi \in H_p^{2\mu}(\mathbb{R}^N). \quad (3.5)$$

The sectoriality result follows from Theorem 2.1 in Chapter 3 in [21]. The regularity of $u(t) = S_{\mu,V}(t)u_0$ when $u_0 \in H_p^{2\mu}(\mathbb{R}^N)$ also follows from Chapter 1 in [21]. In this case u satisfies

$$u_t = -(-\Delta)^\mu u - Vu, \quad t > 0$$

and we get the integral representation of u as $Vu \in C([0, \infty), L^p(\mathbb{R}^N))$, see Chapter 4, Section 4.2 in [21].

(ii) Since $S_{\mu,V}(t)$ is order preserving, it is enough to prove the second inequality because for any order preserving semigroup in $L^p(\mathbb{R}^N)$, since $-|u_0| \leq u_0 \leq |u_0|$ we have

$$|S(t)u_0| \leq S(t)|u_0|.$$

First, for $1 < p \leq p_0$, if $0 \leq u_0 \in H_p^{2\mu}(\mathbb{R}^N)$ we know that $V \geq 0$, $u(t) = S_{\mu,V}(t)u_0 \geq 0$ and $S_\mu(t)$ is order preserving and then by (i) $\int_0^t S_\mu(t-s)Vu(s)ds \geq 0$ and

$$u(t) = S_\mu(t)u_0 - \int_0^t S_\mu(t-s)Vu(s)ds \leq S_\mu(t)u_0, \quad t > 0$$

and (ii) holds for such initial data. By density we get the result for $0 \leq u_0 \in L^p(\mathbb{R}^N)$. For initial data in $L^p(\mathbb{R}^N)$ with $p_0 < p < \infty$ the result follows again by density from the case above.

Finally, for $p = \infty$ and $0 \leq u_0 \in L^\infty(\mathbb{R}^N)$, consider $0 \leq \varphi \in L^1(\mathbb{R}^N)$ and then

$$\langle S_{\mu,V}(t)u_0, \varphi \rangle = \langle u_0, S_{\mu,V}(t)\varphi \rangle \leq \langle u_0, S_\mu(t)\varphi \rangle = \langle S_\mu(t)u_0, \varphi \rangle$$

and we get the pointwise inequality.

Now it is clear that the estimates (2.6) apply to $S_{\mu,V}(t)$.

(iii) Again, first for $1 < p \leq p_0$, if $0 \leq u_0 \in H_p^{2\mu}(\mathbb{R}^N)$ and $u_i(t) = S_{\mu,V_i}(t)u_0 \geq 0$, by (i) $u_2(t)$ satisfies

$$(u_2)_t + (-\Delta)^\mu u_2 + V_1(x)u_2 + (V_2(x) - V_1(x))u_2 = 0, \quad x \in \mathbb{R}^N, \quad t > 0$$

and then, since $0 \leq (V_1 - V_2)u_2 \in C([0, \infty), L^p(\mathbb{R}^N))$,

$$u_2(t) = S_{\mu,V_1}(t)u_0 + \int_0^t S_{\mu,V_1}(t-s)(V_1 - V_2)u_2(s)ds \geq S_{\mu,V_1}(t)u_0, \quad t > 0.$$

By density, (iii) is proved for this range of p . For initial data in $L^p(\mathbb{R}^N)$ with $p_0 < p < \infty$ the result follows again by density from the case above.

Finally, for $p = \infty$ and $0 \leq u_0 \in L^\infty(\mathbb{R}^N)$, consider $0 \leq \varphi \in L^1(\mathbb{R}^N)$ and then

$$\langle S_{\mu,V_1}(t)u_0, \varphi \rangle = \langle u_0, S_{\mu,V_1}(t)\varphi \rangle \leq \langle u_0, S_{\mu,V_2}(t)\varphi \rangle = \langle S_{\mu,V_2}(t)u_0, \varphi \rangle$$

and we get the pointwise inequality. ■

In the case of a bounded potential $0 \leq V \in L^\infty(\mathbb{R}^N)$, part (i) in Proposition 3.3 can be somehow improved and for all $1 \leq p \leq \infty$, the semigroup $\{S_{\mu,V}(t)\}_{t \geq 0}$ above in $L^p(\mathbb{R}^N)$ can be represented by

the variation of constants formula for all initial data. Observe that actually the sign condition on V is not needed in the proof below.

Proposition 3.4. *Assume $0 \leq V \in L^\infty(\mathbb{R}^N)$. Then for $1 \leq p < \infty$ the operator $-(-\Delta)^\mu - V$ with domain $D((-\Delta)^\mu) = H_p^{2\mu}(\mathbb{R}^N)$ is the generator of the semigroup $S_{\mu,V}(t)$.*

For $1 \leq p \leq \infty$ and $u_0 \in L^p(\mathbb{R}^N)$, we have that $u(t) = S_{\mu,V}(t)u_0$ satisfies the variations of constants formula

$$u(t) = S_\mu(t)u_0 - \int_0^t S_\mu(t-s)Vu(s) ds, \quad t > 0.$$

Proof. For $1 \leq p < \infty$ the semigroup $\{S_\mu(t)\}_{t \geq 0}$ is strongly continuous and the multiplication by V is a bounded operator in $L^p(\mathbb{R}^N)$. Hence the result, including the variations of constants formula, follows from Section 3.1 in [21].

Before dealing with the case $p = \infty$ observe first that for $1 \leq p \leq \infty$ and $T_0\|V\|_{L^\infty} < 1$ the mapping

$$\mathcal{F}_p(u)(t) = S_\mu(t)u_0 - \int_0^t S_\mu(t-s)Vu(s) ds$$

defines a contraction in $L^\infty((0, T_0), L^p(\mathbb{R}^N))$ as

$$\|\mathcal{F}_p(u)(t)\|_{L^p} \leq \|u_0\|_{L^p} + \int_0^t \|V\|_{L^\infty}\|u(s)\|_{L^p} ds$$

so $\mathcal{F}_p(u) \in L^\infty((0, T_0), L^p(\mathbb{R}^N))$ and

$$\|\mathcal{F}_p(u)(t) - \mathcal{F}_p(v)(t)\|_{L^p} \leq \int_0^t \|V\|_{L^\infty}\|u(s) - v(s)\|_{L^p} ds \leq T_0\|V\|_{L^\infty} \sup_{0 \leq t \leq T_0} \|u(s) - v(s)\|_{L^p}.$$

Take $u_1(t)$ for $0 \leq t \leq T_0$ the unique fixed point in $L^\infty((0, T_0), L^p(\mathbb{R}^N))$ and consider for $t \geq T_0$,

$$\mathcal{H}_p(u)(t) = S_\mu(t - T_0)u_1(T_0) - \int_{T_0}^t S_\mu(t-s)Vu(s) ds.$$

With similar estimates as above it is immediate to get that this is also a contraction in $L^\infty((T_0, 2T_0), L^p(\mathbb{R}^N))$ and has a unique fixed point $u_2(t)$ for $T_0 \leq t \leq 2T_0$. Then it is easy to get that the function

$$u(t) = \begin{cases} u_1(t), & 0 \leq t \leq T_0 \\ u_2(t), & T_0 \leq t \leq 2T_0 \end{cases}$$

is a fixed point of \mathcal{F}_p in $L^\infty((0, 2T_0), L^p(\mathbb{R}^N))$. Proceeding by induction we get a fixed point of \mathcal{F}_p in $L^\infty((0, kT_0), L^p(\mathbb{R}^N))$ for any $k \in \mathbb{N}$. The uniqueness of the fixed point for \mathcal{F}_p follows from Gronwall's lemma since for two such fixed points and $t > 0$

$$\|u(t) - v(t)\|_{L^p} \leq \int_0^t \|V\|_{L^\infty}\|u(s) - v(s)\|_{L^p} ds.$$

In particular, for $1 \leq p < \infty$ and $T > 0$, $u(t) = S_{\mu,V}(t)u_0$ for $0 \leq t \leq T$ is the unique fixed point of \mathcal{F}_p in $L^\infty((0, T), L^p(\mathbb{R}^N))$ for all $T > 0$.

Hence, if $u_0 \in L^p(\mathbb{R}^N) \cap L^\infty(\mathbb{R}^N)$ with $p > \frac{N}{2\mu}$ then the semigroup solution in $L^p(\mathbb{R}^N)$, $u(t) = S_{\mu,V}(t)u_0$ satisfies, using the variations of constants formula and (2.6),

$$\|u(t)\|_{L^\infty} \leq \|u_0\|_{L^\infty} + \int_0^t \frac{1}{(t-s)^{\frac{N}{2p\mu}}} \|V\|_{L^\infty}\|u(s)\|_{L^p} ds$$

and therefore for any $T > 0$, $S_{\mu,V}(\cdot)u_0 \in L^\infty((0, T), L^\infty(\mathbb{R}^N))$ and so it is the fixed point of \mathcal{F}_∞ in $[0, T]$ in $L^\infty((0, T), L^\infty(\mathbb{R}^N))$.

Now for $u_0 \in L^\infty(\mathbb{R}^N)$ take a sequence $u_0^n \in L^1(\mathbb{R}^N) \cap L^\infty(\mathbb{R}^N) \subset L^p(\mathbb{R}^N)$ with $p > \frac{N}{2\mu}$ such that $u_0^n \rightarrow u_0$ weak-* in $L^\infty(\mathbb{R}^N)$ (for example, take the truncation by zero of u_0 outside the ball $B(0, n) \subset \mathbb{R}^N$). Then for any $\varphi \in L^1(\mathbb{R}^N)$ and $t \geq 0$, as $n \rightarrow \infty$ we have, since $S_{\mu,V}(t)\varphi \in L^1(\mathbb{R}^N)$,

$$\langle S_{\mu,V}(t)u_0^n, \varphi \rangle = \langle u_0^n, S_{\mu,V}(t)\varphi \rangle \rightarrow \langle u_0, S_{\mu,V}(t)\varphi \rangle = \langle S_{\mu,V}(t)u_0, \varphi \rangle$$

i.e. $S_{\mu,V}(t)u_0^n \rightarrow S_{\mu,V}(t)u_0$ weak-* in $L^\infty(\mathbb{R}^N)$.

On the other hand, setting $u^n(t) = S_{\mu,V}(t)u_0^n$ and $u(t) = S_{\mu,V}(t)u_0$, we have,

$$\langle u^n(t), \varphi \rangle = \langle u_0^n, S_\mu(t)\varphi \rangle - \int_0^t \langle u^n(s), VS_\mu(t-s)\varphi \rangle ds.$$

Therefore, for fixed $t > 0$ and for $0 < s < t$ we have, as $n \rightarrow \infty$,

$$\langle u^n(s), VS_\mu(t-s)\varphi \rangle \rightarrow \langle u(s), VS_\mu(t-s)\varphi \rangle$$

and for all $n \in \mathbb{N}$

$$|\langle u^n(s), VS_\mu(t-s)\varphi \rangle| \leq \|VS_\mu(t-s)\varphi\|_{L^1} = g(s)$$

with $g \in L^1(0, t)$. Then Lebesgue's theorem implies that, as $n \rightarrow \infty$, we get

$$\langle u(t), \varphi \rangle = \langle u_0, S_\mu(t)\varphi \rangle - \int_0^t \langle u(s), VS_\mu(t-s)\varphi \rangle ds.$$

Hence, for all $T > 0$, $u(\cdot) = S_{\mu,V}(\cdot)u_0 \in L^\infty((0, T), L^\infty(\mathbb{R}^N))$ satisfies, for all $\varphi \in L^1(\mathbb{R}^N)$,

$$\langle u(t), \varphi \rangle = \langle S_\mu(t)u_0, \varphi \rangle - \int_0^t \langle S_\mu(t-s)Vu(s), \varphi \rangle ds, \quad 0 < t < T$$

and therefore

$$u(t) = S_\mu(t)u_0 - \int_0^t S_\mu(t-s)Vu(s) ds$$

and $u(\cdot) = S_{\mu,V}(\cdot)u_0$ is the unique fixed point of \mathcal{F}_∞ in $[0, T]$ in $L^\infty((0, T), L^\infty(\mathbb{R}^N))$. ■

In view of Propositions 3.3 and 3.4, given $1 \leq p_0 < \infty$ we want to discuss the class of $V \in L_U^{p_0}(\mathbb{R}^N)$ that can be approximated in $L_U^{p_0}(\mathbb{R}^N)$ by bounded functions and to analyze the convergence of the corresponding semigroups. Then we have the following result. Notice that below we use the class $\dot{L}_U^{p_0}(\mathbb{R}^N)$ of functions in $L_U^{p_0}(\mathbb{R}^N)$ such that the translations are continuous in the $L_U^{p_0}(\mathbb{R}^N)$ norm, that is, $V \in \dot{L}_U^{p_0}(\mathbb{R}^N)$ iff

$$\|\tau_y V - V\|_{L_U^{p_0}(\mathbb{R}^N)} \rightarrow 0, \quad \text{as } |y| \rightarrow 0$$

where $y \in \mathbb{R}^N$ and $\tau_y V(x) = V(x - y)$. This is a closed proper subspace of $L_U^{p_0}(\mathbb{R}^N)$, see [3].

Proposition 3.5. (i) For $1 \leq p_0 < \infty$ a function $V \in L_U^{p_0}(\mathbb{R}^N)$ can be approximated in $L_U^{p_0}(\mathbb{R}^N)$ by bounded functions if and only if, defining

$$V_M(x) = \begin{cases} M & \text{if } V(x) > M \\ V(x) & \text{if } -M \leq V(x) \leq M, \\ -M & \text{if } V(x) < -M \end{cases} \quad M > 0,$$

we have $V_M \xrightarrow{M \rightarrow \infty} V$ in $L_U^{p_0}(\mathbb{R}^N)$.

- (ii) If $V \in \dot{L}_U^{p_0}(\mathbb{R}^N)$ then V can be approximated in $L_U^{p_0}(\mathbb{R}^N)$ by bounded functions.
- (iii) Assume V is not too large at infinity in the sense that there exists M_0 such that

$$\lim_{|x| \rightarrow \infty} \int_{B(x,1)} |V(y) - V_{M_0}(y)|^{p_0} dy = 0.$$

Then $V_M \xrightarrow{M \rightarrow \infty} V$ in $L_U^{p_0}(\mathbb{R}^N)$.

- (iv) The class of functions in $L_U^{p_0}(\mathbb{R}^N)$ that can be approximated in $L_U^{p_0}(\mathbb{R}^N)$ by bounded functions is a closed proper subspace of $L_U^{p_0}(\mathbb{R}^N)$.

Proof. (i) Assume V can be approximated in $L_U^{p_0}(\mathbb{R}^N)$ by bounded functions. Then there exists $W_n \in L^\infty(\mathbb{R}^N)$ such that $W_n \xrightarrow{n \rightarrow \infty} V$ in $L_U^{p_0}(\mathbb{R}^N)$. Letting $M_n = \|W_n\|_{L^\infty}$, we have, for every $x \in \mathbb{R}^N$,

$$\int_{B(x,1)} |V(y) - V_{M_n}(y)|^{p_0} dy = \int_{B(x,1) \cap \{V \geq M_n\}} |V(y) - M_n|^{p_0} dy + \int_{B(x,1) \cap \{V \leq -M_n\}} |V(y) + M_n|^{p_0} dy$$

and thus

$$\int_{B(x,1)} |V(y) - V_{M_n}(y)|^{p_0} dy \leq \int_{B(x,1)} |V(y) - W_n(y)|^{p_0} dy.$$

Therefore $V_{M_n} \xrightarrow[n \rightarrow \infty]{} V$ in $L_U^{p_0}(\mathbb{R}^N)$. Since clearly $\|V - V_M\|_{L_U^{p_0}(\mathbb{R}^N)}$ decreases with M , then $V_M \xrightarrow[M \rightarrow \infty]{} V$ in $L_U^{p_0}(\mathbb{R}^N)$.

The converse is immediate.

(ii) If $V \in \dot{L}_U^{p_0}(\mathbb{R}^N)$ from the results in [3] we know that the solution of the heat equation satisfies $S(t)V \xrightarrow[t \rightarrow 0^+]{} V$ in $L_U^{p_0}(\mathbb{R}^N)$ and $S(t)V \in L^\infty(\mathbb{R}^N)$.

(iii) Define the family of bounded continuous functions in \mathbb{R}^N , $H_M(x) = \int_{B(x,1)} |V(y) - V_M(y)|^{p_0} dy$ which are decreasing in M and converge to 0 as $M \rightarrow \infty$ uniformly in bounded sets, by Lebesgue's theorem.

Then the assumption reads $\lim_{|x| \rightarrow \infty} H_{M_0}(x) = 0$. Therefore, given $\varepsilon > 0$ there exists $R > 0$ such that for $|x| > R$ and $M > M_0$ we have

$$0 \leq H_M(x) \leq H_{M_0}(x) \leq \varepsilon.$$

This and the uniform convergence $H_M(x) \xrightarrow[M \rightarrow \infty]{} 0$ for $|x| \leq R$ implies $H_M \xrightarrow[M \rightarrow \infty]{} 0$ uniformly in \mathbb{R}^N and hence $V_M \xrightarrow[M \rightarrow \infty]{} V$ in $L_U^{p_0}(\mathbb{R}^N)$.

(iv) The closedness is immediate as this class is the closure of $L^\infty(\mathbb{R}^N)$ in $L_U^{p_0}(\mathbb{R}^N)$.

To show the subspace is proper, consider a sequence $|x_n| \rightarrow \infty$ with $|x_n - x_m| > 2$ for all $n, m \in \mathbb{N}$. Then define $V(x) = \sum_n n^{\frac{N}{p_0}} \mathcal{X}_{B(x_n, \frac{1}{n})}(x)$. Clearly $\|V\|_{L^{p_0}(B(x,1))} \leq c$ for all $x \in \mathbb{R}^N$ since $B(x, 1)$ contains at most one point of $\{x_n\}_n$, so $V \in L_U^{p_0}(\mathbb{R}^N)$. Also, for any $M > 0$ if $n > M$ we have

$$\|V - V_M\|_{L^{p_0}(B(x_n,1))}^{p_0} \geq c \frac{(n^{\frac{N}{p_0}} - M)^{p_0}}{n^N} \xrightarrow[n \rightarrow \infty]{} c > 0$$

and therefore $\|V - V_M\|_{L_U^{p_0}(\mathbb{R}^N)}^{p_0} \geq c > 0$ for all $M > 0$. ■

Remark 3.6. (i) Assume $V \in L_U^{p_0}(\mathbb{R}^N)$ with $p_0 > 1$ and can be approximated in $L_U^{p_0}(\mathbb{R}^N)$ by bounded functions. Then V can be approximated in $L_U^1(\mathbb{R}^N)$ by bounded functions. To see this just observe that uniform spaces are nested, that is, $L_U^p(\mathbb{R}^N) \subset L_U^q(\mathbb{R}^N)$, if $p > q$.

(ii) Conversely, if $V \in L_U^{p_0}(\mathbb{R}^N)$ with $p_0 > 1$ and can be approximated in $L_U^1(\mathbb{R}^N)$ by bounded functions, then for every $1 < q < p_0$, V can be approximated in $L_U^q(\mathbb{R}^N)$ by bounded functions. To see this, just notice that like in bounded domains, we have the interpolation inequality

$$\|f\|_{L_U^q} \leq \|f\|_{L_U^{p_0}}^{1-\theta} \|f\|_{L_U^1}^\theta, \quad \frac{1}{q} = \frac{1-\theta}{p_0} + \frac{\theta}{1}, \quad 0 < \theta < 1,$$

for any $f \in L_U^{p_0}(\mathbb{R}^N)$.

Now we analyze the approximation of the corresponding semigroups. For this assume now $0 \leq V \in L_U^{p_0}(\mathbb{R}^N)$ with $p_0 > \max\{\frac{N}{2\mu}, 1\}$ and consider the increasing sequence $0 \leq V_M \leq V$. Then, by Proposition 3.3, we have, for $1 \leq p \leq \infty$ and $0 \leq u_0 \in L^p(\mathbb{R}^N)$,

$$0 \leq S_{\mu,V}(t)u_0 \leq S_{\mu,V_M}(t)u_0$$

and for $M_1 \leq M_2$

$$0 \leq S_{\mu,V_{M_2}}(t)u_0 \leq S_{\mu,V_{M_1}}(t)u_0.$$

Hence, the monotonic decreasing limit

$$\tilde{S}_{\mu,V}(t)u_0 \stackrel{\text{def}}{=} \lim_{M \rightarrow \infty} S_{\mu,V_M}(t)u_0$$

exists pointwise and in $L^p(\mathbb{R}^N)$ for all $t > 0$ and satisfies $0 \leq S_{\mu,V}(t)u_0 \leq \tilde{S}_{\mu,V}(t)u_0$, $\tilde{S}_{\mu,V}(t+s)u_0 = \tilde{S}_{\mu,V}(t)\tilde{S}_{\mu,V}(s)u_0$ and $\lim_{t \rightarrow 0^+} \tilde{S}_{\mu,V}(t)u_0 = u_0$ in $L^p(\mathbb{R}^N)$. Using the positive and negative part of $u_0 \in L^p(\mathbb{R}^N)$ we can extend $\tilde{S}_{\mu,V}(t)$ to an order preserving C^0 semigroup of contractions in $L^p(\mathbb{R}^N)$.

Now we prove that $\tilde{S}_{\mu,V}(t) = S_{\mu,V}(t)$ for $t \geq 0$ if V can be approximated in $L_U^{p_0}(\mathbb{R}^N)$ by bounded functions.

Theorem 3.7. *Assume $0 \leq V \in L_U^{p_0}(\mathbb{R}^N)$ with $p_0 > \max\{\frac{N}{2\mu}, 1\}$ can be approximated in $L_U^{p_0}(\mathbb{R}^N)$ by bounded functions.*

Then $\tilde{S}_{\mu,V}(t) = S_{\mu,V}(t)$ for $t \geq 0$, as operators in $L^p(\mathbb{R}^N)$ for $1 \leq p \leq \infty$.

Proof. Take $0 \leq u_0 \in C_c^\infty(\mathbb{R}^N)$ and $u_M(t) = S_{\mu,V_M}(t)u_0$ and $u(t) = S_{\mu,V}(t)u_0$ for $t \geq 0$. Since these semigroups are consistent in the Lebesgue spaces, it is enough to show that $u(t) = \lim_{M \rightarrow \infty} u_M(t)$ for $t > 0$ in some of these spaces. Take $1 < p \leq p_0$. As u_0 belongs to the domain of the generator, $H_p^{2\mu}(\mathbb{R}^N)$, we have, by Propositions 3.3 and 3.4,

$$u_M(t) = S_{\mu}(t)u_0 - \int_0^t S_{\mu}(t-s)V_M u_M(s) ds, \quad u(t) = S_{\mu}(t)u_0 - \int_0^t S_{\mu}(t-s)V u(s) ds, \quad t > 0.$$

Also by general properties of semigroups in [21], we have that $u_M(t), u(t) \in H_p^{2\mu}(\mathbb{R}^N)$ for $t > 0$.

Now notice that (3.4), (3.5) and the closed mapping theorem also imply that the graph norm of $L = (-\Delta)^\mu + V$ in $H_p^{2\mu}(\mathbb{R}^N)$ is equivalent to the $H_p^{2\mu}(\mathbb{R}^N)$ norm, that is,

$$c_0 \|\phi\|_{H_p^{2\mu}} \leq \|L\phi\|_{L^p} + \|\phi\|_{L^p} \leq c_1 \|\phi\|_{H_p^{2\mu}}.$$

This implies, using that $Lu(t) = LS_{\mu,V}(t)u_0 = S_{\mu,V}(t)Lu_0$ and the semigroup is of contractions, that

$$\|u(t)\|_{H_p^{2\mu}} \leq c \|u_0\|_{H_p^{2\mu}}, \quad t > 0.$$

Now using the variations of constants formula and adding and subtracting the term $\int_0^t S_{\mu}(t-s)V_M u(s) ds$, we get that $z_M(t) = u(t) - u_M(t)$ satisfies

$$z_M(t) = - \int_0^t S_{\mu}(t-s)(V - V_M)u(s) ds - \int_0^t S_{\mu}(t-s)V_M z_M(s) ds, \quad t > 0.$$

Hence for $0 < r < 1$, using (2.7),

$$\|z_M(t)\|_{H_p^{2\mu r}} \leq \int_0^t \frac{c}{(t-s)^r} \|(V - V_M)u(s)\|_{L^p} ds + \int_0^t \frac{c}{(t-s)^r} \|V_M z_M(s)\|_{L^p} ds$$

and (3.4) yields

$$\|z_M(t)\|_{H_p^{2\mu r}} \leq \int_0^t \frac{c}{(t-s)^r} \|V - V_M\|_{L_U^{p_0}} \|u(s)\|_{H_p^{2\mu}} ds + \int_0^t \frac{c}{(t-s)^r} \|V_M\|_{L_U^{p_0}} \|z_M(s)\|_{H_p^{2\mu r}} ds$$

and then

$$\|z_M(t)\|_{H_p^{2\mu r}} \leq Ct^{1-r} \|V - V_M\|_{L_U^{p_0}} \|u_0\|_{H_p^{2\mu}} + C \int_0^t \frac{1}{(t-s)^r} \|z_M(s)\|_{H_p^{2\mu r}} ds.$$

Then Henry's singular Gronwall Lemma 7.1.1 in [16] implies that for any $T > 0$

$$\|z_M(t)\|_{H_p^{2\mu r}} \leq C(T) \|V - V_M\|_{L_U^{p_0}} \xrightarrow{M \rightarrow \infty} 0, \quad 0 < t < T.$$

Therefore $\tilde{S}_{\mu,V}(t)u_0 = S_{\mu,V}(t)u_0$ for $t \geq 0$ and $0 \leq u_0 \in C_c^\infty(\mathbb{R}^N)$. By linearity and density we have the result. ■

4. EXPONENTIAL DECAY

In this section we want to characterise certain classes of non negative potentials $0 \leq V \in L_U^{p_0}(\mathbb{R}^N)$ with $p_0 > \frac{N}{2\mu}$ such that the fractional Schrödinger semigroup $\{S_{\mu,V}(t)\}_{t \geq 0}$ decays exponentially in $L^p(\mathbb{R}^N)$. Since the semigroup is of contractions in $L^p(\mathbb{R}^N)$ for $1 \leq p \leq \infty$ then either $\|S_{\mu,V}(t)\|_{\mathcal{L}(L^p(\mathbb{R}^N))} = 1$ for all $t > 0$ or there exists $M \geq 1$, $a > 0$ such that

$$\|S_{\mu,V}(t)\|_{\mathcal{L}(L^p(\mathbb{R}^N))} \leq M e^{-at}, \quad t \geq 0. \quad (4.1)$$

Actually if for $t_0 > 0$, $\alpha \stackrel{\text{def}}{=} \|S_{\mu,V}(t_0)\|_{\mathcal{L}(L^p(\mathbb{R}^N))} < 1$, then writing $t = kt_0 + s$, with $k \in \mathbb{N}$ and $s \in (0, t_0]$ and the semigroup property gives $\|S_{\mu,V}(t)\|_{\mathcal{L}(L^p(\mathbb{R}^N))} \leq \frac{1}{\alpha} e^{-at}$ where $a = \frac{|\ln(\alpha)^{-1}|}{t_0} > 0$, i.e. (4.1). Notice that one obtains $M \geq 1$ even if the semigroup is of contractions.

Then we define the exponential type of the semigroup $\{S_{\mu,V}(t)\}_{t \geq 0}$ in $L^p(\mathbb{R}^N)$, $1 \leq p \leq \infty$, as

$$\omega_p \stackrel{\text{def}}{=} \sup\{a \in \mathbb{R}, \|S_{\mu,V}(t)\|_{\mathcal{L}(L^p(\mathbb{R}^N))} \leq M e^{-at}, \quad t \geq 0, \text{ for some } M \geq 1\},$$

and $\omega_p \geq 0$, since the semigroup is of contractions. The exponential type is related to the spectral bound of the generator $A_p = -(-\Delta)^\mu - V$, since these semigroups are order preserving, then for $1 \leq p < \infty$,

$$-\omega_p = s(A_p) \stackrel{\text{def}}{=} \sup\{Re(\lambda), \lambda \in \sigma(A_p)\},$$

see [24, 25].

Then we have the following result. Notice that below we denote by $\|\cdot\|_{p \rightarrow q} = \|\cdot\|_{\mathcal{L}(L^p, L^q)}$.

Theorem 4.1. *For $0 \leq V \in L_U^{p_0}(\mathbb{R}^N)$ with $p_0 > \max\{\frac{N}{2\mu}, 1\}$, the semigroup $u(t) = S_{\mu,V}(t)u_0$ satisfies the following.*

(i) *For $1 \leq p < \infty$ and $t \geq 0$,*

$$\|S_{\mu,V}(t)\|_{2 \rightarrow 2} \leq \|S_{\mu,V}(t)\|_{p \rightarrow p} = \|S_{\mu,V}(t)\|_{p' \rightarrow p'} \leq \|S_{\mu,V}(t)\|_{\infty \rightarrow \infty} = \|S_{\mu,V}(t)\|_{1 \rightarrow 1}$$

(ii) *For $0 < \mu < 1$ and $1 \leq p \leq \infty$, the exponential type satisfies*

$$\omega_2 \geq \omega_p \geq \omega_\infty \geq \frac{\omega_2}{1 + \frac{N}{4\mu}} \geq 0.$$

(iii) *For $\mu = 1$, the exponential type ω_p is independent of $1 \leq p \leq \infty$.*

In particular, in cases (ii) and (iii) above, either $\omega_p = 0$ for all $1 \leq p \leq \infty$ or they are all positive simultaneously. In such a case there exists a common decay rate independent of p .

Proof. (i) Since for $1 \leq p < \infty$ the semigroup in $L^{p'}(\mathbb{R}^N)$ is the adjoint of the semigroup in $L^p(\mathbb{R}^N)$ we get the equalities in the statement.

Now, if $1 \leq p < 2 < p'$ then by the Riesz-Thorin interpolation theorem we get $\|S_{\mu,V}(t)\|_{2 \rightarrow 2} \leq \|S_{\mu,V}(t)\|_{p \rightarrow p} = \|S_{\mu,V}(t)\|_{p' \rightarrow p'}$.

(ii) From (i) we immediately get $\omega_2 \geq \omega_p = \omega_{p'} \geq \omega_\infty = \omega_1$. Since also $\omega_\infty = \omega_1 \geq 0$ then if $\omega_2 = 0$ then the result is proved. Therefore we just need to consider the case $\omega_2 > 0$, that is, the semigroup decays exponentially in $L^2(\mathbb{R}^N)$.

Since the semigroup in $L^\infty(\mathbb{R}^N)$ is order preserving, then for $t > 0$, $\|S_{\mu,V}(t)\|_{\infty \rightarrow \infty} = \|S_{\mu,V}(t)\mathbf{1}\|_{L^\infty(\mathbb{R}^N)}$ so we estimate below the sup norm of the nonnegative solution $S_{\mu,V}(t)\mathbf{1}$. For this fix $x \in \mathbb{R}^N$ and denote by \mathcal{X}_R the characteristic function of the ball $B(x, R)$ and then

$$(S_{\mu,V}(t)\mathbf{1})(x) = (S_{\mu,V}(t)\mathcal{X}_R)(x) + (S_{\mu,V}(t)(1 - \mathcal{X}_R))(x). \quad (4.2)$$

Now since from part (ii) in Proposition 3.3 the semigroup $S_{\mu,V}(t)$ satisfies (2.6) we have, for $t \geq 1$,

$$(S_{\mu,V}(t)\mathcal{X}_R)(x) = (S_{\mu,V}(1)S_{\mu,V}(t-1)\mathcal{X}_R)(x)$$

and therefore

$$0 \leq (S_{\mu,V}(t)\mathcal{X}_R)(x) \leq \|S_{\mu,V}(1)\|_{2 \rightarrow \infty} \|S_{\mu,V}(t-1)\|_{2 \rightarrow 2} \|\mathcal{X}_R\|_2 = C \|S_{\mu,V}(t-1)\|_{2 \rightarrow 2} R^{\frac{N}{2}}.$$

Now, for the second term in (4.2), using again part (ii) in Proposition 3.3, the properties of the fractional heat kernel in (2.8), (2.9), (2.10) and changing variables, we have

$$0 \leq (S_{\mu,V}(t)(1 - \mathcal{X}_R))(x) \leq (S_{\mu}(t)(1 - \mathcal{X}_R))(x) = \int_{\mathbb{R}^N} k_{0,\mu}(z)(1 - \mathcal{X}_R(x - t^{\frac{1}{2\mu}}z)) dz,$$

that is,

$$0 \leq (S_{\mu,V}(t)(1 - \mathcal{X}_R))(x) \leq \int_{|z| > \frac{R}{t^{\frac{1}{2\mu}}}} k_{0,\mu}(z) dz \sim \int_{|z| > \frac{R}{t^{\frac{1}{2\mu}}}} \frac{1}{(1 + |z|^2)^{\frac{N+2\mu}{2}}} dz.$$

Using polar coordinates we get

$$0 \leq (S_{\mu,V}(t)(1 - \mathcal{X}_R))(x) \lesssim \int_{\frac{R}{t^{\frac{1}{2\mu}}}}^{\infty} \frac{1}{r^{2\mu+1}} dr \sim \frac{t}{R^{2\mu}}$$

and therefore, for $t \geq 1$ and $R > 0$,

$$\|S_{\mu,V}(t)\|_{\infty \rightarrow \infty} \leq c\|S_{\mu,V}(t-1)\|_{2 \rightarrow 2} R^{\frac{N}{2}} + c \frac{t}{R^{2\mu}}.$$

Minimizing this in $R > 0$ we get, for $t \geq 1$,

$$\|S_{\mu,V}(t)\|_{\infty \rightarrow \infty} \leq c\|S_{\mu,V}(t-1)\|_{2 \rightarrow 2}^{\frac{1}{1+\frac{N}{4\mu}}} t^{\frac{N}{N+4\mu}}.$$

Therefore

$$\omega_{\infty} \geq \frac{\omega_2}{1 + \frac{N}{4\mu}}.$$

(iii) For the case $\mu = 1$ we proceed as in case (ii) but since the kernel is the Gaussian we get

$$0 \leq (S_{1,V}(t)(1 - \mathcal{X}_R))(x) \lesssim \int_{|z| > \frac{R}{t^{\frac{1}{2}}}} e^{-|z|^2} dz = \int_{\frac{R}{t^{\frac{1}{2}}}}^{\infty} r^{N-1} e^{-r^2} dr \lesssim \int_{\frac{R}{t^{\frac{1}{2}}}}^{\infty} r e^{-\varepsilon r^2} dr \sim e^{-\varepsilon \frac{R^2}{t}}$$

with $\varepsilon < 1$, so choosing $R = At$ with A large, we get a decay of order $e^{-\varepsilon At}$ as fast as we want. Also, from the estimate in case (ii) above, we have

$$0 \leq (S_{1,V}(t)\mathcal{X}_R)(x) \leq c\|S_{1,V}(t-1)\|_{2 \rightarrow 2} t^{\frac{N}{2}}$$

which is only a polynomial correction of the decay of $\|S_{1,V}(t)\|_{2 \rightarrow 2}$. Therefore $\omega_{\infty} = \omega_2$ and the result is proved. ■

Remark 4.2. (i) *The independence of the exponential type when $\mu = 1$ holds even without assuming $V \geq 0$, see Theorem B.5.1 in [23] and references therein.*

(ii) *When $0 < \mu < 1$ the same proof of part (ii) in Theorem 4.1 but using $\|S_{\mu,V}(t-1)\|_{p \rightarrow p}$ with $1 \leq p < \infty$ leads to*

$$\omega_p \geq \omega_{\infty} \geq \frac{\omega_p}{1 + \frac{N}{2p\mu}}$$

and this combined with $\omega_p = \omega_{p'}$ give

$$\lim_{p \rightarrow 1} \omega_p = \omega_1 = \omega_{\infty} = \lim_{p \rightarrow \infty} \omega_p.$$

(iii) *Observe that for $1 < p < \infty$, if $\|S_{\mu,V}(t)\|_{\mathcal{L}(L^p(\mathbb{R}^N))} = 1$ for all $t > 0$ then there are solutions that converge to 0 in $L^p(\mathbb{R}^N)$ arbitrarily slow.*

More precisely, for any continuous function $g : [0, \infty) \rightarrow (0, 1]$ such that $\lim_{t \rightarrow \infty} g(t) = 0$, there exists $u_0 \in L^p(\mathbb{R}^N)$ such that $\|S_{\mu,V}(t)u_0\|_{L^p} \rightarrow 0$ as $t \rightarrow \infty$ and

$$\limsup_{t \rightarrow \infty} \frac{\|S_{\mu,V}(t)u_0\|_{L^p}}{g(t)} = \infty.$$

To see this, observe first that $(-\Delta)^{\mu}$ is injective in $L^p(\mathbb{R}^N)$ for $1 \leq p < \infty$. Actually, from [15], Chapter 2, Corollary 2.42, page 125, we have that for a tempered distribution such that $(-\Delta)^{\gamma}\phi = 0$ then the Fourier transform $\hat{\phi}$ has support in $\{0\}$ and therefore ϕ is a polynomial.

Then using Corollary 1.1.4 in [20] we get that if $1 < p < \infty$ then $(-\Delta)^\mu$ has dense range in $L^p(\mathbb{R}^N)$. Since the semigroup $\{S_\mu(t)\}_{t \geq 0}$ is analytic and bounded in $L^p(\mathbb{R}^N)$ for $1 < p < \infty$, this in turn implies, by Remark 1.5 in [2], that for all $u_0 \in L^p(\mathbb{R}^N)$ we have $\|S_\mu(t)u_0\|_{L^p} \rightarrow 0$ as $t \rightarrow \infty$. From part (ii) in Proposition 3.3 we also have $\|S_{\mu,V}(t)u_0\|_{L^p} \rightarrow 0$ as $t \rightarrow \infty$.

Now assume by contradiction that for all $u_0 \in L^p(\mathbb{R}^N)$ there exists $C = C(u_0)$ such that

$$\frac{\|S_{\mu,V}(t)u_0\|_{L^p}}{g(t)} \leq C(u_0), \quad t \geq 0.$$

Then, the uniform bounded principle implies that, for some $M > 0$,

$$\frac{\|S_{\mu,V}(t)\|_{\mathcal{L}(L^p)}}{g(t)} \leq M.$$

But then for large enough t we have $\|S_{\mu,V}(t)\|_{\mathcal{L}(L^p)} \leq Mg(t) < 1$ which is a contradiction.

We also have the following result that implies that if there is exponential decay, then we can take $M = 1$ in (4.1).

Proposition 4.3. *Either $\|S_{\mu,V}(t)\|_{\mathcal{L}(L^2(\mathbb{R}^N))} = 1$ for all $t > 0$ or there exists $a > 0$ such that for all $1 \leq p \leq \infty$*

$$\|S_{\mu,V}(t)\|_{\mathcal{L}(L^p(\mathbb{R}^N))} \leq e^{-at}, \quad t \geq 0.$$

Proof. Again, as above, if for some $t_0 > 0$ we have $\|S_{\mu,V}(t_0)\|_{\mathcal{L}(L^2(\mathbb{R}^N))} < 1$, then we have (4.1) in $L^2(\mathbb{R}^N)$. From this and the results in Section 3 we see that the semigroup $\{e^{at}S_{\mu,V}(t)\}_{t \geq 0}$ in $L^2(\mathbb{R}^N)$ is bounded, analytic, C^0 , and is generated by the selfadjoint operator $\mathcal{A} = -(-\Delta)^\mu - V + a$ with the same domain $D(\mathcal{A})$ as $-(-\Delta)^\mu - V$. In particular, $D(\mathcal{A}) = H^{2\mu}(\mathbb{R}^N)$ if $p_0 \geq 2$.

In particular, using [21, Remark 5.4, Chapter 1], the right half space in \mathbb{C} is in the resolvent of \mathcal{A} and therefore the spectrum $\sigma(\mathcal{A})$ must be contained in $(-\infty, 0]$. Then [22, Proposition 12.8, p. 270] implies

$$\langle -\mathcal{A}\phi, \phi \rangle_{L^2(\mathbb{R}^N)} \geq 0 \quad \text{for all } \phi \in D(\mathcal{A})$$

as otherwise $-\mathcal{A}$ would have to possess a negative eigenvalue.

This in turn implies, using [21, Corollary 4.4] that this operator generates a semigroup of contractions, $\{e^{at}S_{\mu,V}(t)\}_{t \geq 0}$. That is, we have $M = 1$ in (4.1) when $p = 2$. See also [17].

But since

$$\langle -\mathcal{A}\phi, \phi \rangle_{L^2(\mathbb{R}^N)} = \int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}}\phi|^2 + \int_{\mathbb{R}^N} (V - a)|\phi|^2 \geq 0, \quad \phi \in D(\mathcal{A})$$

this also implies that we can use Proposition 3.1 with the potential $V - a$ to conclude that $-\mathcal{A} = -(-\Delta)^\mu - V + a$ defines an order preserving semigroup of contractions in $L^p(\mathbb{R}^N)$ for any $1 \leq p \leq \infty$, that is, $\{e^{at}S_{\mu,V}(t)\}_{t \geq 0}$ in $L^p(\mathbb{R}^N)$ is of contractions for any $1 \leq p \leq \infty$. ■

This result combined with Theorem 4.1 gives the following.

Corollary 4.4. *Either $\|S_{\mu,V}(t)\|_{\mathcal{L}(L^p(\mathbb{R}^N))} = 1$ for all $t > 0$ and all $1 \leq p \leq \infty$ or there exists $a > 0$ such that for all $1 \leq p \leq \infty$*

$$\|S_{\mu,V}(t)\|_{\mathcal{L}(L^p(\mathbb{R}^N))} \leq e^{-at}, \quad t \geq 0.$$

Proof. If for some $1 \leq p_0 \leq \infty$ and $t_0 > 1$ we have $\|S_{\mu,V}(t_0)\|_{\mathcal{L}(L^{p_0}(\mathbb{R}^N))} < 1$ then by Theorem 4.1 we get $\omega_2 > 0$ and Proposition 4.3 gives the result. ■

Now we give conditions on the potential to have exponential decay. Following [1] we consider the following

Definition 4.5. *The class \mathcal{G} consists of all open subsets of \mathbb{R}^N containing arbitrarily large balls, that is, the sets such that for any $r > 0$ there exists $x_0 \in \mathbb{R}^N$ such that the ball of radius r around x_0 is included in this set.*

Theorem 4.6. Assume that $0 \leq V \in L_U^{p_0}(\mathbb{R}^N)$ with $p_0 > \max\{\frac{N}{2\mu}, 1\}$ and there exists $M > 0$ such that $0 \leq V_M(x) = \min\{V(x), M\}$ satisfies

$$\int_G V_M(x) dx = \infty \quad \text{for all } G \in \mathcal{G}.$$

Then there exists $a > 0$, independent of $1 \leq p \leq \infty$, such that

$$\|S_{\mu,V}(t)\|_{\mathcal{L}(L^p(\mathbb{R}^N))} \leq e^{-at}, \quad t \geq 0.$$

Proof. Since $0 \leq V_M(x) \leq V(x)$, by Proposition 3.3, it is enough to prove the decay for $S_{\mu,V_M}(t)$. Therefore we can assume V is bounded.

From Corollary 4.4, to prove the result it is enough to find $t > 0$ such that

$$\|S_{\mu,V}(t)\|_{\infty \rightarrow \infty} = \|S_{\mu,V}(t)\mathbf{1}\|_{L^\infty(\mathbb{R}^N)} < 1.$$

For this, notice that with $u_0 = \mathbf{1}$, we have in particular that, from Proposition 3.4, $u(t) = S_{\mu,V}(t)\mathbf{1}$ satisfies

$$u(t) = S_\mu(t)\mathbf{1} - \int_0^t S_\mu(t-s)Vu(s) ds = \mathbf{1} - \int_0^t S_\mu(t-s)Vu(s) ds.$$

Substituting the expression above for $u(s)$ inside the integral term we get

$$u(t) = \mathbf{1} - \int_0^t S_\mu(t-s)V ds + \int_0^t S_\mu(t-s)V \int_0^s S_\mu(s-r)Vu(r) dr ds$$

We use that $0 \leq u(s) \leq S_\mu(s)\mathbf{1} = \mathbf{1}$ and $0 \leq V \leq \|V\|_\infty \mathbf{1}$ and then

$$\begin{aligned} \int_0^t S_\mu(t-s)V \int_0^s S_\mu(s-r)Vu(r) dr ds &\leq \|V\|_\infty \int_0^t S_\mu(t-s)V \int_0^s S_\mu(s-r)\mathbf{1} dr ds \\ &= \|V\|_\infty \int_0^t s S_\mu(t-s)V ds = \|V\|_\infty \int_0^t (t-s)S_\mu(s)V ds \leq \|V\|_\infty t \int_0^t S_\mu(s)V ds. \end{aligned}$$

Therefore

$$u(t) \leq \mathbf{1} - \int_0^t S_\mu(t-s)V ds + \|V\|_\infty t \int_0^t S_\mu(s)V ds = \mathbf{1} + (\|V\|_\infty t - 1) \int_0^t S_\mu(t-s)V ds.$$

From the results in Section 2

$$(S_\mu(s)V)(x) = \int_{\mathbb{R}^N} k_\mu(s, x, y)V(y) dy$$

where $k_\mu(s, x, y) = \frac{1}{t^{\frac{N}{2\mu}}} k_{0,\mu}\left(\frac{x-y}{t^{\frac{1}{2\mu}}}\right)$ is as in (2.2) and (2.9). Using [1, Proposition 1.4] we see in turn that there exist $c > 0$ and $r > 0$ such that

$$\int_{B(x,r)} V(y) dy \geq c \quad \text{for all } x \in \mathbb{R}^N.$$

For any $x \in \mathbb{R}^N$ and $s > 0$ we have

$$\begin{aligned} S_\mu(s)V(x) &= \int_{\mathbb{R}^N} \frac{1}{s^{\frac{N}{2\mu}}} k_{0,\mu}\left(\frac{x-y}{s^{\frac{1}{2\mu}}}\right) V(y) dy \\ &\geq \int_{B(x,r)} \frac{1}{s^{\frac{N}{2\mu}}} k_{0,\mu}\left(\frac{x-y}{s^{\frac{1}{2\mu}}}\right) V(y) dy \\ &\geq \inf_{|z| \leq r} \frac{1}{s^{\frac{N}{2\mu}}} k_{0,\mu}\left(\frac{z}{s^{\frac{1}{2\mu}}}\right) \int_{B(x,r)} V(y) dy \\ &\geq c \left\{ (4\pi s)^{-\frac{N}{2}} e^{-\frac{r^2}{4s}}, \quad \mu = 1 \right. \\ &\quad \left. \frac{s}{(s^\mu + r^2)^{\frac{N+2\mu}{2}}}, \quad 0 < \mu < 1 \right\} =: g_\mu(s) \end{aligned}$$

where we have used the estimate of fractional heat equation kernel from (2.2) and (2.9)-(2.10).

Denoting $\mathcal{G}_\mu(t) \stackrel{\text{def}}{=} \int_0^t g_\mu(s) ds$, we get, for $0 < t < \frac{1}{\|V\|_{L^\infty(\mathbb{R}^N)}}$ and $x \in \mathbb{R}^N$

$$(S_{\mu,V}(t)\mathbf{1})(x) \leq \mathbf{1} - (1 - t\|V\|_{L^\infty(\mathbb{R}^N)})\mathcal{G}_\mu(t) < 1.$$

■

In case V can be approximated by bounded potentials, we have the following result.

Proposition 4.7. *Assume $V \in L^1_U(\mathbb{R}^N)$ can be approximated in $L^1_U(\mathbb{R}^N)$ by bounded functions. Then, they are equivalent*

- (i) $\int_G V_M = \infty$ for all $G \in \mathcal{G}$ with some M .
- (ii) $\int_G V = \infty$ for all $G \in \mathcal{G}$.

Proof. Since $0 \leq V_M \leq V$, clearly (i) implies (ii).

Conversely, from [1, Proposition 1.4], (ii) is equivalent to the existence of $c > 0$ and $r > 0$ such that

$$\inf_{x \in \mathbb{R}^N} \int_{B(x,r)} V \geq c.$$

By a simple covering argument, independent of x , we get that $\sup_{x \in \mathbb{R}^N} \|V_M - V\|_{L^1(B(x,r))} \leq C\|V_M - V\|_{L^1_U(\mathbb{R}^N)} \rightarrow 0$ as $M \rightarrow \infty$ and therefore, for some M we have $\sup_{x \in \mathbb{R}^N} \|V_M - V\|_{L^1(B(x,r))} < \frac{c}{2}$. This implies that for all $x \in \mathbb{R}^N$

$$\int_{B(x,r)} V_M = \int_{B(x,r)} V + \int_{B(x,r)} (V_M - V) > c - \frac{c}{2} = \frac{c}{2}.$$

Again [1, Proposition 1.4] gives (i). ■

Remark 4.8. *Now we give an example that shows that in general in Proposition 4.7 (ii) does not imply (i). That is, we show that may have for all $M > 0$ and $r > 0$*

$$\inf_{x \in \mathbb{R}^N} \int_{B(x,r)} V_M = 0$$

but for some $r > 0$,

$$\inf_{x \in \mathbb{R}^N} \int_{B(x,r)} V > 0.$$

Denote by $\{Q_i\}$ the family of cubes centered at points $i \in \mathbb{Z}^N$ of integer coordinates in \mathbb{R}^N and with edges of length 1 parallel to the axes. Then notice that $B(i, 1) \subset Q_i$ and define in Q_i for $i \neq 0$,

$$V(x) = c|i|^N \mathcal{X}_{B(i, \frac{1}{|i|})}$$

and $V = \frac{1}{c} \mathcal{X}_{B(0,1)}$ in Q_0 where c is the measure of the unit ball.

Then

- (i) For every i , $\int_{Q_i} V = 1$ so $V \in L^1_U(\mathbb{R}^N)$.
- (ii) There exists an $r > 0$ such that for every $x \in \mathbb{R}^N$, $B(x, r)$ contains at least one cube Q_i . Then

$$\int_{B(x,r)} V \geq 1$$

- (iii) For every $M > 0$ and every $r > 0$ there exists $m(r) \in \mathbb{N}$ such that $B(x, r)$ is contained in $m(r)$ cubes and if $|x|$ is large then the centers of these cubes satisfy $|i| \geq |x| - 2r > 0$. For any such cubes

$$\int_{Q_i} V_M \leq \frac{cM}{|i|^N}$$

and then if we take $|x| \rightarrow \infty$ we see that

$$\int_{B(x,r)} V_M \leq m(r) \frac{cM}{(|x| - 2r)^N} \rightarrow 0.$$

Our next results characterises the exponential type of the fractional Schrödinger semigroup.

Theorem 4.9. *With the notations in Theorem 4.6, assume $0 < \mu \leq 1$ and $0 \leq V \in L_U^{p_0}(\mathbb{R}^N)$ with $p_0 > \max\{\frac{N}{2\mu}, 1\}$.*

Then

$$\omega_2 = a_* \stackrel{\text{def}}{=} \inf \left\{ \int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} \phi|^2 + \int_{\mathbb{R}^N} V |\phi|^2 : \phi \in C_c^\infty(\mathbb{R}^N), \|\phi\|_{L^2(\mathbb{R}^N)} = 1 \right\} \geq 0.$$

Moreover, if $a_ > 0$ then*

$$\int_G V(x) dx = \infty \quad \text{for all } G \in \mathcal{G}. \quad (4.3)$$

Conversely, if V can be approximated in $L_U^1(\mathbb{R}^N)$ by bounded functions, then (4.3) implies $a_ > 0$.*

Proof. From Proposition 4.3, for any $a < \omega_2$ we have that the semigroup $\{e^{at} S_{\mu,V}(t)\}_{t \geq 0}$ in $L^2(\mathbb{R}^N)$ is of contractions and the operator $\mathcal{A} = -(-\Delta)^\mu - V + a$ in $L^2(\mathbb{R}^N)$ satisfies

$$\langle -\mathcal{A} \phi, \phi \rangle_{L^2(\mathbb{R}^N)} \geq 0 \quad \text{for all } \phi \in C_c^\infty(\mathbb{R}^N) \subset D(\mathcal{A}).$$

In particular, taking $\phi \in C_c^\infty(\mathbb{R}^N)$ with $\|\phi\|_{L^2(\mathbb{R}^N)} = 1$ then $\int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} \phi|^2 + \int_{\mathbb{R}^N} V |\phi|^2 \geq a > 0$. Hence $a_* \geq a$ and therefore $a_* \geq \omega_2$.

Conversely, if $a < a_*$ we take $f(t) = e^{2at} \|u(t)\|_{L^2(\mathbb{R}^N)}^2$ with $u(t) = S_{\mu,V}(t)\phi$ and then

$$f'(t) = 2ae^{2at} \int_{\mathbb{R}^N} u(t)^2 + 2e^{2at} \int_{\mathbb{R}^N} u(t) \frac{d}{dt} u(t) = 2ae^{2at} \int_{\mathbb{R}^N} u(t)^2 + 2e^{2at} \int_{\mathbb{R}^N} u(t) (-(-\Delta)^\mu u(t) - V u(t))$$

and then

$$\frac{e^{-2at}}{2} f'(t) = - \int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} u(t)|^2 + V u(t)^2 + a \int_{\mathbb{R}^N} u(t)^2 \leq (a - a_*) \int_{\mathbb{R}^N} u(t)^2 \leq 0$$

which yields $f'(t) \leq 0$ and then $\|S_{\mu,V}(t)\|_{\mathcal{L}(L^2(\mathbb{R}^N))} \leq e^{-at}$. Therefore $\omega_2 \geq a_*$.

Now, if $a_* > 0$, assume that (4.3) fails, that is, there exists $G \in \mathcal{G}$ such that

$$\int_G V < \infty.$$

Then we choose a positive sequence satisfying $r_n > n^{\frac{1}{\mu}}$ and a sequence of points $\{x_n\} \subset \mathbb{R}^N$ such that $B(x_n, r_n) \subset G$ for each $n \in \mathbb{N}$.

Assume we had a sequence of functions such that $\{\phi_n\} \subset C_c^\infty(B(x_n, r_n))$, $\|\phi_n\|_{L^2(\mathbb{R}^N)} = 1$ such that

$$\lim_{n \rightarrow \infty} \|\phi_n\|_{L^\infty(\mathbb{R}^N)} = 0, \quad \lim_{n \rightarrow \infty} \|(-\Delta)^{\frac{\mu}{2}} \phi_n\|_{L^2(\mathbb{R}^N)} = 0.$$

Then we would have

$$\begin{aligned} 0 < a_* &\leq \int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} \phi_n|^2 + \int_{\mathbb{R}^N} V |\phi_n|^2 = \int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} \phi_n|^2 + \int_G V |\phi_n|^2 \\ &\leq \int_{\mathbb{R}^N} |(-\Delta)^{\frac{\mu}{2}} \phi_n|^2 + \|\phi_n\|_{L^\infty(\mathbb{R}^N)}^2 \int_G V \rightarrow 0 \quad \text{as } n \rightarrow \infty, \end{aligned}$$

which is a contradiction and therefore (4.3) must be true. To construct a sequence as above we take $\psi \in C_c^\infty(B(0, 1))$ such that $\int_{\mathbb{R}^N} |\psi|^2 = 1$ and defining

$$\psi_n(x) = n^{-\frac{N}{2\mu}} \psi(n^{-\frac{1}{\mu}} x), \quad x \in \mathbb{R}^N, \quad n \in \mathbb{N}$$

we see that $\{\psi_n\} \subset C_c^\infty(B(0, r_n))$, $\|\psi_n\|_{L^2(\mathbb{R}^N)} = 1$, $\lim_{n \rightarrow \infty} \|\psi_n\|_{L^\infty(\mathbb{R}^N)} = 0$ and

$$\lim_{n \rightarrow \infty} \|(-\Delta)^{\frac{\mu}{2}} \psi_n\|_{L^2(\mathbb{R}^N)} = \lim_{n \rightarrow \infty} n^{-1} \|(-\Delta)^{\frac{\mu}{2}} \psi\|_{L^2(\mathbb{R}^N)} = 0$$

since $(-\Delta)^{\frac{\mu}{2}}$ is an homogenous operator of degree μ . Then $\phi_n(\cdot) = \psi_n(\cdot + x_n)$, $n \in \mathbb{N}$, satisfies the conditions above since $(-\Delta)^{\frac{\mu}{2}}$ is invariant under translations, see (2.4).

Conversely, if V can be approximated in $L_U^1(\mathbb{R}^N)$ by bounded functions, then from Proposition 4.7 and Theorem 4.6 we get that (4.3) implies $a_* > 0$. ■

REFERENCES

- [1] W. Arendt and C. J. K. Batty. Exponential stability of a diffusion equation with absorption. *Differential Integral Equations*, 6(5):1009–1024, 1993. Cited ↑ in page: [1](#), [14](#), [15](#), [16](#)
- [2] W. Arendt, C. J. K. Batty, and P. Bényan. Asymptotic stability of Schrödinger semigroups on $L^1(\mathbf{R}^N)$. *Math. Z.*, 209(4):511–518, 1992. Cited ↑ in page: [14](#)
- [3] J. M. Arrieta, J. W. Cholewa, T. Dlotko, and A. Rodríguez-Bernal. Linear parabolic equations in locally uniform spaces. *Math. Models Methods Appl. Sci.*, 14(2):253–293, 2004. Cited ↑ in page: [9](#), [10](#)
- [4] J. M. Arrieta, J. W. Cholewa, T. Dlotko, and A. Rodríguez-Bernal. Asymptotic behavior and attractors for reaction diffusion equations in unbounded domains. *Nonlinear Anal.*, 56(4):515–554, 2004. Cited ↑ in page: [6](#)
- [5] K. Bogdan, T. Grzywny, T. Jakubowski, and D. Pilarczyk. Fractional Laplacian with Hardy potential. *Communications in Partial Differential Equations*, 44(1):20–50, 2019. Cited ↑ in page: [4](#)
- [6] M. Bonforte, Y. Sire, and J.L. Vázquez. Optimal existence and uniqueness theory for the fractional heat equation. *Nonlinear Analysis: Theory, Methods & Applications*, 153:142–168, 2017. Cited ↑ in page: [4](#)
- [7] T. Cazenave and A. Haraux. *An introduction to semilinear evolution equations*, volume 13 of *Oxford Lecture Series in Mathematics and its Applications*. The Clarendon Press, Oxford University Press, New York, 1998. Translated from the 1990 French original by Yvan Martel and revised by the authors. Cited ↑ in page: [2](#)
- [8] J. W. Cholewa and A. Rodríguez-Bernal. Sharp estimates for homogeneous semigroups in homogeneous spaces. Applications to PDEs and fractional diffusion in \mathbb{R}^N . *Commun. Contemp. Math.*, 24(1):Paper No. 2050070, 56, 2022. Cited ↑ in page: [3](#), [4](#)
- [9] J. W. Cholewa and A. Rodríguez-Bernal. On linear higher-order parabolic equations in Morrey spaces. *Anal. Appl. (Singap.)*, 21(6):1561–1608, 2023. Cited ↑ in page: [3](#), [4](#)
- [10] J. W. Cholewa and A. Rodríguez-Bernal. Self-similarity in homogeneous stationary and evolution problems. *J. Evol. Equ.*, 23(2):Paper No. 42, 39, 2023. Cited ↑ in page: [4](#)
- [11] E. B. Davies. *Heat kernels and spectral theory*, volume 92 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1989. Cited ↑ in page: [6](#)
- [12] A. De Pablo, F. Quirós, A. Rodríguez, and J.L. Vázquez. A general fractional porous medium equation. *Comm. Pure Appl. Math.*, LXV:1242–1284, 2012. Cited ↑ in page: [5](#)
- [13] E. Di Nezza, G. Palatucci, and E. Valdinoci. Hitchhiker’s guide to the fractional Sobolev spaces. *Bull. Sci. Math.*, 136(5):521–573, 2012. Cited ↑ in page: [3](#)
- [14] K.J. Engel and R. Nagel. *One-parameter semigroups for linear evolution equations*, volume 194 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2000. Cited ↑ in page:
- [15] L. Grafakos. *Classical Fourier analysis*, volume 249 of *Graduate Texts in Mathematics*. Springer, New York, third edition, 2014. Cited ↑ in page: [13](#)
- [16] D. Henry. *Geometric theory of semilinear parabolic equations*, volume 840 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1981. Cited ↑ in page: [11](#)
- [17] C. S. Kubrusly and N. Levan. On exponential stability of contraction semigroups. *Semigroup Forum*, 83(3):513–521, 2011. Cited ↑ in page: [14](#)
- [18] M. Kwaśnicki. Ten equivalent definitions of the fractional Laplace operator. *Fract. Calc. Appl. Anal.*, 20(1):7–51, 2017. Cited ↑ in page: [3](#)
- [19] A. Lunardi. *Analytic semigroups and optimal regularity in parabolic problems*. Progress in Nonlinear Differential Equations and their Applications, 16. Birkhäuser Verlag, Basel, 1995. Cited ↑ in page: [4](#)
- [20] C. Martínez Carracedo and M. Sanz Alix. *The theory of fractional powers of operators*, volume 187 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam, 2001. Cited ↑ in page: [14](#)
- [21] A. Pazy. *Semigroups of Linear Operators and Applications to Partial Differential Equations*. Springer-Verlag New York, Inc., 1983. Cited ↑ in page: [7](#), [8](#), [11](#), [14](#)
- [22] K. Schmüdgen. *Unbounded self-adjoint operators on Hilbert space*, volume 265 of *Graduate Texts in Mathematics*. Springer, Dordrecht, 2012. Cited ↑ in page: [14](#)
- [23] B. Simon. Schrödinger semigroups. *Bull. Amer. Math. Soc. (N.S.)*, 7(3):447–526, 1982. Cited ↑ in page: [13](#)
- [24] L. Weis. The stability of positive semigroups on L_p spaces. *Proc. Amer. Math. Soc.*, 123(10):3089–3094, 1995. Cited ↑ in page: [12](#)
- [25] L. Weis. A short proof for the stability theorem for positive semigroups on $L_p(\mu)$. *Proc. Amer. Math. Soc.*, 126(11):3253–3256, 1998. Cited ↑ in page: [12](#)
- [26] K. Yosida. *Functional analysis*. Springer-Verlag, Berlin-New York, fifth edition, 1978. Cited ↑ in page: [3](#), [4](#)