# Generalized cyclic symmetric decompositions for the matrix multiplication tensor

Charlotte Vermeylen<sup>a</sup>, Marc Van Barel<sup>b</sup>

<sup>a</sup>Department of Electrical Engineering (ESAT) KU Leuven, Kasteelpark Arenberg 10, Heverlee, 3001, Vlaams-Brabant, Belgium <sup>b</sup>Department of Computer Science KU Leuven, Celestijnenlaan 200C A, Heverlee, 3001, Vlaams-Brabant, Belgium

# Abstract

In this paper we propose a new generalized cyclic symmetric structure in the factor matrices of polyadic decompositions of matrix multiplication tensors for non-square matrix multiplication to reduce the number of variables in the optimization problem and in this way improve the convergence.

*Keywords:* matrix multiplication, tensors, optimization, canonical polyadic decomposition

# 1. Introduction

In this paper we consider the *fast matrix multiplication* (FMM) problem which rewrites matrix multiplication, which is a set of bilinear equations, i.e.,

$$C(i,j) = \sum_{k=1}^{p} A(i,k)B(k,n),$$
(1)

where C := AB, as a tensor equation:

$$C = \sum_{i,j,k=1}^{m,p,n} \left\langle E_{ik}^{m \times p}, A \right\rangle_{\mathrm{F}} \left\langle E_{kj}^{p \times n}, B \right\rangle_{\mathrm{F}} E_{ij}^{m \times n}$$
$$= \left( \sum_{i,j,k=1}^{m,p,n} E_{ik}^{m \times p} \otimes E_{kj}^{p \times n} \otimes E_{ij}^{m \times n} \right) \cdot_{\mathrm{F}} (A, B), \qquad (2)$$

Preprint submitted to Computational and Applied Mathematics April 26, 2024

where  $E_{ij}^{m \times p}$ , for i = 1, ..., m and j = 1, ..., p, is a basis of matrices in  $\mathbb{R}^{m \times p}$ such that  $E_{ij}^{m \times p}(i_1, i_2) = 1$  if  $(i_1, i_2) = (i, j)$  and zero else, ' $\otimes$ ' denotes the tensor product and  $\langle \cdot \rangle_{\rm F}$  and ' $\cdot_{\rm F}$ ' denote the Frobenius inner product. We add a transpose to (2) such that the tensor that is defined has additional interesting properties, such as *cyclic symmetry (CS)*, which will be discussed in more detail in Section 2.2.2:

$$C^{\top} = \left(\sum_{i,j,k=1}^{m,p,n} E_{ik}^{m \times p} \otimes E_{kj}^{p \times n} \otimes \left(E_{ij}^{m \times n}\right)^{\top}\right) \cdot_{\mathrm{F}} (A, B)$$
$$= \left(\sum_{i,j,k=1}^{m,p,n} E_{ik}^{m \times p} \otimes E_{kj}^{p \times n} \otimes E_{ji}^{n \times m}\right) \cdot_{\mathrm{F}} (A, B) . \tag{3}$$

Remark that the Frobenius inner product is a linear function in the elements of a matrix. The rank of the bilinear equation is the minimal r such that the equation can be written as

$$C^{\top} = \sum_{i=1}^{r} \left\langle U_i, A \right\rangle_{\mathrm{F}} \left\langle V_i, B \right\rangle_{\mathrm{F}} W_i, \tag{4}$$

for some matrices  $U_i \in \mathbb{R}^{m \times p}$ ,  $V_i \in \mathbb{R}^{p \times n}$ , and  $W_i \in \mathbb{R}^{n \times m}$ . It can be shown that minimizing r minimizes the computational complexity of matrix multiplication [1, Proposition 15.1], [2]. More specifically, the number of arithmetic operations needed to compute two matrices in  $\mathbb{R}^{n \times n}$  is  $\mathcal{O}(n^{\omega})$ , where  $\omega := \log_n r$ . Since  $r < n^3$ , it holds that  $\omega < 3$ . Additionally, it is well known that the exponent is bounded from below by two, since you need at least  $n^2$  operations to compute all  $n^2$  elements of the matrix multiplication.

Minimizing the rank of the bilinear equation corresponds to finding a (canonical) polyadic decomposition ((C)PD) of the so called matrix multiplication tensor (MMT) [3] defined implicitly in (3):

$$T_{mpn} := \sum_{i,j,k=1}^{m,p,n} e_{ik}^{mp} \otimes e_{kj}^{pn} \otimes e_{ji}^{mn},$$
(5)

where  $e_{ik}^{mp} := \text{vec}\left(E_{ik}^{m \times p}\right)$  and  $T_{mpn}$  is the MMT of dimension  $mp \times pn \times mn$ .

Consequently, (3) can also be formulated as

$$\operatorname{vec}(C^{\top}) = T_{mpn} \cdot_1 \operatorname{vec}(A) \cdot_2 \operatorname{vec}(B), \tag{6}$$

where vec (·) denotes the column-wise vectorization operator, and  $\cdot_1$  and  $\cdot_2$  denote the multiplication along the first and second dimension of  $T_{mpn}$  respectively.

Because of this definition,  $T_{mpn}$  is a sparse tensor consisting of mpn ones:

$$T_{mpn}(i_1 + (i_2 - 1)m, j_1 + (j_2 - 1)p, k_1 + (k_2 - 1)n) = \begin{cases} 1 & \text{if } i_1 = k_2, i_2 = j_1, j_2 = k_1 \\ 0 & \text{else} \end{cases},$$
(7)

for all  $i_1, k_2 = 1, \ldots, m, i_2, j_1 = 1, \ldots, p$ , and  $j_2, k_1 = 1, \ldots, n$ . Remark that a MMT only depends of the size of the matrices that are multiplied and not on the values.

A PD of  $T_{mpn}$  of rank r is a decomposition of  $T_{mpn}$  into r rank-1 tensors:

$$T_{mpn} = \sum_{i=1}^{r} u_i \otimes v_i \otimes w_i, \tag{8}$$

where ' $\otimes$ ' denotes the tensor product and  $u_i, v_i$ , and  $w_i$  are vectors in  $\mathbb{R}^{n_1}$ ,  $\mathbb{R}^{n_2}$ , and  $\mathbb{R}^{n_3}$  respectively. These vectors are collected in three factor matrices U, V, and  $W: U := [u_1, \dots, u_r], V := [v_1, \dots, v_r]$ , and  $W := [w_1, \dots, w_r]$ . A PD of rank r is written in short as PD<sub>r</sub> and a PD of rank r of a tensor Tas PD<sub>r</sub>(T). The minimal r for which a PD of a certain tensor exists is called the rank  $r^*$  of this tensor or  $r^*(T)$  and a PD of this rank a CPD or PD<sub>r\*</sub>(T).

With a PD<sub>r</sub> (8) of  $T_{mpn}$ , we obtain the following base algorithm to multiply any two matrices  $A \in \mathbb{R}^{m \times p}$  and  $B \in \mathbb{R}^{p \times n}$ :

$$\operatorname{vec}\left(C^{\top}\right) = \sum_{i=1}^{r} \langle u_i, \operatorname{vec}(A) \rangle \langle v_i, \operatorname{vec}(B) \rangle w_i,$$
(9)

which is the same as (4) when  $U_i = \operatorname{reshape}(u_i, [m, p]), V_i = \operatorname{reshape}(v_i, [p, n]),$ and  $W_i = \operatorname{reshape}(w_i, [n, m])$ , for all  $i = 1, \ldots, r$ . The number of multiplications between (linear combinations of) elements of A and B in (9) is reduced to r < mpn, where the upper bound holds for standard matrix multiplication. Such multiplications are called 'active' because, when these algorithms are applied recursively, they determine the asymptotic complexity of matrix multiplication [4].

One of the difficulties is that the rank of a CPD of  $T_{mpn}$ , denoted by  $r^*(T_{mpn})$ , is not known for most combinations of (m, p, n). Counterexamples are  $T_{222}$ , for which the set of PDs is completely understood and  $r^*$  is known to be seven [5],  $T_{223}$ , for which the rank is known to be 11 [6], and  $T_{224}$ , for which the rank is known to be 14 [7]. We call PD<sub>7</sub>( $T_{222}$ ) originally discovered by Strassen PD<sub>Strassen</sub> [3]. For other combinations of m, p, and n, lower bounds on the rank exist but none of these lower bounds are attained in practice. For example, the lower bound on  $r^*(T_{333})$  is proven to be 19 [8], and the upper bound is 23 [9, 10, 11, 12]. We call the lowest rank for which an exact PD of  $T_{mpn}$  is known in the literature  $\tilde{r}(T_{mpn})$ . For an overview of  $\tilde{r}(T_{mpn})$  for various m, p, and n, we refer to [11, Table 1] and [13, Fig. 1]. Remark that the rank of  $T_{mpn}$  is equal to the rank of  $T_{pnm}$  or any other permutation of m, p, and n. This is true because we can obtain a PD<sub>r</sub> of  $T_{pnm}$  and  $T_{nmp}$  by cyclically permuting the factors in the PD:

$$T_{pnm} = \sum_{i=1}^{r} v_i \otimes w_i \otimes u_i, \qquad \qquad T_{nmp} = \sum_{i=1}^{r} w_i \otimes u_i \otimes v_i,$$

and additionally, it can be shown that  $T_{mnp} = \sum_{i=1}^{r} \operatorname{vec} (W_i^{\top}) \otimes \operatorname{vec}(V_i^{\top}) \otimes \operatorname{vec}(U_i^{\top})$ , and similarly for  $T_{npm}$  and  $T_{pmn}$ .

To find a PD<sub>r</sub> of  $T_{mpn}$ , we use the following *nonlinear least squares (NLS)* cost function, which is also frequently used in the literature for the FMM problem [11, 14, 10, 15]:

$$\min_{x} \underbrace{\frac{1}{2} \|F(x) - \operatorname{vec}(T_{mpn})\|^{2}}_{=:f(x)},$$
(10)

where  $\|\cdot\|$  denotes the  $\ell_2$ -norm, and F(x) is a vector function defined as

$$F(x) := \operatorname{vec}\left(\sum_{r=1}^{r} u_r \otimes v_r \otimes w_r\right), \qquad x := \begin{bmatrix} \operatorname{vec}(U) \\ \operatorname{vec}(V) \\ \operatorname{vec}(W) \end{bmatrix}.$$
(11)

Remark that  $f(x^*) = 0$ , if and only if  $x^*$  is a PD<sub>r</sub> of  $T_{mpn}$ . However, most

standard optimization algorithms fail to converge to a global minimum of (10). One of the reasons is that PDs of MMT have additional continuous invariances compared to generic PDs, which we will review in Section 2.2.1 [16].

In practice, the minimal length or rank  $r^*$  is approximated experimentally by the lowest length  $\tilde{r}$  for which a solution  $x^*$  with  $f(x^*) = 0$  can be obtained using numerical optimization. However, no globally convergent algorithm for (10) is known to us, and thus we can never be sure that no solution with a lower rank exists.

Since the total number of elements of the factor matrices grows rapidly with the size of the matrices, solving (10) is only feasible for relatively small values of m, p, and n, e.g.,  $m, p, n \leq 5$ . In this paper we give a new structure that can be enforced in PDs of MMTs to reduce the number of variables in the optimization problem and hereby improve the convergence.

To solve (10), in the literature the alternating least squares (ALS) method is frequently used [11, 17, 15, 10] but the convergence is usually slow and a lot of starting points are needed to obtain a solution in a reasonable amount of time. In [14] and [11], a constrained optimization problem and corresponding method are proposed to improve the convergence. More specifically, a Levenberg-Marquardt (LM) method with Lagrange parameters and a quadratic penalty (QP) term in combination with the ALS method are used respectively. However, both methods were still not able to find PDs of rank 49 of MMTs for  $4 \times 4$  matrix multiplication whereas these PDs are known to exist. That is why we proposed in [18] an augmented Lagrangian (AL) method and a new constrained optimization problem to find PDs of MMTs:

$$\min_{x \in \mathcal{T}} f(x), \quad \text{s.t.} \quad h(x) = 0, \quad l \le x \le u.$$
(12)

Different equality constraints h(x) were proposed. The AL method can be considered as a combination of the LM and QP method since the AL objective function is:

$$\min_{x,y_1,y_2,z} \underbrace{f(x) + \langle y_1, h(x) \rangle + \langle y_2, x - z \rangle + \frac{\beta}{2} \left( \|h(x)\|^2 + \|x - z\|^2 \right)}_{=:\mathcal{L}_A(x,y_1,y_2,z;\beta)},$$

subject to (s.t.)  $l \leq z \leq u$ , where  $y_1$  and  $y_2$  are vectors of Lagrange multipliers,  $\beta$  is the regularization parameter, and z is a vector of slack variables

that enables us to rewrite the inequality constraint as an equality constraint. Because f(x) is an NLS function, the AL objective can also be rewritten as an NLS function. The variables in x and the auxiliary parameters are updated successively according to [19, Section 17.4]. We use the LM method for the minimization to x. The pseudo-code of the algorithm is shown in [18, Algorithm 4.2] and the MATLAB implementation is publicly available <sup>1</sup>. The advantage compared to the QP method is that the constraints are satisfied more accurately during optimization. Using this method, we were able to obtain new PD<sub>49</sub> ( $T_{444}$ )s and also different new CS PDs. We will use this method in the numerical experiments in Section 4.

The rest of the paper is organized as follows. In Section 2, we give some preliminaries concerning the FMM problem. In Section 3, the new generalized CS structure is proposed, and in Section 4, numerical experiments are given to demonstrate the use and advantages of including this structure in the optimization problem (10).

# 2. Preliminaries

In this section, we first discuss what we mean with practical PDs and FMM algorithms. Afterwards, we give some well known properties of MMTs.

#### 2.1. Practical algorithms

As discussed in the introduction, a  $\text{PD}_r(T_{mpn})$  can be applied recursively to, e.g., multiply matrices of size  $m^k \times p^k$  and  $p^k \times n^k$ . Such a recursive algorithm requires  $\mathcal{O}(cr^k)$  operations, where the constant c depends on the number of scalar multiplications and nonzeros in the factor matrices. When k is sufficiently large and c sufficiently small, fewer operations are required compared to standard matrix multiplication. To decrease the constant c, we search for sparse PDs, with elements in a discrete set, e.g.,  $\{-1, 0, 1\}$ . This set can be extended with powers of 2 since this is not a costly operation when implemented in hardware. Such PDs are called *practical PDs* as they result in practical FMM algorithms.

Consequently, if we obtain a numerical PD of  $T_{mpn}$  using numerical optimization, it still has to be transformed into a practical one. The *invariance* transformations or *inv-transformations* discovered in [16], which are further

<sup>&</sup>lt;sup>1</sup>https://github.com/CharlotteVermeylen/ALM\_FMM\_stability

discussed in Section 2.2.1, can be used to obtain such PDs [14]. However, not all PDs of  $T_{mpn}$  can be transformed into a practical PD in this way [20]. In [20], this process is called *discretization*.

A disadvantage of (10) is that the solutions have floating point elements and thus extra steps have to be taken or constraints have to be added to the optimization problem to obtain practical PDs. That is why we proposed a new constrained problem formulation in [18] with equality constraint:

$$h_{\text{discr}}(x) := x \cdot (x-1) \cdot (x+1), \tag{13}$$

where  $\cdot$  denotes element-wise multiplication.

#### 2.2. Properties of the matrix multiplication tensor

In this section, first the invariance transformations that are present for all PDs of  $T_{mpn}$  are discussed. Additionally, a short overview of what is known in the literature about the use of these invariance transformations to obtain practical PDs is given. Afterwards, cyclic symmetry and recursive PDs are discussed in more detail.

## 2.2.1. Invariance transformations

The representation of a PD via factor matrices is not unique. For example when permuting the rank-1 tensors, and consequently the columns of the factor matrices according to

$$(U, V, W) \to (U', V', W'),$$

where  $u'_i = u_{\sigma(i)}$ , for all i = 1, ..., r, and  $\sigma = (\sigma(1) \ \sigma(2) \ \sigma(3) \ ... \ \sigma(r))$  is an element of the permutation group of r elements, then the same tensor is obtained:

$$\sum_{i=1}^{r} u_i \otimes v_i \otimes w_i = \sum_{i=1}^{r} u'_i \otimes v'_i \otimes w'_i.$$

Additionally, because of the multi-linearity of the tensor product, the columns of each rank-1 tensor can be scaled as:  $\alpha_i u_i, \beta_i v_i, \frac{1}{\alpha_i \beta_i} w_i$ , for all  $i = 1, \ldots, r$ , and  $\alpha_i, \beta_i$  in  $\mathbb{R}_0$ , without changing the rank-1 tensors because

$$\alpha_i u_i \otimes \beta v_i \otimes \frac{1}{\alpha_i \beta_i} w_i = u_i \otimes v_i \otimes w_i.$$

These two invariances hold for all PDs. Remark that only the scaling invariance is a continuous invariance transformation (of dimension 2r).

For PDs of  $T_{mpn}$ , additional invariances hold [16]. More specifically, it is well known that these PDs are invariant under the following PQR-transformation:

$$u'_i \leftarrow \operatorname{vec}\left(PU_iQ^{-1}\right), \quad v'_i \leftarrow \operatorname{vec}\left(QV_iR^{-1}\right), \quad w'_i \leftarrow \operatorname{vec}\left(RW_iP^{-1}\right), \quad (14)$$

for i = 1, ..., r, and where  $P \in GL(m)$ ,  $Q \in GL(p)$ , and  $R \in GL(n)$ , where GL(i) represents the general linear group of invertible matrices in  $\mathbb{R}^{i \times i}$ . This invariance can easily be proven using the fact that  $C := AB = (P^{\top})^{-1} (P^{\top}A (Q^{\top})^{-1}) (Q^{\top}B (R^{\top})^{-1}) R^{\top}$ . When we substitute this in (9), we obtain:

$$C = \sum_{i=1}^{r} \operatorname{trace} \left( U_i \left( P^{\top} A \left( Q^{\top} \right)^{-1} \right)^{\top} \right)$$
$$\operatorname{trace} \left( V_i \left( Q^{\top} B \left( R^{\top} \right)^{-1} \right)^{\top} \right) \left( P^{\top} \right)^{-1} W_i^{\top} R^{\top}$$
$$= \sum_{i=1}^{r} \operatorname{trace} \left( P U_i Q^{-1} A^{\top} \right) \operatorname{trace} \left( Q V_i R^{-1} B^{\top} \right) \left( R W_i P^{-1} \right)^{\top},$$

where we made use of the fact that the trace is invariant under cyclic permutation of the factors. And thus indeed another base algorithm or  $PD_r$  of  $T_{mpn}$  can be obtained using (14). Additionally, when m = n = p, PDs of  $T_{mmm}$  are invariant under the following *transpose-transformation*:

$$u'_i \leftarrow \operatorname{vec}\left(V_i^{\top}\right), \quad v'_i \leftarrow \operatorname{vec}\left(U_i^{\top}\right), \quad w'_i \leftarrow \operatorname{vec}\left(W_i^{\top}\right),$$

for i = 1, ..., r, which can be proven using the fact that  $C = (B^{\top}A^{\top})^{\top}$ . Lastly, still for m = n = p, the PDs are invariant under cyclic permutation of the factor matrices:

$$\sum_{i=1}^{r} u_i \otimes v_i \otimes w_i = \sum_{i=1}^{r} v_i \otimes w_i \otimes u_i = \sum_{i=1}^{r} w_i \otimes u_i \otimes v_i,$$

which holds because  $T_{mmm}$  is a CS tensor.

Remark that concerning the specific invariances of PDs of  $T_{mpn}$ , only the

PQR-invariance is a continuous invariance and it overlaps with the scaling invariance when P = aI, Q = bI, and R = cI, where I is the identity matrix of the appropriate size and a, b, and c are scaling factors in  $\mathbb{R}_0$ , because in this case the PQR-transformation scales the columns of the factor matrix Uwith  $\frac{a}{b}$ , all columns of V with  $\frac{b}{c}$ , and all columns of W with  $\frac{c}{a}$ , and indeed the product of these factors equals one as for the scaling invariance. Thus, the combination of the scaling and PQR-transformation has at most dimension  $2r + m^2 + p^2 + n^2 - 3$  for all PDs of  $T_{mpn}$ .

We call the combination of the invariance transformations described above *inv-transformations*. Two PDs of  $T_{mpn}$  that can be obtained using inv-transformations are called *inv-equivalent*. Only for  $T_{222}$ , it is known that all PDs are inv-equivalent [5].

#### 2.2.2. Cyclic symmetry

Because of its definition,  $T_{mpn}$  is a structured and more specifically cyclic symmetric (CS) tensor when m = p = n. This means that  $T_{mmm}(i, j, k) = T_{mmm}(j, k, i) = T_{mmm}(k, i, j)$ , for all  $i, j, k = 1, \ldots, m^2$ .

We can make use of this property for constructing a PD by requiring that the rank-1 tensors are symmetric or occur in CS pairs [21]:

$$T_{mmm} = \sum_{i=1}^{s} a_i \otimes a_i \otimes a_i + \sum_{j=1}^{t} \left( b_j \otimes d_j \otimes c_j + c_j \otimes b_j \otimes d_j + d_j \otimes c_j \otimes b_j \right),$$

where  $a_i, b_j, c_j$ , and  $d_j$  are vectors in  $\mathbb{R}^{m^2}$ , for all  $i = 1, \ldots, s$  and  $j = 1, \ldots, t$ , where s and t are parameters denoting respectively the number of symmetric and CS pairs of asymmetric rank-1 tensors. The length then equals r = s+3t. If we define the matrices  $A := [a_1 \cdots a_s], B := [b_1 \cdots b_t], C := [c_1 \cdots c_t],$ and  $D := [d_1 \cdots d_t]$ , the factor matrices can be written in function of these matrices:

$$U = \begin{bmatrix} A & B & C & D \end{bmatrix}, \quad V = \begin{bmatrix} A & D & B & C \end{bmatrix}, \quad W = \begin{bmatrix} A & C & D & B \end{bmatrix}.$$

Consequently, the number of unknowns is reduced by a factor 3, which makes this CS structure very useful for larger problem parameters, e.g.,  $m, p, n \ge 4$ . Furthermore, the cost function can then be changed to

$$\min_{A,B,C,D} \frac{1}{2} \left( \sum_{i=1}^{m^2} \sum_{j=i}^{m^2} \sum_{k=i+1}^{m^2} \left( T_{mmm}(i,j,k) - \sum_{i'=1}^{s} a_{ii'} a_{ji'} a_{ki'} - \sum_{j'=1}^{t} \left( b_{ij'} d_{jj'} c_{kj'} + c_{ij'} b_{jj'} d_{kj'} + d_{ij'} c_{jj'} b_{kj'} \right) \right)^2 + \sum_{i=1}^{m^2} \left( T_{mmm}(i,i,i) - \sum_{i'=1}^{s} a_{ii'}^3 - 3 \sum_{j'=1}^{t} \left( b_{ij'} d_{ij'} c_{ij'} \right) \right)^2 \right),$$
(15)

where  $a_{ii'} := A(i, i')$ ,  $b_{ii'} := B(i, i')$  and similarly for  $c_{ii'}$  and  $d_{ii'}$ , which reduces the number of rows in the Jacobian matrix used in NLS optimization methods to solve (15) from  $m^6$  to  $\frac{1}{3}(m^6 - m^2) + m^2$ . Remark that this cost function only includes one element for each CS pair of points because the CS structure already ensures that these elements are equal. A disadvantage of including this structure may be that by restricting the search space we might not be able to find the most sparse or stable PDs of  $T_{mmm}$ . Furthermore, it is not known if the rank of a PD of  $T_{mmm}$  with this structure equals the canonical rank.

Different practical CS PDs of rank 7 for  $T_{222}$  and of rank 23 for  $T_{333}$  are known in the literature [15, 10]. These PDs were found using an ALS method and further investigated using algebraic geometry and group theory. Strassen's decomposition satisfies s = 1 and t = 2. Although all PD<sub>7</sub> ( $T_{222}$ )s are known to be inv-equivalent, PD<sub>Strassen</sub> can be transformed into a practical PD with CS parameters (s, t) = (4, 1). However, this PD contains more nonzeros than PD<sub>Strassen</sub> and thus is not used in practice.

#### 2.3. Decompositions obtained by recursion

As mentioned in the introduction, an FMM algorithm applies a base algorithm for small m, p, and n, to large matrices, e.g., of size  $m^k \times p^k$ and  $p^k \times n^k$ . In the same way, we can also obtain PDs of rank 49 of  $T_{444}$ using a PD<sub>7</sub> ( $T_{222}$ ) one time recursively. The PD<sub>49</sub> ( $T_{444}$ ) that is obtained using PD<sub>Strassen</sub> is called PD<sup>rec</sup><sub>Strassen</sub> in the rest of the text. In the following proposition, we give the formulas to obtain the factor matrices of a recursive PD from the factor matrices of the original PD. This result is well known, e.g., from the supplementary material of [13], and in [22], and a proof can be found, e.g., in [23, Section 2.2.3]. **Proposition 2.1** (Recursive PD). If U, V, and W are the factor matrices of a  $PD_r(T_{mpn})$ , then the factor matrices U', V', and W', where

$$u'_{i'} := \operatorname{vec} \left( U_{i_1} \otimes U_{i_2} \right), \qquad v'_{i'} := \operatorname{vec} \left( V_{i_1} \otimes V_{i_2} \right), \qquad (16)$$

$$w'_{i'} := \operatorname{vec} \left( W_{i_1} \otimes W_{i_2} \right), \qquad i' := i_2 + (i_1 - 1)r, \qquad (17)$$

for  $i_1, i_2 = 1, \ldots, r$ , are the factor matrices of a  $PD_{r^2}(T_{m^2p^2n^2})$ .

The following corollary gives the CS parameters and factor matrices of a recursive PD as a function of the original PD.

**Corollary 2.2.** If U, V, and W are CS factor matrices of a  $PD_r(T_{mpn})$  with CS parameters s and t, then the factor matrices U', V', and W' of the recursive decomposition  $PD_{r^2}^{rec}(T_{m^2p^2n^2})$  is also CS with parameters  $s' = s^2$  and t' = t(s+r):

$$U' := \begin{bmatrix} A' & B' & C' & D' \end{bmatrix}, \quad V' := \begin{bmatrix} A' & D' & B' & C' \end{bmatrix}, \quad W' := \begin{bmatrix} A' & C' & D' & B' \end{bmatrix},$$

where

$$A'_{i'} := A_{i_1} \otimes A_{i_2}, \qquad i' := i_2 + (i_1 - 1)s, \qquad i_1, i_2 = 1, \dots, s, \tag{18}$$

and

$$B' := \begin{bmatrix} B'_1 & B'_2 \end{bmatrix}, \qquad C' := \begin{bmatrix} C'_1 & C'_2 \end{bmatrix}, \qquad D' := \begin{bmatrix} D'_1 & D'_2 \end{bmatrix}, \qquad (19)$$

where

$$B'_{1,j'} := A_{i_1} \otimes B_j, \qquad C'_{1,j'} := A_{i_1} \otimes C_j, \qquad D'_{1,j'} := A_{i_1} \otimes D_j, \qquad (20)$$
$$B'_{2,k'} := B_j \otimes U_k, \qquad C'_{2,k'} := C_j \otimes W_k, \qquad D'_{2,k'} := D_j \otimes V_k,$$

where  $j' := j + (i_1 - 1)t$  and k' := k + (j - 1)r, for j = 1, ..., t and k = 1, ..., r.

*Proof.* Using Proposition 2.1, we know that  $U'_{i'} = U_{i_1} \otimes U_{i_2}$ ,  $V'_{i'} = V_{i_1} \otimes V_{i_2}$ , and  $W'_{i'} = W_{i_1} \otimes W_{i_2}$ , where  $i' := i_2 + (i_1 - 1)r$ . Thus, the symmetric part of the recursive PD must satisfy:  $U_{i_1} = V_{i_1} = W_{i_1}$  and  $U_{i_2} = V_{i_2} = W_{i_2}$ . Consequently,  $i_1$  and  $i_2$  must be smaller than or equal to s, such that  $U_{i_1} =$  $V_{i_1} = W_{i_1} = A_{i_1}$  and  $U_{i_2} = V_{i_2} = W_{i_2} = A_{i_2}$ . For the other values of  $i_1$  and  $i_2$ , the columns still appear in CS pairs and they can be rearranged as in (20) to satisfy the CS structure. In [23, Appendix A], the different columns that are obtained by using Proposition 2.1 are written out to see this more clearly.  $\hfill \Box$ 

Remark that, since  $PD_{Strassen}$  satisfies s = 1 and t = 2, the parameters of  $PD_{Strassen}^{rec}$  are s = 1 and t = 16. Another CS  $PD_{49}(T_{444})$  can be obtained with (s,t) = (16,11) by using a  $PD_7(T_{222})$  with (s,t) = (4,1) one time recursively.

#### 2.4. Jacobian matrix

In [18], we showed that the Jacobian matrix can be used to investigate the inv-equivalence of PDs of MMTs and to investigate any additional invariances, such as the ones discovered in [18] using parametrizations of decompositions. That is why in Section 4, we will give the size and ranks of the Jacobian matrix at solutions with the structure proposed in the next section.

# 3. Generalization cyclic symmetry

In this section, we give a generalization of the CS structure discussed in Section 2.2.2, to the case when m, n, and p are not equal. We do this to reduce the number of variables and reduce the search space to hopefully speed up the convergence. Note however that similarly as for the CS structure discussed in Section 2.2.2, it is not known if the rank of PDs with this structure equals the canonical rank. Because we can obtain a PD of  $T_{pnm}$  or another permutation of m, p, and n from a PD of  $T_{mpn}$ , we assume in this chapter that  $m \leq p \leq n$ . We know from (7) that:

$$T_{mpn}(i_1 + (i_2 - 1)m, j_1 + (j_2 - 1)p, k_1 + (k_2 - 1)n)$$
  
=  $\sum_{i=1}^r U_i(i_1, i_2)V_i(j_1, j_2)W_i(k_1, k_2) = \begin{cases} 1 & \text{if } i_1 = k_2, i_2 = j_1, j_2 = k_1 \\ 0 & \text{else} \end{cases}$ ,

for all  $i_1, k_2 = 1, ..., m, i_2, j_1 = 1, ..., p$ , and  $j_2, k_1 = 1, ..., n$ . Consequently, as long as  $i_1, i_2, j_1, j_2, k_1, k_2 \leq m$ , it holds that

$$T_{mpn}(i_1 + (i_2 - 1)m, j_1 + (j_2 - 1)p, k_1 + (k_2 - 1)n)$$
  
=  $T_{mpn}(k_1 + (k_2 - 1)m, i_1 + (i_2 - 1)p, j_1 + (j_2 - 1)n)$   
=  $T_{mpn}(j_1 + (j_2 - 1)m, k_1 + (k_2 - 1)p, k_1 + (k_2 - 1)n)$ 

And thus,

$$\sum_{i=1}^{r} U_i(i_1, i_2) V_i(j_1, j_2) W_i(k_1, k_2) = \sum_{i=1}^{r} U_i(k_1, k_2) V_i(i_1, i_2) W_i(j_1, j_2)$$
$$= \sum_{i=1}^{r} U_i(j_1, j_2) V_i(k_1, k_2) W_i(i_1, i_2),$$

which can be used to include a CS structure in a subpart of the factor matrices. An illustration is shown in Figure 1, where the following submatrices are defined in function of the position in the factor matrices:

$$U_{i}(i_{1}, i_{2}) =: \begin{cases} A_{i}(i_{1}, i_{2}) & \text{if } i_{2} \leq m, i \leq s \\ B_{i-s}(i_{1}, i_{2}) & \text{if } i_{2} \leq m, s < i \leq s+t \\ C_{i-s-t}(i_{1}, i_{2}) & \text{if } i_{2} \leq m, s+t < i \leq s+2t \\ D_{i-s-2t}(i_{1}, i_{2}) & \text{if } i_{2} \leq m, s+2t < i \leq r_{\text{CS}} \\ \tilde{U}_{i}(i_{1}, i_{2} - m) & \text{if } i_{2} > m, i \leq r_{\text{CS}} \\ \dot{U}_{i-r_{\text{CS}}}(i_{1}, i_{2}) & \text{if } r_{\text{CS}} < i \end{cases},$$

where  $r_{\rm CS} := s + 3t$ , and for i = 1, ..., r,  $i_1 = 1, ..., m$ , and  $i_2 = 1, ..., p$ ,

$$V_{i}(j_{1}, j_{2}) =: \begin{cases} A_{i}(j_{1}, j_{2}) & \text{if } j_{1}, j_{2} \leq m, i \leq s \\ D_{i-s}(j_{1}, j_{2}) & \text{if } j_{1}, j_{2} \leq m, s < i \leq s+t \\ B_{i-s-t}(j_{1}, j_{2}) & \text{if } j_{1}, j_{2} \leq m, s+t < i \leq s+2t \\ C_{i-s-2t}(j_{1}, j_{2}) & \text{if } j_{1}, j_{2} \leq m, s+2t < i \leq r_{\text{CS}} \\ \hat{V}_{i}(j_{1} - m, j_{2}) & \text{if } j_{1} > m, j_{2} \leq m, i \leq r_{\text{CS}} \\ \tilde{V}_{i}(j_{1}, j_{2} - m) & \text{if } j_{2} > m, i \leq r_{\text{CS}} \\ \dot{V}_{i-r_{\text{CS}}}(j_{1}, j_{2}) & \text{if } r_{\text{CS}} < i \end{cases}$$

for  $j_1 = 1, ..., p$ , and  $j_2 = 1, ..., n$ , and

$$W_{i}(k_{1},k_{2}) =: \begin{cases} A_{i}(k_{1},k_{2}) & \text{if } k_{1} \leq m, i \leq s \\ B_{i-s}(k_{1},k_{2}) & \text{if } k_{1} \leq m, s < i \leq s+t \\ C_{i-s-t}(k_{1},k_{2}) & \text{if } k_{1} \leq m, s+t < i \leq s+2t \\ D_{i-s-2t}(k_{1},k_{2}) & \text{if } k_{1} \leq m, s+2t < i \leq r_{\text{CS}} \\ \hat{W}_{i}(k_{1}-m,k_{2}) & \text{if } k_{1} > m, i \leq r_{\text{CS}} \\ \dot{W}_{i-r_{\text{CS}}}(k_{1},k_{2}) & \text{if } r_{\text{CS}} < i \end{cases},$$

for  $k_1 = 1, ..., n$ , and  $k_2 = 1, ..., m$ , where  $A_i := \text{reshape}(a_i, m \times m)$ , for i = 1, ..., s,  $B_j := \text{reshape}(b_j, m \times m)$ , for j = 1, ..., t, and similarly for  $C_j$  and  $D_j$ , and

$$\begin{split} \tilde{U}_k &:= \operatorname{reshape} \left( \tilde{U}(:,k), \ m \times (p-m) \right), \\ \tilde{V}_k &:= \operatorname{reshape} \left( \tilde{V}(:,k), \ p \times (n-m) \right), \\ \hat{V}_k &:= \operatorname{reshape} \left( \hat{V}(:,k), \ (p-m) \times m \right), \\ \hat{W}_k &:= \operatorname{reshape} \left( \hat{W}(:,k), \ (n-m) \times m \right), \end{split}$$

for  $k = 1, \ldots, r_{\text{CS}}$ , and

$$\begin{split} \dot{U}_{i'} &:= \operatorname{reshape} \big( \dot{U}(:,i'), \ m \times p \big), \\ \dot{V}_{i'} &:= \operatorname{reshape} \big( \dot{V}(:,i'), \ p \times n \big), \\ \dot{W}_{i'} &:= \operatorname{reshape} \big( \dot{W}(:,i'), \ n \times m \big), \end{split}$$

for  $i' = 1, \ldots, (r - r_{\rm CS})$ , and thus  $\tilde{U} \in \mathbb{R}^{m(p-m) \times r_{\rm CS}}$ ,  $\tilde{V} \in \mathbb{R}^{p(n-m) \times r_{\rm CS}}$ ,  $\hat{W} \in \mathbb{R}^{(n-m)m \times r_{\rm CS}}$ ,  $\dot{U} \in \mathbb{R}^{mp \times (r-r_{\rm CS})}$ ,  $\dot{V} \in \mathbb{R}^{pn \times (r-r_{\rm CS})}$ , and  $\dot{W} \in \mathbb{R}^{nm \times (r-r_{\rm CS})}$ . The proportions in Figure 1 are those of  $T_{234}$ , (s,t) := (2,2), and r := 20. We show in the experiments that such PDs indeed exist. Remark that, contrary to the case when m, n, and p are equal, the CS rank  $r_{\rm CS}$  does not equal the rank r, because this would be too restrictive. Also in the experiments that follow we did not find PDs for which both are equal. This is likely because of the interaction of the CS part with the parts  $\tilde{U}$ ,  $\tilde{V}$ ,  $\hat{V}$ , and  $\hat{W}$ . However, including this structure still reduces the number of parameters from (mp + np + mn)r to  $(mp + np + mn)r - 2m^2r_{\rm CS}$ .



Figure 1: Illustration of the generalized CS factor matrices of a  $PD_{20}(T_{234})$ , with (s,t) := (2,2).

# 4. Numerical experiments

In this section we show the results that we obtained with the AL method from [18] to find PDs of MMTs with the generalized CS structure for different combinations of s and t. The AL method takes as input an upper and lower bound on the elements in the factor matrices:  $l \leq x \leq u$ . To show the advantage of the new structure, we set u := -l := 1 and generate 50 random starting points of size  $10^{-2}$  with the built-in function randn of MATLAB:

$$x_0 := 10^{-2} \cdot \text{randn} \left( (mp + np + mn)r - 2m^2 r_{\rm CS} \times 1 \right).$$
 (21)

The number of inner iterations of the LM method was set to 50 and the number of outer iterations to 15. The tolerance on the gradient and on the bound constraint were both set to  $10^{-13}$ .

In Table 1, we show the results that we obtained with the AL method from [18] to find PD<sub>11</sub> ( $T_{223}$ )s for different combinations of s and t. The second to last column (from left to right) indicates the number of exact solutions found for these 50 random starting points. With an exact solution, a solution for which the optimality conditions, i.e., the tolerance on the gradient and constraint, are met, and furthermore for which the cost function is smaller than  $10^{-12}$ . The last column in the table indicates the number of practical solutions that were obtained when starting from the exact numerical PDs and by adding the constraint  $h_{\text{discr}}$  from (13), scaled with a factor 0.1, to the optimization problem. As can be seen, for many combinations, exact and practical PDs exist. An example practical PD<sub>11</sub> ( $T_{223}$ ) with (s, t) := (2, 2) is given by the following equations:

Also the size and minimal, maximal, and number of different ranks of  $J_{CS,gen}$ ,

i.e., the Jacobian matrix of the new generalized CS structure, are given at the solutions. For a list of all the different ranks, we refer to the detailed results of the experiment, which are publicly available <sup>2</sup>. Remark that the second dimension of the Jacobian matrix and the ranks decrease when  $r_{\rm CS}$  increases.

Table 1: Number of exact (numerical and practical)  $PD_{11}(T_{223})$ s that we obtained for different combinations of s and t using the AL method from [18] on 50 random starting points generated as in (21). Also the size and the different ranks of the Jacobian matrix are given.

$r_{ m CS}$	s	t	$\mathrm{size}(J_{\mathrm{CS,gen}})$	<b>rank</b> min	$(J_{CS,g})$ max	$_{ m gen}(x^*)) \ \#$	$\# x^*$	pract.
10	7	1	$144 \times 96$	82	82	1	10	0
9	6	1	$144 \times 104$	88	89	2	32	0
8	2	2	$144 \times 119$	92	97	4	8	1
0	5	1	144 × 112	93	98	5	45	0
	1	2		85	98	7	21	10
7	4	1	$144 \times 120$	87	103	10	46	16
	7	0		101	105	2	15	0
	0	2		91	108	14	29	13
6	3	1	$144 \times 128$	93	107	13	48	36
	6	0		107	111	5	29	0
5	2	1	$144 \times 136$	99	117	13	38	25
0	5	0	$144 \times 150$	112	117	6	44	0
4	1	1	$144 \times 144$	105	123	18	43	30
4	4	0	144 × 144	107	123	16	48	18
2	0	1	$144 \times 159$	111	129	16	39	18
5	3	0	144 × 152	113	129	15	49	29
2	2	0	$144 \times 160$	119	133	14	50	21
1	1	0	$144 \times 168$	123	135	11	49	14
0	0	0	$144 \times 176$	125	135	9	50	15

As can be seen from Table 1, the highest value for  $r_{\rm CS}$  that we were able to obtain is 10 and is obtained for (s,t) := (7,1). We were not able to transform this PD into a practical one using the constraint  $h_{\rm discr}$ .

<sup>&</sup>lt;sup>2</sup>https://github.com/CharlotteVermeylen/ALM\_FMM\_gen\_CS

In the last row, as a reference the results for (s,t) := (0,0), i.e., without the CS structure, are given. Remark that the number of numerical solutions in general decreases when  $r_{\rm CS}$  increases but, on the other hand, the number of practical PDs increases for some combinations of s and t.

In Figure 2, the convergence of the cost function is shown for the experiment with (s,t) := (7,0). As can be seen, 15 out of the 50 starting points converge quickly to a numerically exact PD in less than 100 iterations and the other starting points get stuck in swaps.



Figure 2: Convergence of the cost function in (10) using the AL method from [18] with u := -l := 1 and for 50 random starting points generated with randn of size  $10^{-2}$  to find PD<sub>11</sub> ( $T_{223}$ )s, with (s,t) := (7,0).

On the other hand, the results for  $PD_{14}(T_{224})$ s are shown in Table 2, again for different values of  $r_{CS}$ , s and t. The largest CS rank that we were able to obtain is again  $r_{CS} = 10$  but now both PDs with (s,t) = (4,2)and (s,t) = (7,1) were found. However, again none of these PDs were discretizable using the constraint  $h_{discr}$ .

An example practical  $PD_{14}(T_{224})$  that we obtained for (s,t) = (3,1) has

as submatrices:

Again it can be noticed that by including the generalized CS structure, more practical PDs are obtained and furthermore, more different values of the rank of the Jacobian and thus more non-inv-equivalent PDs are obtained as long as the CS rank is not set too high.

Table 2: Number of exact (numerical and practical)  $PD_{14}(T_{224})$ s that we obtained for different combinations of s and t using the AL method from [18] on 50 random starting points generated as in (21). Also the size and the minimal, maximal, and number of different ranks of the Jacobian matrix are given.

<i>m</i>		+	$\operatorname{size}(\mathbf{I})$	rank	$(J_{ m CS,g})$	# m*	pract	
'CS	8	ι	$SIZe(J_{CS,gen})$	min	max	#	# J	pract.
10	4	2	$256 \times 200$	170	170	1	1	0
10	7	1	230 × 200	172	178	2	6	0
0	3	2	$256 \times 208$	183	183	1	2	0
9	6	1	250 × 208	178	185	5	15	0
	2	2		187	190	3	6	0
8	5	1	$256 \times 216$	185	190	5	38	0
	8	0		184	191	2	3	0
	1	2		178	193	6	22	9
7	4	1	256 × 224	180	197	10	44	14
1	7	0	230 × 224	190	199	5	20	0
	0	2		184	198	9	20	9
6	3	1	$256 \times 232$	184	201	10	41	21
	6	0		196	205	7	20	0
5	2	1	$256 \times 240$	188	212	15	36	16
0	5	0	$230 \times 240$	205	210	6	36	0
1	1	1	256 × 248	196	214	13	30	12
4	4	0	230 × 240	196	217	15	44	12
9	0	1	$256 \times 256$	202	221	11	25	4
5	3	0	$230 \times 230$	202	221	15	42	11
2	2	0	$256 \times 264$	214	227	10	49	4
1	1	0	$256 \times 272$	217	228	9	45	4
0	0	0	$256 \times 280$	217	228	8	50	5

Thirdly, in Table 3, the results for  $T_{233}$ , rank 15 are shown. Now, the maximal value for  $r_{\rm CS}$  that we obtained is 9, and one practical PD is obtained

with parameters (s,t) = (6,1):

Table 3: Number of exact (numerical and practical)  $PD_{15}(T_{233})$ s that we obtained for different combinations of s and t using the AL method from [18] on 50 random starting points generated as in (21). Also the size and the different ranks of the Jacobian matrix are given.

$r_{ m CS}$	s	t	$\mathrm{size}(J_{\mathrm{CS,gen}})$	rank min	$(J_{CS,go})$ max	$_{\mathrm{en}}(x^{*})) \ \#$	$\# x^*$	pract.
0	3	2	294 2 942	216	216	1	2	0
9	6	1	324 × 243	218	218	1	1	1
8	2	2	$394 \times 951$	223	227	2	2	0
0	5	1	$324 \times 201$	225	226	2	4	2
7	1	2	$324 \times 250$	227	228	2	13	2
1	4	1	$324 \times 209$	229	234	5	15	5
	0	2		232	234	3	10	1
6	3	1	$324 \times 267$	234	239	6	25	10
	6	0		237	238	2	2	1
5	2	1	$394 \times 975$	240	244	4	12	7
0	5	0	324 × 273	244	245	2	2	0
4	1	1	204 ~ 282	247	250	3	5	2
4	4	0	324 × 203	249	254	5	11	6
2	0	1	$294 \times 201$	252	256	3	6	2
5	3	0	$324 \times 291$	254	259	6	18	6
2	2	0	$324 \times 299$	254	262	6	20	4
1	1	0	$324 \times 307$	260	264	5	30	0
0	0	0	$324 \times 315$	262	264	3	29	0

Fourthly, in Table 4, the results for  $T_{225}$ , rank 18, are shown. The maximal CS rank is 10 as in Table 1 and Table 2. However, the highest value of the CS rank for which a practical PD is obtained is 9 and is obtained for (s,t) = (6,1). The CS submatrices of the PD that we obtained are:

$$A = \begin{bmatrix} 1 & -1 & 1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 & 0 \\ 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & -1 \end{bmatrix}, B = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, C = \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix}, D = \begin{bmatrix} 0 \\ 0 \\ -1 \\ 0 \end{bmatrix},$$

	ΓO	0	0	0	0	0	0	0	0]					
	0	0	0	-1	0	1	0	0	0					
$\tilde{V}$	0	0	0	0	0	0	0	0	0	ũ		п	ŵ	п
$V \equiv$	0	0	0	-1	0	1	0	0	0	, 0	=	[],	V :	= [],
	0	0	0	0	0	0	0	0	0					
	0	0	0	0	-1	1	0	0	0					
	ΓO	0	0	0	0	0	0	0	0					
	0	0	0	0	-1	1	0	0	0					
TÎ 7	0	0	0	-1	0	1	0	0	0					
W =	0	0	0	0	0	0	0	0	0	,				
	0	0	0	0	1	-1	0	0	0					
	0	0	0	1	0	-1	0	0	0					
	ΓO	1	0	1	0	0 -	-1 -	-1	1	]				
÷.	0	0	0	0	1	1	0	0	1					
U =	1	0	1	-1	0	0	1	0	-1	,				
	1	0	1	-1	0	0	0	0	-1					
	ΓO	0	0	0	0	0	0		0	0]				
		0	0	0	0	0	0		0	0				
	0	0	0	0	0	0	0		0	0				
	0	0	0	0	0	0	0		0	0				
Ţ,	0	1	0	0	-1	0	0		0	0				
V =	0	1	0	-1	0	0	-1		0	0,				
	0	0	1	1	-1	0	0		0	1				
	0	0	1	0	0	0	0		0	0				
	0	0	1	1	0	-1	0	_	1	1				
	$\lfloor -1 \rfloor$	0	1	0	0	0	0		0	0				
	Γ 0	0	0	0	0	0	0		0	0]				
	0	0	0	0	0	0	0		0	0				
	0	1	0	0	0	0	-1		0	0				
	1	0	1	1	0	0	1	_	1	0				
Ŵ.	-1	0	0	0	0	0	0		1	0				
<i>vv</i> —	0	0	0	0	0	0	0		0	0	•			
	0	0	0	0	0	0	0		0	0				
	0	0	0	1	-1	-1	1		0 -	-1				
	0	0	0	-1	0	1	-1		0	1				
	L 0	0	0	0	0	-1	0		0	0				

$r_{\rm CS}$	s	t	$size(J_{CS,ron})$	rank	$(J_{ m CS,g})$	$_{ m gen}(x^*))$	$\# x^*$	pract.
- 05			(° C5,gen)	min	max	#	11	<b>F</b>
10	4	2	400 × 259	302	315	9	13	0
10	7	1	$400 \times 552$	298	310	10	34	0
0	3	2	400 × 260	309	319	9	17	0
9	6	1	$400 \times 300$	288	316	11	44	1
	2	2		308	326	11	19	0
8	5	1	$400 \times 368$	313	323	9	49	0
	8	0		308	325	12	23	0
	1	2		292	331	18	33	4
7	4	1	$400 \times 376$	293	327	15	46	3
	7	0		314	332	17	39	0
	0	2		297	339	14	23	3
6	3	1	$400 \times 384$	299	338	23	46	10
	6	0		319	339	19	41	0
5	2	1	400 × 302	305	341	19	47	8
5	5	0	400 × 392	329	341	13	49	0
4	1	1	400 × 400	313	352	23	49	4
4	4	0	400 × 400	312	351	21	48	5
3	0	1	$400 \times 408$	327	357	20	46	1
5	3	0	400 × 408	322	354	21	48	5
2	2	0	$400 \times 416$	330	354	17	49	2
1	1	0	$400 \times 424$	332	356	16	49	2
0	0	0	$400 \times 432$	336	358	15	50	1

Table 4: Number of exact (numerical and practical)  $PD_{18}(T_{225})$ s for different combinations of *s* and *t* using the AL method from [18] on 50 random starting points generated as in (21). Also the size and the different ranks of the Jacobian matrix are given.

Then in Table 5, the results for  $T_{234}$ , rank 20, are shown. The parameters s and t for which solutions are found are similar to the previous experiments but the number of solutions is on average smaller because the problem becomes more difficult as p, n, and the rank r increase. Also, fewer different values of the rank of the Jacobian matrix are found. Remark that the number of different values of the rank in general increases for moderate values of  $r_{\rm CS}$ . Furthermore, including the CS structure again helps to find practical solutions compared to the generic case ((s,t) = (0,0)) for different combinations of s and t.

Table 5: Number of exact (numerical and practical)  $PD_{20}(T_{234})s$  that we obtained for different combinations of (s,t) using the AL method from [18] on 50 random starting points generated as in (21). Also the size and the different ranks of the Jacobian matrix are given.

naa	6	+	$sizo(I_{aa})$	rank	$(J_{ m CS,g})$	$_{ m en}(x^*))$	$\# x^*$	pract	
		ι	Size( <i>J</i> CS,gen)	min	max	#	# x	pract.	
10	7	1	$576 \times 440$	401	401	1	1	0	
9	6	1	$576 \times 448$	403	408	2	2	0	
8	2	2	$576 \times 456$	409	409	1	2	0	
0	5	1	570 × 450	408	414	5	4	1	
7	1	2	576 × 464	412	415	3	5	0	
1	4	1	570 × 404	410	419	6	5	1	
6	0	2	$576 \times 479$	420	420	1	1	0	
0	3	1	510 × 412	421	424	4	10	0	
5	2	1	$576 \times 480$	425	432	6	7	1	
0	5	0	570 × 400	424	434	3	4	0	
1	1	1	$576 \times 488$	433	438	3	6	0	
4	4	0	570 × 400	427	438	3	4	1	
3	0	1	$576 \times 406$	441	443	2	4	0	
0	3	0	570 × 450	438	447	7	17	0	
2	2	0	$576 \times 504$	441	447	6	15	0	
1	1	0	$576 \times 512$	442	448	7	18	0	
0	0	0	$576 \times 520$	444	448	4	21	0	

Lastly in Table 6, the results for  $T_{245}$  and rank 33 are shown. Again the problem has become more difficult because p, n, and r have increased. Still, we find more numerical solutions for several CS parameters compared to the generic case ((s,t) = (0,0)).

Table 6: Number of exact (numerical and practical)  $PD_{33}(T_{245})$ s that we obtained for different combinations of s and t using the AL method from [18] on 50 random starting points generated as in (21). Also the size and the different ranks of the Jacobian matrix are given.

<b>m</b>	6	+	$size(I_{ab})$	rank	$(J_{ m CS,ge})$	$(x^*))$	# m*	pract	
'CS	3	ι	SIZe(JCS,gen)	min	max	#	# J	pract.	
10	4	2	$1600 \times 1174$	1091	1091	1	1	0	
	3	2		1088	1096	2	2	0	
9	6	1	$1600 \times 1182$	1089	1099	3	5	0	
	9	0		1096	1096	1	1	0	
0	5	1	$1600 \times 1100$	1096	1105	3	3	0	
0	8	0	$1000 \times 1190$	1106	1106	1	1	0	
	1	2		1094	1098	2	2	0	
7	4	1	$1600 \times 1198$	1102	1104	2	3	0	
	7	0		1105	1105	1	1	0	
6	3	1	$1600 \times 1206$	1102	1116	5	5	0	
0	6	0	$1000 \times 1200$	1115	1115	1	2	0	
5	2	1	$1600 \times 1914$	1118	1122	3	3	0	
0	5	0	$1000 \times 1214$	1118	1123	4	4	0	
1	1	1	1600 × 1999	1115	1120	4	4	0	
4	4	0	1000 × 1222	1118	1121	3	3	0	
2	0	1	1600 × 1920	1123	1127	3	5	0	
5	3	0	$1000 \times 1230$	1119	1131	5	7	0	
2	2	0	$1600 \times 1238$	1123	1128	4	7	0	
1	1	0	$1600 \times 1246$	1125	1134	3	6	0	
0	0	0	$1600 \times 1254$	1127	1130	3	4	0	

Remark that we can also choose the CS rank smaller than r when m = p = n. For example, we are able to enforce 11 symmetric rank-1 tensors in a PD<sub>23</sub> ( $T_{333}$ ). The other 12 rank-1 tensors do not have to admit a structure in this case. One of the practical PDs that we found for these parameters has

as symmetric part:

and as asymmetric part:

	$\left\lceil -1 \right\rceil$	0	0	0	0	-1	0	0	0	0	0	0	
	1	0	0	1	0	0	-1	0	-1	0	0	0	
$\dot{U} =$	0	0	0	1	0	-1	0	0	-1	-1	0	0	
	-1	0	0	0	0	-1	0	0	0	0	-1	1	
	1	0	0	1	0	0	0	0	0	0	0	0	,
	0	0	0	1	0	-1	0	1	0	0	-1	0	
	1	0	1	0	1	1	0	0	0	0	0	0	
	-1	1	-1	-1	0	0	0	0	0	0	0	0	
	0	0	0	-1	0	1	0	0	0	0	0	0	
	<b>[</b> 0	0	0	0	0	0	-1	0	0	1	0	0	
	$\begin{bmatrix} 0\\ 0 \end{bmatrix}$	0 0	$\begin{array}{c} 0 \\ 0 \end{array}$	0 0	0 0	$\begin{array}{c} 0 \\ 0 \end{array}$	$^{-1}_{1}$	$\begin{array}{c} 0 \\ 1 \end{array}$	0 0	$\begin{array}{c} 1 \\ 0 \end{array}$	$0 \\ -1$	$\begin{bmatrix} 0\\1 \end{bmatrix}$	
	$\begin{bmatrix} 0\\0\\0 \end{bmatrix}$	0 0 0	0 0 0	0 0 0	0 0 1	0 0 0	-1 1 0	0 1 1	0 0 0	$egin{array}{c} 1 \\ 0 \\ 1 \end{array}$	$0 \\ -1 \\ -1$	$\begin{bmatrix} 0\\1\\0 \end{bmatrix}$	
	$\begin{bmatrix} 0\\0\\0\\1 \end{bmatrix}$	0 0 0 0	$\begin{array}{c} 0 \\ 0 \\ 0 \\ -1 \end{array}$	0 0 0 0	0 0 1 0	0 0 0 0	$-1 \\ 1 \\ 0 \\ -1$	0 1 1 0	0 0 0 0	1 0 1 1	$0 \\ -1 \\ -1 \\ 0$	$\begin{bmatrix} 0\\1\\0\\0\end{bmatrix}$	
$\dot{V} =$	$\begin{bmatrix} 0\\0\\0\\1\\0 \end{bmatrix}$	0 0 0 0	$\begin{array}{c} 0 \\ 0 \\ 0 \\ -1 \\ 0 \end{array}$	0 0 0 0	0 0 1 0 0	0 0 0 0 0	$-1 \\ 1 \\ 0 \\ -1 \\ 1$	0 1 1 0 1	0 0 0 0 0	$     \begin{array}{c}       1 \\       0 \\       1 \\       1 \\       0     \end{array} $	$\begin{array}{c} 0 \\ -1 \\ -1 \\ 0 \\ 0 \end{array}$	$\begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$	,
$\dot{V} =$	$\begin{bmatrix} 0\\0\\1\\0\\0 \end{bmatrix}$	$egin{array}{c} 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{array}$	$\begin{array}{c} 0 \\ 0 \\ 0 \\ -1 \\ 0 \\ -1 \end{array}$	0 0 0 0 0	0 0 1 0 0 0	0 0 0 0 0 0	$-1 \\ 1 \\ 0 \\ -1 \\ 1 \\ 0$	$egin{array}{c} 0 \\ 1 \\ 1 \\ 0 \\ 1 \\ 1 \end{array}$	0 0 0 0 0 0	$     \begin{array}{c}       1 \\       0 \\       1 \\       1 \\       0 \\       1     \end{array} $	$\begin{array}{c} 0 \\ -1 \\ -1 \\ 0 \\ 0 \\ 0 \end{array}$	$ \begin{array}{c} 0\\1\\0\\0\\0\\0\\0\\0\end{array} \end{array} $	,
$\dot{V} =$	$ \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} $	0 0 0 0 1 0	$\begin{array}{c} 0 \\ 0 \\ 0 \\ -1 \\ 0 \\ -1 \\ 0 \end{array}$	0 0 0 0 0 0 0	0 0 1 0 0 0 0	$egin{array}{c} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ -1 \end{array}$	$-1 \\ 1 \\ 0 \\ -1 \\ 1 \\ 0 \\ 1$	0 1 1 0 1 1 0	0 0 0 0 0 0 1	$ \begin{array}{c} 1 \\ 0 \\ 1 \\ 1 \\ 0 \\ 1 \\ -1 \end{array} $	$\begin{array}{c} 0 \\ -1 \\ -1 \\ 0 \\ 0 \\ 0 \\ 0 \end{array}$	$ \begin{array}{c} 0\\1\\0\\0\\0\\0\\0\\0\\0\end{array} \end{array} $	,
$\dot{V} =$	$ \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} $	0 0 0 0 1 0 0	$egin{array}{c} 0 \\ 0 \\ -1 \\ 0 \\ -1 \\ 0 \\ 0 \end{array}$	$egin{array}{ccc} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ -1 \end{array}$	0 0 1 0 0 0 0 0	$egin{array}{c} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ -1 \\ 0 \end{array}$	$     \begin{array}{c}       -1 \\       1 \\       0 \\       -1 \\       1 \\       0 \\       1 \\       -1     \end{array} $	$egin{array}{c} 0 \\ 1 \\ 0 \\ 1 \\ 1 \\ 0 \\ -1 \end{array}$	$egin{array}{c} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ -1 \end{array}$	$egin{array}{c} 1 \\ 0 \\ 1 \\ 1 \\ 0 \\ 1 \\ -1 \\ 0 \end{array}$	$\begin{array}{c} 0 \\ -1 \\ -1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{array}$	$ \begin{array}{c} 0\\1\\0\\0\\0\\0\\0\\0\\0\\0\\0\end{array} \end{array} $	,

	0	0	0	0	1	0	0	0	0	0	0	1	1
	-1	1	-1	0	0	0	0	0	0	0	0	-1	
	0	1	-1	0	1	1	0	0	0	0	0	0	
	0	0	0	0	1	0	1	0	-1	0	0	1	
$\dot{W} =$	0	1	0	0	0	0	0	0	0	0	0	-1	.
	0	1	0	-1	1	0	0	0	-1	0	0	0	
	0	0	0	0	-1	0	0	0	0	-1	1	-1	
	0	-1	0	0	0	0	0	1	0	0	-1	1	
	0	-1	0	0	-1	0	0	0	0	0	0	0	

## 5. Conclusion

The proposed generalized cyclic symmetric structure can be useful to include in the optimization problem for the *fast matrix multiplication (FMM)* problem to reduce the number of parameters and to improve the convergence to numerically exact and practical *polyadic decompositions (PDs)* of *matrix multiplication tensors (MMTs)*. Several new *practical* PDs, i.e., sparse PDs with elements that are either one, minus one, or zero, with this new structure are discovered for various problem parameters, i.e., sizes of the MMT.

#### 6. Acknowledgement

This work was funded by (1) the Flemish Government: this research received funding under the AI Research Program. Charlotte Vermeylen is affiliated to Leuven.AI - KU Leuven institute for AI, B-3000, Leuven, Belgium. (2) KU Leuven Internal Funds: iBOF/23/064, C14/22/096 and IDN/19/014. Marc Van Barel was supported by the Research Council KU Leuven, C1-project C14/17/073 and by the Fund for Scientific Research–Flanders (Belgium), EOS Project no 30468160 and project G0B0123N.

#### 7. Competing interests

Declarations of interest: none.

#### References

 P. Bürgisser, M. Clausen, M. A. Shokrollahi, Algebraic complexity theory, Vol. 315, Springer Science & Business Media, 2013.

- [2] J. M. Landsberg, Tensors: Geometry and Applications, Vol. 128, American Mathematical Soc., 2011.
- [3] V. Strassen, Gaussian elimination is not optimal, Numerische Mathematik 13 (1969) 354–356. doi:10.1007/BF02165411.
- [4] J. M. Landsberg, Geometry and Complexity Theory, Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, 2017. doi:10.1017/9781108183192.
- H. F. de Groote, On varieties of optimal algorithms for the computation of bilinear mappings II. Optimal algorithms for 2 × 2-matrix multiplication, Theoretical Computer Science 7 (1978) 127–148. doi:https: //doi.org/10.1016/0304-3975(78)90045-2.
- [6] V. B. Alekseyev, On the complexity of some algorithms of matrix multiplication, Journal of Algorithms 6 (1) (1985) 71-85. doi:10.1016/ 0196-6774(85)90019-7.
- [7] V. B. Alekseev, A. V. Smirnov, On the exact and approximate bilinear complexities of multiplication of 4×2 and 2×2 matrices, Vol. 282, 2013, pp. 123-139.
   URL https://api.semanticscholar.org/CorpusID:120039387
- [8] M. Bläser, On the complexity of the multiplication of matrices of small formats, Journal of Complexity 19 (1) (2003) 43-60. doi:10.1016/S0885-064X(02)00007-9.
- J. D. Laderman, A non commutative algorithm for multiplying 3 × 3 matrices using 23 multiplications, Bulletin of the American Mathematical Society 82 (1) (1976) 126–128. doi:10.1090/S0002-9904-1976-13988-2.
- [10] G. Ballard, C. Ikenmeyer, J. M. Landsberg, N. Ryder, The geometry of rank decompositions of matrix multiplication II: 3 × 3 matrices, Journal of Pure and Applied Algebra 223 (8) (2019) 3205–3224. doi:10.1016/ j.jpaa.2018.10.014.
- [11] A. V. Smirnov, The bilinear complexity and practical algorithms for matrix multiplication, Computational Mathematics and Mathematical Physics 53 (12) (2013) 1781–1795. doi:10.1134/S0965542513120129.

- [12] M. J. Heule, M. Kauers, M. Seidl, New ways to multiply 3×3-matrices, Journal of Symbolic Computation 104 (2021) 899–916. doi:10.1016/ j.jsc.2020.10.003.
- [13] A. Fawzi, M. Balog, A. Huang, T. Hubert, B. Romera-Paredes, M. Barekatain, A. Novikov, F. J. R. Ruiz, J. Schrittwieser, G. Swirszcz, D. Silver, D. Hassabis, P. Kohli, Discovering faster matrix multiplication algorithms with reinforcement learning, Nature 610 (7930) (2022) 47–53. doi:10.1038/s41586-022-05172-4.
- [14] P. Tichavský, A. H. Phan, A. Cichocki, Numerical CP decomposition of some difficult tensors, Journal of Computational and Applied Mathematics 317 (2017) 362–370. doi:10.1016/j.cam.2016.12.007.
- [15] L. Chiantini, C. Ikenmeyer, J. Landsberg, G. Ottaviani, The geometry of rank decompositions of matrix multiplication I: 2 × 2 matrices, Experimental Mathematics 28 (2019) 322–327. arXiv:1610.08364.
- [16] H. F. de Groote, On varieties of optimal algorithms for the computation of bilinear mappings I. The isotropy group of a bilinear mapping, Theoretical Computer Science 7 (1) (1978) 1-24. doi:10.1016/0304-3975(78)90038-5.
- [17] R. W. Johnson, A. M. McLoughlin, Noncommutative bilinear algorithms for 3 × 3 matrix multiplication, SIAM Journal on Computing 15 (2) (1986) 595–603. doi:10.1137/0215043.
- [18] C. Vermeylen, M. V. Barel, Stability improvements for fast matrix multiplication, Numerical Algorithms (2024). doi:10.1007/s11075-024-01806-y.
- [19] J. Nocedal, S. J. Wright, Numerical optimization, Springer, New York, 1999. doi:10.1007/978-0-387-40065-5.
- [20] G. O. Berger, P.-A. Absil, L. De Lathauwer, R. M. Jungers, M. Van Barel, Equivalent polyadic decompositions of matrix multiplication tensors, Journal of Computational and Applied Mathematics 406 (2022) 113941. doi:j.cam.2021.113941.
- [21] G. Ballard, Discovering fast matrix multiplication algorithms via tensor decomposition, in: SIAM Conference on Computational Science and

Engineering, Vol. 30, 2017. URL https://users.wfu.edu/ballard/pdfs/CSE17.pdf

- [22] J. Huang, L. Rice, D. A. Matthews, R. A. van de Geijn, Generating families of practical fast matrix multiplication algorithms, in: 2017 IEEE International Parallel and Distributed Processing Symposium (IPDPS), 2017, pp. 656–667. doi:10.1109/IPDPS.2017.56.
- [23] C. Vermeylen, Tensor methods: Fast matrix multiplication and rank adaptation, Ph.D. thesis (2024-03-29).
   URL https://lirias.kuleuven.be/4144594&lang=en