THE DIRICHLET PROBLEM WITH ENTIRE DATA FOR NON-HYPERBOLIC QUADRATIC HYPERSURFACES

J. M. ALDAZ AND H. RENDER

ABSTRACT. We show that for all homogeneous polynomials f_m of degree m, in d variables, and each $j = 1, \ldots, d$, we have

$$\left\langle x_{j}^{2}f_{m}, f_{m}\right\rangle_{L^{2}(\mathbb{S}^{d-1})} \geq \frac{\pi^{2}}{4\left(m+d+1\right)^{2}}\left\langle f_{m}, f_{m}\right\rangle_{L^{2}(\mathbb{S}^{d-1})}$$

This result is used to establish the existence of entire harmonic solutions of the Dirichlet problem, when the data are given by entire functions of order sufficiently low on nonhyperbolic quadratic hypersurfaces.

1. INTRODUCTION

The Dirichlet problem has been solved for very general domains of \mathbb{R}^d . S. J. Gardiner proved that solutions exist, and not just for continuous boundary data on unbounded domains, but even for locally bounded Borel data (cf. the characterization given in [8, Theorem 1] and the comments following it). A different question that naturally arises is whether harmonic solutions can be extended beyond the boundary when the data is sufficiently regular, perhaps yielding global solutions. Additional information on these matters can be found in reference [9].

One way of ensuring the existence of global solutions is to prove that if the data is given by a polynomial, or more generally, by an entire function, the solution is also entire. Following a line of research started in [16], D. Khavinson and H. S. Shapiro showed that for every entire function f on \mathbb{C}^d , the solution h of the Dirichlet problem for the ellipsoid with data function f (restricted to the boundary) has a harmonic continuation to \mathbb{R}^d , and hence it extends to an entire function on \mathbb{C}^d , see [12, Theorem 1]. They also asked whether this property characterized ellipsoids, cf. [12, Page 460], a problem that is still open in full generality, though under some additional assumptions a positive answer is known (see [13, Theorem 27]). An equivalent version of the Khavinson-Shapiro problem appears in [14].

²⁰²⁰ Mathematics Subject Classification: Primary: 35A20, Secondary: 35A01.

Key words and phrases: Dirichlet problem, entire harmonic function, nonhyperbolic quadratic surface.

The first named author was partially supported by Grant PID2019-106870GB-I00 of the MICINN of Spain, by ICMAT Severo Ochoa project CEX2019-000904-S (MICINN), and by the Madrid Government (Comunidad de Madrid - Spain) V PRICIT (Regional Programme of Research and Technological Innovation), 2022-2024.

The aim of the present paper is to show that results related to the Khavinson-Shapiro Theorem hold for Dirichlet problems on some specific, not necessarily bounded domains in \mathbb{R}^d , namely, on nonhyperbolic quadratic hypersurfaces. The Dirichlet problem for polynomial data on such hypersurfaces was studied in [5], where the authors present an algorithm to compute polynomial harmonic solutions. Next we recall [5, Definition 4].

Definition 1. Let $a_1, \ldots, a_d, b_1, \ldots, b_d$ and c, be real numbers. A nonhyperbolic quadratic polynomial q has the form

$$q(x) := \sum_{j=1}^{d} a_j^2 x_j^2 + \sum_{j=1}^{d} b_j x_j + c,$$

where at least one $a_i \neq 0$. A nonhyperbolic quadratic hypersurface is the zero set $\{q = 0\}$.

Writing $q = P_2 + P_1 + P_0$, where P_i denotes the homogeneous part of q of degree i, we set $\beta = 1$ if $P_1 \neq 0$ and $\beta = 0$ otherwise. Then we have

Theorem 2. Let $d \ge 2$, and let q be a nonhyperbolic quadratic polynomial. If f is an entire function of order $\rho < 1 - \beta/2$, then then there is a harmonic entire function of order bounded by ρ , which coincides with f on $\{q = 0\}$.

As special cases we mention paraboloids, where $q(x) = a_2^2 x_2^2 + \cdots + a_d^2 x_d^2 - x_1$ and $\rho < 1/2$, ellipsoidal cylinders, where $q(x) = a_2^2 x_2^2 + \cdots + a_d^2 x_d^2 - 1$ and $\rho < 1$, and slabs, where $q(x) = a_d^2 x_d^2 - 1$ and $\rho < 1$. The case of the ellipsoidal cylinders had already been treated in [10] by potential-theoretic methods, under the same hypothesis $\rho < 1$, but without reaching the conclusion that the harmonic solution has finite order bounded by ρ . For slabs, the existence of harmonic entire solutions is obtained in [11], under no assumptions on the order of the entire data functions. Finally, in the (bounded) case of ellipsoids, we have already mentioned the Khavinson-Shapiro Theorem for arbitrary entire functions. They also note that if the data function is of exponential type, so is the harmonic solution (cf. [12, P. 459 Remarks]). In [3, Theorem 1] D. Armitage shows that the order of the solution is bounded by the order of the data function. Thus, for ellipsoids Theorem 2 only yields strictly weaker results: since via a rotation any ellipsoid can be expressed as $q(x) = a_1^2 x_1^2 + \cdots + a_d^2 x_d^2 - 1 = 0$, we take $\beta = 0$ and thus, $\rho < 1$.

The results presented here are obtained by generalizing to arbitrary dimensions Theorems 3, 4 and 5 from [15], proven there only for the case d = 2. Theorem 2 is obtained via the following estimate, together with [15, Theorem 1]).

Theorem 3. For all homogeneous polynomials f_m of degree m in d variables, and each $j = 1, \ldots, d$, we have

$$\langle x_j^2 f_m, f_m \rangle_{L^2(\mathbb{S}^{d-1})} \ge \frac{\pi^2}{4(m+d+1)^2} \langle f_m, f_m \rangle_{L^2(\mathbb{S}^{d-1})}.$$

 $\mathbf{2}$

2. Jacobi polynomials

In the following we need the Jacobi polynomials $P_n^{(\alpha,\alpha)}(x)$ of degree $n \ge 0$ and parameter α , see [17]. Here $x \in \mathbb{R}$. For $n \ge 1$ the Jacobi polynomials $P_n^{(\alpha,\alpha)}(x)$ satisfy the recurrence relation

$$x P_n^{(\alpha,\alpha)}(x) = a_n(\alpha) P_{n+1}^{(\alpha,\alpha)} + g_n(\alpha) P_{n-1}^{(\alpha,\alpha)}(x)$$

with coefficients $a_n(\alpha)$ and $g_n(\alpha)$ defined by

$$a_n(\alpha) = \frac{(n+1)(n+2\alpha+1)}{(2n+2\alpha+1)(n+\alpha+1)},$$

$$g_n(\alpha) = \frac{n+\alpha}{2n+2\alpha+1},$$

and initial conditions $P_0^{(\alpha,\alpha)}(x) = 1$ and $P_1^{(\alpha,\alpha)}(x) = (\alpha + 1)x$. It is easy to see that

(1)
$$x^{2}P_{n}^{(\alpha,\alpha)}(x) = \widetilde{a}_{n}(\alpha)P_{n+2}^{(\alpha,\alpha)}(x) + \widetilde{b}_{n}(\alpha)P_{n}^{(\alpha,\alpha)}(x) + \widetilde{g}_{n}(\alpha)P_{n-2}^{(\alpha,\alpha)}(x),$$

where $\tilde{a}_n = a_n a_{n+1}$, $\tilde{g}_n = g_n g_{n-1}$, and $\tilde{b}_n = a_n g_{n+1} + g_n a_{n-1}$. Write $0 \le y := x^2$ and u := 2n. Note that since $P_{2n}^{(\alpha,\alpha)}(x)$ is even (cf. [17, (4.1.3)]) the function $r_u(y) := P_{2n}^{(\alpha,\alpha)}(\sqrt{y})$ is a polynomial on $[0,\infty)$, and furthermore,

(2)
$$y r_{u}(y) = \widetilde{a}_{u}(\alpha) r_{u+1}(y) + \widetilde{b}_{u}(\alpha) r_{u}(y) + \widetilde{g}_{u}(\alpha) r_{u-1}(y).$$

Next we need an estimate of the first positive root of $P_{2n}^{(\alpha,\alpha)}(x)$. Based on the recent work of \hat{A} . Elbert and D. Siafarikas (see [6]) we can derive quickly the following result:

Theorem 4. Let $x_{2n,1}(\alpha)$ be the first positive zero of $P_{2n}^{(\alpha,\alpha)}(x)$, where $n \geq 3$ and $\alpha \geq -1/2$. Then

$$x_{2n,1}\left(\alpha\right) \geq \frac{\pi}{4\sqrt{\left(\alpha + \frac{1}{2} + n\right)\left(n + 2\right)}}$$

Proof. It is well known that $P_{2n}^{(\alpha,\alpha)}(x)$ is equal to a multiple of the Gegenbauer (or ultraspherical) polynomial $C_{2n}^{\lambda}(x)$ of degree 2n with parameter $\lambda = \alpha + 1/2$. Denote the first positive zero of C_{2n}^{λ} by $y_{2n,1}(\lambda)$, so $x_{2n,1}(\alpha) = y_{2n,1}(\alpha + \frac{1}{2})$. In [6, p. 32], the authors show that

$$f_{2n}(\lambda) := \sqrt{\lambda + \frac{8n^2 + 1}{8n + 2}} y_{2n,1}(\lambda)$$

is an increasing function of λ on the open interval $\left(-\frac{1}{2},\infty\right)$. Hence

$$f_{2n}(\lambda) \ge f_{2n}(0) = \sqrt{\frac{8n^2 + 1}{8n + 2}} y_{2n,1}(0).$$

Now $y_{2n,1}(0)$ is the first positive zero of the Gegenbauer polynomial $C_{2n}^0(x)$. Recalling that $C_{2n}^0(x)$ is equal to the Chebyshev polynomial $T_{2n}(x)$ of the first kind, and that the zeros of T_{2n} are given by $\cos\left(\frac{2k-1}{2n}\frac{\pi}{2}\right)$ for $k = 1, \ldots, 2n$, we conclude that

$$y_{2n,1}(0) = \cos\left(\frac{2n-1}{2n}\frac{\pi}{2}\right) = \sin\left(\frac{\pi}{2} - \frac{2n-1}{2n}\frac{\pi}{2}\right) \\ = \sin\frac{\pi}{4n} \ge \frac{\pi}{4n+2},$$

where we have used that $\cos x = \sin \left(\frac{\pi}{2} - x\right)$, and for the last inequality, [15, Lemma 14]. If $\alpha \ge -\frac{1}{2}$ then $\lambda = \alpha + 1/2 \ge 0$, so

$$\sqrt{\alpha + \frac{1}{2} + \frac{8n^2 + 1}{8n + 2}} y_{2n} \left(\alpha + \frac{1}{2} \right) = f_{2n} \left(\lambda \right) \ge f_{2n} \left(0 \right).$$

It follows that

$$\left(\alpha + \frac{1}{2} + \frac{8n^2 + 1}{8n + 2}\right) (x_{2n,1}(\alpha))^2 \ge \frac{8n^2 + 1}{8n + 2} \frac{\pi^2}{(4n + 2)^2}.$$

Since $n \ge \frac{8n^2+1}{8n+2}$ and $\frac{8n^2+1}{8n+2}\frac{1}{(4n+2)^2} \ge \frac{1}{16(n+2)}$ we obtain the estimate

$$\left(\alpha + \frac{1}{2} + n\right) \left(x_{2n,1}\left(\alpha\right)\right)^2 \ge \frac{1}{16(n+2)}\pi^2.$$

3. Proof of Theorem 3

Denote the unit sphere by $\mathbb{S}^{d-1} = \{\theta \in \mathbb{R}^d : |\theta| = 1\}$, and its surface area measure by $d\theta$. The $L^2(\mathbb{S}^{d-1})$ inner product and its associated norm, are defined in the usual way:

(3)
$$\langle f,g\rangle_{L^2(\mathbb{S}^{d-1})} := \int_{\mathbb{S}^{d-1}} f(\theta) \overline{g(\theta)} d\theta \text{ and } \|f\|_{L^2(\mathbb{S}^{d-1})} = \sqrt{\langle f,f\rangle_{L^2(\mathbb{S}^{d-1})}}.$$

We shall need some results from the theory of spherical harmonics. It is well known that one can construct *orthogonal* spherical harmonics by induction over the dimension d. In order to emphasize the dependence on d, we shall denote an orthogonal basis of the space $\mathcal{H}_k(\mathbb{R}^d)$ of all homogeneous harmonic polynomials of degree k in d variables, by

$$Y_{k,l}^d(x_1, \dots, x_d)$$
 for $l = 1, \dots, a_k^{(d)}$,

where we have written $a_k^{(d)} := \dim \mathcal{H}_k(\mathbb{R}^d)$. We also use $x = (x_1, \ldots, x_d)$ and $|x| = \sqrt{x_1^2 + \cdots + x_d^2}$. The following result can be found, for instance, in [1, p. 460] or in [7].

Theorem 5. Let $d \ge 2$, let $k \in \mathbb{N}$, and for each $s = 0, 1, 2, \ldots, k$, let

$$\{Y_{s,l}^{d-1}: l = 1, \dots, a_s^{(d-1)}\}$$

be an orthogonal basis of $\mathcal{H}_s(\mathbb{R}^{d-1})$. Then the system of polynomials given by

(4)
$$Y_{k,(s,l)}^{d}(x) := Y_{s,l}^{d-1}(x_1, \dots, x_{d-1}) \cdot |x|^{k-s} P_{k-s}^{(s+(d-3)/2, s+(d-3)/2)}\left(\frac{x_d}{|x|}\right),$$

where s = 0, 1, 2, ..., k and $l = 1, ..., a_s^{(d-1)}$, forms an orthogonal basis of $\mathcal{H}_k(\mathbb{R}^d)$.

Using the abreviation $\alpha_s := s + (d-3)/2$, for $k \ge 1$ identity (4) becomes

$$Y_{k,(s,l)}^{d}(x) = Y_{s,l}^{d-1}(x_1, \dots, x_{d-1}) \cdot |x|^{k-s} P_{k-s}^{(\alpha_s, \alpha_s)}\left(\frac{x_d}{|x|}\right)$$

Note that the same factor $Y_{s,l}^{d-1}(x_1,\ldots,x_{d-1})$ appears in the preceding expression for different values of k, when |x| = 1. The recurrence relation (1) together with (4) imply that

$$x_{d}^{2}Y_{k,(s,l)}^{d}(x) = \widetilde{a}_{k-s}(\alpha_{s})Y_{k+2,(s,l)}^{d}(x) + \widetilde{b}_{k-s}(\alpha_{s})Y_{k,(s,l)}^{d}(x) + \widetilde{g}_{k-s}(\alpha_{s})Y_{k-2,(s,l)}^{d}(x).$$

Define the normalized spherical harmonic by

$$\widetilde{Y}_{k,(s,l)}^d = d_{k,(s,l)} \cdot Y_{k,(s,l)}^d$$
 where $d_{k,(s,l)} := \left\| Y_{k,(s,l)}^d \right\|_{\mathbb{S}^{d-1}}^{-1}$.

It follows that

(5)
$$x_{d}^{2} \widetilde{Y}_{k,(s,l)}^{d}(x) = \widetilde{a}_{k-s}(\alpha_{s}) \frac{d_{k,(s,l)}}{d_{k+2,(s,l)}} \widetilde{Y}_{k+2,(s,l)}^{d}(x) + \widetilde{b}_{k-s}(\alpha_{s}) \widetilde{Y}_{k,(s,l)}^{d}(x)$$

(6)
$$+ \widetilde{g}_{k-s}\left(\alpha_s\right) \frac{d_{k,(s,l)}}{d_{k-2,(s,l)}} \widetilde{Y}^d_{k-2,(s,l)}\left(x\right).$$

The following result, cf. [15, Lemma 12], shows that to prove Theorem 3 we only need to consider homogeneous polynomials f_{2m} of even degree.

Lemma 6. Let P_{2k} be a homogeneous polynomial of degree 2k > 0. Suppose that for each $m \in \mathbb{N}$, there exists a constant $C_{2m} > 0$ such that for all homogeneous polynomials f_{2m} of degree 2m, we have

$$\langle P_{2k}f_{2m}, f_{2m} \rangle_{L^2(\mathbb{S}^{d-1})} \ge C_{2m} \langle f_{2m}, f_{2m} \rangle_{L^2(\mathbb{S}^{d-1})}.$$

Then for all homogeneous polynomials f_{2m+1} of degree 2m + 1,

$$\langle P_{2k}f_{2m+1}, f_{2m+1} \rangle_{L^2(\mathbb{S}^{d-1})} \ge C_{2m+2} \langle f_{2m+1}, f_{2m+1} \rangle_{L^2(\mathbb{S}^{d-1})}$$

Let us recall Theorem 3.

Theorem 7. For all homogeneous polynomials f_m of degree m in d variables, and each $j = 1, \ldots, d$, we have

$$\langle x_j^2 f_m, f_m \rangle_{L^2(\mathbb{S}^{d-1})} \ge \frac{\pi^2}{4(m+d+1)^2} \langle f_m, f_m \rangle_{L^2(\mathbb{S}^{d-1})}$$

Proof. Without loss of generality we set j = d. By the preceding Lemma it suffices to show that for all even indices we can take

$$C_{2m} := \frac{\pi^2}{4 \left(2m + 2d\right)^2}.$$

By the Gauss decomposition (cf. Theorem 5.5 in [4] or Theorem 5.7 in the 2001 edition) there exist homogeneous harmonic polynomials h_{2k} of degree 2k for $k = 0, \ldots, m$, such that

(7)
$$f_{2m}(x) = \sum_{k=0}^{m} |x|^{2m-2k} h_{2k}(x).$$

Each harmonic polynomial h_{2k} of degree 2k for $k = 0, \ldots, m$ is a linear combination of the orthonormal spherical harmonics $\widetilde{Y}_{2k,(s,l)}^d$ with $s = 0, 1, 2, \ldots, 2k$ and $l = 1, \ldots, a_s^{(d-1)}$. Thus, we can write

$$h_{2k} = \sum_{s=0}^{2k} \sum_{l=1}^{a_s^{(d-1)}} c_{k,s,l}^d \widetilde{Y}_{2k,(s,l)}$$

for suitable complex coefficients $c_{k,s,l}^d$.

Recall that $\lfloor x \rfloor$ denotes the integer part of x. By reordering it is easy to see that

$$f_{2m}(x) = \sum_{k=0}^{m} |x|^{2m-2k} h_{2k}(x) = \sum_{s=0}^{2m} \sum_{l=1}^{a_s^{(d-1)}} \sum_{k=\lfloor \frac{s+1}{2} \rfloor}^{m} c_{k,s,l} |x|^{2m-2k} \widetilde{Y}_{2k,(s,l)}^d(x).$$

By orthogonality

(8)
$$\langle f_{2m}, f_{2m} \rangle_{L^2(\mathbb{S}^{d-1})} = \sum_{s=0}^{2m} \sum_{l=1}^{a_s^{(d-1)}} \sum_{k=\lfloor \frac{s+1}{2} \rfloor}^m |c_{k,s,l}|^2,$$

(since we are integrating over the unit sphere). Now $\langle x_d^2 f_{2m}, f_{2m} \rangle_{L^2(\mathbb{S}^{d-1})}$ is equal to

$$\sum_{s_1=0}^{2m} \sum_{l_1=1}^{a_s^{(d-1)}} \sum_{k_1=\left\lfloor\frac{s+1}{2}\right\rfloor}^m \sum_{s_2=0}^{2m} \sum_{l_2=1}^{a_s^{(d-1)}} \sum_{k_2=\left\lfloor\frac{s+1}{2}\right\rfloor}^m c_{k_1,s_1,l_1} \overline{c_{k_2,s_2,l_2}} \left\langle x_d^2 \widetilde{Y}_{2k_1,(s_1,l_1)}, \widetilde{Y}_{2k_2,(s_2,l_2)} \right\rangle_{L^2(\mathbb{S}^{d-1})},$$

so by orthogonality (using the relation (5-6) for $x_d^2 \widetilde{Y}_{2k,(s,l)}$) we obtain

(9)
$$\left\langle x_{d}^{2}f_{m}, f_{m}\right\rangle_{\mathbb{S}^{d-1}} = \sum_{s=0}^{2m} \sum_{l=1}^{a_{s}^{(d-1)}} \sum_{k_{1}=\left\lfloor \frac{s+1}{2} \right\rfloor}^{m} \sum_{k_{2}=\left\lfloor \frac{s+1}{2} \right\rfloor}^{m} c_{k_{1},s,l} \overline{c_{k_{2},s,l}} A_{k_{1},k_{2}}\left(s,l\right)$$

where

$$A_{k_{1},k_{2}}(s,l) := \left\langle x_{d}^{2} \widetilde{Y}_{2k_{1},(s,l)}, \widetilde{Y}_{2k_{2},(s,l)} \right\rangle_{L^{2}(\mathbb{S}^{d-1})}.$$

In view of (8) and (9), it suffices to show that there exists a D > 0 (independent of m) such that for each s = 0, ..., 2m and each $l = 1, ..., a_s^{(d-1)}$,

(10)
$$\sum_{k_1 = \lfloor \frac{s+1}{2} \rfloor}^{m} \sum_{k_2 = \lfloor \frac{s+1}{2} \rfloor}^{m} c_{k_1, s, l} \overline{c_{k_2, s, l}} A_{k_1, k_2}(s, l) \ge \frac{\pi^2}{4 \left(2m + D\right)^2} \sum_{k = \lfloor \frac{s+1}{2} \rfloor}^{m} |c_{k, s, l}|^2$$

The integer s is either even or odd, so we can write $s = 2\sigma + \delta$ with $\delta \in \{0, 1\}$. Then

$$\left\lfloor \frac{s+1}{2} \right\rfloor = \left\lfloor \frac{2\sigma + 1 + \delta}{2} \right\rfloor = \sigma + \delta.$$

Let us define the symmetric $(m - \sigma - \delta + 1) \times (m - \sigma - \delta + 1)$ -matrix

$$A_{m}(s,l) := (A_{k_{1},k_{2}}(s,l))_{k_{1},k_{2}=\sigma+\delta,\dots,m}$$

Writing $c^t = \left(c_{\lfloor \frac{s+1}{2} \rfloor, s, l}, \dots, c_{m, s, l}\right)$, it is clear that the left hand side of (10) equals $c^t A_m(s, l) c$,

so it is enough to show that the smallest eigenvalue $\lambda_{m,s,l}^{*}$ of $A_{m}(s,l)$ satisfies the inequality

$$\lambda_{m,s,l}^* \ge \pi^2/4 \left(2m + 2d\right)^2.$$

Since $A_m(2\sigma + 1, l)$ is the submatrix of $A_m(2\sigma, l)$ obtained by deleting the first row and the first column, its lowest eigenvalue is at least as large as $\lambda_{m,2\sigma,l}^*$. Thus, it suffices to prove (10) for $A_m(2\sigma, l)$, or equivalently, for $\delta = 0$.

We now use the recurrence relation (5 - 6) (with 2k instead of k)

$$x_{d}^{2} \widetilde{Y}_{2k,(s,l)}^{d}(x) = \widetilde{a}_{2k-s}(\alpha_{s}) \frac{d_{2k,(s,l)}}{d_{2k+2,(s,l)}} \widetilde{Y}_{2k+2,(s,l)}^{d}(x) + \widetilde{b}_{2k-s}(\alpha_{s}) \widetilde{Y}_{2k,(s,l)}^{d}(x) + \widetilde{g}_{2k-s}(\alpha_{s}) \frac{d_{2k,(s,l)}}{d_{2k-2,(s,l)}} \widetilde{Y}_{2k-2,(s,l)}^{d}(x) .$$

From the orthonormality relations we see that

$$A_{k,k}\left(s,l\right) = \left\langle x_{d}^{2}\widetilde{Y}_{2k,(s,l)}, \widetilde{Y}_{2k,(s,l)} \right\rangle_{L^{2}(\mathbb{S}^{d-1})} = \widetilde{b}_{2k-s}\left(\alpha_{s}\right),$$

$$A_{k,k+1}(s,l) = \left\langle x_d^2 \widetilde{Y}_{2k,(s,l)}, \widetilde{Y}_{2k+2,(s,l)} \right\rangle_{L^2(\mathbb{S}^{d-1})} = \widetilde{a}_{2k-s}(\alpha_s) \frac{d_{2k,(s,l)}}{d_{2k+2,(s,l)}},$$

and

$$A_{k,k-1}(s,l) = \left\langle x_d^2 \widetilde{Y}_{2k,(s,l)}, \widetilde{Y}_{2k-2,(s,l)} \right\rangle_{L^2(\mathbb{S}^{d-1})} = \widetilde{g}_{2k-s}(\alpha_s) \frac{d_{2k,(s,l)}}{d_{2k-2,(s,l)}}$$

for $k = \sigma, \ldots, m$ and $s = 2\sigma$, with all the other entries being equal to zero. It follows that $A_m(s, l)$ is an $(m - \sigma + 1) \times (m - \sigma + 1)$ -matrix of the form

$$\begin{pmatrix} \widetilde{b}_{0} & \widetilde{a}_{0} \frac{d_{2\sigma,(s,l)}}{d_{2\sigma+2,(s,l)}} \\ \widetilde{g}_{2} \frac{d_{2\sigma+2,(s,l)}}{d_{2\sigma,(s,l)}} & \widetilde{b}_{2} & \ddots \\ & \ddots & \ddots & \widetilde{a}_{2m-2-2\sigma} \frac{d_{2m-2,(s,l)}}{d_{2m,(s,l)}} \\ & & \widetilde{g}_{2m-2\sigma} \frac{d_{2m,(s,l)}}{d_{2m-2,(s,l)}} & \widetilde{b}_{2m-2\sigma} \end{pmatrix}$$

We recall some known facts about tridiagonal $n \times n$ -matrices of the form

$$J_n = \begin{pmatrix} \beta_0 & \alpha_0 & & \\ \gamma_1 & \beta_1 & \ddots & \\ & \ddots & \ddots & \alpha_{n-2} \\ & & \gamma_{n-1} & \beta_{n-1} \end{pmatrix}$$

where all the entries α_k and γ_k are non-zero. In order to compute the characteristic polynomial det $(J_n - \lambda I_n)$ of J_n , we define a sequence of polynomials $p_0(x) := 1$, $p_1(x) := (x - \beta_0) / \alpha_0$, and for 0 < k < n, we use the recurrence relation

(11)
$$xp_{k}(x) = \gamma_{k}p_{k-1}(x) + \beta_{k}p_{k}(x) + \alpha_{k}p_{k+1}(x).$$

Then an inductive argument for $n \ge 1$, together with the expansion of the determinant along the last column, yields that $p_n(x)$ is a constant multiple of the characteristic polynomial of J_n , and more specifically, that

$$\det \left(J_n - \lambda I_n\right) = \left(-1\right)^n \alpha_0 \dots \alpha_{n-1} p_n\left(\lambda\right).$$

Thus the eigenvalues of J_n are the zeros of the polynomials $p_n(x)$ defined by (11). Next we consider the special form $A_m(s, l)$ of J_n that appears in our argument, with an additional simplification. Replacing p_k with $q_k := d_k p_k$ (where $d_k \neq 0$) in (11) we obtain the recurrence relation

$$xq_{k}(x) = \alpha_{k} \frac{d_{k}}{d_{k+1}} q_{k+1}(x) + \beta_{k}q_{k}(x) + \gamma_{k} \frac{d_{k}}{d_{k-1}} q_{k-1}(x),$$

with the initial conditions transformed to $q_0(x) = d_0$ and $q_1(x) = d_1(x - \beta_0) / \alpha_0$.

It follows the matrix $A_m(s, l)$, for $s = 2\sigma$, has the same eigenvalues as the matrix

$$\begin{pmatrix} \widetilde{b}_0 & \widetilde{a}_0 & & \\ \widetilde{g}_2 & \widetilde{b}_2 & \ddots & \\ & \ddots & \ddots & \widetilde{a}_{2m-2-2\sigma} \\ & & \widetilde{g}_{2m-2\sigma} & \widetilde{b}_{2m-2\sigma} \end{pmatrix}.$$

Next we replace the even indices 2n appearing in the preceding matrix with u = 2n, and we recognize it as the Jacobi matrix whose eigenvalues are computed via the polynomials appearing in the recurrence relation (2). Denote by y_u the first positive zero of $r_u(y) = P_{2n}^{(\alpha,\alpha)}(\sqrt{y})$, with $n = m - \sigma + 1$ and $\alpha = s + (d-3)/2$. Then the smallest eigenvalue $\lambda_{m,s,l}^{(\alpha,\alpha)}$ of $A_m(s,l)$ is y_u . Note that σ ranges from 0 to m, so $n = m - \sigma + 1$ ranges from 1 to m + 1. Furthermore,

$$\alpha = s + (d-3)/2 = 2(m+1-n) + (d-3)/2$$

By Theorem 4,

$$\left(\alpha + \frac{1}{2} + n\right)(n+2)y_u \ge \frac{\pi^2}{16}.$$

For $n = m - \sigma + 1$ we estimate

$$\left(\alpha + \frac{1}{2} + n \right) (n+2) \leq \max_{n=1,\dots,m+1} \left(2 \left(m + 1 - n \right) + \left(d - 3 \right) / 2 + 1 / 2 + n \right) \cdot (n+2)$$

$$\leq (m+d)^2 \,,$$

using the fact that $d \ge 2$, which in particular entails that for every $\sigma \ge 0$ we have $3\sigma - \sigma^2 - (d\sigma)/2 \le 1$.

It follows that

$$\lambda_{m,s,l}^* = y_u \ge \frac{\pi^2}{16 (m+d)^2}.$$

4. Proof of Theorem 2

The techniques of proof used here are based on the Fischer decomposition of polynomials, cf. [15], a framework that allows a unified treatment of all nonhyperbolic quadratic hypersurfaces. We use the special case of [15, Theorem 1] where $\alpha = 2$ and k = 1.

Theorem 8. Let P_2 be a homogeneous polynomial of degree 2 such that there exist C > 0and D > 0 with

(12)
$$\langle P_2 f_m, f_m \rangle_{L^2(\mathbb{S}^{d-1})} \ge \frac{1}{C \left(m+D\right)^2} \langle f_m, f_m \rangle_{L^2(\mathbb{S}^{d-1})}$$

for all homogeneous polynomials f_m of degree m. Let $q = P_2 + P_1 + P_0$, where P_i denotes the homogeneous part of q of degree i. Set $\beta = 1$ if $P_1 \neq 0$ and $\beta = 0$ otherwise. Then for every entire function f of order $\rho < (2 - \beta)/2$, there exist entire functions s and r of order $\leq \rho$ such that

$$f = q \cdot s + r$$
 and $\Delta r = 0$.

Now Theorem 3 allows us to apply the preceding result to any nonhyperbolic polynomial q, so Theorem 2 immediately follows, since clearly r is harmonic and satisfies r = f on $\{q = 0\}$.

References

- [1] G. E. Andrews, R. Askey, R. Roj, Special Functions, Cambridge University Press 1999.
- [2] D. H. Armitage, S. J. Gardiner, *Classical Potential Theory*, Springer, London 2001.
- [3] D. Armitage, The Dirichlet problem when the boundary function is entire, J. Math. Anal. 291 (2004), 565–577.
- [4] S. Axler, P. Bourdon, W. Ramey, Harmonic Function Theory, Springer, New York 1992.
- [5] S. Axler, P. Gorkin, K. Voss, The Dirichlet problem on quadratic surfaces, Math. Comp. 73 (2004), 637–651.
- [6] Á. Elbert, P.D. Siafarikas, Monotonicity properties of the zeros of ultraspherical polynomials, J. Approx. Theory 97 (1999), 31–39.
- [7] A. Fryant, Integral operators for harmonic functions, in "The mathematical heritage of C.F. Gauss", edited by G.M. Rassias, World Scientific, Singapore 1991, 304–320.
- [8] S.J. Gardiner, The Dirichlet problem with non-compact boundary, Math. Z. 213 (1993), 163–170.
- D. Khavinson, E. Lundberg, Linear holomorphic partial differential equations and classical potential theory. Amer. Math. Soc. Providence, RI (2018)
- [10] D. Khavinson, E. Lundberg, H. Render, Dirichlet's problem with entire data posed on an ellipsoidal cylinder, Potential Anal. 46 (2017), 55–62.
- [11] D. Khavinson, E. Lundberg, H. Render, The Dirichlet problem for the slab with entire data and a difference equation for harmonic functions, Canad. Math. Bull. 60 (2017), 146–153.
- [12] D. Khavinson, H.S. Shapiro, Dirichlet's problem when the data is an entire function, Bull. London Math. Soc. 24 (1992), 456–468.
- [13] H. Render, Real Bargmann spaces, Fischer decompositions and sets of uniqueness for polyharmonic functions, Duke Math. J. 142 (2008), 313–352.
- [14] H. Render, A characterization of the Khavinson-Shapiro conjecture via Fischer operators, Potential Anal. 45 (2016), 539–543.
- [15] Render, H., Aldaz, J. M.: Fischer decompositions for entire functions and the Dirichlet problem for parabolas. Anal. Math. Phys. 12, no. 6, Paper No. 150, 29 pp. (2022)
- [16] H.S. Shapiro, An algebraic theorem of E. Fischer and the Holomorphic Goursat Problem, Bull. London Math. Soc. 21 (1989), 513–537.
- [17] Szegö, G. Orthogonal Polynomials, Vol. 23 American Mathematical Society, 1975 432 pages.

H. RENDER: SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY COLLEGE DUBLIN, DUBLIN 4, IRELAND.

Email address: hermann.render@ucd.ie

10

J.M. Aldaz: Instituto de Ciencias Matemáticas (CSIC-UAM-UC3M-UCM) and Departamento de Matemáticas, Universidad Autónoma de Madrid, Cantoblanco 28049, Madrid, Spain.

Email address: jesus.munarriz@uam.es

Email address: jesus.munarriz@icmat.es