

A NOTE ON THE ORDER OF THE TATE–SHAFAREVICH GROUP MODULO SQUARES

ALEXANDROS KONSTANTINOU

ABSTRACT. We investigate the order of the Tate–Shafarevich group of abelian varieties modulo rational squares. Our main result shows that every square-free natural number appears as the non square-free part of the Tate–Shafarevich group of some abelian variety, thereby validating a conjecture of W. Stein.

CONTENTS

1. Introduction	1
2. Expression for III mod squares	2
3. III of non-square order	5
References	7

1. INTRODUCTION

In the following note, we study the order of the Tate–Shafarevich group $\text{III}(A/K)$ of an abelian variety A defined over a number field K . Its non-zero elements correspond to principal homogeneous spaces for A/K with a rational point over all completions of K though lacking a point in K itself. Despite its central role in the study of rational points, several foundational questions regarding $\text{III}(A/K)$ remain open. These include the Tate–Shafarevich conjecture which stipulates its finiteness. In addition, a conjecture of Stein focuses on the possible values that the order of the Tate–Shafarevich group can take (if finite) modulo rational squares. In the following note, we validate Stein’s conjecture, which can be phrased as follows.

Theorem 1.1 (=Corollary 3.6). *For every positive square-free integer n , there exists an abelian variety A/\mathbb{Q} with finite Tate–Shafarevich group of order nm^2 for some integer $m \geq 1$.*

1.1. Previous work. In the case of principally polarised abelian varieties, the Cassels–Tate pairing,

$$\langle, \rangle : \text{III}(A/K) \times \text{III}(A/K) \rightarrow \mathbb{Q}/\mathbb{Z},$$

2010 *Mathematics Subject Classification.* 11G10 (11G40).

lays down significant constraints on the structure of the Tate–Shafarevich group. In particular, for elliptic curves, this is known to be alternating, and therefore the order of the Tate–Shafarevich group of an elliptic curve must be a square, if finite.

The work of Flach [4] establishes the (slightly weaker) skew symmetric condition $\langle x, y \rangle = -\langle y, x \rangle$ for principally polarised abelian varieties. From this, one can *only* deduce that the Tate–Shafarevich group is either a square or twice a square. Building upon this, Poonen and Stoll present a criterion for deducing if the Tate–Shafarevich group (provided it’s finite) is of square order purely based on the vanishing of $\langle c, c \rangle$ for an element $c \in \text{III}(A/K)[2]$ canonically associated to A . They then use this criterion to present the first example of an abelian variety for which the order of its Tate–Shafarevich group is twice a square.

Subsequently, Stein [13] presents the first examples of abelian varieties defined over \mathbb{Q} in which the order of their Tate–Shafarevich groups satisfy $|\text{III}(A/\mathbb{Q})| = pm^2$ for $p < 25000$ an odd prime. His construction relies on the existence of an abelian variety defined over \mathbb{Q} which becomes isomorphic to a product E^{p-1} of elliptic curves over a cyclic extension. Following this, Keil’s work [6] presents examples of abelian surfaces, given as quotients of a product of two elliptic curves, for which $|\text{III}| = pm^2$ for $p \leq 13$ an odd prime.

1.2. Acknowledgements. I would like to extend my thanks to Vladimir Dokchitser for his guidance and many constructive discussions during the course of this work. In addition, I would like to thank Adam Morgan for many helpful suggestions on a draft version of this paper and for many useful discussions that came before that.

2. EXPRESSION FOR III MOD SQUARES

Our proof relies on breaking down the Weil restriction of scalars of an abelian variety up to isogeny followed by an application of a formula of Cassels–Tate. This allows us to express the size of the Tate–Shafarevich group modulo squares in terms of Birch–Swinnerton-Dyer constants.

2.1. Cassels–Tate formula. In the following subsection, we briefly recall a result of Cassels and Tate which proves invariance of the Birch–Swinnerton-Dyer conjecture under isogeny. We first fix some notation.

Notation 2.1. We let K be a number field. Given a K -rational isogeny $\phi : A_1 \rightarrow A_2$, we write

$$Q(\phi) = |\text{coker}(\phi : A_1(K)/A_1(K)_{\text{tors}} \rightarrow A_2(K)/A_2(K)_{\text{tors}})| \\ \times |\ker(\phi : \text{III}(A_1/K)_{\text{div}} \rightarrow \text{III}(A_2/K)_{\text{div}})|,$$

where III_{div} denotes the divisible part of III .

For an abelian variety A/K with a fixed non-zero global exterior form ω , we denote the Birch–Swinnerton-Dyer periods by

$$\Omega_{\mathbb{C}}(A, \omega) = 2^{\dim A} \int_{A(\mathbb{C})} |\omega \wedge \bar{\omega}|, \quad \Omega_{\mathbb{R}}(A, \omega) = \int_{A(\mathbb{R})} |\omega|.$$

We write $\Omega(A/K, \omega) = \prod_{v|\infty} \Omega_{K_v}(A, \omega)$ where $K_v = \mathbb{R}$ (resp. \mathbb{C}) if v is real (resp. complex). In addition, for a non-archimedean place v , we write $w_{A,v}^{\min}$ for a Néron minimal exterior form on A , $c_v(A/K)$ for its Tamagawa number and $C(A/K, \omega) = \prod_{v \nmid \infty} c_v(A/K) |\omega / w_{A,v}^{\min}|_v$. We write $\text{III}_0 = \text{III} / \text{III}_{\text{div}}$ and \square to denote the square of a rational number.

The following theorem is a version of the Cassels–Tate formula as found in [3, Theorem 4.3].

Theorem 2.2 (Cassels–Tate). *Let $\phi : A_1 \rightarrow A_2$ be a K -rational isogeny. Then, for any non-zero global exterior forms ω_1 and ω_2 on A_1 and A_2 ,*

$$\frac{Q(\phi^{\vee})}{Q(\phi)} = \frac{|A_2(K)_{\text{tors}}| |A_2^{\vee}(K)_{\text{tors}}|}{|A_1(K)_{\text{tors}}| |A_1^{\vee}(K)_{\text{tors}}|} \frac{C(A_1/K, \omega_1) \Omega(A_1/K, \omega_1)}{C(A_2/K, \omega_2) \Omega(A_2/K, \omega_2)} \prod_{p|\deg(\phi)} \frac{|\text{III}_0(A_1/K)[p^{\infty}]|}{|\text{III}_0(A_2/K)[p^{\infty}]|}.$$

Given a field extension L/K , we write ϕ_L for the base change of ϕ to L . The following proposition shows that for carefully chosen L and working modulo squares, the Cassels–Tate formula simplifies.

Lemma 2.3. *Let L/K be a quadratic extension such that all bad primes of A_1/K and A_2/K and all places $v \mid \deg(\phi)$ split in L . In addition, suppose that $\deg(\phi)$ is a square. Then,*

$$\frac{Q(\phi_L^{\vee})}{Q(\phi_L)} = \frac{|A_2(L)_{\text{tors}}| |A_2^{\vee}(L)_{\text{tors}}|}{|A_1(L)_{\text{tors}}| |A_1^{\vee}(L)_{\text{tors}}|} \prod_{p|\deg(\phi)} \frac{|\text{III}_0(A_1/K)[p^{\infty}]|}{|\text{III}_0(A_2/K)[p^{\infty}]|} \pmod{\square}.$$

Proof. We note that $C(A_1/L, \omega_1) \Omega(A_1/L, \omega_1)$ is independent of ω_1 . By our assumptions on L/K , the only terms in this product we have to consider are complex places of L which lie above real places of K and non-archimedean places $w \nmid \deg(\phi)$. As in the proof of [9, Theorem 7.3], when w is complex, we have $\Omega_{\mathbb{C}}(A_1, \phi^* \omega_2) / \Omega_{\mathbb{C}}(A_2, \omega_2) = \deg(\phi) = \square$. For a non-archimedean place $w \nmid \deg(\phi)$, $|(\phi^* \omega_2 / w_{A_1,w}^{\min}) / (\omega_2 / w_{A_2,w}^{\min})|_w = 1$ (see [12, pp. 12]). To conclude, $\frac{C(A_1/L, \omega_1) \Omega(A_1/L, \omega_1)}{C(A_2/L, \omega_2) \Omega(A_2/L, \omega_2)} = \square$ giving the required result. \square

2.2. A decomposition for the Weil-restriction of scalars. In what follows, we let A/K be a principally polarised abelian variety defined over a number field K . We let p be an odd prime, F/K a cyclic extension of degree p and $\text{Res}_K^F A$ the restriction of scalars from F to K . For brevity, we write $B = \text{Res}_K^F A$. This is an F/K -twist of A^p , and the product polarisation on A^p descends to a principal polarisation on B/K . Then, there are K -morphisms

$$i : A \rightarrow B, \quad \text{Tr} : B \rightarrow A.$$

These are the inclusion and the trace map respectively.

After identifying the base change $B_F = A^p$ as principally polarised abelian varieties, the base changed morphisms i_F and Tr_F coincide with the diagonal inclusion and the summation map respectively. In addition, writing $i^\vee : B \rightarrow A$ for the dual of i (having suppressed the canonical principal polarisations on both sides), we have $i^\vee = \text{Tr}$. Finally, we write R to denote the kernel of the trace map. This is an abelian variety defined over K and is an F/K -twist of A^{p-1} (see [13, Proposition 2.4]).

Lemma 2.4. *Let $\phi : R \times A \rightarrow B$ be given by $(x, y) \mapsto x + i(y)$. Then, ϕ is a K -rational isogeny with $\ker(\phi) \cong A[p]$.*

Proof. It suffices to show $\ker(\phi) \cong A[p]$. After base change to \overline{K} , we identify $R_{\overline{K}} = A^{p-1}$ and $B_{\overline{K}} = A^p$. Then, on \overline{K} -points ϕ is given by $(x_1, \dots, x_{p-1}, y) \mapsto (y + x_1, \dots, y + x_{p-1}, y - \sum_{i=1}^{p-1} x_i)$. Thus, restricting the projection $R \times A \rightarrow A$ to $\ker(\phi)$ allows us to deduce that $\ker(\phi) \cong A[p]$. \square

2.3. An expression for $\text{III}(R/L)$ modulo squares. In this subsection, we relate $|\text{III}(R/L)[p^\infty]| \bmod \square$ to the torsion subgroup of R and the rank of A/L .

Lemma 2.5. *Let L/K be a number field extension such that $L \otimes_K F$ is a field. Then, the following hold.*

- (1) *For all odd primes $q \neq p$, $|\text{III}_0(R/L)[q^\infty]| = \square$.*
- (2) *If, in addition, A is an elliptic curve, $|\text{III}_0(R/L)[2^\infty]| = \square$.*

Proof. By Lemma 2.4, $\deg(\phi) = p^{2\dim(A)}$. When $q \neq p$, this isogeny induces an isomorphism $\text{III}_0(A/L)[q^\infty] \times \text{III}_0(R/L)[q^\infty] \xrightarrow{\sim} \text{III}_0(B/L)[q^\infty]$. Since A and B are principally polarised, [10, Theorem 8] asserts that $\text{III}_0(A/L)[q^\infty]$ and $\text{III}_0(B/L)[q^\infty]$ are both of square order for all odd q . This proves (1). Further, by [2, Proposition A.5.2], $B_L \cong \text{Res}_L^{L \otimes_K F} A$, and by combining [8, pp. 178(a)] with Shapiro's Lemma, $\text{III}_0(B/L)[2^\infty] \cong \text{III}_0(A/L \otimes_K F)[2^\infty]$. Therefore, if A is an elliptic curve, both $\text{III}_0(A/L)[2^\infty]$ and $\text{III}_0(B/L)[2^\infty]$ are of square order by [10, Theorem 8, Corollary 12]. Claim (2) follows. \square

In what follows, we write $\text{rk}_p(A/K)$ for the \mathbb{Z}_p -corank of the p^∞ -Selmer group $\text{Sel}_{p^\infty}(A/K) := \varinjlim_{n \geq 1} \text{Sel}_{p^n}(A/K)$. This coincides with the Mordell–Weil rank of A plus the multiplicity of $\mathbb{Q}_p/\mathbb{Z}_p$ in its Tate–Shafarevich group.

Proposition 2.6. *Let L/K be a quadratic extension such that all bad primes of A/K and all primes dividing either $\deg(\phi)$ or the relative discriminant $\Delta(F/K)$ split in L . Then, the following hold.*

- (1) *Up to rational squares,*

$$|\text{III}_0(R/L)[p^\infty]| \equiv \frac{Q(\phi_L)}{Q(\phi_L^\vee)} \frac{|R(L)[p^\infty]|}{|R^\vee(L)[p^\infty]|} \bmod \square.$$

- (2) *If, in addition, $\text{rk}_p(A/L) = \text{rk}_p(A/L \otimes_K F)$, then*

$$|\text{III}_0(R/L)[p^\infty]| \equiv p^{\text{rk}_p(A/L)} \frac{|R(L)[p^\infty]|}{|R^\vee(L)[p^\infty]|} \bmod \square.$$

Proof. Let v be a place of bad reduction for B/K . By [8, Proposition 1], this can only happen when A/K has bad reduction at v or when $v \mid \Delta(F/K)$. This shows that all bad primes of B/K split in L . Since A/K is principally polarised, $|A(L)[p^\infty]| = |A^\vee(L)[p^\infty]|$ and $|\text{III}_0(A/L)[p^\infty]| = |\text{III}_0(A^\vee/L)[p^\infty]|$. The same holds for B . Part (1) follows by combining Lemma 2.3 and Lemma 2.4. For the second claim, it suffices to show that if $\text{rk}_p(A/L) = \text{rk}_p(A/L \otimes_K F)$, then $Q(\phi_L)/Q(\phi_L^\vee) = p^{\text{rk}_p(A/L)}$. After suppressing the principal polarisations $A = A^\vee, B = B^\vee$, we view ϕ^\vee as a morphism $B \rightarrow A \times R^\vee$. Therefore, $\frac{Q(\phi_L)}{Q(\phi_L^\vee)} \equiv Q(\phi_L^\vee \circ \phi_L : A \times R \rightarrow A \times R^\vee) \pmod{\square}$. Since the degree of $\phi^\vee \circ \phi$ is a power of p , it induces an isomorphism $\text{III}(A \times R/L)_{\text{div}}[q^\infty] \xrightarrow{\sim} \text{III}(A \times R^\vee/L)_{\text{div}}[q^\infty]$ for $q \neq p$. From this, we deduce $\ker(\phi_L^\vee \circ \phi_L \mid \text{III}_{\text{div}}) = \ker(\phi_L^\vee \circ \phi_L \mid \text{III}_{\text{div}}[p^\infty])$. Since $B_L \cong \text{Res}_L^{L \otimes_K F} A$ (see [2, Proposition A.5.2]), then by combining [8, pp. 178(a)] with Shapiro's Lemma gives $\text{Sel}_{p^\infty}(B/L) \cong \text{Sel}_{p^\infty}(A/L \otimes_K F)$. By assumption, $\text{rk}_p(A/L) = \text{rk}_p(A/L \otimes_K F)$, from which we deduce $\text{rk}_p(R/L) = \text{rk}_p(R^\vee/L) = 0$. As a result, $Q(\phi_L^\vee \circ \phi_L)$ coincides with $Q(i_L^\vee \circ i_L)$, where we view $i_L^\vee \circ i_L$ as an endomorphism of A_L . Using the description of i and i^\vee from §2.2, $i^\vee \circ i = [p]_A$. This gives the required result. \square

3. III OF NON-SQUARE ORDER

We retain all the notation from §2. In this section, we restrict to the case when $A = E$ is an elliptic curve, and so R is an abelian variety of dimension $p - 1$. In addition, we fix the following notation.

Notation 3.1. We write K_∞ to denote the cyclotomic \mathbb{Z}_p -extension of \mathbb{Q} , and $K_n = \mathbb{Q}(\zeta_{p^n} + \zeta_{p^n}^{-1})$ for the n^{th} layer of this extension. For d a positive, square-free integer, we let $L_n = K_n \cdot \mathbb{Q}(\sqrt{-d})$.

We first show that for n sufficiently large, $\text{III}(R/L_n)$ is finite for any choice of d . Further, having carefully selected d , we show that $|\text{III}(R/L_n)| \equiv p \pmod{\square}$. We begin by proving a lemma involving torsion on R and R^\vee .

Lemma 3.2. *Let E be an elliptic curve defined over \mathbb{Q} . Then, for any $n \geq 0$ and any prime $p \geq 11$ both $R(L_n)[p^\infty]$ and $R^\vee(L_n)[p^\infty]$ are trivial.*

Proof. As abelian varieties over L_{n+1} , both R and R^\vee are isomorphic to E^{p-1} . It therefore suffices to show that $E(L_{n+1})[p^\infty]$ is trivial for $p \geq 11$. Let E_{-d} be the quadratic twist of E by $-d$. Then, there exists a map $E(L_{n+1}) \rightarrow E(K_{n+1}) \times E_{-d}(K_{n+1})$ whose kernel and cokernel are 2-groups. Then, for any odd prime p , this induces an isomorphism

$$E(L_{n+1})[p^\infty] \xrightarrow{\sim} E(K_{n+1})[p^\infty] \times E_{-d}(K_{n+1})[p^\infty].$$

By [1, Theorem 1.1], $E(K_\infty)[p^\infty] = E(\mathbb{Q})[p^\infty]$ (similarly for E_{-d}) for any $p \geq 5$. If $p \geq 11$, Mazur's theorem [7, Theorem 2] asserts that $E(\mathbb{Q})[p^\infty]$ and $E_{-d}(\mathbb{Q})[p^\infty]$ are both trivial. Therefore, $E(L_{n+1})[p^\infty]$ is trivial. \square

The following result is a formal consequence of a theorem by Kato and Rohrlich.

Theorem 3.3. *Let E be an elliptic curve defined over \mathbb{Q} . Then, there exists a positive integer n such that*

- (1) $\text{ord}_{s=1} L(E/L_n, s) = \text{ord}_{s=1} L(E/L_{n+1}, s)$,
- (2) $\text{III}(R/L_n)$ is finite,
- (3) $\text{rk}_p(E/L_{n+1}) = \text{rk}_p(E/L_n)$.

Proof. As E is defined over \mathbb{Q} , $\text{Res}_{K_n}^{L_n} E$ is isogenous to $E \times E_{-d}$ giving an equality of L -functions

$$L(E/L_n, s) = L(E/K_n, s) L(E_{-d}/K_n, s),$$

By Rohrlich [11, pp. 409], the sequence $\{\text{ord}_{s=1} L(E/K_n, s)\}_{n \geq 1}$ stabilises (similarly for E_{-d}), and so $\text{ord}_{s=1} L(E/L_n, s) = \text{ord}_{s=1} L(E/L_{n+1}, s)$ for n sufficiently large. This proves (1). Part (2) is a consequence of a result by Kato [5]. Following the notation introduced there, for G an abelian group, $\tau \in \widehat{G}$, and a G -module M , we let $M^{(\tau)} := \{x \in M : I_\tau \cdot x = 0\}$ where $I_\tau \subseteq \mathbb{Z}[G]$ denotes the kernel of the map $\mathbb{Z}[G] \rightarrow \mathbb{C}^\times$ induced by τ . We apply this to $G = \text{Gal}(L_{n+1}/\mathbb{Q}) \cong \mathbb{Z}/2p^{n+1}\mathbb{Z}$. For brevity, we write $H = \text{Gal}(L_{n+1}/L_n) \cong \mathbb{Z}/p\mathbb{Z}$. Further, the composition $i \circ \text{Tr}$ from §2.2 is equal to $N_H := \sum_{h \in H} h$ viewed as a K -endomorphism of B . It follows that the composition $R \hookrightarrow B \xrightarrow{N_H} B$ is the zero map, which in turn gives a homomorphism $\text{III}(R/L_n) \rightarrow \ker(N_H \mid \text{III}(E/L_{n+1}))$. We claim this has finite kernel. To see this, we look at the short exact sequence of abelian varieties

$$0 \rightarrow R \rightarrow B \xrightarrow{\text{Tr}} E \rightarrow 0.$$

Taking Galois cohomology gives an exact sequence

$$0 \rightarrow E(L_n)/\text{Norm}_{L_n}^{L_{n+1}} E(L_{n+1}) \rightarrow H^1(L_n, R) \rightarrow H^1(L_n, B).$$

By the Mordell–Weil theorem $E(L_n)/pE(L_n)$ is finite, from which we deduce that $E(L_n)/\text{Norm}_{L_n}^{L_{n+1}} E(L_{n+1})$ must be finite. Therefore, the induced map $\text{III}(R/L_n) \rightarrow \ker(N_H \mid \text{III}(E/L_{n+1}))$ has finite kernel. For a primitive character $\tau \in \widehat{G}$, we have $N_H \in I_\tau$. Therefore, for such τ , $\ker(N_H \mid \text{III}(E/L_{n+1})) \subseteq \text{III}(E/L_{n+1})^{(\tau)}$. By Artin formalism,

$$L(E/L_{n+1}, s) = L(E/L_n, s) \prod_{\substack{\chi \in \widehat{H} \\ \chi \neq \text{id}}} L(E/L_n, \chi, s).$$

We let n be as in part (1) and χ a non-trivial character of H . Then, $L(E/L_n, \chi, s) = L(E/\mathbb{Q}, \text{Ind}_H^G \chi, 1) \neq 0$. Therefore, $L(E/\mathbb{Q}, \tau, 1) \neq 0$ for all primitive characters $\tau \in \widehat{G}$. Then, by [5, Corollary 14.3], the τ -part $(\text{III}_{n+1})^{(\tau)}$ is finite for all primitive $\tau \in \widehat{G}$. This completes the proof of (2). For (3), it suffices to show that $R(L_n)$ is finite. This follows in the same way as (2). \square

Theorem 3.4. *Let $p \geq 11$ be a prime. Then, there exists an abelian extension L/\mathbb{Q} and an abelian variety R/L such that $|\text{III}(R/L)| = p \cdot \square$.*

Proof. We let E/\mathbb{Q} be an elliptic curve defined over \mathbb{Q} and d a positive integer such that p and all bad primes of E/\mathbb{Q} split in $\mathbb{Q}(\sqrt{-d})$. For the remainder of this proof, we fix n as in Theorem 3.3 so that $\text{rk}_p(E/L_{n+1}) = \text{rk}_p(E/L_n)$ and $\text{III}(R/L_n)$ is finite. In view of Lemma 2.5(1)-(2), we deduce that $|\text{III}(R/L_n)| \equiv |\text{III}_0(R/L_n)[p^\infty]|$. Combining this with Proposition 2.6(2) (applied with $K = K_n, F = K_{n+1}, L = L_n$) and Lemma 3.2, we get

$$(\dagger) \quad |\text{III}(R/L_n)| \equiv p^{\text{rk}_p(E/L_n)} \pmod{\square}.$$

Since E is defined over \mathbb{Q} , its global root number $\omega(E/\mathbb{Q}(\sqrt{-d})) = -1$. This is because the only contribution in the root number computation comes from the unique archimedean place of $\mathbb{Q}(\sqrt{-d})$. Since $\text{Res}_{\mathbb{Q}}^{\mathbb{Q}(\sqrt{-d})} E$ is isogenous to $E \times E_{-d}$, we deduce that

$$(-1)^{\text{rk}_p(E/\mathbb{Q}(\sqrt{-d}))} = (-1)^{\text{rk}_p(E/\mathbb{Q})} (-1)^{\text{rk}_p(E_{-d}/\mathbb{Q})} \stackrel{[3, \text{Thm 1.4}]}{=} w(E/\mathbb{Q})w(E_{-d}/\mathbb{Q}).$$

In view of the isogeny, $w(E/\mathbb{Q}(\sqrt{-d})) = w(E/\mathbb{Q})w(E_{-d}/\mathbb{Q})$, from which we deduce that $\text{rk}_p(E/\mathbb{Q}(\sqrt{-d})) \equiv 1 \pmod{2}$. Since the parity of p^∞ -Selmer ranks remains unchanged in odd degree Galois extensions [3, Corollary 4.15], then $\text{rk}_p(E/L_n) \equiv \text{rk}_p(E/\mathbb{Q}(\sqrt{-d})) \equiv 1 \pmod{2}$. Combining this with (\dagger) gives the required result. \square

Theorem 3.5. *Let p be a rational prime. Then, there exists an abelian variety R'/\mathbb{Q} such that $|\text{III}(R'/\mathbb{Q})| = p \cdot \square$.*

Proof. For $p \geq 11$, we consider $R' = \text{Res}_{\mathbb{Q}}^L R$ where R is the abelian variety from Theorem 3.4. Since $\text{III}(R'/\mathbb{Q}) \cong \text{III}(R/L)$ (combine Shapiro's lemma with [8, pp. 178(a)]), the result follows by Theorem 3.4. For $p \leq 7$, see [10] and [13]. \square

Corollary 3.6. *For every positive square-free integer n , there exists an abelian variety defined over \mathbb{Q} with finite Tate-Shafarevich group of order $n \cdot \square$.*

Proof. Given abelian varieties A_1, A_2 , $\text{III}(A_1 \times A_2) \cong \text{III}(A_1) \times \text{III}(A_2)$. The result follows from Theorem 3.5. \square

REFERENCES

- [1] M. Chou, H. B. Daniels, I. Krijan, F. Najman, *Torsion groups of elliptic curves over the \mathbb{Z}_p -extensions of \mathbb{Q}* , New York Journal of Mathematics 27, (2021), 99–123.
- [2] B. Conrad, O. Gabber, G. Prasad, *Pseudo-reductive groups*, 2nd ed. Cambridge University Press, (2015).
- [3] T. Dokchitser, V. Dokchitser, *On the Birch-Swinnerton-Dyer quotients modulo squares*, Annals of Mathematics, (2010), 567–596.
- [4] M. Flach, *A generalisation of the Cassel-Tate pairing*. Journal für die reine und angewandte Mathematik 412, (1990), 113–127.
- [5] K. Kato, *p -adic Hodge theory and values of zeta functions of modular forms*, Astérisque 295, (2004), 117–290.

- [6] S. Keil, *Examples of non-simple abelian surfaces over the rationals with non-square order Tate–Shafarevich group*, Journal of Number Theory 144, (2014), 25-69.
- [7] B. Mazur (with an appendix by D. Goldfeld) *Rational isogenies of prime degree*, Inventiones mathematicae 44, (1978), 129-162.
- [8] J. S. Milne, *On the arithmetic of abelian varieties*, Inventiones mathematicae 17, (1972), 177-190.
- [9] J. S. Milne, *Arithmetic Duality Theorems*, Academic Press, Inc. (1986).
- [10] B. Poonen, M. Stoll, *The Cassels–Tate pairing on polarized abelian varieties*, Annals of Mathematics, (1999), 1109-1149.
- [11] D. Rohrlich, *On L -functions of elliptic curves and cyclotomic towers*, Inventiones mathematicae 75.3, (1984), 409-423.
- [12] E. F. Schaefer, *Class groups and Selmer groups*. Journal of Number Theory 56.1, (1996), 79-114
- [13] W. A. Stein, *Shafarevich–Tate groups of nonsquare order*, in Modular curves and Abelian Varieties, Progr. Math 224, Birkhäuser, Basel (2004), 277–289.

UNIVERSITY COLLEGE LONDON, LONDON WC1H 0AY, UK
Email address: alexandros.konstantinou.16@ucl.ac.uk