

REASSESSING RELATIONALITY FOR BIPOLAR DATA

MANUEL CUERNO, FERNANDO GALAZ-GARCÍA, SERGIO GALAZ-GARCÍA,
AND TELMO PÉREZ-IZQUIERDO

ABSTRACT. Methods for clustering people into construals—social affinity groups of individuals who share similarities in how they organize their outlooks on a collection of issues—have recently gained traction. Relational Class Analysis (RCA) is currently the most commonly used method for construal clustering. RCA has been applied to identify affinity groups in social spheres as varied as politics, musical preferences, and attitudes towards science. In this study, we highlight limitations in RCA’s ability to accurately identify the number and underlying structure of construals. These limitations stem from RCA’s mathematical underpinnings and its insensitivity to the bipolar structure of the survey items, which require respondents to place themselves in a support or rejection space and then express the intensity of their support or rejection. We develop an alternative method, which we call Bipolar Class Analysis (BCA), that aims to address this foundational limitation. BCA conceptualizes people’s attitudinal positions as moving along support/rejection semispaces and assesses similarity in opinion organization by taking into account position switches across these semispaces. We conduct extensive simulation analyses, with data organized around different construals, to demonstrate that BCA clusters individuals more accurately than RCA and other available alternatives. We also replicate previous analyses to show that BCA leads to substantively different empirical results than those produced by RCA in its original and later versions, and by Correlational Clustering Analysis (CCA), a method that has been proposed as an alternative to RCA.

arXiv:2404.17042v1 [cs.SI] 25 Apr 2024

. (M. Cuerno) DEPARTMENT OF QUANTITATIVE METHODS, CUNEF UNIVERSIDAD, SPAIN
(F. Galaz-García) DEPARTMENT OF MATHEMATICAL SCIENCES, DURHAM UNIVERSITY, UNITED KINGDOM
(S. Galaz-García) SOCIAL SCIENCES DEPARTMENT / INSTITUTO JUAN MARCH, UNIVERSIDAD CARLOS III-MADRID, SPAIN
(T. Pérez-Izquierdo) BCAM - BASQUE CENTER FOR APPLIED MATHEMATICS, SPAIN
. *E-mail addresses:* manuel.mellado@cunef.edu, fernando.galaz-garcia@durham.ac.uk, sergio.galaz@uc3m.es, tperezizquierdo@bcamath.org.
. *Date:* April 29, 2024.
. *Key words and phrases.* Quantitative Methodology, Construal Clustering, Relational Class Analysis, Correlational Class Analysis, Bipolar Class Analysis, Simulation Analysis .

SECTION 1. INTRODUCTION

Scholarly interest in Construal Clustering Methods (CCMs) has notably grown in recent years. These methods analyze responses to attitudinal surveys to estimate the number, relative population, and underlying characteristics of “construals”—groups of people that use similar shared meanings to interpret a social domain (DiMaggio et al. 2018). CCMs identify construals by clustering people by “relationality”, a term that describes how closely respondents’ opinions co-vary in terms of direction and distance. Increasingly used in empirical sociological analysis, CCMs have been applied to identify construals for issues as diverse as musical tastes, political attitudes, and beliefs about science and religion (Goldberg 2011; Baldassarri and Goldberg 2014; Boutyline 2017; DiMaggio and Goldberg 2018; Sotoudeh and DiMaggio 2023, p. 1842).

Fruitful methodological developments have also taken place alongside these empirical applications. New versions of Relational Class Analysis (RCA), the first CCM made available for social researchers (Goldberg 2011) have been developed (Goldberg and Stein 2018; Sotoudeh and DiMaggio 2023). Boutyline (2017) introduced an alternative method, Correlational Class Analysis (CCA), and showed that this method performs better than RCA when the variation in opinions within the same construal is assumed to be linear. More recently, Sotoudeh and DiMaggio (2023) proposed several variants of RCA and found they match CCA’s capacity to accurately identify respondents’ construal membership.

These developments attest to the growing recognition of CCMs as crucial instruments for addressing what Sotoudeh and DiMaggio (2023, p. 1842) call the “heterogeneity problem” in inferring people’s understandings from survey data: social analysts may only validly infer the meaning of someone’s position towards a specific issue by taking into account her opinions on other issues. Yet, our investigation shows that CCMs cannot effectively tackle this heterogeneity problem with the bipolar survey items they typically use. In this paper, we introduce Bipolar Class Analysis (BCA), a method for construal clustering that overcomes the limitations of current research models in generating valid construal mappings.

Bipolar survey items require respondents to express how much they agree or disagree with a specific question statement (Krosnick 1999; Malhotra and Krosnick 2007; Ostrom 1987). These types of questions are among the most common in social surveys and are the predominant form used for construal clustering. Bipolar questions demand that respondents consider whether they position themselves in a state of rejection or support and also rate the intensity of this position (Ostrom, Betz, and Skowronski 1992). Consequently, a meaningful measure of relationality based

on bipolar data should prioritize identifying construals according to how people’s opinions transition between rejection and support spaces. Neither CCA nor RCA in any of its variants addresses this aspect. They conceptualize co-variation only in terms of local shifts in direction; these calculations of directions are indifferent to whether they result in shifts in positions across rejection or support spaces.

Previous methods used by CCMs to compute relationality, in terms of co-variation in distance of opinion shifts, are problematic due to the varying number of opinion expressions available to respondents across survey items. This issue is commonly understood in terms of response point heterogeneity, yet this interpretation is reductive. It fails to acknowledge that variations in the number of opinion options represent, primarily, variations in the meanings available for respondents to choose from. The approach CCMs use to calculate distances is problematic because it involves mapping survey items into a continuous scale (the real line), potentially assigning identical numerical values to opinion expressions that are in fact different. The calculation of co-variation in distance becomes even more problematic in bipolar data because the number of intensity expressions available for respondents to choose from decreases once they position themselves within a rejection or support semispace. Consequently, the opinions available are limited to those within that specific semispace (Ostrom, Betz, and Skowronski 1992). This reduction in the number of effective choices complicates interpreting opinion responses as continuous cardinal variables, an approach that current CCMs use to compute co-variations in distance.

BCA measures relationality based on how respondents toggle back and forth between rejection or support semispaces. A key aspect of BCA is the introduction of a *polarity function*, which captures the differences in position, relative to rejection or support semispaces, between the answers given by two respondents. Additionally, we propose two adjusted measures of the magnitude of rejection/support intensities. The first measure disregards these intensities, focusing solely on the position of answers within the semispaces. The second measure computes the magnitude based on the distance (number of steps) of the answer to each question relative to the corresponding *neutrality* option. This last magnitude is adjusted to account for disparities in the number of available opinion expressions.

We test BCA’s performance against RCA and CCA using a wide variety of simulated data. Previous data generation processes (DGPs) directly simulate survey answers for synthetic respondents and model construals as linear variations between them (Boutyline 2017; Sotoudeh and DiMaggio 2023). In contrast, our DGP is based on ordered choice and stochastic multivariate dependency. In

this model, respondents hold a specific latent position towards each question, which they use to select from a finite set of opinion alternatives. Respondents' positions across questions are constrained through a global, non-linear dependency structure. This dependency, which can be interpreted as a construal, is explicitly known by design. Our approach more closely approximates the cognitive processes by which people choose among available opinion expressions. Additionally, our DGP avoids the linear modeling of constraints between variables, a method we find to be unrealistic given the existing results in the survey item measurement literature concerning how respondents make their choices.

We show that BCA outperforms CCA and all variations of RCA in accurately identifying the construal membership of respondents, as indicated by higher NMI values after they are adjusted by their upward bias when a large number of construals is estimated (Amelio and Pizzuti 2015; Boutyline 2017, 388). In addition, we show that BCA performs better in two other important indicators: the percentage of times a method estimates the correct number of construals and correlational dissimilarity, an original measure of qualitative accuracy. Correlational dissimilarity quantifies how well the dependency structure of the estimated construals approximates that of real ones.

We make three interrelated and complementary contributions. First, we introduce a method capable of producing more valid and accurate estimations of construals for a dominant type of survey items. This approach not only improves estimation accuracy but also highlights the need for CCMs that address heterogeneities preceding those related to the interpretation of shared meanings, specifically in the metric structure of survey items and the cognitive processes people use to express opinions. Second, we propose a new DGP strategy to more accurately capture these cognitive processes. Third, we enhance the performance evaluation of CCMs by using an adjusted Normalized Mutual Information (NMI) scale and introducing new performance instruments for qualitative accuracy.

The paper is structured as follows: First, in Section 2, we describe RCA and CCA, questioning their ability to produce valid construal descriptions from bipolar data. In Section 3, we introduce BCA as a new construal clustering method, highlighting its differences relative to previous methods. Section 4 reports the results of extensive simulation analyses that examine BCA's accuracy in describing construal structures relative to RCA and CCA. In Section 5, we discuss the differences in the results of construals that BCA produces in data where RCA and CCA have been previously applied. Finally, in Section 6, we recap the strengths of BCA and discuss further avenues for research in construal clustering.

SECTION 2. CONSTRUAL CLUSTERING METHODS: PROCEDURES AND LIMITATIONS

Bipolar survey items require respondents to express either support (affinity/preference) or opposition (disliking/rejection) to a specific question statement. These items compel respondents to first decide whether they position themselves in a positive or negative space relative to the statement, and second, to express the intensity of their stance within that space (Ostrom, Betz, and Skowronski 1992). The range of support or opposition expressions can vary from a single option in binary ‘agree’ or ‘disagree’ formats to potentially infinite responses, though in practice, the number rarely exceeds 50—this is typical for 0–100 “thermometer” scores. Bipolar questions feature a neutral position, which may be presented explicitly (e.g., “neither agree nor disagree”) or remain implicit, such as a latent midpoint in an “agree or disagree” scale.

Bipolar survey items are prevalent in social survey research and are the predominant type of question used in applied CCM research. For instance, 54% of the survey items analyzed in the multiple-dataset study by Soutodeh and DiMaggio (2023) are bipolar. This figure jumps to 97–100% for datasets related to key social research topics such as cultural consumption, and attitudes towards politics, science, religion, and the influence of the market in social life (DiMaggio et al. 2018; DiMaggio and Goldberg 2018). This section will introduce the currently available CCMs and discuss their performance in handling bipolar data.

Relational Class Analysis (RCA) was the first construal clustering method available for sociological research (Goldberg 2011). Since its introduction, RCA has rapidly become the most widely used method of its type for sociological analysis, as evidenced by its frequent application in a range of studies (Goldberg 2011; Baldassarri and Goldberg 2014; DiMaggio and Goldberg 2018; DiMaggio et al. 2018; Sotoudeh and DiMaggio 2023). Additionally, Sotoudeh and DiMaggio (Sotoudeh and DiMaggio 2023) have recently updated RCA to improve its performance.

RCA calculates the *relationality* between two respondents, a number ranging from -1 to 1 measuring the similarity of their respective construals. If this number is close to 1 in absolute value, the respondents are considered to belong to the same construal. After computing relationalities for all respondent pairs, RCA uses a graph partitioning algorithm (Newman 2004, 2006) to obtain the number of construals.

To estimate relationality, RCA first maps the opinion answers of each respondent to the interval $[0, 1]$, as a preliminary step before calculations begin (Goldberg 2011, 1406). The method then considers the differences between the answers a respondent provides to every pair of questions,

termed *pair differences*. RCA computes relationality between two respondents in two steps ¹. First, RCA computes a magnitude for each question pair by subtracting the absolute value of the corresponding pair differences of the two respondents. This magnitude is multiplied by -1 if the pair differences have opposite signs. Then, RCA averages these magnitudes across all question pairs to produce a value between -1 and 1 .

More recently, Boutyline introduced Correlational Class Analysis (CCA) as an alternative to RCA for construal clustering (Boutyline 2017). CCA computes the Pearson correlation coefficient between the answers of a pair of respondents. For this purpose, Boutyline maps each respondent’s answers to the real line. In the final step, the absolute values of these correlation coefficients are used as inputs into Newman’s partitioning algorithm, similarly to RCA’s approach.²

RCA and CCA have offered innovative approaches to capturing similarities in the organization of social opinions. Nevertheless, we contend that the strategies these methods use to extract construals have several limitations which compromise their ability to yield accurate results.

We illustrate the challenges that RCA and CCA face in clustering respondents into construals through a motivating example from politics. This example includes issues that are currently highly divisive in the United States’ political landscape.

In our example, three people participate in a survey requiring them to express their support or opposition to three contentious public opinion items. The items are: “Undocumented migrants should be deported even without due process” (question 1), “Gun control should be free of background checks” (question 2), and “No further Medicaid expansion is needed” (question 3). Respondents can choose from seven options reflecting their level of agreement: **Strongly Disagree**, **Disagree**, **Somewhat Disagree**, **Neither Agree nor Disagree**, **Somewhat Agree**, **Agree**, and **Strongly Agree**.

Drawing on the findings from (Baldassarri and Goldberg 2014), it is known that the community where these three individuals reside has two distinct political construals. The first is termed “traditional ideology”, wherein opinions on the three contentious issues are positively related. Members of this group align along an opinion axis that opposes “uncompromising progressives” and “uncompromising conservatives”. Uncompromising progressives are characterized by their strong rejection of migrant deportation, easy access to guns, and freezes in further health care provision. Conversely, uncompromising conservatives advocate strongly for immediate deportation

1. For a more detailed description of RCA, refer to Goldberg’s foundational paper (Goldberg 2011), as well as Boutyline’s explanation in (Boutyline 2017, Section: Relationality)

2. For further details on improvements to CCA and RCA, refer to (Sotoudeh and DiMaggio 2023, Appendix A: Improving CCA and RCA), which proposes switching from Newman’s partition algorithm to the Louvain group detection algorithm (Blondel et al. 2011). Sotoudeh and DiMaggio note that “this change of algorithm improved NMI for both RCA and CCA by 2%, from 73% to 75% and from 88% to 91% respectively”.

of migrants, unrestricted gun ownership, and no further expansion of Medicaid. Individual A is a moderate conservative within the “traditional ideology” construal. Her responses to the survey items are as follows.

$$A = (\text{Somewhat Agree}, \text{Somewhat Agree}, \text{Somewhat Agree}).$$

Respondent B belongs to the second construal, which we term “alternative ideology” for terminological consistency (Baldassarri and Goldberg 2014). In this construal, items 1 and 2 are positively related to one another but negatively related to item 3. At one end of this construal are “redistributive conservatives” who strongly support migrant deportation and the status quo on gun control, yet oppose halting Medicaid expansion. At the other end are “fiscally conservative progressives”, who outrightly reject migrant deportation and easy access to guns, and strongly oppose further expansions of Medicaid. Individual B, representing a political alternative, can be characterized as a consistently moderate redistributive conservative. Her responses are as follows:

$$B = (\text{Somewhat Agree}, \text{Somewhat Agree}, \text{Somewhat Disagree}).$$

Individual C, like individual A, aligns with the “traditional ideology” construal as a conservative. However, her opinions vary in intensity across the survey items, influenced by personal experiences. Her expressed opinions are as follows:

$$C = (\text{Somewhat Agree}, \text{Somewhat Agree}, \text{Strongly Agree}).$$

Figure 1 plots the absolute values yielded by RCA and CCA for each respondent pair, highlighting a clear discrepancy between the RCA and CCA values and the actual construal memberships. It is important to recall that values close to one indicate shared construals.

RCA and CCA assign the same value to the respondent pairs (A, B) and (A, C), even though only respondents A and C belong to the same construal. Furthermore, RCA assigns a larger relationality to respondents B and C, who are from different construals, than to respondents A and C, who share the same construal. Notably, CCA assigns the maximum value of 1 to the respondent pair (B, C). As a result, RCA and CCA might assign high values to pairs of respondents in different construals. Since the clustering algorithm uses these high values as a criterion to group respondents into estimated construals, RCA and CCA may produce invalid construal estimates.

The primary reason for these discrepancies is the failure of both RCA and CCA to account for the bipolar structure of the data. The opinion space of the three survey items can be conceptualized as the union of two semispaces: a positive opinion semispace and a negative opinion semispace.




FIGURE 1. Top: Answers of each respondent. Bottom: RCA, CCA and BCA scores for the pair (A, B) [left], (A, C) [center], and (B, C) [right].

The positive semispace consists of the options

$\{\text{Neither Agree nor Disagree, Somewhat Agree, Agree, Strongly Agree}\},$

representing neutrality and varying degrees of endorsement. Conversely, the negative opinion semispace comprises the options

$\{\text{Strongly Disagree, Disagree, Somewhat Disagree, Neither Agree nor Disagree}\},$

reflecting neutrality and different levels of opposition.

We consider that the response **Neither Agree nor Disagree** belongs to both semispaces as it represents a *neutrality* position relative to the statement in the corresponding question. A crucial aspect is that respondents typically align themselves within one of these semispaces when evaluating each survey item (Ostrom, Betz, and Skowronski 1992). Therefore, the positioning of each answer relative to these semispaces is crucial to understanding how respondents organize their opinions. Notably, neither RCA nor CCA effectively exploit this information.

As discussed above, RCA computes relationality based solely on differences in opinion answers among pairs of questions. This calculation method does not consider whether answers fall in the same semispace. Consequently, RCA assigns the same value to respondent pairs (A,B) and (A,C). Respondent B’s opinion about Medicaid expansion (**Strongly Agree**) is “two opinion points” higher than respondent A’s (**Somewhat Agree**), while respondent C’s opinion (**Somewhat Disagree**) is “two opinion points” lower than respondent A’s. Despite these opinions falling in different semispaces, RCA treats these relative positions as equivalent. A similar reasoning shows why RCA assigns a large relationality for respondents B and C: both respondents’ opinions regarding Medicaid expansion are at the same “opinion point distance” from their views on the other two issues.

CCA computes the correlation coefficient between the opinions of two respondents. Therefore, what matters for CCA is the deviation of each respondent’s opinion on each issue from their “mean opinion”. Importantly, a respondent’s “mean opinion” may not correspond to the point where the neutral answer is mapped, and it might not even correspond to an existing option. Consequently, being above or below the “mean opinion” generally does not correspond to being in distinct opinion semispaces. For example, the “mean opinion” of respondent B lies somewhere between **Somewhat Agree** and **Neither Agree nor Disagree**. This approach may lead to misleading results.

The issue is illustrated by CCA’s value for respondents B and C, which is the largest possible. When seen as deviations from each respondent’s “mean opinion”, the answers of respondent C mirror those of respondent B. Thus, the correlation between the two is -1 . CCA uses the absolute value of this correlation to cluster respondents into construals. As a result, respondents B and C may be incorrectly clustered together in the same construal, despite their contrasting positions.

In summary, RCA and CCA fail to effectively use the bipolar structure of the data, potentially leading to incorrect construal assignments. Beyond the issues highlighted in our motivating example, we have identified further concerns with the methods developed by Goldberg and Boutyline (2011; 2017). We conclude this section by discussing these concerns.




FIGURE 2. RCA’s rescaling of a 3-point agreement-disagreement question and a 5-point strong-agreement to strong-disagreement item.

First, CCA encounters a problem when clustering respondents who do not deviate from their “mean opinion”—those with the same answer across all items. Boutyline (2017) offers two ways to address this issue: either by removing these respondents from the sample or, as shown in Figure 1, by setting CCA’s value between “constant respondents” and “non-constant respondents” to zero. Note that, as acknowledged by Boutyline (2017, Appendix D), the latter option results in the creation of a “null construal” containing all respondents with the same answer across all items.

Second, when the number of response points across questions varies, RCA’s procedure to standardize answers can cause disparate opinion values to appear identical. As illustrated in Figure 2, this method assigns the value 1 to an **Agree** response from a 3-response point question. In a 5-response item, in turn, a **Strongly Agree** response is also mapped to 1. Given that one response incorporates an adverb of degree, signifying greater intensity, while the other does not, there are substantial grounds to contend that these answers, in principle, do not reflect opinions of identical intensity.

Finally, it is important to note that both RCA and CCA map responses into the real line, assigning them numerical values. This approach treats the responses as cardinal continuous variables, presupposing inherent distances between options in the opinion space. Although there may be methods to measure distances between potential answers to a given item (see Section 3), RCA and CCA require more. These methods need a way to compare responses across different survey

items—they need a “universal ruler”. This is a stringent condition: expressions of support or rejection intensity are often incomparable across different questions, particularly when they do not share the same number of response options. For example, simple “agreement” or “disagreement” from binary bipolar questions do not straightforwardly map into the answer space of a 5-point questions that differentiates between “completely” or “somewhat” agreeing or disagreeing ³. Therefore, we advocate for a method that does not rely on this universal ruler.

SECTION 3. BIPOLAR CLASS ANALYSIS

We introduce a new method, Bipolar Class Analysis (BCA), to address the limitations of RCA and CCA discussed in Section 2. Similar to Goldberg’s algorithm, BCA analyzes pairs of responses to survey items from two respondents. As previously discussed, the structure of semispaces is crucial for bipolar opinion items. Throughout this section, we assume that the opinion space can be split into a negative (rejection) and positive (support) opinion semispaces, intersecting at a neutral element.⁴ Tailored specifically for bipolar data, our method leverages the decomposition into opinion semispaces to achieve its analytical goals.


BCA has two main components: a computation based on a polarity function ω and a magnitude function μ , and a partitioning algorithm similar to those used in RCA and CCA. Like these methods, we use Newman’s partitioning algorithm to determine construals. We will discuss BCA in detail in the remainder of this section. For a formal development of the method, please refer to Appendix A.

The rationale behind the polarity function ω is to compare the permanence or change between opinion semispaces among pairs of different respondents’ answers. Functioning as a sign function, ω takes values in $\{-1, 0, 1\}$ based on the following criteria: If both pairs of answers remain within the same opinion semispace, change polarity in the same direction (either both from negative to positive or positive to negative semispace), one pair is neutrally static (both answers are the neutral element) while the other remains in the same semispace, or both pairs are static at the neutral element, then ω is assigned a value of 1. If both pairs of answers are in different opinion semispaces or change polarity in opposite directions, ω is assigned a value of -1 . In the remaining cases, ω is assigned a value of 0. We illustrate each case in Figure 3. This approach enables us to effectively use the bipolar structure of the data.

3. Goldberg 2011, in Appendix E, argues that “ordinal scales that use five or more categories strongly approximate continuous variables”. However, the related research (O’Brien and Homer 1987; Bollen and Barb 1981) primarily examines how well the correlation coefficient is approximated. There is no clear evidence that these findings extend to RCA computations, and the studies are limited to jointly normal variables. Thus, we find the evidence insufficient even for CCA’s computations.

4. The neutrality element may be explicit, as illustrated in Section 2, or implicit, as in questions offering only choices in favor of or against, where neutrality exists but is not an option.

FIGURE 3. Polarity function for BCA.



Notes: Black dots represent answers of respondent u to question items k and l ; white ones represent answers of a hypothetical respondent v for the same question.

As discussed in Section 2, computing meaningful distances between opinions presents challenges. First, survey questions may differ in the number of available answers; second, these answers are typically ordered only by the relative intensity of feeling. Constructing distances with desirable properties warrants detailed consideration in further research. With this in mind, BCA is designed around a general magnitude term for which we propose two approaches.

The first and simplest approach is to set the magnitude term, denoted as μ_1 , equal to one for all possible cases. In practice, this choice makes the computation of the first step of BCA solely dependent on how respondents move back and forth across semispaces. Given the bipolar structure of the data, this option is always feasible without requiring additional structural impositions. Therefore, we consider it a suitable benchmark. Depending on the data, one might favor alternative magnitude functions. Comparing results obtained using μ_1 with those from other functions can help assess their robustness.

The second approach for calculating the magnitude in BCA is based on the distance (number of steps) from each possible answer to the neutrality element. To compute the corresponding magnitude function, denoted by μ_2 , we require two conditions: (i) *neutrality* must be an explicit option, and (ii) there must be a defined order among the answers to each question. Then, μ_2 is constructed similarly to RCA’s magnitude. Unlike RCA, which uses distances between answers to different questions, BCA considers the variation in distance from each answer to its corresponding neutrality value. This method avoids comparing answers across different questions and provides an implicit normalization. This approach allows BCA to address more effectively the variation in response numbers compared to RCA. For more detailed information, please refer to Appendix A.

When we apply BCA to the example presented in the previous section, we observe that it accurately identifies whether the respondents come from the same construal (see Figure 1). Unlike the results from RCA and CCA, BCA yields higher values for respondents A and C (who are from the same construal) than for respondents A and B or B and C. In this straightforward case, the only difference between the two magnitude functions is the lower value that μ_2 assigns to the relationality between respondents A and C. This difference stems from their differing degrees of agreement with not expanding Medicaid.

Even though our example helps clarify ideas, several questions still need to be answered. For example, does BCA outperform RCA and CCA when applied to larger datasets? Are there substantive differences between using one magnitude over another? To address these questions, we simulate artificial data to systematically evaluate each method.

SECTION 4. SIMULATIONS

We conduct simulations to assess how BCA performs as a construal clustering method. In each simulation, synthetic data sets are generated, where N respondents express opinions on Q questions. These simulations adhere to a modeling strategy that incorporates ordered choice and multivariate dependency. Respondents hold a specific latent *position* towards each question, which they use to select *opinion* values from a finite set of alternatives. The positions of respondents across questions are constrained through a specific dependency structure. This structure, interpretable as a construal, is predetermined by design.

The data sets we simulate vary along several dimensions: (1) the number of respondents, (2) the number of construals; (3) the relative population of each construal; (4) the number of opinion points for each question; (5) the skewness of opinions towards the disagreement or agreement semispaces; and (6) the presence or absence of individual noise in the mapping of positions into opinions.

We generate these data sets following three general steps. First, we define the number of respondents (N) and questions (Q); second, we generate an $N \times Q$ matrix with latent positions X^* ; third, we map the latent positions to an $N \times Q$ -matrix of observed opinions X . The first step is self-explanatory. We elaborate on the second and third steps below and provide a more detailed description of these procedures in Appendix B.

Modeling positions

Respondents' positions on questions are modeled as continuous random variables ranging from 0 (full disagreement) to 1 (full agreement). These positions are not independent across questions: they are constrained by a dependency structure. This structure is encapsulated in the multivariate distribution of positions $F(x_1^*, x_2^*, \dots, x_Q^*)$. Using Sklar's Theorem (Theorem 2.10.9 in Nelsen 2006), the multivariate distribution can be decomposed as follows:

$$F(x_1^*, x_2^*, \dots, x_Q^*) = C(F_1(x_1^*), F_2(x_2^*), \dots, F_Q(x_Q^*)),$$

where the functions $F_q(\cdot)$ are the marginal distributions for each question q and $C(\cdot)$ is a copula function modeling the dependency structure. Sklar's Theorem thus allows us to separate question or individual specific variation (captured in the marginal distributions) from the dependency structure among questions (captured in the copula). The copula function $C(\cdot)$ quantifies the likelihood of finding an individual with a specific position on question 1, given their positions on the other issues. There is a wide variety of copulas that accommodate different dependency structures. We use

Gaussian copulas in our simulation due to their computational efficiency and ease of interpretation. The shape of these copulas is determined by a correlation matrix Σ .⁵

Modeling opinions

Respondents map their latent position onto an observable *opinion* by selecting from a finite (and typically small) number of ordered opinion categories. A respondent chooses a specific opinion value based on where her position lies relative to specific thresholds.⁶ Our model of the opinion spaces includes a “skewness parameter” that determines the number of responses falling in the positive (agreement) semispace relative to the number of responses in the negative (disagreement) semispace. For example, if this parameter is 0.2, then the proportion of the population with agreeing responses is 20% larger than that with disagreeing answers. We also allow for individual-specific random variation in the threshold values used to map positions onto opinions.

Experiments

We use the modeling processes described above to simulate opinion data. Each dataset from these simulations consists of 10 questions and includes 2, 3, or 4 construals, each populated by between 100 and 500 respondents. The specific structure for the k -th construal is predetermined by design and is governed by a correlation matrix Σ^k , where correlations between positions are randomly set. We report the results of three different experiments, each consisting of 1,000 simulated datasets.⁷

In Experiment 1, *Fixed Opinion Parameters*, all questions feature 7-opinion points. The skewness parameter is set to 0, and the thresholds used to convert positions into observable opinions are fixed and identical for all respondents.

Experiment 2, *Heterogeneous Opinion Points*, introduces variability in the response options: Questions are randomly assigned 3, 5, or 7 response alternatives. Skewness remains at 0, and position-mapping thresholds are homogeneous across respondents. The distance between thresholds varies with the number of points: for each question, partitioning the $[0, 1]$ position space into subintervals of equal length.

5. Appendix E provides a brief description of Gaussian copulas.

6. For instance, suppose that a respondent has a latent position of 0.7 about the first issue. If for this issue the “Somewhat Agree” threshold is 0.6 and the “Strongly Agree” threshold is 0.8, the respondent will choose the opinion “Somewhat Agree”.

7. In Appendix D, we provide additional results from four more experiments that apply RCA and CCA variants aligned with changes in clustering procedures observed in later works. These results are consistent with those reported here.

Experiment 3, *Heterogeneous Opinion Points and Thresholds*, introduces additional complexity: Questions continue to be randomly assigned 3, 5, or 7 response alternatives. However, skewness varies randomly between -0.3 and 0.3 , and ordered-choice thresholds are individual-specific.

Performance measures

We cluster respondents into construals for each simulated data set using BCA, RCA, and CCA.⁸ Then, we compare the results with the known structure of the construals present in each dataset. We use four measures of performance:

1. *Construal Number Accuracy*: This is the rate at which a method correctly estimates the number of construals present in the data.
2. *Normalized Mutual Information (NMI)*: This is a measure of accuracy in construal identification (Boutyline 2017, p. 365). NMI values range from zero to one, with higher values indicating greater accuracy in assigning the correct construal affiliation to each respondent. Recent studies have noted that NMI tends to favor methods that estimate a larger number of groups (Amelio and Pizzuti 2015; Boutyline 2017, p. 388; Mahmoudi and Jemielniak 2024).
3. *Scaled NMI*: To address the bias observed in NMI, following recommendations from the literature (Amelio and Pizzuti 2015), we also report scaled NMI, which adjusts NMI values downward when the estimated number of groups significantly exceeds the correct number.⁹
4. *Correlational Dissimilarity*: Discussed in detail in Appendix C, this metric quantifies how well a method estimates the underlying correlation structure of all the construals present in the data. Correlational dissimilarity values are based on comparing estimated correlation matrices with true correlation matrices. compares the estimated correlation matrices with the true matrices to quantify how accurately a method estimates the underlying correlation structure of the construals. Lower correlational dissimilarity values indicate a more accurate estimation of the underlying structure. If the estimated and true correlation matrices are equal, the correlational dissimilarity is zero, indicating no dissimilarity.

Results

We report the results of the simulation analyses in Table 1. Column 1 presents mean values for performance indicators when the value of the magnitude term in BCA is fixed at one ($\mu = \mu_1$).

8. We used the R implementation of both RCA and CCA. The packages are available for download at CRAN: the `RCA` package implements RCA and the `corclass` package implements CCA. Package versions were 2.0 for `RCA` and 0.2.1 for `corclass`, respectively.

9. This adjustment is analogous to the adjusted- R^2 in regression analysis.

We refer to this variant by BCA_1 . Column 2 corresponds to BCA using the distance-to-neutrality magnitude ($\mu = \mu_2$). We will refer to this variant by BCA_2 . Columns 3 and 4 report performance values for Goldberg’s original RCA (2011) and Boutyline’s CCA (2017), respectively.

	BCA_1	BCA_2	RCA	CCA
<i>1. Fixed Opinion Parameters</i>				
Construal number accuracy	0.387	0.405	0.028	0.061
Mean NMI	0.362	0.385	0.446	0.568
Mean Scaled NMI	0.292	0.312	0.187	0.284
Mean correlational dissimilarity	6.668	6.565	11.149	13.631
<i>2. Heterogeneous Opinion Points</i>				
Construal number accuracy	0.388	0.386	0.065	0.065
Mean NMI	0.307	0.318	0.377	0.435
Mean Scaled NMI	0.246	0.253	0.158	0.198
Mean correlational dissimilarity	4.986	5.018	7.605	10.533
<i>3. Het. Opinion Points and Thresholds</i>				
Construal number accuracy	0.410	0.397	0.046	0.203
Mean NMI	0.264	0.271	0.329	0.347
Mean Scaled NMI	0.214	0.217	0.136	0.220
Mean correlational dissimilarity	4.012	4.028	5.891	5.424

TABLE 1. Simulation results for BCA_1 with one-unit magnitude, BCA_2 with variable magnitude, RCA (Goldberg 2011), and CCA (Boutyline 2017).

Panel A reports performance results for Experiment 1, Fixed Opinion Parameters. The results show that RCA and CCA are relatively ineffective in accurately predicting the number of construals, achieving accurate predictions only 2.8% and 6.1% of the time, respectively. These rates are at least five times smaller than those observed for BCA methods. The performance of the two BCA variants is quite similar: BCA_1 accurately predicts the number of construals in 38.7% of cases, while BCA_2 does so in 40.5% of the instances. Additionally, BCA clustering procedures exhibit, on average, correlational dissimilarity values that are at least 40% smaller than those of RCA and CCA, indicating that BCA more effectively captures the underlying structures. NMI is the only performance measure where RCA and CCA exhibit higher values than those of BCA. At first glance, this might suggest that RCA and CCA more accurately identify construal adscription. However, the results for the Mean Scaled NMI tell a different story. Mean values for this indicator are consistently higher for BCA. These findings strongly suggest that the higher NMI scores for RCA and CCA are due to their tendency to divide respondents into more groups than BCA.

Panels B and C report performance results for Experiments 2 and 3. Experiment 2 involves varying the number of opinion points across questions, while in Experiment 3 opinion thresholds and data skewness also vary. Mean *scaled* NMI scores indicate that all clustering methods experience

a loss in classification accuracy with more variable data structures, with BCAs showing a slightly more pronounced decline. Consequently, in Experiments 2 and 3, BCA methods exhibit mean scaled NMI values similar to that of CCA (around 0.220), whereas RCA’s value is significantly lower at 0.136.

Regarding *construal number accuracy*, BCA methods consistently maintain a value of around 0.4. RCA values remain stable but significantly lower, at approximately 0.04. For CCA, however, construal number accuracy improves in Experiment 3, although its capacity to accurately estimate the number of groups is still roughly half that of BCA’s.

Finally, concerning *correlational dissimilarity*, all methods show improved (i.e., lower) values in Experiment 2 and further improvement in Experiment 3. In this latter experiment, RCA and CCA achieve mean values of 5.8 and 5.4, respectively, while BCA methods perform even better, with values of 4.028 for BCA_1 and 4.012 for BCA_2 .

The results strongly indicate that both variants of BCA are superior in capturing construal structures compared to RCA and CCA. BCA methods are more accurate in describing the correlation structure of construals and more capable of accurately estimating the number of construals. They are also just as effective as RCA and, particularly, CCA, in correctly classifying respondents’ construal ascription.

While there are performance differences between BCA_1 and BCA_2 , these differences are not systematic. BCA_2 , which uses a variable magnitude term, achieves better scaled NMI values, while BCA_1 , which sets the magnitude to 1, exhibits better values in construal number accuracy and mean correlational dissimilarity. However, these differences are marginal.

The comparable performance of BCA_1 to BCA_2 in capturing construal structures suggests that the ability to track how individuals toggle between rejection and support spaces is a robust predictor of these structures when analyzing actual data.

SECTION 5. EMPIRICAL RESULTS

The results of our simulations suggest that BCA captures construal structures more accurately than RCA and CCA. However, the critical question remains: Does this increased accuracy have substantive significance? How do the results produced by BCA with real data differ from those obtained using RCA and CCA?

To address these questions, we reanalyzed data sets from three empirical studies that previously employed construal clustering methods. The first data set, structurally the simplest, consists of responses from 1,532 respondents to a 7-point question asking whether people like or dislike 17

different musical genres. This data, drawn from the 1993 edition of the General Social Survey, was used by Goldberg to analyze musical tastes in the US (2011). Boutyline also applied CCA to this data for the same purpose (2017). The second data set comprises answers from 1,481 respondents on 15 variables related to opinions on science, religion, and spiritualism (SRS), with opinion points ranging from 2 to 6. This data originates from the 1998 General Social Survey and was analyzed in (DiMaggio et al. 2018). The third data set comprises responses from 611 individuals to 29 public opinion questions asked between 1984 and 2002, analyzed by Baldassarri and Goldberg (2014) to categorize respondents into different political ideological groups. This data set features 2, 3, 4, and 7-point opinion items.¹⁰

We explore differences across methods by examining the number of construals each method estimated and the relative population size of each construal. Results are presented in Table 2, which reports the number of construals yielded by BCA_1 , BCA_2 , RCA, and CCA for each data set, followed by the population of each construal, ordered by relative size. It is important to note that the construals reported in the same cell are *not* qualitatively comparable.

In the case of the data on science and spirituality, BCA methods yield results similar to those of RCA, with all estimating three construals of roughly equal size. In contrast, CCA estimates four construals with very unequal populations. For the music data, however, BCA_1 and BCA_2 produce notably different results, calculating two groups of roughly equal size, compared to RCA’s three similarly-sized construals and CCA’s five construals of very heterogeneous sizes. Regarding the ANES data, BCA methods estimate two construals of similar size, while RCA estimates three and CCA four.

SECTION 6. CONCLUSIONS

Construal clustering methods have emerged as a powerful methodological tool, paving the way for the rapid development of a new and promising line of empirical research dedicated to studying social affinity structures. This paper aims to advance this research agenda by introducing Bipolar Class Analysis (BCA), a method designed to address the mismatch between previous methods and the questions they analyze. In essence, BCA clusters respondents based on the similarity of their movements between rejection and support zones across different opinions. Our paper examines

10. To clean the data, we followed as closely possible and to the best of our abilities, the coding decisions outlined in the methodological appendices for these papers. All the cleaned data sets are available upon request.

11. Due to the various versions of RCA used in different studies (Goldberg 2011; Baldassarri and Goldberg 2014; DiMaggio et al. 2018), we opted to use the version of RCA in the *R*-package available at <https://cran.r-project.org/web/packages/RCA/index.html> for our computations. After analyzing all data sets with this RCA package, we noticed some discrepancies with the results presented in these articles. Consequently, the RCA column in this table consists of the results provided by the original authors.

	BCA ₁	BCA ₂	RCA ¹¹	CCA
GSS 1998: Science and Religion				
Number of construals	3	3	3	4
<i>1st largest</i>	42.42%	42.29%	40%	54.28%
<i>2nd largest</i>	31.29%	30.80%	39%	45.72%
<i>3rd largest</i>	26.28%	26.89%	21%	0.12%
<i>4th largest</i>				0.12%
GSS 1993: Music				
Number of construals	2	2	3	5
<i>1st largest</i>	50.2%	50.8%	44.5%	31.52%
<i>2nd largest</i>	49.8%	49.2%	30.7%	24.08%
<i>3rd largest</i>			24.8%	23.56%
<i>4th largest</i>				20.56%
<i>5th largest</i>				0.26%
ANES 2004: Public Opinion				
Number of construals	2	2	3	4
<i>1st largest</i>	54.17%	54.66%	39.93%	48.77%
<i>2nd largest</i>	45.82%	45.33%	31.58%	26.02%
<i>3rd largest</i>			28.47%	25.20%

TABLE 2. Groups are qualitatively incomparable across construal clustering techniques. BCA₁: Bipolar Class Analysis with 1-unit magnitude term; BCA₂: Bipolar Class Analysis with variable magnitude term; RCA: RCA presented in its full methodological discussion in Goldberg 2011; CCA: Correlational Class analysis.

the clustering performance of two variations of this method. In one variation, the calculation of relationality between pairs of respondents depends solely on their movements between rejection and support zones. In the other, BCA also incorporates a variable magnitude based on the distance to neutrality of responses.

We find that BCA, in both of its variations, significantly outperforms RCA and CCA in accurately detecting simulated construals. These simulations mimic the structure of real-life survey items where construal structures are predetermined. BCA is as effective as previous methods in accurately classifying the construal ascription of respondents and is superior in correctly identifying the number of construals in the data. Furthermore, BCA estimates correlational structures for these construals more closely approximating those observed in real datasets. Interestingly, we find no substantive difference in performance between the two variations of BCA. This finding suggests that construal clustering techniques focused solely on assessing how similarly people move across rejection/support

semispaces are capable of describing construal structures, regardless of the magnitude of these movements.

Additionally, the superior accuracy of BCA variants has tangible substantive implications when applied to real-world datasets previously analyzed. We discover that BCA estimates a different number and relative population of construals in these datasets.

We contend that Bipolar Class Analysis (BCA) marks a significant advancement towards fully leveraging the analytical potential of construal clustering in empirical sociological research. The introduction of BCA not only enhances current methodologies but also opens several avenues for further methodological refinement. A critical challenge remains in generating appropriate measures of opinion amid varying metrics and response scales. In response, this paper proposes a measure of magnitude that mitigates the influence of larger response domains in estimating distances. Further exploration of this measure's performance, along with the development of alternative measures, represents a promising direction for future research.

Supplementary material¹²

Technical exposition of the mathematical framework.

12. This is a technical exposition of our methods. As we believe that it may be applied to a wide range of problems, in the following we use generic terms to describe the data structure.

APPENDIX A. TECHNICAL EXPOSITION OF BIPOLAR CLASS ANALYSIS (BCA)

Bipolar Data Structure

Opinion polls aim to reflect a certain level of people’s agreement or disagreement about some statements pertaining, for example, to values or political issues. There exist several ways to represent the intensity of feeling in the answers to the questions posed in opinion polls, ranging from the classical *strongly agree–strongly disagree* scale to the thermometer one, usually measured on a 0 to 100 scale, where low numbers represent “cold”, i.e., disapproval, and high numbers represent “hot”, i.e., approval.

In this study, we have concentrated on questions involving the explicit or implicit existence of a *bipolar structure*. Let \mathcal{A} be the set of answers to a specific question in an opinion poll. For example, in an *in favor–against* question, $\mathcal{A} = \{\text{in favor, against}\}$ or, in a *strongly agree–strongly disagree* question,

$$\begin{aligned} \mathcal{A} = \{ & \text{strongly disagree, disagree, slightly disagree,} \\ & \text{neither agree nor disagree,} \\ & \text{slightly agree, agree, strongly agree}\}. \end{aligned}$$

To give structure to \mathcal{A} , we express it as a union $\mathcal{A} = \mathcal{N} \cup \mathcal{P}$ of two distinct non-empty subsets \mathcal{N} and \mathcal{P} , called, respectively, the *negative opinion semispace* and the *positive opinion semispace* of the question. We will assume that \mathcal{N} and \mathcal{P} are either disjoint or have exactly one element in common.

This restriction presents two scenarios. In the first, \mathcal{N} and \mathcal{P} have a unique common element, which we call the *neutrality element*, denoted by n . This element will represent a neutral answer to the given question. For example, a *strongly agree–strongly disagree* question has, as a possible answer, the option “neither agree nor disagree”, which acts as a neutral answer, serving as the neutrality element in \mathcal{A} . In this case, $\mathcal{P} = \{\text{neither agree nor disagree, slightly agree, agree, strongly agree}\}$ and $\mathcal{N} = \{\text{strongly disagree, disagree, slightly disagree, neither agree nor disagree}\}$. In the second scenario, \mathcal{N} and \mathcal{P} are disjoint, indicating no explicit neutrality element in \mathcal{A} . Nevertheless, we consider the implicit existence of a neutrality element n in \mathcal{A} , even though it does not arise as a possible answer to the question. This occurs, for example, in an *in favor–against* question, where $\mathcal{N} = \{\text{against}\}$ and $\mathcal{P} = \{\text{favor}\}$.

Remark A.1. While it is not strictly required for the opinion semispaces \mathcal{P} and \mathcal{N} of a given question to have an equal number of elements, we typically do not encounter such scenarios.¹³ It is worth noting, however, that our decomposition of the answer set \mathcal{A} accommodates this possibility. If a situation arises where \mathcal{P} and \mathcal{N} have a different number of elements, we will inform the reader accordingly; otherwise, we assume that both sets \mathcal{P} and \mathcal{N} contain the same number of answers.

The bipolar structure also applies when considering the set of responses to a poll of $Q \geq 1$ questions. Here, we have Q answer sets $\mathcal{A}_1, \dots, \mathcal{A}_Q$, and we may express the set \mathcal{A} of responses to the poll as the Cartesian product $\mathcal{A} = \mathcal{A}_1 \times \dots \times \mathcal{A}_Q$. We call an element in \mathcal{A} a *response* and \mathcal{A} the *response set*. Each individual answer set can be expressed as $\mathcal{A}_q = \mathcal{N}_q \cup \mathcal{P}_q$.

Relationality

Consider a family of $Q \geq 1$ questions with response set $\mathcal{A} = \mathcal{A}_1 \times \dots \times \mathcal{A}_Q$. A key insight in Goldberg’s definition of relationality (Goldberg 2011) is to compare answers to pairs of questions. To do so, Goldberg’s definition of relationality requires each answer set \mathcal{A}_q to be in a one-to-one correspondence with a strictly increasing finite sequence of equidistant numbers in the unit interval $[0, 1]$, starting with 0 and ending with 1. With this identification in place, we may assume that each response is a tuple of Q numbers between zero and one, allowing for the arithmetical manipulation of answers to different questions. Let $u = (u_1, \dots, u_Q)$ and $v = (v_1, \dots, v_Q)$ be responses in \mathcal{A} and define $u_{kl} = (u_k, u_l)$ and $v_{kl} = (v_k, v_l)$ for $1 \leq k < l \leq Q$. Goldberg’s relationality is a function $R_G: \mathcal{A} \times \mathcal{A} \rightarrow \mathbb{R}$ given by

$$(A.1) \quad R_G(u, v) = \frac{2}{Q(Q-1)} \sum_{k=1}^{Q-1} \sum_{l=k+1}^Q \lambda(u_{kl}, v_{kl}) \delta(u_{kl}, v_{kl}),$$

where

$$(A.2) \quad \lambda: (\mathcal{A}_k \times \mathcal{A}_l) \times (\mathcal{A}_k \times \mathcal{A}_l) \rightarrow \{-1, 1\}$$

$$(u_{kl}, v_{kl}) \mapsto \lambda(u_{kl}, v_{kl}) = \begin{cases} 1, & \text{if } (u_k - u_l)(v_k - v_l) \geq 0 \\ -1, & \text{otherwise,} \end{cases}$$

13. An exception is Question 254 in Eurobarometer 21, which asks people if they “completely agree”, “broadly agree”, “agree if anything”, “disagree to some extent”, or “disagree completely” on whether their country had an excess of migrant workers.

and

$$(A.3) \quad \delta: (\mathcal{A}_k \times \mathcal{A}_l) \times (\mathcal{A}_k \times \mathcal{A}_l) \rightarrow [0, 1]$$

$$(u_{kl}, v_{kl}) \mapsto \delta(u_{kl}, v_{kl}) = 1 - ||u_k - u_l| - |v_k - v_l||.$$

Although Goldberg does not explicitly mentions it, one may interpret the function λ as a “polarity function” and the function δ as a “magnitude” one. Our method’s construction is based on this interpretation of the component functions of R_G .

Polarity. We believe that our redefinition of relationality better captures the structure of the response set compared to Goldberg’s approach. The strength of our method lies in its focus on the polarity function in the above expression. For that reason, we will first define it.

The crux of our approach hinges on whether the same respondent switches between opinion semispaces when answering different questions. For instance, consider a simple poll with two questions, both of the *strongly agree-strongly disagree* type. If a respondent answers “agree” to the first question and “strongly disagree” to the second one, they are situated in the positive opinion semispace for the first question and the negative opinion semispace for the second. Thus, we say that this respondent “crosses the neutrality line” between questions 1 and 2.

Let $u = (u_1, \dots, u_Q)$ be a response in \mathcal{A} . We encounter four distinct scenarios regarding displacement between the opinion semispaces when considering pairs of answers $u_{kl} = (u_k, u_l)$ for $1 \leq k, l \leq Q$ and $k \neq l$. We formalize these scenarios through the introduction of four *movement functions* as follows:

$$(A.4) \quad M_{++}(u_{kl}) = \begin{cases} 1, & \text{if } u_k \in \mathcal{P}_k, u_l \in \mathcal{P}_l \\ 0, & \text{otherwise;} \end{cases}$$

$$(A.5) \quad M_{--}(u_{kl}) = \begin{cases} 1, & \text{if } u_k \in \mathcal{N}_k, u_l \in \mathcal{N}_l \\ 0, & \text{otherwise;} \end{cases}$$

$$(A.6) \quad M_{-+}(u_{kl}) = \begin{cases} 1, & \text{if } u_k \in \mathcal{N}_k^\circ, u_l \in \mathcal{P}_l^\circ \\ 0, & \text{otherwise;} \end{cases}$$

$$(A.7) \quad M_{+-}(u_{kl}) = \begin{cases} 1, & \text{if } u_k \in \mathcal{P}_k^\circ, u_l \in \mathcal{N}_l^\circ \\ 0, & \text{otherwise.} \end{cases}$$

Here, \mathcal{N}° consists of all the elements in \mathcal{N} except for the neutrality one, i.e., $\mathcal{N}^\circ = \mathcal{N} \cap \mathcal{P}^c$, where \mathcal{P}^c denotes the complement of \mathcal{P} . Similarly, we let $\mathcal{P}^\circ = \mathcal{P} \cap \mathcal{N}^c$. We call \mathcal{N}° and \mathcal{P}° the *strictly negative* and *strictly positive* opinion semispaces, respectively.

Intuitively, these movement functions serve as indicators for the four potential displacements a pair of answers could exhibit. Additionally, it is worth noting that at least one of the functions listed above must take the value one for any u_{kl} . Furthermore, with the exception of the case $u_{kl} = (n_k, n_l)$, exactly one of the functions will be non-zero.

For clarity in defining the polarity function, we will establish *product movement functions* between pairs of distinct responses. Let (u_1, \dots, u_Q) and (v_1, \dots, v_Q) be two responses in \mathcal{A} and, as before, let $u_{kl} = (u_k, u_l)$ and $v_{kl} = (v_k, v_l)$ be pairs of answers in u and v , respectively, with $k \neq l$. For ease of notation, we will write $n_{kl} = (n_k, n_l)$, where n_i is the neutrality element in \mathcal{A}_i . The *semispace shift function* $f_{A,B}$ assigns to the pair (u_{kl}, v_{kl}) the value 0 or 1 and is given by

$$(A.8) \quad \begin{aligned} f_{A,B}: (\mathcal{A}_k \times \mathcal{A}_l) \times (\mathcal{A}_k \times \mathcal{A}_l) &\rightarrow \{0, 1\} \\ (u_{kl}, v_{kl}) &\mapsto M_A(u_{kl})M_B(v_{kl}), \end{aligned}$$

where $A, B \in \{++, --, -+, +-\}$. Now, we define the *polarity function* ω , which assigns to the pair (u_{kl}, v_{kl}) the values 0, -1, or 1, as follows. First, if $u_{kl}, v_{kl} \neq n_{kl}$, we let

$$(A.9) \quad \omega(u_{kl}, v_{kl}) = \begin{cases} 1, & \text{if } f_{A,A}(u_{kl}, v_{kl}) = 1 \text{ for at least one } A; \\ -1, & \text{if } f_{A,A^*}(u_{kl}, v_{kl}) = 1 \text{ for at least one } A; \\ 0, & \text{otherwise.} \end{cases}$$

Here, $*$ performs the following transformations

$$(A.10) \quad (++)^* = --, (--)^* = ++, (-+)^* = +- , \text{ and } (+-)^* = -+ .$$

To define ω for the three special cases in the presence of neutrality, we let

$$(A.11) \quad \omega(n_{kl}, v_{kl}) = \begin{cases} 1, & \text{if } M_{++}(v_{kl}) = 1; \text{ or } M_{--}(v_{kl}) = 1; \\ 0; & \text{otherwise;} \end{cases}$$

$$(A.12) \quad \omega(u_{kl}, n_{kl}) = \begin{cases} 1, & \text{if } M_{++}(u_{kl}) = 1 \text{ or } M_{--}(u_{kl}) = 1; \\ 0; & \text{otherwise;} \end{cases}$$

and

$$(A.13) \quad \omega(n_{kl}, n_{kl}) = 1.$$

Magnitude. As discussed in Section 2, the selection of the magnitude function in the definition of relationality presents a delicate decision point in the methodology. One may justifiably assert that no universal magnitude function can be applied for defining relationality due to disparities in the number of possible answers for different questions, as well as variations in scales among them, among other considerations. Nonetheless, researchers must make a selection in practice to conduct analyses. We contend that the methodological challenge of incorporating the “distance” between non-cardinal opinion structures warrants further investigation in subsequent studies.

In this article, we have opted for two different magnitudes. The first, denoted as $\mu_1(u_{kl}, v_{k,l})$, is a constant magnitude, defined as

$$(A.14) \quad \mu_1(u_{kl}, v_{kl}) = 1, \text{ for every } u_{kl}, v_{kl} \in \mathcal{A}_k \times \mathcal{A}_l.$$

One may see μ_1 as the simplest magnitude we can endow $(\mathcal{A}_k \times \mathcal{A}_l) \times (\mathcal{A}_k \times \mathcal{A}_l)$ with, allowing us to focus on studying relationality between responses solely in terms of polarity. One of the principal advantages of μ_1 is that it enables us to compute our relationality solely in terms of the bipolar structure, which is one of the key components of our method.

The second magnitude arises from studying the concrete structure of the set of answers to a given opinion poll. Before defining this magnitude function, let us provide more insight into the construction of the opinion semispaces of a given question. First, we observe that in our framework, each answer set \mathcal{A}_q to an individual question is a *totally ordered set*. That is, there exists a binary relation \leq_q on \mathcal{A}_q satisfying the following conditions:

- *Reflexivity:* $a \leq_q a$ for all $a \in \mathcal{A}_q$.
- *Transitivity:* Given $a, b, c \in \mathcal{A}_q$, if $a \leq_q b$ and $b \leq_q c$, then $a \leq_q c$.
- *Anti-symmetry:* Given $a, b \in \mathcal{A}_q$ with $a \neq b$, if $a \leq_q b$, then $b \not\leq_q a$.
- *Strong connectedness:* For any $a, b \in \mathcal{A}_q$, either $a \leq_q b$ or $b \leq_q a$.

We call \leq_q a *total order* on \mathcal{A}_q .

It is straightforward to verify that, for example, the answer set of questions of *strongly agree–strongly disagree* supports a natural total order given by

$$(A.15) \quad \text{strongly disagree} \leq \text{disagree} \leq \text{slightly disagree} \leq \text{neither agree nor disagree} \\ \leq \text{slightly agree} \leq \text{agree} \leq \text{strongly agree}.$$

Using the total order, we may write an answer set \mathcal{A}_q with explicit neutrality element n_q as the union $\mathcal{A}_q = \mathcal{P}_q \cup \mathcal{N}_q$ with

$$(A.16) \quad \mathcal{P}_q = \{a \in \mathcal{A}_q : n_q \leq a\}$$

and

$$(A.17) \quad \mathcal{N}_q = \{a \in \mathcal{A}_q : a \leq n_q\}.$$

We define μ_2 following the above structure of the answer sets \mathcal{A}_q and the symmetry induced by the corresponding total order \leq_q . While the polarity function ω aims to capture whether answers move between opinion semispaces, μ_2 aims to measure the variation in the relative position with respect to the neutrality element in pairs of answers from two different responses and attempts to homogenize the disparity in the number of options between different spaces of answers.

To define μ_2 , we first consider a response $u = (u_1, \dots, u_Q)$ in \mathcal{A} and define the *variation* $\Delta(u_{kl})$ of the pair of answers $u_{kl} = (u_k, u_l)$ by

$$(A.18) \quad \Delta(u_{kl}) = |\text{dist}_k(u_k, n_k) - \text{dist}_l(u_l, n_l)|,$$

where $\text{dist}_i(\cdot, \cdot)$ denotes a metric function that measures the distance from answer u_i to the neutrality element n_i in the answer space A_i . The value $\Delta(u_{kl})$ measures the magnitude of the change in distances to the neutrality element changes from answer k to answer l . Note that $\Delta(u_{kl}) = \Delta(u_{lk})$. Now, let $v = (v_1, \dots, v_Q)$ be another response in \mathcal{A} and define

$$(A.19) \quad \mu_2(u_{kl}, v_{kl}) = 1 - \frac{|\Delta(u_{kl}) - \Delta(v_{kl})|}{\max_q D_q}.$$

Here, $D_q = \max\{\text{dist}(a, n_q) : a \in \mathcal{A}_q\}$, the maximum distance to the neutrality element in the answer set \mathcal{A}_q , and we maximize between all answer sets. This term serves as a way to normalize the distance between the variation of pairs of answers as well as a way to ponderate questions that have a large number of answers between neutrality and the extremes in \mathcal{A}_q .

In this article, we have chosen the same distance for every question, letting

$$(A.20) \quad \text{dist}_q(a, b) = \begin{cases} \#\{c \in \mathcal{A}_q : b \leq_q c \leq_q a \text{ and } c \neq b\} & \text{if } a \geq_q b; \\ \#\{c \in \mathcal{A}_q : a \leq_q c \leq_q b \text{ and } c \neq b\} & \text{if } a \leq_q b \end{cases}$$

for any $a, b \in \mathcal{A}_q$. Here, $\#$ denotes the number of elements in the set. Thus, $\text{dist}_q(a, n_q)$ measures the relative position of $a \in \mathcal{A}_q$ with respect to the neutrality element n_q .

Let us illustrate the use of this distance and how to compute μ_2 with an example. Suppose we have a poll with two questions with answer sets \mathcal{A}_1 and \mathcal{A}_2 , respectively. Suppose the first question is of type *strongly agree–strongly disagree* with five possible answers totally ordered by

$$(A.21) \quad \textit{strongly disagree} \leq \textit{disagree} \leq \textit{neither agree nor disagree} \leq \textit{disagree} \leq \textit{strongly disagree}.$$

The second one is a question of type *agree–disagree* with three possible answers with total order given by

$$(A.22) \quad \textit{disagree} \leq \textit{neither agree nor disagree} \leq \textit{agree}.$$

The possible values for the distance to the neutrality element in the first answer set are $\text{dist}_1(a, n_1) = 0, 1, \text{ or } 2$, and we have $D_1 = 2$. The distance to the neutrality element in the second answer set takes the values $\text{dist}_2(a, n_2) = 0 \text{ or } 1$ and, hence, $D_2 = 1$. Therefore, $\max_q D_q = 2$. Note that μ_2 has equalized, in terms of distance, the answers *agree* and *disagree* to both questions in terms of their distance to neutrality.

Consider now two responses $u = (\textit{totally agree}, \textit{neither agree nor disagree})$ and $v = (\textit{disagree}, \textit{disagree})$. Then

$$(A.23) \quad \begin{aligned} \Delta(u_{12}) &= |\text{dist}_1(\textit{totally agree}, \textit{neither agree nor disagree}) \\ &\quad - \text{dist}_2(\textit{neither agree nor disagree}, \textit{neither agree nor disagree})| \\ &= |2 - 0| = 2, \\ \Delta(v_{12}) &= |\text{dist}_1(\textit{disagree}, \textit{neither agree nor disagree}) \\ &\quad - \text{dist}_2(\textit{disagree}, \textit{neither agree nor disagree})| \\ &= |1 - 1| = 0. \end{aligned}$$

Thus,

$$(A.24) \quad \mu_2(u, v) = 1 - \frac{|\Delta(u_{12}) - \Delta(v_{12})|}{\max_i D_i} = 1 - \frac{|2 - 0|}{2} = 0.$$

As previously mentioned, one may use other functions to assign magnitude. Note that one may also consider magnitudes that are not necessarily induced by distances in an answer set, as in (A.18) one only needs to assign values to the pairs (a, n_q) with $a \in \mathcal{A}_q$. We have used μ_1 and μ_2 due to the nature of our experiments (μ_2 remains in the spirit of the magnitude used by Baldassarri and Goldberg 2014) and in order to exploit the structure of our questions (μ_1 is based on the bipolar

structure). In other studies, other magnitudes may be defined to address the specific needs of the problems under consideration.

Summarizing, the relationality we have defined is given by

$$(A.25) \quad R(u, v) = \frac{2}{Q(Q-1)} \sum_{k=1}^{Q-1} \sum_{l=k+1}^Q \omega(u_{kl}, v_{kl}) \mu_j(u_{kl}, v_{kl}), \quad j = 1, 2$$

where ω is the polarity function defined in (A.9) adding its three special cases involving neutrality and μ_1 and μ_2 denote the magnitudes defined in (A.14) and (A.19).

APPENDIX B. SIMULATION MODEL

Simulating data with construals. The synthetic datasets consist of matrices $(x_{iq})_{i=1, q=1}^{N, Q}$, where N is the number of respondents and Q is the number of questions in the poll. Here, we present a precise discussion of how the data is generated.

Latent-variable model. In opinion polls, each respondent is asked to choose among a finite number of options. These number of options represent the answer space $\mathcal{A}_q = \{a_1, \dots, a_{H_q}\}$, for $q = 1, \dots, Q$. We allow the number of options to vary across questions. As we have previously discussed, the options in \mathcal{A}_q are totally ordered (see Appendix A). Hence, without loss of generality, we consider that $a_1 \leq_q a_2 \leq_q \dots \leq_q a_{H_q}$.

To choose among these ordered options, we consider that respondent i follows a latent-variable model. That is, we assume that for each respondent i and question q , there is a latent variable $x_{iq}^* \in \mathbb{R}$ that represents the accurate position of i about issue q . The respondent then partitions \mathbb{R} into H_q intervals and chooses their answer $x_{iq} \in \mathcal{A}_q$ depending on the interval where x_{iq}^* falls (for a detailed introduction to the latent-variable model for ordered choice, see Greene and Hensher 2010).

Formally, for each respondent i and question q there are $H_q - 1$ thresholds $(\tau_{iq\eta})_{\eta=1}^{H_q-1}$, with $\tau_{iq1} < \tau_{iq2} < \dots < \tau_{iq, H_q-1}$ that partition \mathbb{R} into H_q intervals. The respondent chooses x_{iq} depending on the interval where x_{iq}^* falls:

$$(B.1) \quad x_{iq} = a_\eta \iff x_{iq}^* \in (\tau_{iq, \eta-1}, \tau_{iq\eta}) \text{ for } \eta = 1, \dots, H_q.$$

In the above equation, we set $\tau_{iq0} = -\infty$ and $\tau_{iq, H_q} = \infty$ for every i and q . Also, since we will consider absolutely continuous distributions for both the latent variables and the thresholds, we note that $P(x_{iq}^* = \tau_{iq\eta}) = 0$. Thus, the fact that we consider open intervals is unimportant.

Construals and copulas. We model construals as a dependence structure on the vector of latent variables $X_i^* = (x_{i1}^*, \dots, x_{iQ}^*)$. That is, the mathematical equivalent to the sentence “respondent i belongs to a certain construal” is “the variables in X_i^* , which give the accurate position of i on each issue, follow a certain joint distribution”. We believe that this model captures the complexity of construals.

To model dependence on the latent variables in X_i^* , we introduce the notion of copulas. Copulas are “a way of studying scale-free measures of dependence” Fisher 1997. A copula is a joint distribution on the unit cube $[0, 1]^Q$ with uniform marginal distributions. Since the marginals are uniform, the copula captures scale-free information about the dependence structure of random vectors. We

refer to Nelsen 2006 for a general introduction to copulas. A relevant result is *Sklar's Theorem* (Th. 2.10.9 in Nelsen 2006). Suppose that X_i^* has absolutely continuous distribution F and let F_q denote the marginal distribution of each variable x_{iq}^* , $q = 1, \dots, Q$. Then there exists a unique copula C such that

$$(B.2) \quad F(x_1, \dots, x_Q) = C(F_1(x_1), \dots, F_Q(x_Q)).$$

This means that we can decompose the joint distribution of X_i^* between (i) its marginal distributions F_q , which model the variability within issue q , and (ii) the copula C , which models the dependence structure on the joint distribution.

We identify a construal with a copula on $[0, 1]^Q$. The space of all copulas is rich enough to capture the many ways in which Q issues may be related in the given construal. For instance, the case where there is no constraint between the issues corresponds to the independence copula $C(u_1, \dots, u_Q) = u_1 \cdot \dots \cdot u_Q$. A popular family, which is suitable for simulations, is the family of Gaussian copulas. Let Φ be the cumulative distribution function (cdf) of a standard normal variable and Φ_Σ be the cdf of a multivariate normal vector with mean zero and covariance matrix Σ . Fix $\Sigma_{qq} = 1$, $q = 1, \dots, Q$, so that the marginal distributions have unit variance. A Gaussian copula with correlation matrix Σ is defined by

$$(B.3) \quad C(u_1, \dots, u_Q; \Sigma) = \Phi_\Sigma(\Phi^{-1}(u_1), \dots, \Phi^{-1}(u_Q)).$$

The family of Gaussian copulas is indexed by the correlation matrices Σ . They generalize the Gaussian-like dependence structure to allow for arbitrary (i.e., non-Gaussian) marginal distributions. Note that when Σ is the identity matrix, we recover the independence copula. We refer to Appendix E for a detailed discussion about the Gaussian family of copulas.

To sum up, say there are K construals on a group of respondents. Each of these construal is modelled by a copula C^k , $k = 1, \dots, K$. If respondent i belongs to construal k , then its latent variable X_i^* follows the distribution F^k given by

$$(B.4) \quad F^k(x_1, \dots, x_Q) = C^k(F_1(x_1), \dots, F_Q(x_Q)),$$

where the F_q are the marginal distributions of each question among all respondents. These marginal distributions are shared by all respondents, independently of the construal they belong to. We note that these distributions determine the consensus that each of the options of question q raises among the group. The distribution F_q characterizes whether respondents tend to agree or disagree about issue q .

In what follows, it is better to “normalize” the accurate position that respondent i has, translating it to the unit cube. Define $u_{iq}^* = F_q^{-1}(x_{iq}^*)$ and $u_i^* = (u_{i1}^*, \dots, u_{iQ}^*)$. If respondent i belongs to construal k , then u_i^* has distribution C^k . That is, u_{iq}^* is uniformly distributed for each question q . An example will help clarify what u_{iq}^* means. Say that the options in question q are ordered from $a_1 = \text{Absolute disagreement}$ to $a_{H_q} = \text{Absolute agreement}$. Then, since u_{iq}^* is uniformly distributed, $u_{iq}^* = 0.7$ means that 70% of the population agrees with the statement of question q less than respondent i . This is what we mean by accurate position of the respondent. The fact that we normalize $u_{iq}^* = F_q^{-1}(x_{iq}^*)$ makes the position of respondents easier to interpret.

We can also “normalize” the thresholds, so we directly work with u_i^* when generating the data. Let $\bar{\tau}_{iq\eta} = F_q^{-1}(\tau_{iq\eta})$. Then, according to equation (B.1), respondent i chooses x_{iq} with the following rule:

$$(B.5) \quad x_{iq} = a_\eta \iff u_{iq}^* \in (\bar{\tau}_{iq,\eta-1}, \bar{\tau}_{iq\eta}) \text{ for } \eta = 1, \dots, H_q.$$

Thresholds and neutrality. As we can see in equation (B.5), respondent i first thinks about her accurate position u_{iq}^* and then decides among the options in \mathcal{A}_q depending on whether u_{iq}^* exceeds certain thresholds. The position u_i^* comes from the distribution C^k and is, therefore, subject to the constraints present in the construal. The random thresholds $(\bar{\tau}_{iq\eta})_{\eta=1}^{H_q}$, in turn, model individual variation (idiosyncratic shocks). These capture individual-specific opinions that originate due to events, readings, family ties, or such experiences. They may induce respondent i not to adhere to the logic prescribed by construal k . Indeed, if the variance of the distribution of the thresholds is high, we will observe loose adherence to each construal.

Appendix A has argued the role of neutrality in opinion polls, even if this is implicitly defined. We implement neutrality and symmetric restrictions to our data-generating process in the procedure to generate thresholds. We consider that, even if it is not an explicit option, there exists a neutral position $\bar{n}_q \in [0, 1]$ for each question. The neutral position models the consensus of the group regarding questions q .

We think of the individual thresholds as departing away from the neutral position. Consider that H_q is odd and that neutrality is explicit, as is the case frequently in opinion pools. We draw $\bar{H}_q = (H_q - 1)/2$ departure distances: $(\omega_{iq\eta})_{\eta=1}^{\bar{H}_q}$. For instance, when $H_q = 5$ we draw 2 distances $(\omega_{iq1}, \omega_{iq2})$. The first distance ω_{iq1} determines the length of the neutrality interval. That is, we set the interval that determines whether $x_{iq} = a_{\bar{H}_q} = n_q$ (neutrality) to

$$(B.6) \quad (\bar{\tau}_{iq\bar{H}_q}, \bar{\tau}_{iq,\bar{H}_q+1}) = (\bar{n}_q - \omega_{iq1}, \bar{n}_q + \omega_{iq1}).$$

The next distances $(\omega_{iq\eta})_{\eta=2}^{\bar{H}_q}$ measure the length of the within threshold intervals. We can recursively define the thresholds that diverge towards 1 by

$$(B.7) \quad \bar{\tau}_{iq, \bar{H}_q + \eta} = \bar{\tau}_{iq, \bar{H}_q + \eta - 1} + \omega_{iq\eta} \text{ for } \eta = 2, \dots, \bar{H}_q.$$

Then, by symmetry with respect to the neutral position, the thresholds that diverge towards 0 are:

$$(B.8) \quad \bar{\tau}_{iq, 1 + \bar{H}_q - \eta} = \bar{\tau}_{iq, 2 + \bar{H}_q - \eta} - \omega_{iq\eta} \text{ for } \eta = 2, \dots, \bar{H}_q.$$

Distances $(\omega_{iq\eta})_{\eta=1}^{\bar{H}_q}$ can be drawn from arbitrary distributions, as long as one guarantees that $0 < \bar{\tau}_{iq1} < \bar{\tau}_{iq2} < \dots < \bar{\tau}_{iq, H_q - 1} < 1$. It is worth highlighting two extreme cases. If the $\omega_{iq\eta}$'s tend to be small, we may say that question q is *divisive*: there will not be many respondents choosing $x_{iq} = n_q$ since the interval $(\bar{n}_q - \omega_{iq1}, \bar{n}_q + \omega_{iq1})$ is short. For divisive questions, respondents tend to choose the extreme positions a_1 or a_{H_q} . On the contrary, if the $\omega_{iq\eta}$ are large, the question is *unimportant*: most respondents will choose the neutrality element n_q , reflecting that there is no cleavage regarding that issue.

Simulation algorithm

We implement a large volume of experiments to test how BCA performs relative to other methods in the literature. To describe them, we first introduce a general algorithm to build synthetic data. The inputs to the algorithm, which vary across experiments (see Section B), are the following:

Set with number of construals	\mathcal{K}	Interval for neutral position	\mathcal{N}
The number of questions	Q	Threshold distances	Fixed, Random question-level, or
Set with number of options	\mathcal{H}		Random individual-level

TABLE 3. Inputs to the data generation algorithm.

The algorithm then proceeds as follows. First, we generate the matrix with **latent accurate positions**:

- (1) Draw the number of construals K from \mathcal{K} .
- (2) Set the number of questions to Q .
- (3) For each construal k , build the construal correlation matrix Σ^k (see below). The copula C^k defining construal k is the Gaussian copula with correlation matrix Σ^k .
- (4) For each construal k , randomly generate the number of respondents adhering to it: $N_k \in \{100, 101, \dots, 499, 500\}$. All cases are equally probable. The total number of respondents is then $N = \sum_{k=1}^K N_k$.

- (5) For each respondent i adhering to construal k , generate the latent accurate position u_i^* following distribution C^k . Precisely, we generate a random draw $(\xi_{iq})_{q=1}^Q$ from a multivariate normal with mean zero and covariance matrix Σ^k . Then, we set $u_{iq}^* = \Phi^{-1}(\xi_{iq})$ for $q = 1, \dots, Q$.

Then, we generate the **synthetic data**:

- (5) For each question q , draw the number of options H_q from \mathcal{H} . The answer set is

$$(B.9) \quad \mathcal{A}_q = \{-(H_q - 1)/2, \dots, -1, 0, 1, \dots, (H_q - 1)/2\}.$$

- (6) For each question q , draw neutral position \bar{n}_q from \mathcal{N} .

- (7) If thresholds vary at individual level, do nothing.

- (8) For each respondent i and question q :

- (a) To generate thresholds, we compute $\bar{H}_q = (H_q - 1)/2$ distances: $(\omega_{iq\eta})_{\eta=1}^{\bar{H}_q}$. How these are obtained varies depending on whether they are set to be random (see below). Once we have the distances, we get the thresholds using equations (B.6), (B.7), and (B.8).
- (b) We set answer x_{iq} according to equation (B.5).

Note that if thresholds do not vary at the individual level, step (7) is replaced by step (8.a), and we only perform step (b) inside point (8). That is, thresholds are generated for each question q , so that $\bar{\tau}_{iq\eta} = \bar{\tau}_{q\eta}$ for every respondent i .

Regarding the **construction of the construal correlation matrix** Σ^k , one needs to obtain *positive definite* random matrices. To do so, we generate a $Q(Q-1)/2$ vector of partial correlations, where each element of the vector is drawn from $2 \cdot \text{Beta}(\alpha_k, \alpha_k) - 1$ (to ensure it is between -1 and 1). The parameter α_k follows an exponential distribution with mean 0.8 . This parameter is used to model how stringent each construal is: large α_k for a lax construal versus small α_k for a stringent construal. We then follow the algorithm in Lewandowski, Kurowicka, and Joe 2009 to build Σ^k from the vector of partial correlations.

The algorithm for **generating fixed (non-random) inter-threshold distances** is the following. Consider a question q with H_q number of options and neutral position \bar{n}_q . Then, to ensure that thresholds are in the interval $(0, 1)$, the largest a distance can be is: $\ell_q = \min\{\bar{n}_q, 1 - \bar{n}_q\}$. We set $\omega_{q1} = \ell_q/H_q$ and $\omega_{q\eta} = 2\ell_q/H_q$ for every $\eta = 2, \dots, \bar{H}_q$. Note that, since thresholds are non-random, all individuals have $\omega_{iq\eta} = \omega_{q\eta}$. That is, thresholds only vary at question-level (if \bar{n}_q and H_q are different). To give a relevant example, consider that the neutral position is centered: $\bar{n}_q = 1/2$. Then, the algorithm partitions $(0, 1)$ in equally sized intervals. In that case $\ell_q = 1/2$, so

that $\omega_{q1} = 1/(2H_q)$ and $\omega_{q\eta} = 1/H_q$ for $\eta \geq 2$. Then, according to equation (B.6):

$$(B.10) \quad \bar{\tau}_{iq, \bar{H}_q} = \frac{1}{2} - \frac{1}{2H_q} \text{ and } \bar{\tau}_{iq, \bar{H}_q+1} = \frac{1}{2} + \frac{1}{2H_q} \text{ for every individual } i.$$

Therefore, $\bar{\tau}_{iq, \bar{H}_q+1} - \bar{\tau}_{iq, \bar{H}_q} = 1/H_q$. The distance between other thresholds is also $1/H_q$ by equations (B.7) and (B.8).

The algorithm for **generating random individual-level inter-threshold distances** is the following. Consider a question q with H_q number of options and neutral position \bar{n}_q . Let $\ell_{q1} = \min\{\bar{n}_q, 1 - \bar{n}_q\}$. We draw a random variable Z_i from a truncated normal distribution with parameters $(\mu, \sigma, a, b) = (1/H_q, 0.05, 0, 1)$.¹⁴ We set $\omega_{iq1} = Z_i \cdot \ell_{q1}$. Then, we define $\ell_{iq2} = \ell_{q1} - \omega_{iq1}$ and draw ω_{iq2} from a truncated normal with parameters $(\mu, \sigma, a, b) = (\ell_{q1}/\bar{H}_q, 0.05, 0, \ell_{iq2})$. We then proceed recursively for $\eta = 3, \dots, \bar{H}_q$: define $\ell_{iq\eta} = \ell_{iq, \eta-1} - \omega_{iq, \eta-1}$ and draw $\omega_{iq\eta}$ from a truncated normal with parameters $(\mu, \sigma, a, b) = (\ell_{q1}/\bar{H}_q, 0.05, 0, \ell_{iq\eta})$.

Regarding the algorithm to **generate random question-level inter-threshold distances**, it parallels the one described above. The only difference is that random variable drawing is done at the question-level, and then all individuals are assigned the same thresholds: $\omega_{iq\eta} = \omega_{q\eta}$. That is, Z_q is drawn for each question and $\omega_{q1} = Z_q \cdot \ell_{q1}$. We then recursively set $\ell_{q\eta} = \ell_{q, \eta-1} - \omega_{q, \eta-1}$ for $\eta = 2, \dots, \bar{H}_q$.

Description of the experiments

Our experiments can be divided into two families. In one, we consider random underlying correlation matrices for each construal. In the other, we simulate data with two construals and fixed correlation matrices. Data in the later family of experiments resembles a plausible scenario when it comes to the organization of political views (in a simplified version).

Table 4 describes the experiments in the first family. Each row represents a different experiment. In columns, we specify the inputs to the data generation algorithm (see Table 3 for a description).

	$\mathcal{K} =$	$Q =$	$\mathcal{H} =$	$\mathcal{N} =$	Threshold distances
E1.1: All fixed 5	{2, 3, 4}	10	{5}	[0.5]	Fixed
E1.2: All fixed 7	{2, 3, 4}	10	{7}	[0.5]	Fixed
E1.3: Random options	{2, 3, 4}	10	{3, 5, 7}	[0.5]	Fixed
E1.4: Random distances	{2, 3, 4}	10	{5}	[0.5]	Random question-level
E1.5: Asymmetric neutrality	{2, 3, 4}	10	{5}	(0.35, 0.65)	Fixed
E1.6: All Random	{2, 3, 4}	10	{3, 5, 7}	(0.35, 0.65)	Random individual-level

TABLE 4. Description of the experiments.

14. If Z_i has a truncated normal distribution with parameters (μ, σ, a, b) , then $Z_i = W_i | a \leq W_i \leq b$ for a random variable $W_i \sim N(\mu, \sigma)$. This means that the support of Z is the interval $[a, b]$.

In the second family of experiments, we generate data with two underlying construals. The number of questions is three (these can be thought of as summaries or indices of three issues). The first and second questions are perfectly co-linear for both construals: individuals have the same latent precise position for them. The two construals differ in the relationship between these two issues and the third: in construal 1 they are positively related, while in construal 2 the relationship is negative. The correlation structure for both construals is:

$$(B.11) \quad \Sigma^1 = \begin{pmatrix} 1 & 1 & 0.7 \\ 1 & 1 & 0.7 \\ 0.7 & 0.7 & 1 \end{pmatrix} \text{ and } \Sigma^2 = \begin{pmatrix} 1 & 1 & -0.7 \\ 1 & 1 & -0.7 \\ -0.7 & -0.7 & 1 \end{pmatrix}.$$

These correlation structures are present across all experiments of the family. Table 5 provides further details about the experiments.

	$\mathcal{K} =$	$Q =$	$\mathcal{H} =$	$\mathcal{N} =$	Threshold distances
E2.1: All fixed 7	{2}	3	{7}	[0.5]	Fixed
E2.2: Varying thresholds	{2}	3	{5}	(0.35, 0.65)	Random question-level
E2.3: Individual thresh.	{2}	3	{5}	(0.35, 0.65)	Random individual-level

TABLE 5. Description of the experiments.

APPENDIX C. CORRELATION DISSIMILARITY

Our simulations consider that the issue constraint for each construal k is given by a Gaussian copula with correlation matrix Σ^k . It is then natural to ask whether each method is able to recover the correlation structure $(\Sigma^k)_{k=1}^K$ from the data. There are two problems to address when answering this question: (i) that the number of construals estimated by the method may differ from the true number of construals and (ii) that the labelling of each construal is arbitrary (i.e., what a certain method labels as “the first construal” may differ from the group that was labelled as “the first construal” when generating the data).

We propose a method to measure how well each method estimates the underlying correlation matrices, which overcomes these issues. The inputs to the method are a collection of “true” and estimated correlation matrices, $(\Sigma^k)_{k=1}^K$ and $(\hat{\Sigma}^k)_{k=1}^{\hat{K}}$, respectively. The number of “true” (K) and estimated (\hat{K}) construals may differ. Additionally, the researcher must provide a measure of dissimilarity between two matrices, which we denote by d . We base our assessment of the methods on the Frobenius distance

$$(C.1) \quad d(A, B) = \sqrt{\text{trace}(A - B)(A - B)'},$$

where A and B are matrices and A' denotes the transpose of A .¹⁵ For more information about the Frobenius distance, see (Cohen 2022).

As we have discussed, our measure of performance must deal with the fact that there is no automatic way to pair an estimated construal with a “true” construal. That is, for a given estimated construal $\hat{k} \in \{1, \dots, \hat{K}\}$ we do not know the corresponding true construal $k \in \{1, \dots, K\}$. Thus, we cannot directly compute $d(\Sigma^k, \hat{\Sigma}^{\hat{k}})$. The problem is exacerbated when the number of estimated construals differs from the number of true construals. To deal with these inconveniences, we will work with all possible pairings between $\{1, \dots, \hat{K}\}$ and $\{1, \dots, K\}$. Our measure of performance will then be the minimum of matrix dissimilarity across possible pairings.

We define a *paring* between $\{1, \dots, \hat{K}\}$ and $\{1, \dots, K\}$ as an injective function $p: \{1, \dots, \hat{K}\} \rightarrow \{1, \dots, K\}$. This function assigns a unique “true” construal $k = p(\hat{k})$ to each estimated construal \hat{k} . Note, however, that no injective function exists when $\hat{K} > K$, that is, when the method overestimates the number of construals. To deal with this, we propose to augment the “true” number of construals by introducing $\hat{K} - K$ artificial construals. The correlation structure of these

¹⁵ The choice of the Frobenius distance is not particularly relevant. Indeed, in case the dissimilarity measure comes from a norm, the ranking of methods will not depend on the chosen norm. This comes from all norms being equivalent.

artificial construals is $\Sigma^k = I_Q$, the Q -dimensional identity matrix. That is, issues for these artificial construals are not constrained, indicating that the construals are not really present. So, when $\hat{K} > K$, we propose to compute the dissimilarity between the collection of estimated correlation matrices $(\hat{\Sigma}^k)_{k=1}^{\hat{K}}$ and the augmented collection of \hat{K} “true” correlation matrices $(\Sigma^1, \dots, \Sigma^K, I_Q, \dots, I_Q)$.¹⁶

To introduce the measure of dissimilarity between $(\hat{\Sigma}^k)_{k=1}^{\hat{K}}$ and $(\Sigma^k)_{k=1}^K$, consider first that $\hat{K} \leq K$. Denote by P the set of all possible pairings between $\{1, \dots, \hat{K}\}$ and $\{1, \dots, K\}$. Given a dissimilarity measure between pairs of matrices d , the dissimilarity between collections, denoted \mathcal{D} , is

$$(C.2) \quad \mathcal{D} \left((\hat{\Sigma}^k)_{k=1}^{\hat{K}}, (\Sigma^k)_{k=1}^K \right) := \min_{p \in P} \left\{ \max_{\hat{k}=1, \dots, \hat{K}} d \left(\hat{\Sigma}^{\hat{k}}, \Sigma^{p(\hat{k})} \right) \right\}, \text{ when } \hat{K} \leq K.$$

As we have discussed, when $\hat{K} > K$, we augment the collection of “true” construals by adding construals whose correlation structure is the identity. That is, we define the collection $(\tilde{\Sigma}^k)_{k=1}^{\hat{K}}$, with $\tilde{\Sigma}^k = \Sigma^k$, for $k \leq K$, and $\tilde{\Sigma}^k = I_Q$, for $k = K + 1, \dots, \hat{K}$. In this case, the dissimilarity between the collections is given by

$$(C.3) \quad \mathcal{D} \left((\hat{\Sigma}^k)_{k=1}^{\hat{K}}, (\Sigma^k)_{k=1}^K \right) := \mathcal{D} \left((\hat{\Sigma}^k)_{k=1}^{\hat{K}}, (\tilde{\Sigma}^k)_{k=1}^{\hat{K}} \right), \text{ when } \hat{K} > K.$$

Remark C.1. An alternative to augmenting the number of “true” construals when $\hat{K} > K$ is to consider, in that case, pairings as injective functions $p: \{1, \dots, K\} \rightarrow \{1, \dots, \hat{K}\}$. This leaves, however, some estimated construals unpaired with “true” construals. Thus, we believe that this other approach unfairly benefits methods that overestimate the number of construals. Indeed, a method can trick this alternative measure by estimating a lot of construals, even if some of those are very dissimilar to the “true” construals. This is so since the minimum in equation (C.2) will be achieved when the dissimilar estimated construals are left unpaired.

A Benchmark to asses method performance

One of the problems of the above measure of dissimilarity is that there is no clear notion of whether a method estimated properly the underlying correlation structure. Certainly, when \mathcal{D} is zero, the estimation is perfect and larger values of \mathcal{D} indicate a worse fit. However, there is no guidance on how to read the magnitude of \mathcal{D} . To address this we propose to compare it to a benchmark.

16. An alternative to this approach would be to consider injective functions $p: \{1, \dots, K\} \rightarrow \{1, \dots, \hat{K}\}$ when $\hat{K} > K$. Below we discuss why we believe that this other approach could be misleading in Remark C.1.

Let $(X_i)_{i=1}^N$ be a synthetic data set generated as described in Appendix B. For each synthetic observation i , we know the construal it belongs to. That is, if there are K “true construals” in the synthetic data, we observe the “true” construal membership function $c: \{1, \dots, N\} \rightarrow \{1, \dots, K\}$, where $c(i) = k$ means that observation i belongs to construal k . With this information, we can estimate $\bar{\Sigma}^k$ using only the observations i such that $c(i) = k$. This correlation matrix differs from the “true” correlation matrix for construal k (Σ^k) for two reasons: (i) there is some sample variation and, more importantly, (ii) we only observe answers X_i and not the accurate latent positions X_i^* .

When we estimate the correlation matrices $(\hat{\Sigma}_k)_{k=1}^{\hat{K}}$ using a given method, we also face the above two sources of noise. However, on top of them, each method must estimate the construal membership function. Therefore, we believe that $\mathcal{D}((\bar{\Sigma}^k)_{k=1}^K, (\Sigma^k)_{k=1}^K) := \mathcal{D}^b$ provides a good benchmark to measure the performance of the method when estimating the underlying correlation structure.

Once we compute \mathcal{D}^b , we report

$$(C.4) \quad \frac{\mathcal{D}\left((\hat{\Sigma}_k)_{k=1}^{\hat{K}}, (\Sigma^k)_{k=1}^K\right)}{\mathcal{D}^b},$$

where $(\hat{\Sigma}_k)_{k=1}^{\hat{K}}$ are derived using each method. The magnitude of the above ratio is meaningful. For instance, if the ratio is 2 for a given method, one can say that the “estimation of the underlying correlation structure with the method is twice as bad as if we knew the true membership of each observation”.

APPENDIX D. ADDITIONAL SIMULATION RESULTS

	RCA (1)	RCA (2)	RCA (3)	RCA (4)	CCA (1)	CCA (2)	BCA (1)	BCA (2)
<i>E1.6: All random</i>								
Mean NMI	0.315	0.022	0.329	0.027	0.347	0.233	0.264	0.271
Mean scaled NMI	0.187	0.017	0.136	0.021	0.220	0.187	0.214	0.217
Right # construals (%)	0.189	0.353	0.046	0.358	0.203	0.354	0.410	0.397
Mean Abs. Dev. in # construals	1.858	0.898	3.417	0.863	1.573	0.776	0.699	0.723
Mean Correlational Dissimilarity	5.495	5.555	5.891	5.625	5.424	4.226	4.012	4.028
<i>E1.5: Asymmetric neutrality</i>								
Mean NMI	0.360	0.018	0.371	0.023	0.445	0.309	0.282	0.292
Mean scaled NMI	0.204	0.014	0.145	0.017	0.262	0.254	0.223	0.230
Right # construals (%)	0.161	0.380	0.036	0.381	0.153	0.416	0.367	0.369
Mean Abs. Dev. in # construals	2.051	0.858	3.523	0.855	1.894	0.679	0.740	0.756
Mean Correlational Dissimilarity	6.709	6.593	7.254	6.635	6.979	4.710	4.800	4.842
<i>E2.1: All fixed 7</i>								
Mean NMI	0.186	0.012	0.186	0.013	0.069	0.069	0.176	0.170
Mean scaled NMI	0.068	0.012	0.068	0.012	0.069	0.069	0.169	0.123
Right # construals (%)	0.000	1.000	0.000	0.969	1.000	1.000	0.867	0.363
Mean Abs. Dev. in # construals	2.004	0.000	2.004	0.031	0.000	0.000	0.133	0.778 3
Mean Correlational Dissimilarity	16.918	25.113	16.918	25.083	10.610	10.610	6.421	14.780
<i>E2.3 Individual. thresh.</i>								
Mean NMI	0.156	0.010	0.157	0.011	0.080	0.082	0.156	0.147
Mean scaled NMI	0.108	0.008	0.102	0.009	0.003	0.055	0.147	0.122
Right # construals (%)	0.351	0.937	0.342	0.902	0.000	0.218	0.823	0.594
Mean Abs. Dev. in # construals	0.973	0.063	1.629	0.099	7.012	0.875	0.184	0.562
Mean Correlational Dissimilarity	6.982	9.507	7.330	9.495	11.667	6.551	2.449	4.045
<i>E2.2: Varying thresholds</i>								
Mean NMI	0.165	0.014	0.165	0.016	0.080	0.083	0.154	0.144
Mean scaled NMI	0.097	0.011	0.092	0.012	0.004	0.056	0.146	0.116
Right # construals (%)	0.228	0.887	0.214	0.848	0.000	0.212	0.859	0.560
Mean Abs. Dev. in # construals	1.240	0.115	1.628	0.154	6.505	0.887	0.148	0.601
Mean Correlational Dissimilarity	7.052	9.883	7.303	9.861	12.474	7.381	2.600	4.621
<i>E1.4: Random distances</i>								
Mean NMI	0.392	0.017	0.405	0.023	0.501	0.349	0.325	0.340
Mean scaled NMI	0.224	0.013	0.170	0.016	0.217	0.266	0.259	0.270
Right # construals (%)	0.168	0.336	0.038	0.334	0.048	0.374	0.380	0.371
Mean Abs. Dev. in # construals	2.058	0.959	3.338	1.008	2.780	0.753	0.730	0.745
Mean Correlational Dissimilarity	8.004	7.364	8.632	7.451	11.440	10.284	5.379	5.414
<i>E1.3: Random options</i>								
Mean NMI	0.363	0.033	0.377	0.041	0.435	0.291	0.307	0.318
Mean scaled NMI	0.207	0.023	0.158	0.026	0.198	0.219	0.246	0.253
Right # construals (%)	0.169	0.329	0.065	0.307	0.065	0.360	0.388	0.386
Mean Abs. Dev. in # construals	2.035	0.919	3.385	1.041	2.475	0.776	0.733	0.746
Mean Correlational Dissimilarity	6.932	6.992	7.605	7.106	10.533	9.873	4.986	5.018
<i>E1.1: All fixed 5</i>								
Mean NMI	0.407	0.019	0.420	0.024	0.535	0.364	0.335	0.356
Mean scaled NMI	0.223	0.013	0.172	0.017	0.235	0.276	0.273	0.289
Right # construals (%)	0.152	0.321	0.048	0.334	0.027	0.381	0.398	0.392
Mean Abs. Dev. in # construals	2.190	0.972	3.394	0.978	2.744	0.769	0.712	0.734
Mean Correlational Dissimilarity	9.079	7.905	9.508	7.992	12.777	12.014	5.592	5.577
<i>E1.2: All fixed 7</i>								
Mean NMI	0.435	0.019	0.446	0.024	0.568	0.390	0.362	0.385
Mean scaled NMI	0.246	0.014	0.187	0.016	0.284	0.302	0.292	0.312
Right # construals (%)	0.147	0.321	0.028	0.314	0.061	0.408	0.387	0.405
Mean Abs. Dev. in # construals	2.123	0.984	3.313	1.066	2.294	0.720	0.727	0.725
Mean Correlational Dissimilarity	10.593	9.250	11.149	9.346	13.631	11.825	6.668	6.565

TABLE 6. Simulation results for RCA (1) Goldberg 2011 implementation, (2) without removing small values, (3) Newman’s stopping criterion, (4) Newman’s stopping criterion and without removing small values; for CCA (1) Boutyline 2017 implementation, (2) without removing small values; for BCA (1) without magnitude μ_1 , (2) with magnitude μ_2 .

APPENDIX E. ABOUT THE GAUSSIAN COPULA

This Appendix briefly discusses how the Gaussian copula imposes a correlation structure on the latent positions. To visualize the Gaussian copula, consider the bivariate case where there are two latent positions (u_1, u_2) . Recall that in this case the Gaussian copula is defined by

$$(E.1) \quad C(u_1, u_2; \Sigma) = \Phi_{\Sigma}(\Phi^{-1}(u_1), \Phi^{-1}(u_2)),$$

where Φ_{Σ} is the cdf of a multivariate normal distribution with mean 0 and correlation matrix Σ , and Φ is the cdf of a univariate standard normal. The correlation matrix is

$$(E.2) \quad \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix},$$

being $\rho \in [-1, 1]$ the correlation between the two latent positions. So the family of bivariate Gaussian copulas is determined by a single parameter: ρ .

It is customary to plot the level curves of $C(\cdot, \cdot; \Sigma)$ to describe its behaviour. That is, we draw sets $\{(u_1, u_2) \in [0, 1]^2 : C(u_1, u_2; \Sigma) = \bar{C}\}$ for different values of $\bar{C} \in [0, 1]$. Figure 4 plots the level sets for $\rho = -0.7, 0, 0.7$.




FIGURE 4. Contour diagrams for the bivariate Gaussian copula. Left: $\rho = -0.7$, middle: $\rho = 0$ (independence), and right: $\rho = 0.7$.

A copula is itself the cdf of a random vector (U_1, U_2) with support in $[0, 1]^2$. Following the properties in Nelsen 2006, p. 10 we have that:

$$(E.3) \quad P(U_2 \leq u_2 | U_1 \leq u_1) = \frac{C(u_1, u_2; \Sigma)}{u_1}$$

Then, we can check the value of the copula at, say, point $(0.4, 0.4)$ to understand how knowing that “ U_1 is below 0.4” affects our knowledge about “ U_2 is below 0.4”. Following Figure 4, when $\rho = -0.7$, the copula satisfies $C(0.4, 0.4; \Sigma) < 0.1$. That is, $P(U_2 \leq 0.4 | U_1 \leq 0.4) < 0.1/0.4 = 1/4$.

When $\rho = 0$ and $\rho = 0.7$ the copula satisfies $0.1 < C(0.4, 0.4; \Sigma) < 0.2$ and $C(0.4, 0.4; \Sigma) > 0.2$, respectively. That is: $1/4 < P(U_2 \leq 0.4|U_1 \leq 0.4) < 1/2$ if $\rho = 0$ and $P(U_2 \leq 0.4|U_1 \leq 0.4) > 1/2$ if $\rho = 0.7$. In conclusion: if there is negative correlation between the latent positions (U_1, U_2) , knowing that U_1 is small leads to low probability of U_2 also being small (less than $1/4$). On the other hand, if there is positive correlation between the latent positions, if we know that U_1 is small, there is a large probability of U_2 also being small (larger than $1/2$).

Another way of understanding the relationship structure imposed by the copulas is to take a look of the level curves of their density. According to Xue-Kun Song 2000, the density of the Gaussian family of copulas is:

$$(E.4) \quad c(u_1, u_2; \Sigma) = \frac{1}{\sqrt{\det(\Sigma)}} \exp \left\{ \frac{1}{2} \begin{pmatrix} \Phi^{-1}(u_1) & \Phi^{-1}(u_2) \end{pmatrix} (\mathbb{I} - \Sigma^{-1}) \begin{pmatrix} \Phi^{-1}(u_1) \\ \Phi^{-1}(u_2) \end{pmatrix} \right\},$$

where \mathbb{I} is the 2×2 identity matrix.

When $\rho = 0$ we have that $\Sigma = \mathbb{I}$. In that case, according to the above equation, $c(u_1, u_2; \Sigma) = 1$ for every point (u_1, u_2) . That is, since there is no dependence structure on (U_1, U_2) , its mass is uniformly distributed: all points are equally probable. Figure 5 plots the level-curves of $c(\cdot, \cdot; \Sigma)$ for $\rho = -0.7, 0.7$.




FIGURE 5. Level curves of the density of the bivariate Gaussian copula. Left: $\rho = -0.7$ and right: $\rho = 0.7$.

Take the independence case ($\rho = 0$), where the mass at all points is 1, as a benchmark. Then, when there is a negative correlation between the latent positions ($\rho = -0.7$), more mass is concentrated around the $u_2 = -u_1$ diagonal. Then, the mass goes to zero as one diverts away from the $u_2 = -u_1$ diagonal. In turn, when there is a positive correlation ($\rho = 0.7$), more mass is concentrated around the $u_2 = u_1$ diagonal.

REFERENCES

- Amelio, Alessia, and Clara Pizzuti. 2015. "Is normalized mutual information a fair measure for comparing community detection methods?" In *Proceedings of the 2015 IEEE/ACM international conference on advances in social networks analysis and mining 2015*, 1584–1585.
- Baldassarri, Delia, and Amir Goldberg. 2014. "Neither Ideologues nor Agnostics: Alternative Voters' Belief System in an Age of Partisan Politics." *American Journal of Sociology* 120 (1): 45–95. ISSN: 00029602, 15375390, accessed November 17, 2023.
- Blondel, Vincent D, Jean-Loup Guillaume, Renaud Lambiotte, and Etienne Lefebvre. 2011. "Fast unfolding of communities in large networks." *Journal of Statistical Mechanics: Theory and Experiment* 2008 (10): P10008. <https://doi.org/10.1088/1742-5468/2008/10/P10008>.
- Bollen, Kenneth A, and Kenney H Barb. 1981. "Pearson's r and coarsely categorized measures." *American Sociological Review*, 232–239.
- Bourdieu, Pierre. 1984. "A social critique of the judgement of taste." *Traducido del francés por R. Nice. Londres, Routledge*.
- Boutyline, Andrei. 2017. "Improving the Measurement of Shared Cultural Schemas with Correlational Class Analysis: Theory and Method." *Sociological Science* 4 (15): 353–393. ISSN: 2330-6696. <https://doi.org/10.15195/v4.a15>. <http://dx.doi.org/10.15195/v4.a15>.
- Cohen, Mike X. 2022. *Practical Linear Algebra for Data Science*. " O'Reilly Media, Inc."
- DiMaggio, Paul, and Amir Goldberg. 2018. "Searching for Homo Economicus: Variation in Americans' Construals of and Attitudes toward Markets." *European Journal of Sociology* 59 (2): 151–189. <https://doi.org/10.1017/S0003975617000558>.
- DiMaggio, Paul, Ramina Sotoudeh, Amir Goldberg, and Hana Shepherd. 2018. "Culture out of attitudes: Relationality, population heterogeneity and attitudes toward science and religion in the U.S." *Poetics* 68:31–51. ISSN: 0304-422X. <https://doi.org/https://doi.org/10.1016/j.poetic.2017.11.001>.
- Durkheim, E., and W.D. Halls. 1997. *The Division of Labor in Society*. Free Press paperback. Free Press. ISBN: 9780684836386.

- Fisher, N.I. 1997. "Copulas." In *Encyclopedia of Statistical Sciences, Volume 1*, edited by Samuel Kotz, Narayanaswamy Balakrishnan, Campbell B Read, and Brani Vidakovic. John Wiley & Sons.
- Goldberg, Amir. 2011. "Mapping shared understandings using relational class analysis: The case of the cultural omnivore reexamined." *American journal of sociology* 116 (5): 1397–1436.
- Goldberg, Amir, and Sarah K. Stein. 2018. "Beyond Social Contagion: Associative Diffusion and the Emergence of Cultural Variation." *American Sociological Review* 83 (5): 897–932. <https://doi.org/10.1177/0003122418797576>.
- Greene, William H, and David A Hensher. 2010. *Modeling ordered choices: A primer*. Cambridge University Press.
- Krosnick, Jon A. 1999. "Survey research." *Annual review of psychology* 50 (1): 537–567.
- Lewandowski, Daniel, Dorota Kurowicka, and Harry Joe. 2009. "Generating random correlation matrices based on vines and extended onion method." *Journal of multivariate analysis* 100 (9): 1989–2001.
- Mahmoudi, Amin, and Dariusz Jemielniak. 2024. "Proof of biased behavior of Normalized Mutual Information." *Scientific Reports* 14, no. 1 (April): 9021. ISSN: 2045-2322. <https://doi.org/10.1038/s41598-024-59073-9>. <https://doi.org/10.1038/s41598-024-59073-9>.
- Malhotra, Neil, and Jon A Krosnick. 2007. "The effect of survey mode and sampling on inferences about political attitudes and behavior: Comparing the 2000 and 2004 ANES to Internet surveys with nonprobability samples." *Political Analysis* 15 (3): 286–323.
- Marx, K., and F. Engels. 1848. *The Communist Manifesto*. Classics Series. Penguin Publishing Group. ISBN: 9780140447576.
- Nelsen, Roger B. 2006. *An introduction to copulas*. Springer.
- Newman, Mark EJ. 2004. "Analysis of weighted networks." *Physical review E* 70 (5): 056131.
- . 2006. "Finding community structure in networks using the eigenvectors of matrices." *Physical review E* 74 (3): 036104.
- O'Brien, Robert M, and Pamela Homer. 1987. "Corrections for coarsely categorized measures: LISREL's polyserial and polychoric correlations." *Quality and Quantity* 21 (4): 349–360.

- Ostrom, Thomas M. 1987. "Bipolar survey items: An information processing perspective." In *Social information processing and survey methodology*, 71–85. Springer.
- Ostrom, Thomas M, Andrew L Betz, and John J Skowronski. 1992. "Cognitive representation of bipolar survey items." In *Context effects in social and psychological research*, 297–311. Springer.
- Sotoudeh, Ramina, and Paul DiMaggio. 2023. "Coping With Plenitude: A Computational Approach to Selecting the Right Algorithm." *Sociological Methods & Research* 52 (4): 1838–1882. <https://doi.org/10.1177/00491241211031273>.
- Weber, M. 1904. *The Protestant Ethic and the "Spirit" of Capitalism: and Other Writings*. Penguin twentieth-century classics. Penguin Publishing Group. ISBN: 9781101098479.
- Xue-Kun Song, Peter. 2000. "Multivariate dispersion models generated from Gaussian copula." *Scandinavian Journal of Statistics* 27 (2): 305–320.