UNIFORM BOUNDS AND THE INVISCID LIMIT FOR THE NAVIER-STOKES EQUATIONS WITH NAVIER BOUNDARY CONDITIONS

MUSTAFA SENCER AYDIN AND IGOR KUKAVICA

ABSTRACT. We consider the vanishing viscosity problem for solutions of the Navier-Stokes equations with Navier boundary conditions in the half-space. We lower the currently known conormal regularity needed to establish that the inviscid limit holds. Our requirement for the Lipschitz initial data is that the first four conormal derivatives are bounded along with two for the gradient. In addition, we establish a new class of initial data for the local existence and uniqueness for the Euler equations in the half-space or a channel for initial data in the conormal space without conormal requirements on the gradient.

1. INTRODUCTION

We address the vanishing viscosity limit problem for the three-dimensional incompressible Navier-Stokes equations

$$\partial_t u^{\nu} - \nu \Delta u^{\nu} + u^{\nu} \cdot \nabla u^{\nu} + \nabla p^{\nu} = 0, \qquad \nabla \cdot u^{\nu} = 0, \qquad (x, t) \in \Omega \times (0, T), \tag{1.1}$$

with the Navier boundary conditions

$$u^{\nu} \cdot n = 0, \qquad \left(\frac{1}{2}(\nabla u^{\nu} + \nabla^T u^{\nu}) \cdot n\right)_{\tau} = -\mu u^{\nu}_{\tau}, \qquad (x,t) \in \partial\Omega \times (0,T), \tag{1.2}$$

where $\mu \in \mathbb{R}$ is a constant, n is the outward unit normal vector, and $v_{\tau} = v - (v \cdot n)n$ is the tangential part of v. It is a well-known problem whether the solutions $\{u^{\nu}\}_{\nu}$ for (1.1) converge to a solution u for the Euler equations

$$u_t + u \cdot \nabla u + \nabla p = 0, \qquad \nabla \cdot u = 0, \qquad (x, t) \in \Omega \times (0, T), \tag{1.3}$$

with the slip boundary condition

$$u \cdot n = 0 \text{ on } \partial\Omega \times (0, T), \tag{1.4}$$

as $\nu \to 0$. For the no-slip boundary condition $u^{\nu} = 0$ on $\partial\Omega \times (0, T)$, the inviscid limit problem is to a large extent open due to the presence of strong boundary layers; see [CL+, FTZ, GK+, GMM, K, Ke, KLS, KVW, M, MM, SC1, SC2, TW] to point out only few references. However, substantial progress has been made for the Navier-type boundary conditions (1.2). When $\Omega \subset \mathbb{R}^3$, this problem was first addressed by Iftimie and Planas in [IP], where they consider (1.3)–(1.4) with the initial data $u_0 \in H^s$ for s > 2.5, the condition that guarantees the local-in-time well-posedness of Euler equations. Then, by assuming the L^2 convergence of the initial data u_0^{ν} for (1.1) to u_0 , they prove the $L_t^{\infty} L_x^2$ convergence of the weak Leray solutions to the local strong solution of the Euler equations. A quantitative result is obtained by Gie and Kelliher in [GK], who considered the generalized slip condition

$$u^{\nu} \cdot n = 0, \qquad \left(\frac{1}{2}(\nabla u^{\nu} + \nabla^T u^{\nu}) \cdot n\right)_{\tau} = (Au^{\nu})_{\tau}, \qquad (x,t) \in \partial\Omega \times (0,T), \tag{1.5}$$

where A is a smooth symmetric tensor. Assuming that the initial data belongs to H^5 , the authors establish the $L_t^{\infty} L_x^2$ and $L_t^2 H_x^1$ convergence with the rates $\nu^{\frac{3}{4}}$ and $\nu^{\frac{1}{4}}$, respectively. These rates were improved in [XZ2] in the case of the generalized vorticity-slip condition

$$u^{\nu} \cdot n = 0,$$
 $(\operatorname{curl} u^{\nu}) \times n = (Bu^{\nu})_{\tau},$ $(x, t) \in \partial \Omega \times (0, T),$

which is equivalent to (1.5) for some smooth symmetric tensor A. Their initial data belongs to H^3 and satisfies (1.7). Under these assumptions, the authors establish, for any s > 0 and $2 \le p < \infty$, the convergence in $L_t^{\infty} L_x^2, L_t^{\infty} H_x^1, L_t^{\infty} W_x^{1,p}, L_{t,x}^{\infty}$, with the rates $\nu^{1-s}, \nu^{\frac{1-s}{2}}, \nu^{\frac{1-s}{p}}, \nu^{\frac{2}{5}(1-s)}$, respectively. We note that the inviscid limit

problem has also been studied with several other Navier-type boundary conditions such as

$$u^{\nu} \cdot n = 0, \qquad \begin{cases} (\operatorname{curl} u) \cdot n = 0 \\ (\operatorname{curl} u) \cdot n = 0 = \Delta u \times n \\ (\operatorname{curl} u)_{\tau} = 0 \\ (\operatorname{curl} u) \times n = 0 \\ ((-pI + \frac{\nu}{2}(\nabla u^{\nu} + \nabla^{T} u^{\nu})) \cdot n) \cdot \tau = 0; \end{cases}$$
(1.6)

we refer the reader to [BdVC1, BdVC2, BdVC3, BdVC4, BS1, BS2, CQ, DN, IS, NP, TWZ, WXZ, X, XX, XZ1] for the corresponding results.

In [MR1], Masmoudi and Rousset derived uniform-in- ν estimates for the Navier-Stokes system in the functional framework of Sobolev conormal spaces. In particular, they assumed that

$$(u, \nabla u)|_{t=0} \in H^7_{co} \times (H^6_{co} \cap W^{1,\infty}_{co}),$$
 (1.7)

(see the definitions (2.3) and (2.4) below) and established the local well-posedness of Euler equations in the same functional setting. Moreover, the unique solution for (1.3)–(1.4) is realized as the $L_{x,t}^{\infty} \cap L_t^{\infty} L_x^2$ limit of u^{ν} as $\nu \to 0$. After establishing the uniform estimates, they use a compactness argument to prove qualitatively that the inviscid limit holds.

In this work, we lower the regularity assumption (1.7) in [MR1] for the case of the three-dimensional half-space. In particular, we address the initial data

$$(u, \nabla u)|_{t=0} \in (H^4_{\mathrm{co}} \cap W^{2,\infty}_{\mathrm{co}}) \times (H^2_{\mathrm{co}} \cap L^{\infty})$$

$$(1.8)$$

and establish the vanishing viscosity limit for $\mu \in \mathbb{R}$; see Theorem 2.2. Then, assuming $\mu \ge 0$, we obtain the inviscid limit under the condition

$$(u, \nabla u)|_{t=0} \in (H^4_{co} \cap W^{2,\infty}_{co}) \times (H^1_{co} \cap L^{\infty});$$
(1.9)

see Theorem 2.3(ii). To prove both of these results, we first obtain uniform-in- ν estimates, stated in Theorems 2.1 and 2.3(i).

We now briefly describe the key steps allowing us to establish the inviscid limit with the data specified in (1.8) and (1.9). First, we note that in (1.7) the condition $\nabla u|_{t=0} \in W_{co}^{1,\infty}$ is required. To maintain this, it seems that we need at least six and five conormal derivatives on u and ∇u , respectively, as demonstrated in [MR2] for the free boundary Navier-Stokes equations. Due to the derivative loss in Euler equations, the need to control $\nabla u \in W_{co}^{1,\infty}$ arises from requiring $\nabla u \in H_{co}^{m-1}$ when $u \in H_{co}^m$. Namely, in [MR1], the authors derive the uniform-in- ν estimate

$$\frac{d}{dt} \|\nabla u^{\nu}\|_{H^{m-1}_{\infty}}^{2} \lesssim \|\nabla p^{\nu}\|_{H^{m-1}_{\infty}} \|\nabla u^{\nu}\|_{H^{m-1}_{\infty}} + (1 + \|\nabla u^{\nu}\|_{W^{1,\infty}_{\infty}})(\|\nabla u^{\nu}\|_{H^{m-1}_{\infty}}^{2} + \|u^{\nu}\|_{H^{m}_{\infty}}^{2}),$$

for all integer $m \ge 7$, and it is challenging to eliminate $\|\nabla u^{\nu}\|_{W^{1,\infty}_{co}}$ without introducing more than six conormal derivatives of u^{ν} . We overcome this in Proposition 4.1 by establishing an estimate that may be simplified to

$$\frac{d}{dt} \|\nabla u^{\nu}\|_{H^m_{co}}^2 \lesssim \|\nabla p^{\nu}\|_{H^m_{co}} \|\nabla u^{\nu}\|_{H^m_{co}} + (1 + \|u^{\nu}\|_{W^{2,\infty}_{co}} + \|\nabla u^{\nu}\|_{L^{\infty}})(\|\nabla u^{\nu}\|_{H^m_{co}}^2 + \|u^{\nu}\|_{H^m_{co}}^2),$$

for m = 1, 2, which does not require $\nabla u^{\nu} \in W_{co}^{1,\infty}$. However, we still need to ensure that $u^{\nu} \in H_{co}^4$ is bounded uniformly in ν , which cannot be achieved by repeating the arguments in [MR1]. This leads to

$$\frac{d}{dt} \|u^{\nu}\|_{H^m_{\infty}}^2 \lesssim \|\nabla p^{\nu}\|_{H^{m-1}_{\infty}} \|u^{\nu}\|_{H^m_{\infty}} + (1 + \|\nabla u^{\nu}\|_{L^{\infty}})(\|\nabla u^{\nu}\|_{H^{m-1}_{\infty}}^2 + \|u^{\nu}\|_{H^m_{\infty}}^2),$$
(1.10)

for all integer $m \ge 0$. In our case, (1.10) dictates $\nabla u^{\nu} \in H^3_{co}$, which is undesirable. For this reason, we revise the conormal estimates and obtain

$$\frac{d}{dt}\|u(t)\|_{4}^{2} \lesssim \|\nabla p^{\nu}\|_{H^{3}_{\infty}}\|u^{\nu}\|_{H^{4}_{\infty}} + (1 + \|\nabla u^{\nu}\|_{L^{\infty}} + \|u^{\nu}\|_{W^{2,\infty}_{\infty}})\|u^{\nu}\|_{H^{4}_{\infty}}^{2};$$
(1.11)

for the precise statement, see Proposition 3.1. Observe that (1.11) does not contain $\|\nabla u^{\nu}\|_{H^m_{co}}$, and this suggests that conormal derivatives of u do not explicitly depend on conormal derivatives of ∇u . This dependence is only by the term $\|u^{\nu}\|_{W^{2,\infty}}$. Similarly, our pressure estimates (5.1) also remove the dependence between the conormal derivatives of p

and of ∇u present in [MR1]. We remark that all the results are possible by relying on $||u||_{W^{2,\infty}_{co}}$ rather than $||\nabla u^{\nu}||_{H^m_{co}}$ or $||\nabla u^{\nu}||_{W^{1,\infty}_{co}}$. The requirement $u_0 \in W^{2,\infty}_{co}$ is much weaker than (1.7). Indeed, from (2.12) we can obtain

$$\|u\|_{W^{2,\infty}_{\infty}}^{2} \lesssim \|\nabla u\|_{H^{2+s_{1}}_{\infty}} \|u\|_{H^{2+s_{2}}_{\infty}}$$

for $s_1 + s_2 = 2+$. This means that we can control $||u||_{W^{2,\infty}_{co}}$ with $||u||_5$ and $||\nabla u||_2$. However, controlling $||u||_{W^{2,\infty}_{co}}$ under the assumptions (1.8) and (1.9) requires a different approach. This is where the cases (1.8) and (1.9) differ from each other depending on the sign of the friction parameter μ .

After establishing the uniform estimates, we pass to the limit by showing that $\{u^{\nu}\}$ is a Cauchy sequence in $L_{t,x}^{\infty} \cap L_t^{\infty} L_x^2$, from where we obtain a unique solution for the Euler equations with an explicit convergence rate for the Navier-Stokes solutions given in (2.6). We emphasize that both of these inviscid limit results are new existence results for the Euler equations. Moreover, they are sharper than the previously mentioned results in the sense that our assumptions on the normal derivative of u are minimal. The earlier works on the inviscid limit problem under Navier-type boundary conditions either assume that a strong solution for the Euler equations exists or that the initial data belongs to a space where the local well-posedness for the Euler equations is already known, such as $H^{2.5+\delta}$. However, here, as can be seen from (1.8) and (1.9), we require the boundedness of only a single normal derivative at the expense of several conormal derivatives, which is sharp from the perspective of inviscid limits, but it is further improvable when we disregard the vanishing viscosity approach.

In Theorem 2.4, we obtain a well-posedness result for the Euler equations under the assumption

$$[u, \nabla u)|_{t=0} \in (H^4_{\rm co} \cap W^{2,\infty}_{\rm co}) \times L^{\infty}, \tag{1.12}$$

which completely removes the differentiability requirement on ∇u in the conormal sense. The a priori estimates in this case follow from repeating the proof of Proposition 2.3 for $\nu = 0$. However, the inviscid limit approach does not apply as a construction method. This is due to the Laplacian term when we perform $W_{co}^{2,\infty}$ estimates. To remedy this, we may employ Theorem 2.2 or Theorem 2.3 to obtain an approximate sequence of solutions with a lifespan independent of the approximation. We elaborate more on this in Section 9.

In addition to our main results, our uniform estimates given for example in (2.17)–(2.20) are also sharp in integervalued Sobolev conormal spaces. In particular, when we consider (1.8), (1.9), or (1.12) with fewer integer conormal derivatives on either u or ∇u , we lose control of the key quantity $||u||_{W^{2,\infty}_{co}}$. We elaborate more on this in Section 6. We also emphasize that, in the view of (4.1), (4.3), and (4.6), the assumption $\nabla u_0 \in L^{\infty}$ in (1.8), (1.9), and (1.12) can be replaced by $(\omega_1, \omega_2) \in L^{\infty}$. In other words, our results hold when we assume that the tangential part of the initial vorticity is bounded, and we may remove the requirement that the initial velocity is Lipschitz.

2. PRELIMINARIES AND THE MAIN RESULTS

We denote $\Omega = \mathbb{R}^3_+$, and for $x \in \Omega$ we write $x = (x_h, z) = (x_1, x_2, z) \in \mathbb{R}^2 \times \mathbb{R}_+$. We consider

$$\partial_t u^{\nu} - \nu \Delta u^{\nu} + u^{\nu} \cdot \nabla u^{\nu} + \nabla p^{\nu} = 0, \qquad \nabla \cdot u^{\nu} = 0, \qquad (x,t) \in \Omega \times (0,T), \tag{2.1}$$

with the boundary conditions

$$u_3^{\nu} = 0, \qquad \partial_z u_{\rm h}^{\nu} = 2\mu u_{\rm h}^{\nu}, \qquad (x,t) \in \{z=0\} \times (0,T),$$
(2.2)

where $u_h = (u_1, u_2)$. To introduce Sobolev conormal spaces, let $\varphi(z) = z/(1+z)$, and denote $Z_1 = \partial_1$, $Z_2 = \partial_2$, and $Z_3 = \varphi \partial_z$. Then define

$$H^m_{\rm co}(\Omega) = \{ f \in L^2(\Omega) : Z^{\alpha} f \in L^2(\Omega), \alpha \in \mathbb{N}^3_0, 0 \le |\alpha| \le m \}$$

$$W^{m,\infty}_{\rm co}(\Omega) = \{ f \in L^{\infty}(\Omega) : Z^{\alpha} f \in L^{\infty}(\Omega), \alpha \in \mathbb{N}^3_0, 0 \le |\alpha| \le m \}.$$
 (2.3)

These are Hilbert and Banach spaces, respectively, endowed with the norms

$$\|f\|_{H^{m}_{\infty}(\Omega)}^{2} = \|f\|_{m}^{2} = \sum_{|\alpha| \le m} \|Z^{\alpha}f\|_{L^{2}(\Omega)}^{2}$$

$$\|f\|_{W^{m,\infty}_{\infty}(\Omega)} = \|f\|_{m,\infty} = \sum_{|\alpha| \le m} \|Z^{\alpha}f\|_{L^{\infty}(\Omega)}.$$
(2.4)

When m = 0, we write $||f||_{L^2}$ and $||f||_{L^{\infty}}$ for $||f||_0$ and $||f||_{0,\infty}$. We are now in a position to state our main results.

Theorem 2.1 (Existence and Uniqueness). Assume that $\nu \in (0,1]$ and $\mu \in \mathbb{R}$. Let $u_0 \in H^4_{co}(\Omega) \cap W^{2,\infty}_{co}(\Omega)$, with $\nabla u_0 \in H^2_{co}(\Omega) \times L^{\infty}(\Omega)$, be such that div $u_0 = 0$, and $u_0 \cdot n = 0$ on $\partial\Omega$. Then, for some T > 0, there exists a unique solution $u^{\nu} \in C([0,T]; H^2_{co}(\Omega)) \cap L^{\infty}(0,T; H^4_{co}(\Omega) \cap W^{2,\infty}_{co}(\Omega))$ to (2.1)–(2.2) on [0,T], and M > 0 independent of ν such that

$$\sup_{[0,T]} \left(\|u^{\nu}(t)\|_{4}^{2} + \|u^{\nu}(t)\|_{2,\infty}^{2} + \|\nabla u^{\nu}(t)\|_{2}^{2} + \|\nabla u^{\nu}(t)\|_{L^{\infty}}^{2} \right) + \nu \int_{0}^{T} \left(\|\nabla u^{\nu}(s)\|_{5}^{2} + \|D^{2}u^{\nu}(s)\|_{2}^{2} \right) ds \leq M, \quad (2.5)$$

where M > 0 depends on the norms of the initial data.

As pointed out in the introduction, the requirement $\nabla u_0 \in H^2_{co}(\Omega) \times L^{\infty}(\Omega)$ can be replaced by $\nabla \omega_0 \in H^2_{co}(\Omega) \times L^{\infty}(\Omega)$ The analogous remark also applies to Theorems 2.2, 2.3, and 2.4. Upon establishing the existence of u^{ν} on a time interval [0, T] which does not depend on ν , the estimate (2.5) is sufficient to pass to the limit in (2.1) to obtain (1.3). Namely, we have the following inviscid limit statement.

Theorem 2.2 (Inviscid Limit). Assume that $\mu \in \mathbb{R}$. Let u_0 and u^{ν} be as in Theorem 2.1. Then, there exists a unique solution $u \in L^{\infty}(0,T; H^4_{co}(\Omega) \cap W^{2,\infty}_{co}(\Omega))$ with $\nabla u \in L^{\infty}(0,T; H^2_{co}(\Omega) \cap L^{\infty}(\Omega))$ to the Euler equations (1.3) and M > 0 independent of ν such that

$$\sup_{[0,T]} (\|u^{\nu} - u\|_{L^{2}}^{2} + \|u^{\nu} - u\|_{L^{\infty}}^{2}) \le M\nu^{\frac{2}{5}},$$
(2.6)

where $\nu \in [0, 1]$ and M > 0 depends on the norms of the initial data.

When the friction parameter μ is non-negative, we obtain a sharper result by imposing conditions on one less conormal derivative on ∇u .

Theorem 2.3 (A sharper result for non-negative friction). Assume that $\nu \in (0,1]$ and $\mu \ge 0$. Let $u_0 \in H^4_{co}(\Omega) \cap W^{2,\infty}_{co}(\Omega)$, and $\nabla u_0 \in H^1_{co}(\Omega) \times L^{\infty}(\Omega)$ be such that div $u_0 = 0$ and $u_0 \cdot n = 0$ on $\partial \Omega$.

i. (Existence and Uniqueness) For all $\nu \in (0,1]$ and $\mu \ge 0$ fixed, there exists a unique solution $u^{\nu} \in C([0,T]; H^1_{co}(\Omega)) \cap L^{\infty}(0,T; H^4_{co}(\Omega) \cap W^{2,\infty}_{co}(\Omega))$ to (2.1)–(2.2) on [0,T], for some T > 0. Moreover, there exists M > 0 independent of ν such that

$$\sup_{[0,T]} (\|u^{\nu}(t)\|_{4}^{2} + \|u^{\nu}(t)\|_{2,\infty}^{2} + \|\nabla u^{\nu}(t)\|_{1}^{2} + \|\nabla u^{\nu}(t)\|_{L^{\infty}}^{2}) + \nu \int_{0}^{T} (\|\nabla u^{\nu}(s)\|_{4}^{2} + \|D^{2}u^{\nu}(s)\|_{1}^{2}) \, ds \leq M,$$

where M depends on the norms of the initial data.

ii. (Inviscid Limit) There exists a unique solution $u \in L^{\infty}(0,T; H^4_{co}(\Omega) \cap W^{2,\infty}_{co}(\Omega))$ with $\nabla u \in L^{\infty}(0,T; H^1_{co}(\Omega) \cap L^{\infty}(\Omega))$ to the Euler equations (1.3) such that (2.6) holds for some constant M > 0 depending on the norms of the initial data and independent of ν .

Furthermore, it is possible to improve on the Euler equations part of Theorem 2.3.

Theorem 2.4 (Euler equations in Sobolev conormal spaces). Let $u_0 \in H^4_{co}(\Omega) \cap W^{2,\infty}_{co}(\Omega)$, and $\nabla u_0 \in L^{\infty}(\Omega)$ be such that div $u_0 = 0$, and $u_0 \cdot n = 0$ on $\partial\Omega$. For some T > 0, there exists a unique solution $u \in L^{\infty}(0,T; H^4_{co}(\Omega) \cap W^{2,\infty}_{co}(\Omega))$ with $\nabla u \in L^{\infty}(\Omega \times (0,T))$ to the Euler equations (1.3) and M > 0 independent of ν such that

$$\sup_{[0,T]} (\|u(t)\|_4^2 + \|u(t)\|_{2,\infty}^2 + \|\nabla u(t)\|_{L^{\infty}}^2) \le M,$$
(2.7)

where M > 0 depends on the norms of the initial data.

When we perform integration by parts in the term involving the Laplacian,

$$-\nu \int_{\Omega} \Delta Z^{\alpha} u^{\nu} Z^{\alpha} u^{\nu} = \nu \int_{\Omega} |\nabla Z^{\alpha} u^{\nu}|^2 + 2\nu \mu \int_{\partial \Omega} |Z^{\alpha} u^{\nu}_{\mathbf{h}}|^2,$$

we obtain a boundary term that is coercive when $\mu \ge 0$. When estimating $\|u^{\nu}\|_{2,\infty}$ with $\mu \in \mathbb{R}$ a boundary term similar to the one above is only controllable when $\nabla u^{\nu} \in H^2_{co}(\Omega)$; see (6.4)–(6.6) for more details. On the other hand, when $\mu \ge 0$, we can estimate $\|u\|_{2,\infty}$ using $\|\nabla u\|_1$. This is why in Theorem 2.3 we only consider non-negative μ . We also remark that the time of existence obtained in Theorems 2.1, 2.3(i), and 2.4 need not be the same. However, for convenience, we denote the time by T. Note that one of the difficulties is that, unlike Z_1 and Z_2 , the operator Z_3 does not commute with ∂_z . Therefore, we analyze the conormal derivatives with or without Z_3 separately, and when we do, we write

$$Z^{\alpha} = Z_{h}^{\tilde{\alpha}} Z_{3}^{k}, \qquad \alpha = (\tilde{\alpha}, k) \in \mathbb{N}_{0}^{2} \times \mathbb{N}_{0}.$$
(2.8)

We shall use the following commutator identities.

Lemma 2.1. Let f be a smooth function. Then there exist smooth, bounded $\{c_{j,\varphi}^k\}_{j=0}^k$ and $\{\tilde{c}_{j,\varphi}^k\}_{j=0}^k$ of z, for $k \in \mathbb{N}$, depending on φ such that

$$(i) Z_{3}^{k} \partial_{z} f = \sum_{j=0}^{k} c_{j,\varphi}^{k} \partial_{z} Z_{3}^{j} f = \partial_{z} Z_{3}^{k} f + \sum_{j=0}^{k-1} c_{j,\varphi}^{k} \partial_{z} Z_{3}^{j} f,$$

$$(ii) \partial_{z} Z_{3}^{k} f = \sum_{j=0}^{k} \tilde{c}_{j,\varphi}^{k} Z_{3}^{j} \partial_{z} f = Z_{3}^{k} \partial_{z} f + \sum_{j=0}^{k-1} \tilde{c}_{j,\varphi}^{k} Z_{3}^{j} \partial_{z} f,$$

$$(iii) Z_{3}^{k} \partial_{zz} f = \sum_{j=0}^{k} \sum_{l=0}^{j} \left(c_{l,\varphi}^{j} c_{j,\varphi}^{k} \partial_{zz} Z_{3}^{l} f + (c_{l,\varphi}^{j})' c_{j,\varphi}^{k} \partial_{z} Z_{3}^{l} f \right),$$

$$(iv) \partial_{zz} Z_{3}^{k} f = \sum_{j=0}^{k} \sum_{l=0}^{j} \tilde{c}_{l,\varphi}^{j} \tilde{c}_{j,\varphi}^{k} Z_{3}^{l} \partial_{zz} f + \sum_{j=0}^{k} (\tilde{c}_{j,\varphi}^{k})' Z_{3}^{j} \partial_{z} f,$$

$$(2.9)$$

where $\tilde{c}_{k,\varphi}^k = 1 = c_{k,\varphi}^k$, and the prime indicates the derivative with respect to the variable z.

The proof is by induction and is based on

$$Z_3\partial_z f = \partial_z Z_3 f - \varphi' \partial_z f$$

We also take advantage of the well-known product estimate given in the next lemma.

Lemma 2.2 (An anisotropic Sobolev product estimate). Let f and g be sufficiently smooth functions and $p_i, q_i \in [2, \infty]$, for i = 1, 2. Then

for any $H_h^{r_i}(\mathbb{R}^2) \subset L_h^{p_i}(\mathbb{R}^2)$ and $H_z^{s_i}(\mathbb{R}_+) \subset L_z^{q_i}(\mathbb{R}_+)$, where i = 1, 2 and $\frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{q_1} + \frac{1}{q_2} = \frac{1}{2}$.

We shall also use the inequalities

$$\|Z^{\alpha_1} f Z^{\alpha_2} g\|_{L^2} \lesssim \|f\|_{L^{\infty}} \|g\|_k + \|f\|_k \|g\|_{L^{\infty}}, \qquad |\alpha_1| + |\alpha_2| = k, \qquad f, g \in L^{\infty} \cap H^k_{\text{co}}$$
(2.11)

and

$$\|f\|_{L^{\infty}}^{2} \lesssim \|(I - \Delta_{h})^{\frac{s_{1}}{2}} \partial_{z} f\|_{L^{2}} \|(I - \Delta_{h})^{\frac{s_{2}}{2}} f\|_{L^{2}}, \qquad s_{1}, s_{2} \ge 0, \qquad s_{1} + s_{2} > 2.$$
(2.12)

For the proof of (2.11), see [G], while (2.12) is proven in [MR2]. Next, using the slip boundary condition, the Hardy inequality and the divergence-free condition, we obtain

$$\left\|\frac{u_{3}^{\nu}}{\varphi}\right\|_{W_{co}^{k,p}} \lesssim \|Z_{h}u_{h}^{\nu}\|_{W_{co}^{k,p}},$$
(2.13)

for all $p \in [1, \infty]$, with $W_{co}^{k,p}$ defined analogously to (2.3). An additional way that we utilize the divergence-free condition is by means of Lemma 2.2. Since for any conormal derivative Z^{α} we have

$$Z^{\alpha}\partial_z u_3^{\nu}\|_{L^2} \lesssim \|u^{\nu}\|_{|\alpha|+1}, \qquad \alpha \in \mathbb{N}_0^3, \tag{2.14}$$

the expression $||Z^{\alpha}u_{3}^{\nu}||_{L_{h}^{4}L_{z}^{\infty}}$ may be estimated by writing

$$\begin{aligned} \|Z^{\alpha}u_{3}^{\nu}\|_{L_{h}^{4}L_{z}^{\infty}} &\lesssim \|Z^{\alpha}u_{3}^{\nu}\|_{H_{h}^{\frac{1}{2}}H_{z}^{\frac{1}{2}+}} &\lesssim \|Z_{h}Z^{\alpha}u_{3}^{\nu}\|_{L^{2}} + \|Z^{\alpha}u_{3}^{\nu}\|_{L^{2}} + \|\partial_{z}Z^{\alpha}u_{3}^{\nu}\|_{L^{2}} + \|\partial_{z}Z_{h}Z^{\alpha}u_{3}^{\nu}\|_{L^{2}} \\ &\lesssim \|u^{\nu}\|_{|\alpha|+2}. \end{aligned}$$

$$(2.15)$$

The same reasoning gives

$$\|Z^{\alpha}f\|_{L^{4}_{\mathfrak{h}}L^{\infty}_{z}} \lesssim \|f\|_{|\alpha|+1} + \|\partial_{z}f\|_{|\alpha|+1},$$
(2.16)

for every $f \in H^{|\alpha|+1}_{\mathrm{co}}(\Omega)$ such that $\partial_z f \in H^{|\alpha|+1}_{\mathrm{co}}(\Omega)$.

Now, we fix $\nu \in (0, 1]$, and denote by (u, p), instead of (u^{ν}, p^{ν}) , the smooth solution to the Navier Stokes system with the viscosity ν .

Proposition 2.3 (A priori estimates). Any smooth solution u to (2.1)–(2.2) defined on [0, T], satisfies the following:

i. If $\mu \in \mathbb{R}$, we have

$$N^{2}(t) = \left(\|u(t)\|_{4} + \|u(t)\|_{2,\infty} + \|\nabla u(t)\|_{2} + \|\nabla u(t)\|_{L^{\infty}}\right)^{2} \le C\left(N^{2}(0) + \int_{0}^{t} N^{3}(s) \, ds\right)$$
(2.17)

and

$$\nu \int_0^t (\|\nabla u\|_4^2 + \|D^2 u\|_2^2) \, ds \le C \left(N^2(0) + \int_0^t N^3(s) \, ds \right), \tag{2.18}$$

for $t \in [0, T]$.

ii. If $\mu \ge 0$, we have

$$\bar{N}^{2}(t) = \left(\|u(t)\|_{4} + \|u(t)\|_{2,\infty} + \|\nabla u(t)\|_{1} + \|\nabla u(t)\|_{L^{\infty}}\right)^{2} \le C\left(\bar{N}^{2}(0) + \int_{0}^{t} \bar{N}^{3}(s) \, ds\right)$$
(2.19)

and

$$\nu \int_0^t (\|\nabla u\|_4^2 + \|D^2 u\|_1^2) \, ds \le C \left(\bar{N}^2(0) + \int_0^t \bar{N}^3(s) \, ds \right), \tag{2.20}$$

for $t \in [0, T]$.

We prove Proposition 2.3 in four steps. In Sections 3, 4, and 5, we derive estimates for conormal derivatives of u, ∇u , and ∇p . Then, in Section 6, we perform L^{∞} estimates for u and ∇u . We conclude the proof of Proposition 2.3, Theorem 2.1, and Theorem 2.3(i) in Section 7. Next, we establish the inviscid limit in Section 8 by proving Theorems 2.2 and 2.3(ii). Finally, in Section 9, we sketch the proof of Theorem 2.4.

3. CONORMAL DERIVATIVE ESTIMATES

In this section, we estimate $||u||_4$.

Proposition 3.1. Let u be a smooth solution of (2.1)-(2.2) on [0,T]. Then we have the inequality

$$\|u(t)\|_{4}^{2} + c_{0}\nu \int_{0}^{t} \|\nabla u(s)\|_{4}^{2} ds \lesssim \|u_{0}\|_{4}^{2} + \int_{0}^{t} \left(\|u(s)\|_{4}^{2} (\|u(s)\|_{W^{1,\infty}} + \|u(s)\|_{2,\infty} + 1) + \|u(s)\|_{4} \|\nabla p(s)\|_{3}\right) ds,$$
(3.1)

where $c_0 > 0$ is independent of ν .

Proof of Proposition 3.1. We proceed by induction on $0 \le |\alpha| \le 4$ and first consider $|\alpha| = 0$, which coincides with the usual L^2 estimate

$$\frac{1}{2}\frac{d}{dt}\|u\|_{L^2}^2 + \nu\|\nabla u\|_{L^2}^2 = -2\mu\nu\|u_h\|_{L^2(\partial\Omega)}^2,$$
(3.2)

where the boundary term results from $-\nu \int u \Delta u \, dx = \nu \|\nabla u\|_{L^2}^2 + \int_{\partial \Omega} \partial_z u u \, d\sigma$, since $u \cdot n = 0$ and n = (0, 0, -1) on $\partial \Omega$. Using (2.2), it follows that

$$-\nu \int u\Delta u \, dx = \nu \|\nabla u\|_{L^2}^2 + 2\mu\nu \|u\|_{L^2(\partial\Omega)}^2, \tag{3.3}$$

which is a coercive term, for $\mu \ge 0$. On the other hand, when $\mu < 0$ we use the trace theorem and Young's inequality to obtain

$$2|\mu|\nu||u||_{L^{2}(\partial\Omega)}^{2} \leq \epsilon\nu||\nabla u||_{L^{2}}^{2} + C_{\epsilon}\mu^{2}\nu||u||_{L^{2}}^{2} \leq \epsilon\nu||\nabla u||_{L^{2}}^{2} + C_{\epsilon}||u||_{L^{2}}^{2},$$
(3.4)

where $\epsilon > 0$ is arbitrarily small. Therefore, combining (3.2) and (3.4) and integrating in time, we get

$$\|u(t)\|_{L^{2}} + c_{0}\nu \int_{0}^{t} \|\nabla u\|_{L^{2}}^{2} ds \lesssim \|u_{0}\|_{L^{2}} + \int_{0}^{t} \|u(s)\|_{L^{2}}^{2} ds,$$
(3.5)

concluding the base step. For the remaining part of the induction, we only present the step from $|\alpha| = 3$ to $|\alpha| = 4$. Thus, we assume that

$$\|u\|_{3}^{2} + c_{0}\nu \int_{0}^{t} \|\nabla u\|_{3}^{2} ds \lesssim \|u_{0}\|_{3}^{2} + \int_{0}^{t} \left(\|u\|_{3}^{2} (\|u\|_{W^{1,\infty}}^{2} + \|u\|_{2,\infty}^{2} + 1) + \|u\|_{3} \|\nabla p\|_{2} \right) ds$$
(3.6)

and aim to show (3.1). We first consider the case when all four conormal derivatives are horizontal, i.e., $Z^{\alpha} = Z_{h}^{\tilde{\alpha}}$ for $|\alpha| = 4$, and then the general case. Applying Z^{α} to $(2.1)_1$, we obtain

$$\partial_t Z_{\mathbf{h}}^{\tilde{\alpha}} u - \nu \Delta Z_{\mathbf{h}}^{\tilde{\alpha}} u + u \cdot \nabla Z_{\mathbf{h}}^{\tilde{\alpha}} u + \nabla Z_{\mathbf{h}}^{\tilde{\alpha}} p = u \cdot \nabla Z_{\mathbf{h}}^{\tilde{\alpha}} u - Z_{\mathbf{h}}^{\tilde{\alpha}} (u \cdot \nabla u),$$
(3.7)

since $Z_h \nabla = \nabla Z_h$. We test (3.7) with $Z_h^{\tilde{\alpha}} u$ and write

$$\frac{1}{2}\frac{d}{dt}\|Z_{h}^{\tilde{\alpha}}u\|_{L^{2}}^{2} + \nu\|\nabla Z_{h}^{\tilde{\alpha}}u\|_{L^{2}}^{2} + 2\mu\nu\|Z_{h}^{\tilde{\alpha}}u\|_{L^{2}(\partial\Omega)}^{2} = -(u \cdot \nabla Z_{h}^{\tilde{\alpha}}u - Z_{h}^{\tilde{\alpha}}(u \cdot \nabla u), Z_{h}^{\tilde{\alpha}}u).$$
(3.8)

Note that to obtain (3.8), we used incompressibility, while as in (3.3), the diffusion term leads to

$$-\nu \int \Delta Z_{\mathbf{h}}^{\tilde{\alpha}} u Z_{\mathbf{h}}^{\tilde{\alpha}} u \, dx = \nu \|\nabla Z_{\mathbf{h}}^{\tilde{\alpha}} u\|_{L^{2}}^{2} + \int_{\partial\Omega} \partial_{z} Z_{\mathbf{h}}^{\tilde{\alpha}} u Z_{\mathbf{h}}^{\tilde{\alpha}} u \, d\sigma = \nu \|\nabla Z_{\mathbf{h}}^{\tilde{\alpha}} u\|_{L^{2}}^{2} + 2\mu\nu \|Z_{\mathbf{h}}^{\tilde{\alpha}} u\|_{L^{2}(\partial\Omega)}^{2}, \tag{3.9}$$

where the last equality follows by applying $Z_h^{\tilde{\alpha}}$ to (2.2). For the boundary term, as in (3.4), we use the trace theorem and Young's inequality to obtain,

$$\|\mu\|\nu\|Z_{h}^{\tilde{\alpha}}u\|_{L^{2}(\partial\Omega)}^{2} \leq \epsilon\nu\|\nabla Z_{h}^{\tilde{\alpha}}u\|_{L^{2}}^{2} + C_{\epsilon}\|u\|_{4}^{2},$$
(3.10)

where $\epsilon > 0$ is arbitrarily small. Therefore, we only have to estimate the commutator term, which we expand as

$$u \cdot \nabla Z_{\mathbf{h}}^{\tilde{\alpha}} u - Z_{\mathbf{h}}^{\tilde{\alpha}} (u \cdot \nabla u) = -\sum_{1 \le |\tilde{\beta}| \le 4} {\tilde{\alpha} \choose \tilde{\beta}} (Z_{\mathbf{h}}^{\tilde{\beta}} u_{\mathbf{h}} \cdot \nabla_{\mathbf{h}} Z_{\mathbf{h}}^{\tilde{\alpha} - \tilde{\beta}} u + Z_{\mathbf{h}}^{\tilde{\beta}} u_{3} \partial_{z} Z_{\mathbf{h}}^{\tilde{\alpha} - \tilde{\beta}} u).$$
(3.11)

For the terms involving $\nabla_h = (\partial_1, \partial_2)$, we take the uniform norm of the factor with the lower number of derivatives, which gives

$$\|Z_{\mathbf{h}}^{\tilde{\beta}}u_{\mathbf{h}} \cdot \nabla_{\mathbf{h}} Z_{\mathbf{h}}^{\tilde{\alpha}-\tilde{\beta}}u\|_{L^{2}} \lesssim \|u\|_{2,\infty} \|u\|_{4}.$$
(3.12)

For the remaining terms in (3.11), when $|\hat{\beta}| \neq 4$, we first divide and multiply by φ to write

$$\left(\left\| Z\frac{u_3}{\varphi} \right\|_{L^{\infty}} \|u\|_4 \lesssim \|u\|_{2,\infty} \|u\|_4, \qquad |\beta| = 1 \right)$$

$$\left\| Z_{\mathbf{h}}^{\tilde{\beta}} \frac{u_{3}}{\varphi} Z_{3} Z_{\mathbf{h}}^{\tilde{\alpha} - \tilde{\beta}} u \right\|_{L^{2}} \lesssim \begin{cases} \|Z_{\varphi}\|_{L^{\infty}} \|u\|^{4} \sim \|u\|_{2,\infty} \|u\|_{4}^{4}, & \|\beta\| = 1 \\ \|Z_{\mathbf{h}} \frac{u_{3}}{\varphi}\|_{L^{\infty}} \|Z_{\mathbf{h}} Z_{3} u\|_{2} + \|Z_{\mathbf{h}} \frac{u_{3}}{\varphi}\|_{2} \|Z_{\mathbf{h}} Z_{3} u\|_{L^{\infty}} \lesssim \|u\|_{4} \|u\|_{2,\infty}, & |\tilde{\beta}| = 2 \\ \|Z_{\mathbf{h}} \frac{u_{3}}{\varphi}\|_{2} \|Z_{\mathbf{h}} Z_{3} u\|_{L^{\infty}} \lesssim \|u\|_{4} \|u\|_{2,\infty}, & |\tilde{\beta}| = 3, \end{cases}$$

$$(3.13)$$

where we have used (2.11) upon taking $(f,g) = (Z_h \frac{u_3}{\omega}, Z_h Z_3 u)$ for $|\tilde{\beta}| = 2$, and (2.13) to bound $Z_h \frac{u_3}{\omega}$. Finally, when $|\tilde{\beta}| = 4$, we immediately have

$$\|Z_{\mathbf{h}}^{\tilde{\beta}}u_{3}\partial_{z}Z_{\mathbf{h}}^{\tilde{\alpha}-\tilde{\beta}}u\|_{L^{2}} \lesssim \|u\|_{4}\|\partial_{z}u\|_{L^{\infty}}, \qquad |\tilde{\beta}| = 4.$$

$$(3.14)$$

Now, combining (3.8)–(3.14) and absorbing the factor of $\nu \|\nabla Z_h^{\tilde{\alpha}} u\|_{L^2}^2$, we obtain

$$\frac{d}{dt} \|Z_{\mathbf{h}}^{\tilde{\alpha}}u\|_{L^{2}}^{2} + c_{0}\nu\|\nabla Z_{\mathbf{h}}^{\tilde{\alpha}}u\|_{L^{2}}^{2} \lesssim \|u\|_{4}(\|u\|_{4} + \|u \cdot \nabla Z_{\mathbf{h}}^{\tilde{\alpha}}u - Z_{\mathbf{h}}^{\tilde{\alpha}}(u \cdot \nabla u)\|_{L^{2}}) \\
\lesssim \|u\|_{4}^{2}(\|u\|_{2,\infty} + \|u\|_{W^{1,\infty}} + 1),$$
(3.15)

where $c_0 > 0$ is independent of ν . Integrating (3.15) implies

$$\|Z_{\mathbf{h}}^{\tilde{\alpha}}u(t)\|_{L^{2}}^{2} + c_{0}\nu \int_{0}^{t} \|\nabla Z_{\mathbf{h}}^{\tilde{\alpha}}u(s)\|_{L^{2}}^{2} ds \lesssim \|Z_{\mathbf{h}}^{\tilde{\alpha}}u_{0}\|_{L^{2}}^{2} + \int_{0}^{t} \|u(s)\|_{4}^{2} (\|u(s)\|_{2,\infty} + \|u(s)\|_{W^{1,\infty}} + 1) ds,$$

$$t \in [0, T]$$

for
$$t \in [0, T]$$
.

Proceeding to the general case, we consider $Z^{\alpha} = Z_{h}^{\tilde{\alpha}} Z_{3}^{k}$ where $1 \le k \le 4$, and apply it to (2.1) obtaining

$$Z^{\alpha}u_{t} - \nu\Delta Z^{\alpha}u + u \cdot \nabla Z^{\alpha}u = u \cdot \nabla Z^{\alpha}u - Z^{\alpha}(u \cdot \nabla u) - Z^{\alpha}\nabla p + \nu Z^{\alpha}\Delta u - \nu\Delta Z^{\alpha}u.$$
(3.16)

Upon testing with $Z^{\alpha}u$, the left-hand side of (3.16) gives

$$\frac{1}{2}\frac{d}{dt}\|Z^{\alpha}u\|_{L^{2}}^{2} + \nu\|\nabla Z^{\alpha}u\|_{L^{2}}^{2}.$$
(3.17)

Observe that there is no boundary term arising from $-\nu\Delta Z^{\alpha}u$ since $Z_3 = \varphi\partial_z = 0$ on $\partial\Omega$. As for the right-hand side of (3.16), we first expand the convection term as

$$\begin{aligned} u \cdot \nabla Z^{\alpha} u - Z^{\alpha} (u \cdot \nabla u) &= u \cdot \nabla Z^{\alpha} u - u \cdot Z^{\alpha} \nabla u - \sum_{1 \le |\beta| \le |\alpha|} \binom{\alpha}{\beta} Z^{\beta} u \cdot Z^{\alpha - \beta} \nabla u \\ &= u_3 \partial_z Z^{\alpha} u - u_3 Z^{\alpha} \partial_z u - \sum_{1 \le |\beta| \le |\alpha|} \binom{\alpha}{\beta} Z^{\beta} u \cdot Z^{\alpha - \beta} \nabla u \\ &= -\sum_{j=0}^{k-1} \tilde{c}_{j,\varphi}^k u_3 \partial_z Z_{h}^{\tilde{\alpha}} Z_{3}^j u - \sum_{1 \le |\beta| \le |\alpha|} \binom{\alpha}{\beta} Z^{\beta} u \cdot Z^{\alpha - \beta} \nabla u \\ &= I_1 + I_2. \end{aligned}$$
(3.18)

To estimate I_1 , we divide and multiply by φ , obtaining

$$\sum_{j=0}^{k-1} \left\| \tilde{c}_{j,\varphi}^{k} \frac{u_{3}}{\varphi} Z_{h}^{\tilde{\alpha}} Z_{3}^{j+1} u \right\|_{L^{2}} \lesssim \left\| \frac{u_{3}}{\varphi} \right\|_{L^{\infty}} \|u\|_{4} \lesssim \|\partial_{z} u\|_{L^{\infty}} \|u\|_{4}, \tag{3.19}$$

using (2.13). As for I_2 , we rewrite

$$I_{2} = -\sum_{1 \le |\beta| \le |\alpha|} \binom{\alpha}{\beta} (Z^{\beta} u_{\mathbf{h}} \cdot \nabla_{\mathbf{h}} Z^{\alpha-\beta} u + Z^{\beta} u_{3} Z^{\alpha-\beta} \partial_{z} u) = I_{21} + I_{22}.$$
(3.20)

The sum I_{21} is estimated using the bounds in (3.12). Next, to bound I_{22} , we proceed as in (3.13)–(3.14), which gives lower order terms when $|\beta| \neq 4$. In particular, since

$$\frac{Z_3 u_3}{\varphi} = \nabla_{\mathbf{h}} \cdot u_{\mathbf{h}},$$

and $1/\varphi$ commutes with Z_h , we only commute ∂_z and Z_3 as in (2.9)(i), obtaining lower order terms. Omitting further details, we conclude

$$(u \cdot \nabla Z^{\alpha} u - Z^{\alpha} (u \cdot \nabla u), Z^{\alpha} u) \lesssim \|u\|_{4} (\|u\|_{2,\infty} + \|u\|_{W^{1,\infty}}).$$
(3.21)

Returning to the right-hand side of (3.16), it remains to estimate the pressure term and the commutator term including the Laplacian. We start by testing the pressure term against $Z^{\alpha}u$, obtaining

$$(Z^{\alpha}\nabla p, Z^{\alpha}u) = (Z^{\alpha}\nabla_{\mathbf{h}}p, Z^{\alpha}u_{\mathbf{h}}) + (Z^{\alpha}\partial_{z}p, Z^{\alpha}u_{3}).$$
(3.22)

Since Z_3 and ∂_z do not commute, we aim to introduce lower order terms and gain a derivative using incompressibility. To achieve this, we first use Lemma 2.1(i) and write

$$(Z_{3}^{k}\partial_{z}Z_{h}^{\tilde{\alpha}}p, Z^{\alpha}u_{3}) = (\partial_{z}Z_{3}^{k}Z_{h}^{\tilde{\alpha}}p, Z^{\alpha}u_{3}) + \sum_{j=0}^{k-1} (c_{j,\varphi}^{k}\partial_{z}Z_{3}^{j}Z_{h}^{\tilde{\alpha}}p, Z^{\alpha}u_{3}).$$
(3.23)

Note that each term under the sum is bounded by $\|\nabla p\|_3 \|u\|_4$. For the leading order term, we integrate by parts and write

$$(\partial_{z} Z_{3}^{k} Z_{h}^{\tilde{\alpha}} p, Z^{\alpha} u_{3}) = (Z^{\alpha} p, \partial_{z} Z^{\alpha} u_{3}) = (Z^{\alpha} p, Z^{\alpha} \partial_{z} u_{3}) - \sum_{j=0}^{k-1} (c_{j,\varphi}^{k} Z^{\alpha} p, Z_{3}^{j} Z_{h}^{\tilde{\alpha}} \partial_{z} u_{3}),$$
(3.24)

as there is no boundary term due to $Z_3 f = 0$ on $\partial\Omega$ for any sufficiently smooth function f. Moreover, each term under the sum is bounded by $\|\nabla p\|_3 \|u\|_4$. Since $Z^{\alpha} \nabla_h \cdot u_h + Z^{\alpha} \partial_z u_3 = 0$, (3.22)–(3.24) imply

$$(Z^{\alpha}\nabla p, Z^{\alpha}u) \lesssim \|\nabla p\|_3 \|u\|_4, \tag{3.25}$$

leaving us only the commutator term for the Laplacian. Using Lemma 2.1, we rewrite

$$\nu Z^{\alpha} \Delta u - \nu \Delta Z^{\alpha} u = \nu Z^{\alpha} \partial_{zz} u - \nu \partial_{zz} Z^{\alpha} u = \nu Z_{h}^{\tilde{\alpha}} (Z_{3}^{k} \partial_{zz} u - \partial_{zz} Z_{3}^{k} u)$$

$$= \nu \sum_{j=0}^{k-1} \sum_{l=0}^{j} \left(c_{l,\varphi}^{j} c_{j,\varphi}^{k} \partial_{zz} Z_{3}^{l} Z_{h}^{\tilde{\alpha}} u + (c_{l,\varphi}^{j})' c_{j,\varphi}^{k} \partial_{z} Z_{3}^{l} Z_{h}^{\tilde{\alpha}} u \right)$$

$$+ \nu \sum_{l=0}^{k-1} \left(c_{l,\varphi}^{k} \partial_{zz} Z_{3}^{l} Z_{h}^{\tilde{\alpha}} u + (c_{l,\varphi}^{k})' \partial_{z} Z_{3}^{l} Z_{h}^{\tilde{\alpha}} u \right).$$
(3.26)

Note that the highest-order term in (3.26), i.e., (j, l) = (k - 1, j) for the first sum and l = k - 1 for the second sum, is $2\nu c_{k-1,\varphi}^k \partial_{zz} Z_3^{k-1} Z_{\mathbf{h}}^{\tilde{\alpha}} u,$

and when we integrate it against $Z^{\alpha}u$, we obtain

$$2\nu \int c_{k-1,\varphi}^{k} \partial_{zz} Z_{3}^{k-1} Z_{h}^{\tilde{\alpha}} u \, ds = -2\nu \int \partial_{z} Z_{3}^{k-1} Z_{h}^{\tilde{\alpha}} u \left(c_{k-1,\varphi}^{k} \partial_{z} Z^{\alpha} u + (c_{k-1,\varphi}^{k})' Z^{\alpha} u \right) \, ds$$

$$\lesssim \nu \| \partial_{z} Z_{3}^{k-1} Z_{h}^{\tilde{\alpha}} u \|_{L^{2}} (\| \partial_{z} Z^{\alpha} u \|_{L^{2}} + \| u \|_{4})$$

$$\leq C_{\epsilon} \nu \| \partial_{z} Z_{3}^{k-1} Z_{h}^{\tilde{\alpha}} u \|_{L^{2}}^{2} + \epsilon \nu (\| \partial_{z} Z^{\alpha} u \|_{L^{2}}^{2} + \| u \|_{4}^{2}), \qquad (3.27)$$

where $\epsilon > 0$ is an arbitrarily small parameter. Repeating this procedure for the remaining terms in (3.26) yields

$$(\nu Z^{\alpha} \Delta u - \nu \Delta Z^{\alpha} u, Z^{\alpha} u) \le \epsilon \nu \|\partial_z Z^{\alpha} u\|_{L^2}^2 + 2C_{\epsilon} \nu \|\nabla u\|_3^2 + \|u\|_4^2.$$
(3.28)

Therefore, collecting (3.16), (3.17), (3.21), (3.25), (3.28), absorbing all the factors of $\nu \|\partial_z Z^{\alpha} u\|_{L^2}^2$, and integrating in time, we obtain

$$\begin{aligned} \|Z^{\alpha}u(t)\|_{L^{2}}^{2} + c_{0}\nu \int_{0}^{t} \|\nabla Z^{\alpha}u(s)\|_{L^{2}}^{2} ds \\ \lesssim \|Z^{\alpha}u_{0}\|_{L^{2}}^{2} + \int_{0}^{t} (\nu\|\nabla u(s)\|_{3}^{2} + \|u(s)\|_{4}^{2} (\|u(s)\|_{W^{1,\infty}} + \|u(s)\|_{2,\infty} + 1) + \|u(s)\|_{4} \|\nabla p(s)\|_{3}) ds. \end{aligned}$$
(3.29)

Note that the inductive hypothesis (3.6) implies

$$\nu \int_{0}^{t} \|\nabla u(s)\|_{3}^{2} ds \lesssim \int_{0}^{t} \left(\|u(s)\|_{3}^{2} (\|u(s)\|_{W^{1,\infty}} + \|u(s)\|_{2,\infty} + 1) + \|u(s)\|_{3} \|\nabla p(s)\|_{2} \right) ds.$$
tly, using this in (3.29) and summing with (3.6) yields (3.1).

Consequently, using this in (3.29) and summing with (3.6) yields (3.1).

4. NORMAL DERIVATIVE ESTIMATES

In this section, our goal is to estimate $\partial_z u$ in $H^2_{co}(\Omega)$. However, as in [MR1], instead of studying this term directly, it is more convenient to work with

$$\eta = \omega_{\rm h} - 2\mu u_{\rm h}^{\perp},\tag{4.1}$$

where $\omega = \operatorname{curl} u$ and $u_{\mathbf{h}}^{\perp} = (-u_2, u_1)^T$. Observe that η solves

$$\eta_t - \nu \Delta \eta + u \cdot \nabla \eta = \omega \cdot \nabla u_h + 2\mu \nabla_h^{\perp} p, \qquad x \in \Omega$$

$$\eta = 0, \qquad z = 0,$$
(4.2)

where $\nabla_{\mathbf{h}}^{\perp} p = (-\partial_2, \partial_1)^T$. The main advantage of analyzing η , rather than ω , is that it vanishes on the boundary. Moreover,

$$\|\partial_z u\|_m \lesssim \|\eta\|_m + \|u\|_{m+1}, \qquad \|\partial_z u\|_{L^{\infty}} \lesssim \|\eta\|_{L^{\infty}} + \|u\|_{1,\infty}, \tag{4.3}$$

showing that the first-order normal derivative of u is controlled by the conormal derivatives of u and η . Due to (4.3), we may replace the term $||u||_{W^{1,\infty}}$ in (3.1) by $||\eta||_{L^{\infty}}$. We are now ready to state the main result of this section.

Proposition 4.1. Let u be a smooth solution of (2.1)-(2.2) on [0,T]. Then we have the inequality

$$\begin{aligned} \|\eta(t)\|_{m}^{2} + c_{0}\nu \int_{0}^{t} \|\nabla\eta(s)\|_{m}^{2} ds \\ \lesssim \|\eta_{0}\|_{m}^{2} + \int_{0}^{t} (\|\eta\|_{m}^{2}(\|u\|_{2,\infty} + \|\eta\|_{L^{\infty}}) + \|\eta\|_{m}(\|u\|_{m+2}\|\eta\|_{L^{\infty}} + \|u\|_{m+1}\|u\|_{1,\infty} + \|\nabla p\|_{m}) ds \end{aligned}$$

$$(4.4)$$

on [0,T] for m = 1, 2, where $c_0 > 0$ is independent of ν .

We note that (4.4) for m = 2 is needed to prove the first part of Proposition 2.3, while m = 1 is needed for the second part.

Proof of Proposition 4.1. We proceed by induction on $0 \le |\alpha| \le 2$. When $|\alpha| = 0$, the standard L^2 estimates yield

$$\|\eta(t)\|_{L^{2}}^{2} + \nu \int_{0}^{t} \|\nabla\eta\|_{L^{2}}^{2} ds \lesssim \|\eta_{0}\|_{L^{2}}^{2} + \int_{0}^{t} (\|\omega(s)\|_{L^{\infty}} \|\nabla u(s)\|_{L^{2}} + \|\nabla p(s)\|_{L^{2}}) \|\eta(s)\|_{L^{2}} ds.$$
(4.5)

Using (4.1), we obtain

$$\|\omega_{\mathbf{h}}\|_{L^{\infty}} \lesssim \|\eta\|_{L^{\infty}} + \|u\|_{L^{\infty}} \text{ and } \|\omega_{\mathbf{h}}\|_{m} \lesssim \|\eta\|_{m} + \|u\|_{m}, \qquad m \in \mathbb{N}_{0},$$

$$(4.6)$$

while for the normal component of the vorticity we have

$$\|\omega_3\|_{L^{\infty}} \lesssim \|u\|_{1,\infty} \text{ and } \|\omega_3\|_m \lesssim \|u\|_{m+1}, \qquad m \in \mathbb{N}_0,$$

$$(4.7)$$

since $\omega_3 = \partial_1 u_2 - \partial_2 u_1$. Therefore, using (4.3), (4.6), and (4.7) in (4.5), we conclude

$$\begin{aligned} &\|\eta(t)\|_{L^{2}}^{2} + \nu \int_{0}^{t} \|\nabla\eta\|_{L^{2}}^{2} ds \\ &\lesssim \|\eta_{0}\|_{L^{2}}^{2} + \int_{0}^{t} (\|\eta\|_{L^{2}}^{2} (\|\eta\|_{L^{\infty}} \|u\|_{1,\infty}) + \|\eta\|_{L^{2}} (\|u\|_{1} \|\eta\|_{L^{\infty}} + \|u\|_{1} \|u\|_{1,\infty} + \|\nabla p\|_{L^{2}}) ds. \end{aligned}$$

It remains to establish similar inequalities for $Z^{\alpha}\eta$, where $|\alpha| = 1, 2$. We only present our estimates for $|\alpha| = 2$, assuming (4.4) for m = 1. We first consider the case with only tangential derivatives.

Step 1. The case $Z^{\alpha} = Z_{\rm h}^{\tilde{\alpha}}$.

Applying $Z_{\rm h}^{\tilde{\alpha}}$ to (4.2)₁, we obtain

$$Z_{h}^{\tilde{\alpha}}\eta_{t} - \nu\Delta Z_{h}^{\tilde{\alpha}}\eta + u \cdot \nabla Z_{h}^{\tilde{\alpha}}\eta = Z_{h}^{\tilde{\alpha}}(\omega \cdot \nabla u_{h}) + 2\mu Z_{h}^{\tilde{\alpha}}\nabla_{h}^{\perp}p + u \cdot \nabla Z_{h}^{\tilde{\alpha}}\eta - Z_{h}^{\tilde{\alpha}}(u \cdot \nabla \eta).$$

$$(4.8)$$

Upon testing with $Z_{\rm h}^{\tilde{\alpha}}\eta$, the left-hand side of (4.8) gives

$$\frac{1}{2} \|Z_{h}^{\tilde{\alpha}}\eta\|_{L^{2}}^{2} + \nu \|\nabla Z_{h}^{\tilde{\alpha}}\eta\|_{L^{2}}^{2}$$
(4.9)

since there is no boundary term thanks to $(4.2)_2$. For the right-hand side of (4.8), the pressure term yields

$$(2\mu Z_{\mathbf{h}}^{\bar{\alpha}} \nabla_{\mathbf{h}}^{\perp} p, Z_{\mathbf{h}}^{\bar{\alpha}} \eta) \lesssim \|\nabla p\|_2 \|\eta\|_2, \tag{4.10}$$

since $|\alpha| = 2$. For the term involving vorticity, we use (2.11) and write

$$\begin{aligned} (Z_{\mathbf{h}}^{\tilde{\alpha}}(\omega \cdot \nabla u_{\mathbf{h}}), Z_{\mathbf{h}}^{\tilde{\alpha}}\eta) \lesssim & \|Z_{\mathbf{h}}^{\tilde{\alpha}}(\omega \cdot \nabla u_{\mathbf{h}})\|_{L^{2}} \|\eta\|_{2} \lesssim (\|\omega\|_{L^{\infty}} \|\nabla u_{\mathbf{h}}\|_{2} + \|\omega\|_{2} \|\nabla u_{\mathbf{h}}\|_{L^{\infty}}) \|\eta\|_{2} \\ \lesssim & \|\eta\|_{2}^{2} (\|\eta\|_{L^{\infty}} + \|u\|_{1,\infty}) + \|\eta\|_{2} \|u\|_{3} (\|\eta\|_{L^{\infty}} + \|u\|_{1,\infty}), \end{aligned}$$

$$(4.11)$$

where, in the last inequality we have used (4.6) and (4.7). Finally, for the commutator term in (4.8), we as in (3.11) first separate the normal and tangential derivatives. Then, we divide and multiply by φ the terms involving ∂_z obtaining

$$I_{\rm com} = u \cdot \nabla Z_{\rm h}^{\tilde{\alpha}} \eta - Z_{\rm h}^{\tilde{\alpha}} (u \cdot \nabla \eta) = \sum_{1 \le |\tilde{\beta}| \le 2} \begin{pmatrix} \tilde{\alpha} \\ \tilde{\beta} \end{pmatrix} \left(Z_{\rm h}^{\tilde{\beta}} u_{\rm h} \cdot \nabla_{\rm h} Z_{\rm h}^{\tilde{\alpha} - \tilde{\beta}} \eta + Z_{\rm h}^{\tilde{\beta}} \frac{u_3}{\varphi} Z_3 Z_{\rm h}^{\tilde{\alpha} - \tilde{\beta}} \eta \right).$$
(4.12)

Once again, we use (2.11) and estimate the sum in (4.12)

$$(I_{\text{com}}, Z_{h}^{\tilde{\alpha}}\eta) \lesssim \left(\|Z_{h}u\|_{L^{\infty}} + \left\|Z_{h}\frac{u_{3}}{\varphi}\right\|_{L^{\infty}} \right) \|\eta\|_{2}^{2} + \left(\|Z_{h}u\|_{2} + \left\|Z_{h}\frac{u_{3}}{\varphi}\right\|_{2} \right) \|\eta\|_{L^{\infty}} \|\eta\|_{2} \lesssim \|\eta\|_{2}^{2} \|u\|_{2,\infty} + \|\eta\|_{2} \|u\|_{4} \|\eta\|_{L^{\infty}},$$

$$(4.13)$$

where we have also used (2.13). Now, collecting (4.9)–(4.13) and integrating in time, it follows that

$$\begin{aligned} \|Z_{\mathbf{h}}^{\tilde{\alpha}}\eta(t)\|_{L^{2}}^{2} + \nu \int_{0}^{t} \|\nabla Z_{\mathbf{h}}^{\tilde{\alpha}}\eta\|_{L^{2}}^{2} ds \\ \lesssim \int_{0}^{t} (\|\eta\|_{2}^{2}(\|\eta\|_{L^{\infty}} + \|u\|_{2,\infty}) + \|\eta\|_{2}(\|u\|_{4}\|\eta\|_{L^{\infty}} + \|u\|_{3}\|u\|_{1,\infty} + \|\nabla p\|_{2})) ds, \end{aligned}$$

$$(4.14)$$

concluding Step 1.

Step 2. We now consider $Z^{\alpha}=Z_{\mathrm{h}}^{\tilde{\alpha}}Z_{3}^{k},$ for $1\leq k\leq |\alpha|\leq 2.$

Applying Z^{α} to the (4.2)₁, we obtain

$$Z^{\alpha}\eta_{t} - \nu\Delta Z^{\alpha}\eta + u \cdot \nabla Z^{\alpha}\eta = Z^{\alpha}(\omega \cdot \nabla u_{h}) + 2\mu Z^{\alpha} \nabla^{\perp}_{h} p + u \cdot \nabla Z^{\alpha}\eta - Z^{\alpha}(u \cdot \nabla \eta) - \nu Z^{\alpha} \Delta \eta + \nu\Delta Z^{\alpha} \eta.$$
(4.15)

Upon testing with $Z^{\alpha}\eta$, the left-hand side of the above equation gives

$$\frac{1}{2}\frac{d}{dt}\|Z^{\alpha}\eta\|_{L^{2}}^{2} + \nu\|\nabla Z^{\alpha}\eta\|_{L^{2}}^{2}, \qquad (4.16)$$

since there is no boundary term due to the presence of Z_3 . Now, for the vorticity and the pressure terms, we repeat (4.10) and (4.11), obtaining

$$(Z^{\alpha}(\omega \cdot \nabla u_{\mathbf{h}}) + 2\mu Z^{\alpha} \nabla_{\mathbf{h}}^{\perp} p, Z^{\alpha} \eta) \lesssim \|\eta\|_{2}^{2} (\|\eta\|_{L^{\infty}} + \|u\|_{1,\infty}) + \|\eta\|_{2} (\|u\|_{3}\|\eta\|_{L^{\infty}} + \|u\|_{3}\|u\|_{1,\infty} + \|\nabla p\|_{2}).$$

$$(4.17)$$

The first commutator term in (4.15) is expanded as in (3.18) and (3.20) by writing

$$\begin{aligned} u \cdot \nabla Z^{\alpha} \eta - Z^{\alpha} (u \cdot \nabla \eta) &= u_{3} \partial_{z} Z^{\alpha} \eta - u_{3} Z^{\alpha} \partial_{z} \eta - \sum_{1 \leq |\beta| \leq |\alpha|} \binom{\alpha}{\beta} Z^{\beta} u \cdot Z^{\alpha - \beta} \nabla \eta \\ &= -\sum_{j=0}^{k-1} \tilde{c}_{j,\varphi}^{k} u_{3} \partial_{z} Z_{h}^{\tilde{\alpha}} Z_{3}^{j} \eta - \sum_{1 \leq |\beta| \leq |\alpha|} \binom{\alpha}{\beta} Z^{\beta} u \cdot Z^{\alpha - \beta} \nabla \eta \\ &= -\sum_{j=0}^{k-1} \tilde{c}_{j,\varphi}^{k} u_{3} \partial_{z} Z_{h}^{\tilde{\alpha}} Z_{3}^{j} \eta - \sum_{1 \leq |\beta| \leq |\alpha|} \binom{\alpha}{\beta} Z^{\beta} u_{h} \cdot \nabla_{h} Z^{\alpha - \beta} \eta \\ &- \sum_{1 \leq |\beta| \leq |\alpha|} \binom{\alpha}{\beta} Z^{\beta} u_{3} Z^{\alpha - \beta} \partial_{z} \eta \\ &= I_{1} + I_{2} + I_{3}. \end{aligned}$$
(4.18)

We divide and multiply I_1 by φ and estimate it as

$$(I_1, Z^{\alpha} \eta) \lesssim \left\| \frac{u_3}{\varphi} \right\|_{L^{\infty}} \|\eta\|_2^2 \lesssim \|\eta\|_2^2 \|u\|_{1,\infty}.$$

$$(4.19)$$

Next, we bound I_2 as

$$(I_2, Z^{\alpha} \eta) \lesssim \|\eta\|_2^2 \|u\|_{2,\infty}.$$
(4.20)

Lastly, for I_3 , we first divide and multiply by φ and introduce lower order terms. It follows that

$$Z^{\beta}u_{3}Z^{\alpha-\beta}\partial_{z}\eta = Z^{\beta}\frac{u_{3}}{\varphi}Z^{\alpha-\beta}Z_{3}\eta + I_{l}, \qquad (4.21)$$

where I_l stands for the lower order terms. We now proceed as in (4.13), obtaining through (2.11)

$$(I_3, Z^{\alpha}\eta) \lesssim \|\eta\|_2^2 \|u\|_{2,\infty} + \|\eta\|_2 \|u\|_4 \|\eta\|_{L^{\infty}}.$$
(4.22)

Collecting (4.18)–(4.22), we have the bound

$$(u \cdot \nabla Z^{\alpha} \eta - Z^{\alpha} (u \cdot \nabla \eta), Z^{\alpha} \eta) \lesssim \|\eta\|_{2}^{2} \|u\|_{2,\infty} + \|\eta\|_{2} \|u\|_{4} \|\eta\|_{L^{\infty}}.$$
(4.23)

Now, it only remains to estimate the commutator term involving the Laplacian. We use Lemma 2.1 to write

$$\nu Z^{\alpha} \Delta \eta - \nu \Delta Z^{\alpha} \eta = \nu \sum_{j=0}^{k-1} \sum_{l=0}^{j} \left(c_{l,\varphi}^{j} c_{j,\varphi}^{k} \partial_{zz} Z_{3}^{l} Z_{h}^{\tilde{\alpha}} \eta + (c_{l,\varphi}^{j})' c_{j,\varphi}^{k} \partial_{z} Z_{3}^{l} Z_{h}^{\tilde{\alpha}} \eta \right) + \nu \sum_{l=0}^{k-1} \left(c_{l,\varphi}^{k} \partial_{zz} Z_{3}^{l} Z_{h}^{\tilde{\alpha}} \eta + (c_{l,\varphi}^{k})' \partial_{z} Z_{3}^{l} Z_{h}^{\tilde{\alpha}} \eta \right),$$

$$(4.24)$$

with the highest-order term

$$2\nu c_{k-1,\varphi}^k \partial_{zz} Z_3^{k-1} Z_h^{\tilde{\alpha}} \eta$$

When we integrate this term against $Z^{\alpha}\eta$ and repeat the steps in (3.27), we obtain

$$(2\nu c_{k-1,\varphi}^{k}\partial_{zz}Z_{3}^{k-1}Z_{h}^{\tilde{\alpha}}\eta, Z^{\alpha}\eta) \leq C_{\epsilon}\nu \|\partial_{z}Z_{3}^{k-1}Z_{h}^{\tilde{\alpha}}\eta\|_{L^{2}}^{2} + \epsilon\nu(\|\partial_{z}Z^{\alpha}\eta\|_{L^{2}}^{2} + \|\eta\|_{2}^{2}),$$
(4.25)

with $\epsilon > 0$ arbitrarily small. Dealing with the remaining terms in (4.24), we get

$$(\nu Z^{\alpha} \Delta \eta - \nu \Delta Z^{\alpha} \eta, Z^{\alpha} \eta) \le \epsilon \nu \|\partial_{z} Z^{\alpha} \eta\|_{L^{2}}^{2} + C_{\epsilon}(\|\partial_{z} \eta\|_{1}^{2} + \|\eta\|_{2}^{2}).$$
(4.26)

Now combining (4.16), (4.17), (4.23), (4.26) and integrating in time, we obtain

$$\begin{aligned} \|Z^{\alpha}\eta(t)\|_{L^{2}}^{2} + c_{0}\nu \int_{0}^{t} \|\nabla\eta\|_{L^{2}}^{2} ds &\lesssim \|Z^{\alpha}\eta_{0}\|_{L^{2}}^{2} + \nu \int_{0}^{t} \|\partial_{z}\eta\|_{1}^{2} ds \\ &+ \int_{0}^{t} (\|\eta\|_{2}^{2}(\|\eta\|_{L^{\infty}} + \|u\|_{2,\infty}) + \|\eta\|_{2}(\|u\|_{4}\|\eta\|_{L^{\infty}} + \|u\|_{3}\|u\|_{1,\infty} + \|\nabla p\|_{2}) ds. \end{aligned}$$

$$(4.27)$$

Consequently, using the inductive hypothesis, i.e., (4.4) for m = 1 to control the normal derivative of η , we obtain (4.4).

5. PRESSURE ESTIMATES

In this section, we estimate the pressure term. We note that $\|\nabla p\|_3$ is needed for the conormal and normal estimates, while $\|D^2p\|_3$ is required to bound $\|u\|_{2,\infty}$.

Proposition 5.1. Let (u, p) be a smooth solution of (2.1)–(2.2). Then we have the inequality

$$\|D^2p\|_3 + \|\nabla p\|_3 \lesssim \|u\|_4 (\|u\|_{2,\infty} + \|\eta\|_{L^{\infty}} + 1) + \nu \|\nabla u\|_4,$$
(5.1)

where the implicit constant is independent of ν .

Observe that upon applying the divergence to (2.1), we obtain

$$-\Delta p = \partial_i u_j \partial_j u_i,\tag{5.2}$$

while on the boundary, using (2.2), we have

$$\nabla p \cdot n = -\partial_z p = -\nu \partial_{ii} u \cdot n = \nu \partial_{ii} u_3 = \nu \partial_{zz} u_3 = -\nu \partial_z \nabla_h \cdot u_h = -2\mu \nu \nabla_h \cdot u_h.$$
(5.3)

We start by treating $Z_{\rm h}^{\tilde{\alpha}}p$ and then we use the induction on k to cover the general case, recalling that $Z^{\alpha} = Z_{\rm h}^{\tilde{\alpha}} Z_3^k$.

Lemma 5.2. Let (u, p) be a smooth solution of (2.1)–(2.2). Then we have the inequality

$$\|D^{2}Z_{h}^{\alpha}p\|_{L^{2}} + \|\nabla Z_{h}^{\alpha}p\|_{L^{2}} \lesssim \|u\|_{4}(\|u\|_{2,\infty} + \|\eta\|_{L^{\infty}} + 1) + \nu\|\nabla u\|_{|\tilde{\alpha}|+1},$$
(5.4)

where $0 \leq |\tilde{\alpha}| \leq 3$.

Proof of Lemma 5.2. We first present our estimates for $|\tilde{\alpha}| = 3$. We start by applying $Z_{h}^{\tilde{\alpha}}$ to (5.2) and (5.3). It follows that $Z_{h}^{\tilde{\alpha}}p$ solves

$$-\Delta Z_{\mathbf{h}}^{\tilde{\alpha}} p = Z_{\mathbf{h}}^{\tilde{\alpha}} (\partial_i u_j \partial_j u_i),$$

with the boundary condition

$$\nabla Z_{\mathbf{h}}^{\tilde{\alpha}} p \cdot n = -2\mu\nu Z_{\mathbf{h}}^{\tilde{\alpha}} \nabla_{\mathbf{h}} \cdot u_{\mathbf{h}}.$$

Applying the H^2 elliptic estimate for the Neumann problem (see [G]), we obtain

$$\|D^{2}Z_{h}^{\tilde{\alpha}}p\|_{L^{2}} + \|\nabla Z_{h}^{\tilde{\alpha}}p\|_{L^{2}} \lesssim \|Z_{h}^{\tilde{\alpha}}(\partial_{i}u_{j}\partial_{j}u_{i})\|_{L^{2}} + 2\|\mu\|\nu\|Z_{h}^{\tilde{\alpha}}\nabla_{h}\cdot u_{h}\|_{H^{1}}.$$
(5.5)

Note that to estimate $Z_{h}^{\tilde{\alpha}}(\partial_{i}u_{j}\partial_{j}u_{i})$ it suffices to consider $Z_{h}^{\tilde{\alpha}}(Z_{h}u_{h}Z_{h}u_{h})$ and $Z_{h}^{\tilde{\alpha}}(Z_{h}u_{3}\partial_{z}u_{h})$ since we may use the divergence-free condition to eliminate the normal derivatives in $\partial_{z}u_{3}\partial_{z}u_{3}$. For the first case, we use (2.11) and obtain

$$|Z_{h}^{\alpha}(Z_{h}u_{h}Z_{h}u_{h})||_{L^{2}} \lesssim ||u||_{4}||u||_{1,\infty}$$

while for the second case, we first expand by writing

$$Z_{h}^{\tilde{\alpha}}(Z_{h}u_{3}\partial_{z}u_{h}) = \sum_{0 \le |\tilde{\beta}| \le |\tilde{\alpha}|} {\alpha \choose \beta} Z_{h}^{\tilde{\beta}}Z_{h}u_{3}Z_{h}^{\tilde{\alpha}-\tilde{\beta}}\partial_{z}u_{h}.$$
(5.6)

When $|\tilde{\beta}| = 3$, we obtain

$$\|Z_{\mathbf{h}}^{\tilde{\beta}}Z_{\mathbf{h}}u_{3}\partial_{z}Z_{\mathbf{h}}^{\tilde{\alpha}-\tilde{\beta}}u_{\mathbf{h}}\|_{L^{2}} \lesssim \|u\|_{4}(\|\eta\|_{L^{\infty}}+\|u\|_{1,\infty}).$$
(5.7)

For the values of $|\tilde{\beta}| \neq 3$, we divide and multiply by φ and write

$$\left| Z_{\mathbf{h}}^{\tilde{\beta}} Z_{\mathbf{h}} \frac{u_{3}}{\varphi} Z_{3} Z_{\mathbf{h}}^{\tilde{\alpha} - \tilde{\beta}} u_{\mathbf{h}} \right\|_{L^{2}} \lesssim \begin{cases} \left\| Z_{\mathbf{h}} \frac{u_{3}}{\varphi} \right\|_{L^{\infty}} \| Z_{\mathbf{h}}^{\alpha - \beta} Z_{3} u \|_{L^{2}} \lesssim \| u \|_{4} \| u \|_{2,\infty}, & |\beta| = 0 \\ \left(\left\| Z_{\mathbf{h}} \frac{u_{3}}{\varphi} \right\|_{L^{\infty}} \| Z_{\mathbf{h}} Z_{3} u \|_{2} + \left\| Z_{\mathbf{h}} \frac{u_{3}}{\varphi} \right\|_{2} \| Z_{\mathbf{h}} Z_{3} u \|_{L^{\infty}} \right) \lesssim \| u \|_{4} \| u \|_{2,\infty}, & |\tilde{\beta}| = 1 \quad (5.8) \\ \| u \|_{4} \| u \|_{2,\infty}, & |\tilde{\beta}| = 2, \end{cases}$$

where, for $|\tilde{\beta}| = 1$, we have used (2.11) by taking f and g as $Z_h \frac{u_3}{\varphi}$ and $Z_h Z_3 u$, respectively. Now, it only remains to estimate the last term in (5.5), where we have

$$\nu \| Z_{h}^{\alpha} \nabla_{h} \cdot u_{h} \|_{H^{1}} \lesssim \nu \| u \|_{4} + \nu \| \nabla u \|_{|\tilde{\alpha}|+1}.$$
(5.9)

As a consequence, combining (5.5)–(5.9) gives (5.4), for $|\tilde{\alpha}| = 3$. When we consider a lower order conormal derivative, i.e., $|\tilde{\alpha}| \leq 2$, the only difference in our estimates is the treatment of the boundary term. Therefore, upon performing (5.5)–(5.9), for $|\tilde{\alpha}| \leq 2$, we may conclude (5.4).

Now, we move on to the general case.

Proof of Proposition 5.1. We present our estimates for $||D^2 Z^{\alpha} p||_{L^2}$ and $||\nabla Z^{\alpha} p||_{L^2}$ with $|\alpha| = 3$, using induction on $0 \le k \le 3$ satisfying $k + |\tilde{\alpha}| = 3$. The case k = 0 follows by using Lemma 5.2. Thus, we assume

$$\begin{split} \|D^{2}Z_{h}^{\beta}Z_{3}^{k}p\|_{L^{2}} + \|\nabla Z_{h}^{\beta}Z_{3}^{k}p\|_{L^{2}} \lesssim \|u\|_{4}(\|u\|_{2,\infty} + \|\eta\|_{L^{\infty}} + 1) \\ + \nu\|\nabla \nabla_{h}Z_{h}^{\tilde{\beta}}u\|_{L^{2}}, \qquad |\tilde{\beta}| + k = 3 \end{split}$$
(5.10)

and aim to prove that

$$\begin{split} \|D^{2} Z_{h}^{\tilde{\alpha}} Z_{3}^{k+1} p\|_{L^{2}} + \|\nabla Z_{h}^{\tilde{\alpha}} Z_{3}^{k+1} p\|_{L^{2}} \lesssim \|u\|_{4} (\|u\|_{2,\infty} + \|\eta\|_{L^{\infty}} + 1) \\ + \nu \|\nabla \nabla_{h} Z_{h}^{\tilde{\alpha}} u\|_{L^{2}}, \qquad |\tilde{\alpha}| + k + 1 = 3. \end{split}$$
(5.11)

Now, applying $Z^{\alpha} = Z_{h}^{\tilde{\alpha}} Z_{3}^{k+1}$ to (5.2), it follows that

$$-\Delta Z^{\alpha} p = Z^{\alpha} (\partial_i u_j \partial_j u_i) + Z^{\alpha} \Delta p - \Delta Z^{\alpha} p.$$

To find the boundary condition, we write

$$\nabla Z^{\alpha} p \cdot n = -\partial_z Z_{\mathbf{h}}^{\tilde{\alpha}} Z_3^{k+1} p = -Z_{\mathbf{h}}^{\tilde{\alpha}} \sum_{j=0}^k \tilde{c}_{j,\varphi}^{k+1} Z_3^j \partial_z p = -\tilde{c}_{0,\varphi}^{k+1} Z_{\mathbf{h}}^{\tilde{\alpha}} \partial_z p = -2\mu\nu \tilde{c}_{0,\varphi}^{k+1} Z_{\mathbf{h}}^{\tilde{\alpha}} \nabla_{\mathbf{h}} \cdot u_{\mathbf{h}},$$

where we have used (5.3) and the fact that Z_3 vanishes on the boundary. Applying the elliptic estimates, as in (5.9), we obtain

$$\|D^{2}Z^{\alpha}p\|_{L^{2}} + \|\nabla Z^{\alpha}p\|_{L^{2}} \lesssim \|Z^{\alpha}(\partial_{i}u_{j}\partial_{j}u_{i})\|_{L^{2}} + \|Z^{\alpha}\Delta p - \Delta Z^{\alpha}p\|_{L^{2}} + \|\mu|\nu\|Z^{\tilde{\alpha}}_{h}\nabla_{h} \cdot u_{h}\|_{H^{1}}.$$
(5.12)

For the first term on the right-hand side of the above inequality, we introduce lower order terms when necessary and use the estimates in (5.7) and (5.8) to conclude that

$$\|Z^{\alpha}(\partial_{i}u_{j}\partial_{j}u_{i})\|_{L^{2}} + |\mu|\nu\|Z_{h}^{\tilde{\alpha}}\nabla_{h} \cdot u_{h}\|_{H^{1}} \lesssim \|u\|_{4}(\|u\|_{2,\infty} + \|\eta\|_{L^{\infty}} + 1) + \nu\|\nabla\nabla_{h}Z_{h}^{\tilde{\alpha}}u\|_{L^{2}}.$$
(5.13)

Therefore, it only remains to estimate the commutator term for the pressure. Using Lemma 2.1, we may expand this term as

$$\begin{split} Z^{\alpha}\Delta p - \Delta Z^{\alpha}p &= Z^{\alpha}\partial_{zz}p - \partial_{zz}Z^{\alpha}p = Z^{\alpha}\partial_{zz}p - \sum_{j=0}^{k+1}\sum_{l=0}^{j}\tilde{c}_{l,\varphi}^{j}\tilde{c}_{j,\varphi}^{k+1}Z_{3}^{l}Z_{h}^{\tilde{\alpha}}\partial_{zz}p + \sum_{j=0}^{k+1}(\tilde{c}_{j,\varphi}^{k+1})'Z_{3}^{j}Z_{h}^{\tilde{\alpha}}\partial_{z}p \\ &= -\sum_{j=0}^{k}\sum_{l=0}^{j}\tilde{c}_{l,\varphi}^{j}\tilde{c}_{j,\varphi}^{k+1}Z_{3}^{l}Z_{h}^{\tilde{\alpha}}\partial_{zz}p - \sum_{l=0}^{k}\tilde{c}_{l,\varphi}^{k+1}Z_{3}^{l}Z_{h}^{\tilde{\alpha}}\partial_{zz}p + \sum_{j=0}^{k}(\tilde{c}_{j,\varphi}^{k+1})'Z_{3}^{j}Z_{h}^{\tilde{\alpha}}\partial_{z}p \\ &= I_{1}+I_{2}+I_{3}, \end{split}$$

where we have used that $(\tilde{c}_{k+1,\varphi}^{k+1})' = (1)' = 0$. We shall only treat the highest-order terms in the above equation. Observe that for I_3 this term is $\partial_z Z_h^{\tilde{\alpha}} Z_3^k p$ which is of lower order and is thus bounded as in (5.10). The highest-order terms of I_1 and I_2 are both equal to $\tilde{c}_{k,\varphi}^{k+1} Z_h^{\tilde{\alpha}} Z_3^k \partial_{zz} p$. Using (5.2), we rewrite this term as

$$\tilde{c}_{k,\varphi}^{k+1} Z_{\mathbf{h}}^{\tilde{\alpha}} Z_{\mathbf{h}}^{3} \partial_{zz} p = -\tilde{c}_{k,\varphi}^{k+1} Z_{\mathbf{h}}^{\tilde{\alpha}} Z_{\mathbf{h}}^{3} \Delta_{\mathbf{h}} p - \tilde{c}_{k,\varphi}^{k+1} Z_{\mathbf{h}}^{\tilde{\alpha}} Z_{\mathbf{h}}^{3} (\partial_{i} u_{j} \partial_{j} u_{i})$$

Since $Z_{h}^{\tilde{\alpha}}Z_{3}^{k}(\partial_{i}u_{j}\partial_{j}u_{i})$ is of lower order, it may be estimated as in (5.7) and (5.8). For $Z_{h}^{\tilde{\alpha}}Z_{3}^{k}\Delta_{h}p$, observe that

$$\|Z_{h}^{\tilde{\alpha}}Z_{3}^{k}\Delta_{h}p\|_{L^{2}} \lesssim \|D^{2}Z_{h}^{\tilde{\alpha}}Z_{3}^{k}p\|_{L^{2}} \lesssim \|u\|_{4}(\|u\|_{2,\infty} + \|\eta\|_{L^{\infty}} + 1) + \nu\|\nabla\nabla_{h}Z_{h}^{\tilde{\alpha}}u\|_{L^{2}},$$

thanks to the inductive hypothesis (5.10). Repeating these estimates for the remaining terms in I_1 , I_2 , and I_3 , we conclude that

$$\|Z^{\alpha}\Delta p - \Delta Z^{\alpha}p\|_{L^{2}} \lesssim \|u\|_{4}(\|u\|_{2,\infty} + \|\eta\|_{L^{\infty}} + 1) + \nu\|\nabla u\|_{4}.$$
(5.14)

Therefore, by (5.12), (5.13), and (5.14), we obtain (5.11), concluding the proof of Proposition 5.1.

6. L^{∞} estimates

In this section, we estimate $\|\eta\|_{L^{\infty}}$ and $\|u\|_{2,\infty}$.

Proposition 6.1. Let u be a smooth solution of (2.1)–(2.2). Then we have the following statements.

i. If $\mu \in \mathbb{R}$, we have

$$\begin{aligned} \|u(t)\|_{2,\infty}^{2} + \|\eta(t)\|_{L^{\infty}}^{2} \lesssim \|u_{0}\|_{2,\infty}^{2} + \|\eta_{0}\|_{L^{\infty}}^{2} + \int_{0}^{t} (\|u\|_{2,\infty} + \|\eta\|_{L^{\infty}} + 1)^{3} + \|u\|_{2,\infty} \|D^{2}p\|_{3} \, ds \\ + \nu \int_{0}^{t} (\|\nabla\eta\|_{2}^{2} + \|\nabla u\|_{4}^{2}) \, ds, \end{aligned}$$

$$(6.1)$$

where the implicit constant is independent of ν .

ii. If $\mu \geq 0$, we have

$$\begin{aligned} \|u(t)\|_{2,\infty}^{2} + \|\eta(t)\|_{L^{\infty}}^{2} \lesssim \|u_{0}\|_{2,\infty}^{2} + \|\eta_{0}\|_{L^{\infty}}^{2} + \int_{0}^{t} (\|u\|_{2,\infty} + \|\eta\|_{L^{\infty}} + 1)^{3} + \|u\|_{2,\infty} \|D^{2}p\|_{3} \, ds \\ + \nu \int_{0}^{t} (\|\nabla\eta\|_{1}^{2} + \|\nabla u\|_{4}^{2}) \, ds, \end{aligned}$$

$$(6.2)$$

where the implicit constant is independent of ν .

Proof of Proposition 6.1. First, the maximum principle for (4.2) implies

$$\frac{d}{dt} \|\eta\|_{L^{\infty}} \lesssim \|\omega\|_{L^{\infty}} \|\nabla u\|_{L^{\infty}} + \|\nabla_{\mathbf{h}}p\|_{L^{\infty}} \lesssim (\|\eta\|_{L^{\infty}} + \|u\|_{1,\infty})^{2} + \|p\|_{1,\infty},$$
(6.3)

where we have used (4.3), (4.6), and (4.7) in the last inequality. Note that the inequality independent of the sign of μ . Next, to treat the term $||u||_{2,\infty}$, we only consider the highest order derivative and note that lower order derivatives are bounded using the same methods. Now let $Z^{\alpha} = Z_{h}^{\tilde{\alpha}} Z_{3}^{k}$ be a conormal derivative of order two, i.e., $|\tilde{\alpha}| + k = 2$. We apply Z^{α} to (2.1) and write

$$Z^{\alpha}u_{t} - \nu\Delta Z^{\alpha}u + u \cdot \nabla Z^{\alpha}u = u \cdot \nabla Z^{\alpha}u - Z^{\alpha}(u \cdot \nabla u) - Z^{\alpha}\nabla p + \nu Z^{\alpha}\Delta u - \nu\Delta Z^{\alpha}u.$$
(6.4)

We complete the proof of Proposition 6.1 in two steps. First, we take $p \ge 4$, test the above equation with $Z^{\alpha}u|Z^{\alpha}u|^{p-2}$, and then pass to the limit as $p \to \infty$. In the second step, we estimate the terms that emerge in the limit.

Step 1. Test (6.4) with $Z^{\alpha}u|Z^{\alpha}u|^{p-2}$.

Upon integrating against $Z^{\alpha}u|Z^{\alpha}u|^{p-2}$, the left-hand side of (6.4) yields

$$\frac{1}{p}\frac{d}{dt}\|Z^{\alpha}u\|_{L^{p}}^{p} + \nu \int \left(\frac{(p-2)}{4}|\nabla|Z^{\alpha}u|^{2}|^{2}|Z^{\alpha}u|^{p-4} + |\nabla Z^{\alpha}u|^{2}|Z^{\alpha}u|^{p-2}\right)dx + 2\mu\nu\delta_{0k}\|Z^{\alpha}u_{h}\|_{L^{p}(\partial\Omega)}^{p}, \quad (6.5)$$

since the boundary term appears only for k = 0. When μ is non-negative, the last term in (6.5) is coercive. This is the only instance where the sign of μ affects the regularity assumption on η_0 . In other words, when $\mu \ge 0$, we are able to

close the estimates using $\eta_0 \in H^1_{co}(\Omega)$. On the other hand, when $\mu < 0$, we use the trace theorem and interpolation to obtain

$$\nu \| Z^{\alpha} u_{\mathbf{h}} \|_{L^{p}(\partial\Omega)}^{p} \lesssim \nu \| Z^{\alpha} u_{\mathbf{h}} \|_{W^{\frac{1}{p},p}}^{p} \lesssim \nu \| Z^{\alpha} u_{\mathbf{h}} \|_{L^{p}}^{\frac{(p+4)p}{p+6}} \| \nabla Z^{\alpha} u_{\mathbf{h}} \|_{L^{6}}^{\frac{2p}{6+p}}.$$
(6.6)

Recalling that $|\alpha| = 2$, it is evident from $\|\nabla Z^{\alpha} u_{h}\|_{L^{6}}$ that we need to control at least two conormal derivatives of η . Proceeding with the right hand side of (6.4), the first three terms give

$$(u \cdot \nabla Z^{\alpha} u - Z^{\alpha} (u \cdot \nabla u) - Z^{\alpha} \nabla p, Z^{\alpha} u | Z^{\alpha} u|^{p-2}) \lesssim \| u \cdot \nabla Z^{\alpha} u - Z^{\alpha} (u \cdot \nabla u) - Z^{\alpha} \nabla p \|_{L^{p}} \| Z^{\alpha} u \|_{L^{p}}^{p-1}, \quad (6.7)$$

which leaves us with the commutator term for the Laplacian, for which we have

$$\nu Z^{\alpha} \Delta u - \nu \Delta Z^{\alpha} u = \begin{cases} 0, & k = 0\\ -\nu (2\varphi' \partial_{zz} Z_{\mathbf{h}} u + \varphi'' \partial_{z} Z_{\mathbf{h}} u), & k = 1\\ -\nu (4\varphi' \partial_{zz} Z_{3} u - 4(\varphi')^{2} \partial_{zz} u + 4\varphi'' \partial_{z} Z_{3} u - 5\varphi' \varphi'' \partial_{z} u + \varphi''' Z_{3} u), & k = 2. \end{cases}$$
(6.8)

Note that when $k \ge 1$ the leading order term is of the form $-\nu \varphi' \partial_{zz} Zu$, and to treat this term we use

$$\partial_{zz}f = \frac{1}{\varphi}Z_3\partial_z f = \frac{1}{\varphi}\partial_z Z_3 f - \frac{\varphi'}{\varphi}\partial_z f.$$
(6.9)

Upon letting f = Zu, we obtain

$$-\nu \int \varphi' \partial_{zz} Z u Z_3 Z u |Z^{\alpha} u|^{p-2} dx = -\nu \int \frac{\varphi'}{\varphi} \partial_z Z_3 Z u Z_3 Z u |Z^{\alpha} u|^{p-2} dx + \nu \int \frac{(\varphi')^2}{\varphi} \partial_z Z u Z_3 Z u |Z^{\alpha} u|^{p-2} dx = I_1 + I_2.$$

$$(6.10)$$

We point out that $I_1 \leq 0$, and to show this, we write

$$I_{1} = -\frac{\nu}{p} \int \frac{\varphi'}{\varphi} \partial_{z} |Z^{\alpha}u|^{p} dx = \frac{\nu}{p} \int \frac{\varphi''\varphi - (\varphi')^{2}}{\varphi^{2}} |Z^{\alpha}u|^{p} dx$$

$$= \frac{\nu}{p} \int (\varphi''\varphi - (\varphi')^{2}) |\partial_{z}Zu|^{2} |Z^{\alpha}u|^{p-2} dx \leq 0,$$
(6.11)

where the last inequality follows from

$$\varphi''(z)\varphi(z) - (\varphi'(z))^2 < 0, \qquad z > 0$$

We proceed with I_2 by writing

$$I_{2} = \nu \int (\varphi')^{2} |\partial_{z} Z u|^{2} |Z^{\alpha} u|^{p-2} dx \lesssim \nu \|\partial_{z} Z u\|_{L^{p}}^{2} \|Z^{\alpha} u\|_{L^{p}}^{p-2}.$$
(6.12)

Going back to (6.8), the next term we treat is $\partial_{zz}u$. Upon applying (6.9) for f = u, it follows that

$$\nu \int (\varphi') \partial_{zz} u Z_3^2 u |Z_3^2 u|^{p-2} dx = \nu \int \frac{(\varphi')^2}{\varphi} \partial_z Z_3 u Z_3^2 u |Z_3^2 u|^{p-2} dx - \nu \int \frac{(\varphi')^3}{\varphi} \partial_z u \partial_z Z_3 u |Z_3^2 u|^{p-2} dx$$

$$= \nu \int (\varphi')^2 |\partial_z Z_3 u|^2 |Z_3^2 u|^{p-2} dx - \nu \int (\varphi')^3 \partial_z u \partial_z Z_3 u |Z_3^2 u|^{p-2} dx$$

$$\lesssim \nu (\|\partial_z Z_3 u\|_{L^p}^2 + \|\partial_z u\|_{L^p} \|\partial_z Z_3 u\|_{L^p}) \|Z^\alpha u\|_{L^p}^{p-2}.$$

(6.13)

We remark that the remaining terms in (6.8) are of lower order, and we estimate them using Hölder's inequality. Therefore, collecting (6.8), (6.10), (6.11), (6.12), and (6.13), it follows that

$$\nu \int (Z^{\alpha} \Delta u - \nu \Delta Z^{\alpha} u) Z^{\alpha} u |Z^{\alpha} u|^{2} dx \lesssim I_{1} + \nu \|Z^{\alpha} u\|_{L^{p}}^{p-2} (\|\partial_{z} Z u\|_{L^{p}}^{2} + \|\partial_{z} u\|_{L^{p}}\|\partial_{z} Z_{3} u\|_{L^{p}}) + \nu \|Z^{\alpha} u\|_{L^{p}}^{p-1} (\|\partial_{z} Z u\|_{L^{p}} + \|\partial_{z} u\|_{L^{p}} + \|Z_{3} u\|_{L^{p}}).$$
(6.14)

Now, thanks to (6.4)–(6.7), (6.14) and since $I_1 < 0$ we arrive at

$$\frac{1}{p} \frac{d}{dt} \| Z^{\alpha} u \|_{L^{p}}^{p} \lesssim \| Z^{\alpha} u \|_{L^{p}}^{p-1} (\| u \cdot \nabla Z^{\alpha} u - Z^{\alpha} (u \cdot \nabla u) - Z^{\alpha} \nabla p \|_{L^{p}}) \\
+ \nu \| Z^{\alpha} u \|_{L^{p}}^{p-1} (\| \partial_{z} Z u \|_{L^{p}} + \| \partial_{z} u \|_{L^{p}} + \| Z_{3} u \|_{L^{p}}) \\
+ \nu \| Z^{\alpha} u \|_{L^{p}}^{p-2} (\| \partial_{z} Z u \|_{L^{p}}^{2} + \| \partial_{z} u \|_{L^{p}} \| \partial_{z} Z_{3} u \|_{L^{p}}) \\
+ \nu \delta_{k0} \| Z^{\alpha} u_{h} \|_{L^{p}}^{\frac{(p+4)p}{p+6}} \| \nabla Z_{h}^{\tilde{\alpha}} u_{h} \|_{L^{6}}^{\frac{2p}{2}},$$

from where, canceling $\|Z^{\alpha}u\|_{L^p}^{p-2}$, letting $p \to \infty$, and integrating in time, we obtain

$$\begin{aligned} \|Z^{\alpha}u(t)\|_{L^{\infty}}^{2} \lesssim \|u_{0}\|_{2,\infty} + \int_{0}^{t} \|u\|_{2,\infty} (\|u \cdot \nabla Z^{\alpha}u - Z^{\alpha}(u \cdot \nabla u)\|_{L^{\infty}}) + \|Z^{\alpha}\nabla p\|_{L^{\infty}} + \|u\|_{W^{1,\infty}}) \\ + \nu \|u\|_{2,\infty} \|\partial_{z}u\|_{1,\infty} + \nu (\|\partial_{z}u\|_{1,\infty}^{2} + \|u\|_{W^{1,\infty}} \|\partial_{z}u\|_{1,\infty} + \delta_{k0} \|\nabla Z_{h}^{\tilde{\alpha}}u_{h}\|_{L^{6}}^{2}) \, ds. \end{aligned}$$

$$(6.15)$$

Step 2. We estimate the terms on the right hand side of (6.15).

We start with $\nu \|\partial_z u\|_{1,\infty}^2$. Utilizing (2.12), for $(s_1, s_2) = (0, 3)$, it follows that

$$\nu \int_0^t \|\partial_z u(s)\|_{1,\infty}^2 \, ds \lesssim \nu \int_0^t (\|\nabla \eta(s)\|_1^2 + \|\nabla u(s)\|_4^2) \, ds.$$
(6.16)

Next, for the term involving the horizontal velocity, recalling that it is present only when $\mu < 0$, we write

$$\nu \int_0^t \|\nabla Z_{\mathbf{h}}^{\tilde{\alpha}} u_{\mathbf{h}}(s)\|_{L^6}^2 \, ds \lesssim \nu \int_0^t (\|\nabla \eta\|_2^2 + \|\nabla u_{\mathbf{h}}(s)\|_4^2) \, ds, \tag{6.17}$$

where we have used the embedding $H^1 \subset L^6$. Next, we consider the commutator term by writing

$$u \cdot \nabla Z^{\alpha} u - Z^{\alpha} (u \cdot \nabla u) = -(1 - \delta_{k0}) \sum_{j=0}^{k-1} \tilde{c}_{j,\varphi}^k \frac{u_3}{\varphi} Z_3 Z_h^{\tilde{\alpha}} Z_3^j u - \sum_{1 \le |\beta| \le |\alpha|} \binom{\alpha}{\beta} Z^{\beta} u \cdot Z^{\alpha-\beta} \nabla u = J_1 + J_2, \quad (6.18)$$

from where it follows that J_1 satisfies

$$\|J_1\|_{L^{\infty}} \lesssim \|u\|_{1,\infty} \|u\|_{2,\infty},\tag{6.19}$$

while for J_2 we have

$$J_{2} = \sum_{1 \le |\tilde{\beta}| \le |\alpha|} \binom{\alpha}{\beta} (Z^{\beta} u_{\mathbf{h}} \cdot \nabla_{\mathbf{h}} Z^{\alpha-\beta} u + Z^{\beta} u_{3} Z^{\alpha-\beta} \partial_{z} u) = J_{21} + J_{22} \lesssim \|u\|_{1,\infty} \|u\|_{2,\infty} + J_{22}.$$
(6.20)

Upon introducing lower order terms when necessary, we obtain

$$\|Z^{\beta}u_{3}Z^{\alpha-\beta}\partial_{z}u\|_{L^{\infty}} \lesssim \begin{cases} \left\|Z^{\beta}\frac{u_{3}}{\varphi}\right\|_{\infty} \|Z^{\alpha-\beta}Z_{3}u\|_{L^{\infty}} \lesssim \|u\|_{2,\infty}^{2}, & |\beta| = 1\\ \|Z^{\beta}u_{3}\|_{L^{\infty}} \|Z^{\alpha-\beta}\partial_{z}u\|_{L^{\infty}} \lesssim \|u\|_{2,\infty}(\|\eta\|_{L^{\infty}} + \|u\|_{1,\infty}), & |\beta| = 2, \end{cases}$$
(6.21)

from where, by collecting (6.18)–(6.21) and integrating in time, it follows that

$$\int_{0}^{t} \|u(s) \cdot \nabla Z^{\alpha} u(s) - Z^{\alpha}(u(s) \cdot \nabla u(s))\|_{L^{\infty}} \|u\|_{2,\infty} ds \lesssim \int_{0}^{t} \|u\|_{2,\infty}^{2} (\|\eta\|_{L^{\infty}} + \|u\|_{2,\infty}) ds.$$
(6.22)

Going back to the right hand side of (6.15), it only remains to estimate the pressure term. Another application of (2.12) yields

$$\|\nabla p\|_{2,\infty} \lesssim \|\partial_z p\|_{2,\infty} + \|p\|_{3,\infty} \lesssim \|D^2 p\|_3^{\frac{1}{2}} \|\nabla p\|_4^{\frac{1}{2}} + \|D^2 p\|_3 \lesssim \|D^2 p\|_3.$$
(6.23)

To conclude, we multiply (6.3) by $\|\eta\|_{L^{\infty}}$ and integrate in time. Then, combining with (6.15) and considering (6.16), (6.17), (6.22), and (6.23) we obtain (6.1). When $\mu \ge 0$, we omit (6.17) and arrive at (6.2).

7. CONCLUDING THE A PRIORI ESTIMATES AND THE PROOFS OF THEOREMS 2.1 AND 2.3(I)

Note that combining (3.1) and (4.4) for m = 2 gives

$$\begin{aligned} u(t)\|_{5}^{2} + \|\eta(t)\|_{2}^{2} + c_{0}\nu \int_{0}^{t} (\|\nabla u(s)\|_{5}^{2} + \|\nabla \eta(s)\|_{2}^{2}) \, ds \\ \lesssim \|u_{0}\|_{5}^{2} + \|\eta_{0}\|_{2}^{2} + \int_{0}^{t} N^{3} \, ds + \int_{0}^{t} (\|\nabla p\|_{3}\|u\|_{4} + \|\nabla p\|_{2}\|\eta\|_{2}) \, ds, \end{aligned}$$

recalling that N is as in (2.17). Now, using the pressure estimates (5.1), we have

$$\begin{aligned} \|u(t)\|_{4}^{2} + \|\eta(t)\|_{2}^{2} + c_{0}\nu \int_{0}^{t} (\|\nabla u(s)\|_{4}^{2} + \|\nabla \eta(s)\|_{2}^{2}) \, ds \\ \lesssim N^{2}(0) + \int_{0}^{t} N^{3} \, ds + \int_{0}^{t} \nu \|\nabla u\|_{4} (\|u\|_{4} + \|\eta\|_{2}) \, ds. \end{aligned}$$

Therefore, upon absorbing the factors of $\nu \|\nabla u\|_4$, we obtain

$$\|u(t)\|_{5}^{2} + \|\eta(t)\|_{2}^{2} + c_{0}\nu \int_{0}^{t} (\|\nabla u(s)\|_{5}^{2} + \|\nabla \eta(s)\|_{2}^{2}) ds \lesssim N^{2}(0) + \int_{0}^{t} N^{3} ds.$$

$$(7.1)$$

Now, we use (7.1) in (6.1) so that we have

ŀ

$$\|u\|_{2,\infty}^2 + \|\eta\|_{L^{\infty}}^2 \lesssim N^2(0) + \int_0^t N^3 \, ds + \int_0^t \|u\|_{2,\infty} \|D^2 p\|_3 \, ds.$$

We may again use the pressure estimates (5.1) and absorb the emerging factors of $\nu \|\nabla u\|_4$. Then, we combine the resulting inequality with (7.1) which yields (2.17) and (2.18). To establish (2.19) and (2.20), we repeat these steps using (3.1), (4.4) for m = 1, (5.1), and (6.2). This concludes the proof of Proposition 2.3.

Now, having established the a priori estimates, we may prove Theorems 2.1 and 2.3(i) simultaneously. First, we may smoothen the initial data and use well-posedness results to obtain solutions to which Proposition 2.3 applies. Then, we can pass to the limit in (2.1)–(2.2) showing that the sequence of solutions is Cauchy in $L^{\infty}(0, T; L^{2}(\Omega))$. This step does not require $\nabla u \in H^{2}_{co}(\Omega)$ and hence it applies in both cases, i.e., when $\nabla u \in H^{1}_{co}(\Omega)$ and $\nabla u \in H^{1}_{co}(\Omega)$. Moreover, for each ν we have a unique solution thanks to the Lipschitz continuity of the initial data. Finally, employing the a priori estimates (2.17)–(2.20), the lifespans of both solutions, which may be different, are independent of ν .

8. PROOFS OF THEOREMS 2.2 AND 2.3(II)

We first prove that, under weaker regularity and boundedness assumptions, the sequence of solutions $\{u^{\nu}\}_{\nu}$ is Cauchy in $L^{\infty}(0,T; L^{2}(\Omega))$.

Proposition 8.1 (Compactness of $\{u^{\nu}\}_{\nu}$). Let $u_0 \in H^1(\Omega) \cap W^{1,\infty}(\Omega)$ be such that $\operatorname{div} u_0 = 0$, and $u_0 \cdot n = 0$ on $\partial\Omega$. Assume that there exists a T > 0 such that for all $\nu \in (0, 1)$ and $\mu \in \mathbb{R}$ fixed there exists a unique solution $u^{\nu} \in L^{\infty}(0, T; H^1(\Omega) \cap W^{1,\infty}(\Omega))$ to (2.1)–(2.2) on [0, T]. Moreover, assume that there exists a constant M > 0such that for all $\nu \in (0, 1)$

$$\sup_{[0,T]} (\|u^{\nu}(t)\|_{H^1}^2 + \|u^{\nu}(t)\|_{W^{1,\infty}}^2) + \nu \int_0^T \|D^2 u^{\nu}(s)\|_{L^2}^2 \, ds \le M.$$
(8.1)

Then, $\{u^{\nu}\}_{\nu}$ is a Cauchy sequence in $L^{\infty}(0,T;L^{2}(\Omega))$ with

$$\sup_{[0,T]} \|u^{\nu_1} - u^{\nu_2}\|_{L^2}^2 \lesssim \nu_1 + \nu_2, \tag{8.2}$$

where the implicit constant is independent of ν_1 and ν_2 .

Proof of Proposition 8.1. Let $\nu_1, \nu_2 \in (0, 1)$, and denote by (u^1, p^1) and (u^2, p^2) two solutions to (2.1)–(2.2), emanating from the same initial data, with the parameters ν_1 and ν_2 respectively. Upon letting $U = u_1 - u_2$, and recalling that $\mu \ge 0$ is fixed, we observe that the difference of solutions U satisfies

$$U_t - (\nu_1 - \nu_2)\Delta u^1 - \nu_2\Delta U + U \cdot \nabla u^1 + u^2 \cdot \nabla U + \nabla P = 0, \text{ and } \operatorname{div} U = 0,$$
(8.3)

where $P = p^1 - p^2$. This equation is coupled with the boundary conditions

$$U_3 = 0 \text{ and } \partial_z U_h = 2\mu U_h, \qquad (x,t) \in \{z=0\} \times (0,T).$$
(8.4)

Testing (8.3) with U and using (8.4), we obtain

$$\frac{1}{2}\frac{d}{dt}\|U\|_{L^{2}}^{2} + \nu_{2}\|\nabla U\|_{L^{2}}^{2} = (\nu_{2} - \nu_{1})\left(\int \nabla u^{1}\nabla U\,dx + 2\mu\int_{\partial\Omega}u_{h}^{1}U_{h}\,d\sigma\right) \\ - 2\mu\nu_{2}\|U_{h}\|_{L^{2}(\partial\Omega)}^{2} - \int U\cdot\nabla u^{1}U\,dx.$$
(8.5)

As a consequence of the uniform bounds, we may allow the implicit constant to depend on the upper bound C in (8.1) for the rest of this section. With this convention, (8.5) implies

$$\frac{d}{dt} \|U\|_{L^2}^2 \lesssim (\nu_1 + \nu_2)(\|U\|_{H^1} \|u^1\|_{H^1} + \|U\|_{H^1}^2) + \|U\|_{L^2}^2 \|\nabla u^1\|_{L^\infty} \lesssim (\nu_1 + \nu_2) + \|U\|_{L^2}^2,$$

from where, applying the Gronwall inequality on (0, T), we get

$$\sup_{[0,T]} \|U\|_{L^2}^2 \lesssim \nu_1 + \nu_2,$$

establishing (8.2).

We note that Proposition 8.1 implies that the sequence of solutions constructed in Theorems 2.1 and 2.3(i) is Cauchy in $L^{\infty}(0,T; L^2(\Omega))$. This is true since Proposition 8.1 holds under weaker regularity and boundedness assumptions and there is no restriction on the sign of μ . To show that (2.6) holds, we use that $\nabla u^{\nu} \in L^{\infty}$ is uniformly bounded and interpolate by writing

$$\|u^{1} - u^{2}\|_{L^{\infty}}^{2} \lesssim \|\nabla(u^{1} - u^{2})\|_{L^{\infty}}^{\frac{6}{5}} \|u^{1} - u^{2}\|_{L^{2}}^{\frac{4}{5}} \lesssim (\nu_{1} + \nu_{2})^{\frac{2}{5}}$$

where u^1 and u^2 are as in the proof of Proposition 8.1. Finally, passing to a subsequence concludes the proofs of Theorems 2.2 and 2.3(ii). The uniqueness follows from the Lipschitz continuity of the data.

9. EULER EQUATIONS IN SOBOLEV CONORMAL SPACES

In this section, we sketch the proof of Theorem 2.4. For some T > 0, we consider a smooth solution for (1.3)–(1.4) on $\Omega \times [0, T]$. We first note that, upon repeating the estimates leading to Proposition 3.1, and to Proposition 5.1 for $\nu = 0$, we obtain

$$\|u(t)\|_{4}^{2} \lesssim \|u_{0}\|_{4}^{2} + \int_{0}^{t} \left(\|u\|_{4}^{2}(\|\partial_{z}u\|_{W^{1,\infty}} + \|u\|_{2,\infty} + 1) + \|u\|_{4}\|\nabla p\|_{3}\right) ds,$$

$$(9.1)$$

and

$$\|D^2 p\|_3 + \|\nabla p\|_3 \lesssim \|u\|_4 (\|u\|_{2,\infty} + \|\partial_z u\|_{L^{\infty}} + 1),$$
(9.2)

where we have considered $\partial_z u$ instead of η . Next, to estimate the Lipschitz norm of u, we consider the vorticity formulation

$$\omega_t + u \cdot \nabla \omega = \omega \cdot \nabla u,$$

from where the maximum principle leads to

$$\frac{d}{dt} \|\omega\|_{L^{\infty}}^2 \lesssim \|\omega\|_{L^{\infty}}^2 \|\nabla u\|_{L^{\infty}}$$

Next, we repeat the remaining estimates in Section 6, recalling that $\nu = 0$, yielding a bound on $||u||_{2,\infty}$. Namely, we arrive at

$$\|u(t)\|_{2,\infty}^{2} + \|\omega(t)\|_{L^{\infty}}^{2} \lesssim \|u_{0}\|_{2,\infty}^{2} + \|\omega_{0}\|_{L^{\infty}}^{2} + \int_{0}^{t} \left((\|u\|_{2,\infty} + \|\omega\|_{L^{\infty}} + 1)^{3} + \|u\|_{2,\infty} \|D^{2}p\|_{3} \right) ds.$$
(9.3)

Since the term $\nu\Delta u$ does not appear, we do not obtain any boundary terms, and thus the assumptions on the sign of the friction parameter μ are not relevant. Now, we note that

$$\|\nabla u\|_{L^{\infty}} \lesssim \|\omega\|_{L^{\infty}} + \|u\|_{1,\infty}.$$
(9.4)

Upon combining, (9.1), (9.2), and (9.3), we employ (9.4) to conclude (2.7).

Having established the a priori estimates, we may construct the unique solution in the following manner. We first approximate our initial data with a sequence satisfying (1.9) and employ Theorem 2.3(ii) to obtain a sequence of approximate solutions. Note that the a priori estimate established above applies to this sequence for some T > 0. Next, we show that the same T, which is independent of the approximation, is a lower bound on maximum time

ACKNOWLEDGMENTS

The authors were supported in part by the NSF grant DMS-2205493. The research was performed during the program "Mathematical Problems in Fluid Dynamics, Part 2," held during the summer of 2023 by the Simons Laufer Mathematical Sciences Institute (SLMath), which is supported by the National Science Foundation (Grant No. DMS-1928930).

REFERENCES

- [BdVC1] H. Beirão da Veiga, and F. Crispo, Sharp inviscid limit results under Navier type boundary conditions. An L^p theory, J. Math. Fluid Mech., 12 (2010), no. 3, 397–411.
- [BdVC2] H. Beirão da Veiga, and F. Crispo, Concerning the W^{k,p}-inviscid limit for 3-D flows under a slip boundary condition, J. Math. Fluid Mech., 13 (2011), no. 1, 117–135.
- [BdVC3] H. Beirão da Veiga, and F. Crispo, A missed persistence property for the Euler equations and its effect on inviscid limits, Nonlinearity, **25** (2012), no. 6, 1661–1669.
- [BdVC4] H. Beirão da Veiga, and F. Crispo, The 3-D inviscid limit result under slip boundary conditions. A negative answer, J. Math. Fluid Mech., 14 (2012), no. 1, 55–59.
- [BS1] L.C. Berselli, and S. Spirito, On the vanishing viscosity limit of 3D Navier-Stokes equations under slip boundary conditions in general domains, Comm. Math. Phys., 316 (2012), no. 1, 171–198.
- [BS2] L.C. Berselli, and S. Spirito, An elementary approach to the inviscid limits for the 3D Navier-Stokes equations with slip boundary conditions and applications to the 3D Boussinesq equations, NoDEA Nonlinear Differential Equations Appl., 21 (2014), no. 2, 149–166.
- [CQ] G.Q. Chen, and Z. Qian, A study of the Navier-Stokes equations with the kinematic and Navier boundary conditions, Indiana Univ. Math. J., **59** (2010), no. 2, 721–760.
- [CL+] P. Constantin, M.C. Lopes Filho, H.J. Nussenzveig Lopes, and V. Vicol, Vorticity measures and the inviscid limit, Arch. Ration. Mech. Anal. 234 (2019), no. 2, 575–593.
- [DN] T.D. Drivas, and H.Q. Nguyen, *Remarks on the emergence of weak Euler solutions in the vanishing viscosity limit*, J. Nonlinear Sci., **29** (2019), no. 2, 709–721.
- [FTZ] M. Fei, T. Tao, and Z. Zhang, On the zero-viscosity limit of the Navier-Stokes equations in \mathbb{R}^3_+ without analyticity, J. Math. Pures Appl. (9) **112** (2018), 170–229.
- [GMM] Optimal Prandtl expansion around concave boundary layer, arXiv:2005.05022.
- [GK+] G.-M. Gie, J.P. Kelliher, M.C. Lopes Filho, A.L. Mazzucato, and H.J. Nussenzveig Lopes, *The vanishing viscosity limit* for some symmetric flows, Ann. Inst. H. Poincaré C Anal. Non Linéaire **36** (2019), no. 5, 1237–1280.
- [G] P. Grisvard, *Elliptic problems in nonsmooth domains*, Classics in Applied Mathematics, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2011.
- [GK] G.M. Gie, and J.P. Kelliher, *Boundary layer analysis of the Navier-Stokes equations with generalized Navier boundary conditions*, J. Differential Equations **253** (2012), no. 6, 1862–1892.
- [IP] D. Iftimie, and G. Planas, Inviscid limits for the Navier-Stokes equations with Navier friction boundary conditions, Nonlinearity, 19 (2006), no. 4, 899–918.
- [IS] D. Iftimie, and F. Sueur, Viscous boundary layers for the Navier-Stokes equations with the Navier slip conditions, Arch. Ration. Mech. Anal. 199 (2011), no. 1, 145–175.
- [K] T. Kato, Remarks on zero viscosity limit for nonstationary Navier-Stokes flows with boundary, Seminar on nonlinear partial differential equations (Berkeley, Calif., 1983), Math. Sci. Res. Inst. Publ., vol. 2, Springer, New York, 1984, pp. 85–98.
- [Ke] J.P. Kelliher, Vanishing viscosity and the accumulation of vorticity on the boundary, Commun. Math. Sci. 6 (2008), no. 4, 869–880.
- [KLS] I. Kukavica, M.C. Lombardo, and M. Sammartino, Zero viscosity limit for analytic solutions of the primitive equations, Arch. Ration. Mech. Anal. 222 (2016), no. 1, 15–45.
- [KVW] I. Kukavica, V. Vicol, and F. Wang, *The inviscid limit for the Navier-Stokes equations with data analytic only near the boundary*, Arch. Ration. Mech. Anal. 237 (2020), no. 2, 779–827.
- [M] Y. Maekawa, On the inviscid limit problem of the vorticity equations for viscous incompressible flows in the half-plane, Comm. Pure Appl. Math. 67 (2014), no. 7, 1045–1128.
- [MM] Y. Maekawa and A. Mazzucato, *The inviscid limit and boundary layers for Navier-Stokes flows*, Handbook of mathematical analysis in mechanics of viscous fluids, Springer, Cham, 2018, pp. 781–828.

- [MR1] N. Masmoudi, and F. Rousset, Uniform Regularity for the Navier–Stokes Equation with Navier Boundary Condition, Arch. Ration. Mech. Anal. 203 (2012), no. 2, 529–575.
- [MR2] N. Masmoudi, and F. Rousset, Uniform regularity and vanishing viscosity limit for the free surface Navier-Stokes equations, Arch. Ration. Mech. Anal. 223 (2017), no. 1, 301–417.
- [NP] J. Neustupa, and P. Penel, Approximation of a solution to the Euler equation by solutions of the Navier-Stokes equation, J. Math. Fluid Mech., 15 (2013), no. 1, 179–196.
- [SC1] M. Sammartino and R.E. Caflisch, Zero viscosity limit for analytic solutions, of the Navier-Stokes equation on a halfspace. I. Existence for Euler and Prandtl equations, Comm. Math. Phys. 192 (1998), no. 2, 433–461.
- [SC2] M. Sammartino and R.E. Caflisch, Zero viscosity limit for analytic solutions of the Navier-Stokes equation on a halfspace. II. Construction of the Navier-Stokes solution, Comm. Math. Phys. 192 (1998), no. 2, 463–491.
- [TWZ] T. Tao, W. Wang, and Z. Zhang, Zero-viscosity limit of the Navier-Stokes equations with the Navier friction boundary condition, SIAM J. Math. Anal. 52 (2020), no. 2, 1040–1095.
- [TW] R. Temam and X. Wang, *On the behavior of the solutions of the Navier-Stokes equations at vanishing viscosity*, vol. 25, 1997, Dedicated to Ennio De Giorgi, pp. 807–828.
- [WXZ] L. Wang, Z. Xin, and A. Zang, Vanishing viscous limits for 3D Navier-Stokes equations with a Navier-slip boundary condition, J. Math. Fluid Mech., 14 (2012), no. 4, 791–825.
- [X] Z. Xin, Vanishing viscosity limits for the 3D Navier-Stokes equations with a slip boundary condition, Proc. Amer. Math. Soc., 145 (2017), no. 4, 1615–1628.
- [XX] Y. Xiao, and Z. Xin, A new boundary condition for the three-dimensional Navier-Stokes equation and the vanishing viscosity limit, J. Math. Phys., 53 (2012), no. 11, 115617, 15.
- [XZ1] Y. Xiao, and Z. Xin, On the vanishing viscosity limit for the 3D Navier-Stokes equations with a slip boundary condition, Comm. Pure Appl. Math., 60 (2007), no. 7, 1027–1055.
- [XZ2] Y. Xiao, and Z. Xin, On the inviscid limit of the 3D Navier-Stokes equations with generalized Navier-slip boundary conditions, Commun. Math. Stat., 1 (2013), no. 3, 259–279.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA 90089 *Email address*: maydin@usc.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA 90089 *Email address*: kukavica@usc.edu