# The pro-Nisnevich topology

Klaus Mattis<sup>\*</sup>

April 29, 2024

#### Abstract

We construct the pro-Nisnevich topology, an analog of the pro-étale topology. We then show that the Nisnevich  $\infty$ -topos embeds into the pro-Nisnevich  $\infty$ -topos, and that the pro-Nisnevich  $\infty$ -topos is locally of homotopy dimension 0.

### Introduction

In [BS14], Bhatt and Scholze constructed the pro-étale topology, and showed that the étale topos over a scheme embeds into the pro-étale topos.

In this short note, we do a similar construction for the Nisnevich topology. Let S be a qcqs scheme of finite Krull-dimension (e.g. a field), and write  $Sm_S$  for the category of smooth quasi-compact S-schemes, equipped with the Nisnevich topology [BH17, Appendix A]. Write  $Shv_{nis}(Sm_S)$  for the  $\infty$ -topos of sheaves on this site. We will construct a site ( $ProSm_S$ , pronis), called the *pro-Nisnevich* site. Write  $Shv_{pronis}^h(ProSm_S)$  for the  $\infty$ -topos of hypersheaves on this site. This  $\infty$ -topos is called the *pro-Nisnevich topos*. We prove the following results:

**Theorem A** (Theorem 2.10). There is a full subcategory  $W \subset \operatorname{ProSm}_S$  and an equivalence  $\operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{h}}(\operatorname{ProSm}_S) \cong \mathcal{P}_{\Sigma}(W)$ . In particular, the pro-Nisnevich topos is locally of homotopy dimension 0.

**Theorem B** (Theorem 2.12). There is a geometric morphism of  $\infty$ -topoi

 $\nu^*$ : Shv<sub>nis</sub>(Sm<sub>S</sub>)  $\rightleftharpoons$  Shv<sup>h</sup><sub>pronis</sub>(ProSm<sub>S</sub>):  $\nu_*$ ,

where  $\nu_*$  is given by restriction and  $\nu^*$  is fully faithful.

Moreover, an n-truncated sheaf  $F \in \operatorname{Shv}_{\operatorname{pronis}}^{h}(\operatorname{ProSm}_{S})$  is in the essential image of  $\nu^*$  if and only if for all  $U \in \operatorname{ProSm}_{S}$  and all presentations of U as a cofiltered limit  $U \cong \lim_i U_i$  (with the  $U_i \in \operatorname{Sm}_S$ ) the canonical comparison map  $\operatorname{colim}_i F(U_i) \to F(U)$  is an equivalence.

<sup>\*</sup>klaus.mattis@uni-mainz.de

#### Acknowledgement

I thank Tom Bachmann for helpful discussions, and Timo Weiß for reading a draft of this article.

The author acknowledges support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through the Collaborative Research Centre TRR 326 *Geometry and Arithmetic of Uniformized Structures*, project number 444845124.

### 1 The Pro-Nisnevich Topology

In this section, we develop an analog of the pro-étale topology from [BS14], adapted for the Nisnevich topology. We show that every affine scheme can be covered in this topology by a cdw-contractible ring, an analog of w-contractible rings from [BS14].

Recall the following definition from [BS14, Section 2.1 and 2.2], especially the alternative characterizations of w-local spaces from [BS14, Lemma 2.1.4]:

**Definition 1.1.** Let A be a ring. We say that A is w-local if the set of closed points  $\operatorname{Spec}(A)^c$  is closed in  $\operatorname{Spec}(A)$ , and the map  $\operatorname{Spec}(A)^c \to \operatorname{Spec}(A) \to \pi_0(\operatorname{Spec}(A))$  is bijective.

Recall the following definitions:

**Definition 1.2.** Let  $f: A \to B$  be a ring map. We say that

- 1. f is a local isomorphism if for every prime ideal  $\mathfrak{q} \subset B$  there exists a  $g \in B$ ,  $g \notin \mathfrak{q}$  such that  $A \to B_g$  induces an open immersion  $\operatorname{Spec}(B_g) \to \operatorname{Spec}(A)$ ,
- 2. f is an *ind-Zariski localization* if f can be written as a filtered colimit of local isomorphisms  $f_i: A \to B_i$ , and
- 3. f is an *ind-Zariski cover* if f is a faithfully flat ind-Zariski localization.

We have the following results about w-local rings:

**Lemma 1.3.** Let A be a ring. Then there exists an ind-Zariski cover  $A \to A^Z$  with  $A^Z$  w-local, such that  $\operatorname{Spec}(A^Z)^c \to \operatorname{Spec}(A)$  is bijective.

Proof. Combine [BS14, Lemma 2.2.4] and [BS14, Lemma 2.1.10].

**Lemma 1.4.** For a ring A and a map  $T \to \pi_0(\operatorname{Spec}(A))$  of totally disconnected compact Hausdorff spaces, there is an ind-Zariski localization  $A \to B$  such that  $\operatorname{Spec}(B) \to \operatorname{Spec}(A)$  gives rise to the given map  $T \to \pi_0(\operatorname{Spec}(A))$  on applying  $\pi_0$ .

Proof. This is [BS14, Lemma 2.2.8].

#### 1.1 Absolutely flat rings

Recall the following definitions from [BS14]:

**Definition 1.5.** Let A be a ring. We say that A is *absolutely flat* if A is reduced and has dimension 0.

Let  $f: A \to B$  be a ring map. We say that f is *weakly étale* if f is flat with flat diagonal. We say that f is *ind-étale* if f can be written as a filtered colimit of étale morphisms  $f_i: A \to B_i$ .

**Lemma 1.6.** Let  $f: A \to B$  be a morphism of rings with A absolutely flat. Suppose that f is weakly étale. Then B is absolutely flat.

Proof. See [Sta23, Tag 092I].

**Lemma 1.7.** Let A be a ring of dimension 0 (e.g. A absolutely flat). Then Spec(A) is totally disconnected compact Hausdorff.

*Proof.* Clearly Spec(A) is compact. It is Hausdorff, see e.g. [Sta23, Tag 0CKV], which applies since Spec(A) is affine and hence separated. Since Spec(A) is spectral, it has a basis of compact open subsets. In Hausdorff spaces, compact subsets are closed, thus Spec(A) has a basis of clopen subsets. This implies that A is totally disconnected.

Corollary 1.8. Let A be absolutely flat. Then A is w-local.

*Proof.* We know that Spec(A) is totally disconnected compact Hausdorff from Lemma 1.7. In particular, all points of Spec(A) are closed and  $\text{Spec}(A) \cong \pi_0(\text{Spec}(A))$ .

**Lemma 1.9.** Let  $A = \operatorname{colim}_i A_i$  be a filtered colimit of absolutely flat rings. Then A is absolutely flat.

*Proof.* Being reduced and of dimension 0 are properties of rings that are stable under filtered colimits: For the first claim, note that a nilpotent element in a ring is the same as a ring map from  $\mathbb{Z}[X]/(X^n)$  for some  $n \ge 2$ . But since  $\mathbb{Z}[X]/(X^n)$ is of finite presentation over  $\mathbb{Z}$ , we see that  $\operatorname{Hom}(\mathbb{Z}[X]/(X^n), \operatorname{colim}_i A_i) \cong$  $\operatorname{colim}_i \operatorname{Hom}(\mathbb{Z}[X]/(X^n), A_i) = 0$ , since the  $A_i$  are reduced. For the second claim, use [Sta23, Tag 01YW and Tag 01YY (2)].

**Definition 1.10.** Let  $f: \operatorname{Spec}(B) \to \operatorname{Spec}(A)$  be a map of affine schemes. We define the *completely decomposed locus* of f as the subset

 $cd(f) \coloneqq \{x \in Spec(B) \mid k(f(x)) \to k(x) \text{ is an isomorphism}\} \subset Spec(B).$ 

We say that f is completely decomposed if f(cd(f)) = Spec(A), i.e. over every point of Spec(A) there is a point in Spec(B) that induces an isomorphism on residue fields.

**Lemma 1.11.** Let  $A = \operatorname{colim}_{i \in I} A_i$  be a filtered colimit of rings. Suppose wlog I has an initial element 0. Denote by  $f_i: \operatorname{Spec}(A_i) \to \operatorname{Spec}(A_0)$  the map induced by  $0 \to i$ , and by  $f_{\infty}: \operatorname{Spec}(A) \to \operatorname{Spec}(A_0)$  the projection.

Then  $\operatorname{cd}(f_{\infty}) = \lim_{i} \operatorname{cd}(f_{i}).$ 

*Proof.* Let  $p_i: \operatorname{Spec}(A) \to \operatorname{Spec}(A_i)$  be the projections. Let  $x_{\infty} \in \operatorname{Spec}(A)$ , and  $x_i := p_i(x_{\infty}) \in \operatorname{Spec}(A_i)$ .

Suppose  $x_{\infty} \in \operatorname{cd}(f_{\infty})$  and thus  $k(x_0) \cong k(x_{\infty})$ . Thus, for all *i* we have a factorization of this isomorphism  $k(x_0) \to k(x_i) \to k(x_{\infty})$ . Hence  $k(x_i) \cong k(x_0)$  and we get  $x_i \in \operatorname{cd}(f_i)$ . This implies  $x_{\infty} \in \lim_i \operatorname{cd}(f_i)$ .

On the other hand, suppose that  $x_{\infty} \in \lim_{i} \operatorname{cd}(f_{i})$ . This implies that  $x_{i} \in \operatorname{cd}(f_{i})$ . By [Sta23, Lemma 0CUG], we know that  $k(x_{\infty}) = \operatorname{colim}_{i} k(x_{i})$ . But  $k(x_{0}) \cong k(x_{i})$  and thus  $k(x_{\infty}) \cong \operatorname{colim}_{i} k(x_{0}) \cong k(x_{0})$ . Hence  $x_{\infty} \in \operatorname{cd}(f_{\infty})$ .  $\Box$ 

**Lemma 1.12.** Let A an absolutely flat ring. Write  $A \cong \operatorname{colim}_i A_i$  as a filtered colimit of its finitely generated subrings. Let  $x \in \operatorname{Spec}(A)$ . For each i write  $p_i: \operatorname{Spec}(A) \to \operatorname{Spec}(A_i)$  for the projection, and  $Z_i(x) \subset \operatorname{Spec}(A_i)$  for the connected component of  $p_i(x)$ . Then  $\{x\} \cong \lim_i Z_i(x)$ .

Proof. Clearly  $x \in \lim_i Z_i(x)$ . Suppose on the other hand that  $y \in \operatorname{Spec}(A)$ , with  $y \neq x$ . Since  $\operatorname{Spec}(A)$  is totally disconnected compact Hausdorff by Lemma 1.7, there exists a clopen subset  $U \subset \operatorname{Spec}(A)$  into clopen subsets with  $x \in U$  and  $y \notin U$ . By descent, there is an index j, and  $U_j \subset \operatorname{Spec}(A_j)$  clopen such that  $U = \operatorname{Spec}(A) \times_{\operatorname{Spec}(A_j)} U_j$ . But then  $Z_i(x) \subseteq \operatorname{Spec}(A_i) \times_{\operatorname{Spec}(A_j)} U_j$  for all  $i \geq j$ , and hence  $\lim_i Z_i(x) \subseteq U$ . Since  $y \notin U$ , we conclude  $y \notin \lim_i Z_i(x)$ .

**Lemma 1.13.** Let  $f: A \to B$  be an étale ring map, with A absolutely flat. Then  $cd(f) \subset Spec(B)$  is closed.

Proof. Write  $A = \operatorname{colim}_{i \in I} A_i$  as a filtered colimit of its finitely generated subrings. As f is of finite presentation, by descent there is an index  $0 \in I$ , a ring  $B_0$  and an étale morphism  $f_0: B_0 \to A_0$  such that f is the basechange of  $f_0$ . Write  $B_i \coloneqq B_0 \otimes_{A_0} A_i$  for each  $i \geq 0$ , and  $f_i: A_i \to B_i$ . In particular,  $\operatorname{Spec}(B) \cong \lim_{i\geq 0} \operatorname{Spec}(B_i)$ . Note that  $\operatorname{Spec}(B_i)$  is of finite type over  $\operatorname{Spec}(\mathbb{Z})$ (as it is étale over  $\operatorname{Spec}(A_i)$  which is of finite type over  $\operatorname{Spec}(\mathbb{Z})$ ). In particular, the connected components of  $\operatorname{Spec}(B_i)$  are open. Write  $W_i \subset \operatorname{Spec}(B_i)$  for the union of those components  $W_{i,j} \subset \operatorname{Spec}(B_i)$  such that  $f_i|_{W_{i,j}}: W_{i,j} \to \operatorname{Spec}(A_i)$ is an isomorphism onto a component of  $\operatorname{Spec}(A_i)$ . It suffices to prove that  $\operatorname{cd}(f) = \lim_i W_i$ : Indeed, since  $\operatorname{Spec}(B_i)$  has only finitely many components, we see that  $W_i$  is closed. Thus, also  $\operatorname{cd}(f) = \lim_i W_i$  is closed as a limit of closed subsets.

It is clear that  $\lim_i W_i \subseteq \operatorname{cd}(f)$ , as  $W_i \subseteq \operatorname{cd}(f_i)$ . So let  $x \in \operatorname{cd}(f)$  and  $y \coloneqq f(x)$ . We will write  $p_i$ :  $\operatorname{Spec}(A) \to \operatorname{Spec}(A_i)$  for the projection, and  $Z_i(y) \subseteq \operatorname{Spec}(A_i)$  for the component of  $p_i(y)$ . Let  $V_i(y) \coloneqq \operatorname{Spec}(B_i) \times_{\operatorname{Spec}(A_i)}$   $Z_i(y) \subseteq \operatorname{Spec}(B_i)$ , which is a clopen subset. As the inclusion  $\{x\} \hookrightarrow f^{-1}(y)$  is clopen (because  $f^{-1}(y)$  is a discrete set of points), we may assume by descent (possibly by replacing 0 by a larger index), that there exists a  $T_0 \subseteq V_0(y)$  clopen such that  $\{x\} = T_0 \times_{V_0(y)} f^{-1}(y)$ , and set  $T_i \coloneqq \operatorname{Spec}(B_i) \times_{\operatorname{Spec}(B_0)} T_0$  (which is clopen in  $V_i(y)$ ). Note that  $T_i \times_{Z_i(y)} \{y\} \cong T_i \times_{V_i(y)} f^{-1}(y) \cong \{x\}$ . As f is completely decomposed at x by definition, it induces an isomorphism  $\{x\} \xrightarrow{\simeq} \{y\}$ . Thus, again by descent and Lemma 1.12, we conclude that  $T_i \xrightarrow{\simeq} Z_i(y)$ for  $i \gg 0$ . But  $T_i \subseteq V_i(y)$  is clopen, and hence  $T_i \subseteq V_i(y) \subseteq \text{Spec}(B_i)$  is also clopen, and maps isomorphically to  $Z_i(y)$ . Therefore,  $T_i \subseteq W_i$  and we conclude that  $x \in \lim_i W_i$ .

**Corollary 1.14.** Let A be absolutely flat and  $f: A \to B$  an ind-étale algebra. Then cd(f) is closed.

*Proof.* As a limit of closed subsets is closed, we can first reduce to the case that f is étale via Lemma 1.11, and then conclude by Lemma 1.13.

#### **1.2** Weakly Contractible Covers

**Definition 1.15.** Let  $f: A \to B$  be a ring map. We say that f is a Nisnevich cover if f is étale and completely decomposed; and that f is an *ind-Nisnevich* cover if f is ind-étale and completely decomposed.

**Definition 1.16.** A ring A is called *cdw-contractible* if every ind-Nisnevich-cover  $A \rightarrow B$  splits.

**Definition 1.17.** Let A be a ring. We write  $\operatorname{Ind}_{et}(A)$  for the category of indétale A-algebras.

Recall the following definition from [BS14, Definition 2.2.10]:

**Definition 1.18.** Let  $A \to B$  be a ring map. We have a base-change functor  $\operatorname{Ind}_{et}(A) \to \operatorname{Ind}_{et}(B)$ . Define  $\operatorname{Hens}_A(-)$ :  $\operatorname{Ind}_{et}(B) \to \operatorname{Ind}_{et}(A)$  via

$$B' \mapsto \operatorname{colim}_{A \xrightarrow{et} A' \to B'} A'$$

This is a right adjoint to the base change functor and called *henselization*. We say that a ring A is *henselian along an ideal* I if  $\text{Hens}_A(A/I) \cong A$ .

**Lemma 1.19.** Let A be a ring, and  $I \subseteq A$  an ideal contained in the Jacobson radical  $I_A$  of A. Let  $\overline{A}$  be an ind-étale A/I-algebra. Then  $B := \text{Hens}_A(\overline{A})$  is henselian along IB with  $B/IB \cong \overline{A}$ .

*Proof.* We first prove the second part. Since  $\overline{A}$  is an ind-étale A/I-algebra, we know that  $\operatorname{Hens}_{A/I}(\overline{A}) \cong \overline{A}$ . Note that  $B/IB = \operatorname{colim}_{A \xrightarrow{et} A' \to \overline{A}} A'/IA'$ . By basechange,  $A/I \to A'/IA'$  is an étale morphism for all such A'. Thus, it suffices to show that every factorization  $A/I \xrightarrow{et} C \to \overline{A}$  is the basechange of a factorization  $A \to A' \to \overline{A}$ . Note that  $A/I \to C$  can be lifted to an étale morphism  $A \to A'$  by [Sta23, Tag 04D1]. But since A'/IA' = C, this fits into a factorization  $A \to A' \to \overline{A}$  as desired.

For the first part we need to show that  $B \xrightarrow{\text{id}} B$  is final under étale morphisms  $B \to B'$  such that there is a factorization  $B \to B' \to B/IB = \overline{A}$ . So let

 $B \to B'$  such an étale morphism. We need to show that it has a section. But  $B = \operatorname{colim}_{A \xrightarrow{et} A' \to \overline{A}} A'$ . By finite presentation of  $B \to B'$  there is thus a factorization  $A \to A' \to \overline{A}$  with  $A \to A'$  étale and an étale morphism  $A' \to \tilde{B}$  such that  $B \otimes_{A'} \tilde{B} \cong B'$ . Since  $A' \to \tilde{B}$  is étale, also  $A \to \tilde{B}$  is étale. Note that there is a factorization  $A \to \tilde{B} \to \overline{A}$ . We then get a canonical map  $\tilde{B} \to B$  by definition of B. But then  $B' \cong B \otimes_{A'} \tilde{B} \to B \otimes_{A'} B \to B$  gives the desired section.

**Lemma 1.20.** Let A be a ring which is henselian along an ideal I. Suppose A/I is cdw-contractible. Then A is cdw-contractible.

*Proof.* Let  $f: A \to B$  be an ind-Nisnevich cover. Then  $f/I: A/I \to B/IB$  is an ind-Nisnevich cover and thus has a section. But

$$\operatorname{Hom}_{A}(B, A) \cong \operatorname{Hom}_{A}(B, \operatorname{Hens}_{A}(A/I)) \cong \operatorname{Hom}_{A/I}(B/IB, A/I).$$

This gives us the desired section of f.

**Definition 1.21.** Let T be a totally disconnected compact Hausdorff space. We say that T is *extremally disconnected* if every surjection  $S \rightarrow T$  from a totally disconnected compact Hausdorff space S has a section.

**Lemma 1.22.** Let A be absolutely flat. Then A is cdw-contractible if and only if  $\text{Spec}(A) = \pi_0(\text{Spec}(A))$  is extremally disconnected.

*Proof.* First we see from Lemma 1.7 that because A is absolutely flat, the topological space Spec(A) is a totally disconnected compact Hausdorff space, so in particular  $\text{Spec}(A) \cong \pi_0(\text{Spec}(A))$ .

Suppose that A is cdw-contractible. Let  $f_0: T \to \operatorname{Spec}(A)$  be a surjective morphism of totally disconnected compact Hausdorff spaces. But  $f_0$  is induced by an ind-Zariski cover  $A \to B$ , see Lemma 1.4, which has a section because A is cdw-contractible and ind-Zariski covers are ind-Nisnevich covers. This gives us the topological section  $\operatorname{Spec}(A) \cong \pi_0(\operatorname{Spec}(A)) \to \pi_0(\operatorname{Spec}(B)) \cong T$ .

Suppose now that  $\operatorname{Spec}(A)$  is extremally disconnected. Let  $f \colon A \to B$  be an ind-Nisnevich cover. Thus, B is absolutely flat, see Lemma 1.6 (as ind-étale maps are weakly étale, see [Sta23, Tag 092N]). Using Corollary 1.14, we see that  $\operatorname{cd}(\operatorname{Spec}(f)) \subseteq \operatorname{Spec}(B)$  is closed. Since f is completely decomposed, we see that  $\operatorname{cd}(\operatorname{Spec}(f)) \to \operatorname{Spec}(B) \to \operatorname{Spec}(A)$  is in fact surjective. Let B' be the B-algebra realizing the closed subset  $\operatorname{cd}(\operatorname{Spec}(f))$  with its reduced scheme structure. Then B' is reduced and thus absolutely flat. By definition of B',  $h \colon \operatorname{Spec}(B') \to \operatorname{Spec}(A)$  induces isomorphisms on all residue fields. Now, h has a topological section s because  $\operatorname{Spec}(A) = \pi_0(\operatorname{Spec}(A))$  is extremally disconnected, and  $\operatorname{Spec}(B')$  is totally disconnected compact Hausdorff by Lemma 1.7. Let  $\operatorname{Spec}(B'')$  be the reduced closed subscheme of  $\operatorname{Spec}(B')$  realizing the image  $s(\operatorname{Spec}(A)) \subset \operatorname{Spec}(B')$  (note that the image of s is closed because s is a map between compact Hausdorff spaces). This maps isomorphically to  $\operatorname{Spec}(A)$ , which gives us the section

$$\operatorname{Spec}(A) \cong \operatorname{Spec}(B'') \subseteq \operatorname{Spec}(B') \subseteq \operatorname{Spec}(B).$$

**Lemma 1.23.** Let A be a ring. Then there is an ind-Nisnevich cover  $A \to \overline{A}$  with  $\overline{A}$  a cdw-contractible ring.

*Proof.* Consider the ind-Zariski cover  $A \to A^Z$  by a w-local ring from Lemma 1.3. Note that  $A^Z/I_{A^Z}$  is absolutely flat, using [BS14, Lemma 2.2.3]. Let T be an extremally disconnected compact Hausdorff space covering  $\operatorname{Spec}(A^Z/I_{A^Z})$ . Let A' be an ind-Zariski cover of  $A^Z/I_{A^Z}$  such that  $\operatorname{Spec}(A') \to \operatorname{Spec}(A^Z/I_{A^Z})$  realizes the map  $T \to \operatorname{Spec}(A^Z/I_{A^Z})$ , see Lemma 1.4. Then A' is absolutely flat, and cdw-contractible by Lemma 1.22. Define  $\overline{A} := \operatorname{Hens}_{A^Z}(A')$ . Then  $\overline{A}$  is henselian along  $I_{A^Z}\overline{A}$ , with quotient  $\overline{A}/I_{A^Z}\overline{A} \cong A'$  (see Lemma 1.19), hence  $\overline{A}$  is cdw-contractible using Lemma 1.20. Now note that  $A^Z \to \overline{A}$  induces isomorphisms of residue fields at the closed points of  $\operatorname{Spec}(A^Z)$ . Note that  $\operatorname{Spec}(A^Z)^c \to \operatorname{Spec}(A)$  is bijective (see Lemma 1.3) and induces isomorphisms of residue fields. Thus,  $\operatorname{Spec}(\overline{A}) \to \operatorname{Spec}(A)$  is completely decomposed. Since  $A \to A^Z$  is ind-Zariski, and  $A^Z \to \overline{A}$  is ind-étale by construction, we conclude that  $A \to \overline{A}$  is an ind-Nisnevich cover. □

### 2 The Pro-Nisnevich $\infty$ -Topos

Let S be a qcqs scheme with finite Krull-dimension.

**Definition 2.1.** Write  $\operatorname{ProSm}_S$  for the category of pro-smooth schemes over S, i.e. morphisms  $X \to S$  such that X can be written as a cofiltered limit  $X \cong \lim_i X_i$  where  $X_i \in \operatorname{Sm}_S$ . Write  $\operatorname{ProSmAff}_S$  for the full subcategory of pro-smooth schemes over S which are affine.

**Definition 2.2.** Let  $\mathcal{U} := \{f_i \colon U_i \to U\}_{i \in I}$  be a family of morphisms in  $\operatorname{ProSm}_S$ . We say that  $\mathcal{U}$  is a *pro-Nisnevich cover* if

- $f_i$  is pro-étale for all  $i \in I$ ,
- the morphism  $\coprod_i U_i \to U$  is completely decomposed, and
- the  $f_i$  form an fpqc-cover [Sta23, Tag 03NW].

Similarly, let  $\mathcal{U} := \{f_i : U_i \to U\}_{i \in I}$  be a family of morphisms in ProSmAff<sub>S</sub>. We say that  $\mathcal{U}$  is a *pro-Nisnevich cover* if it is a pro-Nisnevich cover in ProSm<sub>S</sub>.

Remark 2.3. Let Spec(f):  $\text{Spec}(B) \to \text{Spec}(A)$  be a morphism in  $\text{ProSmAff}_S$ . Then  $\{\text{Spec}(f)\}$  is a pro-Nisnevich cover if and only if  $f: A \to B$  is an ind-Nisnevich cover.

Recall the definition of a site [Sta23, Tag 00VH] and a morphism of sites [Sta23, Tag 00X1].

**Lemma 2.4.** There are sites ( $ProSm_S$ , pronis) and ( $ProSmAff_S$ , pronis), where the covers are given by pro-Nisnevich covers.

*Proof.* The only nontrivial part is the stability of covers under pullbacks. First note that the pullback of a pro-étale cover exists and is again a pro-étale cover, which follows from [Sta23, Lemma 098L]. Since completely decomposed families of morphisms are stable under pullbacks in the category of all S-schemes, the result follows as pullbacks in  $ProSm_S$  along pro-étale morphisms coincide with the underlying pullback in the category of all S-schemes (this can be proven analogously to [Sta23, Tag 098M]).

We can define the following  $\infty$ -topoi:

Definition 2.5. We will write

- $\operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{nh}}(\operatorname{ProSm}_S)$  for the  $\infty$ -topos of sheaves of anima on the site ( $\operatorname{ProSm}_S$ ,  $\operatorname{pronis}$ ),
- $\operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{h}}(\operatorname{ProSm}_{S})$  for the hypercompletion of  $\operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{nh}}(\operatorname{ProSm}_{S})$ ,
- $\operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{nh}}(\operatorname{ProSmAff}_S)$  for the  $\infty$ -topos of sheaves of anima on the site ( $\operatorname{ProSmAff}_S$ ,  $\operatorname{pronis}$ ), and
- $\operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{h}}(\operatorname{ProSmAff}_{S})$  for the hypercompletion of  $\operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{nh}}(\operatorname{ProSmAff}_{S})$ .

**Lemma 2.6.** There are equivalences of  $\infty$ -topoi

$$\operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{nh}}(\operatorname{ProSmAff}_S) \cong \operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{nh}}(\operatorname{ProSm}_S)$$

and

$$\operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{h}}(\operatorname{ProSmAff}_{S}) \cong \operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{h}}(\operatorname{ProSm}_{S})$$

induced by left Kan extension along the inclusion  $\operatorname{ProSmAff}_S \hookrightarrow \operatorname{ProSm}_S$ .

*Proof.* Note that the inclusion u: ProSmAff<sub>S</sub>  $\rightarrow$  ProSm<sub>S</sub> preserves covers by definition, and commutes with limits as limits of affine schemes are affine. Thus, the first equivalence is an easy application of [Hoy15, Lemma C.3]. The second equivalence is just the hypercompletion of the first.

**Definition 2.7.** We write W for the full subcategory of ProSmAff<sub>S</sub> consisting of (spectra of) cdw-contractible rings.

**Lemma 2.8.** The category W is extensive ([BH17, Definition 2.3]), and the category  $\mathcal{P}_{\Sigma}(W)$  is an  $\infty$ -topos.

*Proof.* The category is extensive, as it is a full subcategory of the category of schemes (which is extensive), stable under finite coproducts and summands. The second part is [BH17, Lemma 2.4].  $\Box$ 

Recall the notion of a locally weakly contractible site [Mat24, Definition B.3].

**Lemma 2.9.** The site ( $ProSmAff_S$ , pronis) is locally weakly contractible.

*Proof.* We have seen that the category W is extensive, see Lemma 2.8, and consists by definition of weakly contractible objects. The pro-Nisnevich topology is a Σ-topology, because clopen immersions are in particular pro-étale and induce isomorphisms on the residue fields. The pro-Nisnevich topology on ProSm<sub>S</sub> is finitary (cf. [Lur18, Definition A.3.1.1]) by definition, so every object is quasicompact. Note that every object in ProSmAff<sub>S</sub> has a cover by an object in W, this is the content of Lemma 1.23. This proves the lemma.

**Theorem 2.10.** There is an equivalence of  $\infty$ -topoi

$$\operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{h}}(\operatorname{ProSm}_{S}) \cong \mathcal{P}_{\Sigma}(W).$$

Thus,  $\operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{h}}(\operatorname{ProSm}_{S})$  is locally of homotopy dimension 0 and in particular Postnikov-complete.

*Proof.* There is an equivalence  $\operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{h}}(\operatorname{ProSm}_{S}) \cong \operatorname{Shv}_{\operatorname{pronis}}^{\operatorname{h}}(\operatorname{ProSmAff}_{S})$ , see Lemma 2.6. Thus, the theorem follows from an application of [Mat24, Lemma B.7], because the site (ProSmAff<sub>S</sub>, pronis) is locally weakly contractible, see Lemma 2.9. See [Mat24, Lemma 4.20] for a proof that  $\mathcal{P}_{\Sigma}(W)$  is locally of homotopy dimension 0. The last claim is [Lur09, Proposition 7.2.1.10]. □

**Lemma 2.11.** The  $\infty$ -topos  $Shv_{nis}(Sm_S)$  is Postnikov-complete.

*Proof.* If S is noetherian, then the proof of this result is completely analogous to the proof of [Mat24, Lemma 5.1]. Here we use that S is qcqs of finite Krull dimension (and that schemes in  $\text{Sm}_S$  possess the same property, note that for us, they are by definition quasi-compact). If S is not noetherian, we can still apply the same strategy, but resort to [CM21, Theorem 3.18] instead of [Lur18, Theorem 3.7.7.1].

Theorem 2.12. There is a geometric morphism

 $\nu^*$ : Shv<sub>nis</sub>(Sm<sub>S</sub>)  $\rightleftharpoons$  Shv<sup>h</sup><sub>pronis</sub>(ProSm<sub>S</sub>):  $\nu_*$ ,

where the right adjoint is given by restriction, and the left adjoint is fully faithful.

Moreover, an n-truncated sheaf  $F \in \text{Shv}_{\text{pronis}}^{\text{h}}(\text{ProSm}_S)$  is in the essential image of  $\nu^*$  if and only if for all  $U \in \text{ProSm}_S$  and all presentations of U as cofiltered limit  $U \cong \lim_i U_i$  (with the  $U_i \in \text{Sm}_S$ ) the canonical map  $\text{colim}_i F(U_i) \to F(U)$  is an equivalence.

*Proof.* Both involved  $\infty$ -topoi are Postnikov-complete, see Theorem 2.10 and Lemma 2.11. We want to apply [Mat24, Proposition B.8] to the natural inclusion of sites

$$(Sm_S, nis) \subset (ProSm_S, pronis).$$

In the notation of [Mat24, Proposition B.8], it remains to prove that  $\iota_h j^* F \cong k^* \iota'_h F$  for every *n*-truncated Nisnevich sheaf  $F \in \text{Shv}_{nis}(\text{Sm}_S)$ , i.e. we have to show that the presheaf  $k^* \iota'_h F$  is already a pro-Nisnevich hypersheaf. The proof of this fact is completely analogous to the proof of [Mat24, Theorem B.24].  $\Box$ 

## References

- [BH17] Tom Bachmann and Marc Hoyois. Norms in motivic homotopy theory. Astérisque, 2017.
- [BS14] Bhargav Bhatt and Peter Scholze. The pro-étale topology for schemes, 2014.
- [CM21] Dustin Clausen and Akhil Mathew. Hyperdescent and étale k-theory. Inventiones mathematicae, 225:981–1076, 2021.
- [Hoy15] Marc Hoyois. A quadratic refinement of the grothendieck-lefschetzverdier trace formula. Algebraic & Geometric Topology, 14(6):3603-3658, 2015.
- [Lur09] Jacob Lurie. Higher topos theory, 2009.
- [Lur18] Jacob Lurie. Spectral algebraic geometry. Available online at https://www.math.ias.edu/~lurie/papers/sag-rootfile.pdf, 2018.
- $[{\rm Mat}24]$  Klaus Mattis. Unstable  $p\mbox{-completion}$  in motivic homotopy theory, 2024.
- [Sta23] The Stacks project authors. The stacks project. https://stacks.math.columbia.edu, 2023.