

# Walking behavior induced by $\mathcal{PT}$ symmetry breaking in a non-Hermitian XY model with clock anisotropy

Eduard Naichuk,<sup>1,2</sup> Jeroen van den Brink,<sup>1,3</sup> and Flavio S. Nogueira<sup>1</sup>

<sup>1</sup>*Institute for Theoretical Solid State Physics, IFW Dresden, Helmholtzstr. 20, 01069 Dresden, Germany*

<sup>2</sup>*Bogolyubov Institute for Theoretical Physics, 03143 Kyiv, Ukraine*

<sup>3</sup>*Institute for Theoretical Physics and Würzburg-Dresden Cluster of Excellence ct.qmat, TU Dresden, 01069 Dresden, Germany*

(Dated: April 29, 2024)

A quantum system governed by a non-Hermitian Hamiltonian may exhibit zero temperature phase transitions that are driven by interactions, just as its Hermitian counterpart, raising the fundamental question how non-Hermiticity affects quantum criticality. In this context we consider a non-Hermitian system consisting of an XY model with a complex-valued four-state clock interaction that may or may not have parity-time-reversal ( $\mathcal{PT}$ ) symmetry. When the  $\mathcal{PT}$  symmetry is broken, and time-evolution becomes non-unitary, a scaling behavior similar to the Berezinskii-Kosterlitz-Thouless phase transition ensues, but in a highly unconventional way, as the line of fixed points is absent. From the analysis of the  $d$ -dimensional RG equations, we obtain that the unconventional behavior in the  $\mathcal{PT}$  broken regime follows from the collision of two fixed points in the  $d \rightarrow 2$  limit, leading to walking behavior or pseudocriticality. For  $d = 2 + 1$  the near critical behavior is characterized by a correlation length exponent  $\nu = 3/8$ , a value smaller than the mean-field one. These results are in sharp contrast with the  $\mathcal{PT}$ -symmetric case where only one fixed point arises for  $2 < d < 4$  and in  $d = 1 + 1$  three lines of fixed points occur with a continuously varying critical exponent  $\nu$ .

The possibilities to access non-Hermitian systems [1–3] experimentally significantly expanded in the past years, with possible incarnations ranging from metamaterials to ultracold atoms realizations [4–6], and more recently including electronic systems having applications as sensors [7–9]. While in any realistic physical system interactions play an important role, from a theoretical perspective *interacting* non-Hermitian systems have been little explored. Particularly pertinent is the question how non-Hermiticity affects interaction-driven quantum phase transitions and their universal physical properties also in absence of  $\mathcal{PT}$  symmetry, where time-evolution becomes non-unitary [3, 10–13]. This motivates considering quantum field theoretic models, where the renormalization group (RG) provides the defining paradigm for exploration of the phase structure. In this context it is very important to establish how the critical behavior depends on the spacetime dimensionality  $d = D + 1$ .

In order to address the questions above and understand how the breaking of  $\mathcal{PT}$  symmetry affects the phase structure, we investigate an effective quantum field theory Lagrangian consisting of an XY model with a non-Hermitian,  $\mathcal{PT}$ -symmetric clock interaction in  $d = D + 1$  spacetime dimensions. Its Hermitian counterpart arises in several contexts both in high-energy physics [14, 15] and condensed matter physics [16–21]. From an RG perspective the  $N$ -state clock interaction is an irrelevant operator for  $N > 4$  in spacetime dimensions  $2 < d < 4$ , so one expects a crossover behavior from the XY universality class to  $Z_N$  symmetric models. In particular, the case  $N = 4$  corresponds in the strong coupling clock interaction limit to two decoupled Ising regimes [19, 20]. Since the  $\mathcal{PT}$ -symmetric clock model can be mapped into

its Hermitian counterpart, the fundamental question to answer is how  $\mathcal{PT}$  symmetry breaking changes the phase structure of the system in  $d = D + 1$  dimensions. In particular, we would like to establish whether a well defined phase transition occurs. Here we show that when the  $\mathcal{PT}$  symmetry is broken for  $2 < d < 4$  the RG flow is characterized by two fixed points rather than only one in the  $\mathcal{PT}$ -symmetric case, and that near one of them complex conjugated scaling dimensions emerge. This leads to a phase transition in  $d = 2 + 1$  dimensions, with a near critical behavior characterized by a correlation length exponent  $\nu = 3/8$ . As  $d \rightarrow 2$  the two fixed points collide, leading to a so called ‘walking’ behavior [22], sometimes also referred to as pseudocriticality, a concept that has been in focus in recent years both in high-energy and condensed matter physics [22–36]. The resulting scaling behavior for  $d \rightarrow 2$  is similar to a Berezinskii-Kosterlitz-Thouless (BKT) one. However, the actual phase transition is not a BKT one, since the RG flow features a fixed point rather than a line of fixed points. We obtain that the flow lines walk around the fixed point on RG-invariant surfaces.

*Field theory* — Let us consider a non-Hermitian theory consisting of a  $U(1)$ -invariant Lagrangian with a  $Z_N$  anisotropy perturbation,

$$\begin{aligned} \mathcal{L}_{\text{NH}} = & |\partial_\mu \psi|^2 + m^2 |\psi|^2 + \frac{u}{2} |\psi|^4 \\ & - \frac{1}{2} \left[ (v + w) \psi^N + (v - w) \psi^{*N} \right], \end{aligned} \quad (1)$$

where  $u, v, w > 0$ . The  $Z_N$  anisotropic interaction is irrelevant in the RG sense for  $N > 4$  in case the spacetime dimension lies in the interval  $2 < d < 4$ . For  $w = 0$  this expectation is corroborated by recent Monte Carlo

simulations [20]. Our main focus here will be on the  $N = 4$  case. For  $d = 2$  a special treatment is required, which is one of the main topics in the subsequent discussion. The Hermitian model for  $N = 4$  has been studied in different contexts in the past decades. For example, it arises as an effective theory emerging from breaking the  $SU(N)$  symmetry in a non-Abelian Higgs model [14] and more recently for  $N = 4$  and  $d = 3$  as theories describing the transition to valence-bond solid phases [19].

The connection to the  $N$ -state clock model is easily established after employing the following parametrization of the scalar field,  $\psi = \rho e^{i\theta}/\sqrt{2}$ , in which case the Lagrangian becomes,

$$\begin{aligned} \mathcal{L}_{\text{NH}} = & \frac{1}{2}(\partial_\mu \rho)^2 + \frac{\rho^2}{2}(\partial_\mu \theta)^2 + \frac{m^2}{2}\rho^2 + \frac{u}{8}\rho^4 \\ & - \frac{\rho^N}{2^{N/2}} [v \cos(N\theta) + iw \sin(N\theta)]. \end{aligned} \quad (2)$$

In the representation above we see that this non-Hermitian theory is  $\mathcal{PT}$ -symmetric, where  $\mathcal{P}\theta\mathcal{P}^{-1} = -\theta$  and  $\mathcal{T}i\mathcal{T}^{-1} = -i$ . The amplitude  $\rho$  is both  $\mathcal{P}$ - and  $\mathcal{T}$ -invariant.

Near criticality phase fluctuations are dominant, so we can consider the amplitude field  $\rho$  as being essentially frozen, and we can approximately write  $\rho^2 = \rho_0^2 = -2m^2/u$  for  $m^2 < 0$ . In this case we obtain the effective clock model,

$$\mathcal{L}_{\text{eff}} = \frac{K}{2}(\partial_\mu \theta)^2 - z_r \cos N\theta - iz_i \sin N\theta, \quad (3)$$

where  $K = \rho_0^2$ ,  $z_r = v\rho_0^N/2^{N/2}$ , and  $z_i = w\rho_0^N/2^{N/2}$ . For  $z_i = 0$  the model is well studied in the literature, one seminal paper being the one by José *et al.* [16], where the two-dimensional XY model with clock-type symmetry breaking fields was considered.

*Renormalization group analysis* — A major difference with respect to the theories discussed in Refs. [11] and [12] lies in the fact that the scalar field  $\theta$  is a periodic field. Hence, for  $z_i = 0$  the Lagrangian above yields the effective theory for an XY model with a  $N$ -fold clock symmetry breaking field [16].

Within a path integral formalism we can perform a shift of the field,  $\theta \rightarrow \theta + (i/N)\text{arctanh}(z_i/z_r)$  [11]. This leads us to consider the following equivalent effective Lagrangian,

$$\mathcal{L}_{\text{eff}} = \frac{K}{2}(\partial_\mu \theta)^2 - \tilde{z} \cos N\theta, \quad (4)$$

where  $\tilde{z} = \sqrt{z_r^2 - z_i^2}$ . When the  $\mathcal{PT}$ -symmetry is broken, we simply have  $\tilde{z} \rightarrow i\tilde{z}$ . Thus, we see that the couplings  $z_r$  and  $z_i$  are not actually independent. This leads to the non-renormalization of the ratio  $z_r/z_i$ , which ultimately is what will lead to the walking behavior in the  $\mathcal{PT}$  broken regime.

We will follow a procedure similar to the one by José *et al.* [16] in their analysis of the XY model in the presence of symmetry breaking fields (the simplest case refers to an external in-plane magnetic field). First, we note that when  $\tilde{z} = 0$  the effective Lagrangian (4) for  $d = 2$  is exactly dual to a sine Gordon theory of the form [16, 37],

$$\mathcal{L}_{\text{SG}} = \frac{1}{8\pi^2 K}(\partial_\mu \chi)^2 - z \cos \chi, \quad (5)$$

corresponding to the field theory of a Coulomb gas of point vortices [37]. Hence, defining the dimensionless renormalized couplings  $y$  and  $\kappa$  associated respectively to the bare couplings  $z$  and  $K$ , we obtain the usual RG equations underlying the BKT transition [16, 38].

From now on we focus on the  $N = 4$  and discuss the clock model symmetry breaking perturbation for this case. Let us define the dimensionless couplings  $y_r$  and  $y_i$  associated to  $z_r$  and  $z_i$ , respectively. The RG equations in  $d$  dimensions are given by [39],

$$\frac{d\kappa}{dl} = \frac{4}{\pi^2}(y_r^2 - y_i^2) - \kappa^2 y^2 + (d-2)\kappa, \quad (6)$$

$$\frac{dy}{dl} = [d - f(d)\kappa]y. \quad (7)$$

$$\frac{dy_r}{dl} = \left[ d - \frac{4f(d)}{\pi^2 \kappa} \right] y_r, \quad (8)$$

$$\frac{dy_i}{dl} = \left[ d - \frac{4f(d)}{\pi^2 \kappa} \right] y_i, \quad (9)$$

where  $f(d) = (d-2)\Gamma(d/2-1)/(2\pi^{d/2-2})$ .

We note that  $\ln(y_r/y_i)$  is an RG invariant, so we can write  $y_i = ky_r$ , where  $k$  is a constant. We see that for  $k = 1$  we obviously recover when  $d = 2$  the RG flow equations corresponding to a BKT behavior. In the  $\mathcal{PT}$ -symmetric case ( $y_r > y_i$  or, equivalently,  $k < 1$ ) and  $d = 2$  these equations are easily shown to be equivalent to the ones obtained in Ref. [16]. This can be seen by deriving the RG equation for  $\tilde{y} = \sqrt{y_r^2 - y_i^2}$ , which is obtained directly from Eqs. (8) and (9) as,

$$\frac{d\tilde{y}}{dl} = \left[ d - \frac{4f(d)}{\pi^2 \kappa} \right] \tilde{y}, \quad (10)$$

along with the corresponding  $d = 2$  result,  $d\tilde{y}/dl = [2 - 4/(\pi\kappa)]\tilde{y}$ .

If the  $\mathcal{PT}$  symmetry is not broken we obtain for  $d = 3$  and small  $\tilde{y}$  the usual three-dimensional XY universality class. This critical behavior is governed by the nontrivial fixed point for  $\tilde{y} = 0$  shown in panel (a) of Fig. 1. Our RG equations yield in this case the  $\epsilon$ -expansion ( $\epsilon = d - 2$ ) result  $\nu = 1/(2\sqrt{\epsilon}) + 1/8 + \mathcal{O}(\sqrt{\epsilon})$ , which reproduces for  $\epsilon = 1$  one-loop value,  $\nu \approx 0.625$  [40]. The RG flow for




FIG. 1. RG flows for the non-Hermitian system associated to the Lagrangian of Eq. (3). The RG flows shown here are for  $d = 3$ . In all panels the red dot represents the fixed point. Panels (a) and (b) represent planes featuring nontrivial fixed points arising in panels (c) and (d). In the  $\mathcal{PT}$ -symmetric case of panel (c) the only fixed point governs the XY universality class when the clock perturbation is small. In the broken  $\mathcal{PT}$  symmetry case [panel (d)], both planes of panels (a) and (b) arise, with the fixed point of panel (b) being a highly unstable repulsive spiral. In this case a weakly first-order phase transition occurs.

nonzero and large  $\tilde{y}$  exhibits a critical behavior where the four-state clock interaction becomes very large, as illustrated in panel (c) of Fig. 1. This situation is well known to lead to two decoupled Ising models [16, 20].

When the  $\mathcal{PT}$ -symmetry is broken the situation changes considerably. Since in this case  $y_r < y_i$ , Eq. (6) becomes,

$$\frac{d\kappa}{dl} = -\frac{4}{\pi^2}\tilde{y}^2 - \kappa^2 y^2 + (d-2)\kappa, \quad (11)$$

while Eqs. (7) and (10) remain unchanged. The first thing to note is that now we have two non-trivial fixed points, namely,  $P_1 = (\kappa_1, y_1, \tilde{y}_1) = (d/f(d), \sqrt{(d-2)/\kappa_1}, 0)$  and  $P_2 = (\kappa_2, y_2, \tilde{y}_2) = (4f(d)/(\pi^2 d), 0, (\pi/2)\sqrt{(d-2)\kappa_2})$ . The fixed point  $P_1$  is the same as shown in panel (a) of Fig. 1. The fixed point  $P_2$ , on the other hand, is highly unstable, corresponding to the repulsive spiral shown in panel (b) of Fig. 1, with

the full flow diagram featuring both fixed points shown in panel (d). Although the fixed point  $P_1$  is the same as the one for  $\tilde{y} = 0$  of panel (c) in Fig. 1,  $P_1$  does not govern the three-dimensional XY universality class as it is the case in panel (c). Indeed, after linearization of the RG equations in the  $\mathcal{PT}$  broken regime, we obtain the eigenvalues of the linear system,  $\lambda_0 = d - f(d)\kappa_2$  and  $\lambda_{\pm} = [d - 2 \pm i\sqrt{(d-2)(7d+2)}]/2$ . The complex conjugated eigenvalues  $\lambda_{\pm}$  are associated to the repulsive spiral fixed point  $P_2$ . The critical exponent  $\nu$  is determined by the eigenvalue  $\lambda_0$  as  $\nu = 1/\lambda_0$ . For  $d = 3$  it yields  $\nu = 3/8$  (the  $\epsilon$ -expansion practically yields the same result in this case). We note that this is smaller than the mean-field value  $1/2$ , thus indicating that a near critical behavior, or pseudocriticality, when the  $\mathcal{PT}$  symmetry is broken.

When  $d \rightarrow 2$  the fixed points  $P_1$  and  $P_2$  collide, representing a typical characteristic of walking behavior [22]. In Fig. 2 we compare the scaling behaviors of the  $\mathcal{PT}$ -symmetric with the  $\mathcal{PT}$  broken case. In the  $\mathcal{PT}$ -symmetric case both  $dy/dl$  and  $d\tilde{y}/dl$  vanish for  $\kappa_* = 2/\pi$ , leading to three lines of fixed points, as  $\tilde{y}_*^2 = y_*^2$  provides one more fixed line in addition to the two BKT ones determined by the vanishing of both  $y$  and  $\tilde{y}$ . In panels (a) and (b) of Fig. 2 we illustrate the dual RG BKT flows for the situations where  $\tilde{y} = 0$  and  $y = 0$ , respectively. Overall, however, the phase transition is not a BKT one in this case: the correlation length exhibits a power-law with a continuously varying critical exponent  $\nu(y_*)$  [16]. Panel (c) of Fig. 2 shows a view of the full RG flow diagram. In the  $\mathcal{PT}$  broken regime a BKT flow is still present for  $\tilde{y} = 0$ , but the BKT flow for  $y = 0$  disappears, as it is illustrated in panel (d) of Fig. 2. The resulting RG flow is better understood by introducing the variables  $X = 2 - \pi\kappa$ ,  $Y = 2y/\sqrt{\pi}$ , and  $\tilde{Y} = 2\tilde{y}/\sqrt{\pi}$ , in which case we can rewrite the RG equations in the form,  $dX/dl \approx -X\tilde{Y}^2 + Y^2$ ,  $dY/dl = XY$ , and  $d\tilde{Y}/dl \approx -X\tilde{Y}$ . We obtain the family of hyperboloids,  $X^2 - Y^2 + \tilde{Y}^2 = c^2 = \text{const}$ , are RG-invariants, see Fig. 2(e). At the critical point the constant  $c$  vanishes, thus defining the cone with apex at the fixed point shown in Fig. 2(f) as a separatrix. Using the hyperboloid RG invariant one can proceed to approximately solve the RG flow equations [39] to obtain the correlation length. From the solution,  $X(l) = X(0) \cos(\sqrt{2}cl)$ , the correlation length  $\xi$  is obtained by demanding that  $X(l_*) \sim \mathcal{O}(1)$  for some scale  $l_*$  such that  $\ln(\xi/a) = l_*$ . This leads to a BKT scaling,  $\xi^{-1} \sim \exp(-\text{const}/\sqrt{K - K_c})$ , where  $K_c = \kappa_*$ . The critical behavior is thus essentially governed by the BKT flow diagram of Fig. 2(b). Strictly speaking the phase transition is not exactly a BKT one though. The flow lines do not stop at the fixed line that exists for  $\tilde{Y} = 0$  when the latter is nonzero, no matter how small. They follow parabolic trajectory around the conical separatrix shown in Fig. 2(f). In other words, it is a BKT-like scaling occurring by means of a run-

ning around the fixed point, with no fixed lines present. In view of these results the  $\mathcal{PT}$  broken regime evidences a scaling behavior known as walking or pseudocritical, which was studied in detail by Gorbenko *et al.* [22] (see also Refs. [25, 28, 31, 41], a concept closely related to ‘conformality loss’ [23, 24, 26, 29, 42]).

*Conclusions and outlook* — We have determined the phase structure and critical behavior of a non-Hermitian XY model with a four-state clock perturbation both in 1+1 and 2+1 dimensions. Our RG analysis gives information about both  $\mathcal{PT}$ -symmetric and  $\mathcal{PT}$ -broken regimes, where the former phases are associated with real energy eigenvalues giving rise to a unitary time-evolution and the latter not. In view of the exact equivalence of the  $\mathcal{PT}$ -symmetric Lagrangian to a Hermitian clock model, our RG analysis in this case essentially corroborates existing numerical works for the Hermitian clock model [17, 20, 21]. On the other hand, our RG analysis also gives access to the  $\mathcal{PT}$ -broken regime and we show that its phase structure and critical behavior are very different from its  $\mathcal{PT}$ -symmetric counterpart, as discussed in the main text. The main point we make here is that the  $\mathcal{PT}$  broken regime exhibits walking (or pseudocritical) behavior in the sense defined in Ref. [22]. This analysis paves the way to consider the role of dangerously irrelevant operators [43] in non-Hermitian systems, which are pertinent for non-Hermitian deconfined quantum critical points, where in a number of cases can be qualified as pseudocritical [22, 24, 25, 28]. From an experimental point of view, we expect that the effective theories investigated here may for instance be realized using Josephson junction arrays with an asymmetric current bias [44]. Other interesting experimental possibilities arise when certain non-Hermitian conductance matrices are interpreted as non-Hermitian Hamiltonians, as recently discussed in Refs. [7, 8].

We thank Slava Rychkov for useful comments and discussions on the concept of walking. We acknowledge financial support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), through SFB 1143 project A5 and the Würzburg-Dresden Cluster of Excellence on Complexity and Topology in Quantum Matter-c.qmat (EXC 2147, Project Id No. 390858490).

## Supplemental Material

### Derivation of the RG equations

Given the renormalized counterparts  $\kappa$  and  $y$  of  $K$  and  $z$ , respectively, we obtain the usual RG equations associated to the BKT phase transition [16, 38],

$$\frac{d\kappa^{-1}}{dl} = y^2, \quad (12)$$




FIG. 2. RG flows for the non-Hermitian system associated to the Lagrangian of Eq. (3). The RG flows shown here are for  $d = 2$ . In all panels the red dot represents the fixed point  $\kappa_* = 2/\pi$  and  $\tilde{y} = y = 0$ . Panels (a) and (b) show separately the BKT planes appearing in panel (c) for the  $\mathcal{PT}$ -symmetric case. Note that these planes are dual to each other in the sense that the fixed line is for  $\kappa \geq \kappa_*$  in panel (a) and for  $\kappa \leq \kappa_*$  in panel (b). When the  $\mathcal{PT}$  symmetry is broken only the BKT plane of panel (a) survives, causing the BKT transition seems to dominate the critical behavior, as shown in panel (d). Panel (e) shows the RG-invariant surfaces. In panel (f) the conical separatrix with apex at the fixed point is shown. In both panels (e) and (f) we have defined the variables,  $X = 2 - \pi\kappa$ ,  $Y = 2y/\sqrt{\pi}$ , and  $\tilde{Y} = 2\tilde{y}/\sqrt{\pi}$ .

$$\frac{dy}{dl} = (2 - \pi\kappa)y. \quad (13)$$

Switching on the  $n = 4$  clock model perturbation modifies these RG equations considerably [16]. In order to obtain the result, we simply disregard the periodicity of  $\phi$  in Eq. (4), which makes it equivalent to an ordinary sine-Gordon theory, which is known to be exactly equivalent to a Coulomb gas in  $d$  dimensions [37]. Note that in this case we are not specializing to  $d = 2$ . The reason for this will become clear below. Importantly, we have to adapt the derivation as given in Ref. [45] such that it allows for a non-Hermitian sine-Gordon interaction. Crucial in this regard is the role played by the spatially dependent dielectric constant,

$$\varepsilon(r) = 1 + S_d \chi(r), \quad (14)$$

where  $S_d = 2\pi^{d/2}/\Gamma(d/2)$  is the surface area of the unit sphere in  $d$  dimensions and the electric susceptibility of the Coulomb gas is given by,

$$\chi(r) = S_d \int_a^r ds s^{d-1} \alpha(s) n(s). \quad (15)$$

In the above expression  $\alpha(r)$  represents the polarizability and  $n(r)$  is the local average number of dipoles. It is in this quantity that the couplings  $z_r$  and  $z_i$  appearing in Eq. (3) of the main text will enter. Approximately (i.e., for small dipoles) we have,

$$n(r) \approx (z_r^2 - z_i^2) e^{-U(r)}, \quad (16)$$

where  $U(r)$  is the renormalized electric potential. From the self-consistent analysis of  $U(r)$  the RG equations (8) and (9) in the main text are obtained for the dimensionless couplings  $y_r$  and  $y_i$  associated to  $z_r$  and  $z_i$ . Note that for broken  $\mathcal{PT}$  symmetry where  $z_r < z_i$  the density (16) becomes negative (in this sense  $n(r)$  is best interpreted as a density imbalance), and the dielectric constant  $\varepsilon(r) < 1$ , leading to an anti-screening in the Coulomb gas.

Defining  $\kappa = 4/(\pi^2 \tilde{\kappa})$  and  $\tilde{y}$  as the dimensionless renormalized counterpart of  $\tilde{z}$ , we obtain using the results of Refs. [45–47] the leading order RG equations in  $d$  dimensions in the form,

$$\frac{d\tilde{\kappa}^{-1}}{dl} = \tilde{y}^2 + (d-2)\tilde{\kappa}^{-1}, \quad (17)$$

$$\frac{d\tilde{y}}{dl} = [d - f(d)\tilde{\kappa}] \tilde{y}, \quad (18)$$

where  $f(d) = (d-2)\Gamma(d/2-1)/(2\pi^{d/2-2})$  and  $l = \ln(r/a)$  is the RG logarithmic scale.

In order to obtain the full set of RG equations for  $d > 2$  we have to surmount the technical difficulty imposed by the fact that in this case we are not dealing with an

ensemble of point vortices any longer. In fact, for  $d = 3$  the duality leads to an Abelian Higgs theory describing an ensemble of vortex loops rather than a sine-Gordon theory [48–50]. However, it has been argued in the past [51, 52] that near  $d = 2$  the RG equations for  $d > 2$  simply amount to a modification of Eq. (12) in which a term  $-(d-2)\kappa^{-1}$  is added to its right-hand side. Crucially, we note that this term has a sign opposite to the similar term appearing in Eq. (17) for the Coulomb gas case. The latter leads to the well known result that the Coulomb gas for  $d > 2$  does not undergo a phase transition, reflecting the absence of a fixed point and the runaway flow [45, 46].

Doing an  $\epsilon$ -expansion ( $\epsilon = d-2$ ) in the way just mentioned, which follows the prescription in Refs. [51, 52] has the drawback that it yields a correlation length critical exponent  $\nu = 1/(2\sqrt{\epsilon})[1 + \mathcal{O}(\sqrt{\epsilon})]$ , which does not deviate from the mean-field theory value  $\nu = 1/2$  when  $\epsilon = 1$ . However, this can be improved in the following way. We can argue that the RG equation in  $d$  dimensions for  $y$  has the same form as the one for  $\tilde{y}$  given in Eq. (18), i.e., we do not perform the  $\epsilon$ -expansion in the RG equation. This would lead to the following RG equations in absence of the symmetry breaking field,

$$\frac{d\kappa^{-1}}{dl} = y^2 - (d-2)\kappa^{-1}, \quad (19)$$

$$\frac{dy}{dl} = [d - f(d)\kappa] y. \quad (20)$$

Linearizing the above equations around the fixed point with coordinates  $\kappa_* = d/f(d)$  and  $y_* = \sqrt{(d-2)/\kappa_*}$ , we find the eigenvalues  $a_{\pm} = [2 - d \pm \sqrt{(d-2)(9d-2)}]/2$ , leading to  $1/\nu = a_+$ . While setting  $d = 3$  directly in this expression yields exactly  $\nu = 1/2$ , we have to remember that the procedure assumes that near  $d = 2$  we can still use point vortices as a leading order approximation to vortex loops. Hence, our expression for  $\nu$  should also be considered within the framework of an  $\epsilon$ -expansion. As a matter of fact, we obtain,  $\nu = 1/(2\sqrt{\epsilon}) + 1/8 + \mathcal{O}(\sqrt{\epsilon})$ , which in turns yields  $\nu \approx 0.625$  when we set  $\epsilon = 1$  at the end. This result precisely agrees with the one-loop RG result obtained from a Landau-Ginzburg-Wilson analysis of a  $|\psi|^4$  theory. There are other approaches to improve the strategy followed by Nelson and Fisher [51], like for example dealing more directly with the scaling behavior of vortex loops [53], which ultimately leads to the more accurate result,  $\nu \approx 2/3$  [40]. However, our approach has the virtue of being technically simpler, as it relies more on the original idea of Refs. [51, 52].

After the above preliminary discussion, we are now finally in position to write down the RG equations for the Lagrangian of Eq. (4), which simply amounts to combining the results of Eqs. (17) and (18) with those of Eqs. (19) and (20) to obtain,

$$\frac{d\kappa}{dl} = \frac{4}{\pi^2} \tilde{y}^2 - \kappa^2 y^2 + (d-2)\kappa, \quad (21)$$

$$\frac{d\tilde{y}}{dl} = \left[ d - \frac{4f(d)}{\pi^2\kappa} \right] \tilde{y}, \quad (22)$$

while the flow equation for  $y$  is given by Eq. (20).

### Approximate analysis of the BKT-like scaling in the $\mathcal{PT}$ broken regime

RG equations for  $\mathcal{PT}$  broken region in 1 + 1 dimensions are given by,

$$\frac{d\kappa}{dl} = -\frac{4}{\pi^2}\tilde{y}^2 - \kappa^2 y^2, \quad \frac{dy}{dl} = [2 - \pi\kappa]y, \quad \frac{d\tilde{y}}{dl} = \left[ 2 - \frac{4}{\pi\kappa} \right] \tilde{y}. \quad (23)$$

Let's introduce the new variables,

$$X = 2 - \pi\kappa, \quad Y = \frac{2}{\sqrt{\pi}}y, \quad \tilde{Y} = \frac{2}{\sqrt{\pi}}\tilde{y}. \quad (24)$$

The RG equations become

$$\frac{dX}{dl} = \tilde{Y}^2 + \left(1 - \frac{X}{2}\right)^2 Y^2 \approx \tilde{Y}^2 + Y^2, \quad \frac{dY}{dl} = XY, \quad \frac{d\tilde{Y}}{dl} = 2 \left(1 - \frac{1}{1 - \frac{X}{2}}\right) \tilde{Y} \approx -X\tilde{Y}. \quad (25)$$

Thus, the family of hyperboloid

$$X^2 - Y^2 + \tilde{Y}^2 = c^2 = \text{const} \quad (26)$$

are RG invariants. For  $X^2 + \tilde{Y}^2 < Y^2$ , we parametrize the hyperboloid of two sheets by

$$X = c \sinh u \cos v, \quad Y = c \cosh u, \quad \tilde{Y} = c \sinh u \sin v, \quad u \in \mathbb{R}^2, \quad v \in [0, \pi). \quad (27)$$

It is convenient to define new variable

$$X(l) \approx X_0 + \tilde{X}(l). \quad (28)$$

and rewrite the Eqs. (25) in terms of  $X_0$  and  $\tilde{X}$

$$\frac{d\tilde{X}}{dl} \approx \tilde{Y}^2 + Y^2, \quad \frac{dY}{dl} \approx X_0 Y, \quad \frac{d\tilde{Y}}{dl} \approx -X_0 \tilde{Y}. \quad (29)$$

In this approximation, we can obtain a solution in closed form

$$u(l) = \text{arccosh}(e^{X_0 l}), \quad v(l) = \arcsin\left(\frac{e^{-2X_0 l}}{\sqrt{e^{-2X_0 l} - 1}}\right). \quad (30)$$

In the large-distance limit,  $l$  goes to infinity. Therefore, if  $X_0$  is positive, then  $u(l)$  increases and  $v(l)$  goes to 0. As a result  $X(l)$  and  $Y(l)$  also increases, until the RG trajectory leaves the perturbative regime and  $\tilde{Y}(l)$  goes to 0. By contrast, if  $X_0 < 0$  then  $u(l)$  goes to 0 and  $v(l)$  increases. In this case, our approximation does not work, since  $v$  is limited by the interval  $[0, \pi)$ .

For  $X^2 + \tilde{Y}^2 > Y^2$ , we parametrize the hyperboloid of one sheet by

$$X = c \cosh u \cos v, \quad Y = c \sinh u, \quad \tilde{Y} = c \cosh u \sin v, \quad u \in \mathbb{R}^2, \quad v \in [0, \pi). \quad (31)$$

The solution of Eqs. (29) is then

$$u(l) = \text{arcsinh}(e^{X_0 l}), \quad v(l) = \arcsin\left(\frac{e^{-2X_0 l}}{\sqrt{X_0(e^{-2X_0 l} + 1)}}\right). \quad (32)$$

Therefore, if  $X_0 > 0$ , then we have exactly the same behavior as in the previous case.

Since the critical point is characterized by  $X = 0$ ,  $Y = 0$  and  $\tilde{Y} = 0$ , the constant  $c^2$  has to vanish there. Thus,  $c^2$  controls the distance to the critical point, which we can capture in terms of the parameter  $K$ , according to

$$c^2 = b(K - K_c), \quad (33)$$

where  $b$  is an unimportant prefactor. Using Eqs. (25), we can write

$$\frac{d^2 X}{dl^2} = 2\tilde{Y} \frac{d\tilde{Y}}{dl} + 2Y \frac{dY}{dl} = -2X(-Y^2 + \tilde{Y}^2) = -2X(c^2 - X^2) \approx -2c^2 X, \quad (34)$$

which can be satisfied by the next ansatz

$$X(l) = X(0) \cos(\sqrt{2}cl). \quad (35)$$

We evaluate the left-hand side when  $X(l)$  is of order 1. This happens at a scale  $l = l_*$  which is related to the correlation length  $\xi = ae^{l_*}$ . Equation

$$X(0) \cos(\sqrt{2}cl_*) = 1 \quad (36)$$

has a solution

$$l_* = \frac{1}{\sqrt{2b(K - K_c)}} \arccos[1/X(0)], \quad |X(0)| \geq 1 \quad (37)$$

and finally

$$\xi^{-1} = a^{-1} \exp\left(-\frac{1}{\sqrt{2b(K - K_c)}} \arccos[1/X(0)]\right). \quad (38)$$

- 
- [1] C. M. Bender and S. Boettcher, Real spectra in non-hermitian hamiltonians having  $\mathcal{PT}$  symmetry, *Phys. Rev. Lett.* **80**, 5243 (1998).
- [2] C. M. Bender, Introduction to  $\mathcal{PT}$ -symmetric quantum theory, *Contemporary Physics* **46**, 277 (2005), <https://doi.org/10.1080/00107500072632>.
- [3] Y. Ashida, Z. Gong, and M. Ueda, Non-hermitian physics, *Advances in Physics* **69**, 249 (2020), <https://doi.org/10.1080/00018732.2021.1876991>.
- [4] A. Stegmaier, S. Imhof, T. Helbig, T. Hofmann, C. H. Lee, M. Kremer, A. Fritzsche, T. Feichtner, S. Klemmt, S. Höfling, I. Boettcher, I. C. Fulga, L. Ma, O. G. Schmidt, M. Greiter, T. Kiessling, A. Szameit, and R. Thomale, Topological defect engineering and  $\mathcal{PT}$  symmetry in non-hermitian electrical circuits, *Phys. Rev. Lett.* **126**, 215302 (2021).
- [5] A. Wang, Z. Meng, and C. Q. Chen, Non-hermitian topology in static mechanical metamaterials, *Science Advances* **9**, eadf7299 (2023), <https://www.science.org/doi/pdf/10.1126/sciadv.adf7299>.
- [6] Q. Liang, D. Xie, Z. Dong, H. Li, H. Li, B. Gadway, W. Yi, and B. Yan, Dynamic signatures of non-hermitian skin effect and topology in ultracold atoms, *Phys. Rev. Lett.* **129**, 070401 (2022).
- [7] K. Ochkan, R. Chaturvedi, V. Könye, L. Veyrat, R. Giraud, D. Mailly, A. Cavanna, U. Gennser, E. M. Hankiewicz, B. Büchner, J. van den Brink, J. Dufouleur, and I. C. Fulga, Non-hermitian topology in a multi-terminal quantum hall device, *Nature Physics* [10.1038/s41567-023-02337-4](https://doi.org/10.1038/s41567-023-02337-4) (2024).
- [8] V. Könye, K. Ochkan, A. Chyzykova, J. C. Budich, J. van den Brink, I. C. Fulga, and J. Dufouleur, Non-hermitian topological ohmmeter (2023), [arXiv:2308.11367](https://arxiv.org/abs/2308.11367) [cond-mat.mes-hall].
- [9] J. Li, M. Tang, J. Duan, X. Xu, K. Xu, L. Ma, and J. Wang, Exceptional points in a spiral ring cavity for enhanced biosensing, *Journal of Lightwave Technology* **41**, 2870 (2023).
- [10] N. Hatano and D. R. Nelson, Localization transitions in non-hermitian quantum mechanics, *Phys. Rev. Lett.* **77**, 570 (1996).
- [11] C. M. Bender, H. Jones, and R. Rivers, Dual  $\mathcal{PT}$ -symmetric quantum field theories, *Physics Letters B* **625**, 333 (2005).
- [12] Y. Ashida, S. Furukawa, and M. Ueda, Parity-time symmetric quantum critical phenomena, *Nature Communications* **8** (2016).
- [13] S.-B. Zhang, M. M. Denner, T. c. v. Bzdušek, M. A. Sentef, and T. Neupert, Symmetry breaking and spectral structure of the interacting hatano-nelson model, *Phys. Rev. B* **106**, L121102 (2022).
- [14] G. 't Hooft, On the phase transition towards permanent quark confinement, *Nuclear Physics B* **138**, 1 (1978).
- [15] S. Elitzur, R. B. Pearson, and J. Shigemitsu, Phase structure of discrete abelian spin and gauge systems, *Phys. Rev. D* **19**, 3698 (1979).

- [16] J. V. José, L. P. Kadanoff, S. Kirkpatrick, and D. R. Nelson, Renormalization, vortices, and symmetry-breaking perturbations in the two-dimensional planar model, *Phys. Rev. B* **16**, 1217 (1977).
- [17] J. Hove and A. Sudbø, Criticality versus  $q$  in the  $(2+1)$ -dimensional  $Z_q$  clock model, *Phys. Rev. E* **68**, 046107 (2003).
- [18] M. Oshikawa, Ordered phase and scaling in  $Z_n$  models and the three-state antiferromagnetic potts model in three dimensions, *Phys. Rev. B* **61**, 3430 (2000).
- [19] S. Pujari, F. Alet, and K. Damle, Transitions to valence-bond solid order in a honeycomb lattice antiferromagnet, *Phys. Rev. B* **91**, 104411 (2015).
- [20] H. Shao, W. Guo, and A. W. Sandvik, Monte carlo renormalization flows in the space of relevant and irrelevant operators: Application to three-dimensional clock models, *Phys. Rev. Lett.* **124**, 080602 (2020).
- [21] P. Patil, H. Shao, and A. W. Sandvik, Unconventional  $u(1)$  to  $Z_q$  crossover in quantum and classical  $q$ -state clock models, *Phys. Rev. B* **103**, 054418 (2021).
- [22] V. Gorbenko, S. Rychkov, and B. Zan, Walking, weak first-order transitions, and complex cfts, *Journal of High Energy Physics* **2018**, 108 (2018).
- [23] D. B. Kaplan, J.-W. Lee, D. T. Son, and M. A. Stephanov, Conformality lost, *Phys. Rev. D* **80**, 125005 (2009).
- [24] F. S. Nogueira and A. Sudbø, Deconfined quantum criticality and conformal phase transition in two-dimensional antiferromagnets, *Europhysics Letters* **104**, 56004 (2013).
- [25] A. Nahum, J. T. Chalker, P. Serna, M. Ortuño, and A. M. Somoza, Deconfined quantum criticality, scaling violations, and classical loop models, *Phys. Rev. X* **5**, 041048 (2015).
- [26] F. S. Nogueira, J. van den Brink, and A. Sudbø, Conformality loss and quantum criticality in topological higgs electrodynamics in  $2+1$  dimensions, *Phys. Rev. D* **100**, 085005 (2019).
- [27] R. Ma and C. Wang, Theory of deconfined pseudocriticality, *Phys. Rev. B* **102**, 020407 (2020).
- [28] B. Ihrig, N. Zerf, P. Marquard, I. F. Herbut, and M. M. Scherer, Abelian higgs model at four loops, fixed-point collision, and deconfined criticality, *Phys. Rev. B* **100**, 134507 (2019).
- [29] H. Ma and Y.-C. He, Shadow of complex fixed point: Approximate conformality of  $q > 4$  potts model, *Phys. Rev. B* **99**, 195130 (2019).
- [30] L. Burgelman, L. Devos, B. Vanhecke, F. Verstraete, and L. Vanderstraeten, Contrasting pseudocriticality in the classical two-dimensional heisenberg and  $rp^2$  models: Zero-temperature phase transition versus finite-temperature crossover, *Phys. Rev. E* **107**, 014117 (2023).
- [31] M. Weber and M. Vojta,  $Su(2)$ -symmetric spin-boson model: Quantum criticality, fixed-point annihilation, and duality, *Phys. Rev. Lett.* **130**, 186701 (2023).
- [32] B. Hawashin, A. Eichhorn, L. Janssen, M. M. Scherer, and S. Ray, The nordic-walking mechanism and its explanation of deconfined pseudocriticality from wess-zumino-witten theory (2023), [arXiv:2312.11614 \[cond-mat.str-el\]](https://arxiv.org/abs/2312.11614).
- [33] M. Weber, Tunable quantum criticality and pseudocriticality across the fixed-point annihilation in the anisotropic spin-boson model (2024), [arXiv:2403.02400 \[cond-mat.str-el\]](https://arxiv.org/abs/2403.02400).
- [34] Z. Zhou, L. Hu, W. Zhu, and Y.-C. He, The  $SO(5)$  deconfined phase transition under the fuzzy sphere microscope: Approximate conformal symmetry, pseudo-criticality, and operator spectrum (2024), [arXiv:2306.16435 \[cond-mat.str-el\]](https://arxiv.org/abs/2306.16435).
- [35] J. L. Jacobsen and K. J. Wiese, Lattice realization of complex cfts: Two-dimensional potts model with  $q > 4$  states (2024), [arXiv:2402.10732 \[hep-th\]](https://arxiv.org/abs/2402.10732).
- [36] Y. Tang, H. Ma, Q. Tang, Y.-C. He, and W. Zhu, Reclaiming the lost conformality in a non-hermitian quantum 5-state potts model (2024), [arXiv:2403.00852 \[cond-mat.stat-mech\]](https://arxiv.org/abs/2403.00852).
- [37] J. Fröhlich and T. Spencer, On the statistical mechanics of classical coulomb and dipole gases, *Journal of Statistical Physics* **24**, 617 (1981).
- [38] D. R. Nelson and J. M. Kosterlitz, Universal jump in the superfluid density of two-dimensional superfluids, *Phys. Rev. Lett.* **39**, 1201 (1977).
- [39] See Supplemental Material at [URL will be inserted by publisher] for a derivation of this result, and where Refs. [45–49, 51–53] are cited.
- [40] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena: Fifth Edition*, International series of monographs on physics (Oxford University Press, 2021).
- [41] I. F. Herbut, Chiral symmetry breaking in three-dimensional quantum electrodynamics as fixed point annihilation, *Phys. Rev. D* **94**, 025036 (2016).
- [42] V. A. Miransky and K. Yamawaki, Conformal phase transition in gauge theories, *Phys. Rev. D* **55**, 5051 (1997).
- [43] D. J. Amit and L. Peliti, On dangerous irrelevant operators, *Annals of Physics* **140**, 207 (1982).
- [44] C. G. L. Böttcher, F. Nichele, M. Kjaergaard, H. J. Suominen, J. Shabani, C. J. Palmstrøm, and C. M. Marcus, Superconducting, insulating and anomalous metallic regimes in a gated two-dimensional semiconductor–superconductor array, *Nature Physics* **14**, 1138 (2018).
- [45] F. S. Nogueira and H. Kleinert, Compact quantum electrodynamics in  $2+1$  dimensions and spinon deconfinement: A renormalization group analysis, *Phys. Rev. B* **77**, 045107 (2008).
- [46] J. M. Kosterlitz, The  $d$ -dimensional coulomb gas and the roughening transition, *Journal of Physics C: Solid State Physics* **10**, 3753 (1977).
- [47] H. Kleinert, F. S. Nogueira, and A. Sudbø, Kosterlitz–thouless-like deconfinement mechanism in the  $(2+1)$ -dimensional abelian higgs model, *Nuclear Physics B* **666**, 361 (2003).
- [48] M. E. Peskin, Mandelstam–’t Hooft duality in abelian lattice models, *Ann. Phys. (N. Y.)* **113**, 122 (1978).
- [49] C. Dasgupta and B. I. Halperin, Phase transition in a lattice model of superconductivity, *Phys. Rev. Lett.* **47**, 1556 (1981).
- [50] H. Kleinert, *Gauge Fields in Condensed Matter: Vol. 1: Superflow and Vortex Lines (Disorder Fields, Phase Transitions) Vol. 2: Stresses and Defects (Differential Geometry, Crystal Melting)* (World Scientific, 1989).
- [51] D. R. Nelson and D. S. Fisher, Dynamics of classical XY spins in one and two dimensions, *Phys. Rev. B* **16**, 4945 (1977).
- [52] J. L. Cardy and H. W. Hamber,  $o(n)$  heisenberg model close to  $n = d = 2$ , *Phys. Rev. Lett.* **45**, 499 (1980).
- [53] S. R. Shenoy, Vortex-loop scaling in the three-dimensional xy ferromagnet, *Phys. Rev. B* **40**, 5056 (1989).