# Properties of the complementarity set for the cone of copositive matrices

#### O. I. Kostyukova<sup>a</sup>

<sup>a</sup>Institute of mathematics, Nationaly academy of sciences of Belarus, Minsk, Belarus

#### ARTICLE HISTORY

Compiled April 29, 2024

#### ABSTRACT

For a proper cone K and its dual cone  $K^*$  in  $\mathbb{R}^n$ , the complementarity set of K is defined as  $\mathbb{C}(K) = \{(x, y) : x \in K, y \in K^*, x^\top y = 0\}$ . It is known that  $\mathbb{C}(K)$  is an *n*-dimensional manifold in the space  $\mathbb{R}^{2n}$ . If K is a symmetric cone, points in  $\mathbb{C}(K)$ must satisfy at least *n* linearly independent bi-linear identities. Since this knowledge comes in handy when optimizing over such cones, it makes sense to search for similar relationships for non-symmetric cones.

In this paper, we study properties of the complementarity set for the dual cones of copositive and completely positive matrices. Despite these cones are of great interest due to their applications in optimization, they have not yet been sufficiently studied.

#### **KEYWORDS**

Copositive matrices; completely positive matrices; strict complementarity; complementarity set

**AMS CLASSIFICATION** 49N15, 90C25, 90C33, 90C46

## 1. Introduction

Let  $\mathcal{K}$  be a proper cone in  $\mathbb{R}^n$  (that is, a closed, pointed and convex cone with nonempty interior in  $\mathbb{R}^n$ ) and  $\mathcal{K}^*$  be its dual cone. The set

$$\mathbb{C}(\mathcal{K}) := \{ (x, s) : x \in \mathcal{K}, \ s \in \mathcal{K}^*, \ x^{\top} s = 0 \}$$

$$\tag{1}$$

is called the complementarity set of  $\mathcal{K}$ . It follows from the definition that  $\mathbb{C}(\mathcal{K})$  and  $\mathbb{C}(\mathcal{K}^*)$  are congruent: one can be obtained from the other by exchanging the first and last n coordinates.

The complementarity sets play a very significant role in the context of primal and dual linear optimization problems over a cone [1-7] and in complementarity problems [8,9]. The strict complementarity condition plays a central role in establishing error bounds and quantifying the sensitivity of the solution of conic problems [10].

The following theorem was proved in [11,12]

**Theorem 1.1.** For any proper cone  $\mathcal{K}$  in  $\mathbb{R}^n$ , the complementarity set  $\mathbb{C}(\mathcal{K})$  is an *n*-dimensional manifold.

CONTACT O. I. Kostyukova. Email: kostyukova@im.bas-net.by

In other words, there are *n* linearly independent functions  $f_i(x, s)$  for i = 1, ..., nsuch that in the presence of cone constraints  $x \in \mathcal{K}$ ,  $s \in \mathcal{K}^*$ , one complementarity condition  $x^{\top}s = 0$  is equivalent to *n* conditions  $f_i(x, s) = 0$  for i = 1, ..., n. These *n* equalities are the complementary relations (or conditions). The *n* functions are not unique and are characterized by the cone  $\mathcal{K}$ .

If these functions are known, then solving a conic optimization problem over the cone  $\mathcal{K}$ , we can combine the complementary equations with primal and dual feasibility equations, and get a system of equations with equal number of unknowns and equations. In the absence of various forms of degeneracy, this set of equations determines the primal and dual optimal solutions. Hence, the problem of explicit defining such functions is of great importance.

For some  $\mathcal{K} \subset \mathbb{R}^n$ , the manifold  $\mathbb{C}(\mathcal{K})$  can be described by a set of n rather simple functions  $f_i(x, s)$ , i = 1, ..., n, w.r.t. x and s, namely by n bi-linear functions  $x^{\top}Q_is$ , i = 1, ..., n, where  $\{Q_i, i = 1, ..., n\}$  is a set of linearly independent  $n \times n$ matrices. Such cones are called *perfect* (see [13]). This fact proves to be very useful when optimizing over such cones. It is known [12,13] that a symmetric cone  $\mathcal{K}$  (that is,  $\mathcal{K}$  is self-dual and its automorphism group acts transitively on its interior) is perfect. Some generalizations of the second-order cone, which are perfect, are considered in [9,14].

In general, for non-symmetric cones, the complementarity conditions cannot be represented only by bi-linear functions [12,13]. Hence, it is natural to try to find other types of functions, characterizing the complementarity conditions for non-symmetric cones. Despite the fact that such functions exist, constructing them explicitly is not an easy task. This will explain the fact that to date there are few known works devoted to this problem [12].

In this paper, we examine the complementarity conditions for the cones COP(p) and CP(p) of copositive and completely positive  $p \times p$  matrices. The cones COP(p) and CP(p) are proper cones (i.e. closed, convex, pointed, and full dimensional) and they are dual to each other. These cones are of great importance due to their applications in optimization, especially in creating convex formulations of NP-hard problems [1,15–17].

Despite their popularity in applications and the large number of studies devoted to these cones, they are still not well studied [17]. The cones COP(p) and CP(p) are complicated, they do not possess some "good" properties of the conic sets: they are neither self-dual, nor homogeneous, nor nice, and hence not facially exposed.

Notice that both cones COP(p) and CP(p) have dimension p(p+1)/2. It is known [13] that there exist only p linearly independent bi-linear functions  $f_k(X,U)$ ,  $X \in COP(p)$ ,  $U \in CP(p)$ , k = 1, ..., p, which are sequences of the complementary condition trace(XU) = 0. Hence, for these cones, the complementarity system can never be written as a square system by means of bi-linear functions alone, and it is necessary to look for other types of functions.

In the paper for the cone  $\mathcal{K} \subset \mathbb{R}^{n \times n}$  under consideration, using a pair  $(x^0, s^0) \in \mathbb{C}(\mathcal{K})$ , we introduce a set of *m* bi-linear functions  $\Omega_i(x, w)$ , i = 1, ..., m, and *n* linear functions s = Lw w.r.t. extended set of variables  $x \in \mathbb{R}^n$ ,  $s \in \mathbb{R}^n$ ,  $w \in \mathbb{R}^m$ , that satisfy the condition

$$\operatorname{rank}\left(\frac{\partial\Omega_i(x^0,w^0)}{\partial(x,w)}\right) = m$$
 where  $w^0 = Ls^0$ ,

and allow us to completely describe the set  $\mathbb{C}(\mathcal{K})$  in a neighborhood of a point  $(x^0, s^0)$ 

under some non-degeneracy assumptions.

The paper is structured as follows. Sections 1 and 2 contain an introduction, and basic notations and problem statement. In Section 3, some auxiliary results are formulated and proved. In Section 4, we study properties of the complementarity set for the cone COP(p), and based on these properties we obtain a system of equations called defining equations. This system allows us to describe the complementarity set in a neighborhood of a given point  $(X^0, U^0) \in \mathbb{C}(COP(p))$  under some assumptions. In Section 4, we show that under made assumptions, the defining equations are independent in a neighborhood of the point  $(X^0, U^0)$ . Section 6 contains some examples that show that assumptions made in the paper are essential for our study. Technical propositions are proved in appendix.

## 2. Problem statement

Let p be a positive integer and  $P = \{1, ..., p\}$ . We denote by  $\mathbb{R}^p$  the p-dimensional Euclidean vector space with standard orthogonal basis  $\{e_k, k = 1, ..., p\}$ . We use  $\mathbb{R}^p_+$  to denote the set of element-wise non-negative p-vectors. For  $t = (t_k, k \in P)^\top \in \mathbb{R}^p$ , we

denote its support as by supp $(t) := \{k \in P : t_k \neq 0\}$  and use the norm  $||t||_1 = \sum_{k=1}^p |t_k|$ .

We let  $\mathbb{S}(p)$  denote the real linear space of symmetric  $p \times p$  matrices. For  $U \in \mathbb{S}(p)$ and  $W \in \mathbb{S}(p)$ , an inner product is defined by  $U \bullet W = \text{trace}(UW)$ . For a set  $\mathcal{L} \subset \mathbb{S}(p)$ , we denote by int  $\mathcal{L}$ , ri  $\mathcal{L}$ , conv  $\mathcal{L}$ , cone  $\mathcal{L}$  the interior, relative interior, convex hull, conic hull of  $\mathcal{L}$ , respectively, and by span  $\mathcal{L}$  the space spanned by  $\mathcal{L}$  and by  $\mathcal{L}^{\perp}$  the orthogonal complement of its span. We use  $\mathbb{O}_{n \times p}$ , E(p), and **0** to denote the  $n \times p$  matrix of all zeroes, identity  $p \times p$  matrix, and a null vector. The dimension of the vector will be clear from the context.

For a matrix  $F \in \mathbb{S}(p)$ , we denote by  $F_{ij}$ ,  $i \in P$ ,  $j \in P$ , its elements and define the vector  $\operatorname{svec}(F) \in \mathbb{R}^{p(p+1)/2}$  by the rule (see [18])

$$\operatorname{svec}(F) = (F_{11}, \sqrt{2}F_{21}, ..., \sqrt{2}F_{p1}, F_{22}, \sqrt{2}F_{32}, ..., \sqrt{2}F_{p2}, ..., F_{pp})^{\top}.$$

For a set of symmetric  $p \times p$  matrices F(i),  $i \in I$ , we consider that  $\operatorname{rank}(F(i), i \in I) := \operatorname{rank}(\operatorname{svec}(F(i)), i \in I)$ .

For a cone  $\mathcal{K} \subset \mathbb{S}(p)$ , we denote by  $\mathcal{K}^*$  its dual cone

$$\mathcal{K}^* := \{ U \in \mathbb{S}(p) : U \bullet X \ge 0 \ \forall X \in \mathcal{K} \}.$$

We use  $\mathbb{S}_+(p)$  to denote the cone of symmetric semidefinite  $p \times p$  matrices and  $\mathcal{COP}(p)$  to denote the cone of copositive  $p \times p$  matrices

$$\mathcal{COP}(p) := \{ D \in \mathbb{S}(p) : t^{\top} Dt \ge 0 \ \forall t \in \mathbb{R}^p_+ \}.$$

It is easy to see that the cone  $\mathcal{COP}(p)$  can be equivalently defined as follows:

$$\mathcal{COP}(p) := \{ D \in \mathbb{S}(p) : t^{\top} Dt \ge 0 \ \forall t \in T \} \text{ where } T := \{ t \in \mathbb{R}^p_+ : ||t||_1 = 1 \}.$$

The cone  $\mathcal{COP}(p)$  is not self-dual, its dual cone is the cone of completely positive

 $p \times p$  matrices

$$\mathcal{CP}(p) := (\mathcal{COP}(p))^* = \operatorname{cone}\{t \, t^\top : t \in \mathbb{R}^p_+\}.$$

We say that matrices  $X \in \mathcal{COP}(p)$  and  $U \in \mathcal{CP}(p)$  are complementary if  $X \bullet U = 0$ .

**Definition 2.1.** (see [3]) Let  $X^0 \in COP(p)$  and  $U^0 \in CP(p)$ , we say that  $X^0$  is strictly complementary to  $U^0$  if

$$X^{0} \in \operatorname{ri}(\mathcal{COP}(p) \cap U^{0^{\perp}}), \tag{2}$$

 $U^0$  is strictly complementary to  $X^0$  if

$$U^{0} \in \operatorname{ri}(\mathcal{CP}(p) \cap X^{0^{\perp}}).$$
(3)

For the cone  $\mathcal{COP}(p)$  consider the complementarity set

$$\mathbb{C}(\mathcal{COP}(p)) := \{ (X, U) \in \mathcal{COP}(p) \times (\mathcal{COP}(p))^* : X \bullet U = 0 \} = \{ (X, U) : X \in \mathcal{COP}(p), U \in \mathcal{CP}(p), X \bullet U = 0 \}.$$

The purpose of this paper is to obtain explicit relations in the form of equations that allow us to characterize the complementarity set  $\mathbb{C}(\mathcal{COP}(p))$  in the neighborhood of a given point  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p))$  satisfying some conditions.

## 3. Auxiliary constructions and results

For a given matrix  $X^0 \in \mathcal{COP}(p)$ , denote by  $T_a(X^0)$  the set of its normalized zeroes

$$T_a(X^0) := \{ t \in T : t^\top X^0 t = 0 \}.$$

The set  $T_a(X^0)$  is empty or a union of a finite number of convex bounded polyhedra [19].

Suppose that  $T_a(X^0) \neq \emptyset$ . Denote by  $\tau(j), j \in J \subset \mathbb{N}$ , the set of all vertices of the set conv  $T_a(X)$  and define the corresponding index sets

$$M(j) := \{k \in P : e_k^\top X^0 \tau(j) = 0\}, j \in J.$$
(4)

Having  $\tau(j), M(j), j \in J$ , let us partition the index set J into maximum number of subsets  $J(s), s \in S \subset \mathbb{N}, |S| \ge 1$ , such that

**a)**  $J = \bigcup_{s \in S} J(s)$ , **b)** for any  $s \in S$ , it holds  $\bigcup_{j \in J(s)} \operatorname{supp}(\tau(j)) \subset M(i) \ \forall i \in J(s), \mathbf{c}$ ) if  $|S| \ge 2$ , then for all  $s \in S$ ,  $\bar{s} \in S$ ,  $s \ne \bar{s}$ , we have  $J(s) \setminus J(\bar{s}) \ne \emptyset$ ,  $J(\bar{s}) \setminus J(s) \ne \emptyset$ , and  $\forall i_0 \in J(s) \setminus J(\bar{s}), \exists j_0 \in J(\bar{s}) \setminus J(s) \text{ and } \exists k_0 \in \operatorname{supp}(\tau(i_0)) \text{ such that } k_0 \notin M(j_0).$  The following proposition is proved in [20].

**Proposition 3.1.** Let  $\{J(s), s \in S\}$  be the partition of the set J such that the conditions **a**) - **c**) are satisfied. Then the set  $T_a(X^0)$  can be presented in the form

$$T_a(X^0) = \bigcup_{s \in S} T_a(s, X^0), \text{ where } T_a(s, X^0) := \operatorname{conv}\{\tau(j), j \in J(s)\}, \ \forall s \in S.$$
(5)

Denote

$$P_*(s) = P_*(s, X^0) := \bigcup_{j \in J(s)} \operatorname{supp}(\tau(j)) \subset P, \ s \in S.$$
(6)

It follows from condition **b**) that

$$P_*(s) \subset M(j) \ \forall j \in J(s), \ \forall s \in S.$$

Consider a matrix  $U^0 \in C\mathcal{P}(p)$  such that  $(X^0, U^0) \in \mathbb{C}(CO\mathcal{P}(p))$ . If  $U^0 \in C\mathcal{P}(p)$ and  $X^0 \bullet U^0 = 0$ , then the matrix  $U^0$  admits a presentation

$$U^{0} = \sum_{i \in I^{*}} \alpha_{i} t(i)(t(i))^{\top} \text{ with some } \alpha_{i} > 0, \ t(i) \in T_{a}(X^{0}), \ i \in I^{*}.$$

Consequently,  $U^0$  admits a presentation

$$U^{0} = \sum_{s \in S} U^{0}(s), \ U^{0}(s) \in \mathcal{F}(s) := \operatorname{cone}\{t \, t^{\top}, t \in T_{a}(s, X^{0})\} \subset \mathcal{CP}(p) \ \forall s \in S.$$
(7)

Notice that by construction

$$U_{kq}^{0}(s) = 0 \ \forall k \in P \setminus P_{*}(s), \ \forall q \in P, \ \forall s \in S.$$
(8)

Also notice that, in general, there may exist another set of matrices  $\bar{U}^0(s) \in \mathcal{F}(s), s \in S$ , such that  $U^0$  admits a presentation  $U^0 = \sum_{s \in S} \bar{U}^0(s)$ .

Denote

$$V(I) = \{(i,j) : i \in I, j \in I, i \leq j\} \text{ for } I \subset \mathbb{N},$$
  
$$\boldsymbol{\xi}(i,j) = (\tau(i) + \tau(j)) \text{ for } (i,j) \in V(J(s)), s \in S.$$

For  $s \in S$ , let  $J_b(s) \subset J(s)$  be such that

$$\operatorname{rank}(\tau(j), j \in J(s)) = \operatorname{rank}(\tau(j), j \in J_b(s)) = |J_b(s)|.$$

It is known (see [21]) that for any  $s \in S$ , the matrices  $\boldsymbol{\xi}(i,j)\boldsymbol{\xi}(i,j)^{\top}$ ,  $(i,j) \in V(J_b(s))$  are linearly independent.

One can show that

$$\operatorname{rank}(t \, t^{\top}, t \in T_a(s, X^0)) = \operatorname{rank}(\boldsymbol{\xi}(i, j) \boldsymbol{\xi}(i, j)^{\top}, \ (i, j) \in V(J_b(s))) = |V(J_b(s))|,$$
(9)

and the following lemma holds true.

**Lemma 3.2.** Condition (3) is equivalent to the following one: matrix  $U^0$  admits a presentation (7) where matrices  $U^0(s), s \in S$ , have the form

$$U^{0}(s) = \sum_{(i,j)\in V(J(s))} \alpha_{ij} \boldsymbol{\xi}(i,j) (\boldsymbol{\xi}(i,j))^{\top} + \sum_{i\in I(s)} t(i)(t(i))^{\top}, \quad (10)$$
  
$$\alpha_{ij} > 0, (i,j) \in V(J(s)); \ t(i) \in \operatorname{cone} T_{a}(s, X^{0}), \ i \in I(s), \ s \in S.$$

In what follows, we will say that a pair  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p))$  satisfies

Assumption j), if  $U^0$  is strictly complementary to  $X^0$ ,

Assumption jj), if the following matrices are linearly independent

$$(\tau(i) + \tau(j))(\tau(i) + \tau(j))^{\top}, \ (i,j) \in V(J_b(s)), \ s \in S,$$
(11)

Assumption jjj), if  $M(j) = P_*(s) \ \forall j \in J(s), \ \forall s \in S.$ 

**Remark 1.** One can show that Assumptions j) and jjj) imply that  $X^0$  is strictly complementary to  $U^0$ .

For  $s \in S$ , using given set  $P_*(s)$  and number  $p(s) := |P_*(s)|$ , let us introduce matrix transformations

$$\mathcal{A}(X,s): \mathbb{S}(p) \to \mathbb{S}(p(s)) \text{ and } \mathcal{B}(W,s): \mathbb{S}(p(s)) \to \mathbb{S}(p)$$
 (12)

defined by the rules

$$\mathcal{A}(X,s) =: X(s) = \begin{pmatrix} x_{kq}, q \in P_*(s) \\ k \in P_*(s) \end{pmatrix}, \mathcal{B}(W,s) =: U(s) = \begin{pmatrix} u_{kq}, q \in P \\ k \in P \end{pmatrix},$$
$$u_{kq} = w_{kq} \text{ for } k \in P_*(s) \text{ and } q \in P_*(s), \ u_{kq} = 0 \text{ for } k \in P \setminus P_*(s) \text{ and } q \in P,$$

where  $x_{kq}$  is (k, q)-th element of matrix X and  $w_{kq}$  is (k, q)-th element of matrix W.

**Proposition 3.3.** Suppose that for a pair  $(X^0, U^0) \in \mathbb{C}(COP)$  Assumptions j) and jj) hold true. Then the following conditions are satisfied:

i) there exists the only set of matrices  $U^0(s)$ ,  $s \in S$ , satisfying (7),

ii) for any  $s \in S$ , matrix  $W^0(s) := \mathcal{A}(U^0(s), s)$  admits a presentation

$$W^{0}(s) = \mathcal{M}(s)(\mathcal{M}(s))^{\top} \text{ with some } \mathcal{M}(s) \in \mathbb{R}^{p(s) \times n(s)}, \ \mathcal{M}(s) > \mathbb{O}_{p(s) \times n(s)},$$
(13)

*iii)*  $\mathcal{A}(X^0, s) \in \mathbb{S}_+(p(s)), \mathcal{A}(U^0(s), s) \in \mathbb{S}_+(p(s)), \mathcal{A}(X^0, s) + \mathcal{A}(U^0(s), s) \in \mathbb{N}$ int  $\mathbb{S}_+(p(s)) \forall s \in S$ .

**Proof.** In fact, it is easy to show that Assumption jj) implies condition i).

To prove conditions ii) and iii), let us consider a fixed  $s \in S$  and denote  $X^0(s) := \mathcal{A}(X^0, s), W^0(s) := \mathcal{A}(U^0(s), s)$ . Due to Lemma 3.2, the matrix  $W^0(s)$  admits a presentation

$$W^{0}(s) = \sum_{(i,j)\in V(J(s))} \boldsymbol{\xi}_{*}(i,j)\boldsymbol{\xi}_{*}(i,j)^{\top} + \sum_{i\in I(s)} t_{*}(i)(t_{*}(i))^{\top}$$
(14)  
$$\tau_{*}(j) := (e_{i}^{\top}\tau(j), k \in P_{*}(s)) \in \mathbb{R}^{p(s)}, \ i \in J(s).$$

where

$$\alpha_{ij} > 0, \, \boldsymbol{\xi}_*(i,j) := \sqrt{\alpha_{ij}}(\tau_*(i) + \tau_*(j)) \in \mathbb{R}^{p(s)}_+, \, (i,j) \in V(J(s)),$$
$$t_*(i) \in T_*(s, X^0) := \operatorname{cone}\{\tau_*(j), j \in J(s)\} \subset \mathbb{R}^{p(s)}_+, \, i \in I(s).$$

Let us prove conditions ii). Denote

$$\hat{t} = \sum_{(i,j) \in V(J(s))} \xi_*(i,j) + \sum_{i \in I(s)} t_*(i).$$

By construction,  $\hat{t} > \mathbf{0}$  and  $\hat{t} \in T_*(s, X^0)$ . It is easy to check that

$$\sum_{\substack{(i,j)\in V(J(s))}} (\boldsymbol{\xi}_{*}(i,j) + \theta \hat{t}) (\boldsymbol{\xi}_{*}(i,j) + \theta \hat{t})^{\top} + \sum_{i\in I(s)} (t_{*}(i) + \theta \hat{t}) (t_{*}(i) + \theta \hat{t})^{\top}$$
(15)  
=  $W^{0}(s) + 2\theta \hat{t} \hat{t}^{\top} + \theta^{2} \gamma \hat{t} \hat{t}^{\top}, \ \gamma := |V(J(s))| + |I(s)|.$ 

Note that equalities (9) imply the equalities

$$\operatorname{rank}(tt^{\top}, t \in T_*(s, X^0)) = \operatorname{rank}(\boldsymbol{\xi}_*(i, j)\boldsymbol{\xi}_*(i, j)^{\top}, (i, j) \in V(J_b(s))) = |V(J_b(s))|.$$
(16)

Hence, for sufficiently small  $\theta > 0$ , we have

$$\operatorname{rank}(tt^{\top}, t \in T_*(s, X^0)) = \operatorname{rank}\left((\boldsymbol{\xi}_*(i, j) + \theta \hat{t})(\boldsymbol{\xi}_*(i, j) + \theta \hat{t})^{\top}, (i, j) \in V(J_b(s))\right).$$

Taking into account this equality, equalities (16) and the inclusion  $\hat{t} \in T_*(s, X^0)$ , we obtain that

$$\hat{t}\hat{t}^{\top} = \sum_{(i,j)\in V(J_b(s))} \beta_{ij}(\theta) (\boldsymbol{\xi}_*(i,j) + \theta\hat{t}) (\boldsymbol{\xi}_*(i,j) + \theta\hat{t})^{\top}, \qquad (17)$$
$$\hat{t}\hat{t}^{\top} = \sum_{(i,j)\in V(J_b(s))} \beta_{ij} \boldsymbol{\xi}_*(i,j) \boldsymbol{\xi}_*(i,j)^{\top}, \ \beta_{ij}(\theta) = \beta_{ij} + O(\theta), \ (i,j) \in V(J_b(s)).$$

Hence, it follows from (15) and (17) that

$$W^{0}(s) = \sum_{(i,j)\in V(J(s))} (\boldsymbol{\xi}_{*}(i,j) + \theta \hat{t}) (\boldsymbol{\xi}_{*}(i,j) + \theta \hat{t})^{\top} + \sum_{i\in I(s)} (t_{*}(i) + \theta \hat{t}) (t_{*}(i) + \theta \hat{t})^{\top} - (2\theta + \theta^{2}\gamma) \sum_{(i,j)\in V(J_{b}(s))} \beta_{ij}(\theta) (\boldsymbol{\xi}_{*}(i,j) + \theta \hat{t}) (\boldsymbol{\xi}_{*}(i,j) + \theta \hat{t}) (\boldsymbol{\xi}_{*}(i,j) + \theta \hat{t})^{\top} = \sum_{(i,j)\in V(J(s))} \mu_{ij}(\theta) (\boldsymbol{\xi}_{*}(i,j) + \theta \hat{t}) (\boldsymbol{\xi}_{*}(i,j) + \theta \hat{t})^{\top} + \sum_{i\in I(s)} (t_{*}(i) + \theta \hat{t}) (t_{*}(i) + \theta \hat{t})^{\top}$$

where  $\mu_{ij}(\theta) = 1 - (2\theta + \theta^2 \gamma)\beta_{ij}(\theta), (i, j) \in V(J_b(s)), \ \mu_{ij}(\theta) = 1, \ (i, j) \in V(J(s)) \setminus V(J_b(s))$ . Notice that for sufficiently small  $\theta > 0$  the following inequalities hold true

$$\mu_{ij}(\theta) > 0, \, \boldsymbol{\xi}_*(i,j) + \theta \hat{t} > \mathbf{0} \,\,\forall (i,j) \in V(J(s)); \,\, t_*(i) + \theta \hat{t} > \mathbf{0} \,\,\forall i \in I(s).$$

Hence we have shown that the matrix  $W^0(s)$  admits a presentation (13) with

$$\mathcal{M}(s) = \left(\sqrt{\mu_{ij}(\theta)}(\boldsymbol{\xi}_*(i,j) + \theta\hat{t}), (i,j) \in V(J(s)); \ t_*(i) + \theta\hat{t}, i \in I(s)\right).$$

Let us prove conditions iii). By construction,  $W^0(s) \in \mathcal{CP}(p(s)), X^0(s) \in \mathcal{COP}(p(s))$ and  $X^0(s)\bar{t} = \mathbf{0}$  where  $\bar{t} = \sum_{j \in J(s)} \tau_*(j) \in \mathbb{R}^{p(s)}, \bar{t} > \mathbf{0}$ . Hence

$$X^{0}(s) \in \mathbb{S}_{+}(p(s)), \ W^{0}(s) \in \mathbb{S}_{+}(p(s)) \implies X^{0}(s) + W^{0}(s) \in \mathbb{S}_{+}(p(s)).$$

Let  $\tilde{t} \in \mathbb{R}^{p(s)}$  be such that  $\tilde{t}^{\top}(X^0(s) + W^0(s))\tilde{t} = 0$ . This implies

$$\tilde{t}^{\top} X^0(s) \tilde{t} = 0, \ \tilde{t}^{\top} W^0(s) \tilde{t} = 0 \implies X^0(s) \tilde{t} = \mathbf{0}, \ W^0(s) \tilde{t} = \mathbf{0}.$$

If  $X^0(s)\tilde{t} = \mathbf{0}$ , then  $\tilde{t} \in H(X^0(s)) := \{t \in \mathbb{R}^{p(s)} : X^0(s)t = \mathbf{0}\}$ . Note that  $X^0(s) \in \mathbb{S}_+(p(s))$  and  $X^0(s)\bar{t} = \mathbf{0}, \bar{t} > \mathbf{0}$ . Hence it follows from Proposition A.4 that

$$H(X^{0}(s)) = \operatorname{span}\{\tau_{*}(j), j \in J(s)\}.$$
(18)

This implies that  $\tilde{t} = \sum_{j \in J(s)} \beta_j \tau_*(j)$ . Then taking into account presentation (14), we obtain  $0 = \tilde{t}^\top W^0(s) \tilde{t} = \sum_{(i,j) \in V(J(s))} \alpha_{ij} (\tilde{t}^\top (\tau_*(i) + \tau_*(j)))^2 + \sum_{i \in I(s)} (\tilde{t}^\top t_*(i))^2$ . Consequently,

$$\tilde{t}^{\top}(\tau_*(i) + \tau_*(j)) = 0 \ \forall (i,j) \in V(J(s)) \implies \tilde{t}^{\top}\tau_*(j) = 0 \ \forall j \in J(s).$$

It follows from the latter equalities and (18) that  $\tilde{t} \in (H(X^0(s)))^{\perp}$ . This inclusion and inclusion  $\tilde{t} \in (H(X^0(s)))$  proved above imply the equality  $\tilde{t} = \mathbf{0}$ . Thus we have proved that condition iii) holds true.

To prove the next proposition, we will use the following lemma proved in [22].

**Lemma 3.4.** Let  $\mathcal{M}, \mathcal{L} \in \mathbb{R}^{p \times m}$ . Then  $\mathcal{L}\mathcal{L}^{\top} = \mathcal{M}\mathcal{M}^{\top}$  iff there exists an orthogonal matrix  $\Omega \in \mathbb{R}^{m \times m}$  such that  $\mathcal{L}\Omega = \mathcal{M}, \mathcal{L} = \mathcal{M}\Omega^{\top}$ .

**Proposition 3.5.** Let  $\varepsilon_0 > 0$  be a sufficiently small number. Suppose that for  $s \in S$ , matrix  $W^0(s) \in C\mathcal{P}(p(s))$  satisfies relations (13) and matrices  $W(s,\varepsilon) \in \mathbb{S}_+(p(s))$ ,  $\varepsilon \in [0,\varepsilon_0]$ , are such that  $W(s,\varepsilon) \to W^0(s)$  as  $\varepsilon \to 0$ . Then for  $s \in S$  and  $\varepsilon \in [0,\varepsilon_0]$ , the matrix  $W(s,\varepsilon)$  admits a presentation

$$W(s,\varepsilon) = \mathcal{M}(s,\varepsilon)(\mathcal{M}(s,\varepsilon))^{\top} \text{ with } \mathcal{M}(s,\varepsilon) \in \mathbb{R}^{p(s) \times n(s)}, \ \mathcal{M}(s,\varepsilon) > \mathbb{O}_{p(s) \times n(s)}.$$
(19)

**Proof.** Let us consider a fixed  $s \in S$ . By assumption  $W(s,\varepsilon) \in \mathbb{S}_+(p(s)), \varepsilon \in [0,\varepsilon_0]$ . Hence there exist matrices  $B(s,\varepsilon) \in \mathbb{R}^{p(s) \times p(s)}$  such that  $||B(s,\varepsilon)|| < \infty$ ,  $W(s,\varepsilon) = B(s,\varepsilon)B^{\top}(s,\varepsilon)$  for  $\varepsilon \in [0,\varepsilon_0]$ . Let B(s,0) be any limit point of the sequence  $B(s,\varepsilon)$  as  $\varepsilon \to 0$ . Then taking into account that  $W(s,\varepsilon) \to W^0(s)$  as  $\varepsilon \to 0$ , we obtain that  $W^0(s) = B(s,0)B^{\top}(s,0)$ .

On the other hand, it follows from relations (13) that  $W^0(s) = \mathcal{M}(s)\mathcal{M}^{\top}(s)$ , where  $\mathcal{M}(s) \in \mathbb{R}^{p(s) \times n(s)}$ ,  $\mathcal{M}(s) > \mathbb{O}_{p(s) \times n(s)}$ . Without loss of generality, we may suppose that  $n(s) \ge p(s)$  since otherwise one can extend the matrix  $\mathcal{M}(s)$  with columns  $\mathcal{M}_i(s)$ , i = 1, ..., n(s), by replacing one column  $\mathcal{M}_1(s)$  with p(s) - n(s) + 1 identical columns  $\mathcal{M}_1(s)/\sqrt{p(s) - n(s) + 1}$ .

Thus we have

$$W(s,\varepsilon) = \bar{B}(s,\varepsilon)\bar{B}(s,\varepsilon)^{\top}, \ W^{0}(s) = \bar{B}(s,0)\bar{B}(s,0)^{\top} = \mathcal{M}(s)\mathcal{M}(s)^{\top} \text{ where} \\ \bar{B}(s,\varepsilon) := [B(s,\varepsilon), \mathbb{O}_{p(s)\times(n(s)-p(s))}] \text{ if } n(s) > p(s), \bar{B}(s,\varepsilon) := B(s,\varepsilon) \text{ if } n(s) = p(s).$$

Applying Lemma 3.4, we conclude that there exists orthogonal matrix  $\Omega \in \mathbb{R}^{n(s) \times n(s)}$ such that  $\bar{B}(s, 0)\Omega = \mathcal{M}(s)$ .

Let us set  $\mathcal{M}(s,\varepsilon) := B(s,\varepsilon)\Omega$ . Since  $\mathcal{M}(s) > \mathbb{O}_{p(s)\times n(s)}$  and  $\mathcal{M}(s,\varepsilon) = \mathcal{M}(s) + O(\varepsilon)$ , we conclude that  $\mathcal{M}(s,\varepsilon) > \mathbb{O}_{p(s)\times n(s)}$  for sufficiently small  $\varepsilon \geq 0$ . Relations (19) follows from the equalities  $W(s,\varepsilon) = \bar{B}(s,\varepsilon)\bar{B}(s,\varepsilon)^{\top} = \bar{B}(s,\varepsilon)\Omega\Omega^{\top}\bar{B}(s,\varepsilon)^{\top} = \mathcal{M}(s,\varepsilon)\mathcal{M}(s,\varepsilon)^{\top}$ .

## 4. System of defining equations for complementarity set

The aim of this section is to obtain a system of equations that allows us to describe the complementarity set for the cone  $\mathcal{COP}(p)$  in a neighborhood of a given point  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p)).$ 

**Theorem 4.1.** Suppose that for a pair  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p))$  Assumptions j)-jjjhold true. Then for any  $(X(\varepsilon), U(\varepsilon)) \in \mathbb{C}(\mathcal{COP}(p))$  such that  $X(\varepsilon) \to X^0$ ,  $U(\varepsilon) \to U^0$  as  $\varepsilon \to 0$ , there exist  $\varepsilon_0 > 0$  and matrix functions  $W(s, \varepsilon) \in \mathbb{S}(p(s))$ ,  $s \in S$ ,  $\varepsilon \in [0, \varepsilon_0]$ , such that

$$\mathcal{A}(X(\varepsilon), s)W(s, \varepsilon) + W(s, \varepsilon)\mathcal{A}(X(\varepsilon), s) = \mathbb{O}_{p(s) \times p(s)} \ \forall s \in S,$$
(20)

$$U(\varepsilon) = \sum_{s \in S} \mathcal{B}(W(s,\varepsilon), s).$$
<sup>(21)</sup>

**Proof.** For  $A \subset \mathbb{R}^p$ ,  $B \subset \mathbb{R}^p$ , and  $t \in \mathbb{R}^p$ , denote

$$\rho(t,B) := \min_{\tau \in B} ||t - \tau||, \ \delta(A,B) := \max_{t \in A} \rho(t,B)$$

It is not difficult to show that Assumption jjj) implies that

$$T_a(s, X^0) \cap T_a(\bar{s}, X^0) = \emptyset \ \forall s \in S, \ \forall \bar{s} \in S, \ s \neq \bar{s},$$
$$\nu_{kj} := e_k^\top X^0 \tau(j) > 0 \ \forall k \in P \setminus P_*(s), \ \forall j \in J(s), \ \forall s \in S.$$
(22)

Hence there exists  $\gamma > 0$  such that

$$\{t \in T : \rho(t, T_a(s, X^0)) \le \gamma\} \cap \{t \in T : \rho(t, T_a(\bar{s}, X^0)) \le \gamma\} = \emptyset \ \forall s \in S, \ \forall \bar{s} \in S, \ s \neq \bar{s}.$$
(23)

Set

$$\delta(\varepsilon) := \delta(T_a(X(\varepsilon)), T_a(X^0)), \ t(\varepsilon) := \arg \max_{t \in T_a(X(\varepsilon))} \rho(t, T_a(X^0)).$$
(24)

By construction,

$$\delta(\varepsilon) = \rho(t(\varepsilon), T_a(X^0)), \ t(\varepsilon) \in T_a(X(\varepsilon)) \subset T.$$
(25)

Let  $\varepsilon_k, k = 1, 2, ..., be$  a sequence such that  $\varepsilon_k \to 0$  as  $k \to \infty$ . Let  $\delta_*$  and  $t^*$  be any limit points of the sequences  $\delta(\varepsilon_k), t(\varepsilon_k), k = 1, 2, ...,$  respectively. Notice that such points exist. Then it follows from (24), (25) and condition  $X(\varepsilon) \to X^0$  as  $\varepsilon \to 0$  that  $\delta_* = \rho(t^*, T_a(X^0))$  and  $t^* \in T_a(X^0)$ . It follows from these relations that  $\delta_* = 0$ . Thus we have shown that for any sequence  $\varepsilon_k, k = 1, 2, ...,$  such that  $\varepsilon_k \to 0$  as  $k \to \infty$  we have

$$\delta(\varepsilon_k) \to 0 \text{ as } k \to \infty.$$
 (26)

Denote

$$T_{a}(\varepsilon) := T_{a}(X(\varepsilon)), \ T_{a}(s,\varepsilon) := \{t \in T_{a}(\varepsilon) : \rho(t,T_{a}(s,X^{0})) \le \delta(\varepsilon)\}, \ s \in \bar{S}, \quad (27)$$
$$\bar{S} := \{s \in S : T_{a}(s,\varepsilon) \ne \emptyset\}.$$

Due to (23), (24) and (26), for sufficiently small  $\varepsilon \geq 0$ , we obtain

$$T_a(\varepsilon) = \bigcup_{s \in S} T_a(s,\varepsilon), \ T_a(s,\varepsilon) \cap T_a(\bar{s},\varepsilon) = \emptyset \ \forall s \in S, \ \forall \bar{s} \in S, \ s \neq \bar{s}.$$
(28)

Let us fix  $s \in \overline{S}$  and consider any vector  $\eta(s,\varepsilon) \in T_a(s,\varepsilon)$ . By assumption,  $X(\varepsilon) \in COP(p)$  and  $\eta(s,\varepsilon) \in T_a(\varepsilon)$ , hence

$$(\eta(s,\varepsilon))^{\top}X(\varepsilon)\eta(s,\varepsilon) = 0 \implies e_k^{\top}X(\varepsilon)\eta(s,\varepsilon) \begin{cases} = 0 \text{ if } e_k^{\top}\eta(s,\varepsilon) > 0, \\ \ge 0 \text{ if } e_k^{\top}\eta(s,\varepsilon) = 0, \end{cases} \forall k \in P.$$
(29)

Suppose that there exist  $k_0 \in P \setminus P_*(s)$  and a sequence  $\varepsilon_k, k = 1, 2, ...,$  such that  $\varepsilon_k \to 0$  as  $k \to \infty$  and  $e_{k_0}^\top \eta(s, \varepsilon_k) > 0 \ \forall k = 1, 2, ...$  Then due to (29) we have

$$e_{k_0}^{\top} X(\varepsilon_k) \eta(s, \varepsilon_k) = 0 \ \forall k = 1, 2, \dots$$

This implies that

$$e_{k_0}^{\top} X^0 \eta^*(s) = 0 \tag{30}$$

where  $\eta^*(s)$  is a limit point of the sequence  $\eta(s, \varepsilon_k)$ , k = 1, 2, ... It follows from (26), (27) that  $\eta^*(s) \in T_a(s, X^0)$  and, consequently,

$$\eta^*(s) = \sum_{j \in J(s)} \alpha_j \tau(j), \ \alpha_j \ge 0, \ j \in J(s), \ \sum_{j \in J(s)} \alpha_j = 1.$$

Taking into account these relations, inequalities (22), and equality (30), we obtain

$$0 = e_{k_0}^{\top} X^0 \eta^*(s) = \sum_{j \in J(s)} \alpha_j e_{k_0}^{\top} X^0 \tau(j) \ge \min_{j \in J(s)} \nu_{k_0 j} > 0.$$

We have obtained the contradictory relationships.

Thus we have shown that for sufficiently small  $\varepsilon \geq 0$ , the following equalities hold true

$$e_k^{\top}\eta(s,\varepsilon) = 0 \ \forall k \in P \setminus P_*(s), \ \forall \eta(s,\varepsilon) \in T_a(s,\varepsilon), \ \forall s \in \bar{S}.$$
(31)

By construction,  $(X(\varepsilon), U(\varepsilon)) \in \mathbb{C}(\mathcal{COP}(p))$ , hence matrix  $U(\varepsilon)$  admits a presentation

$$U(\varepsilon) = \sum_{i=1}^{p_*} \alpha_i(\varepsilon) t(i,\varepsilon) (t(i,\varepsilon))^\top, \, \alpha_i(\varepsilon) \ge 0, \, t(i,\varepsilon) \in T_a(\varepsilon), \, i = 1, ..., p_*, \, p_* = \frac{p(p+1)}{2}.$$

Then taking into account (27) (28), we conclude that  $U(\varepsilon)$  admits a presentation  $U(\varepsilon) = \sum_{s \in S} U(s, \varepsilon)$  where  $U(s, \varepsilon) \in \mathcal{F}(s, \varepsilon) := \operatorname{cone}\{t t^{\top}, t \in T_a(s, \varepsilon)\} \in \mathcal{CP}(p), \ s \in \overline{S}, \ U(s, \varepsilon) = \mathbb{O}_{p \times p}, \ s \in S \setminus \overline{S}.$  It follows from (31) that by construction, for these matrices, the following equalities hold true:

$$U_{kq}(s,\varepsilon) = 0 \ \forall k \in P \setminus P_*(s), \ \forall q \in P, \ \forall s \in S.$$

For  $s \in S$ , using matrices  $U(s, \varepsilon) \in \mathcal{CP}(p)$  and  $U^0(s) \in \mathcal{CP}(p)$  let us define matrices

$$W(s,\varepsilon) := \mathcal{A}(U(s,\varepsilon),s) \in \mathcal{CP}(p(s)), \ W^0(s) := \mathcal{A}(U^0(s),s) \in \mathcal{CP}(p(s)).$$

It is evident that relations (21) hold true with the matrices  $W(s,\varepsilon)$ ,  $s \in S$ .

Now let us show that  $\overline{S} = S$  and equalities (20) hold true. It follows from Assumptions j), jj) and Proposition 3.3 that conditions i) and ii) hold true. Hence, for all  $s \in S$ ,  $W(s, \varepsilon) \to W^0(s)$  as  $\varepsilon \to 0$  and matrix  $W^0(s)$  admits a presentation (13). Then  $W(s, \varepsilon) \neq \mathbb{O}_{p(s) \times p(s)}$  for all  $s \in S$ , and consequently  $\overline{S} = S$ . Moreover, we have shown that for sufficiently small  $\varepsilon_0 > 0$ ,  $W(s, \varepsilon) \in \mathcal{CP}(p(s)) \subset \mathbb{S}_+(p(s)) \forall \varepsilon \in [0, \varepsilon_0]$ . Consequently, it follows from Proposition 3.5 that for  $s \in S$ , the matrices  $W(s, \varepsilon)$ ,  $\varepsilon \in [0, \varepsilon_0]$ , admit presentations (19).

By assumption,  $(X(\varepsilon), U(\varepsilon)) \in \mathbb{C}(\mathcal{COP}(p))$ , hence

$$X(s,\varepsilon) := \mathcal{A}(X(\varepsilon),s) \in \mathcal{COP}(p(s)) \ \forall s \in S,$$

$$X(\varepsilon) \bullet U(\varepsilon) = 0 \implies X(\varepsilon) \bullet U(s,\varepsilon) = 0 \ \forall s \in S, \iff$$

$$X(s,\varepsilon) \bullet W(s,\varepsilon) = 0 \iff \sum_{i=1}^{n(s)} (\mathcal{M}_i(s,\varepsilon))^\top X(s,\varepsilon) \mathcal{M}_i(s,\varepsilon) = 0 \ \forall s \in S,$$
(32)
$$(32)$$

where for  $s \in S$ ,  $\mathcal{M}_i(s,\varepsilon)$  is *i*-th column of the matrix  $\mathcal{M}(s,\varepsilon)$ . It follows from the latter equalities and inclusions (32) that

$$(\mathcal{M}_i(s,\varepsilon))^\top X(s,\varepsilon)\mathcal{M}_i(s,\varepsilon) = 0, \ \mathcal{M}_i(s,\varepsilon) > 0 \ \forall i = 1, ..., n(s), \ \forall s \in S \Longrightarrow$$
$$X(s,\varepsilon) \in \mathbb{S}_+(p(s)) \ \forall s \in S.$$
(34)

Equalities (20) follow from (33), (34) and inclusions  $W(s,\varepsilon) \in \mathcal{CP}(p(s)) \subset \mathbb{S}_+(p(s))$  for all  $s \in S$ .

**Theorem 4.2.** Suppose that for a pair  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p))$  Assumptions *j*)-*jjj*) hold true. Let  $X(\varepsilon) \in \mathbb{S}(p)$  and  $W(s, \varepsilon) \in \mathbb{S}(p(s)), s \in S$ , be such that

$$X(\varepsilon) \to X^0, \ W(s,\varepsilon) \to W^0(s) := \mathcal{A}(U^0(s),s) \ as \ \varepsilon \to 0 \ \forall s \in S$$

where  $U^0(s)$ ,  $s \in S$ , are as in (7), and equality (20) holds. Then for  $\varepsilon \in [0, \varepsilon_0]$ , where  $\varepsilon_0 > 0$  is a sufficiently small number, the following conditions are fulfilled

$$X(\varepsilon) \in \mathcal{COP}(p), \ W(s,\varepsilon) \in \mathcal{CP}(p(s)) \ \forall s \in S,$$
(35)

$$U(\varepsilon) := \sum_{s \in S} \mathcal{B}(W(s,\varepsilon), s) \in \mathcal{CP}(p), \ X(\varepsilon) \bullet U(\varepsilon) = 0,$$
(36)

and consequently  $(X(\varepsilon), U(\varepsilon)) \in \mathbb{C}(\mathcal{COP}(p)).$ 

**Proof.** Notice that it follows from Assumptions j)-jj) and Proposition 3.3 that conditions i)-iii) hold true.

First, let us set  $X(s,\varepsilon) = \mathcal{A}(X(\varepsilon),s)$  and prove the following relations

$$X(s,\varepsilon)W(s,\varepsilon) = \mathbb{O}_{p(s)\times p(s)},\tag{37}$$

$$X(s,\varepsilon) \in \mathbb{S}_+(p(s)), \ W(s,\varepsilon) \in \mathbb{S}_+(p(s)) \ \forall s \in S, \ \forall \varepsilon \in [0,\varepsilon_0].$$
(38)

By condition iii), we have  $X^0(s) + W^0(s) \in \operatorname{int} \mathbb{S}_+(p(s)) \ \forall s \in S$  and by assumption  $X(s,\varepsilon) \to X^0(s), \ W(s,\varepsilon) \to W^0(s) \ \forall s \in S \text{ as } \varepsilon \to 0.$  This implies that

$$X(s,\varepsilon) + W(s,\varepsilon) \in \operatorname{int} \mathbb{S}_+(p(s)) \ \forall s \in S, \ \forall \varepsilon \in [0,\varepsilon_0].$$

It follows from these inclusions, equalities (20) and Proposition A.1 that relations (37), (38) hold true.

It follows from condition ii) that for all  $s \in S$ , matrix  $W^0(s)$  admits a presentation (13), and by assumption  $W(s,\varepsilon) \to W^0(s)$  as  $\varepsilon \to 0$ , and we have shown that  $W(s,\varepsilon) \in \mathbb{S}_+(p(s)), \varepsilon \in [0,\varepsilon_0]$ . Consequently, it follows from Proposition 3.5 that for any  $s \in S$ , the matrices  $W(s,\varepsilon), \varepsilon \in [0,\varepsilon_0]$ , admit presentations (19) and hence the latter inclusions in (35) hold true. This implies that

$$U(\varepsilon) := \sum_{s \in S} \mathcal{B}(W(s,\varepsilon),s) = \sum_{s \in S} U(s,\varepsilon) \in \mathcal{CP}(p) \text{ with } U(s,\varepsilon) := \mathcal{B}(W(s,\varepsilon),s) \ \forall s \in S.$$

Notice that due to specific structure of matrix  $U(s,\varepsilon)$  we have that  $X(\varepsilon) \bullet U(s,\varepsilon) =$  $X(s,\varepsilon) \bullet W(s,\varepsilon)$  for all  $s \in S$ . Also notice that it was proved above (see (37)) that  $X(s,\varepsilon) \bullet W(s,\varepsilon) = 0$  for all  $s \in S$ . Taking into account these observations, we conclude that  $X(\varepsilon) \bullet U(\varepsilon) = 0$  and relations (36) are proved.

Now let us show that  $X(\varepsilon) \in \mathcal{COP}(p)$  for sufficiently small  $\varepsilon \geq 0$ . Denote

$$\tau(\varepsilon) := \arg\{\min \tau^\top X(\varepsilon)\tau, \text{ s.t. } \tau \in T\}.$$

Suppose that there exists a sequence  $\varepsilon_n$ ,  $n = 1, 2, ..., \varepsilon_n \to 0$  as  $n \to \infty$ , such that  $X(\varepsilon_n) \notin \mathcal{COP}(p)$ , hence

$$\tau(\varepsilon_n)^\top X(\varepsilon_n) \tau(\varepsilon_n) < 0.$$
(39)

To simplify the presentation, we will denote  $\varepsilon_n$  by  $\varepsilon$  and suppose that  $\varepsilon \to 0$ . Let  $\tau^*$  be a limit point of the sequence  $\tau(\varepsilon)$  as  $\varepsilon \to 0$ . Hence, we obtain that

$$0 \le \tau^{*\top} X^0 \tau^* = \lim_{\varepsilon \to 0} \tau(\varepsilon)^\top X(\varepsilon) \tau(\varepsilon) \le 0 \implies \tau^{*\top} X^0 \tau^* = 0.$$

This implies that  $\tau^* \in T_a(s, X^0)$  for some  $s \in S$ . Below we will consider this fixed s. Let us partition vectors  $\tau(\varepsilon) = (\tau_k(\varepsilon), k \in P)^\top$  and  $\tau^* = (\tau_k^*, k \in P)^\top$ , and matrices  $X(\varepsilon), \Delta X(\varepsilon) := X(\varepsilon) - X^0$  and  $X^0$  w.r.t. the partition  $P_*(s) \cup (P \setminus P_*(s))$  of the set P as follows

$$\tau_*(\varepsilon) := (\tau_k(\varepsilon), k \in P_*(s)), \ \tau_0(\varepsilon) := (\tau_k(\varepsilon), k \in P \setminus P_*(s)), \ \tau_*^* := (\tau_k^*, k \in P_*(s)),$$

$$X(\varepsilon) = \begin{pmatrix} X_*(\varepsilon) & X_{*0}(\varepsilon) \\ (X_{*0}(\varepsilon))^\top & X_0(\varepsilon) \end{pmatrix}, \Delta X(\varepsilon) = \begin{pmatrix} \Delta X_*(\varepsilon) & \Delta X_{*0}(\varepsilon) \\ (\Delta X_{*0}(\varepsilon))^\top & \Delta X_0(\varepsilon) \end{pmatrix},$$

$$X^{0} = \begin{pmatrix} X^{0}_{*} & X^{0}_{*0} \\ (X^{0}_{*0})^{\top} & X^{0}_{0} \end{pmatrix}.$$

Then

$$\tau(\varepsilon) = \begin{pmatrix} \tau_*(\varepsilon) \\ \tau_0(\varepsilon) \end{pmatrix}, \tau^* = \begin{pmatrix} \tau^*_* \\ \mathbf{0} \end{pmatrix}, \ \tau_*(\varepsilon) \to \tau^*_*, \ \tau_0(\varepsilon) \to \mathbf{0} \text{ as } \varepsilon \to 0.$$

Notice that here  $X_*(\varepsilon) = X(s,\varepsilon)$  with s under consideration. Using this notation, let us calculate

$$\tau(\varepsilon)^{\top}X(\varepsilon)\tau(\varepsilon) = \begin{pmatrix} \tau_{*}(\varepsilon) \\ \mathbf{0} \end{pmatrix}^{\top}X(\varepsilon) \begin{pmatrix} \tau_{*}(\varepsilon) \\ \mathbf{0} \end{pmatrix}^{\top}X(\varepsilon) \begin{pmatrix} \mathbf{0} \\ \tau_{0}(\varepsilon) \end{pmatrix} =$$
$$\tau_{*}(\varepsilon)^{\top}X_{*}(\varepsilon)\tau_{*}(\varepsilon) + 2(\tau_{*}^{*} + \Delta\tau_{*}(\varepsilon))^{\top}(X_{*0}^{0} + \Delta X_{*0}(\varepsilon))\tau_{0}(\varepsilon) + \tau_{0}(\varepsilon)^{\top}X_{0}(\varepsilon)\tau_{0}(\varepsilon) =$$
$$\tau_{*}(\varepsilon)^{\top}X_{*}(\varepsilon)\tau_{*}(\varepsilon) + \tau_{0}(\varepsilon)^{\top}X_{0}(\varepsilon)\tau_{0}(\varepsilon) +$$
$$2\tau_{*}^{*\top}X_{*0}^{0}\tau_{0}(\varepsilon) + 2\tau_{*}^{*\top}\Delta X_{*0}(\varepsilon)\tau_{0}(\varepsilon) + 2\Delta\tau_{*}(\varepsilon)^{\top}X_{*0}^{0}\tau_{0}(\varepsilon) + 2\Delta\tau_{*}(\varepsilon)^{\top}\Delta X_{*0}(\varepsilon)\tau_{0}(\varepsilon),$$

where  $\Delta \tau_*(\varepsilon) = \tau_*(\varepsilon) - \tau_*^*$ .

If  $\tau_0(\varepsilon) \equiv 0$  for all  $\varepsilon \to 0$ , then, taking into account that  $X_*(\varepsilon) = X(s, \varepsilon) \in \mathbb{S}_+(p(s))$ (see (38)), we have  $\tau(\varepsilon)^{\top} X(\varepsilon) \tau(\varepsilon) = \tau_*(\varepsilon)^{\top} X_*(\varepsilon) \tau_*(\varepsilon) \ge 0$ . But this contradicts (39).

Now suppose that  $\tau_0(\varepsilon) \neq 0$ . Hence, there exists

$$\Delta \tau_0 = \lim_{\varepsilon \to 0} \frac{\tau_0(\varepsilon)}{||\tau_0(\varepsilon)||_1}, \ \Delta \tau_0 \ge \mathbf{0}, \ \sum_{k \in P \setminus P_*(s)} \Delta \tau_{0k} = 1.$$

Notice that  $\tau_0(\varepsilon) \to \mathbf{0}, \ \Delta \tau_*(\varepsilon) \to \mathbf{0}, \ \Delta X(\varepsilon) \to \mathbb{O}_{p \times p}$  as  $\varepsilon \to 0$ . Then

$$\lim_{\varepsilon \to 0} \frac{\tau(\varepsilon)^\top X(\varepsilon)\tau(\varepsilon)}{||\tau_0(\varepsilon)||_1} = \lim_{\varepsilon \to 0} \frac{\tau_*(\varepsilon)^\top X_*(\varepsilon)\tau_*(\varepsilon)}{||\tau_0(\varepsilon)||_1} + 2\tau_*^{*\top} X_{*0}^0 \Delta \tau_0.$$

Since  $X_*(\varepsilon) = X(s,\varepsilon) \in \mathbb{S}_+(p(s))$ , we have  $\lim_{\varepsilon \to 0} \frac{\tau_*(\varepsilon)^\top X_*(\varepsilon)\tau_*(\varepsilon)}{||\tau_0(\varepsilon)||_1} \ge 0$ , and

$$\tau_*^{*\top} X^0_{*0} \Delta \tau_0 = \sum_{k \in P \setminus P_*(s)} \Delta \tau_{0k} e_k^\top X^0 \tau^* \ge \mu(s) \sum_{k \in P \setminus P_*(s)} \Delta \tau_{0k} = \mu(s),$$

where  $\mu(s) := \min\{\nu_{kj}, k \in P \setminus P_*(s), j \in J(s)\} > 0, \nu_{kj}$  are defined in (22). Hence we obtain that  $\tau(\varepsilon)^\top X^0(\varepsilon)\tau(\varepsilon) \ge 0$  for sufficiently small  $\varepsilon \ge 0$ , but this contradicts (39). Thus, we have proved that  $X^0(\varepsilon) \in \mathcal{COP}(p)$  for sufficiently small  $\varepsilon \geq 0.$ 

**Theorem 4.3.** Suppose that for a pair  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p))$  Assumptions j)-jjj) hold true. Then there exists  $\varepsilon_0 > 0$  such that a pair  $(X, U) \in \mathbb{S}(p) \times \mathbb{S}(p)$ ,  $||(X, U) - \mathbb{S}(p) \times \mathbb{S}(p)| = 0$  $|(X^0, U^0)|| \leq \varepsilon_0$ , belongs to the complementarity set  $\mathbb{C}(\mathcal{COP}(p))$  iff there exist matrices  $W(s) \in \mathbb{S}(p(s)), s \in S, such that$ 

$$\mathcal{A}(X,s)W(s) + W(s)\mathcal{A}(X,s) = \mathbb{O}_{p(s) \times p(s)} \ \forall s \in S,$$
(40)

$$U = \sum_{s \in S} \mathcal{B}(W(s), s).$$
(41)

**Proof** follows from Theorems 4.1 and 4.2.

Thus we showed that in a neighborhood of a given point  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p))$ satisfying Assumptions j)-jjj), the complementarity set  $\mathbb{C}(\mathcal{COP}(p))$  is uniquely characterized by system of bi-linear (40) and linear (41) equations. It is natural to call this system as a system of defining equations for cone  $\mathbb{C}(\mathcal{COP}(p))$ .

Note that this system of defining equations itself is uniquely defined by the sets  $P_*(s), s \in S$ , which in turn are uniquely determined by the matrix  $X^0$ .

Above we supposed that  $T_a(X^0) \neq \emptyset$ . Let us analyze the case when  $T_a(X^0) = \emptyset$ , and consequently  $X^0 \in \operatorname{int} \mathcal{COP}(p)$ . It is evident that if  $X \in \operatorname{int} \mathcal{COP}(p)$ , then condition  $(X,U) \in \mathbb{C}(\mathcal{COP}(p))$  implies the equality  $U = \mathbb{O}_{p \times p}$ , and there exists  $\varepsilon_0 > 0$  such that the inequality  $||X - X^0|| \leq \varepsilon_0$  implies the inclusion  $X \in \operatorname{int} \mathcal{COP}(p)$ . Hence, if  $T_a(X^0) = \emptyset$ , then there exists  $\varepsilon_0 > 0$  such that Theorem 4.3 holds true with  $U = U^0 = \mathbb{O}_{p \times p}$  and all  $X \in \mathcal{COP}(p)$  satisfying the inequality  $||X - X^0|| \leq \varepsilon_0$ .

Let us end this section with a small example illustrating Theorem 4.3.

Set 
$$a = (1, -1, 1)^{\top}$$
,  $b = (1, 1, 0)^{\top}$ ,  $c = (0, 1, 1)^{\top}$ , and consider a pair  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(3))$  where  $X^0 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} + a a^{\top} \in \mathcal{COP}(3), \ U^0 = bb^{\top} + cc^{\top} \in \mathcal{CP}(3).$ 

For these matrices, we have  $T_a(X^0) = \{\tau(1) = b/2, \tau(2) = c/2\}, J = \{1, 2\}, S = \{1, 2\}, J(1) = \{1\}, P_*(1) = \{1, 2\} = M(1), J(2) = \{2\}, P_*(2) = \{2, 3\} = M(2), U^0(1) = bb^{\top}, U^0(2) = cc^{\top}$ . One can check that for  $(X^0, U^0)$  all Assumptions j)-jjj) are fulfilled.

Using the sets  $P_*(1)$  and  $P_*(2)$ , let us form the system of defining equations (40). For this example the system has the form

$$\begin{pmatrix} x_{11} & x_{12} \\ x_{12} & x_{22} \end{pmatrix} \begin{pmatrix} w_{11}(1) & w_{12}(1) \\ w_{12}(1) & w_{22}(1) \end{pmatrix} + \begin{pmatrix} w_{11}(1) & w_{12}(1) \\ w_{12}(1) & w_{22}(1) \end{pmatrix} \begin{pmatrix} x_{11} & x_{12} \\ x_{12} & x_{22} \end{pmatrix} = \mathbb{O}_{2 \times 2}, \quad (42)$$

$$\begin{pmatrix} x_{22} & x_{23} \\ x_{23} & x_{33} \end{pmatrix} \begin{pmatrix} w_{22}(2) & w_{23}(2) \\ w_{23}(2) & w_{33}(2) \end{pmatrix} + \begin{pmatrix} w_{22}(2) & w_{23}(2) \\ w_{23}(2) & w_{33}(2) \end{pmatrix} \begin{pmatrix} x_{22} & x_{23} \\ x_{23} & x_{33} \end{pmatrix} = \mathbb{O}_{2 \times 2}. \quad (43)$$

It is a system w.r.t. variables

 $x_{11}, x_{12}, x_{22}, x_{23}, x_{33}, w_{11}(1), w_{12}(1), w_{22}(1), w_{22}(2), w_{23}(2), w_{33}(2).$ 

According to Theorem 4.3, there exists  $\varepsilon_0 > 0$  such that a pair  $(X, U) \in \mathbb{S}(3) \times \mathbb{S}(3)$ ,  $||(X, U) - (X^0, U^0)|| \le \varepsilon_0$ , belongs to the complementarity set  $\mathbb{C}(\mathcal{COP}(3))$  iff these matrices X and U have the forms

$$X = \begin{pmatrix} x_{11}^* & x_{12}^* & x_{13} \\ x_{12}^* & x_{22}^* & x_{23}^* \\ x_{13}^* & x_{23}^* & x_{33}^* \end{pmatrix}, U = \begin{pmatrix} w_{11}^*(1) & w_{12}^*(1) & 0 \\ w_{12}^*(1) & w_{22}^*(1) + w_{22}^*(2) & w_{23}^*(2) \\ 0 & w_{23}^*(2) & w_{33}^*(2) \end{pmatrix}$$

where  $(x_{11}^*, x_{12}^*, x_{22}^*, x_{23}^*, x_{33}^*, w_{11}^*(1), w_{12}^*(1), w_{22}^*(1), w_{23}^*(2), w_{23}^*(2), w_{33}^*(2))$  is a solution to system (42), (43).

## 5. Independence of bi-linear functions forming the system of defining equations

In Section 4, we obtained a system of defining equations (40), (41), which allows one to uniquely characterize the complementarity set  $\mathbb{C}(\mathcal{COP}(p))$  in a neighborhood of a given point  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p))$  satisfying Assumptions j)-jjj). In this system, the main role play equations (40) since equations (41) are linear and needed only for defining matrix U using matrices W(s),  $s \in S$ , satisfying (40). In this section, we show that Jacobian matrix of the system of equations (40), calculated at point  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p))$ , has full row rank.

Given a pair  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p))$  and a presentation of  $U^0$  in the form (7), let us denote  $p_*(s) = p(s)(p(s)) + 1)/2, s \in S, p_* = p(p+1)/2, m = \sum_{s \in S} p_*(s)$ . Using  $X \in \mathbb{S}(p)$  and  $W(s) \in \mathbb{S}(p(s)), s \in S$ , as parameters, let us form vectors and **bi-linear** vector-functions

$$z = \begin{pmatrix} \operatorname{svec}(X) \\ \operatorname{svec}(W(s)) \\ s \in S \end{pmatrix} \in \mathbb{R}^{p_* + m}, \ z^0 = \begin{pmatrix} \operatorname{svec}(X^0) \\ \operatorname{svec}(\mathcal{A}(U^0(s), s)) \\ s \in S \end{pmatrix} \in \mathbb{R}^{p_* + m},$$
(44)

$$\Omega(z,s) := \operatorname{svec}(\mathcal{A}(X,s)W(s) + W(s)\mathcal{A}(X,s)) \in \mathbb{R}^{p_*(s)}, s \in S,$$
$$\Omega(z) := (\Omega^\top(z,s), s \in S)^\top \in \mathbb{R}^m.$$

It is easy to see that system (40) can be rewritten as  $\Omega(z) = 0$ .

**Proposition 5.1.** Suppose that for a pair  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p))$  Assumption jj) and condition iii) from Proposition 3.3 are satisfied. Let  $U^0(s), s \in S$ , be matrices satisfying (7), vector  $z^0$  and function  $\Omega(z)$  be as defined above. Then the matrix  $\frac{\partial \Omega(z^0)}{\partial z} \in \mathbb{R}^{m \times (p_* + m)}$  has full rank:  $\operatorname{rank} \frac{\partial \Omega(z^0)}{\partial z} \in \mathbb{R}^{m \times (p_* + m)} = m$ .

**Proof.** For given  $M, N \in \mathbb{R}^{p \times p}$ , denote by  $M \otimes_s N$  a symmetric Kronecker product of these matrices, see [18] for exact definition and properties. Notice that  $M \otimes_s N$  is  $p_* \times p_*$  matrix,  $M \otimes_s N = N \otimes_s M$ , it is symmetric if M and N are symmetric, and for  $X, U \in \mathbb{S}(p)$  we have

$$(X \otimes_s E(p))\operatorname{svec}(U) = \frac{1}{2}\operatorname{svec}(XU + UX), \tag{45}$$

where E(p) is identity  $p \times p$  matrix. Denote

$$X^{0}(s) := \mathcal{A}(X^{0}, s) \in \mathbb{S}(p(s)), \ W^{0}(s) := \mathcal{A}(U^{0}(s), s) \in \mathbb{S}(p(s)), \ s \in S,$$

$$\mathcal{K}(s) := W^0(s) \otimes_s E(p(s)) \in \mathbb{S}(p_*(s)), \ \mathcal{L}(s) := X^0(s) \otimes_s E(p(s)) \in \mathbb{S}(p_*(s)), \ s \in S.$$

Let  $\mathcal{K}_{ij}(s) \in \mathbb{R}^{p_*(s)}$ ,  $(i, j) \in V(P_*(s))$ , be columns of the matrix  $\mathcal{K}(s)$  for  $s \in S$ , and

let  $S = \{1, ..., s_*\}$ . Then the matrix  $\frac{\partial \Omega(z^0)}{\partial z}$  has the form

$$\frac{\partial \Omega(z^0)}{\partial z} = \begin{pmatrix} \mathcal{Q}(1) & \mathcal{L}(1) & \mathbb{O} & \mathbb{O} & \dots & \mathbb{O} \\ \mathcal{Q}(2) & \mathbb{O} & \mathcal{L}(2) & \mathbb{O} & \dots & \mathbb{O} \\ \dots & \dots & \dots & \dots & \dots \\ \mathcal{Q}(s_*) & \mathbb{O} & \mathbb{O} & \mathbb{O} & \dots & \mathcal{L}(s_*) \end{pmatrix}$$

where for  $s \in S$ ,  $\mathcal{Q}(s)$  is  $p_*(s) \times p_*$  matrix with columns  $\mathcal{Q}_{ij}(s), (i, j) \in V(P)$ :

$$\mathcal{Q}_{ij}(s) = \mathbf{0} \ \forall (i,j) \in V(P) \setminus V(P_*(s)), \ \mathcal{Q}_{ij}(s) = \mathcal{K}_{ij}(s) \ \forall (i,j) \in V(P_*(s)).$$

Suppose that  $\operatorname{rank} \frac{\partial \Omega(z^0)}{\partial z} \in \mathbb{R}^{m \times (p_* + m)} < m$ . Then there exists a set of matrices  $Y(s) \in \mathbb{S}(p(s)), \ s \in S, \text{ such that } \sum_{s \in S} ||Y(s)|| > 0 \text{ and } \left( \frac{\operatorname{svec}(Y(s))}{s \in S} \right)^{\top} \frac{\partial \Omega(z^0)}{\partial z} = \mathbf{0}^{\top}.$ The latter equality involves the second sec The latter equality implies the equalities

$$\sum_{s \in S} (\operatorname{svec}(Y(s)))^\top \mathcal{Q}(s) = \mathbf{0}^\top, \ (\operatorname{svec}(Y(s)))^\top \mathcal{L}(s) = \mathbf{0}^\top \ \forall s \in S.$$

Then it follows from these equalities and (45) that

$$\sum_{s \in S} \mathcal{B}(Z(s), s) = \mathbb{O}_{p \times p}, \ X^0(s)Y(s) + Y(s)X^0(s) = \mathbb{O}_{p(s) \times p(s)} \ \forall s \in S,$$

where  $Z(s) = W^0(s)Y(s) + Y(s)W^0(s)$  for all  $s \in S$ . It follows from statement a) in

Proposition A.2 that  $X^0(s)Z(s) = \mathbb{O}_{p(s)\times p(s)}$   $\forall s \in S$ . It follows from statement a) in By construction, for all  $s \in S$ ,  $X^0(s) \in \mathbb{S}_+(p(s))$ ,  $X^0(s)t(s) = \mathbf{0}$  where  $t(s) = \sum_{j \in J(s)} \tau_*(j) > \mathbf{0}$ ,  $\tau_*(j) = (e_k^\top \tau(j), k \in P_*(s))^\top$ ,  $j \in J(s)$ . Hence it follows from Proposition A.3 that the vectors  $\tau_*(j), j \in J_b(s)$ , form a basis of the subspace  $\{t \in \mathbb{R}^{p(s)} : X^0(s)t = 0\}$ . Then applying Proposition A.4 we obtain that matrix Z(s) can be written in the form

$$Z(s) = \sum_{(i,j)\in V(J_b(s))} \beta_{ij}(s)(\tau_*(i) + \tau_*(j))(\tau_*(i) + \tau_*(j))^\top.$$

This implies that  $\mathcal{B}(Z(s), s) = \sum_{(i,j) \in V(J_b(s))} \beta_{ij}(s)(\tau(i) + \tau(j))(\tau(i) + \tau(j))^{\top}$  and hence

$$\mathbb{O}_{p \times p} = \sum_{s \in S} \mathcal{B}(Z(s), s) = \sum_{s \in S} \sum_{(i,j) \in V(J_b(s))} \beta_{ij}(s) (\tau(i) + \tau(j)) (\tau(i) + \tau(j))^{\top}.$$

It follows from this equality and Assumption jj) that  $\beta_{ij}(s) = 0$  for all  $(i, j) \in V(J_b(s))$ and all  $s \in S$ , and consequently  $Z(s) = \mathbb{O}_{p(s) \times p(s)} \forall s \in S$ . Taking into account these equalities, condition iii) and statement b) in Proposition A.2, we obtain the equalities  $Y(s) = \mathbb{O}_{p(s) \times p(s)} \forall s \in S$ . But this contradicts the condition  $\sum_{s \in S} ||Y(s)|| > 0$ .  $\Box$ 

Let  $\Omega_i(z)$ , i = 1, ..., m, be bi-linear functions forming vector-function  $\Omega(z)$  from the system of defining equations. It follows from proposition proved above that the functions  $\Omega_i(z)$ , i = 1, ..., m, are independent in a neighborhood of a point  $z^0$  constructed on the base of a point  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p))$  by rules (44).

Let us consider example from Section 4. In this example, we have p = 3,  $p_* = 6$ , p(s) = 2,  $p_*(s) = 3$ , for  $s \in S = \{1, 2\}$ , and system (40) has the form (42), (43). It is a system of  $m := p_*(1) + p_*(2) = 6$  bi-linear equations w.r.t. to vector

 $z = (x_{11}, x_{12}, x_{13}, x_{22}, x_{23}, x_{33}, w_{11}(1), w_{12}(1), w_{22}(1), w_{23}(2), w_{33}(2))^{\top} \in \mathbb{R}^{p_* + m}.$ 

One can check that, as it is stated in Proposition 5.1, the bi-linear functions forming this system are independent in a neighborhood of a point  $z^0$  constructed on the base of  $X^0$ ,  $U^0(1)$ ,  $U^0(2)$  by rules (44).

#### 6. Some comments on Assumptions j)-jjj)

In this section, we will show that none of the assumptions j)-jjj) can be omitted in the formulations of the theorems Theorems 4.1, 4.2 and Proposition 5.1.

Let us start with Assumption jj). One can show that under Assumption jj), for a given  $U^0 \in C\mathcal{P}(p)$  such that  $X^0 \bullet U^0 = 0$  there is a unique presentation in the form (7). Note that this assumption does not exclude the situation when for some  $s \in S$ , matrix  $U^0(s)$  from unique presentation in the form (7) may have several different cp factorizations.

If Assumption j) is fulfilled but Assumption jj) is not satisfied, then there exist several sets of matrices, for example,  $\mathbb{U}^0 = \{U^0(s), s \in S\}$  and  $\overline{\mathbb{U}} = \{\overline{U}(s), s \in S\}$  such that  $\mathbb{U}^0 \neq \overline{\mathbb{U}}$  and  $U^0 = \sum_{s \in S} U^0(s) = \sum_{s \in S} \overline{U}(s), \ U^0(s), \ \overline{U}(s) \in \mathcal{F}(s), \ \forall s \in S$ . Notice

that now we can not guarantee the fulfillment of the condition ii) for matrices from both sets  $\mathbb{U}^0$  and  $\overline{\mathbb{U}}$ . But as it will be shown below the fulfillment of this condition is essential.

Also note that if Assumption jj) is violated, then  $\operatorname{rank} \frac{\partial \Omega(z^0)}{\partial z} < m$  and as a result statement of Proposition 5.1 does not hold true.

Violation of Assumption j) in Theorem 4.1. Set p = 5 and for parameter vector  $\theta = (\theta_j, j = 1, ..., 5)$  define a matrix

$$H(\theta) = \begin{pmatrix} 1 & -\cos(\theta_4) & \cos(\theta_4 + \theta_5) & \cos(\theta_2 + \theta_3) & -\cos(\theta_3) \\ -\cos(\theta_4) & 1 & -\cos(\theta_5) & \cos(\theta_1 + \theta_5) & \cos(\theta_4 + \theta_3) \\ \cos(\theta_4 + \theta_5) & -\cos(\theta_5) & 1 & -\cos(\theta_1) & \cos(\theta_1 + \theta_2) \\ \cos(\theta_3 + \theta_2) & \cos(\theta_1 + \theta_5) & -\cos(\theta_1) & 1 & -\cos(\theta_2) \\ -\cos(\theta_3) & \cos(\theta_3 + \theta_4) & \cos(\theta_1 + \theta_2) & -\cos(\theta_2) & 1 \end{pmatrix}$$

and vectors

$$a(\theta) = (\cos(\theta_4 + \theta_5), -\cos(\theta_5), 1, -\cos(\theta_1), \cos(\theta_1 + \theta_2))^{\top}, b(\theta) = (\sin(\theta_4 + \theta_5), -\sin(\theta_5), 0, \sin(\theta_1), -\sin(\theta_1 + \theta_2))^{\top}.$$

Consider a vector  $\theta^* = (\theta_j^*, j = 1, ..., 5)$  such that  $\theta_j^* > 0, j = 1, ..., 5, \sum_{j=1}^5 \theta_j^* = \pi$ . Let us fix this vector and set  $X^0 = H(\theta^*)$ .

It is known (see [23]) that  $H(\hat{\theta^*}) = a(\theta^*)(a(\theta^*))^\top + b(\theta^*)(b(\theta^*))^\top, X^0 \in \mathbb{S}_+(5) \subset \mathbb{S}_+(5)$ 

 $\mathcal{COP}(5),\,T_a(X^0)=\mathrm{conv}\{\tau(j,\theta^*),j=1,...,5\}$  where

$$\tau(1,\theta) = \begin{pmatrix} \sin(\theta_5) \\ \sin(\theta_4 + \theta_5) \\ \sin(\theta_4) \\ 0 \\ 0 \end{pmatrix}, \tau(2,\theta) = \begin{pmatrix} 0 \\ \sin(\theta_1) \\ \sin(\theta_1 + \theta_5) \\ \sin(\theta_5) \\ 0 \end{pmatrix}, \tau(3,\theta) = \begin{pmatrix} 0 \\ 0 \\ \sin(\theta_2) \\ \sin(\theta_1 + \theta_2) \\ \sin(\theta_1) \end{pmatrix},$$

$$\tau(4,\theta) = \begin{pmatrix} \sin(\theta_2) \\ 0 \\ 0 \\ \sin(\theta_3) \\ \sin(\theta_3 + \theta_2) \end{pmatrix}, \tau(5,\theta) = \begin{pmatrix} \sin(\theta_4 + \theta_3) \\ \sin(\theta_3) \\ 0 \\ 0 \\ \sin(\theta_4) \end{pmatrix}$$

Hence  $S = \{1\}$ ,  $P_*(1) = \{1, ..., 5\}$ , and it is evident that Assumptions jj) and jjj) are satisfied.

Let us set  $U^0 = U(\theta^*)$  where  $U(\theta) := \sum_{j=1}^{5} \tau(j,\theta) (\tau(j,\theta))^{\top}$ . By construction  $U(\theta^*) \in U(\theta)$ 

 $\mathcal{CP}(5)$  and  $H(\theta^*) \bullet U(\theta^*) = 0$ , consequently  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(5))$ . One can check that  $U^0$  is not strictly complementary to  $X^0$ , i.e., Assumption j) does not hold true. For  $0 < \varepsilon < \theta_1^*$ , let us set  $\theta(\varepsilon) = (\theta_1^* - \varepsilon, \theta_j^*, j = 2, ..., 5)$  and consider matrices

$$X(\varepsilon) := H(\theta(\varepsilon)), \ U(\varepsilon) := U(\theta(\varepsilon)).$$

It is evident that  $X(\varepsilon) \to X^0$  and  $U(\varepsilon) \to U^0$ . It is known (see [23]) that  $X(\varepsilon) \in COP(p), U(\varepsilon) \in CP(p)$  and  $X(\varepsilon) \bullet U(\varepsilon) = 0$ . It is also known that  $X(\varepsilon) \notin \mathbb{S}_+(5)$ .

Let us show that  $X(\varepsilon)U(\varepsilon) + U(\varepsilon)X(\varepsilon) \neq \mathbb{O}_{5\times 5}$ . Do to this, first let us show that  $X^0 + U^0 \in \operatorname{int} \mathbb{S}_+(5)$ .

Remind that by construction  $X^0 = a(\theta^*)(a(\theta^*))^\top + b(\theta^*)(b(\theta^*))^\top$ . Since  $X^0 \in \mathbb{S}_+(5)$ and  $U^0 \in \mathbb{S}_+(5)$ , hence it is evident that  $X^0 + U^0 \in \mathbb{S}_+(5)$ . Let  $t \in \mathbb{R}^5$  be such that

$$t^{\top} (X^0 + U^0) t = 0 \implies t^{\top} X^0 t = 0, \ t^{\top} U^0 t = 0, \implies t^{\top} a(\theta^*) = 0, \ t^{\top} b(\theta^*), \ t^{\top} \tau(j, \theta^*) = 0, \ j = 1, \dots, 5.$$

It was shown in [23] that this system has only trivial solution t = 0. Thus we have shown that  $X^0 + U^0 \in int \mathbb{S}_+(5)$ . Consequently

$$X(\varepsilon) + U(\varepsilon) \in \operatorname{int} \mathbb{S}_+(5)$$
 for sufficiently small  $\varepsilon > 0$ .

Now let us suppose that for sufficiently small  $\varepsilon > 0$ , the equality  $X(\varepsilon)U(\varepsilon) + U(\varepsilon)X(\varepsilon) = \mathbb{O}_{5\times 5}$  holds true. Then it follows from Proposition A.1 that  $X(\varepsilon) \in \mathbb{S}_+(p)$ , but this contradicts the known condition  $X(\varepsilon) \notin \mathbb{S}_+(p)$ .

Thus we have shown that all conditions of Theorem 4.1, except Assumption j), are fulfilled, but statements of this theorem do not hold true. Hence the assumption is essential in this theorem.

Violation of assumption jjj in Theorem 4.1. Let us set p = 3 and consider

$$X^{0} = \begin{pmatrix} 1 & -1 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = a a^{\top} + b b^{\top}, U^{0} = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} = \tau \tau^{\top},$$
$$a^{\top} = (1, -1, 0), \ b^{\top} = (0, 0, 1), \ \tau^{\top} = (1, 1, 0).$$

It is easy to see that  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p)), T_a(X^0) = \tau = T_a(1, X^0)$ . Hence  $J = \{1\}, \tau(1) = \tau, S = \{1\}, J(1) = J, P_*(1) = \{1, 2\}, M(j) = \{1, 2, 3\}, j \in J, U^0(1) = U^0, X^0(1) = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}, W^0(1) = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ . Assumption j) and j) are fulfiled, Assumption jjj) is not fulfilled.

Set

$$\begin{aligned} a^{\top}(\varepsilon) &= (1, -1, \varepsilon), \ b^{\top}(\varepsilon) = (0, -\varepsilon, 1), \ \tau^{\top}(\varepsilon) = (1 - \varepsilon^2, 1, \varepsilon), \\ X(\varepsilon) &= a(\varepsilon) \ a^{\top}(\varepsilon) + b(\varepsilon) \ b^{\top}(\varepsilon), \ U(\varepsilon) = \tau(\varepsilon) \tau^{\top}(\varepsilon). \end{aligned}$$

Hence

$$X(1,\varepsilon) = \begin{pmatrix} 1 & -1 \\ -1 & 1+\varepsilon^2 \end{pmatrix}, W(1,\varepsilon) = \begin{pmatrix} (1-\varepsilon^2)^2 & (1-\varepsilon^2) \\ (1-\varepsilon^2) & 1 \end{pmatrix}.$$

By construction  $X(\varepsilon) \to X^0$ ,  $U(\varepsilon) \to U^0$  as  $\varepsilon \to 0$ ,  $X(\varepsilon) \bullet U(\varepsilon) = 0$  and  $X(\varepsilon) \in \mathbb{S}_+(3) \subset \mathcal{COP}(3)$ ,  $U(\varepsilon) \in \mathcal{CP}(3)$  for all  $\varepsilon \in [0, 1]$ .

Thus we have shown that all conditions of Theorem 4.1 except Assumption jjj) are satisfied. But the statements of the theorem is not true since

$$X(1,\varepsilon)W(1,\varepsilon) + U(1,\varepsilon)X(1,\varepsilon) \neq \mathbb{O}_{2\times 2}$$
 and  $U_{3q}(\varepsilon) \neq 0 \ \forall q = 1, 2, 3.$ 

Note that in this example, violation of Assumption jjj) implies that  $X^0$  is not strictly complementary to  $U^0$ .

Violation of Assumption j) in Theorem 4.2. Let us set p = 4,

$$X^{0} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, U^{0} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{pmatrix} = (e_{2} + e_{3} + e_{4})(e_{2} + e_{3} + e_{4})^{\top}.$$

It is easy to see that  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p)), T_a(X^0) = \operatorname{conv}\{e_2, e_3, e_4\} = T_a(1, X^0).$ Hence  $J = \{1, 2, 3\}, \tau(1) = e_2, \tau(2) = e_3, \tau(3) = e_4, S = \{1\}, J(1) = J, P_*(1) = \{2, 3, 4\}, M(j) = \{2, 3, 4\}, j \in J, U^0(1) = U^0, X^0(1) = \mathbb{O}_{3\times 3}, W^0(1) = (1, 1, 1)(1, 1, 1)^{\top}.$  Hence we see that Assumptions jj) and jjj) are fulfilled but Assumption j) is not fulfilled. Notice that in this example despite the violation of Assumption j), conditions i) and ii) are satisfied. Consider

,

$$X(\varepsilon) = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 0 & -\varepsilon & \varepsilon \\ 1 & -\varepsilon & 0 & \varepsilon \\ 1 & \varepsilon & \varepsilon & -2\varepsilon \end{pmatrix}, W(1,\varepsilon) = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \implies X(1,\varepsilon) = \begin{pmatrix} 0 & -\varepsilon & \varepsilon \\ -\varepsilon & 0 & \varepsilon \\ \varepsilon & \varepsilon & -2\varepsilon \end{pmatrix}$$

It is easy to see that  $X(\varepsilon) \to X^0$ ,  $W(1,\varepsilon) \to W^0(1)$  as  $\varepsilon \to 0$  and  $X(1,\varepsilon)W(1,\varepsilon) +$  $U(1,\varepsilon)X(1,\varepsilon) = \mathbb{O}_{3\times 3} \ \forall \varepsilon \in \mathbb{R}.$ 

Thus all conditions of Theorem 4.2 except Assumption j) are satisfied. But the statements of the theorem is not true since  $X(\varepsilon) \notin COP(4)$  for  $\varepsilon > 0$ .

Violation of Assumption jjj in Theorem 4.2. Let us p = 3 and consider

$$X^{0} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, U^{0} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} = e_{2}e_{2}^{\top} + e_{3}e_{3}^{\top} + (e_{2} + e_{3})(e_{2} + e_{3})^{\top}.$$

It is easy to see that  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p)), T_a(X^0) = \operatorname{conv}\{e_2, e_3\} = T_a(1, X^0).$ Hence  $J = \{1, 2\}, \tau(1) = e_2, \tau(2) = e_3, S = \{1\}, J(1) = J, P_*(1) = \{2, 3\}, M(j) = \{1, 2, 3\}, j \in J, U^0(1) = U^0, X^0(1) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, W^0(1) = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$ . By construction (see Lemma 3.2),  $U^0$  is strictly complementary to  $X^0$ , hence Assumption j) is fulfilled.

Assumption jj) is also fulfilled as  $S = \{1\}$ . Since  $P_*(1) \neq M(j), j \in J$ , we conclude that Assumption jjj) is not fulfilled.

Consider 
$$X(\varepsilon) = \begin{pmatrix} 1 & -\varepsilon & 0 \\ -\varepsilon & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$
,  $W(1,\varepsilon) = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \implies X(1,\varepsilon) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$ .  
It is easy to see that

is easy to see that

$$X(\varepsilon) \to X^{0}, W(1,\varepsilon) \to W^{0}(1) \text{ as } \varepsilon \to 0;$$

$$X(1,\varepsilon)W(1,\varepsilon) + U(1,\varepsilon)X(1,\varepsilon) = \mathbb{O}_{2\times 2} \,\forall \varepsilon \in \mathbb{R}.$$

$$(46)$$

Thus we have shown that all conditions of Theorem 4.2 except Assumption jjj) are satisfied. But the statements of the theorem is not true since  $X(\varepsilon) \notin COP(3)$  for  $\varepsilon > 0$ .

Note that in this example, violation of Assumption jjj) leads to the fact that  $X^0$  is not strictly complementary to  $U^0$ .

Violation of condition ii) in Theorem 4.2. Notice that Theorem 4.2 is formulated under Assumptions j)-jjj). But in fact we use only Assumption jjj) and conditions i)-iii) which are consequence of Assumptions j) and jj). In an example below, Assumption jij) and conditions i) and iii) are fulfilled, but the only condition ii) is violated. As a result statements of Theorem 4.2 do not hold true.

Let us set p = 3 and consider

$$X^{0} = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, U^{0} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

It is easy to see that  $(X^0, U^0) \in \mathbb{C}(\mathcal{COP}(p)), T_a(X^0) = \operatorname{conv}\{e_2, e_3\} = T_a(1, X^0).$ 

Hence  $J = \{1, 2\}, \tau(1) = e_2, \tau(2) = e_3, S = \{1\}, J(1) = J, P_*(1) = \{2, 3\}, M(j) = \{2, 3\}, j \in J, U^0(1) = U^0, X^0(1) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, W^0(1) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ . Since rank $(X^0(1) + W^0(1)) = 2$ , we have  $(X^0(1) + W^0(1)) \in \operatorname{int} \mathbb{S}_+(2)$ . Thus in this example Assumptions jj), jjj) and conditions i) and iii) are fulfilled but condition ii) is violated.

Let us set

$$X(\varepsilon) = X^0 \implies X^0(1) = X(1,\varepsilon) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, \ W(1,\varepsilon) = \begin{pmatrix} 1 & -\varepsilon \\ -\varepsilon & 1 \end{pmatrix}.$$

Then by construction, relations (46) hold true.

Thus we have shown that all conditions (under which Theorem 4.2 is proved) except condition ii) are satisfied. But the statements of the theorem is not true. In fact, in the example we have  $W(1,\varepsilon) \notin C\mathcal{P}(2)$  for  $\varepsilon > 0$  and as result  $U(\varepsilon) := \sum_{s \in S} \mathcal{B}(W(s,\varepsilon),s) =$ 

 $\mathcal{B}(W(1,\varepsilon),1) \notin \mathcal{CP}(p).$ 

In this section, we have shown that none of the assumptions j)-jjj) can be omitted in the formulations of the theorems Theorems 4.1, 4.2 and Proposition 5.1 and none of the assumptions is a consequence of other assumptions. However, we guess that in Theorems 4.1, 4.2 the assumptions jj), jjj) can be replaced by some less restrictive ones.

## 7. Conclusion

The main motivation for these studies is our desire to explore parametric conic problems over the cones of copositive and completely positive matrices. This led us to examine for these cones the complementarity relations, which play a central role in the study of optimization problem. Based on the analysis of the properties of the elements of the complementarity set under consideration, we obtained a system of bi-linear and linear equations w.r.t. extended set of variables, which allow us to completely describe the complementarity set in a neighborhood of a point from this set under some non-degeneracy assumptions. This result will be the basis for our further research related to the sensitivity of the solution to copositive programming problems.

Also, these results can serve as the basis for further research of the properties of the complementarity set, which will allow us to relax the assumptions made in the paper.

#### References

- Anjos MF, Lasserre JB. Handbook on semidefinite, conic and polynomial optimization. Vol. 166. Springer Science & Business Media; 2011.
- [2] Letchford AN, Parkes AJ. A guide to conic optimisation and its applications. RAIRO-Operations Research. 2018;52(4-5):1087–1106.
- [3] Pataki G. The geometry of semidefinite programming. Handbook of semidefinite programming: Theory, Algorithms, and Applications. 2000;:29–65.
- [4] Ramírez C H, Roshchina V. Refining the partition for multifold conic optimization problems. Optimization. 2020;69(11):2489–2507.
- [5] Mohammad-Nezhad A, Terlaky T. Parametric analysis of semidefinite optimization. Optimization. 2020;69(1):187-216.

- [6] Bonnans JF, Ramírez C H. Perturbation analysis of second-order cone programming problems. Mathematical Programming. 2005;104(2-3):205-227.
- [7] Kostyukova O, Tchemisova T, Dudina O. Immobile indices and cq-free optimality criteria for linear copositive programming problems. Set-Valued and Variational Analysis. 2020; 28:89–107.
- [8] Cottle RW, Pang JS, Stone RE. The linear complementarity problem. SIAM; 2009.
- Gao Y, Németh SZ, Sznajder R. The monotone extended second-order cone and mixed complementarity problems. Journal of Optimization Theory and Applications. 2022; 193(1-3):381–407.
- [10] Ding L, Udell M. A strict complementarity approach to error bound and sensitivity of solution of conic programs. Optimization Letters. 2023;17(7):1551–1574.
- [11] Alizadeh F, Schmieta S. Optimization with semi-definite, quadratic and linear constraints, report 23-97. Rutgers Center for Operations Research, Rutgers University. 1997;.
- [12] Rudolf G, Noyan N, Papp D, et al. Bilinear optimality constraints for the cone of positive polynomials. Mathematical programming. 2011;129(1):5–31.
- [13] Gowda MS, Tao J. On the bilinearity rank of a proper cone and lyapunov-like transformations. Mathematical Programming. 2014;147:155–170.
- [14] Sznajder R. The lyapunov rank of extended second order cones. Journal of Global Optimization. 2016;66:585–593.
- [15] Bomze IM. Copositive optimization-recent developments and applications. European Journal of Operational Research. 2012;216(3):509–520.
- [16] Bomze IM, Schachinger W, Uchida G. Think co (mpletely) positive! matrix properties, examples and a clustered bibliography on copositive optimization. Journal of Global Optimization. 2012;52(3):423–445.
- [17] Berman A, Dur M, Shaked-Monderer N. Open problems in the theory of completely positive and copositive matrices. The Electronic Journal of Linear Algebra. 2015;29:46– 58.
- [18] Alizadeh F, Haeberly JPA, Overton ML. Primal-dual interior-point methods for semidefinite programming: convergence rates, stability and numerical results. SIAM Journal on Optimization. 1998;8(3):746–768.
- [19] Kostyukova O, Tchemisova T. On equivalent representations and properties of faces of the cone of copositive matrices. Optimization. 2022;71(11):3211–3239.
- [20] Kostyukova O, Tchemisova T. Structural properties of faces of the cone of copositive matrices. Mathematics. 2021;9(21):2698.
- [21] Dickinson PJ. Geometry of the copositive and completely positive cones. Journal of Mathematical Analysis and Applications. 2011;380(1):377–395.
- [22] Groetzner P, Dür M. A factorization method for completely positive matrices. Linear Algebra and its Applications. 2020;591:1–24.
- [23] Zhang Q. Completely positive cones: are they facially exposed? Linear Algebra and its Applications. 2018;558:195–204.

#### Appendix A. Proofs of auxiliary statements

**Proposition A.1.** Suppose that  $U \in \mathbb{S}(p)$ ,  $X \in \mathbb{S}(p)$ ,  $UX + XU = \mathbb{O}_{p \times p}$  and  $X + U \in$ int  $\mathbb{S}_+(p)$ . Then  $X \in \mathbb{S}_+(p)$ ,  $U \in \mathbb{S}_+(p)$  and  $UX = \mathbb{O}_{p \times p}$ .

**Proof.** If  $U \in \mathbb{S}(p)$  and  $X \in \mathbb{S}(p)$ , then there exist matrices Q and G such that

$$U = Q\Lambda Q^{\top}, \ X = GMG^{\top}, \ Q^{\top}Q = E(p), \ G^{\top}G = E(p) \text{ where}$$
$$\Lambda = \operatorname{diag}(\lambda_k, k = 1, ..., p), \ M = \operatorname{diag}(\mu_k, k = 1, ..., p),$$

 $\lambda_k, k = 1, ..., p$ , are eigenvalues of matrix  $U, \mu_k, k = 1, ..., p$ , are eigenvalues of matrix X. Then conditions  $UX + XU = \mathbb{O}_{p \times p}$  and  $X + U \in \operatorname{int} \mathbb{S}_+(p)$  take the form

$$\mathbb{O}_{p \times p} = Q \Lambda Q^{\top} G M G^{\top} + G M G^{\top} Q \Lambda Q^{\top}, \ Q \Lambda Q^{\top} + G M G^{\top} \in \operatorname{int} \mathbb{S}_{+}(p)$$

It follows from these relations and condition  $\det Q \neq 0$  that

$$\Lambda F + F\Lambda = \mathbb{O}, \quad \Lambda + F \in \operatorname{int} \mathbb{S}_+(p) \text{ where } F := Q^\top G M G^\top Q.$$

This implies that  $2\lambda_k F_{kk} = 0$ ,  $\lambda_k + F_{kk} > 0 \ \forall k = 1, ..., p$ , wherefrom we obtain that  $\lambda_k \ge 0, F_{kk} \ge 0$  for k = 1, ..., p.

If  $\lambda_k \geq 0$  for all k = 1, ..., p, then it is evident that  $U = Q\Lambda Q^\top \in \mathbb{S}_+(p)$ . Notice that  $F_{kk} = \mu_k ||\omega_k||^2$ , where  $\omega_k = G^\top Qe_k$ . If det  $Q \neq 0$  and det  $G \neq 0$ , then  $||\omega_k||^2 > 0$  for all k = 1, ..., p. These inequalities and relations  $F_{kk} \ge 0, F_{kk} = \mu_k ||\omega_k||^2$ , k = 1, ..., p, imply the inequalities  $\mu_k \geq 0$  for all k = 1, ..., p and, hence,  $X = GMG^{\top} \in I$  $\mathbb{S}_+(p)$ . Thus we have proved that  $U \in \mathbb{S}_+(p)$  and  $X \in \mathbb{S}_+(p)$ .

It follows from the condition  $UX + XU = \mathbb{O}_{p \times p}$  that  $0 = \operatorname{trace}(UX + XU) =$  $2 \operatorname{trace}(UX)$ . It is easy to show that the relations  $U \in S_+(p), X \in S_+(p)$  and trace(UX) = 0 imply the equality  $UX = \mathbb{O}_{p \times p}$ . 

**Proposition A.2.** Let  $X \in \mathbb{S}_+(p)$ ,  $W \in \mathbb{S}_+(p)$ , and  $Y \in \mathbb{S}(p)$  be such that WX = $\mathbb{O}_{p \times p},$ 

$$XY + YX = \mathbb{O}_{p \times p}.\tag{A1}$$

Denote  $Z := YW + WY \in \mathbb{S}(p)$ . Then a)  $XZ = \mathbb{O}_{p \times p}$ , b) the equality  $Z = \mathbb{O}_{p \times p}$ implies the equality  $Y = \mathbb{O}_{p \times p}$  if  $(X + W) \in \operatorname{int} \mathbb{S}_+(p)$ .

**Proof.** It follows from inclusions  $X \in \mathbb{S}_+(p), W \in \mathbb{S}_+(p)$ , and equality  $WX = \mathbb{O}_{p \times p}$ that there exist a matrix  $Q \in \mathbb{R}^{p \times p}$  and numbers  $w_k \geq 0, x_k \geq 0, k = 1, ..., p$ , such that  $Q^{\top}Q = E(p)$  and

$$QXQ^{\top} = \text{diag}(x_k, k = 1, ..., p), \ QWQ^{\top} = \text{diag}(w_k, k = 1, ..., p), \ w_k x_k = 0 \ \forall \ k = 1, ..., p$$

Denote F := XZ, then it follows from equality  $WX = \mathbb{O}_{p \times p}$  that F = XYW. Hence

$$QFQ^{\top} = QXQ^{\top}QYQ^{\top}QWQ = \operatorname{diag}(x_k, k = 1, ..., p)\overline{Y}\operatorname{diag}(w_k, k = 1, ..., p) \quad (A2)$$

where  $\bar{Y} := QYQ^{\top}$ . On the other hand, equality (A1) implies that

$$\mathbb{O}_{p \times p} = Q(XY + YX)Q^{\top} = QXQ^{\top}QYQ^{\top} + QYQ^{\top}QXQ^{\top} \Longrightarrow \\
\mathbb{O}_{p \times p} = \operatorname{diag}(x_k, k = 1, ..., p)\bar{Y} + \bar{Y}\operatorname{diag}(x_k, k = 1, ..., p) \Longrightarrow$$

$$\operatorname{diag}(x_k, k = 1, ..., p)\bar{Y} = -\bar{Y}\operatorname{diag}(x_k, k = 1, ..., p).$$
(A3)

It follows from this equality, and equalities  $w_k x_k = 0 \ \forall k = 1, ..., p$ , and (A2) that

$$QFQ^{\top} = -\bar{Y}\text{diag}(x_k, k = 1, ..., p)\text{diag}(w_k, k = 1, ..., p) = \mathbb{O}_{p \times p}$$

Taking into account that  $\det Q \neq 0$  we conclude that  $F = \mathbb{O}_{p \times p}$ .

Now suppose that  $(X + W) \in \operatorname{int} \mathbb{S}_+(p)$  and equality  $Z = \mathbb{O}_{p \times p}$  holds true. Then the set  $P = \{1, ..., p\}$  can be partition us follows

$$P_x := \{k \in P : x_k > 0\}, \ P_y := \{k \in P : y_k > 0\}, \ P = P_x \cup P_y, \ P_x \cap P_y = \emptyset.$$

The equality  $Z = \mathbb{O}_{p \times p}$  implies the equality  $YW + WY = \mathbb{O}_{p \times p}$  wherefrom we obtain that

$$\mathbb{O}_{p \times p} = \operatorname{diag}(w_k, k = 1, ..., p)\overline{Y} + \overline{Y}\operatorname{diag}(w_k, k = 1, ..., p).$$

It follows from this equality and equality (A3) that

$$\begin{split} \bar{Y}_{kq}x_k + \bar{Y}_{kq}x_q &= 0 \ \forall k \in P_x, \ q \in P_x; \ \bar{Y}_{kq}x_k = 0 \ \forall k \in P_x, \ q \in P_y; \\ \bar{Y}_{kq}y_k + \bar{Y}_{kq}y_q &= 0 \ \forall k \in P_y, \ q \in P_y; \ \bar{Y}_{kq}y_q = 0 \ \forall k \in P_x, \ q \in P_y. \end{split}$$

These equalities imply that  $\bar{Y}_{kq} = 0$  for all  $k \in P$  and  $q \in P$  and hence  $Y = \mathbb{O}_{p \times p}$ .

**Proposition A.3.** Let  $X \in \mathbb{S}_+(p)$ . Suppose that there exists  $\overline{t} \in H(X) := \{t \in \mathbb{R}^p : Xt = 0\}$  such that  $\overline{t} > 0$ . Then  $H(X) = \operatorname{span}\{\tau(j), j \in J\}$ , where  $\tau(j), j \in J$ , are vertices of the polyhedron  $T_*(X) := H(X) \cap \{t \in \mathbb{R}^p : t \ge 0, \sum_{k=1}^p t_k = 1\}$ .

**Proof.** Note that by construction, H(X) is a subspace,  $T_*(X)$  is a convex bounded polyhedron, and  $\tau(j) \ge 0, j \in J$ .

Since  $\tau(j) \in H(X), j \in J$ , we have  $\operatorname{span}\{\tau(j), j \in J\} \subset H(X)$ . Let us show that  $H(X) \subset \operatorname{span}\{\tau(j), j \in J\}$ .

Let  $t(i), i \in I_b$ , be a basis of the subspace H(X). Then

$$\bar{t} = \sum_{i \in I_b} \alpha_i t(i), \ \sum_{i \in I_b} |\alpha_i| > 0.$$

It is evident that for sufficiently large  $\theta$  > we have

$$\bar{t}(i) := t(i) + \theta \bar{t} > 0, \ i \in I_b.$$

Consider an index  $i_0 \in I_b$  such that  $\alpha_{i_0} \neq 0$  and show that the vectors  $\bar{t}(i), \in I_b \setminus i_0, \bar{t}$ , are linearly independent.

Suppose the contrary: there exist numbers  $\beta_i, i \in I_b \setminus i_0$ ,  $\beta_0$  such that  $\sum_{i \in I_b \setminus i_0} \beta_i \overline{t}(i) + \beta_0 \overline{t} = 0$ ,  $\sum_{i \in I_b \setminus i_0} |\beta_i| + |\beta_0| > 0$ . This implies that

$$\sum_{i \in I_b \setminus i_0} \beta_i t(i) + \bar{\beta}_0 \bar{t} = 0 \text{ where } \bar{\beta}_0 = \beta_0 + \theta \sum_{i \in I_b \setminus i_0} \beta_i.$$

The following cases are possible: A)  $\bar{\beta}_0 \neq 0$ , B)  $\bar{\beta}_0 = 0$ ,  $\sum_{i \in I_b \setminus i_0} |\beta_i| \neq 0$ .

Consider case A). Then  $\bar{t} = \sum_{i \in I_b \setminus i_0} \bar{\beta}_i t(i)$  where  $\bar{\beta}_i = -\beta_i/\bar{\beta}_0, i \in I_b \setminus i_0$ . Hence

$$\sum_{i \in I_b} \alpha_i t(i) = \sum_{i \in I_b \setminus i_0} \bar{\beta}_i t(i) \implies \alpha_{i_0} t(i_0) = \sum_{i \in I_b \setminus i_0} (\bar{\beta}_i - \alpha_i) t(i) \text{ where } \alpha_{i_0} \neq 0.$$

But this contradicts the assumption that the vectors t(i),  $i \in I_b$ , are linearly independent.

In case B), we have  $\sum_{i \in I_b \setminus i_0} \beta_i t(i) = 0$ ,  $\sum_{i \in I_b \setminus i_0} |\beta_i| \neq 0$ . But this contradicts the

assumption that the vectors t(i),  $i \in I_b$ , are linearly independent.

Thus we have proved that the vectors  $\overline{t}(i) \in H(X), i \in I_b \setminus i_0$ ,  $\overline{t}$  are linearly independent. Let us set

$$\hat{t}(i) := \bar{t}(i)/||\bar{t}(i)||_1, \ i \in I_b \setminus i_0; \ \hat{t}(i_0) = \bar{t}/||\bar{t}||_1.$$

By construction, the vectors  $\hat{t}(i), i \in I_b$ , form a basis of the subspace H(X) and satisfy the conditions:  $\hat{t}(i) \in T_a(X), i \in I_b$ . Notice that it follows from the latter inclusions that  $\hat{t}(i) = \sum_{j \in J} \alpha_{ij} \tau(j), \sum_{j \in J} \alpha_{ij} = 1, \alpha_{ij} \ge 0, j \in J$ . Then for any  $t \in H(X)$ , we have

$$t = \sum_{i \in I_b} \beta_i \hat{t}(i) = \sum_{i \in I_b} \beta_i \sum_{j \in J} \alpha_{ij} \tau(j) = \sum_{j \in J} \bar{\alpha}_j \tau(j) \text{ where } \bar{\alpha}_j = \sum_{i \in I_b} \beta_i \alpha_{ij}, \ j \in J.$$

This implies that  $t \in \text{span}\{\tau(j), j \in J\}$  and consequently  $H(X) \subset \text{span}\{\tau(j), j \in J\}$ .

**Proposition A.4.** Let  $X \in \mathbb{S}_+(p)$  and  $Z \in \mathbb{S}(p)$  be such that  $XZ = \mathbb{O}_{p \times p}$ . Let  $\tau(j), j \in J_b$ , be a basis of the subspace  $H(X) := \{t \in \mathbb{R}^m : Xt = 0\}$ . Then the matrix Z can be presented in the form

$$Z = \sum_{(i,j)\in V(J_b)} \beta_{ij}(\tau(i) + \tau(j))(\tau(i) + \tau(j))^{\top}, \ \beta_{ij} \in \mathbb{R} \ \forall (i,j) \in V(J_b).$$
(A4)

**Proof.** If  $Z \in \mathbb{S}(p)$ , then there exists an orthogonal matrix  $Q \in \mathbb{R}^{p \times p}$  with columns q(k), k = 1, ..., p, such that

$$Z = Q \text{diag}(z_k, k = 1, ..., p) Q^{\top} = \sum_{k=1}^{p} z_k q(k) q^{\top}(k)$$
(A5)

where  $z_k, k = 1, ..., p$ , are the eigenvalues of Z. Consequently

$$\mathbb{O}_{p \times p} = XZ = XQ \operatorname{diag}(z_k, k = 1, ..., p)Q^\top \implies \mathbb{O}_{p \times p} = XQ \operatorname{diag}(z_k, k = 1, ..., p)$$
$$\implies Xq(k) = 0 \ \forall k \in I := \{k \in \{1, ..., p\} : z_k \neq 0\}.$$

Thus, for  $k \in I$ , we obtain  $q(k) \in H(X)$  and consequently  $q(k) = \sum_{j \in J_b} \alpha_j(k) \tau(j)$ .

This implies that, for  $k \in I$ , we have

$$q(k)q^{\top}(k) = \left(\sum_{j \in J_b} \alpha_j(k)\tau(j)\right)\left(\sum_{i \in J_b} \alpha_i(k)\tau(j)\right)^{\top}$$
$$= \sum_{j \in J_b} \alpha_j^2(k)\tau(j)(\tau(j))^{\top} + \sum_{(i,j) \in V_*(J_b)} \alpha_i(k)\alpha_j(k)(\tau(j)(\tau(i))^{\top} + \tau(i)(\tau(j))^{\top})$$

where  $V_*(J_b) = V(J_b) \setminus \{(j, j), j \in J_b\}$ . Taking into account that

$$\tau(j)(\tau(i))^{\top} + \tau(i)(\tau(j))^{\top} = (\tau(i) + \tau(j))(\tau(i) + \tau(j))^{\top} - \tau(j)(\tau(j))^{\top} - \tau(i)(\tau(i))^{\top}$$

we obtain that  $q(k)q^{\top}(k) = \sum_{(i,j)\in V(J_b)} \bar{\beta}_{ij}(k)(\tau(i) + \tau(j))(\tau(i) + \tau(j))^{\top}$  with some  $\bar{\beta}_{ij}(k) \in \mathbb{R}, \ (i,j) \in V(J_b)$ . It follows from the latter equality and (A5) that presentation (A4) holds true with  $\beta_{ij} = \sum_{k\in I} z_k \bar{\beta}_{ij}(k), \ (i,j) \in V(J_b)$ .