

# [O I] fine structure line profiles in Mon R2 and M17 SW: the puzzling nature of cold foreground material identified by [<sup>12</sup>C II] self-absorption

C. Guevara<sup>1</sup>, J. Stutzki<sup>1</sup>, V. Ossenkopf-Okada<sup>1</sup>, U. Graf<sup>1</sup>, Y. Okada<sup>1</sup>, N. Schneider<sup>1</sup>, P.F. Goldsmith<sup>2</sup>, J.P. Pérez-Beaupuits<sup>3,5,6</sup>, S. Kabanovic<sup>1</sup>, M. Mertens<sup>1</sup>, N. Rothbart<sup>4</sup>, and R. Güsten<sup>5</sup>

<sup>1</sup> I. Physikalisches Institut, Universität zu Köln, Zùlpicher Str. 77, 50937 Köln, Germany  
e-mail: guevara@ph1.uni-koeln.de

<sup>2</sup> Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 91109-8099, USA

<sup>3</sup> European Southern Observatory, Alonso de Córdova 3107, Vitacura, Santiago, Chile

<sup>4</sup> Institute of Optical Sensor Systems, German Aerospace Center (DLR), Rutherfordstr. 2, 12489 Berlin, Germany

<sup>5</sup> Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany

<sup>6</sup> Centro de Astro-Ingeniería UC, Instituto de Astrofísica, Pontificia Universidad Católica de Chile, Avda Vicuña Mackenna 4860, Macul, Santiago, Chile

Received date / Accepted date

## ABSTRACT

**Context.** Recent studies of the optical depth comparing [<sup>12</sup>C II] and [<sup>13</sup>C II] line profiles in Galactic star-forming regions revealed strong self-absorption in [<sup>12</sup>C II] by low excitation foreground material, implying a large column density of C<sup>+</sup> corresponding to an equivalent A<sub>V</sub> of a few, up to about 10 mag.

**Aims.** As the nature and origin of such a large column of cold C<sup>+</sup> foreground gas are difficult to explain, it is essential to constrain the physical conditions of this material.

**Methods.** We conducted high-resolution observations of [O I] 63 μm and [O I] 145 μm lines in M17 SW and Mon R2. The [O I] 145 μm transition traces warm PDR-material, while the [O I] 63 μm line traces foreground material as manifested by absorption dips.

**Results.** Comparison of both [O I] line profiles with [C II] isotopic lines confirm warm PDR-origin background emission and a significant column of cold foreground material causing self-absorption visible in [<sup>12</sup>C II] and [O I] 63 μm profiles. In M17 SW, the C<sup>+</sup> and O<sup>0</sup> column densities are comparable for both layers. Mon R2 exhibits larger O<sup>0</sup> columns compared to C<sup>+</sup>, indicating additional material where the carbon is neutral or in molecular form. Small scale spatial variation of the foreground absorption profiles and the large column density (~ 10<sup>18</sup> cm<sup>-2</sup>) of the foreground material suggest emission from high-density regions associated with the cloud complex, not a uniform diffuse foreground cloud.

**Conclusions.** The analysis confirms that the previously detected intense [C II] foreground absorption is attributable to a large column of low excitation dense atomic material, where carbon is ionized, and oxygen is in neutral atomic form.

**Key words.** ISM:clouds – ISM:individual objects: M17 – photon-dominated region (PDR) – ISM:individual objects: MonR2

## 1. Introduction

The fine structure emission lines of [O I], together with the [C II] 158 μm emission line and high-J CO lines, are the main cooling lines for photodissociation regions (PDRs, [Tielens & Hollenbach 1985](#); [Hollenbach & Tielens 1999](#)) in the interstellar medium (ISM). The spin-orbit coupling in neutral atomic oxygen leads to three fine-structure levels and, correspondingly, to two [O I] fine structure transitions. The lower transition, [O I] <sup>3</sup>P<sub>1</sub> → <sup>3</sup>P<sub>2</sub>, has a wavelength of 63.2 μm, corresponding to an energy of its upper level of 227.7 K; the collisional rate coefficients with H give a critical density at 77 K of 7.8×10<sup>5</sup> cm<sup>-3</sup>, and with H<sub>2</sub> as collision partner the critical density is 5×10<sup>5</sup> cm<sup>-3</sup> ([Goldsmith 2019](#)). The upper transition, [O I] <sup>3</sup>P<sub>0</sub> → <sup>3</sup>P<sub>1</sub>, has a wavelength of 145.5 μm. Its upper state energy is 326.6 K above the ground state, or 98.9 K above the mid-level, with a critical density of 5.8×10<sup>6</sup> cm<sup>-3</sup> for H<sub>2</sub> as collision partner ([Goldsmith 2019](#)). The [O I] fine structure emission from PDRs is thus bright if the gas is dense. It is, therefore, together with the [C II] 158 μm fine structure line, commonly used for tracing the warm and dense gas in

star-forming regions, locally and out to high redshift galaxies. The [O I] 63 μm transition rapidly reaches a high optical depth in low-temperature gas (T ≪ 230 K), so high that the intensity is no longer a measure of the column density. The fine structure lines of the oxygen isotopes, in particular <sup>18</sup>O, are so close in frequency that they blend with the Doppler-broadened main isotope transition in astronomical sources, so that the isotopic line ratios cannot be used to determine the [O I] optical depth. Early indications of high optical depth have been reported by [Stacey et al. \(1983\)](#) and [Boreiko & Betz \(1996\)](#) through comparison of the integrated line intensity ratio of the [O I] lines, but the optical depth could be only inferred indirectly. High signal-to-noise line profiles are required to measure the fine structure line optical depth directly, showing saturation or self-absorption. With the high spectral resolution and high signal-to-noise available with the GREAT ([Heyminck et al. 2012](#)) instrument on SOFIA (Stratospheric Observatory for Infrared Astronomy, [Young et al. 2012](#); [Temi et al. 2018](#)), several authors [Leurini et al. \(2015\)](#); [Schneider et al. \(2018\)](#); [Mookerjea et al. \(2019\)](#); [Kirsanova et al.](#)

(2020); Goldsmith et al. (2021) have recently reported velocity resolved [O I] 63  $\mu\text{m}$  observations showing complex line profiles. A recent survey of [O I] 63  $\mu\text{m}$  line profiles towards 12 star-forming cores in the Milky Way by Goldsmith et al. (2021) showed self-absorption by foreground material for half of the sources. In addition, high optical depth and self-absorption in [O I] are often invoked to explain the observed too-low [O I] 63  $\mu\text{m}$  intensities in comparison to model predictions from other observed lines.

In a recent study of several Galactic star-forming regions comparing the [ $^{12}\text{C II}$ ] and [ $^{13}\text{C II}$ ] line profiles at high signal-to-noise ratio (Guevara et al. 2020), we have shown that the [C II] 158  $\mu\text{m}$  emission is optically thick in a wide range of physical conditions. The two sources, Monoceros R2 (Mon R2) and M17 SW, were studied and show deep and narrow self-absorption in the line profiles against the bright and broad background line emission. The analysis of those spectra showed that the bright background line emission, derived from the optically thin [ $^{13}\text{C II}$ ] line profile and compatible with originating in PDRs, is absorbed by a large column density of ionized carbon in cold foreground material, with a low excitation temperature ( $T_{\text{ex}} \lesssim 25$  K). A lower limit for the foreground material, derived under the assumption that all carbon is in the form of  $\text{C}^+$ , gives a corresponding visual extinction of several, up to ten, magnitudes. The amount of material can be considerably larger if the carbon is only partially ionized and the foreground gas contains more carbon in the form of neutral atoms or bound in molecules, particularly CO. However, there is no correlation between the self-absorption features in the ionized carbon line profiles and low-J CO line profiles. To present knowledge, ionized carbon is present mainly in PDR layers at temperatures above 60. Hence, the nature of such large amounts of ionized carbon in low excitation layers of gas is very puzzling. Low excitation may imply low density (well below the critical density of [C II]), but such diffuse gas would need to fill large volumes to explain the observed total column density. Kabanovic et al. (2022) have presented a scenario for the source RCW 120, where the carbon is present in a diffuse envelope with neutral hydrogen.

Mon R2 is a star-forming region at a distance of 778 pc (Zucker et al. 2019). The region contains a reflection nebula, and the UCH II region is surrounded by several PDRs with different physical conditions. The molecular cloud associated with Mon R2 has a hub-filament structure and contains clumps of density up to  $10^6$   $\text{cm}^{-3}$ . The radiation field in the interface region between the H II region and the cloud is about  $10^5$   $G_0$  (Pilleri et al. 2012). The sources have been studied through several atomic and molecular tracers (e.g. Pilleri et al. 2012, 2013; Treviño-Morales et al. 2014; Rayner et al. 2017; Treviño-Morales et al. 2019).

M17 is one of the brightest and most massive star-forming regions in the Galaxy, located at a distance of 1.9 kpc (Wu et al. 2019). M17 SW is the sub-region located in the southwest, where an H II region is localized and is associated with a giant molecular cloud and PDR interface. M17 SW is considered a prototype of an edge-on PDR. The H II region is ionized by a highly obscured (with a visual extinction  $A_V > 10$  mag) cluster of many ( $>100$ ) OB stars (Hoffmeister et al. 2008). The gas near the H II region is distributed in high density clumps ( $n \leq 10^5$   $\text{cm}^{-3}$ ) embedded in interclump material ( $n \sim 10^3$   $\text{cm}^{-3}$ ) surrounded by diffuse gas ( $n \sim 300$   $\text{cm}^{-3}$ ), irradiated by a strong UV field of  $G_0 = 5.6 \times 10^4$  (Meixner et al. 1992). Recently, Klein et al. (2023) using non-velocity resolved observations of several infrared lines such as [C II], both [O I] lines, [O III] or [N III] from FIFI-LS (Colditz et al. 2018; Fischer et al. 2018) derived hydrogen nuclei

density and UV radiation maps, with an average hydrogen density of  $10^{5.9}$   $\text{cm}^{-3}$  for the molecular cloud. They found that the ionization and photodissociation fronts are nearly merged with a sharp density jump from the ionized region to the neutral one.

To characterize the nature of the absorbing layer in these two sources, Mon R2 and M17 SW, we followed up on the investigation we started in Guevara et al. (2020) by observing the two sources in both [O I] fine structure lines with the upGREAT<sup>1</sup> instrument (Risacher et al. 2016) on board SOFIA along the same lines-of-sight as previously observed in [C II].

The paper is organized as follows: Section 2 details the observational setup and data reduction. Section 3 shows the observed [O I] 145  $\mu\text{m}$  and [O I] 63  $\mu\text{m}$  line profiles and a comparison against [ $^{12}\text{C II}$ ] and [ $^{13}\text{C II}$ ] profiles. Section 4 presents a multi-component double-layered Gaussian profile analysis using both [O I] lines to estimate oxygen column densities and excitation temperature. Section 5 discusses the results and the possible nature of the background and foreground layers. Finally, Section 6 summarizes the present work.

## 2. Observations and data reduction

We observed the [O I] 63  $\mu\text{m}$  and [O I] 145  $\mu\text{m}$  fine-structure lines in Mon R2 and M17 SW with the 7-pixel High-Frequency Array (HFA) and the 7x2-pixel Low-Frequency Array (LFA) arrays of the upGREAT receiver on board SOFIA. Mon R2 was observed in December 2018 and M17 SW in June 2019 and April 2022. For Mon R2, the LFA was tuned to observe the [O I] 145  $\mu\text{m}$  in the H-polarization array (LFAH) and [C II] 158  $\mu\text{m}$  line in the V-polarization (LFAV) simultaneously. The single-sideband (SSB) system temperatures at the source velocity ( $T_{\text{sys}}$ ) were 3500 and 3050 K, respectively. In parallel, the HFA was tuned to [O I] 63  $\mu\text{m}$ , with a  $T_{\text{sys}}$ (SSB) of 4700 K. The parameters are summarized in Table 1. As the former [C II] spectra (Guevara et al. 2020) were observed with the old single-pixel configuration of the GREAT instrument at two discrete positions, we had no information on the spatial variation of the [C II] line profiles. Hence, we first observed a small, fully sampled map (for the three lines) of  $180'' \times 180''$  extent in on-the-fly mode with a grid separation of  $3''$  in the horizontal and vertical directions. Then, we performed deep integrations in total power mode of 5 min ON-source time for the two positions previously observed in [C II] (Fig. 1). Appendix A lists the coordinates of each position in detail and the OFF positions used. The two [C II] positions were selected to follow the peak emission. The average precipitable water vapor column was 7  $\mu\text{m}$ . The OFF position presented some weak contamination of  $\sim 2$  K, corrected through the same procedure for [C II] as described by Guevara et al. (2020), Appendix B, fitting a Gaussian profile to the OFF position spectra and then adding the fitted profile into the ON spectra to recover the lost emission.

In June 2019, we observed a fully sampled small [O I] map for M17 SW using upGREAT in OTF mode with a grid separation of  $3''$  in horizontal and vertical directions. The LFA was tuned to simultaneously observe the [O I] 145  $\mu\text{m}$  in LFAH and [C II] 158  $\mu\text{m}$  in LFAV, with a  $T_{\text{sys}}$ (SSB) at the velocity of the source of 3000 and 2000 K, respectively. The map has an extent of  $220'' \times 220''$ . The average precipitable water vapor was 8  $\mu\text{m}$ . The HFA was tuned to [O I] 63  $\mu\text{m}$ , but the telluric line was located in the central velocities of the emission profile, rendering

<sup>1</sup> upGREAT is a development by the MPI für Radioastronomie and KOSMA/Universität zu Köln, in cooperation with the DLR Institut für Optische Sensorsysteme.

Table 1: Observational parameters

Sources	RA (h:m:s)	DEC (°:':")	LFAH	T <sub>sys</sub> (K)	LFAV	T <sub>sys</sub> (K)	HFA	T <sub>sys</sub> (K)	<pwv> <sup>a</sup> (μm)
Mon R2	06:07:46.2	-06:23:08.0	[O I] 145 μm	3700	[C II] 158 μm	3150	[O I] 63 μm	4800	7
M17 SW <sub>June2019</sub>	18:20:27.6	-16:12:00.9	[O I] 145 μm	3000	[C II] 158 μm	2100	-	-	8
M17 SW <sub>April2022</sub>	18:20:27.6	-16:12:00.9	-	-	-	-	[O I] 63 μm	5200	6.5

<sup>a</sup> Average over all pixels of the precipitable water vapor column.




Fig. 1: Mon R2 [C II] integrated intensity map for the velocity range between 0 and 30 km s<sup>-1</sup> with the two [C II] single pointings (in black) from Guevara et al. (2020). [O I] 63 μm integrated intensity map in blue contours covering the same velocity range of [C II] (levels at 100, 150, 200, 240, 270 and 300 K km s<sup>-1</sup>). The white circle represents the FWHM beam size of both maps, 15''.

the observations useless. Hence, we repeated the same observations in April 2022 at a different time to avoid the telluric line contamination for the [O I] 63 μm emission. The observations were successful, but this time [O I] 145 μm was the line affected by the telluric feature in the central velocities. The average precipitable water vapor was 6.5 μm with a T<sub>sys</sub>(SSB) at 63 μm of 5100 K. The [C II] integrated intensity map shown in Fig. 2 is the combination of the observations referred above and a map observed within the SOFIA Feedback Legacy Project<sup>2</sup> (Schneider et al. 2020) (The full [C II] map will be discussed in a separate publication<sup>3</sup>). The positions for the analysis below were selected following the [C II] peak emission along the ridge. The seven positions were previously observed in [<sup>12</sup>C II] and [<sup>13</sup>C II], already analyzed in Guevara et al. (2020). The OFF position was apparently free of emission.

The data were calibrated to an intensity scale in main beam brightness temperature, T<sub>mb</sub>, with the *kalibrate* task (Guan et al. 2012), part of the standard GREAT pipeline. Then, we subtracted

<sup>2</sup> <https://feedback.astro.umd.edu/>

<sup>3</sup> For details on the observing scheme, see Schneider et al. (2020).




Fig. 2: M17 SW [C II] integrated intensity map for the velocity range between 0 and 40 km s<sup>-1</sup> with the original [C II] single pointings (in black) from Guevara et al. (2020) (equivalent to a LFA array pointing). [O I] 63 μm integrated intensity map in blue contours covering the same velocity range of [C II] (levels at 40, 70, 100, 120, 140 and 160 and 200 K km s<sup>-1</sup>). The black circle shows the FWHM beam size for both maps, 15''.

baselines with the CLASS 90 package, part of the GILDAS<sup>4</sup> software, and resampled the data to 0.3 km s<sup>-1</sup> channel width, the same as used for the [C II] data.

For Mon R2, the atmospheric 63 μm atomic oxygen line at the time of the observation was located at -10 km s<sup>-1</sup> LSR-velocity, sufficiently far away from the emission of the source, and hence not affecting the analysis of the line profile. For inter-comparison, we have convolved the three maps to a 15'' beam size from the nominal 6.3'' for the [O I] 63 μm and 14.1'' for the [O I] 145 μm and [C II] lines. For M17 SW, the maps observed at different times of the year allowed us to avoid the telluric line. Hence, both OI maps were convolved to a 15'' beam size on a 5'' grid with a velocity resolution of 0.3 km s<sup>-1</sup>. All the maps were gridded and convolved into a joint resolution to compare them in identical conditions, avoiding the difference in intensities given by the dissimilar beam sizes.

<sup>4</sup> <https://www.iram.fr/IRAMFR/GILDAS/>




Fig. 3: Mon R2 [O I] 63  $\mu\text{m}$ , [O I] 145  $\mu\text{m}$ , [ $^{12}\text{C II}$ ] and [ $^{13}\text{C II}$ ] spectra for the single pointings previously observed in [C II]. All spectra have been convolved to a beam size of 15". The vertical dashed lines at the 8 and 11 km s<sup>-1</sup> absorption dip are present in both spectra, below the horizontal dashed line at 0 K. The [ $^{13}\text{C II}$ ] has been smoothed in velocity for a matter of display.

### 3. [O I] Line profiles

The [C II] and [O I] emission in Mon R2 is compact (see Fig. 1), extending over  $\approx 60''$  in all three lines, in contrast to M17 SW, where the emission in both lines is extended along the PDR edge (see Fig. 2). Fig. 3 compares the line profiles of the newly observed [O I] spectra in Mon R2 obtained in deep integration and the previously observed [ $^{12}\text{C II}$ ] and [ $^{13}\text{C II}$ ] spectra, the [ $^{13}\text{C II}$ ] profiles shown are the average of the two outer hfs-satellites, as explained in Guevara et al. (2020). The [O I] 145  $\mu\text{m}$  profiles are very similar to those of the [ $^{13}\text{C II}$ ] line, though the latter, being very weak, have lower signal-to-noise. The line profiles agree in peak position and width but also show similarities in the detailed profile structure, which is composed of two Gaussian components overlapping in velocity and added together, as expected for optically thin emission. In contrast, the [O I] 63  $\mu\text{m}$  profile shows deep absorption notches at 8 and 12 km s<sup>-1</sup> velocity and possibly a number of additional weaker ones. The overlay with the [ $^{12}\text{C II}$ ] profiles shows that the [O I] absorption matches in center velocity and line width with the self-absorption notches visible in the [ $^{12}\text{C II}$ ] line.

The match in the absorption notches between [ $^{12}\text{C II}$ ] and [O I] 63  $\mu\text{m}$  is present not only in the two positions mentioned above. As shown in Fig. 4, the self-absorption notches are strong across the source and show spatial variation over distances as close as 10" to 20", corresponding to 0.6 to 0.12 pc at the distance of Mon R2, (see Sect. 5.4 for a discussion of this variation).

The absorption to negative intensities (after baseline subtraction) at 11 km s<sup>-1</sup> can be explained by a weak 63  $\mu\text{m}$  dust continuum. Additional diffuse gas is present in the line of sight. Pilleri et al. (2013) found foreground absorption through observations of small hydrocarbons at the referenced velocity, with an equivalent  $A_V$  of 1 magnitude, detached from the source. The absorption by this component is insufficient to explain the absorption dips but may weakly contribute to the absorption at this velocity (see Sect. 3 for an analysis of the absorption dips). The self-absorption notches strongly reduce the [O I] 63  $\mu\text{m}$  integrated intensity (Fig. 5). The [O I] 63  $\mu\text{m}$  emission is almost completely absorbed out between 10 and 12 km s<sup>-1</sup>. Consequently, the [O I]

63  $\mu\text{m}$  line shows a more fragmented structure in the channel maps.

The comparison of the line profiles shows a similar behavior for M17 SW (Fig. 6): the [O I] 145  $\mu\text{m}$  line profile is very similar to the formerly observed [ $^{13}\text{C II}$ ] profiles (where, in this case, the similarity can be confirmed in detail, because of the high S/N of the deep [ $^{13}\text{C II}$ ] integrations). The [O I] 63  $\mu\text{m}$  profiles are heavily affected by self-absorption, and the absorption features are well correlated in position and width with those identified in [ $^{12}\text{C II}$ ] (see the next section for more details). Only the lower velocity component at around 11 km s<sup>-1</sup> in the [ $^{12}\text{C II}$ ] line is not reproduced in [O I] 63  $\mu\text{m}$ . As discussed in Guevara et al. (2020), this component is associated with the [N II] emission and also visible in H I, thus presumably associated with diffuse and ionized gas. It is also not traced by the low-J CO lines (Pérez-Beaupuits et al. 2015a,b).

To check for positional variations of the [O I] self-absorption profiles at small angular scales, we plot a mosaic of the observed line profiles on a grid with 10" separation, smoothed to the common spatial resolution of 15". Figure 7 shows the spectra in [O I] 63  $\mu\text{m}$ , [O I] 145  $\mu\text{m}$  and [C II]. The [C II] spectra show spatial variations in their profiles from position to position, namely across 10", resp. 0.10 pc, both in the absorption depth and the velocity of the absorption dips. The variations are similarly present in the [O I] 63  $\mu\text{m}$  spectra, although the deep absorption down to close to the zero intensity level makes the variations less pronounced in intensity. The velocities of the absorption dips are well correlated between the [O I] 63  $\mu\text{m}$  and the [C II] emission (same as above, see Sect. 5.4). In contrast, the [O I] 145  $\mu\text{m}$  shows a relatively simple profile of smoothly superimposed emission components. Fig. 8 shows the channel maps for both [O I] lines in M17 SW. Like in Mon R2, the [O I] 63  $\mu\text{m}$  seems to disappear in some areas at the velocity of the absorption dips, in particular at 22 and 24 km s<sup>-1</sup>. We find a third absorption component located south of M17 SW at 16 km s<sup>-1</sup>. Future papers will study the whole maps, for the moment we will focus on the positions already observed in [C II].

Table 2 compares the peak and integrated intensities between both [O I] line transitions for both sources. In general, the [O I] 145  $\mu\text{m}$  peak intensity overshoots the [O I] 63  $\mu\text{m}$  peak intensity by a factor of a few due to the self-absorption effect, except for M17 SW position 4. The latter shows a comparable peak intensity but at different velocities, again a result of self-absorption. The difference is more evident in the integrated intensity, with relatively larger [O I] 145  $\mu\text{m}$  integrated intensities, giving a 145/63 integrated intensity ratio between 1.2 and 7. The self-absorption effects thus significantly enhance the integrated intensity ratio. This effect must be considered when analyzing data, particularly non-resolved data without velocity information, such as high-redshift extra-galactic observations. Although this self-absorption effect has already been observed under different environments of Galactic molecular clouds (such as Leurini et al. 2015; Gusdorf et al. 2017; Kristensen et al. 2017; Schneider et al. 2018, and references there in), it has also been observed in the large scale [O I] 63  $\mu\text{m}$  spectra of extra-galactic sources like Arp 220, NGC 4945, NGC 4418, and even in the [O III] 88  $\mu\text{m}$  of IRAS17208-0014 (for example González-Alfonso et al. 2012; Fernández-Ontiveros et al. 2016).

### 4. Gaussian Multi-component Analysis

For a quantitative comparison, we utilize in a two-layer radiative transfer model (Gaussian multi-component fit to the [O I] profiles) similar to what was introduced in our previous [ $^{12}\text{C II}$ ] and




Fig. 4: Mon R2 [O I] 63  $\mu\text{m}$ , [O I] 145  $\mu\text{m}$  and [C II] mosaic spectra for the central area of the source. The dotted lines are located at 5, 10, 15 and 20  $\text{km s}^{-1}$  respectively. The angular resolution is the same for [C II], [O I] 145  $\mu\text{m}$  and [O I] 63  $\mu\text{m}$  (15''), with a grid spacing of 10''. The offset coordinates of each spectrum with respect to the source position are given in the top left corner of each box.




Fig. 5: Mon R2 channel maps of the velocity-binned emission of [O I] 145  $\mu\text{m}$  in contours (with levels at 15, 24, 33, 42, 51, 60 and 70  $\text{K km s}^{-1}$ ) and [O I] 63  $\mu\text{m}$  in color. The angular resolution for both lines is 15'' with a grid separation of 5''. The velocity bin for the channels is 1  $\text{km s}^{-1}$ , therefore we can express the integrated intensity units of  $\text{K km s}^{-1}$  as purely K.

Table 2: Mon R2 and M17 SW [O I] 145  $\mu\text{m}$  and [O I] 63  $\mu\text{m}$  integrated and peak intensity.

Positions	Peak Intensity			Integrated Intensity		
	[O I] <sub>145</sub> (K)	[O I] <sub>63</sub> (K)	$\frac{[\text{O I}]_{145}}{[\text{O I}]_{63}}$	[O I] <sub>145</sub> (K km s <sup>-1</sup> )	[O I] <sub>63</sub> (K km s <sup>-1</sup> )	$\frac{[\text{O I}]_{145}}{[\text{O I}]_{63}}$
MonR2 1	65	12	5.4	394	111	3.5
MonR2 2	85	53	1.6	495	228	2.2
M17SW 0	60	23	2.6	260	119	2.2
M17SW 1	41	6	6.8	197	30	6.6
M17SW 2	42	4	10.5	233	33	7.1
M17SW 3	26	18	1.4	199	77	2.6
M17SW 4	71	71	1.0	421	363	1.2
M17SW 5	11	7	1.6	74	46	1.7
M17SW 6	66	27	2.4	339	129	2.6




Fig. 6: Same as Fig. 3 but for M17 SW. The dotted lines mark the two absorption dips present for both fine structure lines at 21 and 24 km s<sup>-1</sup>.

[<sup>13</sup>C II] study (Guevara et al. 2020). The formal solution of the radiative transfer equation for a two-layer model with  $b = 1 \dots B$  number of background components and  $f = 1 \dots F$  number of foreground components gives for the brightness temperature of the line transition  $t$  ( $t = 145, 63$  denoting the [O I] 145  $\mu\text{m}$  and the [O I] 63  $\mu\text{m}$  transition respectively):

$$T_{\text{mb},t}(v) = \left[ \mathcal{J}_v(T_{\text{ex},t}) \left( 1 - \exp \left( - \sum_b \tau_{b,t}(v) \right) \right) \right] \exp \left( - \sum_f \tau_{f,t}(v) \right) + \mathcal{J}_v(T_{\text{ex},f}) \left( 1 - \exp \left( - \sum_f \tau_{f,t}(v) \right) \right) \quad (1)$$

Here,  $\mathcal{J}_v(T) = \frac{h\nu}{k} \left( e^{h\nu/k_b T} - 1 \right)^{-1}$  is the expression for the equivalent brightness temperature of a blackbody emission at temperature  $T$ . The optical depth of each transition  $t$  and component  $i = b, f$  is given by:

$$\tau_{i,t}(v) = \frac{c^3}{8\pi\nu_t^3} \phi_i(v) N_{u_i} A_t \left( \exp \left( h\nu_t / (k_b T_{\text{ex},t}) \right) - 1 \right) \quad (2)$$

with the upper and lower state of transition  $t$  given by  $u_t$  and  $l_t$  respectively, its upper state column density by  $N_{u_i}$ , and its Einstein-A-coefficients  $A_t = A_{u_i, l_t}$ . The numerical values of the latter are  $A_{145} = 1.75 \times 10^{-5}$  and  $A_{63} = 8.91 \times 10^{-5} \text{ s}^{-1}$  (Wiese & Fuhr 2007). The line profile of each component is:

$$\phi_i(v) = \left( \frac{4 \ln(2)}{\pi} \right)^{1/2} \frac{1}{\Delta v_{\text{FWHM},i}} \exp \left( \frac{-(v - v_{\text{LSR},i})^2 4 \ln(2)}{\Delta v_{\text{FWHM},i}^2} \right) \quad (3)$$

where  $v_{\text{LSR},i}$  the local standard of rest (LSR) velocity and  $\Delta v_{\text{FWHM},i}$  the full-width-half-maximum velocity line width of the line from component  $i$ . The upper and lower state column densities of a particular transition are related through the Boltzmann-equation by the excitation temperature of the transition:

$$\frac{N_{u_i}}{N_{l_i}} = \frac{g_{u_i}}{g_{l_i}} \exp \left( \frac{\Delta E_{t,i}}{k_b T_{\text{ex},t}} \right) \quad (4)$$

where  $\Delta E_t = h\nu_t$ . The optical depth of transition  $t$  and component  $i$  can thus be alternatively expressed by its lower state column density as:




Fig. 7: M17 SW [O I] 63  $\mu\text{m}$ , [O I] 145  $\mu\text{m}$  and [C II] mosaic. The two dotted vertical lines per spectrum are located at 22 and 24  $\text{km s}^{-1}$  respectively, added to guide the eye to the absorption dips. The left upper corner in each box gives the offsets of each spectrum.

$$\tau_{i,t}(v) = \frac{c^3}{8\pi\nu_i^3} \phi_i(v) \frac{g_{u_i}}{g_l} N_{l,i} A_t (1 - \exp(-h\nu_i/k_b T_{\text{ex},i})) \quad (5)$$

The parameters describing the line ( $t = 145, 63$ ) intensity of each background ( $i = b$ ) and foreground ( $i = f$ ) component are thus four parameters: its excitation temperature,  $T_{\text{ex},i}$ , either its upper-state,  $N_{u_i}$ , or lower state,  $N_{l,i}$  column density, and its line width,  $\Delta v_{\text{FWHM},i}$  and its line center position,  $v_{\text{LSR},i}$ .

We assume that the excitation temperatures of all background components is the same (sufficient to obtain good fits, see below), so that we have one common value for the excitation temperature of the [O I] 145  $\mu\text{m}$  transition:  $T_{\text{ex},b,145} \equiv T_{\text{ex},145,bg}$ ;  $b = 1 \dots B$  and another common one for the [O I] 63  $\mu\text{m}$  transition:  $T_{\text{ex},b,63} \equiv T_{\text{ex},63,bg}$ ;  $b = 1 \dots B$ , and correspondingly for the [O I] 63  $\mu\text{m}$  foreground lines. Note that for the three-level [O I] system, the upper-state column density of [O I] 63  $\mu\text{m}$  is the same as the lower-state column density of the [O I] 145  $\mu\text{m}$  line. Thus, three parameters, namely the column density in any one of the three states (or the total column density) and the two excitation temperatures of both transitions, are sufficient to completely describe each component.

#### 4.1. Fitting the background emission

The observed line profiles are then fitted by a two-layer model with a background in emission ( $b$ ) composed of several components ( $N_b$ ) and an absorbing foreground ( $f$ ) with a different num-

ber of components ( $N_f$ ), each with the fit parameters above. As we do not know the details of the excitation of the three levels (two excitation temperatures) of the atomic oxygen fine structure level system, we have used the peak main beam brightness temperature of [O I] 63  $\mu\text{m}$ , shining through in between the absorption notches, for the estimation of the background excitation temperature of [O I] 63  $\mu\text{m}$  by applying the inverse Rayleigh-Jeans correction. Then, we derive the background [O I] 145  $\mu\text{m}$  excitation temperature via the analytic solution to the balance equation between collisional excitation and de-excitation with spontaneous emission. See Appendix B for a detailed description of this procedure. The outcome is almost independent of the assumed density. The resulting kinetic temperatures in the background layer are given in Table 3 for a density of  $10^6 \text{ cm}^{-3}$ . In Section 5 below, we further investigate how sensitive the derived column densities of the background and foreground layers are against variations of the assumed density and kinetic temperature of the background layer.

We start with the fit of the observed [O I] 145  $\mu\text{m}$  line profiles. First, we fit the background emission in the [O I] 145  $\mu\text{m}$  line profile by several Gaussian emission line background components. The excitation temperature is fixed to the common value,  $T_{\text{ex},b,145} \equiv T_{\text{ex},145,bg}$ , as discussed above. The fit parameters are the upper state column density of each background component  $b$ ,  $N_{u_{145},b}$ , their LSR velocity  $v_{\text{LSR},b}$ , and their line width  $\Delta v_{\text{FWHM},b}$ , using the lowest possible number of Gaussian components. The process is iterative, and we increase the number of components one by one until there is no substantial decrease in




Fig. 8: M17 SW channel maps of the integrated intensity emission of [O I] 145  $\mu\text{m}$  in contours (with levels at 6, 13, 19, 26, 32, 39, 45, 52, 58, 65, 71 and 78  $\text{K km s}^{-1}$ ) and [O I] 63  $\mu\text{m}$  in color. The angular resolution for both lines. Same as Fig. 5, we can express the units of the map as K.

Table 3: Mon R2 and M17 SW background excitation temperatures for both transitions and their respective kinetic temperatures.

Positions	$T_{\text{ex},63}^{\text{a}}$ (K)	$T_{\text{ex},145}^{\text{b}}$ (K)	$T_{\text{kin}}^{\text{b}}$ (K)
MonR2 1	78	144	90
MonR2 2	140	227	152
M17SW 0	95	200	114
M17SW 1	65	113	74
M17SW 2	70	125	80
M17SW 3	86	170	102
M17SW 4	145	240	158
M17SW 5	67	117	77
M17SW 6	100	222	122

<sup>a</sup> RJ-corrected peak  $T_{\text{mb},63}$  shining through.

<sup>b</sup> Derived from 3-level rate equation solution, see App. B, at  $n_{\text{H}_2} = 10^6 \text{cm}^{-3}$ .

the chi-square of the fit with additional components, hence we stop the iteration. This way, the simple, close to Gaussian profiles of [O I] 145  $\mu\text{m}$  require one or two Gaussian components,  $B = B_{145}$ , for the fitting.

Next, we identify the contribution of these  $B_{145}$  components to the line wing emission of the lower transition, [O I] 63  $\mu\text{m}$ . The [O I] 63  $\mu\text{m}$  upper-level column density is fixed at the value derived from the upper state column density of the [O I] 145  $\mu\text{m}$  line, converted using the excitation temperature of the background via the Boltzmann-relation (Eq. 4). The line center position,  $v_{\text{LSR},b}$ , is also fixed within a narrow range (10% of the value) for each component, namely to the value fitted for the [O I] 145  $\mu\text{m}$  profile, and the [O I] 63  $\mu\text{m}$  excitation temperature is fixed to the value of  $T_{\text{ex},63,b}$  derived above. The fit results (see tables C.1, C.2, and C.3 to C.9) show that the fitted width is typically less than 10% smaller for the [O I] 63  $\mu\text{m}$  line, and thus consistent with the width fitted for the upper line. The fitting is restricted to the [O I] 63  $\mu\text{m}$  line wings not affected by foreground absorption; the relevant velocity ranges are from 2 to 5 and 14 to 18  $\text{km s}^{-1}$  for Mon R2 and from 12 to 15 and 25 to 28  $\text{km s}^{-1}$  for M17S SW respectively.

The thus fitted [O I] 63  $\mu\text{m}$  line profiles, resulting from the  $b = 1 \dots B_{145}$  background components, show (outside of the core emission, which is heavily blended with the self-absorption) weak fit residuals requiring  $B_{63}$  additional background velocity components, which we number by  $b = B_{145} + 1, \dots, B_{145} + B_{63}$ . We fit these by fixing the excitation temperature to the value used for the background emission. However, we now allow the column densities of the additional [O I] 63  $\mu\text{m}$  emission components,  $N_{u_{63},b}$ , to vary freely, as well as their line widths and


central velocities,  $\Delta v_{\text{FWHM},b}$ ,  $v_{\text{LSR},b}$ . These components are not visible in [O I] 145  $\mu\text{m}$  due to their low column density as derived from the fit. This effect is verified by calculating the corresponding [O I] 145  $\mu\text{m}$  background emission, which turns out to be of the order of 0.1 K or lower. We follow a step-by-step procedure instead of a simultaneous fitting of both [O I] lines because the weak background components are only visible in [O I] 63  $\mu\text{m}$  while the main components come from [O I] 145  $\mu\text{m}$ . Therefore, we do not expect a one-to-one correlation between the two [O I] lines.

The total column densities of oxygen are listed in Table 4, obtained from the upper and lower [O I] transitions derived from fitting the background emission profiles for the [O I] 145  $\mu\text{m}$  transition (background only) and the [O I] 63  $\mu\text{m}$  transition. These column densities are obtained by adding up all velocity components, where, however, the total column density is dominated by the one to two components  $b = 1 \dots B_{145}$  dominating the [O I] 145  $\mu\text{m}$  emission. The detailed fit parameters of all components and levels are listed in the Appendix C.


#### 4.2. Fitting of the foreground component

In the next step, we deal with the foreground absorption visible in the [O I] 63  $\mu\text{m}$  line. The column densities of the foreground absorption components are derived by fitting the absorption profiles against the fitted line profile of the [O I] 63  $\mu\text{m}$  background emission. The intensity in the self-absorption notches drops to very low values. This requires correspondingly low [O I] 63  $\mu\text{m}$  excitation temperatures of the absorbing material. Due to the high energy of the [O I] 63  $\mu\text{m}$  transition, the resulting optical depth hardly depends on the exact value of  $T_{\text{ex}}$ . Therefore, we fix the foreground excitation temperature to a single and low value of  $T_{\text{ex},f} = 20$  K. The population of the [O I] 63  $\mu\text{m}$  upper level is very small at these low excitation temperatures, so the total column density also is very insensitive to the fixed  $T_{\text{ex},f}$ . In Section 5.2, we quantify this minimal sensitivity of the fitted foreground column density on the exact value of the assumed value of  $T_{\text{ex},f}$ . Because of the low excitation temperature of the [O I] 63  $\mu\text{m}$  transition the [O I] 63  $\mu\text{m}$  upper level, providing the lower level of the [O I] 145  $\mu\text{m}$  transition, is hardly populated so that the foreground does not contribute to the observed [O I] 145  $\mu\text{m}$  line profile at all, neither in emission nor in absorption, independent of its assumed excitation temperature. Therefore, the foreground fit is done to the [O I] 63  $\mu\text{m}$  profile alone. This approach is backed up by the lack of any self-absorption nodges in the observed [O I] 145  $\mu\text{m}$  line profile.

With the excitation temperature fixed to this low value, this leaves the foreground column densities,  $N_{l_{63},f}$ , line width,  $\Delta v_{\text{FWHM},f}$ , and line center,  $v_{\text{LSR},f}$  as the fitting parameters of the foreground components. As the [C II] and [O I] absorption dips are very similar in line center velocity and width, we use the line center position and width of the [C II] foreground components fitted by Guevara et al. (2020) as initial guesses for the foreground [O I] fitting of the most substantial absorption components. We keep the velocity of the [C II] components fixed, allowing for a minor variation in width, and we freely vary the oxygen column density. An example of the fitting is shown in Fig. 9, which shows the results for position 2 in Mon R2 for both [O I] lines. The thus fitted column densities of the foreground are listed in Table 5. In the tables in Appendix C, we have marked in bold letters the column density of the components that significantly contribute to the total column density at every position. We also added a column with the [C II] column densities per component (extracted from Guevara et al. 2020) for the compo-



(a) Mon R2 [O I] 145  $\mu\text{m}$  fitting for position 2. *Top Left* - This panel represents the fitted model in blue and the observed spectra in red. *Bottom Left* - Residual curve between the observed spectra and the model. *Right* - Composition of each fitted component in cyan.



(b) Mon R2 [O I] 63  $\mu\text{m}$  fitting for position 2. *Top Left* - This panel represents the fitted model in blue and the observed spectra in red. *Bottom Left* - Residual curve between the observed spectra and the model. *Top Right* - Composition of each fitted component in cyan and the fitted [O I] 63  $\mu\text{m}$  background profile in orange. *Bottom Right* - Foreground optical depth for the [O I] 63  $\mu\text{m}$  line of each Gaussian component in pink.

Fig. 9: Mon R2 [O I] fits for position 2 for both transitions. The background fit needs 7 components, the 2 coming from the [O I] 145  $\mu\text{m}$  line carrying the bulk of the column density, plus 5 additional weak ones. The foreground fit needs 6 components.

nents where we have a match in velocity between the [O I] and [C II].

Some Gaussian components have unphysically narrow line widths, well below  $1 \text{ km s}^{-1}$ . We can attribute these components to the constraints of the fit. We are fitting emission profiles that are not necessarily Gaussian with Gaussian shapes. This creates a need to cover some gaps in the fit with these narrow components. They all have small amplitudes so that they do not contribute significantly to the column densities derived.

Table 4: Mon R2 and M17 SW [O I] 145  $\mu\text{m}$  and [O I] 63  $\mu\text{m}$  background column density per position for the two layer model.

Positions	[O I] <sub>145</sub>		[O I] <sub>63</sub>		$N_{bg}(\text{O}^0)^b$ ( $\text{cm}^{-2}$ )
	# Back. Comp.	$\tau_{bg}^*$ <sup>a</sup>	# Back. Comp.	$\tau_{bg}^*$ <sup>a</sup>	
MonR2 1	3	0.8	7	42.1	5.1E+19
MonR2 2	2	0.3	7	6.3	1.3E+19
M17SW 0	3	0.3	7	11.1	1.3E+19
M17SW 1	1	0.8	2	72.1	5.0E+19
M17SW 2	1	0.7	2	48.5	3.9E+19
M17SW 3	2	0.2	3	9.7	1.3E+19
M17SW 4	2	0.3	3	5.4	9.8E+18
M17SW 5	3	0.1	3	6.6	1.3E+19
M17SW 6	2	0.4	2	13.5	1.5E+19

<sup>a</sup>  $\tau^*$  corresponds to the peak optical depth of the peak component, calculated from the fitted parameters according to eq. 2.

<sup>b</sup> Total column density, calculated from the sum of the three levels.

Table 5: Mon R2 and M17 SW [O I] 63  $\mu\text{m}$  foreground column density per position for the two layer model.

Positions	#	$\tau_{fg}^*$ <sup>a</sup>	$N_{fg}(\text{O}^0)$
	Fore. Comp.		( $\text{cm}^{-2}$ )
MonR2 1	4	12.1	4.6E+18
MonR2 2	6	5.4	3.9E+18
M17SW 0	7	3.6	1.8E+18
M17SW 1	4	5.7	3.3E+18
M17SW 2	6	3.1	3.4E+18
M17SW 3	5	3.9	3.0E+18
M17SW 4	8	2.8	2.5E+18
M17SW 5	5	3.0	3.9E+18
M17SW 6	6	1.8	2.5E+18

<sup>a</sup> Same as in Table 4, resp. eq. 2.

<sup>b</sup> Same as in Table 4.

### 4.3. Foreground and background column densities

The derived total column density of the [O I] background and foreground, given by the sum of the individual velocity components are listed in Tables 6 and 7, respectively. For easier comparison, the column densities are also converted to hydrogen column densities,  $N_H$ , and to *equivalent visual extinctions*, assuming that all oxygen is in the form of  $\text{O}^0$ . From Wakelam & Herbst (2008, their table 1, for high-metal elemental abundances), we use an O/H abundance ratio of  $2.56 \times 10^{-4}$ . We consider the canonical conversion factor between the total hydrogen column density and visual extinction of  $1.87 \times 10^{21} \text{ cm}^{-2}/A_V$  (Bohlin et al. 1978). Tables 6 and 7 also compare the total oxygen column densities of the foreground and background components to the ones derived previously for [C II]. The  $\text{C}^+$  column density also has been converted to an equivalent visual extinction using a C/H abundance value from Wakelam & Herbst (2008, same as above, table 1) of  $1.2 \times 10^{-4}$  and the canonical conversion factor listed above to convert to magnitudes. The tables also list the ratio of the [O I] and [C II] column densities and the ratio of the equivalent extinctions. The abundance ratio between oxygen and carbon is 2.1, when using the elemental abundances relative to hydrogen quoted above.

In order to easier visualize the multiple foreground and background components, both for the new [O I] and the former [C II] data, and in particular also for the discussion of the position-to-position variations below (see Section 5.4), we compare the fitted parameters for different components in the foreground and background, namely their velocities, velocity widths and column densities, in Figs. 13 and 14. For the same purpose, we visualize these fitted parameters for both species and for both the background and foreground components in Figs. 17 to 16 by showing each component in a bar representation.

We notice that the background column density ratio of [O I] to [C II] is close to the elemental abundance in most positions; correspondingly, the equivalent  $A_V$  ratio is close to unity. This is expected if all material is fully excited. However, due to the higher critical density of the [O I] transitions it is likely that more  $\text{C}^+$  is traced in emission. Therefore, we expect to measure a background column density ratio between oxygen and ionized carbon where the elemental abundance is the upper limit. However the value for Mon R2, position 1, in the background layer, clearly sticks out; the other two cases for an [O I] excess in the background layer well above the elemental ratio are positions 1 and 5 in M17 SW. We note that for these positions, we use a relatively low  $T_{ex,63,bg}$ , derived from the relatively low peak brightness of the [O I] 63  $\mu\text{m}$  line. The latter is a lower limit, which holds in the optically thick case; higher values for  $T_{ex}$  are perfectly feasible and would result in a correspondingly lower [O I] column density, as discussed in Section 5.1. On the other hand, positions 0, 4, and 6 for M17 SW have a lower oxygen column density compared to the other positions. As discussed above, these values could be an effect of a too-low density of the collision partners to excite the oxygen; see Sect 5.1.1 for an explanation of how density affects the results.

For the foreground layer, we expect an opposite situation. In cold material some atomic oxygen may be left in overall molecular material where most carbon is bound in CO. Therefore we should measure a [O I] to [C II] abundance ratio higher than the elemental abundance in absorption. Mon R2 presents the higher ratios for both sources. Given the likely scenarios for the structure of Mon R2 (see Sect. 5.4), it is expected that this excess is due to the inherent structure of the source. M17 SW shows a different behavior. The values that deviate from what is expected are found in positions 0, 2, 4, and 6. Interestingly, the affected positions are closer to the main ridge (see Fig. 2). It is clear that a particular spatial configuration plays a role here. But with the foreground layer being invisible and only being detected through the absorption profile of bright atomic lines, it is hard to speculate without much information about it. Future studies over the whole map for the foreground layer in both lines could help to resolve this issue.

## 5. Discussion

We have estimated the background and foreground column densities for neutral atomic oxygen under the assumption of a lower limit for the excitation temperature in the background layer along the line of reasoning discussed in App. B. With the estimated range of excitation temperatures for [O I] 63  $\mu\text{m}$  between 65 and 145 K, we derive column densities from the fit to the observed line profiles between  $8.5$  and  $49 \times 10^{18} \text{ cm}^{-2}$  for the background layer and between  $1.8$  and  $4.6 \times 10^{18} \text{ cm}^{-2}$  for the foreground layer. In the following, we first evaluate how sensitively the results are affected by the necessary simplifying assumptions we made. Then, we turn to a discussion of the characteristics of the foreground layer.

Table 6: Mon R2 and M17 SW comparison between oxygen and ionized carbon column densities and equivalent extinctions for the background columns.

Positions	Background					
	$N_{bg}(O^0)$ ( $\text{cm}^{-2}$ )	$N_{bg}(C^+)^a$ ( $\text{cm}^{-2}$ )	$\frac{N(O^0)}{N(C^+)}$	$A_{V,bg} O^{0b}$ (mag.)	$A_{V,bg} C^{+a}$ (mag.)	$\frac{A_{V,bg} O^0}{A_{V,bg} C^+}$
MonR2 1	5.1E+19	4.2E18	12.1	106	19	5.6
MonR2 2	1.3E+19	4.7E18	2.8	28	21	1.3
M17SW 0	1.3E+19	9.2E18	1.4	28	41	0.7
M17SW 1	5.0E+19	8.0E18	6.3	105	36	2.9
M17SW 2	3.9E+19	5.6E18	3.8	82	25	1.8
M17SW 3	1.3E+19	4.4E18	7	28	20	1.4
M17SW 4	9.8E+18	7.6E18	1.3	20	34	0.6
M17SW 5	1.3E+19	3.0E18	4.3	27	13	2.1
M17SW 6	1.5E+19	7.7E18	1.9	31	34	0.9

<sup>a</sup> [ $^{12}\text{C II}$ ] column densities from Guevara et al. (2020).

<sup>b</sup> Equivalent visual extinction for the [O I] and [C II] column densities (factors for abundance and conversion from  $N(\text{H})$  as discussed in the text), with  $N(\text{H}) = 1.87 \times 10^{21} \text{ cm}^{-2} A_V$ .

Table 7: Mon R2 and M17 SW comparison between oxygen and ionized carbon column densities and equivalent extinctions for the foreground columns.

Positions	Foreground					
	$N_{fg}(O^0)$ ( $\text{cm}^{-2}$ )	$N_{fg}(C^+)^a$ ( $\text{cm}^{-2}$ )	$\frac{N(O^0)}{N(C^+)}$	$A_{V,fg} O^{0b}$ (mag.)	$A_{V,fg} C^{+a}$ (mag.)	$\frac{A_{V,fg} O^0}{A_{V,fg} C^+}$
MonR2 1	4.6E+18	8.3E17	5.5	9.7	3.7	2.6
MonR2 2	3.9E+18	6.4E17	5.9	8.1	2.9	2.8
M17SW 0	1.8E+18	2.0E18	1.0	3.8	9.2	0.4
M17SW 1	3.3E+18	1.7E18	1.9	6.8	7.6	0.9
M17SW 2	3.4E+18	3.0E18	1.1	7.1	13	0.5
M17SW 3	3.0E+18	7.7E17	4.0	6.2	3.5	1.8
M17SW 4	2.5E+18	1.3E18	1.8	5.1	5.8	0.9
M17SW 5	3.9E+18	3.9E17	10	8.2	1.7	4.8
M17SW 6	2.5E+18	1.8E18	1.4	5.3	8.0	0.7

<sup>a</sup> [ $^{12}\text{C II}$ ] column densities from Guevara et al. (2020).

<sup>b</sup> Equivalent visual extinction of the total [O I] and [C II] column densities (factors for abundance and conversion from  $N(\text{H})$  as discussed in the text), with  $N(\text{H}) = 1.87 \times 10^{21} \text{ cm}^{-2} A_V$ .

### 5.1. Effect of Variations in the background physical parameters

The choice of the value for the excitation temperature of the background layer in both [O I] transitions is a critical element of the study. As presented above, we have taken the excitation temperature of the [O I] 63  $\mu\text{m}$  transition in the background as the lower limit derived from the peak brightness temperature of this transition observed at each position; with the assumption of high density, necessary because of the high critical density of the [O I] 145  $\mu\text{m}$  transition, we then use the analytical solution for the population of the three-level [O I] fine structure system (see App. B) to derive the excitation temperature for the upper transition (assuming  $10^6 \text{ cm}^{-3}$  for the density). In order to estimate how sensitive the derived column densities for background layer components and the foreground components are to these assumptions, we now analyze how the result depends on varying first the density and then the kinetic temperature of the background components (along the algorithm presented in App. B).

#### 5.1.1. Effects of varying the density of the background layer

First, let us start with the density. As the default, we have assumed a density of  $10^6 \text{ cm}^{-3}$  for deriving the upper transition excitation temperature from lower transition one, estimated from the peak brightness of the [O I] 63  $\mu\text{m}$  line. Fig. 10 shows how changes in the assumed collision partner density, and hence different values for the derived  $T_{\text{ex},145}$ , result in different best-fit results for the background (dashed line) and foreground column density (solid line). We have selected two positions for each of the sources to be shown for this analysis. The other positions are affected in the same way.

The analysis shows that an increase in density leads to a corresponding increase in the estimated background column density and vice versa. This proportionality can be explained according to Fig. B.2, which illustrates the effect that the upper [O I] line shows a regime of population inversion. Any increase in density results in a decrease in the [O I] 145  $\mu\text{m}$  excitation temperature (and vice versa) and, consequently, an increase in the [O I] column density necessary to maintain the observed intensity. In comparison, the [O I] 63  $\mu\text{m}$  excitation temperature remains constant. Note that M17 SW position 4 can only be fit-




Fig. 10: Best fit column densities compared for selected positions against a range in the assumed density. The background is indicated by dashed lines and the foreground in solid ones.

ted with densities higher than  $10^5 \text{ cm}^{-3}$ . For lower densities, the [O I]  $145 \mu\text{m}$  peak intensity can not be reached. It is worth noting that position 4 has the highest [O I]  $145 \mu\text{m}$  intensity and it is located at the [O I] peak, suggesting that the density here should be much higher than the other positions.

The foreground column density necessary for the deep absorption features depends very little on the background layer's density variation. This insensibility is evident as the [O I]  $63 \mu\text{m}$  absorption is highly optically thick so that the increase in the background intensity to be absorbed does not need to be compensated by an enhanced foreground absorption. The modest variations visible in the fit result originate in second-order effects. Depending on the details of the line profiles, the optically thinner wings of the absorbing layers are more or less sensitive to an increase or decrease in the background intensity resulting from the density variations.

### 5.1.2. Effects of varying the kinetic temperature of the background layer

Now, let us continue analyzing the sensitivity of the derived background and foreground column densities against the second physical parameter, the kinetic temperature of the background, keeping the density fixed at  $10^6 \text{ cm}^{-3}$ . Above, we have assumed the kinetic temperature as derived from the upward Rayleigh-Jeans corrected peak brightness temperature of the [O I]  $63 \mu\text{m}$  emission shining through in between the absorption dips, giving a lower limit to the excitation temperature. As the kinetic temperature is always larger than the excitation temperature (see Fig. B.2) for any density, this is also a lower limit to the kinetic temperature of the background. We now vary the kinetic temperature of the background from this minimum value up to higher values. We calculate the excitation temperatures for both [O I] lines from the analytic formulas at the given temperature and density (see App. B), and perform the least-squares fit of the line profiles with these parameters.




Fig. 11: Best fit column densities compared for selected positions against a range in the assumed kinetic temperatures.

We stop increasing the kinetic temperature of the background once the background column density drops to the level of the foreground, breaking our initial assumption of having most of the material concentrated in the warm background layer. This upper limit gives the maximum kinetic temperature used in the fitting, which turns out to be roughly 50 K above our minimum value. Therefore, we have a reduced range of about 50 K for valid kinetic temperatures where our assumptions remain accurate.

Fig. 11 shows the thus fitted column densities for both layers against the range of background kinetic temperatures for selected positions fitted in Mon R2 and M17 SW. The figure shows that in order to fit the observed brightness of the background emission, the necessary background column density is roughly inversely proportional to the excitation temperature and, hence, the kinetic temperature.

The foreground column density is only slightly affected by the changes in the background. Increases in the background kinetic temperature lead to a rise in the background excitation and main beam temperatures, reducing the optical depth due to the severe reduction in column density. Then, the higher background intensity requires more foreground material to absorb the background emission for obtaining the observed self-absorbed profile. In any case, the range of variation is lower than an order of magnitude, keeping the foreground density well constrained.

To conclude, the discussion on the effect of varying the physical parameters of the background material, namely density (in Section 5.1.1) and kinetic temperature (in this Section), shows that the resulting background column densities span a range in physical scenarios of warm PDR material with correlated temperatures; they reach to about 10 times larger column densities for the background compared to the minimum derived for the nominal parameters and cannot be constrained further. However, the important result about the foreground column density is that it is very well constrained independent of the assumptions about the physical parameters of the background.




Fig. 12: Best fit foreground column densities compared for selected positions against a range in the assumed foreground excitation temperatures.

### 5.2. Effect of variations in the foreground excitation temperature

We now analyze how changes in the foreground excitation temperature assumed for the [O I] 63  $\mu\text{m}$  line affect the foreground column density. For the previous analysis, we have assumed a temperature of 20 K. Figure 12 shows how the foreground column density derived from the line profile changes when the foreground's excitation temperature varies. The foreground excitation temperature was varied from 15 K to 45 K. We have selected 15 K as a lower limit because this value is the lowest possible for [C II] from the energy balance (Kabanovic et al. 2022), a reasonable assumption where both atoms should coexist. There is insufficient material to absorb the background in some positions for lower values than this, establishing a lower limit to the foreground excitation temperature. The upper value of 45 K is an appropriate upper limit for the foreground gas to still produce visible absorption nodes; higher values come too close to the background values.

The column density is not much affected by the assumed value for the foreground excitation temperature, changing monotonically by not more than 20% over this temperature range. Thus, we conclude that the foreground column density of the absorption components, derived from the fit to the complex line profiles, is a relatively robust result.

### 5.3. Column Densities of the different components of the Background Layer

One of the central motivations behind this work is to study the nature of the foreground layer identified through the self-absorption traced by both cooling lines, [C II] and [O I]. Nevertheless, we also discuss the nature of the background layer. In Fig. 13, we plot the column density for each fitted Gaussian velocity component for both [C II] and [O I] for each position analyzed. [C II] is characterized by diamonds and [O I] by circles. Figs. 17 and 15 compare the fitted background components for each position in a column density bar plot.

One or two components dominate the background column density contribution for both atoms around the systemic velocity, 10  $\text{km s}^{-1}$  for Mon R2 and 21  $\text{km s}^{-1}$  for M17 SW with more than 80% of contribution from these main components. We also see that the additional components tracing the line wings contribute only marginally to the column density, with Mon R2 showing a red-shifted line wing and M17 SW showing emission at both sides skewed to the blue side of the spectra.

We notice that main background emission, both in [C II] and in [O I], is fitted by a small number of well overlapping Gaussian components, resulting in a smooth overall profile. Sometimes even only one single component dominates, in particular in [C II], where the bulk background emission is defined by the [<sup>13</sup>C II] line profile. The [O I] background emission, having to match both the [O I] 145  $\mu\text{m}$  profile and the line wings of heavily self-absorbed [O I] 63  $\mu\text{m}$  profile, sometimes needs several more Gaussian components, which are, however heavily overlapping and sum-up to a similar total column density than the single [C II] component (an example are the three components at M17 SW pos. 0, or the two components in [O I] matching the single component in [C II] at M17 SW pos. 3; Fig. 17).

Nonetheless, the dominant contribution in column density comes from components, where both are present, and which have similar velocities and line-width in [C II] and [O I]. We can clearly see that the background is not fully uniform and shows some variations from position to position. To conclude, the background, at the different positions, shows a smooth line structure dominated by one or two Gaussian components and consistent velocities around the systemic velocity.

### 5.4. Physical Properties of the Foreground Layer

The significant result from the previous observations and analysis (Guevara et al. 2020) of the [C II] self-absorption notches was that they require a large foreground column of [C II] with very low  $T_{\text{ex}}$  material ( $\leq 20$  K). The present study shows that the [O I] 63  $\mu\text{m}$  line also shows deep self-absorption notches, which agree well in general terms in velocity and line width with the [C II] self-absorption when both are present and correspond to similar total column densities.

Similar to the background emission components analysis shown in Fig. 13, we display the column densities, velocities, and line widths of the individual foreground components in Fig. 14. The figure demonstrates that the self-absorption notches in [C II] and [O I] 63  $\mu\text{m}$  have similar velocities and narrow line widths. Moreover, the different components at the different positions can be grouped around common velocities, with 8 and 12  $\text{km s}^{-1}$  for Mon R2, and 16, 18, 21, and 24  $\text{km s}^{-1}$  for M17 SW. However, the exact velocity for each component fluctuates around these average values, the magnitude of these fluctuations being on the order of the line widths.

The total column density at each position is distributed between different velocities, and the column density of a given velocity component varies from position to position. This is best seen in Figs. 18 and 16. As discussed above, most velocity components in the foreground are seen in both species, [O I] and [C II], with similar fit parameters. In several cases, the foreground absorption being fitted by a single component in one species, requires two velocity components with different widths in the other. Examples are M17 SW, pos. 0, where the [O I] 63  $\mu\text{m}$  absorption at 24.1  $\text{km s}^{-1}$  is fitted by a single component with a width of 0.6  $\text{km s}^{-1}$  (see Table C.3), whereas the corresponding [C II] absorption at 24.1  $\text{km s}^{-1}$  needs two components, one with a wider line width of 2  $\text{km s}^{-1}$  and a narrower one with a width




Fig. 13: Compilation of background Gaussian parameters for all positions from the multi-component analysis from Section 3. Each position is plotted in columns along the velocity axis, showing both atomic lines. Circles symbolize the [O I] Gaussian components for each position and triangles the [C II] ones (from Guevara et al. (2020)). The color of each symbol represents its column density. The length of the vertical bars denotes half of the line width value for a better visualization.

of also  $0.6 \text{ km s}^{-1}$ . A similar situation is shown for the  $21.3 \text{ km s}^{-1}$  center velocity component in M17 SW, pos. 3, where the [C II] absorption is fitted by a single component of width  $2.2 \text{ km s}^{-1}$  (Table F.4 in Guevara et al. (2020)) and the [O I] absorption needs two fit components, one with width  $2 \text{ km s}^{-1}$ , the other one with a width of  $1.1 \text{ km s}^{-1}$  (see Table C.6). These difference are presumably due to the oversimplifying assumption of purely Gaussian absorption line profiles; the different fit components nevertheless can safely be counted as contributions to the same absorption feature.

There are very few occasions where an absorption component only shows up in [O I], and not in [C II]. This only occurs in the line wings and for features with relatively low column density; examples are the  $25.9 \text{ km s}^{-1}$  and the  $16.9 \text{ km s}^{-1}$  component in M17 SW, pos. 3 (see Table C.6), or the  $15.5 \text{ km s}^{-1}$  component in M17 SW, pos. 6 (see Table C.9). In all these cases the

significance of the absorbing foreground fit components may be spurious, as the absorption is fitted against the weak and somewhat noisy line wings of the background [O I]  $145 \mu\text{m}$  emission line profiles.

In contrast, there are several cases where the [C II] line shows clear absorption components, but where the corresponding [O I]  $63 \mu\text{m}$  absorption shows much lower column densities or is almost absent. This is typically the case for the [C II] absorption on the low velocity side of the line profiles in M17 SW. Examples are the  $16.4$  and  $19.1 \text{ km s}^{-1}$  components at pos. 0, see Table F4 in Guevara et al. (2020), and the corresponding [O I]  $63 \mu\text{m}$  absorption component at  $17.4 \text{ km s}^{-1}$ , or the  $16.6$  and  $19.9 \text{ km s}^{-1}$  [C II] components in pos. 4 (see Table F4 as above) and the corresponding [O I]  $63 \mu\text{m}$  component at  $19.6 \text{ km s}^{-1}$  (see Table C.7). As the [O I] emission does not show the weak wing emission below  $17 \text{ km s}^{-1}$ , these features cannot show up in absorption in

[O I]  $63 \mu\text{m}$ , even if they might be present in the foreground, as they lack background emission to absorb against.

Both cases discussed above occur in the wings of the main line emission. We conclude that the quantitative comparison of [O I] and [C II] column densities in these components might be affected by the slightly different form of the emission line profiles from the background. Similarly, the quantitative comparison of the absorbing column densities from position to position, as discussed in the next paragraph, will be affected, for the features in the line wings, by corresponding variations in the line profile of the background emission.

#### 5.4.1. Spatial variations and density estimate

We can obtain a lower limit estimate for the density of the foreground material from the variations in the column density between adjacent positions. Based on the uncertainties for column density values of the foreground absorption against the background line wings, as discussed above, we focus for the analysis of the variation in column density of the foreground components from position to position, on the absorbing components in the line cores, namely the features at velocities between 18 to 25  $\text{km s}^{-1}$ . Comparing the absorption feature near 24.1  $\text{km s}^{-1}$  at pos. 0 and pos. 1 of M17 SW, i.e. between a position on the bright interface ridge to one further into the molecular cloud, which has a total H I-equivalent column density of  $2.8 \times 10^{21} \text{ cm}^{-2}$  at pos. 0 and drops to  $0.8 \times 10^{21} \text{ cm}^{-2}$  at pos. 1, hence shows a decrease by  $2.0 \times 10^{21} \text{ cm}^{-2}$  in [C II], and from  $4.5 \times 10^{21} \text{ cm}^{-2}$  to  $0.6 \times 10^{21} \text{ cm}^{-2}$ , hence a decrease of  $3.9 \times 10^{21} \text{ cm}^{-2}$  in [O I]. Comparing the column density variations between positions along the ridge, i.e. pos. 3 and pos. 0, the velocity components at about 21.3 to 21.7  $\text{km s}^{-1}$  drop in [O I] from  $(3.5 + 0.6 \times 10^{21} \text{ cm}^{-2})$  to  $2.0 \times 10^{21} \text{ cm}^{-2}$ , i.e. a drop by 2.1, whereas it increases from 4.2 to 6.2, i.e. an increase by 2.0 in [C II]. The increase in [C II] versus the decrease in [O I] in this case indicates a change in the chemical composition along the line of sight from positions to position.

Repeating this comparison for all pairs of adjacent positions (using the fit results as displayed in Figs. 18 and 16), we see that the variations from position, both in [O I] and [C II], are typically of order 1 to  $2 \times 10^{21} \text{ cm}^{-2}$  and peak variations are up to about  $4 \times 10^{21} \text{ cm}^{-2}$ . This includes the variations in column density between the two positions measured in Mon R2.

In order to understand to what degree the variations in the absorbing column densities from position to position might be an artifact of the ambiguity of the foreground absorption fit with several Gaussian components, we checked how the resulting fit looks like, if we fix the foreground components to the parameters from the adjacent position, first in all three parameters: line center, width and column density. We have done this test for position 1 in Mon R2 with the parameters of position 2 and position 6 in M17 SW with the parameters of position 0. Fig. 19 shows the resulting fit for position 6 as an example. In a second test, we allow each component's column density and line width to vary, keeping the velocity (and excitation temperature) fixed (Fig. 20 shows an example of the resulting fit). The resulting fits are listed from Tables C.10 to C.13. Both attempts were unsuccessful, as demonstrated by the significant residuals well above the noise level, with (obviously) slightly better results when we allowing for more variations. Considering that the foreground absorption is multiplicative,  $e^{-\tau_{fg}}$ , implying that the optical depth is determined as a proportion of the background intensity and hence independent of its actual intensity and background intensity variation, this clearly demonstrates that the variation in column den-

sity, and hence optical depth, between adjacent positions derived from the full fits of the two-layer model as discussed above, is a significant result. Keeping the column density constant across adjacent positions, as would be the case for a smooth and homogeneous foreground component, is in contradiction with the observed line profiles.

Thus, we derive the variations in H I equivalent column density for each species to be between 1-2 and up to  $4 \times 10^{21} \text{ cm}^{-2}$ . These are either variations of the total column density, i.e. decrease or increase in both species; or they indicate variation of the chemical composition along each line of sight, in case the variation is in the opposite sense for both species. The spatial scales associated with the rapid spatial variation at the distance of the M17 SW positions correspond to 0.27 pc and 0.12 pc for Mon R2.

As discussed in Guevara et al. (2020), assuming that the components observed are not all pencil like directed along the line of sight, the magnitude of the column density variation, combined with the linear scale corresponding to the angular separation at the distance of the source, gives a constraint on the variation of the density of the foreground material, and hence a lower limit estimate of the density of the foreground material. With the numbers quoted above, this results in minimum hydrogen volume densities of 1.2 to 2.4, maximum about  $5 \times 10^3 \text{ cm}^3$  for M17 SW, and correspondingly higher, according to closer distance and hence smaller spatial scale of the distance between the observed positions, i.e. 2.8 to  $5.6 \times 10^3 \text{ cm}^3$ , for Mon R2.

We emphasize that these densities are a lower limit due to several factors: first, the densities are derived from beam averaged column densities. Thus, if the source has structure on scales smaller than the beam, the column densities, and hence the derived densities, will be accordingly higher. Secondly, the densities are derived from the smallest observed distance, the pixel spacing of the upGREAT array; similar changes on smaller scales, as suggested by the profile changes shown for the fully sampled [C II] map and the regridded [O I] map in Fig. 7 would give correspondingly larger densities. In addition, the densities can be higher if the absorbing material is a surface layer significantly smaller than the spatial size of the variations. And lastly, the derivation of the H I column densities assumes that all the material is in the form of  $\text{O}^0$  and  $\text{C}^+$ ; they would be correspondingly higher, and similarly for the volume densities, if only a fraction of the material is in the form of  $\text{O}^0$  or  $\text{C}^+$ , a possibility that is already indicated by the variation of the  $\text{O}^0$  and  $\text{C}^+$  column density ratios for the different components and positions.

As a side note, we can estimate the kinetic temperatures for these densities that would result in the excitation temperature we have used for our analysis, 20 K, from the excitation temperature plots in Appendix B. This gives a foreground kinetic temperature of 30 K for Mon R2 and 38 K for M17 SW. The total column density of the low  $T_{\text{ex}}$ , absorbing foreground is very large as shown in Table 7, corresponding to an equivalent visual extinction of several  $A_V$  or more. The agreement in velocity implies that it is associated with the source. Based on the arguments above about the spatial (in projection) variation of the derived column densities, we estimate it to have a relatively high density. Both the column and the volume densities are lower limits, assuming that all its carbon is in the form of  $\text{C}^+$  and all its oxygen is in the form of  $\text{O}^0$ . These arguments together strongly suggest that the material is not a diffuse, smooth foreground layer, but is directly associated with the background material, i.e. the dense, strongly UV illuminated PDR material.

The nature and origin of this material are rather puzzling: one would need a mechanism that produces such a large column




Fig. 14: Same as Fig. 13 for the foreground parameters.

of low  $T_{\text{ex}}$ , dense material, but where the carbon stays ionized although the high density should favor rapid recombination, as is firmly predicted by PDR scenarios. At the same time, the oxygen stays in atomic form despite the significant extinction and high density that would tend to transform  $\text{O}^0$  to CO, depleting all the available carbon and also forming OH or  $\text{H}_2\text{O}$ .

#### 5.4.2. Chemical composition

Assuming that carbon might be only partially ionized, with a part as neutral atomic carbon or in a molecular form such as CO, would increase the amount of material not visible in any other tracer. In parallel, additional oxygen would have to be present in molecular form, presumably water ice, frozen out on dust. The attractive scenario of putting the additional carbon and oxygen into carbon monoxide can be ruled out observationally, as the low-J CO lines in particular those from rare isotopologues do not show a comparable column of cold CO. This analysis holds for the positions in M17 SW (Pérez-Beaupuits et al. 2015a), where

the  $\text{O}^0$  and  $\text{C}^+$  column densities are consistent with the elemental abundances.

For the positions where the derived column density ratio of oxygen to carbon is higher than the elemental abundance, namely positions 3 and 5 in M17 SW and both positions in Mon R2, we can assume that carbon in the foreground absorbing material is indeed only partially ionized, with a fractional ionization of 20 to 50%. Alternatively, the absorbing layer may be composed of two chemically different components, where a column density of about 20 to 50% of the total would have to be low  $T_{\text{ex}}$ , fully ionized [C II] material. For the Mon R2 lines of sight, the assumption of additional absorbing material is perfectly reasonable. It is known that Mon R2 is an embedded source with several young stellar objects (YSO, Beckwith et al. 1976) inside the molecular cloud illuminated from the backside. This darkening could help to explain the large discrepancy between the  $\text{O}^0$  and  $\text{C}^+$  column densities, compared to M17 SW, but does not explain the correlation between the [O I] and [C II] emission, because we would not expect the presence of ionized carbon in the dense molecular foreground material. Pilleri et al. (2012) pro-




Fig. 15: The fitted components of the background emission in Mon R2 at each position observed with high S/N, both for [O I] (this paper; top) and [C II] (from the former study by Guevara et al. (2020), bottom) at different scales. The components are shown as bars with an amplitude given in column density per velocity width on an equivalent  $N_{\text{H}1}$  column density scale. The width of each bar is the FWHM width of the component. The (integrated) column density of each component is indicated by the number at the top.




Fig. 16: Same as Fig. 15, but for the foreground components at the two positions observed for Mon-R2.

posed in their model for Mon R2, apart from the dense PDR, a surrounding envelope with densities of  $5 \times 10^4 \text{ cm}^{-3}$ , molecular hydrogen column density of  $5 \times 10^{22} \text{ cm}^{-2}$  and kinetic temperature of 35 K, similar values that we find here for the foreground layer and its estimated density from the differences between positions, confirming our results. Hence, we have for Mon R2 a foreground separated in two phases, with a temperature gradient, where one part of the gas contains only  $\text{O}^0$  coexisting with molecular material, while in the other, outer layer  $\text{O}^0$  and  $\text{C}^+$  coexist.

M17 SW shows a different picture. The foreground column density ratios between [O I] and [C II] are lower than the elemental ratio inside the main ridge and higher outside, showing some spatial correlation playing a role. As discussed above in Sect 4.3, studies over the whole map are needed. Single pointings are insufficient for a proper analysis of spatial effects. More-

over, Hoffmeister et al. (2008) found an optical foreground extinction of 2 mag, a significantly lower value than the foreground layer derived here. Therefore, we propose a scenario of a dense atomic foreground layer where ionized carbon and oxygen coexist with a density of  $10^3 \text{ cm}^{-3}$ , not ruling out the existence of a density gradient towards the exterior, leading to an additional atomic diffuse layer.

### 5.5. Comparison Between Gas and Dust

We compare the column densities derived above from the [O I] and [C II] fine-structure lines, both for the background and foreground components, with the total gas column densities derived from the sub-mm-wavelength dust emission. The column density maps were derived from Herschel/PACS and SPIRE (Griffin et al. 2010) 250, 350, and 500  $\mu\text{m}$  for Mon R2 (Rayner et al. 2017) and M17 SW (Schneider, N., priv. comm.), using the method described by Palmeirim et al. (2013). The method estimates the total gas column density from the optically thin dust emission for both sources (Table 8) by fitting the spectral energy distributions (SEDs) and computes the gas surface density distribution of the region pixel by pixel. Note that this method deals with only a single temperature component of the dust emission and is thus biased towards the bulk column density of the lower temperature gas of the molecular clouds. It also traces parts of the warm PDR material, which has bright dust emission, but the analysis weighted it down by constraining the dust emission wavelength bands to longward of 160  $\mu\text{m}$ . Also, the values estimated are upper values in a 18'' beam because there is line-of-sight contamination, with  $A_V \sim 10$  mag for M17 and  $A_V \sim 2$  mag for Mon R2 (Schneider et al., in prep.).

For M17 SW, the gas column densities are between 54 and 336 magnitudes. Background equivalent extinction values derived from [O I] at all positions are 1 to 10 times smaller than the ones derived from dust. This difference is unsurprising as the dust emission also traces the dense molecular material of the bulk molecular cloud. The warm PDR traced by the background emission in the fine structure lines contributes 10 to 25% of the total column density traced by the dust emission (Table 8). We also note that the column density of the warm PDR background derived here is larger than the one derived from the mid- and high-J CO lines (Harris et al. 1987; Pérez-Beaupuits et al. 2015a) and quoted in previous studies of the [C II] emission (Stutzki et al. 1988). The low-velocity resolution of the previous [C II] observations ignored that the foreground absorption significantly weakens the [ $^{12}\text{C II}$ ] emission. We note that the cold and dense molecular cloud material responsible for the bulk dust emission is not traced by the column densities derived from the [O I] lines, observed in emission. This disparity is plausible as oxygen will be mainly in molecular form (CO,  $\text{H}_2\text{O}$ ) and partially frozen out onto dust grains.

For Mon R2, the situation is different. The equivalent extinctions derived from dust for both positions are much closer to those derived from the [O I] fine structure lines. This closeness could be interpreted as neutral oxygen that coexists with dense foreground molecular material, as has been suggested before by several authors (Pilleri et al. 2012; Ginard et al. 2012; Treviño-Morales et al. 2019). As discussed above in Sect 5.4, the dust strengthens the scenario of a double-layered foreground, with a component in molecular and another one in atomic form.

The comparison of the column densities derived from the emission from the warm background in [C II] and [O I] shows that both are consistent with the emission originating in standard PDR material, with excitation temperatures between 60 and




Fig. 17: Same as Fig. 15, but for the background components fitted at the M17 SW positions.

Table 8: Mon R2 and M17 SW dust equivalent extinction comparison.

Component	$A_V$ Dust (mag.)	Background				Foreground			
		$A_{V,bg} O^0$ <sup>a</sup> (mag.)	$\frac{A_V^{Dust}}{A_{V,bg} O^0}$	$A_{V,bg} C^{+a}$ (mag.)	$\frac{A_V^{Dust}}{A_{V,bg} C^+}$	$A_{V,fg} O^{0b}$ (mag.)	$\frac{A_V^{Dust}}{A_{V,fg} O^0}$	$A_{V,fg} C^{+b}$ (mag.)	$\frac{A_V^{Dust}}{A_{V,bg} C^+}$
MonR2 1	73	106	0.7	19	3.8	9.7	7.5	3.7	20
MonR2 2	40	28	1.4	21	2.9	8.1	4.9	2.9	14
M17SW 0	253	28	9.0	41	12	3.8	64	9.2	28
M17SW 1	336	105	3.2	36	9.3	6.8	49	7.6	44
M17SW 2	282	82	3.4	25	11.3	7.1	40	7.6	44
M17SW 3	328	28	12	20	16	6.2	53	3.5	94
M17SW 4	167	20	8.4	34	4.9	5.1	33	5.8	29
M17SW 5	54	27	2.0	13	4.2	8.2	6.6	1.7	32
M17SW 6	88	31	2.8	34	2.6	5.3	17	8.0	11

<sup>a</sup> Extracted from Table 6.

<sup>b</sup> Extracted from Table 7.

140 K for the [O I] and between 150 and 250 K for the [C II]. As discussed in our previous analysis, the amount of material is significantly larger than expected for a single PDR layer, so we have to invoke multiple PDR layers within the beam with several tens of magnitudes in  $A_p$ . The column densities derived here refer only to the bright PDR material in the fine structure line emission. A portion of the warm, dense PDR material will contribute

to the mid- and high-J CO emission. These lines can trace the material because we have worked under the assumption that all [O I] emission is in the form of atomic material, not considering the molecular contribution.




Fig. 18: Same as Fig. 17, but for the foreground components fitted at the M17 SW positions.

### 5.6. Comparison with other models

Are there alternative explanations for the nature of the foreground layer? Goldsmith et al. (2021) discuss the [O I] profile toward W3, which shows strong absorption, similar to what this paper discusses. At first sight, the situation looks similar to the Mon R2 case discussed here. They find similar values for the neutral oxygen foreground column density, from 2 to  $7 \times 10^{18} \text{ cm}^{-2}$ , reinforcing the idea of a foreground layer of atomic material with similar column densities. They attribute its foreground absorption to geometry effects because the source is heated from the backside, allowing for additional oxygen in atomic form in cold molecular layers, together with CO, between the PDR and the observer. However, such a scenario cannot explain the sizeable  $\text{C}^+$  column densities together with more prominent  $\text{O}^0$  column densities at the same velocity and line width. The [C II] absorption dips, requiring a large column of [C II] much colder than feasible in a PDR layer, rule out a scenario similar to the one discussed for W3 by Goldsmith et al. (2021). The analysis continues in Goldsmith et al. (2023), with the modeling, through the Meudon code (Le Petit et al. 2006), of the foreground layer in W3A. Their model identifies foreground hydrogen column densities of  $2 \times 10^{22} \text{ cm}^{-2}$ , or equivalently a oxygen column density of  $\sim 5 \times 10^{22} \text{ cm}^{-2}$ , with a density of

$n(\text{H}) = 250 \text{ cm}^{-3}$  over a spatial extension of 26 pc. Where the interpretation of the absorption profile in [O I] resulting from a smooth foreground maybe a valid scenario in the case of W3A, the [O I] absorption observed in Mon R2 and M17 SW must be due to high density material, based on the differences in foreground column densities between positions over small angular distances. Mapping observations in [O I] are not available for W3A and hence a decision between these two different scenarios is not possible with the presently available data. In addition, a  $^{13}\text{C II}$  profile with sufficient S/N is not available in the case of W3A, so that the background column density of [C II] is not well constrained. W3A might behave like our sources, but maps in [O I] and [C II] would be needed to distinguish between the scenarios. Regrettably, new observations in these atomic cooling lines are not feasible in the near future.

Kabanovic et al. (2022) present a similar scenario for RCW120 for [C II] only. [C II] observations show correlations between the velocity of foreground components derived from a similar multi-component double-layered Gaussian analysis and atomic hydrogen absorption dips in the line profile. Through a H I self-absorption (HISA) analysis, they compared the H I absorption line profile with that of [C II]. They found correlations in line width and LSR velocity for some components. They attributed the origin of the foreground layer causing the [C II] self-




Fig. 19: Unsuccessful (note the significantly increased level of the residuals) fit for M17 SW pos. 6, using the foreground parameters fitted to position 0.




Fig. 20: Same as Fig. 19, using the foreground parameters fitted to position 0, but allowing for variations in width and column density.

absorption to diffuse and extended H I that coexists with C<sup>+</sup>. [O I] 63  $\mu\text{m}$  was observed in the source (Kabanovic, S., priv. comm.), but there is only faint emission in the southwestern area. Besides, no [O I] 145  $\mu\text{m}$  observations are available, making it hard to verify any self-absorption effects in the [O I] 63  $\mu\text{m}$  profiles. New observations in both oxygen lines at high resolution and S/N would be needed for an analysis similar to ours.

## 6. Summary

The newly-observed high spectral resolution observations of the [O I] 145  $\mu\text{m}$  and the [O I] 63  $\mu\text{m}$  lines towards M17 SW and Mon R2 show strong absorption notches in the [O I] 63  $\mu\text{m}$  line and allow for a detailed multi-component Gaussian analysis, separating the emission into a background and foreground layer. The background emission is consistent with warm, dense PDR-material emission in both sources, well-known and characterized

by many previous studies. This material shows smooth Gaussian emission profiles towards which we formerly observed intense [<sup>12</sup>C II] self-absorption in both sources and now newly detected [O I] 63  $\mu\text{m}$  foreground absorption. The deep foreground self-absorption features require significant column densities of low  $T_{\text{ex}}$  atomic oxygen in the foreground material.

Comparison with the previously observed [<sup>12</sup>C II] and [<sup>13</sup>C II] profiles towards both Mon R2 and M17 SW show that the [O I] 145  $\mu\text{m}$  emission line profiles, tracing the warm background, are entirely consistent with the [<sup>13</sup>C II] emission profiles from the background material, and have similar total column densities. The intense self-absorption notches visible in [O I] 63  $\mu\text{m}$  match well with the previously observed [<sup>12</sup>C II] absorption features, both in their central velocities and widths. Hence, we conclude that, for the majority, they trace the same material, which has to be very cold (in terms of excitation temperature) due to the deep absorption. The column densities in the absorption layer are also consistent for both species and the particularities for each source. The large column density of the foreground material and the small spatial scales derived from the sudden variation of the central velocity and depth of the absorption features constrain the material to have significant density, between  $10^3$  and  $10^4 \text{ cm}^{-3}$ . The nature of this foreground layer, with fully ionized carbon and atomic oxygen at relatively high densities and low excitation temperature, is very puzzling.

We emphasize that the foreground absorption is only visible at high spectral and spatial resolution, available with the upGREAT instrument on SOFIA and in Galactic sources. Spatial averaging, i.e., observations of sources at a larger distance, namely in nearby or further out galaxies, resulting in lower spectral resolution observations, smooth out the absorption features in the line profiles, leading to reduced integrated intensity. Thus, the absorbing foreground absorption component of unidentified origin may be, at least partially, responsible for the [O I] and [C II] line intensity deficiencies quoted for such sources. Large scale mapping observations with high signal to noise in both [C II] and [O I] fine structure lines and their analysis in galactic clouds are necessary to resolve this issue. We analyzed one of the best datasets that combines both [C II] isotopes together with both [O I] transitions, and without access to new observations at the same resolution and S/N in the near future, it will be hard to disentangle this scenario. Still, we can affirm, without doubt, that there is a cold, relatively dense foreground layer associated with the main PDR composed of ionized carbon and neutral oxygen absorbing the emission of the background.

*Acknowledgements.* This work is based on observations made with the NASA/DLR Stratospheric Observatory for Infrared Astronomy (SOFIA). SOFIA was jointly operated by the Universities Space Research Association, Inc. (USRA), under NASA contract NNA17BF53C, and the Deutsches SOFIA Institut (DSI) under DLR contract 50 OK 0901 to the University of Stuttgart. The work was carried out within the Collaborative Research Centre (CRC) 956 sub-projects A4 and C1 (project ID 184018867) and CRC 1601 sub-projects A6 and B2, funded by the Deutsche Forschungsgemeinschaft (DFG), project ID 500700252. This research was carried out in part at the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration (80NM0018D0004).

## References

- Beckwith, S., Evans, N. J., I., Becklin, E. E., & Neugebauer, G. 1976, ApJ, 208, 390  
 Bohlin, R. C., Savage, B. D., & Drake, J. F. 1978, ApJ, 224, 132  
 Boreiko, R. T. & Betz, A. L. 1996, ApJ, 464, L83  
 Colditz, S., Beckmann, S., Bryant, A., et al. 2018, Journal of Astronomical Instrumentation, 7, 1840004

- Fernández-Ontiveros, J. A., Spinoglio, L., Pereira-Santaella, M., et al. 2016, *ApJS*, 226, 19
- Fischer, C., Beckmann, S., Bryant, A., et al. 2018, *Journal of Astronomical Instrumentation*, 7, 1840003
- Ginard, D., González-García, M., Fuente, A., et al. 2012, *A&A*, 543, A27
- Goldsmith, P. F. 2019, *ApJ*, 887, 54
- Goldsmith, P. F., Langer, W. D., Seo, Y., et al. 2021, *ApJ*, 916, 6
- Goldsmith, P. F., Langer, W. D., Seo, Y., et al. 2023, *ApJ*, 952, 102
- Goldsmith, P. F., Yıldız, U. A., Langer, W. D., & Pineda, J. L. 2015, *ApJ*, 814, 133
- González-Alfonso, E., Fischer, J., Graciá-Carpio, J., et al. 2012, *A&A*, 541, A4
- Griffin, M. J., Abergel, A., Abreu, A., et al. 2010, *A&A*, 518, L3
- Guan, X., Stutzki, J., Graf, U. U., et al. 2012, *A&A*, 542, L4
- Guevara, C., Stutzki, J., Ossenkopf-Okada, V., et al. 2020, *A&A*, 636, A16
- Gusdorf, A., Anderl, S., Lefloch, B., et al. 2017, *A&A*, 602, A8
- Harris, A. I., Stutzki, J., Genzel, R., et al. 1987, *ApJ*, 322, L49
- Heyminck, S., Graf, U. U., Güsten, R., et al. 2012, *A&A*, 542, L1
- Hoffmeister, V. H., Chini, R., Scheyda, C. M., et al. 2008, *ApJ*, 686, 310
- Hollenbach, D. J. & Tielens, A. G. G. M. 1999, *Reviews of Modern Physics*, 71, 173
- Kabanovic, S., Schneider, N., Ossenkopf-Okada, V., et al. 2022, *A&A*, 659, A36
- Kirsanova, M. S., Ossenkopf-Okada, V., Anderson, L. D., et al. 2020, *MNRAS*, 497, 2651
- Klein, R., Reedy, A., Fischer, C., et al. 2023, *ApJ*, 945, 29
- Kristensen, L. E., Gusdorf, A., Mottram, J. C., et al. 2017, *A&A*, 601, L4
- Le Petit, F., Nehmé, C., Le Bourlot, J., & Roueff, E. 2006, *ApJS*, 164, 506
- Laurini, S., Wyrowski, F., Wiesemeyer, H., et al. 2015, *A&A*, 584, A70
- Lique, F., Klos, J., Alexander, M. H., Le Picard, S. D., & Dagdigan, P. J. 2018, *MNRAS*, 474, 2313
- Meixner, M., Haas, M. R., Tielens, A. G. G. M., Erickson, E. F., & Werner, M. 1992, *ApJ*, 390, 499
- Mookerjee, B., Sandell, G., Güsten, R., et al. 2019, *A&A*, 626, A131
- Palmeirim, P., André, P., Kirk, J., et al. 2013, *A&A*, 550, A38
- Pérez-Beaupuits, J. P., Güsten, R., Spaans, M., et al. 2015a, *A&A*, 583, A107
- Pérez-Beaupuits, J. P., Stutzki, J., Ossenkopf, V., et al. 2015b, *A&A*, 575, A9
- Pilleri, P., Fuente, A., Cernicharo, J., et al. 2012, *A&A*, 544, A110
- Pilleri, P., Treviño-Morales, S., Fuente, A., et al. 2013, *A&A*, 554, A87
- Rayner, T. S. M., Griffin, M. J., Schneider, N., et al. 2017, *A&A*, 607, A22
- Risacher, C., Güsten, R., Stutzki, J., et al. 2016, *A&A*, 595, A34
- Schneider, N., Röllig, M., Simon, R., et al. 2018, *A&A*, 617, A45
- Schneider, N., Simon, R., Guevara, C., et al. 2020, *PASP*, 132, 104301
- Stacey, G. J., Smyers, S. D., Kurtz, N. T., & Harwit, M. 1983, *ApJ*, 265, L7
- Stutzki, J., Stacey, G. J., Genzel, R., et al. 1988, *ApJ*, 332, 379
- Tem, P., Hoffman, D., Ennico, K., & Le, J. 2018, *Journal of Astronomical Instrumentation*, 7, 1840011
- Tielens, A. G. G. M. & Hollenbach, D. 1985, *ApJ*, 291, 722
- Treviño-Morales, S. P., Fuente, A., Sánchez-Monge, Á., et al. 2019, *A&A*, 629, A81
- Treviño-Morales, S. P., Pilleri, P., Fuente, A., et al. 2014, *A&A*, 569, A19
- Wakelam, V. & Herbst, E. 2008, *ApJ*, 680, 371
- Wiese, W. L. & Fuhr, J. R. 2007, *Journal of Physical and Chemical Reference Data*, 36, 1287
- Wu, Y. W., Reid, M. J., Sakai, N., et al. 2019, *ApJ*, 874, 94
- Young, E. T., Becklin, E. E., Marcum, P. M., et al. 2012, *ApJ*, 749, L17
- Zucker, C., Speagle, J. S., Schlafly, E. F., et al. 2019, *ApJ*, 879, 125

## Appendix A: Observational parameters

In this appendix, we provide the observational parameters of the observed positions in detail. The coordinates of each position are the same as the original [C II] observations from Guevara et al. (2020). The tables contain the absolute and relative coordinates (with respect to the source coordinate, see Table 1) for each observed position and their OFF position.

Table A.1: Monoceros R2 and M17 SW positions

	RA	DEC	Rel. Offset	Rel. Offset
	(J2000)	(J2000)	RA	DEC
	(h:m:s)	(°:′:″)	(″)	(″)
<b>Mon R2</b>				
Pos. 1	6:07:46.21	-6:23:03.01	0	5
Pos. 2	6:07:44.87	-6:23:03.01	-20	5
OFF	6:07:19.28	-6:33:08.72	-400	-600
<b>M17 SW</b>				
Pos. 0	18:20:23.46	-16:12:02.01	-60.6	-1.1
Pos. 1	18:20:22.45	-16:12:30.20	-75.1	-29.3
Pos. 2	18:20:21.34	-16:12:05.00	-91.1	-4.1
Pos. 3	18:20:22.32	-16:11:35.89	-76.9	25.0
Pos. 4	18:20:24.61	-16:11:34.82	-44.0	26.1
Pos. 5	18:20:25.70	-16:12:02.77	-28.2	-1.9
Pos. 6	18:20:24.66	-16:12:29.23	-43.3	-28.3
OFF	16:21:04.87	-16:13:07.35	537	-67

## Appendix B: Excitation temperature estimate for the [O I] line

We have estimated the excitation temperature for both [O I] transitions in a three step process. We do not have previous information about the excitation temperature of the oxygen; therefore, as a starting point we assume that [O I] 63  $\mu\text{m}$  is optically thick so that the brightness temperature traces the excitation temperature (including the Rayleigh-Jeans correction), and at least at one velocity, the background brightness temperature shines through the foreground absorption. Hence, we have selected as a lower limit for the [O I] 63  $\mu\text{m}$  the  $T_{\text{mb}}$  peak emission and then converted to  $T_{\text{ex}}$  given by the Rayleigh-Jeans correction under the assumption of optically thick emission as:

$$T_{\text{mb}} = \mathcal{J}_\nu(T_{\text{ex}}) = \frac{h\nu}{k} \left( e^{h\nu/kT_{\text{ex}}} - 1 \right)^{-1} \quad (\text{B.1})$$

With the excitation temperature of the [O I] 63  $\mu\text{m}$  line, we can estimate the excitation temperature of the [O I] 145  $\mu\text{m}$  line, though the balance between collisional excitation and de-excitation and spontaneous emission when assuming a gas density. Then, the collisional rate  $C_{ij}$  from level  $i$  to  $j$  is given by:

$$C_{ij} = R_{ij}n \quad (\text{B.2})$$

with  $n$  the density of the collision partner ( $\text{cm}^{-3}$ ) and  $R_{ij}$  the collisional rate coefficient ( $\text{cm}^{-3} \text{s}^{-1}$ ). The oxygen coefficients at different kinetic temperatures for electrons, hydrogen, and molecular hydrogen were estimated by Lique et al. (2018). Now, the ratio between adjacent levels for a three-level system is (Goldsmith et al. 2015; Goldsmith 2019):




Fig. B.1: Excitation temperature of [O I] 63  $\mu\text{m}$  for a combination of kinetic temperature and density of the gas. We have added contour levels for excitation temperatures of 20, 50, 100 and 150 K.

$$\frac{n_2}{n_1} = \frac{C_{12}(C_{01} + C_{02}) + C_{02}(A_{10} + C_{10})}{(A_{21} + C_{21} + C_{20})(C_{01} + C_{02}) - C_{20}C_{02}} \quad (\text{B.3})$$

$$\frac{n_1}{n_0} = \frac{(A_{21} + C_{21} + C_{20})(C_{01} + C_{02}) - C_{20}C_{02}}{(A_{21} + C_{21} + C_{20})(A_{10} + C_{10}) + C_{20}C_{12}} \quad (\text{B.4})$$

The excitation temperature of a transition is related to the ratio of the level population through the Boltzmann equation by:

$$\frac{n_j}{n_i} = \frac{g_j}{g_i} \exp\left[-(E_j - E_i)/k_b T_{\text{ex}ij}\right] \quad (\text{B.5})$$

with  $g$  the statistical weight and  $E$  energy of a level. Upward and downward collision rates are related by:

$$C_{ji} = C_{ij} \frac{g_i}{g_j} \exp(-h\nu_{ij}/k_b T_{\text{kin}}) \quad (\text{B.6})$$

Here, we selected molecular hydrogen as main collision partner with an orto-to-para ratio of 3:1. The resulting excitation temperatures for [O I] 63  $\mu\text{m}$  and [O I] 145  $\mu\text{m}$  as a function of kinetic temperature and density are shown in Fig. B.1 and B.2, respectively.

Using these dependencies we can obtain the [O I] 145  $\mu\text{m}$  excitation temperature from the [O I] 63  $\mu\text{m}$  excitation temperature if we either assume a particular gas density or kinetic temperature. We can simply look up the parameters in Fig. B.1 that produce the [O I] 63  $\mu\text{m}$  excitation temperature measured in the first step and go with these parameters into Fig. B.2 to obtain the [O I] 145  $\mu\text{m}$  excitation temperature. Here, we assumed a density of  $10^6 \text{ cm}^{-3}$ , as an intermediate value given the critical density of both transitions. A different method would be to fix the kinetic temperature and then search for a density. The methods are




Fig. B.2: Excitation temperature of [O I] 145  $\mu\text{m}$  for a combination of kinetic temperature and density of the gas. We have added contour levels for excitation temperatures of 100, 150, 200 and 250 K.

equivalent and the actual choice of neither the method nor the value is critical for the final output. Actually, it is not critical to precisely know the [O I] 145  $\mu\text{m}$  excitation temperature because the quantity entering the radiative transfer is the emissivity ( $B(T_{\text{ex}}) \times \tau_{145}$ ), not  $T_{\text{ex}}$  as such. In Fig. B.3, we have plotted in colors the [O I] 145  $\mu\text{m}$  emissivity ( $B(T_{\text{ex}}) \times \tau_{145}$ ) as a function of the kinetic temperature and column density on a Rayleigh-Jeans scale. We can see that the [O I] 145  $\mu\text{m}$  emissivity and the [O I] 63  $\mu\text{m}$  excitation temperature go almost in parallel. Thus, if we have a given  $T_{\text{ex},63}$ , we can read the [O I] 145  $\mu\text{m}$  emissivity with reasonable accuracy. We do not need to know the exact excitation temperature of the [O I] 145  $\mu\text{m}$  line and the corresponding gas density and temperature. We see that for a  $T_{\text{ex},63}$  of 75 K and below, the [O I] 145  $\mu\text{m}$  emissivity stays almost constant along the contours, it is independent of the actual density. Hence, we have no significant uncertainty in the background column from the method. At higher  $T_{\text{ex},63}$  we can conclude that we are still safe against variations towards higher densities, but that, for example, for  $T_{\text{ex},63} = 100$  K, we underestimate the emissivity by a factor 2 if the actual gas density is  $10^5 \text{ cm}^{-3}$  instead of  $10^6 \text{ cm}^{-3}$  so that we overestimate the background component column by this factor of two.

Altogether, we have estimated the excitation temperature in a 3-step process: 1) selection of an excitation temperature from [O I] 63  $\mu\text{m}$  main beam temperature peak, 2) estimation of the kinetic temperature of oxygen from Fig. B.1 under the assumption of a density of  $10^6 \text{ cm}^{-3}$ , together with the excitation temperature from 1), 3) estimate of the excitation temperature for [O I] 145  $\mu\text{m}$  from Fig. B.2. The derived excitation temperatures for both transitions and the kinetic ones are shown in Table 3.




Fig. B.3: [O I] 145  $\mu\text{m}$  emissivity in colors for a given column density. The contours are the [O I] 63  $\mu\text{m}$  excitation temperature like in Fig. B.1.

### Appendix C: Parameters of individual components

In this appendix, we list the fitted parameters for each Gaussian component for all the positions according to equation 1, 2 and 3. Tables C.1 and C.2 describe Mon R2 and Tables C.3 to C.9 describe M17 SW. Tables C.10 to C.13 describe the attempts of using foreground parameters of one position into another. The main components, providing more than 90% of the background and foreground material are indicated by numbers in bold print.

Table C.1: Mon R2 position 1 parameters.

Component	$N_u$ [OI] ( $\text{cm}^{-2}$ )	V ( $\text{km s}^{-1}$ )	$\Delta V$ ( $\text{km s}^{-1}$ )	$N_l$ [OI] <sup>a</sup> ( $\text{cm}^{-2}$ )	$N$ [C II] <sup>b</sup> ( $\text{cm}^{-2}$ )
T <sub>ex</sub> Background [OI] <sub>145</sub> = 144.6 K					
Comp. 1 [O] <sub>145</sub>	5.8E+16	7.8	2.1	3.4E+17	
Comp. 2 [O] <sub>145</sub>	1.8E+17	10.0	4.7	1.1E+18	
Comp. 3 [O] <sub>145</sub>	2.4E+16	11.7	11.0	1.4E+17	
T <sub>ex</sub> Background [OI] <sub>63</sub> = 77.5 K					
Comp. 1 [O] <sub>63</sub> <sup>c</sup>	3.4E+17	7.8	1.7	<b>1.1E+19</b>	
Comp. 2 [O] <sub>63</sub>	3.3E+13	9.5	0.2	1.0E+15	
Comp. 3 [O] <sub>63</sub> <sup>c</sup>	1.1E+18	10.1	3.8	<b>3.3E+19</b>	
Comp. 4 [O] <sub>63</sub> <sup>c</sup>	1.4E+17	11.8	9.6	<b>4.4E+18</b>	
Comp. 5 [O] <sub>63</sub>	2.3E+15	15.9	2.4	7.2E+16	
Comp. 6 [O] <sub>63</sub>	3.1E+15	20.7	4.1	9.6E+16	
Comp. 7 [O] <sub>63</sub>	3.1E+15	23.9	7.3	9.6E+16	
T <sub>ex</sub> Foreground = 19.9 K					
Comp. 1 [O] <sub>63</sub>	6.5E+12	2.2	2.9	<b>1.0E+18</b>	
Comp. 2 [O] <sub>63</sub>	9.5E+12	8.6	2.4	<b>1.5E+18</b>	
Comp. 3 [O] <sub>63</sub> <sup>d</sup>	1.0E+06	8.8	3.8	1.6E+11	5.9E+17
Comp. 4 [O] <sub>63</sub> <sup>d</sup>	1.4E+13	11.4	1.4	<b>2.2E+18</b>	2.1E+17

<sup>a</sup> Column density of the lower level population of the transition, calculated from  $N_u$ [OI] and T<sub>ex</sub>.

<sup>b</sup> [C II] foreground column densities from Guevara et al. (2020) when the velocities match.

<sup>c</sup> Components derived from the [O I] 145  $\mu\text{m}$  Gaussian fitting.

<sup>d</sup> Parameters of the components taken from the [C II] multi-component analysis from Guevara et al. (2020).

Table C.2: Mon R2 position 2 parameters.

Component	$N_u$ [OI] ( $\text{cm}^{-2}$ )	V ( $\text{km s}^{-1}$ )	$\Delta V$ ( $\text{km s}^{-1}$ )	$N_l$ [OI] <sup>a</sup> ( $\text{cm}^{-2}$ )	$N$ [C II] <sup>b</sup> ( $\text{cm}^{-2}$ )
T <sub>ex</sub> Background [OI] <sub>145</sub> = 227.0 K					
Comp. 1 [O] <sub>145</sub>	2.0E+17	9.9	6.3	9.3E+17	
Comp. 2 [O] <sub>145</sub>	8.9E+16	11.5	2.8	4.1E+17	
T <sub>ex</sub> Background [OI] <sub>63</sub> = 140.0 K					
Comp. 1 [O] <sub>63</sub>	1.8E+15	1.6	2.0	1.5E+16	
Comp. 2 [O] <sub>63</sub> <sup>c</sup>	9.3E+17	10.0	5.1	<b>7.9E+18</b>	
Comp. 3 [O] <sub>63</sub> <sup>c</sup>	4.1E+17	11.6	3.4	<b>3.5E+18</b>	
Comp. 4 [O] <sub>63</sub>	1.1E+16	15.7	2.1	9.0E+16	
Comp. 5 [O] <sub>63</sub>	9.1E+15	17.5	3.5	7.7E+16	
Comp. 6 [O] <sub>63</sub>	5.6E+15	20.7	5.0	4.8E+16	
Comp. 7 [O] <sub>63</sub>	2.7E+15	25.3	6.3	2.3E+16	
T <sub>ex</sub> Foreground = 20.0 K					
Comp. 1 [O] <sub>63</sub> <sup>d</sup>	5.7E+12	7.3	1.8	<b>8.9E+17</b>	9E+16
Comp. 2 [O] <sub>63</sub>	7.0E+12	7.6	4.1	<b>1.1E+18</b>	
Comp. 3 [O] <sub>63</sub> <sup>d</sup>	2.5E+07	9.7	3.7	4.0E+12	1.4E+17
Comp. 4 [O] <sub>63</sub>	1.4E+12	9.9	1.2	2.2E+17	
Comp. 5 [O] <sub>63</sub> <sup>d</sup>	4.4E+12	11.1	1.1	<b>6.8E+17</b>	2.5E+17
Comp. 6 [O] <sub>63</sub> <sup>d</sup>	6.5E+12	12.0	1.1	<b>1.0E+18</b>	1.6E+17

<sup>a</sup> Column density of the lower level population of the transition, calculated from  $N_u$ [OI] and T<sub>ex</sub>.

<sup>b</sup> [C II] foreground column densities from Guevara et al. (2020) when the velocities match.

<sup>c</sup> Components derived from the [O I] 145  $\mu\text{m}$  Gaussian fitting.

<sup>d</sup> Parameters of the components taken from the [C II] multi-component analysis from Guevara et al. (2020).

Table C.3: M17 SW position 0 parameters.

Component	$N_u$ [O I] ( $\text{cm}^{-2}$ )	V ( $\text{km s}^{-1}$ )	$\Delta V$ ( $\text{km s}^{-1}$ )	$N_l$ [O I] <sup>a</sup> ( $\text{cm}^{-2}$ )	$N$ [C II] <sup>b</sup> ( $\text{cm}^{-2}$ )
T <sub>ex</sub> Background [OI] <sub>145</sub> = 200.1 K					
Comp. 1 [O] <sub>145</sub>	7.0E+16	20.3	3.2	3.4E+17	
Comp. 2 [O] <sub>145</sub>	5.7E+16	20.7	5.2	2.8E+17	
Comp. 3 [O] <sub>145</sub>	1.0E+16	21.6	2.5	5.2E+16	
T <sub>ex</sub> Background [OI] <sub>63</sub> = 94.9 K					
Comp. 1 [O] <sub>63</sub>	3.6E+15	12.1	3.5	6.5E+16	
Comp. 2 [O] <sub>63</sub>	2.6E+15	14.5	1.7	4.8E+16	
Comp. 3 [O] <sub>63</sub> <sup>c</sup>	3.4E+17	20.3	2.6	<b>6.3E+18</b>	
Comp. 4 [O] <sub>63</sub> <sup>c</sup>	2.8E+17	20.7	4.1	<b>5.1E+18</b>	
Comp. 5 [O] <sub>63</sub> <sup>c</sup>	5.1E+16	21.5	2.0	9.4E+17	
Comp. 6 [O] <sub>63</sub>	1.3E+15	26.3	1.2	2.3E+16	
Comp. 7 [O] <sub>63</sub>	1.0E+15	27.9	1.1	1.9E+16	
T <sub>ex</sub> Foreground = 21.0 K					
Comp. 1 [O] <sub>63</sub> <sup>d</sup>	3.4E+10	17.0	0.1	2.9E+15	6.5E+17
Comp. 2 [O] <sub>63</sub>	2.3E+12	17.4	2.1	<b>1.9E+17</b>	
Comp. 3 [O] <sub>63</sub> <sup>d</sup>	9.2E+06	19.0	3.5	7.8E+11	3.5E+17
Comp. 4 [O] <sub>63</sub> <sup>d</sup>	2.0E+07	21.4	2.3	1.7E+12	7.4E+17
Comp. 5 [O] <sub>63</sub>	6.0E+12	21.7	1.7	<b>5.1E+17</b>	
Comp. 6 [O] <sub>63</sub>	8.2E+11	23.2	0.2	2.9E+16	
Comp. 7 [O] <sub>63</sub> <sup>d</sup>	1.4E+13	24.1	0.6	<b>1.1E+18</b>	3.2E+17

<sup>a</sup> Column density of the lower level population of the transition, calculated from  $N_u$ [O I] and T<sub>ex</sub>.

<sup>b</sup> [C II] foreground column densities from Guevara et al. (2020) when the velocities match.

<sup>c</sup> Components derived from the [O I] 145  $\mu\text{m}$  Gaussian fitting.

<sup>d</sup> Parameters of the components taken from the [C II] multi-component analysis from Guevara et al. (2020).

Table C.4: M17 SW position 1 parameters.

Component	$N_u$ [O I] ( $\text{cm}^{-2}$ )	V ( $\text{km s}^{-1}$ )	$\Delta V$ ( $\text{km s}^{-1}$ )	$N_l$ [O I] <sup>a</sup> ( $\text{cm}^{-2}$ )	$N$ [C II] <sup>b</sup> ( $\text{cm}^{-2}$ )
T <sub>ex</sub> Background [OI] <sub>145</sub> = 112.6 K					
Comp. 1 [O] <sub>145</sub>	1.2E+17	18.9	4.1	8.8E+17	
T <sub>ex</sub> Background [OI] <sub>63</sub> = 65.0 K					
Comp. 1 [O] <sub>63</sub>	1.2E+16	12.5	3.8	6.8E+17	
Comp. 2 [O] <sub>63</sub> <sup>c</sup>	8.7E+17	18.8	3.3	<b>4.8E+19</b>	
T <sub>ex</sub> Foreground = 20.8 K					
Comp. 1 [O] <sub>63</sub> <sup>d</sup>	5.6E+12	16.4	3.3	<b>5.3E+17</b>	5.3E+17
Comp. 2 [O] <sub>63</sub> <sup>d</sup>	1.7E+13	18.7	3.5	<b>1.6E+18</b>	4.5E+17
Comp. 3 [O] <sub>63</sub> <sup>d</sup>	1.0E+13	21.3	2.3	<b>9.8E+17</b>	5.8E+17
Comp. 4 [O] <sub>63</sub> <sup>d</sup>	1.9E+12	24.1	0.5	<b>1.6E+17</b>	9.1E+16

<sup>a</sup> Column density of the lower level population of the transition, calculated from  $N_u$ [O I] and T<sub>ex</sub>.

<sup>b</sup> [C II] foreground column densities from Guevara et al. (2020) when the velocities match.

<sup>c</sup> Components derived from the [O I] 145  $\mu\text{m}$  Gaussian fitting.

<sup>d</sup> Parameters of the components taken from the [C II] multi-component analysis from Guevara et al. (2020).

Table C.5: M17 SW position 2 parameters.

Component	$N_u$ [O I] ( $\text{cm}^{-2}$ )	V ( $\text{km s}^{-1}$ )	$\Delta V$ ( $\text{km s}^{-1}$ )	$N_l$ [O I] <sup>a</sup> ( $\text{cm}^{-2}$ )	$N$ [C II] <sup>b</sup> ( $\text{cm}^{-2}$ )
T <sub>ex</sub> Background [OI] <sub>145</sub> = 125.1 K					
Comp. 1 [O] <sub>145</sub>	1.3E+17	19.7	4.7	8.8E+17	
T <sub>ex</sub> Background [OI] <sub>63</sub> = 70.0 K					
Comp. 1 [O] <sub>63</sub>	7.2E+15	12.6	3.9	3.1E+17	
Comp. 2 [O] <sub>63</sub> <sup>c</sup>	8.8E+17	19.6	3.8	<b>3.8E+19</b>	
T <sub>ex</sub> Foreground = 19.0 K					
Comp. 1 [O] <sub>63</sub> <sup>d</sup>	1.0E+07	14.0	3.0	2.7E+12	5.5E+17
Comp. 2 [O] <sub>63</sub>	1.9E+12	16.0	3.2	<b>5.0E+12</b>	9.1E+17
Comp. 3 [O] <sub>63</sub> <sup>d</sup>	5.6E+10	16.7	0.5	1.5E+16	
Comp. 4 [O] <sub>63</sub> <sup>d</sup>	3.0E+12	19.1	3.1	<b>8.1E+17</b>	3.5E+17
Comp. 5 [O] <sub>63</sub> <sup>d</sup>	2.5E+12	21.2	2.0	<b>6.7E+17</b>	5.5E+17
Comp. 6 [O] <sub>63</sub> <sup>d</sup>	5.2E+12	24.3	2.8	<b>1.4E+18</b>	6.1E+17

<sup>a</sup> Column density of the lower level population of the transition, calculated from  $N_u$ [O I] and T<sub>ex</sub>.

<sup>b</sup> [C II] foreground column densities from Guevara et al. (2020) when the velocities match.

<sup>c</sup> Components derived from the [O I] 145  $\mu\text{m}$  Gaussian fitting.

<sup>d</sup> Parameters of the components taken from the [C II] multi-component analysis from Guevara et al. (2020).

Table C.6: M17 SW position 3 parameters.

Component	$N_u$ [O I] ( $\text{cm}^{-2}$ )	V ( $\text{km s}^{-1}$ )	$\Delta V$ ( $\text{km s}^{-1}$ )	$N_l$ [O I] <sup>a</sup> ( $\text{cm}^{-2}$ )	$N$ [C II] <sup>b</sup> ( $\text{cm}^{-2}$ )
T <sub>ex</sub> Background [OI] <sub>145</sub> = 170.4 K					
Comp. 1 [O] <sub>145</sub>	6.1E+16	20.0	4.6	3.3E+17	
Comp. 2 [O] <sub>145</sub>	3.9E+16	23.7	3.6	2.1E+17	
T <sub>ex</sub> Background [OI] <sub>63</sub> = 86.0 K					
Comp. 1 [O] <sub>63</sub>	2.2E+15	9.5	3.7	5.3E+16	
Comp. 2 [O] <sub>63</sub>	3.2E+17	20.0	3.7	<b>7.6E+18</b>	
Comp. 3 [O] <sub>63</sub>	2.1E+17	23.7	2.9	<b>4.9E+18</b>	
T <sub>ex</sub> Foreground = 19.6 K					
Comp. 1 [O] <sub>63</sub>	9.4E+11	16.9	1.6	1.5E+17	
Comp. 2 [O] <sub>63</sub>	1.0E+12	21.3	1.1	1.5E+17	
Comp. 3 [O] <sub>63</sub> <sup>d</sup>	5.6E+12	21.3	2.0	<b>8.9E+17</b>	5.1E+17
Comp. 4 [O] <sub>63</sub> <sup>d</sup>	7.7E+12	23.7	1.5	<b>1.2E+18</b>	2.6E+17
Comp. 5 [O] <sub>63</sub>	3.6E+12	25.9	2.2	5.7E+17	

<sup>a</sup> Column density of the lower level population of the transition, calculated from  $N_u$ [O I] and T<sub>ex</sub>.

<sup>b</sup> [C II] foreground column densities from Guevara et al. (2020) when the velocities match.

<sup>c</sup> Components derived from the [O I] 145  $\mu\text{m}$  Gaussian fitting.

<sup>d</sup> Parameters of the components taken from the [C II] multi-component analysis from Guevara et al. (2020).

Table C.7: M17 SW position 4 parameters.

Component	$N_u$ [O I] ( $\text{cm}^{-2}$ )	V ( $\text{km s}^{-1}$ )	$\Delta V$ ( $\text{km s}^{-1}$ )	$N_l$ [O I] <sup>a</sup> ( $\text{cm}^{-2}$ )	$N$ [C II] <sup>b</sup> ( $\text{cm}^{-2}$ )
T <sub>ex</sub> Background [OI] <sub>145</sub> = 238.6 K					
Comp. 1 [O] <sub>145</sub>	4.5E+16	20.0	2.5	2.0E+17	
Comp. 2 [O] <sub>145</sub>	1.9E+17	21.3	6.2	8.4E+17	
T <sub>ex</sub> Background [OI] <sub>63</sub> = 145.0 K					
Comp. 1 [O] <sub>63</sub>	2.3E+16	16.5	4.0	1.8E+17	
Comp. 2 [O] <sub>63</sub> <sup>c</sup>	2.0E+17	20.0	3.1	<b>1.6E+18</b>	
Comp. 3 [O] <sub>63</sub> <sup>c</sup>	8.4E+17	21.3	5.0	<b>6.7E+18</b>	
T <sub>ex</sub> Foreground = 20.7 K					
Comp. 1 [O] <sub>63</sub> <sup>d</sup>	1.3E+07	16.6	3.0	1.3E+12	2.7E+17
Comp. 2 [O] <sub>63</sub>	4.6E+07	19.6	2.1	4.6E+12	
Comp. 3 [O] <sub>63</sub> <sup>d</sup>	1.3E+12	19.6	1.2	1.2E+17	2.1E+17
Comp. 4 [O] <sub>63</sub> <sup>d</sup>	4.6E+12	21.4	2.4	<b>4.6E+17</b>	3.8E+17
Comp. 5 [O] <sub>63</sub>	9.9E+12	21.5	0.8	9.8E+16	
Comp. 6 [O] <sub>63</sub> <sup>d</sup>	6.7E+12	23.8	1.1	<b>6.5E+17</b>	1.4E+17
Comp. 7 [O] <sub>63</sub>	1.0E+13	24.6	4.1	<b>1.0E+18</b>	
Comp. 8 [O] <sub>63</sub> <sup>d</sup>	1.4E+12	26.8	3.4	1.4E+17	2.6E+17

<sup>a</sup> Column density of the lower level population of the transition, calculated from  $N_u$ [O I] and T<sub>ex</sub>.

<sup>b</sup> [C II] foreground column densities from Guevara et al. (2020) when the velocities match.

<sup>c</sup> Components derived from the [O I] 145  $\mu\text{m}$  Gaussian fitting.

<sup>d</sup> Parameters of the components taken from the [C II] multi-component analysis from Guevara et al. (2020).

Table C.8: M17 SW position 5 parameters.

Component	$N_u$ [O I] ( $\text{cm}^{-2}$ )	V ( $\text{km s}^{-1}$ )	$\Delta V$ ( $\text{km s}^{-1}$ )	$N_l$ [O I] <sup>a</sup> ( $\text{cm}^{-2}$ )	$N$ [C II] <sup>b</sup> ( $\text{cm}^{-2}$ )
T <sub>ex</sub> Background [OI] <sub>145</sub> = 117.3 K					
Comp. 1 [O] <sub>145</sub>	1.3E+16	13.6	4.0	8.7E+16	
Comp. 2 [O] <sub>145</sub>	6.3E+15	21.1	2.3	4.4E+16	
Comp. 3 [O] <sub>145</sub>	1.7E+16	22.9	7.3	1.2E+17	
T <sub>ex</sub> Background [OI] <sub>63</sub> = 66.9 K					
Comp. 1 [O] <sub>63</sub> <sup>c</sup>	8.7E+16	13.6	3.2	<b>4.3E+18</b>	
Comp. 2 [O] <sub>63</sub> <sup>c</sup>	4.3E+16	21.2	1.9	<b>2.2E+18</b>	
Comp. 3 [O] <sub>63</sub> <sup>c</sup>	1.2E+17	22.8	5.8	<b>5.9E+18</b>	
T <sub>ex</sub> Foreground = 20.0 K					
Comp. 1 [O] <sub>63</sub> <sup>d</sup>	1.1E+13	12.7	2.7	<b>1.6E+18</b>	1.8E+17
Comp. 2 [O] <sub>63</sub> <sup>d</sup>	1.2E+12	18.4	2.0	1.8E+17	1.4E+17
Comp. 3 [O] <sub>63</sub>	3.7E+12	19.9	2.7	<b>5.5E+17</b>	
Comp. 4 [O] <sub>63</sub>	9.0E+12	23.8	6.0	<b>1.4E+18</b>	
Comp. 5 [O] <sub>63</sub> <sup>d</sup>	1.6E+12	23.9	0.8	2.4E+17	7.2E+16

<sup>a</sup> Column density of the lower level population of the transition, calculated from  $N_u$ [O I] and T<sub>ex</sub>.

<sup>b</sup> [C II] foreground column densities from Guevara et al. (2020) when the velocities match.

<sup>c</sup> Components derived from the [O I] 145  $\mu\text{m}$  Gaussian fitting.

<sup>d</sup> Parameters of the components taken from the [C II] multi-component analysis from Guevara et al. (2020).

Table C.9: M17 SW position 6 parameters.

Component	$N_u$ [O I] ( $\text{cm}^{-2}$ )	V ( $\text{km s}^{-1}$ )	$\Delta V$ ( $\text{km s}^{-1}$ )	$N_l$ [O I] <sup>a</sup> ( $\text{cm}^{-2}$ )	$N$ [C II] <sup>b</sup> ( $\text{cm}^{-2}$ )
T <sub>ex</sub> Background [OI] <sub>145</sub> = 221.6 K					
Comp. 1 [O] <sub>145</sub>	9.3E+16	19.8	7.2	4.4E+17	
Comp. 2 [O] <sub>145</sub>	9.4E+16	21.5	2.9	4.4E+17	
T <sub>ex</sub> Background [OI] <sub>63</sub> = 101.0 K					
Comp. 1 [O] <sub>63</sub> <sup>b</sup>	4.4E+17	19.8	5.8	<b>6.9E+18</b>	
Comp. 2 [O] <sub>63</sub> <sup>b</sup>	4.4E+17	21.5	2.3	<b>7.0E+18</b>	
T <sub>ex</sub> Foreground = 20.2 K					
Comp. 1 [O] <sub>63</sub>	1.0E+13	15.5	4.0	<b>1.4E+18</b>	
Comp. 2 [O] <sub>63</sub> <sup>c</sup>	3.5E+12	17.2	3.1	<b>4.7E+17</b>	5.7E+17
Comp. 3 [O] <sub>63</sub> <sup>c</sup>	1.0E+12	19.1	2.2	1.4E+17	3.2E+17
Comp. 4 [O] <sub>63</sub> <sup>c</sup>	2.2E+12	21.6	1.6	2.9E+17	7.6E+17
Comp. 5 [O] <sub>63</sub>	3.0E+08	21.6	2.3	3.9E+13	
Comp. 6 [O] <sub>63</sub> <sup>c</sup>	2.0E+12	24.1	0.7	2.5E+17	1.4E+17

<sup>a</sup> Column density of the lower level population of the transition, calculated from  $N_u$ [O I] and T<sub>ex</sub>.

<sup>b</sup> [C II] foreground column densities from Guevara et al. (2020) when the velocities match.

<sup>c</sup> Components derived from the [O I] 145  $\mu\text{m}$  Gaussian fitting.

<sup>d</sup> Parameters of the components taken from the [C II] multi-component analysis from Guevara et al. (2020).

Table C.10: Mon R2 position 1 parameters using free foreground parameters of position 2.

Component	$N_u$ [O I] ( $\text{cm}^{-2}$ )	V ( $\text{km s}^{-1}$ )	$\Delta V$ ( $\text{km s}^{-1}$ )	$N_l$ [O I] <sup>a</sup> ( $\text{cm}^{-2}$ )
T <sub>ex</sub> Background [OI] <sub>145</sub> = 144.6 K				
Comp. 1 [O] <sub>145</sub>	5.8E+16	7.8	2.1	3.4E+17
Comp. 2 [O] <sub>145</sub>	1.8E+17	10.0	4.7	1.1E+18
Comp. 3 [O] <sub>145</sub>	2.4E+16	11.7	11.0	1.4E+17
T <sub>ex</sub> Background [OI] <sub>63</sub> = 77.5 K				
Comp. 1 [O] <sub>63</sub> <sup>b</sup>	3.4E+17	7.8	1.7	<b>1.1E+19</b>
Comp. 2 [O] <sub>63</sub>	1.6E+16	9.4	1.0	5.0E+17
Comp. 3 [O] <sub>63</sub> <sup>b</sup>	1.1E+18	10.1	3.8	<b>3.3E+19</b>
Comp. 4 [O] <sub>63</sub> <sup>b</sup>	1.4E+17	11.8	9.6	4.4E+18
Comp. 5 [O] <sub>63</sub>	2.3E+15	15.9	2.4	7.3E+16
Comp. 6 [O] <sub>63</sub>	3.1E+15	20.7	4.1	9.7E+16
Comp. 7 [O] <sub>63</sub>	3.0E+15	23.9	7.3	9.5E+16
T <sub>ex</sub> Foreground = 19.9 K				
Comp. 1 [O] <sub>63</sub>	5.7E+12	7.3	1.8	<b>8.9E+17</b>
Comp. 2 [O] <sub>63</sub>	7.0E+12	7.6	4.1	<b>1.1E+18</b>
Comp. 3 [O] <sub>63</sub>	2.5E+07	9.7	3.7	4.0E+12
Comp. 4 [O] <sub>63</sub>	1.4E+12	9.9	1.2	2.2E+17
Comp. 5 [O] <sub>63</sub>	4.4E+12	11.1	1.1	<b>6.8E+17</b>
Comp. 6 [O] <sub>63</sub>	6.5E+12	12.0	1.1	<b>1.0E+18</b>

<sup>a</sup> Column density of the lower level population of the transition, calculated from  $N_u$ [O I] and T<sub>ex</sub>.

<sup>b</sup> Components derived from the [O I] 145  $\mu\text{m}$  Gaussian fitting.

Table C.11: Mon R2 position 1 parameters using free foreground parameters of position 2.

Component	$N_u$ [O I] ( $\text{cm}^{-2}$ )	V ( $\text{km s}^{-1}$ )	$\Delta V$ ( $\text{km s}^{-1}$ )	$N_l$ [O I] <sup>a</sup> ( $\text{cm}^{-2}$ )
T <sub>ex</sub> Background [OI] <sub>145</sub> = 144.6 K				
Comp. 1 [O] <sub>145</sub>	5.8E+16	7.8	2.1	3.4E+17
Comp. 2 [O] <sub>145</sub>	1.8E+17	10.0	4.7	1.1E+18
Comp. 3 [O] <sub>145</sub>	2.4E+16	11.7	11.0	1.4E+17
T <sub>ex</sub> Background [OI] <sub>63</sub> = 77.5 K				
Comp. 1 [O] <sub>63</sub> <sup>b</sup>	3.4E+17	7.8	1.7	<b>1.1E+19</b>
Comp. 2 [O] <sub>63</sub>	1.6E+16	9.4	1.0	5.0E+17
Comp. 3 [O] <sub>63</sub> <sup>b</sup>	1.1E+18	10.1	3.8	<b>3.3E+19</b>
Comp. 4 [O] <sub>63</sub> <sup>b</sup>	1.4E+17	11.8	9.6	4.4E+18
Comp. 5 [O] <sub>63</sub>	2.3E+15	15.9	2.4	7.3E+16
Comp. 6 [O] <sub>63</sub>	3.1E+15	20.7	4.1	9.7E+16
Comp. 7 [O] <sub>63</sub>	3.0E+15	23.9	7.3	9.5E+16
T <sub>ex</sub> Foreground = 20.4 K				
Comp. 1 [O] <sub>63</sub>	2.6E+12	7.3	1.4	<b>3.0E+17</b>
Comp. 2 [O] <sub>63</sub>	9.3E+11	7.6	9.2	1.1E+17
Comp. 3 [O] <sub>63</sub>	2.5E+08	9.7	2.9	2.9E+13
Comp. 4 [O] <sub>63</sub>	1.4E+13	9.9	2.9	<b>1.6E+18</b>
Comp. 5 [O] <sub>63</sub>	7.1E+12	11.1	0.9	<b>8.2E+17</b>
Comp. 6 [O] <sub>63</sub>	3.8E+12	12.0	0.9	<b>4.3E+17</b>

<sup>a</sup> Column density of the lower level population of the transition, calculated from  $N_u$ [O I] and T<sub>ex</sub>.

<sup>b</sup> Components derived from the [O I] 145  $\mu\text{m}$  Gaussian fitting.

Table C.12: M17 SW position 6 parameters using fixed foreground parameters of position 0.

Component	$N_u$ [O I] ( $\text{cm}^{-2}$ )	V ( $\text{km s}^{-1}$ )	$\Delta V$ ( $\text{km s}^{-1}$ )	$N_l$ [O I] <sup>a</sup> ( $\text{cm}^{-2}$ )
T <sub>ex</sub> Background [OI] <sub>145</sub> = 221.6 K				
Comp. 1 [O] <sub>145</sub>	9.3E+16	19.8	7.2	4.4E+17
Comp. 2 [O] <sub>145</sub>	9.4E+16	21.5	2.9	4.4E+17
T <sub>ex</sub> Background [OI] <sub>63</sub> = 101.0 K				
Comp. 1 [O] <sub>63</sub>	4.4E+17	19.8	5.8	<b>6.9E+18</b>
Comp. 2 [O] <sub>63</sub>	4.4E+17	21.5	2.3	<b>7.0E+18</b>
T <sub>ex</sub> Foreground = 21.0 K				
Comp. 1 [O] <sub>63</sub>	5.5E+11	16.9	0.7	4.6E+16
Comp. 2 [O] <sub>63</sub>	1.6E+12	17.8	1.7	1.3E+17
Comp. 3 [O] <sub>63</sub>	4.6E+07	19.0	3.5	4.0E+12
Comp. 4 [O] <sub>63</sub>	8.5E+07	21.4	2.3	7.2E+12
Comp. 5 [O] <sub>63</sub>	6.0E+12	21.7	1.7	<b>5.1E+17</b>
Comp. 6 [O] <sub>63</sub>	8.7E+11	23.2	0.2	2.8E+16
Comp. 7 [O] <sub>63</sub>	1.4E+13	24.1	0.6	<b>1.1E+18</b>

<sup>a</sup> Column density of the lower level population of the transition, calculated from  $N_u$ [O I] and T<sub>ex</sub>.

<sup>b</sup> Components derived from the [O I] 145  $\mu\text{m}$  Gaussian fitting.

Table C.13: M17 SW position 6 parameters using free foreground parameters of position 0.

Component	$N_u$ [OI] ( $\text{cm}^{-2}$ )	$V$ ( $\text{km s}^{-1}$ )	$\Delta V$ ( $\text{km s}^{-1}$ )	$N_l$ [OI] <sup>a</sup> ( $\text{cm}^{-2}$ )
$T_{\text{ex}}$ Background [OI] <sub>145</sub> = 221.6 K				
Comp. 1 [O] <sub>145</sub>	9.3E+16	19.8	7.2	4.4E+17
Comp. 2 [O] <sub>145</sub>	9.4E+16	21.5	2.9	4.4E+17
$T_{\text{ex}}$ Background [OI] <sub>63</sub> = 101.0 K				
Comp. 1 [O] <sub>63</sub>	4.4E+17	19.8	5.8	<b>6.9E+18</b>
Comp. 2 [O] <sub>63</sub>	4.4E+17	21.5	2.3	<b>7.0E+18</b>
$T_{\text{ex}}$ Foreground = 20.0 K				
Comp. 1 [O] <sub>63</sub>	5.5E+12	16.9	3.1	<b>8.1E+17</b>
Comp. 2 [O] <sub>63</sub>	5.0E+12	17.8	6.8	<b>7.3E+17</b>
Comp. 3 [O] <sub>63</sub>	4.6E+06	19.0	35.0	6.8E+11
Comp. 4 [O] <sub>63</sub>	8.5E+06	21.4	22.5	1.2E+12
Comp. 5 [O] <sub>63</sub>	1.3E+12	21.7	1.3	<b>1.8E+17</b>
Comp. 6 [O] <sub>63</sub>	8.7E+10	23.2	0.2	2.7E+15
Comp. 7 [O] <sub>63</sub>	1.7E+12	24.1	0.7	<b>2.4E+17</b>

<sup>a</sup> Column density of the lower level population of the transition, calculated from  $N_u$ [OI] and  $T_{\text{ex}}$ .

<sup>b</sup> Components derived from the [OI] 145  $\mu\text{m}$  Gaussian fitting.