

REAL SPIN BORDISM AND ORIENTATIONS OF TOPOLOGICAL K-THEORY

ZACHARY HALLADAY AND YIGAL KAMEL

ABSTRACT. We construct a commutative orthogonal C_2 -ring spectrum, $\text{MSpin}_{\mathbb{R}}^c$, along with a C_2 - E_∞ -orientation $\text{MSpin}_{\mathbb{R}}^c \rightarrow \text{KU}_{\mathbb{R}}$ of Atiyah's Real K-theory. Further, we define E_∞ -maps $\text{MSpin} \rightarrow (\text{MSpin}_{\mathbb{R}}^c)^{C_2}$ and $\text{MU}_{\mathbb{R}} \rightarrow \text{MSpin}_{\mathbb{R}}^c$, which are used to recover the three well-known orientations of topological K-theory, $\text{MSpin}^c \rightarrow \text{KU}$, $\text{MSpin} \rightarrow \text{KO}$, and $\text{MU}_{\mathbb{R}} \rightarrow \text{KU}_{\mathbb{R}}$, from the map $\text{MSpin}_{\mathbb{R}}^c \rightarrow \text{KU}_{\mathbb{R}}$. We also show that the integrality of the \hat{A} -genus on spin manifolds provides an obstruction for the fixed points $(\text{MSpin}_{\mathbb{R}}^c)^{C_2}$ to be equivalent to MSpin , using the Mackey functor structure of $\underline{\pi}_* \text{MSpin}_{\mathbb{R}}^c$. In particular, the usual map $\text{MSpin} \rightarrow \text{MSpin}^c$ does not arise as the inclusion of fixed points for any C_2 - E_∞ -ring spectrum.

CONTENTS

1. Introduction	1
2. Preliminaries	4
3. The Real spin bordism spectrum	9
4. The fixed points of $\text{MSpin}_{\mathbb{R}}^c$	15
5. A Real orientation of Real spin bordism	17
6. A Real spin orientation of Real K-theory	21
References	26

1. INTRODUCTION

The main result of this paper refines all of the standard orientations of topological K-theory to a single *Real spin orientation*, $\text{MSpin}_{\mathbb{R}}^c \rightarrow \text{KU}_{\mathbb{R}} = \text{KR}$, of Atiyah's Real K-theory.

Theorem 1.1 (Propositions 3.12, 3.13, 4.5, 5.1, and 6.13). There exists a map of C_2 - E_∞ -ring spectra, $\text{MSpin}_{\mathbb{R}}^c \rightarrow \text{KU}_{\mathbb{R}}$, satisfying the following properties.

- (1) The underlying spectrum of $\text{MSpin}_{\mathbb{R}}^c$ is MSpin^c .
- (2) There exists an E_∞ -map $\text{MSpin} \rightarrow (\text{MSpin}_{\mathbb{R}}^c)^{C_2}$.
- (3) There exists a C_2 - E_∞ -map $\text{MU}_{\mathbb{R}} \rightarrow \text{MSpin}_{\mathbb{R}}^c$.

The three standard orientations, $\text{MSpin}^c \rightarrow \text{KU}$, $\text{MSpin} \rightarrow \text{KO}$, and $\text{MU}_{\mathbb{R}} \rightarrow \text{KU}_{\mathbb{R}}$, are recovered as E_∞ -maps from the map $\text{MSpin}_{\mathbb{R}}^c \rightarrow \text{KU}_{\mathbb{R}}$.

In analogy with Real bordism and Real K-theory, we use the name *Real spin bordism* to refer to the C_2 -spectrum $\text{MSpin}_{\mathbb{R}}^c$. The Real spin orientation $\text{MSpin}_{\mathbb{R}}^c \rightarrow \text{KU}_{\mathbb{R}}$ recovers the standard orientations of K-theory as follows. The Real orientation of $\text{KU}_{\mathbb{R}}$ is obtained by precomposing

with $\mathrm{MU}_{\mathbb{R}} \rightarrow \mathrm{MSpin}_{\mathbb{R}}^c$, the spin^c orientation of KU is obtained by taking underlying spectra, and the spin orientation of KO is obtained as the composite

$$\mathrm{MSpin} \rightarrow (\mathrm{MSpin}_{\mathbb{R}}^c)^{C_2} \rightarrow \mathrm{KU}_{\mathbb{R}}^{C_2} \simeq \mathrm{KO}.$$

We construct $\mathrm{MSpin}_{\mathbb{R}}^c$ as follows. Let $\mathbb{R}^{p,q}$ be the C_2 -representation $\mathbb{R}^p \oplus \mathbb{R}^q \sigma$, where σ is the sign representation. Then the complex Clifford algebra $\mathrm{Cl}(\mathbb{R}^{p,q})$ inherits a corresponding conjugate-linear C_2 -action by the universal property of Clifford algebras. This action restricts to $\mathrm{Spin}^c(p,q) \subset \mathrm{Cl}(\mathbb{R}^{p,q})$, and makes the usual representation, $\mathrm{Spin}^c(p,q) \times \mathbb{R}^{p,q} \rightarrow \mathbb{R}^{p,q}$, C_2 -equivariant. Using an appropriate model of $\mathrm{ESpin}^c(p,q)$, this induces a C_2 -action on the Thom space, $\mathrm{MSpin}^c(p,q)$, of the universal $\mathrm{Spin}^c(p,q)$ -vector bundle with fiber $\mathbb{R}^{p,q}$. We define $\mathrm{MSpin}_{\mathbb{R}}^c$ as an orthogonal C_2 -spectrum whose value on the C_2 -representation $\mathbb{R}^{p,q}$ is the C_2 -space $\mathrm{MSpin}^c(p,q)$.

Remark 1.2. Complex conjugation on $\mathrm{U}(1)$ induces a C_2 -action on $\mathrm{MSpin}^c(n)$, which describes the underlying spectrum of $\mathrm{MSpin}_{\mathbb{R}}^c$ with its induced C_2 -action. This spectrum with C_2 -action can be promoted to a genuine C_2 -spectrum, $(\mathrm{MSpin}_{\mathbb{R}}^c)^h$, called the *cofree completion* of $\mathrm{MSpin}_{\mathbb{R}}^c$. Using the completion map $\mathrm{MSpin}_{\mathbb{R}}^c \rightarrow (\mathrm{MSpin}_{\mathbb{R}}^c)^h$ and the fact that $\mathrm{KU}_{\mathbb{R}} \simeq \mathrm{KU}_{\mathbb{R}}^h$, all of the results in this paper apply to $(\mathrm{MSpin}_{\mathbb{R}}^c)^h$ as well. The question of whether or not $\mathrm{MSpin}_{\mathbb{R}}^c$ is equivalent to $(\mathrm{MSpin}_{\mathbb{R}}^c)^h$ amounts to knowing whether or not the fixed points $(\mathrm{MSpin}_{\mathbb{R}}^c)^{C_2}$ is equivalent to the homotopy fixed points $(\mathrm{MSpin}_{\mathbb{R}}^c)^{hC_2} = ((\mathrm{MSpin}_{\mathbb{R}}^c)^h)^{C_2}$, which we do not investigate in this paper.

Given the facts that $\mathrm{Spin}^c(n)^{C_2} = \mathrm{Spin}(n)$ and $\mathrm{KU}_{\mathbb{R}}^{C_2} = \mathrm{KU}_{\mathbb{R}}^{hC_2} = \mathrm{KO}$, one might expect that the C_2 -(homotopy) fixed points of $\mathrm{MSpin}_{\mathbb{R}}^c$ is equivalent to MSpin . The following theorem says that no such equivalence can cover MSO .

Theorem 1.3. There does not exist a genuine C_2 -spectrum E satisfying all of the following conditions.

- (1) The homotopy groups of the underlying spectrum are $E_*^e \cong \mathrm{MSpin}_*^c$;
- (2) The homotopy groups of the C_2 -fixed points are $E_*^{C_2} \cong \mathrm{MSpin}_*$;
- (3) The C_2 -action on E_*^e is via a ring homomorphism on MSpin_*^c ;
- (4) The diagram

$$\begin{array}{ccc} \mathrm{MSpin}_* & \xrightarrow{\mathrm{res}} & \mathrm{MSpin}_*^c \\ & \searrow u_* & \swarrow u_*^c \\ & \mathrm{MSO}_* & \end{array}$$

commutes, where u_* and u_*^c are the forgetful maps that take the underlying oriented bordism class, and $\mathrm{res}: \mathrm{MSpin}_* \cong E_*^{C_2} \rightarrow E_*^e \cong \mathrm{MSpin}_*^c$ is induced by inclusion of fixed points.

Since the maps $\mathrm{MSpin} \rightarrow (\mathrm{MSpin}_{\mathbb{R}}^c)^{C_2}$ and $\mathrm{MSpin} \rightarrow (\mathrm{MSpin}_{\mathbb{R}}^c)^{hC_2}$ from Theorem 1.1 factor the usual map $\mathrm{MSpin} \rightarrow \mathrm{MSpin}^c$, Theorem 1.3 implies that neither map is an equivalence. We do not know whether or not either $(\mathrm{MSpin}_{\mathbb{R}}^c)^{C_2}$ or $(\mathrm{MSpin}_{\mathbb{R}}^c)^{hC_2}$ is equivalent to MSpin , but since Theorem 1.3 implies that the expected comparison map cannot be an equivalence, there is no reason to expect that such an equivalence exists.

Remark 1.4. In light of the maps $MU_{\mathbb{R}} \rightarrow MSpin_{\mathbb{R}}^c \rightarrow KU_{\mathbb{R}}$, it is natural to compare $MSpin_{\mathbb{R}}^c$ to what is known about the various types of fixed points of $MU_{\mathbb{R}}$ and $KU_{\mathbb{R}}$. In particular, in the case of Real bordism, the C_2 -(homotopy) fixed points $MU_{\mathbb{R}}^{C_2} = MU_{\mathbb{R}}^{hC_2}$ is not equivalent to MO , but the *geometric fixed points*, $\Phi^{C_2} MU_{\mathbb{R}}$, is. Despite this analogy, we know that $\Phi^{C_2} MSpin_{\mathbb{R}}^c \not\cong MSpin$, since the functor Φ^{C_2} is lax monoidal, and there does not exist a ring map from MO to $MSpin$. We collect this information in a table in Figure 1.

type of C_2 -fixed points	$MU_{\mathbb{R}}$	$MSpin_{\mathbb{R}}^c$	$KU_{\mathbb{R}}$
genuine $(\)^{C_2}$	\emptyset	?	KO
homotopy $(\)^{hC_2}$	\emptyset	?	KO
geometric $\Phi^{C_2}(\)$	MO	\underline{MSpin}	\underline{KO}

FIGURE 1. Comparing the various types of fixed points of Real bordism, Real spin bordism, and Real K-theory

Remark 1.5. Even though $MU_{\mathbb{R}}$ and $MSpin_{\mathbb{R}}^c$ are both Thom spectra built using the groups $U(n)$ and $Spin^c(n)$ with $U(n)^{C_2} = O(n)$ and $Spin^c(n)^{C_2} = Spin(n)$, the analogy between them breaks down for the following reason. The C_2 -action on $U(n)$ determines the action on the $(\mathbb{R}^{n,n})$ th space of the Real bordism spectrum, while the C_2 -action on $Spin^c(n)$ determines the action on the (\mathbb{R}^n) th space of the Real spin bordism spectrum. The group that corresponds to the $(\mathbb{R}^{n,n})$ th space of $MSpin_{\mathbb{R}}^c$ is $Spin^c(n, n)$, and $Spin^c(n, n)^{C_2} \not\cong Spin(n)$.

Lastly, we point out that the cohomological Thom classes that correspond to the Real spin orientation of KR constructed in this paper were originally constructed by Atiyah in the form of the following generalized Bott periodicity theorem.

Theorem 1.6 (Atiyah [1], Theorem 6.3). Let $G \rightarrow Spin^c(p, q)$ be a C_2 -equivariant homomorphism of groups with C_2 -action, and suppose $p \equiv q \pmod{8}$. Then there is a class $u \in KU_{\mathbb{R}, G}(\mathbb{R}^{p,q})$, such that multiplication by u induces an isomorphism,

$$KU_{\mathbb{R}, G}(X) \cong KU_{\mathbb{R}, G}(\mathbb{R}^{p,q} \times X).$$

As in the case of Theorem 1.1, this theorem contains all of the classical Thom isomorphisms as special cases. For example, letting $G = Spin(8n)$ with trivial C_2 -action, $(p, q) = (8n, 0)$, and taking X to be the total space of a principal $Spin(8n)$ -bundle, this recovers the Atiyah–Bott–Shapiro Thom class in KO -theory for the spin vector bundle associated to X [2]. Similarly, the KU -Thom classes for $spin^c$ vector bundles are obtained from the identity map $Spin^c(8n) = Spin^c(8n, 0)$, and the $KU_{\mathbb{R}}$ -Thom classes for Real vector bundles are obtained from the inclusion $U(n) \rightarrow Spin^c(n, n)$.

1.1. Overview of the paper. In Section 2, we establish some terminology about Real structures and review facts that we need about orthogonal C_2 -spectra. In Section 3, we construct $MSpin_{\mathbb{R}}^c$ as a commutative monoid object in the category of orthogonal C_2 -spectra. The spaces we use to model $MSpin_{\mathbb{R}}^c$ follow the definition of $MSpin^c$ in [7], except that we

introduce (different) C_2 -actions. In section 4, we define the map $\text{MSpin} \rightarrow (\text{MSpin}_{\mathbb{R}}^c)^{C_2}$ and provide a general obstruction for such a map to be an equivalence by proving Theorem 1.3. This obstruction uses the integrality of the \hat{A} -genus on spin manifolds, together with formal properties that the homotopy Mackey functor $\pi_* \text{MSpin}_{\mathbb{R}}^c$ must satisfy. In section 5, we construct the map $\text{MU}_{\mathbb{R}} \rightarrow \text{MSpin}_{\mathbb{R}}^c$ in the category of Real spectra ([5], [6]). To do this, we use a model of $\text{MU}_{\mathbb{R}}$ that is mostly derived from our model of $\text{MSpin}_{\mathbb{R}}^c$ at the regular representations. The desired map then follows from the fact that $U(n) \rightarrow \text{Spin}^c(n, n)$ is C_2 -equivariant. Most of the work in section 5 lies in showing that our definition of $\text{MU}_{\mathbb{R}}$ is actually equivalent to Real bordism, which we do by describing the C_2 -fixed points of the Real vector bundle associated to a Real free $U(n)$ -space (Lemma 5.8). In section 6, we construct the map $\text{MSpin}_{\mathbb{R}}^c \rightarrow \text{KU}_{\mathbb{R}}$ in the category of orthogonal C_2 -spectra. Our model of $\text{KU}_{\mathbb{R}}$ follows the definition of KU in [7] in terms of spaces of graded $*$ -homomorphisms of C^* -algebras, except that we endow all relevant C^* -algebras with Real structures. We show that our model of $\text{KU}_{\mathbb{R}}$ represents Atiyah's Real K-theory by showing that the associated cohomology theory restricted to spaces with trivial C_2 -action is equivalent to KO .

1.2. Acknowledgements. There are several people we would like to thank for helping this work come to fruition: Arun Debray, Cameron Krulewski, Natalia Pacheco-Tallaj, and Luuk Stehouwer, for helping prove an earlier version of Corollary 4.7, which motivated our use of a genus in the proof of Theorem 1.3; Hassan Abdallah, Mike Hill, Kiran Luecke, and Charles Rezk, for helpful discussions relating to various aspects of this paper; and Dan Berwick-Evans, Connor Grady, Cameron Krulewski, and Vesna Stojanoska, for helpful feedback on an earlier draft. Finally, we would like to specially thank our advisors, Vesna Stojanoska and Dan Berwick-Evans, respectively, for their many hours of guidance both before and during the writing of this paper.

2. PRELIMINARIES

First, we fix notation for the categories of spaces and G -spaces that we use in this paper. Let \mathcal{T} be the category of pointed, compactly generated, weak Hausdorff spaces enriched over itself, and let \mathcal{T}^G be the \mathcal{T} -enriched category of G -objects in \mathcal{T} with equivariant maps. Lastly, let \mathcal{T}_G be the \mathcal{T}^G -enriched category with the same objects as \mathcal{T}^G , but with $\mathcal{T}_G(X, Y) = \mathcal{T}(X, Y)$ which is made into a G -space by conjugation. Then we have $\mathcal{T}^G(X, Y) = \mathcal{T}_G(X, Y)^G$.

2.1. Real structures. This section introduces what the word Real refers to in various contexts. In short, *Real* will always mean “equipped with a C_2 -action” when applied to objects, and “respects C_2 -actions” when applied to morphisms. We use the word Real to restrict to the context in which \mathbb{C} is always endowed with its fixed C_2 -action given by complex conjugation. For example, this forces Real structures on complex vector spaces to be conjugate-linear. We often denote a Real structure (or C_2 -action) by $x \mapsto \bar{x}$, when it is clear which action we are referring to.

Definition 2.1. The category of *Real groups* is the category of group objects in \mathcal{T}^{C_2} . If G is a Real group, a *Real G -space* is a G -module object in \mathcal{T}^{C_2} . We will also call the objects and morphisms of \mathcal{T}^{C_2} *Real spaces* and *Real maps*, respectively.

Remark 2.2. Given a Real group G , a Real G -space is the same data as an object of $\mathcal{T}^{C_2 \times G}$, where $C_2 \times G$ is formed from the Real structure on G .

The following definition is more general than is needed here; we include it to indicate how Real linear algebra over the complex numbers fits into a framework similar to the one above.

Definition 2.3. Let Ab^{C_2} be the symmetric monoidal category of C_2 -objects in abelian groups under tensor product. A *Real ring* is a monoid object in Ab^{C_2} . If R is a Real ring, a *Real R -module* is an R -module object in Ab^{C_2} .

Example 2.4. Complex conjugation gives \mathbb{C} the structure of a Real ring. With this Real structure, a Real \mathbb{C} -module is a complex vector space together with a complex conjugate-linear involution.

From this point on, the term *Real vector space* will mean Real \mathbb{C} -module, where \mathbb{C} is given the fixed real structure of Example 2.4. Similarly, in any \mathbb{C} -linear setting, such as C^* -algebras, \mathbb{C} is given this fixed Real structure.

Remark 2.5. Let $X, Y \in \mathcal{T}_{C_2}$. Recall that C_2 acts on $\mathcal{T}_{C_2}(X, Y)$ by conjugation, $\overline{f}(x) = \overline{f(\overline{x})}$. This turns $\text{Aut}_{\mathcal{T}_{C_2}}(X)$ into a Real group and makes X into a Real $\text{Aut}_{\mathcal{T}_{C_2}}(X)$ -space. In this paper, we often use conjugation to define Real structures on spaces of specific kinds of functions between Real spaces. In each case, the resulting action is well-defined. The main observation is that in \mathbb{C} -linear contexts, conjugation by C_2 respects \mathbb{C} -linearity while complex conjugation of values does not.

Proposition 2.6. Let V, W be Real vector spaces. Then $V \otimes W$ is a Real vector space via $\overline{v \otimes w} = \overline{v} \otimes \overline{w}$. If V, W are also Real G -spaces, on which G acts by linear maps, then $V \otimes W$ is a Real G -space via $g \cdot (v \otimes w) = (gv) \otimes (gw)$.

Proof. We have $\overline{g \cdot (v \otimes w)} = \overline{gv} \otimes \overline{gw} = \overline{g} \overline{v} \otimes \overline{g} \overline{w} = \overline{g} \cdot (\overline{v \otimes w})$. \square

Proposition 2.7. Let V be a Real vector space. Then $V \cong V^{C_2} \otimes \mathbb{C}$ as Real vector spaces, where the Real structure on $V^{C_2} \otimes \mathbb{C}$ is given by complex conjugation on the second factor.

Proof. Define $T : V \rightarrow V^{C_2} \otimes \mathbb{C}$ by $T(v) = \frac{1}{2}(v + \overline{v}) \otimes 1 - (\frac{i}{2}(v - \overline{v})) \otimes i$. First, note that T is well-defined, since $(v + \overline{v}), i(v - \overline{v}) \in V^{C_2}$. Next, T is Real, since

$$\begin{aligned} \overline{T(v)} &= \frac{1}{2}(v + \overline{v}) \otimes 1 - \left(\frac{i}{2}(v - \overline{v}) \right) \otimes \overline{i} \\ &= \frac{1}{2}(v + \overline{v}) \otimes 1 + \left(\frac{i}{2}(v - \overline{v}) \right) \otimes i \\ &= \frac{1}{2}(\overline{v} + v) \otimes 1 - \left(\frac{i}{2}(\overline{v} - v) \right) \otimes i = T(\overline{v}). \end{aligned}$$

Lastly, T is an isomorphism, since $T^{-1} : V^{C_2} \otimes \mathbb{C} \rightarrow V$ is given by $T^{-1}(v_1 \otimes 1 + v_2 \otimes i) = v_1 + iv_2$. \square

The following proposition collects various routine facts that we use in this paper.

Proposition 2.8. Let X be a Real G -space, let Y be a Real space, let Z be a Real right G -space, and let $\psi: H \rightarrow G$ be a Real homomorphism.

- (1) The action homomorphism $\varphi: G \rightarrow \text{Aut}_{\mathcal{T}_{C_2}}(X)$ is Real.
- (2) The Real space $\mathcal{T}_{C_2}(X, Y)$ is a Real G -space via $g \cdot f = f \circ g^{-1}$.
- (3) Restriction by ψ makes X into a Real H -space.
- (4) The space $Z \times_G X$ is a Real space via $[\overline{z, x}] = [\overline{z}, \overline{x}]$.

2.2. Equivariant orthogonal spectra. In this section, we recall some facts about C_2 -spectra and fix our choice of model. For more details, we refer the reader to Hill, Hopkins, and Ravenel [6] and [5], or Mandell and May [12]. Throughout, we work in the genuine equivariant context. Our primary choice of model for this is the category of orthogonal C_2 -spectra.

Given finite dimensional inner product spaces V, W , let $\text{O}(V, W)$ be the space of linear isometric embeddings $V \rightarrow W$, and let $W - V$ be the vector bundle on $\text{O}(V, W)$ whose fiber over $\iota: V \rightarrow W$ is the orthogonal complement $W - \iota V$. When V and W are orthogonal C_2 -representations, $\text{O}(V, W)$ and $W - V$ inherit compatible C_2 -actions. Let \mathcal{I}_{C_2} be the \mathcal{T}^{C_2} -enriched category whose objects are finite dimensional orthogonal C_2 -representation and whose C_2 -space of morphisms is the Thom space

$$\mathcal{I}_{C_2}(V, W) = \text{Thom}(\text{O}(V, W), W - V).$$

Composition is then given by applying the Thom construction to the vector bundle map

$$\begin{array}{ccc} (V_3 - V_2) \times (V_2 - V_1) & \longrightarrow & V_3 - V_1 \\ \downarrow & & \downarrow \\ \text{O}(V_2, V_3) \times \text{O}(V_1, V_2) & \longrightarrow & \text{O}(V_1, V_3). \end{array}$$

Definition 2.9. An *orthogonal C_2 -spectrum* is a \mathcal{T}^{C_2} -enriched functor

$$X: \mathcal{I}_{C_2} \rightarrow \mathcal{T}_{C_2}.$$

Let Sp^{C_2} be the category of orthogonal C_2 -spectra and equivariant enriched natural transformations between them. There is an adjunction,

$$\Sigma_{C_2}^\infty: \mathcal{T}^{C_2} \rightleftarrows \text{Sp}^{C_2}: \Omega_{C_2}^\infty,$$

where $\Sigma_{C_2}^\infty X$ is the orthogonal C_2 -spectrum defined by $V \mapsto \Sigma^V X$. We let S_{C_2} denote the C_2 -sphere spectrum $\Sigma_{C_2}^\infty S^0$. Let $\mathbb{R} \in \mathcal{I}_{C_2}$ denote the trivial 1-dimensional representation, $\sigma \in \mathcal{I}_{C_2}$ be the 1-dimensional sign representation, and $\rho = \mathbb{R} \oplus \sigma$ be the regular C_2 -representation.

Definition 2.10. Given an orthogonal C_2 -spectrum X , a subgroup $H \subset C_2$, and $V \in \mathcal{I}_{C_2}$, define the *V th-stable H homotopy group* to be

$$\pi_V^H X = \text{colim}_n \pi_V^H \Omega^{n\rho} X(n\rho).$$

This definition can be extended to virtual representations by defining

$$\pi_{V-W}^H X = \text{colim}_n \pi_V^H \Omega^{n\rho} X(n\rho + W).$$

Genuine equivariance endows these homotopy groups with the additional structure of a Mackey functor $\underline{\pi}_{V-W}X = \underline{\pi}_{V-W}^{(\cdot)}X$, as H varies.

Definition 2.11. A C_2 -Mackey Functor, M , consists of Abelian groups $M(C_2)$ and $M(e)$, together with group homomorphisms

$$(1) \quad \begin{array}{ccc} M(C_2) & & \text{satisfying:} \\ \text{res} \left(\begin{array}{c} \downarrow \\ \uparrow \end{array} \right)_{\text{tr}} & & (1) \overline{\text{res}(x)} = \text{res}(x); \\ M(e) & & (2) \text{tr}(\overline{y}) = \text{tr}(y); \\ \left(\begin{array}{c} \cup \\ \overline{\quad} \end{array} \right) & & (3) \overline{y} = y; \\ & & (4) \text{res}(\text{tr}(y)) = y + \overline{y}. \end{array}$$

An isomorphism of C_2 -Mackey functors is a map of diagrams (1) such that each component is an isomorphism.

Proposition 2.12. The category Sp^{C_2} together with weak equivalences given by $\underline{\pi}_*$ -isomorphisms is a homotopical category¹ which presents the genuine C_2 -stable homotopy category. This can be refined to a model category however we will not need any specifics involving (co)fibrations.

Let Sp^{BC_2} denote the category of orthogonal spectra with C_2 -action (see [15]). Restricting to the subcategory of trivial C_2 -representations in \mathcal{I}_{C_2} induces an equivalence of 1-categories [5, Proposition A.19],

$$(\)^e: \text{Sp}^{C_2} \rightarrow \text{Sp}^{BC_2}.$$

Definition 2.13. Let X be an orthogonal C_2 -spectrum. The image, X^e , under the map $(\)^e: \text{Sp}^{C_2} \rightarrow \text{Sp}^{BC_2}$ is called the *underlying spectrum* of X .

Taking homotopy groups of the underlying spectrum recovers the trivial subgroup homotopy groups of X , $\pi_*(X) \cong \pi_*(X^e)$, along with the induced C_2 -action. More care is required when discussing the C_2 -homotopy groups $\pi_*^{C_2}(X)$. The *categorical fixed points* functor

$$F^{C_2}: \text{Sp}^{C_2} \rightarrow \text{Sp},$$

is given by

$$(F^{C_2}X)(n) = (X(\mathbb{R}^n))^{C_2}.$$

However, categorical fixed points does not preserve weak equivalences, as it can fail to capture the C_2 -homotopy groups of X . This can be fixed by taking an appropriate fibrant replacement.

Definition 2.14. [15, p. 71] Let X be an orthogonal C_2 -spectrum. The C_2 -fixed points of X is the orthogonal spectrum $X^{C_2} \in \text{Sp}$ given by $(X^{C_2})(n) = (\Omega^{n\sigma} X(n\rho))^{C_2}$.

¹Recall that a homotopical category is a category with a class of weak equivalences satisfying the ‘‘two out of six’’ property and containing all identity maps.

Then we have $\pi_*^{C_2}(X) \cong \pi_*(X^{C_2})$ [15, Proposition 7.2]. Similarly, we could have modelled X^e by

$$(X^e)(n) \cong \Omega^{n\sigma} X(n\rho).$$

Levelwise inclusion of fixed points defines a map,

$$X^{C_2} \rightarrow X^e,$$

which upon taking homotopy groups induces the restriction map,

$$\text{res}: \pi_*^{C_2}(X) \rightarrow \pi_*(X),$$

in the homotopy Mackey functor $\underline{\pi}_* X$.

Remark 2.15. The categorical fixed points fit into a Quillen adjunction

$$T_{C_2}: \text{Sp} \rightleftarrows \text{Sp}^{C_2}: F^{C_2},$$

where the left adjoint comes from giving a spectrum the trivial C_2 -action and then passing through the equivalence of 1-categories $\text{Sp}^{BC_2} \simeq \text{Sp}^{C_2}$. Deriving this adjunction allows us to identify the cohomology theory represented by X^{C_2} in terms of X .

Proposition 2.16. Let E^* be the cohomology theory represented by X^{C_2} , and let $Y \in \mathcal{T}$. Then viewing $Y \in \mathcal{T}^{C_2}$ by giving it the trivial C_2 -action, from the above remark we have

$$E^n(Y) \cong [\Sigma_{C_2}^\infty Y, \Sigma^n X]_{C_2}.$$

Now, we consider the multiplicative aspects of our model. Commutative ring structures in the genuine equivariant setting is a rich subject and has been explored by Blumberg and Hill [3], as well as by others. We will be interested in the complete N_∞ -algebras, also called C_2 - E_∞ -ring spectra. The category of C_2 - E_∞ -ring spectra can be modeled using orthogonal C_2 -spectra as follows. Day convolution equips Sp^{C_2} with the structure of a closed symmetric monoidal category with monoidal unit given by S_{C_2} . As discussed in [5, Section 2.1] and [3], the category $\text{CAlg}(\text{Sp}^{C_2})$ of commutative monoid objects in Sp^{C_2} models C_2 - E_∞ -ring spectra. Further, by [14, Proposition 22.1] it follows that $\text{CAlg}(\text{Sp}^{C_2})$ is equivalent to the category of lax symmetric monoidal \mathcal{T}^{C_2} -enriched functors $\mathcal{I}_{C_2} \rightarrow \mathcal{T}_{C_2}$. This is the setting in which we will address multiplicative questions.

Definition 2.17. A *commutative orthogonal C_2 -ring spectrum* is a lax symmetric monoidal \mathcal{T}^{C_2} -enriched functor $\mathcal{I}_{C_2} \rightarrow \mathcal{T}_{C_2}$. A *(C_2 - E_∞ -) map of commutative orthogonal C_2 -ring spectra* is a symmetric monoidal C_2 -equivariant enriched natural transformation.

As is discussed in [15, p. 77], the categorical and genuine fixed points are lax symmetric monoidal functors. Thus, for a commutative orthogonal C_2 -ring spectrum X , the spectra $F^{C_2} X$ and X^{C_2} inherit the structure of commutative monoids in Sp .

Proposition 2.18. There is a symmetric monoidal natural transformation,

$$F^{C_2} \rightarrow ()^{C_2},$$

from the categorical fixed points to the C_2 -fixed points.

Such a natural transformation can be defined levelwise by letting

$$\iota: \mathbb{R}^n \rightarrow n\rho$$

be the inclusion of the trivial submodule inside of $n\rho$, taking the composition

$$S^{n\sigma} \rightarrow \mathcal{I}_{C_2}(\mathbb{R}^n, n\rho) \rightarrow \mathcal{T}_{C_2}(X(\mathbb{R}^n), X(n\rho)),$$

and then passing under adjunction to

$$X(\mathbb{R}^n) \rightarrow \Omega^{n\sigma} X(n\rho),$$

before finally taking fixed points.

3. THE REAL SPIN BORDISM SPECTRUM

In this section, we construct $\text{MSpin}_{\mathbb{R}}^c$ as a commutative monoid in orthogonal C_2 -spectra. This mostly consists of equipping the constructions of Joachim in [7] with Real structures and verifying C_2 -equivariance wherever it is relevant. The extra data comes from the Real structures on the Clifford algebras $\text{Cl}_{p,q}$ mentioned in Section 1, combined with the constructions from Section 2.1.

Given an inner product space V , the Clifford algebra $\text{Cl}(V)$ is characterized by the following universal property. For any linear map $f: V \rightarrow A$ to an \mathbb{R} -algebra A such that² $f(v)^2 = \langle v, v \rangle$, there is a unique extension to an algebra homomorphism \tilde{f} in

$$\begin{array}{ccc} V & \xrightarrow{f} & A \\ \downarrow & \nearrow \tilde{f} & \\ \text{Cl}(V) & & \end{array}$$

Let $\text{Cl}_n = \text{Cl}(\mathbb{R}^n)$, where \mathbb{R}^n is given the standard inner product. The Clifford algebra is a $\mathbb{Z}/2$ -graded algebra by specifying that $V \subset \text{Cl}(V)$ lies in the odd subspace. Accordingly, the symbol \otimes refers to the graded tensor product. In particular, the symmetry isomorphism contains a sign,

$$V \otimes W \xrightarrow{\sim} W \otimes V, \quad v \otimes w \mapsto (-1)^{|v||w|} w \otimes v,$$

for v, w purely even or odd. With this symmetric monoidal structure, there is an isomorphism $\text{Cl}(V \oplus W) \cong \text{Cl}(V) \otimes \text{Cl}(W)$. For $V \in \mathcal{I}_{C_2}$, the Clifford algebra $\text{Cl}(V)$ inherits an induced C_2 -action by the universal property above, where C_2 acts by algebra maps. We extend this action to a Real structure on the complex Clifford algebra $\text{Cl}(V) := \text{Cl}(V) \otimes \mathbb{C}$ by complex conjugation on the \mathbb{C} -factor and Proposition 2.6. Note that this C_2 -action on $\text{Cl}(V)$ preserves the $\mathbb{Z}/2$ -grading (i.e. it is even), since it comes from an action on V . Lastly, fix the $*$ -structure on $\text{Cl}(V)$ to be the one generated by $v^* = v$, for $v \in V$. Let $\text{Cl}_n = \text{Cl}(\mathbb{R}^n)$, where \mathbb{R}^n is the trivial C_2 -representation, and $\text{Cl}_{p,q} = \text{Cl}(\mathbb{R}^{p,q})$, where $\mathbb{R}^{p,q} = \mathbb{R}^p \oplus \mathbb{R}^q \sigma$.

²see Remark 3.1

Remark 3.1 (Clifford sign conventions). Sign conventions for Clifford algebras differ significantly throughout the literature. The conventions we use are chosen to be easily comparable to the ones in [7] and [9], whose results we use in this paper. Forgetting the Real structure, the underlying complex C^* -algebra $\mathbb{Cl}(V)$ agrees with the C^* -algebra denoted \mathbb{Cl}_V in [7]. The Real C^* -algebra $\mathbb{Cl}_{p,q}$ is isomorphic to the Real C^* -algebra denoted $C_{p,q}$ in [9], as follows. Fix generators $v_1, \dots, v_p, w_1, \dots, w_q$ of $\mathbb{Cl}_{p,q}$ and $\varepsilon_1, \dots, \varepsilon_p, e_1, \dots, e_q$ of $C_{p,q}$, with

$$\begin{aligned} v_i^2 &= 1, & w_j^2 &= 1, & \varepsilon_i^2 &= 1, & e_j^2 &= -1, \\ \overline{v_i} &= v_i, & \overline{w_j} &= -w_j, & \overline{\varepsilon_i} &= \varepsilon_i, & \overline{e_j} &= e_j, \\ v_i^* &= v_i, & w_j^* &= w_j, & \varepsilon_i^* &= \varepsilon_i, & e_j^* &= -e_j. \end{aligned}$$

Then the homomorphism $\mathbb{Cl}_{p,q} \rightarrow C_{p,q}$ defined by $v_i \mapsto \varepsilon_i$ and $w_j \mapsto ie_j$ is an isomorphism of Real graded C^* -algebras.

Let $L^2(V)$ be the L^2 -completion of the pre-Hilbert space of continuous functions $V \rightarrow \mathbb{C}$ with compact support. Remark 2.5 defines a Real structure on $L^2(V)$, and Proposition 2.6 defines a Real structure on $\mathbb{Cl}(V) \otimes L^2(V)$. Here, we take $L^2(V)$ to be $\mathbb{Z}/2$ -graded by even and odd functions³. Then the induced Real structure on $\mathbb{Cl}(V) \otimes L^2(V)$ is also even. Let $\mathcal{B}(\mathbb{Cl}(V) \otimes L^2(V))$ be the Real space of bounded operators on $\mathbb{Cl}(V) \otimes L^2(V)$ equipped with the strong $*$ -topology and the Real structure from Remark 2.5.

Definition 3.2. Let $U_V \subset \mathcal{B}(\mathbb{Cl}(V) \otimes L^2(V))$ denote the graded unitary group of $\mathbb{Cl}(V) \otimes L^2(V)$, that is, the group consisting of unitary operators on $\mathbb{Cl}(V) \otimes L^2(V)$ that are either grading preserving or grading reversing, and let $\text{Pin}^c(V) \subset \mathbb{Cl}(V)$ be the group generated under multiplication by unit vectors $S(V) \subset V$ and unit complex numbers $U(1) \subset \mathbb{C}$.

Then the C_2 -actions on $\mathcal{B}(\mathbb{Cl}(V) \otimes L^2(V))$ and $\mathbb{Cl}(V)$ restrict to U_V and $\text{Pin}^c(V)$, respectively, making them into Real groups. Since both C_2 -actions are grading preserving, they restrict to Real structures on the respective even subgroups, $U_V^{\text{even}} \subset U_V$ and $\text{Spin}^c(V) \subset \text{Pin}^c(V)$. By Remark 2.5, the orthogonal group $O(V)$ is a Real group. Twisted conjugation of $\text{Pin}^c(V)$ on $\mathbb{Cl}(V)$ restricts to a homomorphism $\rho_V: \text{Pin}^c(V) \rightarrow O(V)$, defined by $\rho_V(g)(v) = (-1)^{|g|}gv g^{-1}$.

Proposition 3.3. The homomorphism ρ_V is a Real homomorphism.

Proof. Let $g \in \text{Pin}^c(V)$ and $v \in V$. Since the C_2 -action on $\text{Pin}^c(V)$ is even, we have

$$\overline{\rho_V(g)}(v) = \overline{\rho_V(g)(\overline{v})} = \overline{(-1)^{|g|}g\overline{v}g^{-1}} = (-1)^{|g|}\overline{g}v\overline{g}^{-1} = (-1)^{|\overline{g}|}\overline{g}v\overline{g}^{-1} = \rho_V(\overline{g})(v).$$

□

Remark 2.5 also says that V is a Real $O(V)$ -space, so Proposition 2.8 gives $L^2(V)$ the structure of a Real $O(V)$ -space. By Propositions 2.8 and 3.3, $L^2(V)$ becomes a Real $\text{Pin}^c(V)$ -space. Left multiplication also makes $\mathbb{Cl}(V)$ into a Real $\text{Pin}^c(V)$ -space, so Proposition 2.6 gives us a Real $\text{Pin}^c(V)$ -action on $\mathbb{Cl}(V) \otimes L^2(V)$. Since $\text{Pin}^c(V)$ acts on $\mathbb{Cl}(V) \otimes L^2(V)$ by unitary

³Here, *even* means $f(-x) = f(x)$, and *odd* means $f(-x) = -f(x)$.

operators, we get an induced Real homomorphism (Proposition 2.8) $i_V: \text{Pin}^c(V) \rightarrow U_V$. Unwinding the definitions yields the formula,

$$i_V(g)(v \otimes f) = (gv) \otimes (f \circ \rho_V(g)^{-1}),$$

as in [7]. This formula makes it clear that i_V is even, so it restricts to a Real homomorphism

$$i_V: \text{Spin}^c(V) \rightarrow U_V^{\text{even}}.$$

Thus, we get a Real $\text{Spin}^c(V)$ -action on U_V^{even} by left multiplication.

Proposition 3.4. If $V \neq 0$, then the $\text{Spin}^c(V)$ -space U_V^{even} is a model for the universal $\text{Spin}^c(V)$ -principal bundle, $E\text{Spin}^c(V)$.

Proof. Grading preserving unitary operators can be restricted to the even and odd subspaces to obtain $U_V^{\text{even}} \cong U((\text{Cl}(V) \otimes L^2(V))^{\text{even}}) \times U((\text{Cl}(V) \otimes L^2(V))^{\text{odd}})$, which is contractible by Kuiper's theorem [10]. The $\text{Spin}^c(V)$ -action on U_V^{even} is free, since i_V is injective. \square

We will also use $U_V^{\text{even}} / \text{Spin}^c(V)$ to model $B\text{Spin}^c(V)$. By Proposition 2.8, the associated bundle $\gamma_{\text{Spin}^c(V)} = U_V^{\text{even}} \times_{\text{Spin}^c(V)} V$ inherits the structure of a Real space.

Definition 3.5. For $V \in \mathcal{I}_{C_2}$, define the Real space

$$\text{MSpin}_{\mathbb{R}}^c(V) := \text{Thom}(\gamma_{\text{Spin}^c(V)})$$

to be the Thom space of the universal $\text{Spin}^c(V)$ -vector bundle, extending the Real structure on $\gamma_{\text{Spin}^c(V)}$ by fixing ∞ .

In order to give $\text{MSpin}_{\mathbb{R}}^c$ the structure of a C_2 - E_∞ -ring spectrum, we will realize it as a lax symmetric monoidal functor on \mathcal{I}_{C_2} . To do this, we first note that given $V, W \in \mathcal{I}_{C_2}$, we get a canonical (even) map $T_{V,W}: U_V \times U_W \rightarrow U_{V \oplus W}$, by taking the tensor product of operators, since

$$(2) \quad \begin{aligned} \text{Cl}(V \oplus W) \otimes L^2(V \oplus W) &\cong (\text{Cl}(V) \otimes \text{Cl}(W)) \otimes (L^2(V) \otimes L^2(W)) \\ &\cong (\text{Cl}(V) \otimes L^2(V)) \otimes (\text{Cl}(W) \otimes L^2(W)). \end{aligned}$$

Similarly, given an isomorphism $\iota: V \rightarrow W$, we get an isomorphism $\tilde{\iota}: \text{Cl}(V) \otimes L^2(V) \rightarrow \text{Cl}(W) \otimes L^2(W)$, which induces a homomorphism $U_\iota: U_V \rightarrow U_W$, defined by $U_\iota(A) = \tilde{\iota} A \tilde{\iota}^{-1}$. Furthermore, note that these assignments make $U_{(\cdot)}$ into a functor with respect to isomorphisms. Similarly, $T_{(\cdot)}$ is a natural transformation.

Proposition 3.6. Given $V, W \in \mathcal{I}_{C_2}$ of the same dimension, the map $U: \text{O}(V, W) \rightarrow \text{Hom}(U_V, U_W)$ defined above is C_2 -equivariant.

Proof. We have

$$\begin{aligned} \overline{U}_\iota(A)(v \otimes f) &= \overline{U}_\iota(\overline{A})(v \otimes f) \\ &= \overline{\tilde{\iota} A \tilde{\iota}^{-1}}(\overline{v \otimes f}) \\ &= \tilde{\iota} A \tilde{\iota}^{-1}(v \otimes f) \\ &= U_{\tilde{\iota}}(v \otimes f). \end{aligned}$$

\square

Now, let $V, W \in \mathcal{I}_{C_2}$, and $n = \dim W - \dim V \geq 0$. We define the structure maps

$$\mathcal{I}_{C_2}(V, W) \rightarrow \mathcal{T}_{C_2}(\mathrm{MSpin}_{\mathbb{R}}^c(V), \mathrm{MSpin}_{\mathbb{R}}^c(W)),$$

using the adjoint form

$$\mathcal{I}_{C_2}(V, W) \wedge \mathrm{MSpin}_{\mathbb{R}}^c(V) \xrightarrow{\sigma_{V,W}} \mathrm{MSpin}_{\mathbb{R}}^c(W),$$

by taking the Thomification of the vector bundle map

$$\begin{array}{ccc} (W - V) \times (\mathrm{U}_V^{\mathrm{even}} \times_{\mathrm{Spin}^c(V)} V) & \xrightarrow{\tilde{\sigma}_{V,W}} & \mathrm{U}_W^{\mathrm{even}} \times_{\mathrm{Spin}^c(W)} W \\ \downarrow & & \downarrow \\ \mathrm{O}(V, W) \times \mathrm{BSpin}^c(V) & \longrightarrow & \mathrm{BSpin}^c(W). \end{array}$$

Informally, $\tilde{\sigma}_{V,W}$ is the map

$$\begin{array}{ccc} ((\iota, w), [A, v]) \in (W - V) \times (\mathrm{U}_V^{\mathrm{even}} \times_{\mathrm{Spin}^c(V)} V) & & \\ \downarrow & & \downarrow \tilde{\sigma}_{V,W} \\ [T_{W-\iota V, V}(\mathrm{id}, A), w + v] \in \mathrm{U}_W^{\mathrm{even}} \times_{\mathrm{Spin}^c(W)} W. & & \end{array}$$

The issue with this formulation is that $W - \iota V$ is not an object of \mathcal{I}_{C_2} . In order to properly define $\tilde{\sigma}_{V,W}$, we make the C_2 -equivariant identification

$$\begin{array}{ccc} W - V & \xrightarrow{\sim} & \mathrm{O}(\mathbb{R}^n \oplus V, W) \times_{\mathrm{O}(n)} \mathbb{R}^n \\ \downarrow & & \downarrow \\ \mathrm{O}(V, W) & \xrightarrow{\sim} & \mathrm{O}(\mathbb{R}^n \oplus V, W) / \mathrm{O}(n). \end{array}$$

Definition 3.7. Define

$$\tilde{\sigma}_{V,W} : (\mathrm{O}(\mathbb{R}^n \oplus V, W) \times_{\mathrm{O}(n)} \mathbb{R}^n) \times (\mathrm{U}_V^{\mathrm{even}} \times_{\mathrm{Spin}^c(V)} V) \rightarrow \mathrm{U}_W^{\mathrm{even}} \times_{\mathrm{Spin}^c(W)} W,$$

by

$$\tilde{\sigma}_{V,W}([\iota, x], [A, v]) = [\mathrm{U}_\iota(T_{\mathbb{R}^n, V}(\mathrm{id}, A)), \iota(x + v)].$$

Proposition 3.8. The map $\tilde{\sigma}_{V,W}$ is C_2 -equivariant.

Proof. We have

$$\begin{aligned} \sigma_{V,W}(\overline{[\iota, x]}, \overline{[A, v]}) &= \sigma_{V,W}(\overline{[\iota, x]}, \overline{[A, v]}) = [\mathrm{U}_\tau(T_{\mathbb{R}^n, V}(\mathrm{id}, \overline{A})), \tau(\overline{x} + \overline{v})] \\ &= [\overline{(\mathrm{U}_\iota)(T_{\mathbb{R}^n, V}(\mathrm{id}, A))}, \overline{\iota(x + v)}] \\ &= \overline{[\mathrm{U}_\iota(T_{\mathbb{R}^n, V}(\mathrm{id}, A)), \iota(x + v)]} \\ &= \overline{[\mathrm{U}_\iota(T_{\mathbb{R}^n, V}(\mathrm{id}, A)), \iota(x) + \iota(v)]} \\ &= \overline{\sigma_{V,W}([\iota, x], [A, v])}. \end{aligned}$$

□

Definition 3.9. Define $\sigma_{V,W} : \mathcal{I}_{C_2}(V, W) \wedge \mathrm{MSpin}_{\mathbb{R}}^c(V) \rightarrow \mathrm{MSpin}_{\mathbb{R}}^c(W)$ as the Thomification of $\tilde{\sigma}_{V,W}$.

To define the unit, note that $U_0^{\text{even}} = *$, so $\text{MSpin}_{\mathbb{R}}^c(0) = S^0$.

Definition 3.10. Let $\eta_0^{\text{MSpin}_{\mathbb{R}}^c}: S^0 \rightarrow \text{MSpin}_{\mathbb{R}}^c(0)$ be the identity map on S^0 .

Definition 3.11. Let $V, W \in \mathcal{I}_{C_2}$. Define

$$\mu_{V,W}^{\text{MSpin}_{\mathbb{R}}^c}: \text{MSpin}_{\mathbb{R}}^c(V) \wedge \text{MSpin}_{\mathbb{R}}^c(W) \rightarrow \text{MSpin}_{\mathbb{R}}^c(V \oplus W),$$

as the Thomification of the map

$$(U_V^{\text{even}} \times_{\text{Spin}^c(V)} V) \times (U_W^{\text{even}} \times_{\text{Spin}^c(W)} W) \rightarrow U_{V \oplus W}^{\text{even}} \times_{\text{Spin}^c(V \oplus W)} (V \oplus W)$$

defined by $([A, v], [B, w]) \mapsto [T_{V,W}(A, B), v + w]$.

Note that $\mu_{V,W}^{\text{MSpin}_{\mathbb{R}}^c}$ is clearly C_2 -equivariant.

Proposition 3.12. The data $(\text{MSpin}_{\mathbb{R}}^c, \mu^{\text{MSpin}_{\mathbb{R}}^c}, \eta_0^{\text{MSpin}_{\mathbb{R}}^c})$ defines a lax symmetric monoidal enriched functor $\mathcal{I}_{C_2} \rightarrow \mathcal{T}_{C_2}$, and hence, a C_2 - E_{∞} -ring spectrum, $\text{MSpin}_{\mathbb{R}}^c \in \text{CAlg}(\text{Sp}^{C_2})$.

Proof. Verifying functoriality of $\text{MSpin}_{\mathbb{R}}^c$ is equivalent to showing that the following square commutes

$$\begin{array}{ccc} \mathcal{I}_{C_2}(V_2, V_3) \wedge \mathcal{I}_{C_2}(V_1, V_2) \wedge \text{MSpin}_{\mathbb{R}}^c(V_1) & \xrightarrow{\text{id} \wedge \sigma_{V_1, V_2}} & \mathcal{I}_{C_2}(V_2, V_3) \wedge \text{MSpin}_{\mathbb{R}}^c(V_2) \\ \downarrow & & \downarrow \sigma_{V_2, V_3} \\ \mathcal{I}_{C_2}(V_1, V_3) \wedge \text{MSpin}_{\mathbb{R}}^c(V_1) & \xrightarrow{\sigma_{V_1, V_3}} & \text{MSpin}_{\mathbb{R}}^c(V_3). \end{array}$$

This follows from the fact that both compositions are the Thomification of the same map of bundles

$$\begin{array}{ccc} (V_3 - V_2) \times (V_2 - V_1) \times (U_{V_1}^{\text{even}} \times_{\text{Spin}^c(V_1)} V_1) & \xrightarrow{\phi} & U_{V_3}^{\text{even}} \times_{\text{Spin}^c(V_3)} V_3 \\ \downarrow & & \downarrow \\ \text{O}(V_2, V_3) \times \text{O}(V_1, V_2) \times \text{BSpin}^c(V_1) & \longrightarrow & \text{BSpin}^c(V_3), \end{array}$$

where ϕ takes

$$((\iota_2, v_3), (\iota_1, v_2), [A, v_1]) \in (V_3 - V_2) \times (V_2 - V_1) \times (U_{V_1}^{\text{even}} \times_{\text{Spin}^c(V_1)} V_1)$$

to

$$[T_{V_3 - \iota_2 \iota_1 V_1, V_1}(\text{id}, A), v_3 + v_2 + v_1] \in U_{V_3}^{\text{even}} \times_{\text{Spin}^c(V_3)} V_3.$$

Naturality of $\mu^{\text{MSpin}_{\mathbb{R}}^c}$ is equivalent to commutativity of

$$\begin{array}{ccc} (\mathcal{I}_{C_2}(V_1, W_1) \wedge \text{MSpin}_{\mathbb{R}}^c(V_1)) \wedge (\mathcal{I}_{C_2}(V_2, W_2) \wedge \text{MSpin}_{\mathbb{R}}^c(V_2)) & \longrightarrow & \text{MSpin}_{\mathbb{R}}^c(W_1) \wedge \text{MSpin}_{\mathbb{R}}^c(W_2) \\ \downarrow & & \downarrow \\ (\mathcal{I}_{C_2}(V_1, W_1) \wedge \mathcal{I}_{C_2}(V_2, W_2)) \wedge (\text{MSpin}_{\mathbb{R}}^c(V_1) \wedge \text{MSpin}_{\mathbb{R}}^c(V_2)) & & \\ \downarrow & & \downarrow \\ \mathcal{I}_{C_2}(V_1 \oplus V_2, W_1 \oplus W_2) \wedge \text{MSpin}_{\mathbb{R}}^c(V_1 \oplus V_2) & \longrightarrow & \text{MSpin}_{\mathbb{R}}^c(W_1 \oplus W_2). \end{array}$$

Going around the top and then the right is the Thomification of the map

$$(W_1 - V_1) \times (W_2 - V_2) \times (U_{V_1}^{\text{even}} \times_{\text{Spin}^c(V_1)} V_1) \times (U_{V_2}^{\text{even}} \times_{\text{Spin}^c(V_2)} V_2) \rightarrow U_{W_1 \oplus W_2}^{\text{even}} \times_{\text{Spin}^c(W_1 \oplus W_2)} (W_1 \oplus W_2),$$

given by

$$((\iota_1, w_1), (\iota_2, w_2), [A, v_1], [B, v_2]) \mapsto [T_{W_1, W_2}(T_{W_1 - \iota_1 V_1}(\text{id}, A), T_{W_2 - \iota_2 V_2}(\text{id}, B)), w_1 + v_1 + w_2 + v_2].$$

While going around the left and then the bottom is the Thomification of the map

$$(W_1 - V_1) \times (W_2 - V_2) \times (U_{V_1}^{\text{even}} \times_{\text{Spin}^c(V_1)} V_1) \times (U_{V_2}^{\text{even}} \times_{\text{Spin}^c(V_2)} V_2) \rightarrow U_{W_1 \oplus W_2}^{\text{even}} \times_{\text{Spin}^c(W_1 \oplus W_2)} (W_1 \oplus W_2),$$

given by

$$((\iota_1, w_1), (\iota_2, w_2), [A, v_1], [B, v_2]) \mapsto [T_{W_1 \oplus W_2 - \iota_1 \oplus \iota_2 (V_1 \oplus V_2), V_1 \oplus V_2}(\text{id}, T_{V_1, V_2}(A, B)), w_1 + w_2 + v_1 + v_2].$$

These two maps are equal by naturality of T . The associativity, unitality, and symmetry conditions are all clear directly by the constructions. Thus, $\text{MSpin}_{\mathbb{R}}^c$ is a lax symmetric monoidal functor. \square

Proposition 3.13. The underlying spectrum of $\text{MSpin}_{\mathbb{R}}^c$ is MSpin^c .

Proof. We compare our construction to the one in Section 6 of [7]. Since the underlying spectrum is defined by restricting to trivial C_2 -representations we suppose that V is a trivial C_2 -representation. The following two maps of short exact sequences

$$\begin{array}{ccccccc} 1 & \longrightarrow & \text{Spin}^c(V) & \longrightarrow & \text{Pin}^c(V) & \longrightarrow & \mathbb{Z}/2 \longrightarrow 1 \\ & & \downarrow i_V & & \downarrow i_V & & \downarrow = \\ 1 & \longrightarrow & U_V^{\text{even}} & \longrightarrow & U_V & \longrightarrow & \mathbb{Z}/2 \longrightarrow 1 \end{array}$$

and

$$\begin{array}{ccccccc} 1 & \longrightarrow & U(1) & \longrightarrow & \text{Pin}^c(V) & \longrightarrow & O(V) \longrightarrow 1 \\ & & \downarrow = & & \downarrow i_V & & \downarrow j_V \\ 1 & \longrightarrow & U(1) & \longrightarrow & U_V & \longrightarrow & \text{PU}_V \longrightarrow 1 \end{array}$$

allow us to identify $\gamma_{\text{Spin}^c(V)} = U_V^{\text{even}} \times_{\text{Spin}^c(V)} V \cong \text{PU}_V \times_{O(V)} V$, which shows that the underlying space $\text{MSpin}_{\mathbb{R}}^c(V)$ is the same as the one denoted by $\mathbb{M}\text{Spin}_V^c$ in [7]. Using this identification, we get a map

$$k_V: O(V) \times_{O(V)} V \xrightarrow{j_V \times_{O(V)} \text{id}_V} \text{PU}_V \times_{O(V)} V \xrightarrow{\sim} U_V^{\text{even}} \times_{\text{Spin}^c(V)} V.$$

The map $\tilde{\eta}_V: S^V \rightarrow \text{MSpin}_{\mathbb{R}}^c(V)$ obtained by taking the Thom construction of k_V is the V -component of the lax monoidal natural transformation $S \rightarrow \text{MSpin}^c$ playing the role of the unit of the ring spectrum defined in [7]. In our model, the V -component of the ring spectrum unit $\eta_V^{\text{MSpin}_{\mathbb{R}}^c}: S^V \rightarrow \text{MSpin}_{\mathbb{R}}^c(V)$ is obtained by combining the structure maps of

Definition 3.9 with the functor unit $\eta^{\text{MSpin}_{\mathbb{R}}^c}: S^0 \rightarrow \text{MSpin}_{\mathbb{R}}^c(0)$ from Definition 3.10. Specifically, $\eta_V^{\text{MSpin}_{\mathbb{R}}^c}$ is the Thomification of $\tilde{\sigma}_{0,V}: \text{O}(\mathbb{R}^n, V) \times_{\text{O}(n)} \mathbb{R}^n \rightarrow \text{U}_V^{\text{even}} \times_{\text{Spin}^c(V)} V$, defined above by $\tilde{\sigma}_{0,V}([\iota, x]) = [\text{id}, \iota(x)]$. Fixing a specific isomorphism $\varphi: \mathbb{R}^n \xrightarrow{\sim} V$, the diagram

$$\begin{array}{ccc} \text{O}(\mathbb{R}^n, V) \times_{\text{O}(n)} \mathbb{R}^n & \xrightarrow{\tilde{\sigma}_{0,V}} & \text{U}_V^{\text{even}} \times_{\text{Spin}^c(V)} V \\ \tilde{\varphi} \downarrow & & \uparrow \cong \\ \text{O}(V) \times_{\text{O}(V)} V & \xrightarrow{j_V \times_{\text{O}(V)} \text{id}_V} & \text{PU}_V \times_{\text{O}(V)} V, \end{array}$$

where $\tilde{\varphi}([\iota, x]) = [\iota\varphi^{-1}, \varphi(x)]$, commutes, since

$$(j_V \times_{\text{O}(V)} \text{id}_V)([\iota\varphi^{-1}, \varphi(x)]) = [j_V(\iota\varphi^{-1}), \varphi(x)] = [\text{id}, (\iota\varphi^{-1})\varphi(x)] = [\text{id}, \iota(x)].$$

Thus, $\eta_V^{\text{MSpin}_{\mathbb{R}}^c} = \tilde{\eta}_V$. It is also evident that the multiplication maps are the same as those in [7]. Thus, $(\text{MSpin}_{\mathbb{R}}^c)^e = \text{MSpin}^c$. \square

Remark 3.14. Another way to organize the data of a commutative orthogonal ring spectrum is via the formalism of \mathcal{I} -FSP's [13]. This is the language that Joachim [7] uses; however, it is not hard to see that our constructions agree with his when the models are compared.

4. THE FIXED POINTS OF $\text{MSpin}_{\mathbb{R}}^c$

Recall that the inclusion $\text{Spin}(n) \hookrightarrow \text{Spin}^c(n)$ factors through an isomorphism with the C_2 -fixed points of $\text{Spin}^c(n) = \text{Spin}^c(n, 0)$. In light of this, one might hope that the usual map $\text{MSpin} \xrightarrow{u} \text{MSpin}^c$ witnesses MSpin as the C_2 -(homotopy) fixed points of $\text{MSpin}_{\mathbb{R}}^c$.

$$\begin{array}{ccc} \text{Spin}(n) & \hookrightarrow & \text{Spin}^c(n) \\ \cong \searrow & & \nearrow \\ & & \text{Spin}^c(n)^{C_2} \end{array} \quad \rightsquigarrow \quad \begin{array}{ccc} \text{MSpin} & \xrightarrow{u} & \text{MSpin}^c \\ \tilde{u} \dashrightarrow & & \nearrow \\ & & (\text{MSpin}_{\mathbb{R}}^c)^{C_2} \end{array}$$

In this section, we show that u does factor through $\tilde{u}: \text{MSpin} \rightarrow (\text{MSpin}_{\mathbb{R}}^c)^{C_2}$ as above, but that \tilde{u} cannot be an equivalence (Corollary 4.7).

Lemma 4.1. For $V \in \mathcal{I}_{C_2}$ with $\dim V \geq 1$, the space $(\text{U}_V^{\text{even}})^{C_2}$ is contractible.

Proof. Restricting a grading preserving operator to the even and odd subspaces yields,

$$(\text{U}_V^{\text{even}})^{C_2} \cong \text{U}(\mathcal{H}_V^0)^{C_2} \times \text{U}(\mathcal{H}_V^1)^{C_2},$$

where $\mathcal{H}_V^0, \mathcal{H}_V^1$ are the even and odd parts of $\text{Cl}(V) \otimes L^2(V)$, respectively, and $\text{U}(\mathcal{H})^{C_2}$ is the group of Real unitary operators on the Real Hilbert space \mathcal{H} . Since $\mathcal{H} \cong \mathcal{H}^{C_2} \otimes \mathbb{C}$ (Proposition 2.7), $\text{U}(\mathcal{H})^{C_2}$ can be identified with the orthogonal group $\text{O}(\mathcal{H}^{C_2})$ of the real Hilbert space \mathcal{H}^{C_2} . By the real version of Kuiper's theorem [10], $\text{O}(\mathcal{H}^{C_2})$ is contractible, which implies $(\text{U}_V^{\text{even}})^{C_2}$ is contractible. \square

Definition 4.2. Let $\text{MSpin}(n) := \text{Thom}((\text{U}_{\mathbb{R}^n}^{\text{even}})^{C_2} \times_{\text{Spin}(n)} \mathbb{R}^n)$, where $\text{Spin}(n)$ is identified with $\text{Spin}^c(n)^{C_2}$.

Proposition 4.3. The spaces $\text{MSpin}(n)$ form an orthogonal spectrum, with the usual structure maps, equivalent to MSpin .

Proof. By Lemma 4.1, the space $(\text{U}_{\mathbb{R}^n}^{\text{even}})^{C_2}$ is contractible with a free $\text{Spin}(n)$ -action, and thus is a model for $\text{ESpin}(n)$. From there, this is a standard Thom spectrum construction. For example, one can follow the same steps taken in Section 3 with the obvious modifications. \square

Definition 4.4. Let $\tilde{u}(n): \text{MSpin}(n) \rightarrow \text{MSpin}_{\mathbb{R}}^c(\mathbb{R}^n)$ be the Thomification of the composite

$$(\text{U}_{\mathbb{R}^n}^{\text{even}})^{C_2} \times_{\text{Spin}(n)} \mathbb{R}^n \rightarrow \text{U}_{\mathbb{R}^n}^{\text{even}} \times_{\text{Spin}(n)} \mathbb{R}^n \rightarrow \text{U}_{\mathbb{R}^n}^{\text{even}} \times_{\text{Spin}^c(n)} \mathbb{R}^n.$$

induced by the inclusions $(\text{U}_{\mathbb{R}^n}^{\text{even}})^{C_2} \hookrightarrow \text{U}_{\mathbb{R}^n}^{\text{even}}$ and $\text{Spin}(n) \hookrightarrow \text{Spin}^c(n)$.

By construction, the image of $\tilde{u}(n)$ is contained in the C_2 -fixed points of $\text{MSpin}_{\mathbb{R}}^c(\mathbb{R}^n)$.

Proposition 4.5. The maps $\tilde{u}(n)$ define a map of commutative orthogonal ring spectra

$$\tilde{u}: \text{MSpin} \rightarrow (\text{MSpin}_{\mathbb{R}}^c)^{C_2}.$$

Proof. The maps $\tilde{u}(n)$ define a symmetric monoidal natural transformation between commutative monoids in Sp from MSpin to the categorical fixed points $F^{C_2} \text{MSpin}_{\mathbb{R}}^c$. By Proposition 2.18, this induces a symmetric monoidal natural transformation to $(\text{MSpin}_{\mathbb{R}}^c)^{C_2}$. \square

We now show that \tilde{u} is not an equivalence by proving Theorem 1.3.

4.1. Proof of Theorem 1.3. Suppose E is a genuine C_2 -spectrum satisfying conditions (1)-(4) of Theorem 1.3. Under the isomorphisms in (1) and (2), the homotopy Mackey functor $\pi_* E$ consists of the data

$$\text{MSpin}_* \begin{array}{c} \xrightarrow{\text{res}} \\ \xleftarrow{\text{tr}} \end{array} \text{MSpin}_*^c \begin{array}{c} \xrightarrow{\overline{(\quad)}} \\ \xleftarrow{\overline{(\quad)}} \end{array} \overline{(\quad)},$$

where $\overline{(\quad)}$ is the C_2 -action in (3). Given any class $\alpha \in \text{MSpin}_*^c$, the Mackey functor structure (Definition 2.11) implies that the equation,

$$\alpha + \overline{\alpha} = \text{res}(\text{tr}(\alpha)),$$

holds. Then the commutative diagram in (4) implies that for any $\alpha \in \text{MSpin}_*^c$, the oriented bordism class $u_*^c(\alpha + \overline{\alpha})$ must have a spin manifold as a representative. Thus, it is sufficient to show there is some $\alpha \in \text{MSpin}_*^c$ for which $u_*^c(\alpha + \overline{\alpha})$ has no spin representative. Recall the \hat{A} -genus (see e.g. [11]),

$$\begin{array}{ccc} \text{MSpin}_* & \xrightarrow{\hat{A}} & \mathbb{Z} \\ u_* \downarrow & & \downarrow \\ \text{MSO}_* & \xrightarrow{\hat{A}} & \mathbb{Q}, \end{array}$$

and consider the class

$$\gamma = [\mathbb{C}\mathbb{P}^2] \in \text{MSpin}_4^c.$$

We know that $\hat{A}(u_*^c(\gamma)) = -1/8$ ([11]). By the discussion above, the class $\gamma + \overline{\gamma}$ must be in the image of MSpin_4 . Then $\hat{A}(u_*^c(\gamma + \overline{\gamma})) =: n \in \mathbb{Z}$, so

$$\hat{A}(u_*^c(\overline{\gamma})) = \frac{1}{8} + n.$$

Now consider the class

$$\alpha = \gamma^2 \in \mathrm{MSpin}_8^c.$$

By multiplicativity of both the C_2 -action and \hat{A} , we see that

$$\hat{A}(u_*^c(\alpha + \bar{\alpha})) = \frac{1}{64} + \frac{1}{64} + \frac{2n}{8} + n^2 = \frac{8n+1}{32} + n^2,$$

which is not an integer, since 32 never divides $8n+1$. Thus, $\alpha + \bar{\alpha}$ is not in the image of $\mathrm{res}: \mathrm{MSpin}_* \rightarrow \mathrm{MSpin}_*^c$, which is a contradiction. This completes the proof of Theorem 1.3.

Remark 4.6. As can be seen from the proof, condition (4) in Theorem 1.3 can be loosened to just requiring that the map $\mathrm{res}: \mathrm{MSpin}_* \rightarrow \mathrm{MSpin}_*^c$ preserves the \hat{A} -genus.

Corollary 4.7. If there is an equivalence $\mathrm{MSpin} \xrightarrow{\sim} (\mathrm{MSpin}_{\mathbb{R}}^c)^{C_2}$, then the composite,

$$\mathrm{MSpin} \xrightarrow{\sim} (\mathrm{MSpin}_{\mathbb{R}}^c)^{C_2} \rightarrow \mathrm{MSpin}^c,$$

does not take a spin manifold to its underlying spin^c manifold on homotopy groups. It follows that \tilde{u} of Proposition 4.5 is not an equivalence.

Remark 4.8. Recall that the geometric fixed points of Real bordism is unoriented bordism, $\Phi^{C_2} \mathrm{MU}_{\mathbb{R}} \simeq \mathrm{MO}$. In analogy, one might expect an equivalence between $\Phi^{C_2} \mathrm{MSpin}_{\mathbb{R}}^c$ and MSpin , but Proposition 5.1 (below) implies that such an equivalence would induce a ring map $\mathrm{MO} \rightarrow \mathrm{MSpin}$ (by lax monoidality of Φ^{C_2}), which doesn't exist. Similarly, $\mathrm{MSpin} \not\cong \Phi^{C_2}((\mathrm{MSpin}_{\mathbb{R}}^c)^h)$.

5. A REAL ORIENTATION OF REAL SPIN BORDISM

In this section, we show that the complex orientation of MSpin^c can be refined to a Real orientation of $\mathrm{MSpin}_{\mathbb{R}}^c$.

Proposition 5.1. There is a map of C_2 - E_{∞} -ring spectra $\mathrm{MU}_{\mathbb{R}} \rightarrow \mathrm{MSpin}_{\mathbb{R}}^c$, where $\mathrm{MU}_{\mathbb{R}}$ denotes the Real bordism spectrum. Thus, $\mathrm{MSpin}_{\mathbb{R}}^c$ is Real oriented.

The next two subsections are dedicated to proving Proposition 5.1.

5.1. Real spectra. We begin by briefly recalling another model for C_2 -spectra which will be convenient for constructing the Real orientation. This model is not needed anywhere else in this paper. For all details, we refer the reader to the appendix of Hill, Hopkins, and Ravenel [5, Section B.12] or Schwede [15, Chapter 2]. Since a C_2 -spectrum is determined by its values on multiples of a regular representation, we can instead *define* C_2 -spectra by indexing over this smaller sequence of representations. This idea is made precise as follows. Define a \mathcal{T}^{C_2} -enriched category $\mathcal{I}_{\mathbb{R}}$ with objects $\mathbb{R}^n =: n$, and with the C_2 -mapping space $\mathcal{I}_{\mathbb{R}}(n, m)$ defined as the Thom space of the difference bundle associated to the complexifications,

$$\mathcal{I}_{\mathbb{R}}(n, m) = \mathrm{Thom}(U(\mathbb{C}^n, \mathbb{C}^m); \mathbb{C}^m - \mathbb{C}^n),$$

with C_2 -action given by complex conjugation.

Definition 5.2. The category of *Real spectra*, $\mathrm{Sp}_{\mathbb{R}}$, is the category of \mathcal{T}^{C_2} -enriched functors

$$\mathcal{I}_{\mathbb{R}} \rightarrow \mathcal{T}_{C_2}.$$

Given $X \in \mathrm{Sp}_{\mathbb{R}}$, $k \in \mathbb{Z}$, and a subgroup $H \subset C_2$, define the homotopy group

$$\pi_k^H(X) = \mathrm{colim}_n \pi_{k+n\rho}^H(X(n)).$$

Define a weak equivalence between Real spectra to be a map which induces an isomorphism on homotopy groups. This homotopical category structure can be refined to a model category [6]. Define a \mathcal{T}^{C_2} -enriched functor $i: \mathcal{I}_{\mathbb{R}} \rightarrow \mathcal{I}_{C_2}$ by

$$n \mapsto n\rho = \mathbb{R}^n \otimes \rho,$$

and the natural inclusion on mapping C_2 -spaces.

Proposition 5.3. [5, Proposition B.226] There is a Quillen equivalence,

$$i_! : \mathrm{Sp}_{\mathbb{R}} \rightleftarrows \mathrm{Sp}^{C_2} : i^*,$$

given by left Kan extension along i and precomposition by i respectively.

The left adjoint is also strongly symmetric monoidal, so we also get a Quillen equivalence between categories of commutative monoids [5, Proposition B.248],

$$i_! : \mathrm{CAlg}(\mathrm{Sp}_{\mathbb{R}}) \rightleftarrows \mathrm{CAlg}(\mathrm{Sp}^{C_2}) : i^*.$$

Because of this Quillen equivalence, C_2 - E_{∞} -ring maps can be modelled by lax symmetric monoidal natural transformations between lax symmetric monoidal functors out of $\mathcal{I}_{\mathbb{R}}$.

5.1.1. $\mathrm{MU}_{\mathbb{R}}$ and Real orientations. As is discussed in [5, Section B.12] and [15, Chapter 2], the most natural way to construct $\mathrm{MU}_{\mathbb{R}}$ as a C_2 - E_{∞} -ring spectrum is by constructing it as a commutative monoid object in Real spectra and then applying the Quillen equivalence above. This can be done by equipping the spaces $\mathrm{MU}(n)$ with the complex conjugation actions, which defines an object $\mathcal{M}\mathcal{U}_{\mathbb{R}} \in \mathrm{CAlg}(\mathrm{Sp}_{\mathbb{R}})$. As in [5, Definition B.251], $\mathrm{MU}_{\mathbb{R}}$ is then defined as a commutative orthogonal C_2 -ring spectrum by applying $i_!$ to a cofibrant replacement of $\mathcal{M}\mathcal{U}_{\mathbb{R}}$ in $\mathrm{CAlg}(\mathrm{Sp}_{\mathbb{R}})$.

Remark 5.4. For $X \in \mathrm{CAlg}(\mathrm{Sp}^{C_2})$, Real E_{∞} -orientations of X are represented by commutative ring maps $\mathrm{MU}_{\mathbb{R}} \rightarrow X$. By the Quillen equivalence in Proposition 5.3, it is equivalent to consider lax symmetric monoidal natural transformations $\mathcal{M}\mathcal{U}_{\mathbb{R}} \rightarrow i^*(X)$ in $\mathrm{Sp}_{\mathbb{R}}$, where $i^*(X)$ is the Real spectrum obtained from X by restricting to regular representations.

5.2. $\mathrm{MSpin}_{\mathbb{R}}^c$ is Real oriented. Recall that the homomorphism $\iota \times \det: \mathrm{U}(n) \rightarrow \mathrm{SO}(2n) \times \mathrm{U}(1)$ lifts to a homomorphism $\varphi(n): \mathrm{U}(n) \rightarrow \mathrm{Spin}^c(2n)$. Since ι is injective, $\varphi(n)$ is also injective. Furthermore, observe that $\varphi(n)$ is actually a Real homomorphism of Real groups $\varphi(n): \mathrm{U}(n) \rightarrow \mathrm{Spin}^c(n\rho)$. By Proposition 3.4, the $\mathrm{U}(n)$ -space $\mathrm{U}_{n\rho}^{\mathrm{even}}$ is a model for $\mathrm{EU}(n)$ as a Real $\mathrm{U}(n)$ -space. The associated bundle $\gamma_{\mathrm{U}(n)} = \mathrm{U}_{n\rho}^{\mathrm{even}} \times_{\mathrm{U}(n)} \mathbb{C}^n$ then also inherits a Real structure and the identity map on $\mathrm{U}_{n\rho}^{\mathrm{even}} \times \mathbb{C}^n = \mathrm{U}_{n\rho}^{\mathrm{even}} \times (n\rho)$ descends to a Real map of Real vector bundles $\gamma_{\varphi(n)}: \gamma_{\mathrm{U}(n)} \rightarrow \gamma_{\mathrm{Spin}^c(n\rho)}$.

Definition 5.5. Define $\mathcal{M}\mathcal{U}'_{\mathbb{R}}(n)$ to be the Thom space of $\gamma_{\mathrm{U}(n)}$, with its induced Real structure, and let $M\varphi(n): \mathcal{M}\mathcal{U}'_{\mathbb{R}}(n) \rightarrow \mathrm{MSpin}_{\mathbb{R}}^c(n\rho)$ be the Thomification of $\gamma_{\varphi(n)}$.

Proposition 5.6. The Real spaces $\mathcal{MU}'_{\mathbb{R}}(n)$ (with the usual structure maps) define a lax symmetric monoidal enriched functor $\mathcal{MU}'_{\mathbb{R}}: \mathcal{I}_{\mathbb{R}} \rightarrow \mathcal{T}_{C_2}$, and thus, $\mathcal{MU}'_{\mathbb{R}} \in \text{CAlg}(\text{Sp}_{\mathbb{R}})$.

Proof. This is a standard Thom spectrum construction, which is analogous to the construction in Section 3. \square

Proposition 5.7. The maps $M\varphi(n)$ define a map of commutative monoids

$$M\varphi: \mathcal{MU}'_{\mathbb{R}} \rightarrow i^* \text{MSpin}_{\mathbb{R}}^c$$

in $\text{Sp}_{\mathbb{R}}$.

Proof. This follows from another standard Thom spectrum argument, using the fact that the squares,

$$\begin{array}{ccc} \text{U}(n) & \longrightarrow & \text{Spin}^c(n\rho) \\ \downarrow & & \downarrow \\ \text{U}(n+1) & \longrightarrow & \text{Spin}^c((n+1)\rho) \end{array} \quad \text{and} \quad \begin{array}{ccc} \text{U}(n) \times \text{U}(m) & \longrightarrow & \text{Spin}^c(n\rho) \times \text{Spin}^c(m\rho) \\ \downarrow & & \downarrow \\ \text{U}(n+m) & \longrightarrow & \text{Spin}^c((n+m)\rho), \end{array}$$

commute. \square

The last step we need in order to show that $\text{MSpin}_{\mathbb{R}}^c$ is Real oriented is to show that $\mathcal{MU}'_{\mathbb{R}}$ is equivalent to $\mathcal{MU}_{\mathbb{R}}$. By construction, we know that the space $\mathcal{MU}'_{\mathbb{R}}(n)$ is equivalent to $\text{MU}(n)$. However, it is not obvious that the C_2 -action we constructed on $\mathcal{MU}'_{\mathbb{R}}(n)$ is equivalent to the complex conjugation action on $\text{MU}(n)$. The rest of this section is devoted to showing that our construction does model MU with complex conjugation.

Lemma 5.8. If X and Y are (right and left, respectively) Real $\text{U}(n)$ -spaces, with $\text{U}(n)$ acting freely on X , then the C_2 -fixed points of $X \times_{\text{U}(n)} Y$ is given by the space

$$(X \times_{\text{U}(n)} Y)^{C_2} \cong X^{C_2} \times_{\text{O}(n)} Y^{C_2}.$$

Proof. Define the map

$$\phi: X^{C_2} \times_{\text{O}(n)} Y^{C_2} \rightarrow (X \times_{\text{U}(n)} Y)^{C_2},$$

to be the one induced by the inclusions $X^{C_2} \hookrightarrow X$ and $Y^{C_2} \hookrightarrow Y$. That is, we have that $\phi([x, y]_{\text{O}(n)}) = [x, y]_{\text{U}(n)}$. Here the notation $[x, y]_G$ is referring to the class represented by (x, y) in $X \times_G Y$. First, we show injectivity. Suppose $\phi([x_0, y_0]_{\text{O}(n)}) = \phi([x_1, y_1]_{\text{O}(n)})$. Then there exists $U \in \text{U}(n)$, such that

$$(x_0, y_0) = (x_1 U^*, U y_1).$$

But since x_0, x_1 are fixed by C_2 we get

$$x_1 U^* = x_0 = \overline{x_0}, = \overline{x_1 \overline{U}^*} = x_1 \overline{U}^*.$$

Thus, since G acts freely on X , we must have that $U = \overline{U}$. So $U \in \text{O}(n)$ and

$$[x_0, y_0]_{\text{O}(n)} = [x_1, y_1]_{\text{O}(n)}.$$

Next, we show surjectivity. Let $[x, y]_{U(n)} \in (X \times_{U(n)} Y)^{C_2}$. Then there exists $U \in U(n)$ such that

$$(\bar{x}, \bar{y}) = (xU^*, Uy).$$

Observe that from $\bar{x} = xU^*$, we obtain

$$x = \overline{xU^*} = \bar{x}\overline{U^*} = \bar{x}U^T,$$

by taking conjugates. However, if we instead acted by U on both sides of $\bar{x} = xU^*$, we would obtain $\bar{x}U = x$. By freeness of the action of $U(n)$ on X , we must have that $U = U^T$, so U is symmetric. Then we may choose an $S \in U(n)$ such that $S^2 = U$ and $S^T = S$. This can be done by diagonalizing U and choosing a square root for each eigenvalue of U [17, Theorem 5.12]. Now consider the element

$$(w, z) = (xS^*, Sy).$$

By construction, we have that

$$[w, z]_{U(n)} = [x, y]_{U(n)}.$$

Observe that both w and z are fixed points, since

$$\bar{w} = \overline{xS^*} = \bar{x}\overline{S^*} = \bar{x}S = xU^*S = xS^* = w,$$

and

$$\bar{z} = \overline{Sy} = \bar{S}\bar{y} = S^*\bar{y} = S^*Uy = Sy = z.$$

It follows that $[x, y]_{U(n)}$ is in the image of ϕ . Continuity of the inverse is then immediate from the fact that open sets in both the source and the target can be represented by the set of orbits of an open set in $X^{C_2} \times Y^{C_2}$. \square

Proposition 5.9. There is a weak equivalence in $\text{CAlg}(\text{Sp}_{\mathbb{R}})$,

$$\mathcal{MU}'_{\mathbb{R}} \simeq \mathcal{MU}_{\mathbb{R}}.$$

Proof. By construction, we know that we have a nonequivariant homotopy equivalence

$$\mathcal{MU}'_{\mathbb{R}}(n) \simeq \text{MU}(n) = \mathcal{MU}_{\mathbb{R}}(n),$$

but we need to verify that the C_2 -action that we have defined on $\mathcal{MU}'_{\mathbb{R}}(n)$ is equivalent to complex conjugation. Recall that complex conjugation on the universal complex vector bundle $\gamma_n = \text{EU}(n) \times_{U(n)} \mathbb{C}^n$ over $\text{BU}(n)$ defines a universal n -dimensional Real vector bundle [4]. Then the Real bundle $U_{n\rho}^{\text{even}} \times_{U(n)} \mathbb{C}^n \rightarrow U_{n\rho}^{\text{even}} / U(n)$, induces a map of Real vector bundles which fits into a pullback diagram of C_2 -spaces

$$(3) \quad \begin{array}{ccc} U_{n\rho}^{\text{even}} \times_{U(n)} \mathbb{C}^n & \longrightarrow & \text{EU}(n) \times_{U(n)} \mathbb{C}^n \\ \downarrow & & \downarrow \\ U_{n\rho}^{\text{even}} / U(n) & \longrightarrow & \text{BU}(n). \end{array}$$

We define

$$\theta: \mathcal{MU}'_{\mathbb{R}} \rightarrow \mathcal{MU}_{\mathbb{R}},$$

by letting

$$\theta_n: \mathcal{MU}'_{\mathbb{R}}(n) \rightarrow \mathcal{MU}_{\mathbb{R}}(n),$$

be the C_2 -equivariant map given by Thomification of the map of total spaces,

$$\beta_n: \mathbb{U}_{n\rho}^{\text{even}} \times_{\mathbb{U}(n)} \mathbb{C}^n \rightarrow \mathbb{E}\mathbb{U}(n) \times_{\mathbb{U}(n)} \mathbb{C}^n,$$

in our map of Real vector bundles. We will verify that θ_n is an equivalence of C_2 -spaces. To do this, we first observe that θ_n is an equivalence on underlying spaces. Since β_n fits into the pullback diagram of complex vector bundles (3), the map on base spaces must be a homotopy equivalence, since this pullback square classifies that $\mathbb{U}_{n\rho}^{\text{even}} \times_{\mathbb{U}(n)} \mathbb{C}^n$ is a model for the universal n -dimensional complex vector bundle. Thus, the map on Thom spaces must also be an equivalence. On fixed points, we use Lemma 5.8 (once on the total space and once on the base space) to observe that the real vector bundle

$$(\mathbb{U}_{n\rho}^{\text{even}} \times_{\mathbb{U}(n)} \mathbb{C}^n)^{C_2} \rightarrow (\mathbb{U}_{n\rho}^{\text{even}} / \mathbb{U}(n))^{C_2},$$

can be identified with

$$(\mathbb{U}_{n\rho}^{\text{even}})^{C_2} \times_{\mathbb{O}(n)} \mathbb{R}^n \rightarrow (\mathbb{U}_{n\rho}^{\text{even}})^{C_2} / \mathbb{O}(n).$$

Then using the fact that taking fixed points preserves pullback squares, we see that $\beta_n^{C_2}$ fits into a pullback square giving a map of real vector bundles

$$\begin{array}{ccc} (\mathbb{U}_{n\rho}^{\text{even}})^{C_2} \times_{\mathbb{O}(n)} \mathbb{R}^n & \longrightarrow & \mathbb{E}\mathbb{O}(n) \times_{\mathbb{O}(n)} \mathbb{R}^n \\ \downarrow & & \downarrow \\ (\mathbb{U}_{n\rho}^{\text{even}})^{C_2} / \mathbb{O}(n) & \longrightarrow & \mathbb{B}\mathbb{O}(n). \end{array}$$

By Lemma 4.1, $(\mathbb{U}_{n\rho}^{\text{even}})^{C_2}$ is a contractible space with a free $\mathbb{O}(n)$ -action, so this pullback square classifies the universal real vector bundle on the model of $\mathbb{B}\mathbb{O}(n)$ given by $(\mathbb{U}_{n\rho}^{\text{even}})^{C_2} / \mathbb{O}(n)$. Thus, the map on Thom spaces, $\theta_n^{C_2}$, is an equivalence. We conclude that θ_n is an equivalence of C_2 -spaces for every n . Monoidality and naturality of this construction are clear, so the maps θ_n give an equivalence

$$\theta: \mathcal{M}\mathcal{U}'_{\mathbb{R}} \simeq \mathcal{M}\mathcal{U}_{\mathbb{R}}$$

in $\text{CAlg}(\text{Sp}_{\mathbb{R}})$. □

This completes the proof of Proposition 5.1.

6. A REAL SPIN ORIENTATION OF REAL K-THEORY

In this section, we refine the Atiyah–Bott–Shapiro spin^c orientation of KU ([2], [7]) to a C_2 - E_{∞} -map from $\text{MSpin}_{\mathbb{R}}^c$ to Atiyah’s Real K-theory. By Proposition 4.5, this recovers the Atiyah–Bott–Shapiro spin orientation of KO as an E_{∞} -map as well. We start by adapting the construction of KU in [7] (for trivial G) to a model for Real K-theory as a C_2 - E_{∞} -ring spectrum, $\text{KU}_{\mathbb{R}}$. Due to the adaptability of Kasparov’s KK -theory to the setting of Real C^* -algebras, our modifications consist of equipping all relevant C^* -algebras with appropriate Real structures. Following [7], we implicitly use the formalism of \mathcal{I}_{C_2} -FSP’s [13] for ease of comparison of our constructions.

Let $\mathfrak{s} = C_0(\mathbb{R})$ be the $\mathbb{Z}/2$ -graded C^* -algebra of continuous functions $\mathbb{R} \rightarrow \mathbb{C}$ vanishing at infinity, graded by even and odd functions. We equip \mathfrak{s} with a Real structure defined by complex conjugation on values.

Proposition 6.1. The C^* -algebra \mathfrak{s} is generated by the functions $a, b \in \mathfrak{s}$ defined by

$$a(t) = \frac{1}{1+t^2} \quad \text{and} \quad b(t) = \frac{t}{1+t^2}.$$

Given a $\mathbb{Z}/2$ -graded Hilbert space \mathcal{H} , let $\mathcal{K}(\mathcal{H})$ denote the $\mathbb{Z}/2$ -graded C^* -algebra of compact operators on \mathcal{H} . For $V \in \mathcal{I}_{C_2}$, let $\mathcal{K}_V = \mathcal{K}(L^2(V))$. Then Remark 2.5 gives \mathcal{K}_V a Real structure, and Proposition 2.6 gives $\text{Cl}(V) \otimes \mathcal{K}_V$ a Real structure. If A, B are any Real $\mathbb{Z}/2$ -graded C^* -algebras, then Remark 2.5 also makes the space of even $*$ -homomorphisms $C_{\text{gr}}^*(A, B)$ into a Real space.

Definition 6.2. For $V \in \mathcal{I}_{C_2}$, define the Real space

$$\text{KU}_{\mathbb{R}}(V) := C_{\text{gr}}^*(\mathfrak{s}, \text{Cl}(V) \otimes \mathcal{K}_V).$$

Remark 6.3. For Hilbert spaces $\mathcal{H}_1, \mathcal{H}_2$, there is a canonical isomorphism

$$(4) \quad \mathcal{K}(\mathcal{H}_1) \otimes \mathcal{K}(\mathcal{H}_2) \cong \mathcal{K}(\mathcal{H}_1 \otimes \mathcal{H}_2).$$

Combining this with the isomorphism $\text{Cl}(V) \xrightarrow{\sim} \mathcal{K}(\text{Cl}(V))$ given by Clifford multiplication, yields a canonical isomorphism

$$(5) \quad \mathcal{K}(\text{Cl}(V) \otimes L^2(V)) \cong \text{Cl}(V) \otimes \mathcal{K}_V.$$

Each of the isomorphisms above respect the Real structures obtained from the constructions in Section 2.1, so the isomorphism in (5) respects the Real structures as well. Hence, we may identify the Real spaces $\text{KU}_{\mathbb{R}}(V) = C_{\text{gr}}^*(\mathfrak{s}, \mathcal{K}(\text{Cl}(V) \otimes L^2(V)))$.

To construct the multiplication maps $\mu_{V,W}^{\text{KU}_{\mathbb{R}}}$, notice that there is a coassociative and cocommutative comultiplication on \mathfrak{s} ,

$$\Delta: \mathfrak{s} = C_0(\mathbb{R}) \xrightarrow{\tilde{\Delta}} C_0(\mathbb{R}^2) \xrightarrow{m^{-1}} C_0(\mathbb{R}) \otimes C_0(\mathbb{R}) = \mathfrak{s} \otimes \mathfrak{s},$$

where $\tilde{\Delta}(a)(x, y) = \frac{1}{1+x^2+y^2}$, $\tilde{\Delta}(b)(x, y) = \frac{x+y}{1+x^2+y^2}$, and $m(f \otimes g)(x, y) = f(x)g(y)$.

If we endow $C_0(\mathbb{R}^2)$ with the Real structure given by complex conjugation on values, then it is clear that Δ is Real.

Definition 6.4. Given Real $\mathbb{Z}/2$ -graded C^* -algebras A, B , define the Real map

$$\star: C_{\text{gr}}^*(\mathfrak{s}, A) \wedge C_{\text{gr}}^*(\mathfrak{s}, B) \rightarrow C_{\text{gr}}^*(\mathfrak{s}, A \otimes B)$$

to be the composite of Real maps

$$C_{\text{gr}}^*(\mathfrak{s}, A) \wedge C_{\text{gr}}^*(\mathfrak{s}, B) \xrightarrow{\otimes} C_{\text{gr}}^*(\mathfrak{s} \otimes \mathfrak{s}, A \otimes B) \xrightarrow{\Delta^*} C_{\text{gr}}^*(\mathfrak{s}, A \otimes B).$$

Definition 6.5. Let $V, W \in \mathcal{I}_{C_2}$. Define the Real map

$$\mu_{V,W}^{\text{KU}_{\mathbb{R}}}: \text{KU}_{\mathbb{R}}(V) \wedge \text{KU}_{\mathbb{R}}(W) \rightarrow \text{KU}_{\mathbb{R}}(V \oplus W),$$

via the \star in Definition 6.4 with $A = \mathcal{K}(\text{Cl}(V) \otimes L^2(V))$ and $B = \mathcal{K}(\text{Cl}(W) \otimes L^2(W))$, where we use the identifications in (5), (4), and (2).

Next, we will define the unit, $\eta^{\text{KU}_\mathbb{R}}$, using functional calculus.

Definition 6.6. For $V \in \mathcal{I}_{C_2}$, define $\text{fc}_V: V \rightarrow C_{\text{gr}}^*(\mathfrak{s}, \text{Cl}(V))$, as follows: for $v \in V$,

$$\text{fc}_V(v)(a) = \frac{1}{1 + |v|^2} \quad \text{and} \quad \text{fc}_V(v)(b) = \frac{v}{1 + |v|^2}.$$

Proposition 6.7. The map fc_V is C_2 -equivariant.

Proof. On generators,

$$\overline{\text{fc}_V(v)}(a) = \overline{\text{fc}_V(v)(\bar{a})} = \overline{\text{fc}_V(v)(a)} = \text{fc}_V(v)(a) = \text{fc}_V(\bar{v})(a),$$

and

$$\overline{\text{fc}_V(v)}(b) = \overline{\text{fc}_V(v)(\bar{b})} = \overline{\text{fc}_V(v)(b)} = \frac{\bar{v}}{1 + |v|^2} = \frac{\bar{v}}{1 + |\bar{v}|^2} = \text{fc}_V(\bar{v})(b).$$

□

Since $\text{fc}_V(v) \rightarrow 0$ as $v \rightarrow \infty$, we get an induced C_2 -equivariant map

$$\beta_V: S^V \rightarrow C_{\text{gr}}^*(\mathfrak{s}, \text{Cl}(V))$$

of Real spaces. Let $P_{e^{-|\cdot|^2}} \in \mathcal{K}_V$ be the projection on the Gaussian $e^{-|\cdot|^2} \in L^2(V)$.

Definition 6.8. Define $p_V \in C_{\text{gr}}^*(\mathfrak{s}, \mathcal{K}_V)$ by

$$p_V(f) = f(0)P_{e^{-|\cdot|^2}},$$

i.e. the composite

$$\mathfrak{s} \xrightarrow{\text{eval}_0} \mathbb{C} \xrightarrow{P_{e^{-|\cdot|^2}}} \mathcal{K}_V.$$

Note that p_V is invariant with respect to the Real structure of $C_{\text{gr}}^*(\mathfrak{s}, \mathcal{K}_V)$. Then β_V and p_V can be combined to obtain a Real map

$$\eta_V^{\text{KU}_\mathbb{R}}: S^V \rightarrow \text{KU}_\mathbb{R}(V),$$

defined as the composite

$$S^V \cong S^V \wedge S^0 \xrightarrow{\beta_V \wedge p_V} C_{\text{gr}}^*(\mathfrak{s}, \text{Cl}(V)) \wedge C_{\text{gr}}^*(\mathfrak{s}, \mathcal{K}_V) \xrightarrow{\star} C_{\text{gr}}^*(\mathfrak{s}, \text{Cl}(V) \otimes \mathcal{K}_V) = \text{KU}_\mathbb{R}(V).$$

Proposition 6.9. The data $(\text{KU}_\mathbb{R}(V), \mu^{\text{KU}_\mathbb{R}}, \eta^{\text{KU}_\mathbb{R}})$ define a commutative orthogonal C_2 -ring spectrum, $\text{KU}_\mathbb{R}$, with underlying spectrum KU .

Proof. Nonequivariantly, all of our constructions agree with those of [7], so the only additional content here is the fact that $\mu_{V,W}^{\text{KU}_\mathbb{R}}$ and $\eta_V^{\text{KU}_\mathbb{R}}$ are C_2 -equivariant, for $V, W \in \mathcal{I}_{C_2}$. This was shown above as each map was constructed. □

Proposition 6.10. The C_2 -spectrum $\text{KU}_\mathbb{R}$ represents Atiyah's Real K-theory.

Proof. Since we know that the underlying spectrum of $\text{KU}_\mathbb{R}$ is KU it is sufficient to show that the C_2 -fixed points of $\text{KU}_\mathbb{R}$ is KO . Let $X \in \mathcal{T} \subset \mathcal{T}^{C_2}$, and set $S^{p,q} = S^{\mathbb{R}^{p,q}}$. By Lemma 5.8 of [7], there is a homeomorphism of spaces,

$$\begin{aligned}
\mathcal{T}_{C_2}(S^{p,q} \wedge X, \mathrm{KU}_{\mathbb{R}}(\mathbb{R}^{p+n,q})) &= \mathcal{T}_{C_2}(S^{p,q} \wedge X, C_{\mathrm{gr}}^*(\mathfrak{s}, \mathrm{Cl}_{p+n,q} \otimes \mathcal{K}_{\mathbb{R}^{p+n,q}})) \\
&\cong C_{\mathrm{gr}}^*(\mathfrak{s}, C_0(S^{p,q} \wedge X) \otimes \mathrm{Cl}_{p+n,q} \otimes \mathcal{K}_{\mathbb{R}^{p+n,q}}) \\
&\cong C_{\mathrm{gr}}^*(\mathfrak{s}, C_0(X) \otimes C_0(S^{p,q}) \otimes \mathrm{Cl}_{p,q} \otimes \mathrm{Cl}_n \otimes \mathcal{K}_{\mathbb{R}^{p+n,q}}),
\end{aligned}$$

which is also clearly C_2 -equivariant. Recall that the category of Real C^* -algebras is equivalent to the category of real C^* -algebras via fixed points in one direction and complexification in the other [8]. Thus, taking fixed points yields

$$\begin{aligned}
\mathcal{T}^{C_2}(S^{p,q} \wedge X, \mathrm{KU}_{\mathbb{R}}(\mathbb{R}^{p+n,q})) &\cong C_{\mathrm{gr}}^*(\mathfrak{s}, C_0(X) \otimes C_0(S^{p,q}) \otimes \mathrm{Cl}_{p,q} \otimes \mathrm{Cl}_n \otimes \mathcal{K}_{\mathbb{R}^{p+n,q}})^{C_2} \\
&\cong C_{\mathrm{gr}}^*(\mathfrak{s}^{C_2}, C_0(X)^{C_2} \otimes C_0(S^{p,q})^{C_2} \otimes (\mathrm{Cl}_{p,q})^{C_2} \otimes \mathrm{Cl}_n \otimes (\mathcal{K}_{\mathbb{R}^{p+n,q}})^{C_2}).
\end{aligned}$$

Let KKO denote Kasparov's real KK -functor and let KKR denote Kasparov's "Real" KK -functor [9]. By the real version of the main theorem in [16], we have

$$\pi_0 C_{\mathrm{gr}}^*(\mathfrak{s}^{C_2}, A \otimes (\mathcal{K}_{\mathbb{R}^{p+n,q}})^{C_2}) \cong \mathrm{KKO}(A),$$

so taking homotopy classes, we get

$$\begin{aligned}
[S^{p,q} \wedge X, \mathrm{KU}_{\mathbb{R}}(\mathbb{R}^{p+n,q})]_{C_2} &\cong \mathrm{KKO}(C_0(X)^{C_2} \otimes C_0(S^{p,q})^{C_2} \otimes (\mathrm{Cl}_{p,q})^{C_2} \otimes \mathrm{Cl}_n) \\
&\cong \mathrm{KKR}(C_0(X) \otimes C_0(S^{p,q}) \otimes \mathrm{Cl}_{p,q} \otimes \mathrm{Cl}_n),
\end{aligned}$$

using the equivalence of real C^* -algebras and Real C^* -algebras. By Kasparov's Real Bott periodicity [9],

$$\mathrm{KKR}(A) \cong \mathrm{KKR}(C_0(S^V) \otimes \mathrm{Cl}(V) \otimes A),$$

we have

$$\begin{aligned}
\mathrm{KKR}(C_0(X) \otimes C_0(S^{p,q}) \otimes \mathrm{Cl}_{p,q} \otimes \mathrm{Cl}_n) &\cong \mathrm{KKR}(C_0(X) \otimes \mathrm{Cl}_n) \\
&\cong \mathrm{KKO}(C_0(X)^{C_2} \otimes \mathrm{Cl}_n).
\end{aligned}$$

Now, we use the fact that the constructions in [7], when applied to real C^* -algebras, represent KO -theory, i.e. the spaces $\mathrm{KO}(n) := C_{\mathrm{gr}}^*(\mathfrak{s}^{C_2}, \mathrm{Cl}_n \otimes (\mathcal{K}_{\mathbb{R}^n})^{C_2})$ form a spectrum (in a way analogous to how we defined $\mathrm{KU}_{\mathbb{R}}$ representing KO). Since C_2 acts trivially on X , $C_0(X)^{C_2}$ is the algebra of real-valued continuous functions on X , so

$$\begin{aligned}
\mathrm{KKO}(C_0(X)^{C_2} \otimes \mathrm{Cl}_n) &\cong [X, C_{\mathrm{gr}}^*(\mathfrak{s}^{C_2}, \mathrm{Cl}_n \otimes (\mathcal{K}_{\mathbb{R}^n})^{C_2})] \\
&\cong [X, \mathrm{KO}(n)].
\end{aligned}$$

By putting the above isomorphisms together and taking a colimit, we find that

$$\mathrm{KU}_{\mathbb{R}}^*(X) \cong \mathrm{KO}^*(X),$$

for all X with trivial C_2 -action. By Proposition 2.16, this implies $(\mathrm{KU}_{\mathbb{R}})^{C_2} \simeq \mathrm{KO}$, which completes the proof. \square

We proceed to construct the Real spin orientation of $\mathrm{KU}_{\mathbb{R}}$. In the following definition, we use the identification in Remark 6.3 to view $\eta_V^{\mathrm{KU}_{\mathbb{R}}}$ as a map $S^V \rightarrow C_{\mathrm{gr}}^*(\mathfrak{s}, \mathcal{K}(\mathrm{Cl}(V) \otimes L^2(V)))$.

Definition 6.11. Define $\tilde{\alpha}_V: \mathrm{U}_V^{\mathrm{even}} \times V \rightarrow C_{\mathrm{gr}}^*(\mathfrak{s}, \mathcal{K}(\mathrm{Cl}(V) \otimes L^2(V)))$ by

$$\tilde{\alpha}_V(A, v) = (-1)^{|A|} A \eta_V^{\mathrm{KU}_{\mathbb{R}}}(v) A^{-1}.$$

Proposition 6.12. The map $\tilde{\alpha}_V$ in Definition 6.11 induces a C_2 -equivariant map

$$\alpha_V: \text{MSpin}_{\mathbb{R}}^c(V) \rightarrow \text{KU}_{\mathbb{R}}(V)$$

Proof. Let $(A, v) \in \text{U}_V^{\text{even}} \times V$, $g \in \text{Spin}^c(V)$, and $s \in \mathfrak{s}$. Then

$$\begin{aligned} \tilde{\alpha}_V(Ag^{-1}, gv)(s) &= (-1)^{|A \cdot g^{-1}|} (A \cdot g^{-1}) \eta_V^{\text{KU}_{\mathbb{R}}}(g \cdot v)(s) (A \cdot g^{-1})^{-1} \\ &= (-1)^{|A|} A i_V(g^{-1}) \eta_V^{\text{KU}_{\mathbb{R}}}(\rho_V(g)v)(s) i_V(g) A^{-1} \\ &= (-1)^{|A|} A i_V(g^{-1}) \eta_V^{\text{KU}_{\mathbb{R}}}((-1)^{|g|} g v g^{-1})(s) i_V(g) A^{-1}. \end{aligned}$$

Now let $w \otimes f \in \mathbb{C}l(V) \otimes L^2(V)$. For $s = a \in \mathfrak{s}$,

$$\begin{aligned} i_V(g^{-1}) \eta_V^{\text{KU}_{\mathbb{R}}}(g v g^{-1})(a) i_V(g)(w \otimes f) &= i_V(g^{-1}) \eta_V^{\text{KU}_{\mathbb{R}}}(g v g^{-1})(a) (g w \otimes f \circ \rho_V(g)^{-1}) \\ &= i_V(g^{-1}) \left(\frac{1}{1 + |g v g^{-1}|^2} g w \right) \otimes P_{e^{-|\cdot|^2}}(f \circ \rho_V(g)^{-1}) \\ &= g^{-1} \left(\frac{1}{1 + |v|^2} g w \right) \otimes P_{e^{-|\cdot|^2}}(f \circ \rho_V(g)^{-1} \circ \rho_V(g)) \\ &= \left(\frac{1}{1 + |v|^2} w \right) \otimes P_{e^{-|\cdot|^2}}(f) = \eta_V^{\text{KU}_{\mathbb{R}}}(v)(a)(w \otimes f). \end{aligned}$$

So $\tilde{\alpha}_V(Ag^{-1}, gv)(a) = \tilde{\alpha}_V(A, v)(a)$. For $s = b \in \mathfrak{s}$,

$$\begin{aligned} i_V(g^{-1}) \eta_V^{\text{KU}_{\mathbb{R}}}(g v g^{-1})(b) i_V(g)(w \otimes f) &= i_V(g^{-1}) \left(\frac{g v g^{-1}}{1 + |g v g^{-1}|^2} g w \right) \otimes P_{e^{-|\cdot|^2}}(f \circ \rho_V(g)^{-1}) \\ &= g^{-1} \left(\frac{g v g^{-1}}{1 + |v|^2} g w \right) \otimes P_{e^{-|\cdot|^2}}(f \circ \rho_V(g)^{-1} \circ \rho_V(g)) \\ &= \left(\frac{v}{1 + |v|^2} w \right) \otimes P_{e^{-|\cdot|^2}}(f) = \eta_V^{\text{KU}_{\mathbb{R}}}(v)(b)(w \otimes f). \end{aligned}$$

Thus, $\tilde{\alpha}_V(Ag^{-1}, gv) = \tilde{\alpha}_V(A, v)$, which means α_V is well-defined on $\text{U}_V^{\text{even}} \times_{\text{Spin}^c(V)} V$. We can extend α_V to ∞ by 0, since $\eta_V^{\text{KU}_{\mathbb{R}}}(\infty) = 0$; thus, $\alpha_V: \text{MSpin}_{\mathbb{R}}^c(V) \rightarrow \text{KU}_{\mathbb{R}}(V)$ is well-defined. For C_2 -equivariance, we note that $\eta_V^{\text{KU}_{\mathbb{R}}}$ is C_2 -equivariant, so we see that

$$\overline{\alpha([A, v])} = (-1)^{|A|} \overline{A \eta_V^{\text{KU}_{\mathbb{R}}}(v) A^{-1}} = (-1)^{|A|} \overline{A \eta_V^{\text{KU}_{\mathbb{R}}}(\bar{v}) A^{-1}} = \alpha_V(\overline{[A, v]}),$$

for $[A, v] \in \text{MSpin}_{\mathbb{R}}^c(V)$. □

Proposition 6.13. The maps α_V define a map of commutative orthogonal C_2 -ring spectra,

$$\alpha: \text{MSpin}_{\mathbb{R}}^c \rightarrow \text{KU}_{\mathbb{R}}.$$

Proof. By the previous proposition we showed that each component, α_V , is equivariant. It only remains to check naturality and monoidality, but these follow from Joachim [7], since our constructions are identical to his on underlying spaces. □

It is shown in [7] that on underlying spectra, α refines the Atiyah–Bott–Shapiro spin^c orientation of KU [2].

Corollary 6.14. Taking C_2 -fixed points, the map α above induces a map of E_∞ -ring spectra,

$$\mathrm{MSpin} \rightarrow \mathrm{KO},$$

refining the Atiyah–Bott–Shapiro spin orientation of KO [2].

Proof. The map follows from Proposition 6.13, Proposition 4.5, and $(\mathrm{KU}_\mathbb{R})^{C_2} \simeq \mathrm{KO}$. To see that it does indeed refine the Atiyah–Bott–Shapiro orientation, recall that the KO -cohomology class associated to a map $\mathrm{MSpin}(n) \rightarrow \mathrm{KU}_\mathbb{R}(\mathbb{R}^n)$ arises by taking fixed points (= “real points”) in the category of Real C^* -algebras (see the proof of Proposition 6.10). The result then follows from the proof in [7], together with the observation that the spin^c Atiyah–Bott–Shapiro Thom class associated to a spin vector bundle is the complexification of its spin Thom class. \square

This completes the proof of Theorem 1.1.

REFERENCES

- [1] M. F. Atiyah. Bott periodicity and the index of elliptic operators. *The Quarterly Journal of Mathematics*, 19(1):113–140, 01 1968.
- [2] M.F. Atiyah, R. Bott, and A. Shapiro. Clifford modules. *Topology*, 3:3–38, 1964.
- [3] Andrew J. Blumberg and Michael A. Hill. Operadic multiplications in equivariant spectra, norms, and transfers, 2015.
- [4] Allan L. Edelson. Real vector bundles and spaces with free involutions. *Transactions of the American Mathematical Society*, 157:179–188, 1971.
- [5] Michael A. Hill, Michael J. Hopkins, and Douglas C. Ravenel. On the non-existence of elements of Kervaire invariant one. *Ann. of Math (2)*, 184(1):1–262, 2016.
- [6] Michael A. Hill, Michael J. Hopkins, and Douglas C. Ravenel. *Equivariant Stable Homotopy Theory and the Kervaire Invariant Problem*. New Mathematical Monographs. Cambridge University Press, 2021.
- [7] Michael Joachim. *Higher coherences for equivariant K-theory*, page 87–114. London Mathematical Society Lecture Note Series. Cambridge University Press, 2004.
- [8] G. G. Kasparov. Hilbert c^* -modules : Theorems of stinespring and voiculescu. *Journal of Operator Theory*, 4(1):133–150, 1980.
- [9] G G Kasparov. The operator k -functor and extensions of c^* -algebras. *Mathematics of the USSR-Izvestiya*, 16(3):513, jun 1981.
- [10] Nicolaas H. Kuiper. The homotopy type of the unitary group of hilbert space. *Topology*, 3(1):19–30, 1965.
- [11] H. Blaine Lawson and Marie-Louise Michelsohn. *Spin Geometry (PMS-38)*. Princeton University Press, 1989.
- [12] M. A. Mandell, J. P. May, S. Schwede, and B. Shipley. Diagram spaces, diagram spectra, and FSP’s. Preprint, December 8, 1998, K-theory Preprint Archives, <http://www.math.uiuc.edu/K-theory/0319/>.
- [13] M. A. Mandell and J. P. May. Equivariant orthogonal spectra and S-modules. *Memoirs of the American Mathematical Society*, 159, 09 2002.
- [14] M. A. Mandell, J. P. May, S. Schwede, and B. Shipley. Model categories of diagram spectra. *Proceedings of the London Mathematical Society*, 82(2):441–512, 2001.
- [15] Stefan Schwede. Lectures on equivariant stable homotopy theory. 2016.
- [16] Jody Trout. On graded K -theory, elliptic operators and the functional calculus. *Illinois Journal of Mathematics*, 44(2):294 – 309, 2000.
- [17] Fuzhen Zhang. *Unitary Matrices and Contractions*, pages 131–158. Springer New York, New York, NY, 1999.