On the categoricity of complete second order theories*

Tapio Saarinen University of Helsinki Jouko Väänänen University of Helsinki and University of Amsterdam

Hugh Woodin Harvard University

Abstract

We show, assuming PD, that every complete finitely axiomatized second order theory with a countable model is categorical, but that there is, assuming again PD, a complete recursively axiomatized second order theory with a countable model which is non-categorical. We show that the existence of even very large (e.g. supercompact) cardinals does not imply the categoricity of all finite complete second order theories. More exactly, we show that a non-categorical complete finitely axiomatized second order theory can always be obtained by (set) forcing. We also show that the categoricity of all finite complete second order theories with a model of a certain singular cardinality κ of uncountable cofinality can be forced over any model of set theory. Previously, Solovay had proved, assuming V = L, that every complete finitely axiomatized second order theory (with or without a countable model) is categorical, and that in a generic extension of L there is a complete finitely axiomatized second order theory with a countable model which is non-categorical.

^{*}The first and second author would like to thank the Academy of Finland, grant no: 322795. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 101020762).

1 Introduction

A second order theory T is *complete* if it decides, in the semantical sense, every second order sentence ϕ in its own vocabulary i.e. if for every such ϕ either $T \models \phi$ or $T \models \neg \phi$, or equivalently, all models of T are second order equivalent. The question we investigate in this paper is whether every complete second order theory is *categorical* in the sense that all of its models are isomorphic. Already in 1928 Fraenkel [8] mentions this question as a question 'calling for clarification'. Carnap [6] claimed a positive answer, but his proof had an error (see [5]).

For mere cardinality reasons there are always complete non-categorical second order theories. One needs only consider models of the empty vocabulary. Since there are only continuum many different second order theories, there must be two such models of different cardinality with the same (a fortiori complete) second order theory.

Categoricity of complete second order theories would follow if all second order equivalent models were isomorphic, which is not the case again for cardinality reasons. However, if V = L, then countable second order equivalent models are isomorphic [1] and, moreover, every complete finitely axiomatized second order theory is categorical [26]. But if a Cohen real is added to a model of V = L, then there are countable non-isomorphic second order equivalent models [1], and if \aleph_1 Cohen-reals are added to a model of V = L, there is a complete finitely axiomatized second order theory (with a countable model) which is non-categorical [26].

Fraïssé [9, 10] conjectured that countable second order equivalent ordinals are equal. Marek [18, 19] showed that Fraïssé's conjecture is true under the assumption V = L, and false in a forcing extension obtained by collapsing an inaccessible cardinal to ω_1 .

The ambitious goal in the area of this paper is to decide in a definitive way the status of categoricity of complete second order theories. Since we are dealing with a question that cannot be decided in ZFC alone, it is natural to make an assumption such as PD, a consequence of the existence of large cardinals (e.g. infinitely many Woodin cardinals). We offer a partial solution to the full question by solving the case of second order theories with countable models. We have also partial results about theories with uncountable models. In particular, we show that a non-categorical complete finitely axiomatized second order theory can always be obtained by (set) forcing. This shows that large cardinal assumptions cannot imply, as V = L does, the categoricity of

all complete finitely axiomatized second order theories.

Notation: We recall the usual definition of the beth hierarchy: $\beth_0 = \omega$, $\beth_{\alpha+1} = 2^{\beth_{\alpha}}$, and $\beth_{\nu} = \sup_{\alpha < \nu} \beth_{\alpha}$ for limit ν . An ordinal α is called a beth fixed point if $\alpha = \beth_{\alpha}$. If μ is a cardinal, we use $\operatorname{Fn}(I, J, \mu)$ to denote the poset of partial functions $I \to J$ of cardinality $< \mu$, ordered by $p \le q \iff q \subseteq p$. The trivial poset $\operatorname{Fn}(\emptyset, \emptyset, 1)$ is denoted $(\{0\}, =)$.

We denote the second order theory of a structure M by $\operatorname{Th}_2(M)$. A second order theory T is complete if $\operatorname{Th}_2(M) = \operatorname{Th}_2(N)$ for all $M, N \models T$, and T is categorical if $M \cong N$ for all $M, N \models T$. For second order sentences ϕ, ψ we write $\phi \models \psi$ to mean $M \models \phi$ implies $M \models \psi$ for all M, and similarly $T \models T'$ for second order theories T, T', and we say T axiomatizes T'. If T is a finite (resp. recursive) set of sentences and $T \models T'$, we say T' is finitely (resp. recursively) axiomatizable.

A cardinal λ is second order characterizable if there is a second order sentence ϕ in the empty vocabulary such that $N \models \phi$ if and only if $|N| = \lambda$.

2 The case of L[U]

It is already known that if V = L, then every complete finitely axiomatized second order theory is categorical [26]. We now show that this also holds if V = L[U], and we show there are complete recursively axiomatized second order theories that are non-categorical (with very large models).

Assuming V = L[U], we write κ for the sole measurable cardinal, U for the unique normal measure on κ and $<_{L[U]}$ for the canonical well-order. By a L[U]-premouse we mean a structure $(L_{\alpha}[W], \in, W)$ where W is an $L_{\alpha}[W]$ -ultrafilter on some $\gamma < \alpha$. Recall that a premouse $(L_{\alpha}[W], \in, W)$ is iterable (under taking iterated ultrapowers), i.e. that every iterate is well-founded, if and only if every iterate in an iteration of any countable length is well-founded. Observe that every iterate in an iteration of countable length has the same cardinality as the original premouse, so the iterability of a premouse is expressible in second order logic. See for example [12, chapter 20] for more details.

Theorem 1. Assume V = L[U]. Every complete finitely axiomatized second order theory is categorical.

Proof. Suppose ϕ is a complete second order sentence in a vocabulary with a single binary relation symbol R (for simplicity). Note first that ϕ has models

in only one cardinality. If not, let N be a model of ϕ of least cardinality, and M another model with |M| > |N|. Let θ be the sentence

$$\exists P \exists R'(\theta'(P) \land \phi'(P, R'))$$

where

- P is a unary predicate, not occurring in ϕ , and R' is a binary relation symbol not occurring in ϕ .
- $\phi'(P, R')$ is a modification of the sentence ϕ , where the first order quantifiers $\exists x \dots$ are relativised to P as $\exists x (P(x) \land \dots)$, and each occurrence of R is replaced by R'.
- $\theta'(P)$ says that the cardinality of P is smaller than the ambient domain of the model (for example, that there is no injective function with range contained in P).

As ϕ is complete and $M \models \theta$ (by taking (P, R') isomorphic to N), also $N \models \theta$, so there is a model of ϕ of cardinality smaller than that of N, which is a contradiction. Thus all models of ϕ have the same cardinality.

Now let M_0 be the $<_{L[U]}$ -least model of ϕ . Suppose first that $|M_0| > \kappa$: in this case we can mimic the categoricity argument for L as follows. Let θ be the sentence

$$\exists E \exists u \exists m \exists P \exists R'(\theta'(E, u) \land \phi'(P, R') \land \theta_{least}(E, u, m) \land \theta_{isom}(E, m, P, R'))$$

where

- E, R' are binary predicate symbols, P a unary predicate symbol and u, m are first order variables, none occurring in ϕ (the intuition is that E is \in , u is a normal ultrafilter, m is a structure in the vocabulary of ϕ , P is the domain of m and R is the single binary relation of m).
- $\theta'(E, u)$ states E is well-founded and extensional (so that E has a transitive collapse, and the domain of the model equipped with E can be thought of as a transitive set), and its collapse is a level of L[u] having a normal measure u as an element.
- $\phi'(P, R')$ is (as before) a modification of the sentence ϕ where each first order quantifier $\exists x \dots$ is relativised to P as $\exists x (P(x) \land \dots)$, and each occurrence of R is replaced by R'.

- $\theta_{least}(E, u, m)$ says $m <_{L[u]} m'$ for any other m' = (Q, S) also satisfying $\phi'(Q, S)$ (using the formula defining the canonical well-order of L[u] with u as a parameter).
- $\theta_{isom}(E, m, P, R')$ states that m = (P, R'), and that (P, R') is isomorphic to the ambient model (so there is an injection F with range P such that $R(x, y) \leftrightarrow R'(F(x), F(y))$ for all x, y).

If $M \models \theta$ with witnesses E, u and m = (P, R'), and $\pi : (M, E) \to (N, \in)$ is the transitive collapse, then $\pi(u) = U$ is the unique normal measure U on κ , $N = L_{\alpha}[U]$ for some α and $\pi(m)$ is the $<_{L[U]}$ -least model M_0 of ϕ , so M is isomorphic to M_0 .

Conversely, let α be least such that $M_0 \in L_{\alpha}[U]$. Then $\kappa < \alpha < |M_0|^+$ and $U \in L_{\alpha}[U]$, so we may pick a bijection $\pi \colon M_0 \to L_{\alpha}[U]$ and let E, u and m = (P, R') be the preimages of \in , U and M_0 under π to witness $M_0 \models \theta$.

Thus the above sentence θ is such that $M \models \theta$ if and only if M is isomorphic to the $<_{L[U]}$ -least model of ϕ . Now if $M \models \phi$, also $M \models \theta$ by completeness of ϕ , so M is isomorphic to M_0 and ϕ is categorical.

Suppose now that $|M_0| = \lambda < \kappa$. In this case we cannot find a binary relation E on M_0 and $u \in M_0$ such that u is a normal measure in the transitive collapse of (M_0, E) , so we modify the previously produced sentence θ . This argument relies on a straightforward modification of the Δ_3^1 well-order of reals in L[U]. We make the further assumption that the domain of M_0 is a cardinal (and that M_0 is the $<_{L[U]}$ -least among such models), and let θ be the sentence

 $\exists E \exists W \exists m \exists P \exists R' (\theta'(E, W) \land \phi'(E, P, R') \land \theta_{least}(E, W, m) \land \theta_{isom}(E, m, P, R'))$ where

- E, R' are binary and W, P unary predicate symbols, and m a first order variable, none occurring in ϕ .
- $\theta'(E, W)$ states E is well-founded and extensional, whose transitive collapse is an iterable L[U]-premouse $(L_{\alpha}[W], \in, W)$ for some α , where W is a L[W]-ultrafilter on some γ , where γ is an ordinal greater than the cardinality of the ambient model.
- $\phi'(E, P, R')$ is the sentence $\phi'(P, R')$ from before, with the additional stipulation that the extent of the predicate P is a cardinal.

- $\theta_{least}(E, W, m)$ says $m <_{L[W]} m'$ for any other m' = (Q, S) also satisfying $\phi'(E, Q, S)$ (using the formula defining the canonical well-order of L[W] with W as a predicate).
- $\theta_{isom}(E, m, P, R')$ remains unchanged from earlier.

We claim that θ is a sentence such that $M \models \theta$ if and only if M is isomorphic to the $<_{L[U]}$ -least model of ϕ (among models whose domain is a cardinal). So suppose $M \models \theta$ with witnesses E, W and m = (P, R'), and let $\pi : (M, E, W) \to (N, \in, W')$ be the transitive collapse. Then $W' = \pi''(W)$ is a N-ultrafilter on some $\gamma > \lambda$ and $N = L_{\alpha}[W']$ for some $\alpha > \gamma$, and $\pi(m)$ is the $<_{L[W']}$ -least model of ϕ in $L_{\alpha}[W']$, to which M is isomorphic.

To see why $\pi(m)$ is M_0 , let $j: L[U] \to L[F]$ and $k: L_{\alpha}[W'] \to L_{\delta}[F]$ be long enough iterations of L[U] and $L_{\alpha}[W]$ respectively such that they become comparable. Then $\operatorname{crit}(j) = \kappa > \lambda$ and $\operatorname{crit}(k) = \gamma > \lambda$, so $j(M_0) = M_0$ and $k(\pi(m)) = \pi(m)$. By elementarity, both M_0 and $\pi(m)$ are now the $<_{L[F]}$ -least model of ϕ among models whose domain is a cardinal, so $\pi(m) = M_0$ and M is isomorphic to M_0 .

Conversely, to see $M_0 \models \theta$ amounts to finding an appropriate premouse $(L_{\alpha}[W], \in, W)$. Let δ be a large enough cardinal such that $M_0, U \in L_{\delta}[U]$, and that $(L_{\delta}[U], \in, U)$ is an iterable premouse. Then let N be the Skolem hull of $\lambda \cup \{M_0\}$ in $L_{\delta}[U]$ of cardinality λ , and let $\pi: (N, \in, U \cap N) \to (L_{\alpha}[W], \in, W)$ be the transitive collapse. Now $(L_{\alpha}[W], \in, W)$ is also an iterable premouse with $|L_{\alpha}[W]| = \lambda$, W is a $L_{\alpha}[W]$ -ultrafilter on some $\gamma = \pi(\kappa) > \lambda$, and $\pi(M_0) = M_0$, so by elementarity M_0 is the $\langle L_{[W]}$ -least model of ϕ as required. So θ is a sentence such that $M \models \theta$ if and only if M is isomorphic to M_0 , implying as before that ϕ is categorical.

Finally, observe that the case $|M_0| = \kappa$ is impossible, since the measurable cardinal κ is Π_1^2 -indescribable [11]. Thus ϕ is categorical.

It turns out that finite axiomatizability is key for the preceding theorem. For every second order characterizable cardinal $\lambda > \kappa$, we produce a non-categorical recursively axiomatizable theory whose models have cardinality λ .

Theorem 2. Assume V = L[U]. Suppose κ is measurable and λ is second order characterizable with $\lambda > \kappa$. Then there is a recursively axiomatizable theory T with κ many non-isomorphic models of cardinality λ .

Proof. For $\alpha < \kappa$ let $M_{\alpha} = (\lambda + \alpha, <)$, so in a structure of cardinality λ , M_{α} is straightforwardly definable from α (as λ is second order characterizable). These models have the property that $M_{\alpha} \cong M_{\beta}$ implies $M_{\alpha} = M_{\beta}$.

For a second order sentence ϕ in vocabulary (<), let

$$S_{\phi} = \{ \alpha < \kappa : M_{\alpha} \models \phi \},$$

and let T_0 be the set of sentences ϕ such that $S_{\phi} \in U$. As U is an ultrafilter, T_0 is a complete theory (so for any ϕ , exactly one of $\phi \in T_0$ or $\neg \phi \in T_0$ hold), and by the σ -completeness of U the intersection $X = \bigcap \{S_{\phi} : \phi \in T_0\} \in U$ is nonempty. The set X is such that for any $\alpha, \beta \in X$, the structures M_{α} , M_{β} have the same second order theory T_0 , so it remains to see that the theory T_0 is recursively axiomatizable.

For a second order sentence ϕ in vocabulary (<), let E be a binary relation symbol and u a first order variable, neither occurring in ϕ , and let ϕ^+ be the second order sentence

$$\exists E \exists u (\theta'(E, u) \land (\exists x \in u) (\forall \alpha \in x)" M_{\alpha} \models \phi")$$

where $\theta'(E, u)$ says E is well-founded and extensional, and its transitive collapse is a level of L[u] containing λ and having a normal measure u as an element. Note that ϕ^+ is a sentence in the empty vocabulary. Intuitively, ϕ^+ states that $M_{\alpha} \models \phi$ for a U-big set of ordinals $\alpha < \kappa$, so for any structure N with $|N| = \lambda$ we have the equivalences

$$N \models \phi^+ \iff \{\alpha < \kappa : M_\alpha \models \phi\} = S_\phi \in U$$

 $\iff M_\alpha \models \phi \text{ for some } \alpha \in X$
 $\iff \phi \in T_0.$

The import of the vocabulary of ϕ^+ being empty is that for a structure N, the truth of $N \models \phi^+$ depends only on |N|, so we get that for all structures N with $|N| = \lambda$,

$$N \models \phi^+ \iff M_\alpha \models \phi^+ \text{ for some } \alpha \in X \iff \phi^+ \in T_0$$

so also $\phi \leftrightarrow \phi^+ \in T_0$ for all second order sentences ϕ in vocabulary (<). Now define the recursive set of sentences

 $T = \{\phi \leftrightarrow \phi^+ : \phi \text{ is a second order sentence in vocabulary } (<)\}.$

Observe that any model N of the theory T has cardinality λ , since taking θ_{λ} to be the second order characterization of λ , we have $M_{\alpha} \models \theta_{\lambda}$ for all $\alpha < \kappa$, so $N \models \theta_{\lambda}^+$ and thus $N \models \theta_{\lambda}$ since $\theta_{\lambda}^+ \leftrightarrow \theta_{\lambda} \in T$.

To see T axiomatizes T_0 , suppose $N \models T$ so $|N| = \lambda$, and that ϕ is a second order sentence in the vocabulary (<), so either $\phi \in T_0$ or $\neg \phi \in T_0$. In the former case we have $S_{\phi} \in U$ so $N \models \phi^+$, so $N \models \phi$, and in the latter case we have $S_{\neg \phi} \in U$ so $N \models \neg \phi$. Thus $\operatorname{Th}_2(N) = T_0$, so T recursively axiomatizes T_0 as desired.

In conclusion, all complete finitely axiomatizable theories are categorical in L[U] as in L, and in L[U] there are complete recursively axiomatizable second order theories that are non-categorical (whereas this is still unknown in L).

3 Countable models

We already remarked earlier that if V = L, then every complete finitely axiomatized second order theory is categorical [26]. We now show that for theories with a countable model this is a consequence of PD, and therefore a consequence of large cardinals:

Theorem 3. Assume PD. Every complete finitely axiomatized second order theory with a countable model is categorical.

Proof. Suppose ϕ is a complete second order sentence with a countable model. Then by completeness all models of ϕ are countable. Suppose ϕ is on the level Σ_n^1 of second order logic and its vocabulary is, for simplicity, just one binary predicate symbol P. Let R be the Σ_n^1 (lightface) set of real numbers coding models of ϕ . By PD and its consequence, the Projective Uniformization Theorem [22, Theorem 6C5], there is a Σ_{n+1}^1 (even Σ_n^1 if n is even) element r in R. Suppose r codes the model M of ϕ . We show that every model of ϕ is isomorphic to M. Suppose N is a model of ϕ . Let θ be the following second order sentence:

$$\exists Q_{+} \exists Q_{\times} (\theta_{1}(Q_{+}, Q_{\times}) \wedge \exists A(\theta_{2}(Q_{+}, Q_{\times}, A) \wedge \exists B(\theta_{3}(Q_{+}, Q_{\times}, A, B) \wedge \exists F \theta_{4}(F, B)))),$$

where

• $\theta_1(Q_+, Q_\times)$ is the standard second order characterization of $(\mathbb{N}, +, \times)$.

- $\theta_2(Q_+, Q_\times, A)$ says that the set A satisfies the Σ_{n+1}^1 definition of r in terms of Q_+ and Q_\times .
- $\theta_3(Q_+, Q_\times, A, B)$ says in a domain N that (N, B) is the binary structure coded by A in terms of Q_+ and Q_\times .
- $\theta_4(F,B)$ is the second order sentence which says that F is a bijection and

$$\forall x \forall y (P(x,y) \leftrightarrow B(F(x),F(y))).$$

Thus, θ essentially says "I am isomorphic to the model coded by r." Trivially, $M \models \theta$. Recall that $M \models \phi$. Since ϕ is complete, $\phi \models \theta$. Therefore our assumption $N \models \phi$ implies $N \models \theta$ and therefore $N \cong M$.

We make a few remarks about the proof. First, if n=2, then we can use the Novikov-Kondo-Addison Uniformisation Theorem and PD is not needed. Thus we obtain:

Corollary 4. A complete Σ_2^1 -sentence of second order logic with a countable model is always categorical.

In fact, the categorical finite second order axiomatizations of structures such as $(\mathbb{N}, +, \times)$, $(\mathbb{R}, +, \times, 0, 1)$ and $(\mathbb{C}, +, \times, 0, 1)$ (any many other classical structures) are all on the Π^1_1 -level of second order logic.

Second, the above proof gives also the following more general result: Assume $Det(\Delta_{2n}^1)$. Suppose T is a recursively axiomatized theory on the Σ_{2n+2}^1 -level of second order logic, which is complete for sentences on this level of second order logic. Then T is categorical.

An essential ingredient of the proof of Theorem 3 was the assumption that the complete second order theory is finitely axiomatized. The following theorem shows that "finitely" cannot be replaced by "recursively".

Theorem 5. Assume PD. There is a recursively axiomatized complete second order theory with 2^{ω} non-isomorphic countable models.

Proof. For any $x \subseteq \omega$ let

$$M_r = (V_\omega \cup \{y \subseteq \omega : y \equiv_T x\}, \in),$$

where $y \equiv_T x$ means the Turing-equivalence of y and x. We denote the second order theory of M_x by $\text{Th}_2(M_x)$. By construction, $x \equiv_T y$ implies

 $\operatorname{Th}_2(M_x) = \operatorname{Th}_2(M_y)$. On the other hand, if $x \not\equiv_T y$, then clearly $M_x \not\cong M_y$. If ϕ is a second order sentence, then ' $M_x \models \phi$ ' is a projective property of x, closed under \equiv_T , and hence by Turing Determinacy for projective sets [20] has a constant truth value on a cone of reals x. By intersecting the cones we get a cone C of reals x on which $\operatorname{Th}_2(M_x)$ is constant. For any second order ϕ let ϕ^+ be the second order sentence

"
$$M_y \models \phi$$
 for a cone of y"

and $\hat{\phi}$ the sentence $\phi \leftrightarrow \phi^+$. Let us consider the recursive second order theory T consisting of $\hat{\phi}$, when ϕ ranges over second order sentences in the vocabulary of the structures M_x . We may immediately conclude that T is complete, for if a second order sentence ϕ is given, then by the choice of C either $M_x \models \phi$ for $x \in C$ or $M_x \models \neg \phi$ for $x \in C$. In the first case $\hat{\phi} \models \phi$ and in the second case $\hat{\phi} \models \neg \phi$. Therefore, $T \models \phi$ or $T \models \neg \phi$. There are a continuum of non-Turing equivalent reals in the cone C. Hence there are a continuum of non-isomorphic M_x with $x \in C$.

4 Models of cardinality \aleph_1

Next, we show that the (*) axiom (see Definition 4.33 in [28]) has categoricity consequences for theories with a model of cardinality \aleph_1 . Thus these consequences can also be derived from forcing axioms, namely MM^{++} which implies the (*) axiom (as shown in [4]). The following theorem answers a question of Boban Veličković.

Theorem 6. Assume (*). Then there is a complete finitely axiomatizable second order theory with ω_2 (= 2^{ω_1}) non-isomorphic models of cardinality \aleph_1 .

Proof. The pertinent consequence of (*) is the quasihomogeneity of the nonstationary ideal on ω_1 (see Section 5.8 in [28], particularly Definition 5.100). We take "NS $_{\omega_1}$ is quasihomogeneous" to be the following statement: if $X \subseteq \mathcal{P}(\omega_1)$ is ordinal definable from parameters in $\mathbb{R} \cup \{NS_{\omega_1}\}$, and X is closed under equality modulo NS_{ω_1} , and X contains one bistationary subset of ω_1 , then X contains every bistationary subset of ω_1 .

We focus on the ω_1 -like dense linear orders $\Phi(S) = \eta + \sum_{\alpha < \omega_1} \eta_{\alpha}$, where

$$\eta_{\alpha} = \begin{cases} \eta, & \alpha \notin S \\ 1 + \eta, & \alpha \in S, \end{cases}$$

 η is the order type of the rationals, and $S \subseteq \omega_1$ is bistationary. These models have the property that $\Phi(S) \cong \Phi(S')$ if and only if $S \triangle S' \in NS_{\omega_1}$. For a second order sentence ϕ in vocabulary (<), the set

$$X_{\phi} = \{ S \subseteq \omega_1 : S \text{ bistationary, } \Phi(S) \models \phi \}$$

is ordinal definable, and closed under equality modulo NS_{ω_1} , so the quasihomogeneity of NS_{ω_1} implies that X_{ϕ} contains either every bistationary subset of ω_1 , or none of them.

This shows the models $\Phi(S)$ for bistationary $S \subseteq \omega_1$ all have the same complete second order theory, which is thus non-categorical. This theory is axiomatized by the second order sentence in vocabulary (<) expressing "I am isomorphic to $\Phi(S)$ for some bistationary $S \subseteq \omega_1$ ", so it is finitely axiomatizable, as required.

Some categoricity consequences of (*) can already be derived from AD, the axiom of determinacy. As the axiom (*) states that $L[\mathcal{P}(\omega_1)]$ is a homogeneous forcing extension of a model of AD by a forcing that does not add reals, the categoricity consequences of AD for theories with a model of cardinality $\leq \aleph_1$ also follow from (*). (Of course, the existence of recursively axiomatized non-categorical theories under (*) is overshadowed by the existence of even finitely axiomatized such theories.)

Theorem 7. Assume AD. Then there is a complete recursively axiomatized second order theory with at least 2^{\aleph_0} many models of cardinality \aleph_1 .

Proof. By Martin, AD implies $\omega_1 \to (\omega_1)^{\omega}$, and moreover the homogeneous set given by $\omega_1 \to (\omega_1)^{\omega}$ can be taken to be a club (see [14]). We may then intersect ω many homogeneous clubs for ω many colorings to obtain $\omega_1 \to (\omega_1)_{2\omega}^{\omega}$, and the homogeneous subset can still be taken to be a club.

We focus on models of the form $M_X = (\omega_1, <, X)$ for $X \in [\omega_1]^{\omega}$. The second order theory $\operatorname{Th}_2(M_X)$ in the vocabulary (<, X) can be encoded by a real $f(X) \in 2^{\omega}$ consisting of the Gödel numbers of the sentences true in M_X . This gives a coloring $f : [\omega_1]^{\omega} \to 2^{\omega}$, so we find a homogeneous club subset $H_0 \subseteq \omega_1$ such that f(X) does not depend on $X \in [H_0]^{\omega}$. Hence the models M_X with $X \in [H_0]^{\omega}$ all have the same complete second order theory T_0 , which is thus non-categorical.

The theory T_0 is axiomatized by

$$T = \{\phi \leftrightarrow \phi^+ : \phi \text{ is a second order sentence}\}\$$

where for a given second order sentence ϕ in vocabulary (<, X), the sentence ϕ^+ expresses "there exists a club $C \subseteq \omega_1$ such that $M_X \models \phi$ for all $X \in [C]^{\omega}$ ".

For a given second order sentence ϕ , if $M_X \models \phi$ for each $X \in [H_0]^\omega$, then H_0 serves to witness that ϕ^+ holds, so $T \models \phi$. Conversely, if ϕ^+ holds, there is a club C such that $M_X \models \phi$ for every $X \in [C]^\omega$, and taking $X \in [C \cap H_0]^\omega$ we see also that $M_X \models \phi$ for all $X \in [H_0]^\omega$. Thus $T \models \phi$ for exactly those ϕ such that $M_X \models \phi$ for all $X \in [H_0]^\omega$, so we see that T is a recursive axiomatization of the theory T_0 as desired.

The same can be analogously derived from the (*) axiom, as follows:

Corollary 8. Assume (*). Then there is a complete recursively axiomatized second order theory with ω_2 many models of cardinality \aleph_1 .

Proof. Recall (*) states that $L[\mathcal{P}(\omega_1)] = L(\mathbb{R})^{\mathbb{P}_{\text{max}}}$ and AD holds in $L(\mathbb{R})$. As \mathbb{P}_{max} is homogeneous and does not add reals under AD (see Lemmas 4.40 and 4.43 in [28]), $\omega_1 = \omega_1^{L(\mathbb{R})}$ and $[\omega_1]^{\omega} = ([\omega_1]^{\omega})^{L(\mathbb{R})}$.

We again look at models $M_X = (\omega_1, <, X)$ for $X \in [\omega_1]^{\omega}$, and working in $L(\mathbb{R})$, define a coloring $f: [\omega_1]^{\omega} \to 2^{\omega}$ by

$$f(X) = r \iff L(\mathbb{R}) \models \mathbb{P}_{\max} \Vdash "\check{r} \text{ codes } \operatorname{Th}_2(M_{\check{X}})".$$

That f is a well-defined total function relies on the homogeneity of \mathbb{P}_{\max} . By $\mathrm{AD}^{L(\mathbb{R})}$ we find a club $H_0 \in L(\mathbb{R})$, $H_0 \subseteq \omega_1$ homogeneous for f. Stepping out of $L(\mathbb{R})$, we see that the models M_X , $X \in [H_0]^{\omega}$ all have the same complete second order theory T_0 (in $L(\mathbb{R})^{\mathbb{P}_{\max}} = L[\mathcal{P}(\omega_1)]$ and in V both).

Working now in V, we again define

$$T = \{\phi \leftrightarrow \phi^+ : \phi \text{ is a second order sentence}\}\$$

where for a given second order sentence ϕ , the sentence ϕ^+ expresses "there exists a club $C \subseteq \omega_1$ such that $M_X \models \phi$ for all $X \in [C]^{\omega}$ ". The proof concludes analogously to the preceding theorem.

We note that (*) calculates $|\omega_1^{\omega}|$ to be ω_2 , so T_0 has ω_2 many non-isomorphic models as claimed.

Of course, we may also use the fact that the club filter on ω_1 is an ultrafilter under AD to get another complete recursively axiomatized non-categorical second order theory, the difference being that this theory has ω_1 many models instead. The proof, analogous to the proof of Theorem 2, is omitted:

Theorem 9. Assume AD. Then there is a complete recursively axiomatized second order theory with ω_1 many models of cardinality \aleph_1 .

This proof is also easily modified to assume (*) instead:

Corollary 10. Assume (*). Then there is a complete recursively axiomatized second order theory with ω_1 many models of cardinality \aleph_1 .

Thus, under (*), a complete non-categorical theory with a model of cardinality \aleph_1 may have either ω_1 or ω_2 many non-isomorphic models.

5 Forcing non-categoricity

We shall show (Theorem 14) that we can force, over any model of set theory, a finite complete non-categorical second order theory with a model of cardinality \aleph_1 . This shows that large cardinals cannot imply the categoricity of finite complete second order theories in general and, in particular, in the case that the theory has a model of cardinality \aleph_1 . This is in contrast to finite complete second order theories with a countable model where PD implies categoricity (Theorem 3).

Here is an outline of the proof. We start with a preparatory countably closed forcing \mathbb{P} obtaining a generic extension V[G]. Then we add \aleph_1 Cohenreals obtaining a further generic extension V[G][H]. In this model we consider for every $x \subseteq \omega$ the model

$$M_x = (HC^{V[x]}, HC^V, \in). \tag{1}$$

We show that if x is Cohen-generic over V[G], then the complete second order theory of M_x is finitely axiomatizable (in second order logic), and if x and y are mutually Cohen-generic over V[G], then M_x and M_y are second order equivalent but non-isomorphic.

We begin by recalling the following fast club forcing \mathbb{P}_{fast} , due to R. Jensen: Conditions are pairs $p = (c_p, E_p)$ where c_p is a countable closed subset of ω_1 and C_p is club in ω_1 . We define $(c_p, E_p) \leq (c_q, E_q)$ if c_q is an initial segment of c_p , $E_q \subseteq E_p$, and $c_p \setminus c_q \subseteq E_q$. This forcing is countably closed. If we assume CH, this forcing has the \aleph_2 -c.c. It is called fast club forcing because of the following property: Suppose G is \mathbb{P}_{fast} -generic. If C_G is the union of the sets c_p such that $p \in G$, then the following holds: If D is

any club in V, then there is α such that $C_G \setminus \alpha \subseteq D$. The set C_G is called a fast club (over V).

Let \mathbb{Q} be the poset $\operatorname{Fn}(\omega_1 \times \omega, 2, \omega)$ for adding \aleph_1 Cohen reals. We use fast club forcing to build a preparatory iterated forcing in such a way that after forcing with \mathbb{Q} the ground model reals are second order definable from any set $A \subseteq \omega_1$ with a certain second order property. The following lemma is crucial in the iteration:

Lemma 11. Suppose $G \times H$ is $\mathbb{P}_{fast} \times \mathbb{Q}$ -generic over V. Suppose $A \subseteq \omega_1$ is in V[H] and $D \subseteq C_G$ is a club in $V[G \times H]$ such that $V[G \times H]$ satisfies $\forall \alpha < \omega_1(D \cap \alpha \in L[A])$. Then $\mathcal{P}(\omega)^V \subseteq L[A]$.

Proof. We modify a construction from the proof of [30, Lemma 4.33] to our context. Let us call a pair (A, B) of sets of ordinals an *interlace*, if $A \cap B = \emptyset$, above every element of A there is an element of B, and vice versa. Suppose we have disjoint sets $X, Y, Z \subseteq \omega_1$ such that $(X \cup Y, Z)$ is an interlace. Let $z \sim z'$ in Z if $(z, z') \cap (X \cup Y) = \emptyset$. Let $[z_n]$, $n < \omega$, be the first $\omega \sim$ equivalence classes in Z in increasing order. The triple (X, Y, Z) is said to code the set $a \subseteq \omega$ if for all $n < \omega$:

$$n \in a \iff \min\{\alpha \in X \cup Y : [z_n] < \alpha < [z_{n+1}]\} \in X.$$

It suffices to prove that for every $a \subseteq \omega$ in V there is a triple $(X, Y, Z) \in L[A]$ such that $(X \cup Y, Z)$ is an interlace, and (X, Y, Z) codes a. To this end, suppose $a \in \mathcal{P}(\omega)^V$.

Suppose A is a \mathbb{Q} -name for A in $V, \tau \in V$ is a \mathbb{P}_{fast} -name for a \mathbb{Q} -name \dot{D} for D, and \dot{F} a \mathbb{Q} -name for a function $\omega_1 \to \omega_1$ which lists the elements of \dot{D} in increasing order. W.l.o.g. τ is a \mathbb{P}_{fast} -name $\langle \dot{f}_{\alpha} : \alpha < \omega_1 \rangle$ for a sequence of countable partial functions defined on ω_1 such that $\{\dot{f}_{\alpha}(\gamma) : \gamma \in \text{dom}(f_{\alpha})\}$ is a maximal antichain in \mathbb{Q} and $\dot{f}_{\alpha}(\gamma)$ forces $\dot{F}(\alpha) = \gamma$. Suppose (w.l.o.g.) the weakest condition in $\mathbb{P}_{\text{fast}} \times \mathbb{Q}$ forces what is assumed about \dot{A}, \dot{F}, τ and \dot{D} . Since $\mathbb{P}_{\text{fast}} \Vdash$ " $\mathbb{Q} \Vdash \dot{D} \subseteq C_{\dot{G}}$ ", we have $\Vdash \text{dom}(\dot{f}_{\alpha}) \subseteq C_{\dot{G}}$. More generally, if $p \in \mathbb{P}_{\text{fast}}$ decides the countable set $\text{dom}(\dot{f}_{\alpha})$, then

$$p \Vdash \operatorname{dom}(\dot{f}_{\alpha}) \subseteq c_p \setminus \alpha. \tag{2}$$

If $\delta < \omega_2$, let W_{δ} be the set of conditions $p \in \mathbb{P}_{\text{fast}}$ such that p decides $\text{dom}(\dot{f}_{\delta})$. It is easy to see that W_{δ} is dense.

We construct descending ω -sequences (p_n) , (q_n) and (r_n) in \mathbb{P}_{fast} as follows. We let $p_0 = q_0 = r_0$ be the weakest condition in \mathbb{P}_{fast} . Suppose p_n, q_n and r_n have been defined already. Let $\delta_n = \max(c_{r_n} \cup \{0\})$. Now there are two cases:

1. Case $n \in a$:

- (a) Let $p_{n+1} \leq p_n$ such that $\min(c_{p_{n+1}} \setminus c_{p_n}) > \delta_n$ and $p_{n+1} \in W_{\delta_n}$.
- (b) Let $q_{n+1} \leq q_n$ such that $\min(c_{q_{n+1}} \setminus c_{q_n}) > \max(c_{p_{n+1}})$ and $q_{n+1} \in W_{\delta_n}$.
- (c) Let $r_{n+1} \leq r_n$ such that $\min(c_{r_{n+1}} \setminus c_{r_n}) > \max(c_{q_{n+1}})$ and $q_{n+1} \in W_{\delta_n}$.

2. Case $n \notin a$:

- (a) Let $q_{n+1} \leq q_n$ such that $\min(c_{q_{n+1}} \setminus c_{q_n}) > \delta_n$ and $q_{n+1} \in W_{\delta_n}$.
- (b) Let $p_{n+1} \leq p_n$ such that $\min(c_{p_{n+1}} \setminus c_{p_n}) > \max(c_{q_{n+1}})$ and $p_{n+1} \in W_{\delta_n}$.
- (c) Let $r_{n+1} \leq r_n$ such that $\min(c_{r_{n+1}} \setminus c_{r_n}) > \max(c_{p_{n+1}})$ and $r_{n+1} \in W_{\delta_n}$.

Note that if $\delta_n < \alpha < \min(c_{p_{n+1}} \setminus c_{p_n})$, then $p_{n+1} \Vdash \alpha \notin C_{\dot{G}}$, whence $p_{n+1} \Vdash \alpha \notin \tau$. Respectively, if $\delta_n < \alpha < \min(c_{q_{n+1}} \setminus c_{q_n})$, then $q_{n+1} \Vdash \alpha \notin C_{\dot{G}}$, whence $q_{n+1} \Vdash \alpha \notin \tau$, and if $\delta_n < \alpha < \min(c_{r_{n+1}} \setminus c_{r_n})$, then $r_{n+1} \Vdash \alpha \notin C_{\dot{G}}$, whence $r_{n+1} \Vdash \alpha \notin \tau$.

Similarly, if $\max(c_{p_{n+1}}) < \alpha < \delta_{n+1}$, then $p_{n+2} \Vdash \alpha \notin C_{\dot{G}}$, whence $p_{n+2} \Vdash \alpha \notin \tau$. Respectively, if $\max(c_{q_{n+1}}) < \alpha < \delta_{n+1}$, then $q_{n+2} \Vdash \alpha \notin C_{\dot{G}}$, whence $q_{n+2} \Vdash \alpha \notin \tau$.

Finally, if $\alpha \in I = [\min(c_{p_{n+1}}), \max(c_{p_{n+1}})]$, then p_{n+1} may leave the sentence $\alpha \in \tau$ undecided, but still $p_{n+1} \Vdash I \cap \tau \neq \emptyset$, since p_{n+1} decides $\text{dom}(\dot{f}_{\delta_n})$ and we have (2). Respectively, q_{n+1} forces $[\min(c_{q_{n+1}}), \max(c_{q_{n+1}})] \cap \tau \neq \emptyset$, and r_{n+1} forces $[\min(c_{r_{n+1}}), \max(c_{r_{n+1}})] \cap \tau \neq \emptyset$.

Let $p_{\omega} = \inf_{n} p_{n}$, $q_{\omega} = \inf_{n} q_{n}$, $r_{\omega} = \inf_{n} r_{n}$, and let $\delta = \sup\{\delta_{n} : n < \omega\}$. Let $G_{0} \subseteq \mathbb{P}_{\text{fast}}$ be generic over V[H] such that $p_{\omega} \in G_{0}$, $G_{1} \subseteq \mathbb{P}_{\text{fast}}$ generic over V[H] such that $q_{\omega} \in G_{1}$, and $G_{2} \subseteq \mathbb{P}_{\text{fast}}$ generic over V[H] such that $r_{\omega} \in G_{2}$. Lastly, let

$$X = \tau_{G_0 \times H} \cap \delta, \ Y = \tau_{G_1 \times H} \cap \delta, \ Z = \tau_{G_2 \times H} \cap \delta.$$

As $\Vdash_{\mathbb{P}_{\text{fast}}\times\mathbb{Q}} \tau \cap \delta \in L[\dot{A}]$ and $\dot{A}_{G_0\times H} = \dot{A}_H$, we have $V[G_0\times H] \models X \in L[A]$. By absoluteness, $V[H] \models X \in L[A]$. Similarly, $V[H] \models Y, Z \in L[A]$.

Now by construction, $(X \cup Y, Z)$ is an interlace and (X, Y, Z) codes a. Hence $a \in L[A]$.

We need another auxiliary lemma for the iteration:

Lemma 12. Assume G is \mathbb{P}_{fast} -generic over V, $\mathbb{R} \in V[G]$ is a σ -closed forcing, K is \mathbb{R} -generic over V[G], H is \mathbb{Q} -generic over V[G][K], $A \subseteq \omega_1$ is in V[H], and in V[G][K][H], there is a club $D \subseteq C_G$ such that $D \cap \alpha \in L[A]$ for all $\alpha < \omega_1$. Then such a club D must already exist in V[G][H].

Proof. Suppose $\dot{A} \in V$ is a \mathbb{Q} -name for A and $\dot{D} \in V[G]$ is an \mathbb{R} -name for a \mathbb{Q} -name for D. Suppose $\dot{F} \in V[G]$ is an \mathbb{R} -name for a \mathbb{Q} -name for a function $\omega_1 \to \omega_1$ listing the elements of \dot{D} in increasing order. W.l.o.g. \dot{D} is a \mathbb{R} -name $\langle \dot{f}_{\alpha} : \alpha < \omega_1 \rangle$ for a sequence of countable partial functions defined on ω_1 such that $\{\dot{f}_{\alpha}(\gamma) : \gamma \in \text{dom}(f_{\alpha})\}$ is a maximal antichain in \mathbb{Q} and $\dot{f}_{\alpha}(\gamma)$ forces $\dot{F}(\alpha) = \gamma$. Suppose (w.l.o.g.) the weakest condition in $\mathbb{R} \times \mathbb{Q}$ forces what is assumed about \dot{A} , \dot{F} , and \dot{D} . Since $\Vdash \dot{D} \subseteq C_{\dot{G}}$, we have $\Vdash \text{dom}(\dot{f}_{\alpha}) \subseteq C_{\dot{G}}$.

We shall define a descending sequence $(r_{\alpha})_{\alpha<\omega_1}$ in K. For a start, $r_0\in K$ can be arbitrary. Suppose $r_{\alpha}\in K$ has been defined already. Let $r_{\alpha+1}\leq r_{\alpha}$ such that $r_{\alpha+1}\in K$ and $r_{\alpha+1}$ decides $\mathrm{dom}(\dot{f}_{\beta})$ and $\dot{f}_{\alpha}(\gamma)$ for $\beta\leq\alpha$ and $\gamma\in\mathrm{dom}(\dot{f}_{\alpha})$. Let $g_{\alpha}\in V[G]$ such that $r_{\alpha+1}\Vdash\dot{f}_{\alpha}=g_{\alpha}$. Let \dot{S} be a \mathbb{Q} -name in V for a function $\omega_1\to\omega_1$ such that $g_{\alpha}(\gamma)\Vdash\dot{S}(\alpha)=\gamma$. Let $\dot{E}\in V$ be a \mathbb{Q} -name such that $\Vdash\dot{E}=\{\dot{S}(\alpha):\alpha<\omega_1\}$. Now

$$V[K][H] \models \dot{E}_H = \dot{D}_{K \times H} \wedge \dot{E}_H \cap \delta \in L[A],$$

whence $V[H] \models \dot{E}_H \cap \delta \in L[A]$ follows by absoluteness.

Now we can construct the iteration in such a way that after forcing with the iteration and then with \mathbb{Q} the ground model reals, which are the same as the reals after the iteration, are second order definable from any set $A \subseteq \omega_1$ with a certain second order property.

Lemma 13. We assume CH. Suppose \mathbb{P} is the countable support iteration of fast club forcing of length ω_2 . Let G be \mathbb{P} -generic over V. Suppose H is \mathbb{Q} -generic over V[G]. Suppose in V[G][H] there is a set $A \subseteq \omega_1$ such that for every club C, there is a club $D \subseteq C$ such that $D \cap \alpha \in L[A]$ for all $\alpha < \omega_1$. Then $P(\omega)^V \subseteq L[A]$.

Proof. Let $\mathbb{P} = \langle \mathbb{P}_{\alpha} : \alpha < \omega_2 \rangle$ be the countable support iteration of $\langle \dot{Q}_{\alpha} : \alpha < \omega_2 \rangle$, where $\mathbb{P}_{\alpha} \Vdash "\dot{Q}_{\alpha}$ is the fast club forcing \mathbb{P}_{fast} ". Let $G_{\alpha} = G \cap \mathbb{P}_{\alpha}$. Let $\dot{A} \in V[G]$ be an H-name for A. Choose β large enough such that $\dot{A} \in V[\langle G_{\alpha} : \alpha < \beta \rangle]$. Now G_{β} is \mathbb{P}_{fast} -generic over $V[\langle G_{\alpha} : \alpha < \beta \rangle]$. But, V[G] is a generic extension of $V[\langle G_{\alpha} : \alpha < \beta \rangle][G_{\beta}]$ by countably closed forcing and by assumption, in V[G][H], there is a club $D \subseteq C_{G_{\beta}}$ such that $D \cap \eta \in L[A]$ for all $\eta < \omega_1$. We apply Lemma 12 in $V[\langle G_{\alpha} : \alpha < \beta \rangle][G_{\beta}][H]$ such that $D \cap \alpha \in L[A]$ for all $\alpha < \omega_1$. By Lemma 11, $P(\omega)^V \subseteq L[A]$.

Theorem 14. There is a set of forcing conditions that forces the existence of a complete non-categorical finite second order theory with a model of cardinality \aleph_1 .

Proof. Assume w.l.o.g., CH. As said above, we start with some preparatory countably closed forcing \mathbb{P} obtaining a generic extension V[G]. Then we add \aleph_1 Cohen-reals obtaining a further generic extension V[G][H]. In this model we consider for every $x \subseteq \omega$ the model M_x as defined in (1). Clearly, the cardinality of M_x is \aleph_1 . We shall now show that if x is Cohen-generic over V[G], e.g. one of the \aleph_1 many coded by H, then the complete second order theory of M_x is finitely axiomatizable (in second order logic). To end the proof of the theorem, we show that if x and y are mutually Cohen-generic over V[G], then M_x and M_y are second order equivalent but non-isomorphic.

In order to use second order logic over ω_1 to talk about HC^V and Cohengenericity over V we need to be able to decide, by the means offered by second order logic, which reals in V[G][H] are in V (or, equivalently, in V[G]) and which are not. This is precisely the purpose of the preparatory forcing \mathbb{P} .

We denote the starting ground model by V and assume, w.l.o.g., that V satisfies CH. We let the preparatory forcing $\mathbb{P} = \langle \mathbb{P}_{\alpha} : \alpha < \omega_2 \rangle$ be the countable support iteration of $\langle \dot{Q}_{\alpha} : \alpha < \omega_2 \rangle$, where $\mathbb{P}_{\alpha} \Vdash "\dot{Q}_{\alpha}$ is the fast club forcing \mathbb{P}_{fast} ". Let G be \mathbb{P} -generic over V and $G_{\alpha} = G \cap \mathbb{P}_{\alpha}$. In V[G] we force with \mathbb{Q} a generic H. Note that $\aleph_1^{V[G][H]} = \aleph_1^V$ and $\mathcal{P}(\omega)^{V[G]} = \mathcal{P}(\omega)^V$. Working in V[G][H], let the second order sentence $\phi(R, E)$, where R is unary and E is binary, say in a model M:

- (1) E^M is a well-founded relation satisfying ZFC^- + "every set is countable". This should be also true when relativized to R^M .
- $(2) |M| = \aleph_1.$

- (3) If $P' \in R^M$ denotes (in M) the set $\operatorname{Fn}(\omega, 2, \omega)$ of conditions for adding one Cohen real, then there is $K \subseteq P'$ such that K is P'-generic over R^M and $M \models "V = R[K]"$.
- (4) If $a \subseteq \omega$ and the transitive collapse of M is N, then the following conditions are equivalent:
 - (a) $a \in \mathbb{R}^N$.
 - (b) If $A \subseteq \omega_1$ and for every club $C \subseteq \omega_1$ there is a club $D \subseteq C$ such that $D \cap \alpha \in L[A]$ for every $\alpha < \omega_1$, then $a \in L[A]$.

Note that we can express " $D \cap \alpha \in L[A]$ ", or equivalently " $\exists \beta(|\beta| = \aleph_1 \wedge D \cap \alpha \in L_{\beta}[A]$ ", in second order logic on M since second order logic gives us access to all structures of cardinality |M| (= \aleph_1).

Claim: The following conditions are equivalent in V[G][H]:

- (i) $M \models \phi(R, E)$.
- (ii) $M \cong M_x$ for some real x which is Cohen generic over V.
- Proof. (i) implies (ii): Suppose $M \models \phi(R, E)$. Let (N, U, \in) be the transitive collapse of (M, R^M, E^M) . By (3), there is r which is Cohen-generic over U and $N = HC^{U[r]}$. We show that $U = HC^V$. Suppose $a \in \mathcal{P}(\omega)^V$. We use condition (4) to demonstrate that $a \in U$. To this end, let A be as in (4b). By Lemma 13, $a \in L[A]$. Thus (4) implies $a \in U$. On the other hand, suppose $a \in (\mathcal{P}(\omega))^U$. We again use (4) to show that $a \in \mathcal{P}(\omega)^V$. Let $A \subseteq \omega_1$ code $([\omega_1]^\omega)^V$. If C is any club in V[G][H], then, since H is obtained by means of a CCC forcing, there is a club $D \subseteq C$ in V[G]. Now $D \cap \alpha \in V$, whence $D \cap \alpha \in L[A]$, for all $\alpha < \omega_1$. It follows that $a \in L[A]$. Since $A \in V$, we may conclude $a \in \mathcal{P}(\omega)^V$. Hence, $U = HC^V$ and r is Cohen-generic over V. We have proved (ii).
- (ii) implies (i): Suppose $(N, R^N, E^N) = (HC^{V[r]}, HC^V, \in)$, where r is $\operatorname{Fn}(\omega, 2, \omega)$ -generic over V. We show that $(N, R^N, E^N) \models \phi(R, E)$. Conditions (1) and (2) are trivially satisfied. Condition (3) holds by construction. To prove that condition (4) holds, suppose $a \subseteq \omega$ and let A be as in (4). By Lemma 13, $a \in L[A]$. Condition (4) and thereby the Claim is proved.

We continue the proof of Theorem 14. The sentence $\phi(R, E)$ is non-categorical in V[G][H] because if we take two mutually generic (over V[G])

Cohen reals r_0 and r_1 , then M_{r_0} and M_{r_1} are non-isomorphic models of $\phi(R, E)$. To prove that $\phi(R, E)$ is complete, suppose (M, R^M, E^M) and (N, R^N, E^N) are two models of $\phi(R, E)$. W.l.o.g., they are of the form (M, R^M, \in) and (N, R^N, \in) , where M and N are transitive sets. By construction, they are of the form M_{r_0} and M_{r_1} where both r_0 and r_1 are Cohen generic over HC^V , hence over $HC^{V[G]}$. They are subsumed by the generic H. By homogeneity of Cohen forcing $Fn(\omega, 2, \omega)$ the models are second order equivalent.

In fact the forcing gives something stronger. If κ is a cardinal that is second order characterizable in the forcing extension, we may replace the model $M_x = (HC^{V[x]}, HC^V, \in)$, where $x \subseteq \omega$ is Cohen over V, with the model $(\kappa \cup HC^{V[x]}, HC^V, \in)$, and the proof of Theorem 14 goes through mutatis mutandis:

Corollary 15. There is a set of forcing conditions that forces the following: if κ is any second order characterizable cardinal, there is a complete non-categorical finitely axiomatizable second order theory with a model of cardinality κ .

Since the non-isomorphic models above derive from mutually generic Cohen reals, it follows that the non-categorical theories in question have (at most) continuum many non-isomorphic models. We lastly mention how to get non-categorical theories with more models than this.

It is straightforward to see that in theorem 14 and the constructions preceding it, the cardinal \aleph_1 may be replaced with any cardinal μ^+ with μ regular. That is, the ω_2 -length countable support iteration of fast club forcing at ω_1 is replaced by a μ^{++} -length $\leq \mu$ -sized support iteration of fast club forcing at μ^+ , and the forcing to add \aleph_1 many Cohen subsets of ω is replaced by adding μ^+ many Cohen subsets of μ . The model M_x is then taken to be of the form $(H(\mu)^{V[x]}, H(\mu)^V, \in)$ where x is a Cohen subset of μ generic over V.

From this variation, we then get the following corollary.

Corollary 16. Suppose μ is a regular cardinal. There is then a set of forcing conditions that forces the following: if μ is second order characterizable, and if $\kappa \geq \mu$ is any second order characterizable cardinal, there is a complete non-categorical finite second order theory T with a model of cardinality κ . Also, the theory T has between μ^+ and 2^{μ} many models up to isomorphism.

Note that the concern of the second order characterizability of μ and κ in the forcing extension are irrelevant for cardinals with simple definitions such as \aleph_n , $n < \omega$ or \aleph_{ω_1+1} , for example.

In conclusion we cannot hope to prove the categoricity of finite complete second order theories from large cardinals even if we restrict to theories which have a model of regular uncountable cardinality.

6 Forcing categoricity

In [2] (for $\kappa > \omega_1$) and [3] (for $\kappa = \omega_1$), Aspero and Friedman proved the following:

Theorem 17. Suppose κ is the successor of a regular cardinal, and uncountable. Then there is a poset \mathbb{P} such that in a generic extension by \mathbb{P} , there is a lightface first order definable well-order of $H(\kappa^+)$.

Since we can translate a first order lightface definable well-order of $H(\kappa^+)$ into a well-order of $\mathcal{P}(\kappa)$ that is second order definable over any structure of cardinality κ , we obtain the following corollary.

Theorem 18. Suppose κ is the successor of a regular cardinal, uncountable, and that κ is second order characterizable. Then there is a poset \mathbb{P} that forces the following: every finitely axiomatizable second order theory with a model of cardinality κ is categorical.

We are thus left to consider the case of theories with models of limit cardinality, whether regular or singular.

The following theorem shows that the categoricity of complete second order theories with a model of singular cardinality is (relatively) consistent with large cardinals. We are indebted to Boban Veličković for suggesting how to improve an earlier weaker version of this result.

Theorem 19. Suppose κ is a singular strong limit with uncountable cofinality λ . Then there is a forcing notion \mathbb{P} of cardinality κ such that

- 1. \mathbb{P} preserves κ singular strong limit of uncountable cofinality λ .
- 2. \mathbb{P} forces the statement: Every finitely axiomatizable complete second order theory with a model of cardinality κ is categorical.

Proof. W.l.o.g. we assume GCH up to κ . We first force a second order definable well-order of the bounded subsets of κ with a reverse Easton type iteration of length κ described in [21, Theorem 20].

Let $e: \kappa \to \kappa$ be the function which lists the set B of beth fixed points $> \lambda$ in increasing order, and let $S = \langle \kappa_{\xi} : \xi < \lambda \rangle \subseteq B$ be an increasing cofinal sequence in κ such that $\kappa_0 > \lambda$. Let $\pi : \kappa \times \kappa \to \kappa$ be the Gödel pairing function. Let W be a well-order of V_{κ} . Suppose $A \subseteq \mu$, where $\mu \in B$. We write $A \sim V_{\mu}$ if

$$(V_{\mu}, \in) \cong (\mu, \{(\alpha, \beta) : \pi(\alpha, \beta) \in A\}).$$

Let the poset $E(\mu, A)$ be the iteration (product) of the posets \mathbb{R}_{α} , $\alpha < \mu$, where

$$\mathbb{R}_{\alpha} = \begin{cases} \operatorname{Fn}(\aleph_{\mu+\alpha+3} \times \aleph_{\mu+\alpha+1}, 2, \aleph_{\mu+\alpha+1}), & \text{if } \alpha = \omega \cdot \beta \text{ and } \beta \in A \\ \operatorname{Fn}(\aleph_{\mu+\alpha+4} \times \aleph_{\mu+\alpha+2}, 2, \aleph_{\mu+\alpha+2}), & \text{if } \alpha = \omega \cdot \kappa_{\xi} + 1, \, \xi < \lambda \\ (\{0\}, =) & \text{otherwise} \end{cases}$$

with Easton support i.e. $E(\mu, A)$ consists of functions $p \in \prod_{\alpha < \mu} \mathbb{R}_{\alpha}$ such that, denoting the support $\{\alpha : f(\alpha) \neq \emptyset\}$ of f by $\operatorname{supp}(p)$, $|\operatorname{supp}(p) \cap \gamma| < \gamma$ for all regular γ .

We now define an iteration $\langle \mathbb{P}_{\alpha} : \alpha < \kappa \rangle$ with the property that \mathbb{P}_{α} does not change beth fixed points $\beta = \beth_{\beta}$ for any β . We let $\mathbb{P} = \langle \mathbb{P}_{\alpha} : \alpha < \kappa \rangle$ be the following iteration: If α is a limit ordinal, we use direct limits for regular α and inverse limits for singular α . Suppose then $\alpha = \beta + 1$. Let \dot{A} be the W-first \mathbb{P}_{β} -name \dot{A} in V_{κ} such that $\mathbb{P}_{\beta} \Vdash \dot{A} \sim V_{\check{e}(\check{\beta})}$. Then $\mathbb{P}_{\alpha} = \mathbb{P}_{\beta} \star E(\check{e}(\check{\beta}), \dot{A})$. Let G be \mathbb{P} -generic over V and $G_{\alpha} = G \cap \mathbb{P}_{\alpha}$.

In the forcing extension V[G], for every $\mu \in B$ there is a set $A \subseteq \mu$ which codes, via the canonical bijection $\pi : \kappa \times \kappa \to \kappa$, a bijection $f_A : \mu \to (V_\mu)^{V[G]}$. The set A itself satisfies

$$V[G] \models A = \{\alpha < \mu : 2^{\aleph_{\mu + \omega \cdot \alpha + 1}} = \aleph_{\mu + \omega \cdot \alpha + 3}\}$$

and from A we can read off f_A and a well-order $<_{\mu}^*$ of $(V_{\mu})^{V[G]}$:

$$V[G] \models f_A(\alpha) <^*_{\mu} f_A(\beta) \iff \alpha < \beta < \mu.$$

Now working in V[G], fix a collection $\mathcal{F} \subseteq \mathcal{P}(\kappa)$, and we set out to define a well-order not on the whole of \mathcal{F} but a certain subset of it. Define a relation R on \mathcal{F} by

$$XRY \iff X \cap \kappa_{\xi} <_{\kappa_{\xi}}^{*} Y \cap \kappa_{\xi} \text{ for all but boundedly many } \xi < \lambda.$$

As λ is uncountable, R is well-founded, so the set

$$\mathcal{W} = \{ X \in \mathcal{F} : X \text{ is minimal in } R \}$$

is nonempty, and if $X, Y \in \mathcal{W}$ with $X \neq Y$, then both $X \cap \kappa_{\xi} <_{\kappa_{\xi}}^{*} Y \cap \kappa_{\xi}$ and $Y \cap \kappa_{\xi} <_{\kappa_{\xi}}^{*} X \cap \kappa_{\xi}$ occur for unboundedly many $\xi < \lambda$.

To see that $|\mathcal{W}| < \kappa$, suppose to the contrary that $|\mathcal{W}| \ge \kappa$ and define a coloring $c : [\mathcal{W}]^2 \to \lambda$ by $c(\{X,Y\}) = \pi(\xi_1,\xi_2)$ where ξ_1 is the least $\xi < \lambda$ such that $X \cap \kappa_{\xi} <_{\kappa_{\xi}}^* Y \cap \kappa_{\xi}$, and ξ_2 is the least $\xi < \lambda$ such that $Y \cap \kappa_{\xi} <_{\kappa_{\xi}}^* X \cap \kappa_{\xi}$. Since $|\mathcal{W}| \ge \kappa > (2^{\lambda})^+$, by the Erdös-Rado theorem there is a set $H \subseteq \mathcal{W}$ homogeneous for c of color $\pi(\xi_1, \xi_2)$ and cardinality λ^+ . But this is a contradiction, since ordering H in $<_{\kappa_{\xi_1}}^*$ -increasing order yields an infinite decreasing sequence in the well-order $<_{\kappa_{\xi_2}}^*$, so $|\mathcal{W}| < \kappa$.

Now for each $X \in \mathcal{W}$, define $f_Y : \lambda \to \kappa$ such that $f_X(\xi)$ is the index of $X \cap \kappa_{\xi}$ in the well-order $<_{\kappa_{\xi}}^*$. Then the set $\bigcup \{\operatorname{ran}(f_X) : X \in \mathcal{W}\}$ has some cardinality $\gamma < \kappa$, and we can let $h : \bigcup \{\operatorname{ran}(f_X) : X \in \mathcal{W}\} \to \gamma$ be the transitive collapse map.

Then for $X \in \mathcal{W}$, the function $h \circ f_X : \lambda \to \gamma$ can be encoded as a subset of a large enough $\mu \in B$, and obviously $h \circ f_X \neq h \circ f_Y$ if $X \neq Y$, so we can well-order \mathcal{W} by

$$X \lhd Y \iff h \circ f_X <^*_{\mu} h \circ f_Y$$

and all this is second order definable in V[G] in a structure of size κ , if the collection \mathcal{F} is. This allows us to pick a distinguished element of \mathcal{F} as the \triangleleft -least R-minimal element.

Suppose now that ϕ is a complete second order sentence with a model M of cardinality κ , and let \mathcal{F} consist of the set of $X \subseteq \kappa$ encoding a model of ϕ . Note that over a model of cardinality κ we can write a formula $\phi_R(X,Y)$ expressing XRY for $X,Y \in \mathcal{F}$, a formula $\phi_{\mathcal{W}}(X)$ expressing $X \in \mathcal{W}$, and a formula $\phi_{\lhd}(X,Y)$ expressing $X \lhd Y$ if X and Y are R-minimal.

Let $M \models \Phi$ now say that $X \subseteq M$ encodes a model isomorphic to M (and thus satisfies ϕ), and for any $Y \subseteq M$ that also encodes a model of ϕ , $\neg \phi_R(Y, X)$, and moreover if for all $Z \subseteq M$ that encode a model of ϕ also $\neg \phi_R(Z, Y)$, then X = Y or $\phi_{\lhd}(X, Y)$. That is, $X \in \mathcal{W}$ and if also $Y \in \mathcal{W}$ then X = Y or $X \triangleleft Y$, which uniquely specifies X. As the model of ϕ with the least code in this sense satisfies Φ and ϕ is complete, ϕ implies Φ and thus that all models of ϕ are isomorphic, so ϕ is categorical. \square

The method of the preceding proof does not extend to the cases of the limit cardinal κ being regular, or of countable cofinality, so these cases are left open.

In conclusion, no known large cardinal axiom (e.g. the existence of huge cardinals) can decide whether all complete second order theories with a model of singular cardinality are categorical. In particular, such axioms cannot imply that all finite complete second order theories are categorical.

7 Theories with only countably many models

Since under PD we have non-categorical complete recursively axiomatized second order theories, we may ask how badly categoricity can fail in those cases? Echoing Vaught's Conjecture, we may ask whether the number of countable non-isomorphic models of a complete recursively axiomatized second order theory is always countable or 2^{ω} . Leaving this question unresolved, we have the following result which demonstrates the ability of categorical theories to 'capture' (in the sense of [23]) the models of non-categorical theories.

Theorem 20. Assume $AD^{L(\mathbb{R})}$. If T is a recursively axiomatized complete second order theory with only countably many non-isomorphic countable models, then there is a recursively axiomatized categorical second order theory S the unique model of which interprets all the countable models of T.

Proof. Let T be a recursively axiomatized second order theory with only countably many non-isomorphic countable models. Let A be the Π^1_{ω} (i.e. an intersection of a recursively coded family of sets each of which is Π^1_n for some n) set of reals that code a model of T. Since A is a countable union of equivalence classes of the Σ^1_1 -equivalence relation of isomorphism, we may conclude that A is Σ^1_1 .

We wish to show that A is $\Pi_2^1(r_0)$ in a parameter r_0 which is a Π_ω^1 singleton. For this, we mimic a proof of Louveau (Theorem 1 in [17]) to show:

Theorem 21. Assume $AD^{L(\mathbb{R})}$. Every Σ_1^1 set which is Π_{ω}^1 is $\Pi_2^1(r_0)$ for some real r_0 such that $\{r_0\}$ is a $\Delta_{\omega+1}^1$ -singleton.

Proof. Let A be a Σ_1^1 set that is also Π_{ω}^1 , say $A = \bigcap_n A_n$ with each A_n being Π_n^1 . Let also $U \subseteq (\omega^{\omega})^2$ be a universal Σ_1^1 set.

We define for each n a game G_n on ω where players I and II take turns to play the digits of reals α and γ respectively (there is no need to let II

pass turns here). Then II wins a play of G_n if $\alpha \in A \implies \gamma \in U$ and $\alpha \notin A_n \implies \gamma \notin U$.

As in Louveau's proof, II has a winning strategy as follows: since A is Σ_1^1 , we have $A(x) \iff U(y,x)$ for some y, so II wins by playing the digits of $\langle y,\alpha\rangle$ (as I is playing the digits of α). The complexity of the winning set for II in G_n is Σ_{ω}^1 , so by Moschovakis's strategic basis theorem ([22], Theorem 6E.2), II has a winning strategy σ_n that is a $\Delta_{\omega+1}^1$ -singleton. Note that the pointclass Σ_{ω}^1 , i.e. the collection of countable unions of recursively coded families of projective sets, is both adequate and scaled (see Remark 2.2 in [24], essentially Theorem 2.1 in [27]).

Then the set $B_n = \{y \mid (y * \sigma_n)_{II} \in U\}$ is a $\Sigma_1^1(\sigma_n)$ set with $A \subseteq B_n \subseteq A_n$ (where $(y * \sigma_n)_{II}$ denotes the real γ the strategy σ_n produces as I plays $\alpha = y$), so altogether $A = \bigcap_n B_n$ is a $\Pi_2^1(s_0)$ set where $s_0 = \langle \sigma_n \mid n < \omega \rangle$ is a $\Delta_{\omega+1}^1$ -singleton.

We may reduce the complexity of the parameter down to being a Π^1_{ω} singleton by the following theorem of Rudominer:

Theorem 22 (Rudominer [24]). Assume $AD^{L(\mathbb{R})}$. Then every real s_0 which is a $\Sigma^1_{\omega+1}$ singleton, is recursive in a real r_0 which is a Π^1_{ω} singleton.

Therefore the set A is a $\Pi_2^1(r_0)$ set where r_0 is a Π_ω^1 singleton. Let $\eta(r,s)$ be a second order Π_2^1 formula which defines the predicate $s \in A$ on $(\mathbb{N}, +, \times, r_0)$. Let $\theta_1(Q_+, Q_\times)$ be the standard second order characterization of $(\mathbb{N}, +, \times)$, as above in the proof of Theorem 3. Let $\psi_n(Q_+, Q_\times, s)$, $n < \omega$, be second order formulas such that if X_n is the set of reals s satisfying $\psi_n(Q_+, Q_\times, s)$ in $(\mathbb{N}, +, \times)$, then $\{r_0\} = \bigcap_n X_n$. Let P be a new unary predicate symbol and

$$S = \{\theta_1(Q_+, Q_\times)\} \cup \{\psi_n(Q_+, Q_\times, P) : n < \omega\}.$$

Suppose M is a model of S. W.l.o.g. the arithmetic part of M consists of the standard + and \times on \mathbb{N} . Let s be the interpretation of P in M. Then $s = r_0$. Thus S is categorical. The theory S is recursive because the proofs of Theorems 21 and 22 are sufficiently uniform. In conclusion, M is categorically characterized by the recursive second order theory S.

Now the countable models of T are interpretable in S in the following sense: a real s codes a model of T if and only if $M \models \eta(r_0, s)$. We also get a translation of sentences: if ϕ is a second-order sentence in the vocabulary of T, letting $\hat{\phi}$ be the sentence $\exists X(\eta(r_0, X) \land X \models \phi)$, we have that $\phi \in T$ if and only if $\hat{\phi} \in S$.

8 Definable models of categorical theories

Suppose we are given a categorical second order theory T. Naturally, we assume that T has a model, otherwise categoricity is vacuous. But what can be said about the models of T apart from their isomorphism with each other? In particular, can we always find a model which is definable in some reasonable sense, e.g. hereditarily ordinal definable? To emphasize this point, consider the second order sentence which characterizes the structure $(\mathbb{N}, +, \cdot, 0^{\sharp})$. This categorical sentence has no models in L. We ask, can we have a categorical sentence with no models in HOD? Since it could be that V = HOD, we are looking at this question under assumptions stronger than ZFC.

The following result of Kaplan and Shelah is useful for us:

Theorem 23 ([13]). If \mathbb{P} forces the collapse of $|\omega_2|$ to ω , then there is a \mathbb{P} -term τ for a countable model such that

1. If $G_1 \times G_2$ is generic for $\mathbb{P} \times \mathbb{P}$ then

$$V[G_1][G_2] \models M_1 \cong M_2,$$

where M_1 is the interpretation τ^{G_1} of τ by G_1 and M_2 is τ^{G_2} .

2. $\mathbb{P} \Vdash \text{``} \tau \text{ is not isomorphic to } \check{M}\text{''}, \text{ for any } M \text{ in } V.$

We make some observations about the proof. It involves a construction of Laskowski and Shelah:

Theorem 24 ([16]). There is a countable consistent first order theory T, with a predicate V in its vocabulary, having the following property. For any model $M \models T$ and any $A \subseteq V^M$, isolated types are dense over A but the theory $T(A) = \text{Th}(M, a)_{a \in A}$ has an atomic model if and only if $|A| < \omega_2$.

The theory T is as follows. Let L be a countable vocabulary consisting of two unary predicates U, V, one unary function symbol p, as well as binary

relations R_n and binary functions f_n for $n < \omega$ (the functions will not be total, but instead have domain U). Let K be the collection of all finite L-structures satisfying a certain finite list of first order axioms (see [16]). Let \mathcal{B} be the Fraïsse limit of K and let $T = \text{Th}(\mathcal{B})$. The theory T is well defined since \mathcal{B} is unique up to isomorphism.

We then form an uncountable model of the theory T as follows. For an ordinal α let L_{α} be the vocabulary L together with α many new constant symbols c_{β} , $\beta < \alpha$. Using a standard Henkin construction, we form a term model for the theory T together with the additional axioms stating that the new constant symbols name distinct elements. We let $T(A_{\alpha})$ be the theory of this term model in the vocabulary L_{α} . (Although the Henkin construction involves forming the completion of a theory, we can make the choice of which completion to use definable by referring to the well-ordering of the sentences.)

We can also observe that for a countable ordinal α , the class of countable atomic models of $T(A_{\alpha})$ is definable from $T(A_{\alpha})$, which itself is definable from α , and the definitions can be carried out in $H(\omega_1)$. Using these two observations, the following obtains:

Theorem 25 (ZF). Assume ω_2^{HOD} is countable. Then there is a countable model M such that

- 1. The isomorphism class of M is ordinal definable.
- 2. There is no model in HOD which is isomorphic to M.

Moreover, if the property of a linear order of being of order-type ω_2^{HOD} is second order definable in the countably infinite structure of the empty vocabulary, then the second order theory of M is finitely axiomatizable.

Proof. Let $\alpha = \omega_2^{\text{HOD}}$. Let $T(A_{\alpha})$ be the theory constructed above. Finally, let M be a countable atomic model of $T(A_{\alpha})$. Since HOD satisfies $|T(A_{\alpha})| = \omega_2$, the theory $T(A_{\alpha})$ has no atomic model in HOD, but as being an atomic model is absolute, this shows that there is no model in HOD isomorphic to M.

The isomorphism class of M is ordinal definable as the class of countable atomic models of $T(A_{\alpha})$, which is definable from α . Additionally, if α is second order definable in the countably infinite structure of the empty vocabulary, we can define the theories T and $T(A_{\alpha})$ in second order logic expressing "I am isomorphic to a countable atomic model of $T(A_{\alpha})$ " with a single second order sentence. This finitely axiomatizes the second order theory of M.

Of course, the assumption that ω_2^{HOD} is second order definable in the countably infinite structure of the empty vocabulary is somewhat ad hoc. However, it holds, for example, in L[G], where G is P-generic over L for $P = \text{Coll}(\omega, <\omega_3)^L$. This is because the poset P is weakly homogeneous, so $\text{HOD}^{L[G]} = \text{HOD}^L(P) = L$, whence $\omega_2^{\text{HOD}} = \omega_2^L$ is countable and second order definable in any countable model in L[G].

We also obtain the following variation:

Corollary 26. Assume $ZFC + AD^{L(\mathbb{R})} + \text{``}HOD \cap \mathbb{R} = HOD^{L(\mathbb{R})} \cap \mathbb{R}$ " and that ω_2^{HOD} is definable in $HOD^{L(\mathbb{R})} \upharpoonright \Theta^{L(\mathbb{R})}$ and countable. Let M be the countable model of Theorem 25. Let $N = (\Theta^{L(\mathbb{R})}, <, M)$ (w.l.o.g. the domain of M is ω). Then the second order theory of N is finitely axiomatizable and categorical but has no model which belongs to HOD.

Proof. We can use [7, Theorem 3.10, Chapter 23]) to define $\mathrm{HOD}^{L(\mathbb{R})} \upharpoonright \Theta^{L(\mathbb{R})}$ and $L_{\Theta^{L(\mathbb{R})}}(\mathbb{R})$ from $\Theta^{L(\mathbb{R})}$ in second order logic, which then allows us to define ω_2^{HOD} and M as in Theorem 25.

The assumptions of Corollary 26 follow, for example, from $ZFC+AD^{L(\mathbb{R})}+V=L(\mathbb{R})[G]$, where G is \mathbb{P}_{\max} -generic, as then $HOD^{L(\mathbb{R})}=HOD^{L(\mathbb{R})[G]}$ and ω_2^{HOD} is countable.

9 Open questions

The following question was raised by Solovay [26]:

Open Problem 1. Assuming V = L, is every recursively axiomatized complete second order theory categorical?

Our results do not solve this one way or another, and it remains an interesting open question. In L[U] there are recursively axiomatized complete non-categorical second order theories, but we do not know if such theories necessarily have only large models:

Open Problem 2. Suppose V = L[U], κ is the sole measurable cardinal of L[U], and T is a complete recursively axiomatized second order theory that has a model of cardinality $\lambda < \kappa$ such that λ is second order characterizable. Is T categorical?

There are many other open questions related to finite or recursively axiomatized complete second order theories with uncountable models. We showed that we can force categoricity for successor cardinals of regular cardinals, and some singular limit cardinals, but the following two cases were left open:

Open Problem 3. Can we always force the categoricity of all finite complete second order theories with a model of cardinality κ , where κ is either a regular (non-measurable) limit cardinal, or singular of cofinality ω ?

An I_0 -cardinal is a cardinal λ such that there is $j: L(V_{\lambda+1}) \to L(V_{\lambda+1})$ with critical point below λ . Note that then λ is singular of cofinality ω , λ^+ is measurable in $L(V_{\lambda+1})$ ([29]), and the Axiom of Choice fails in $L(V_{\lambda+1})$ ([15]). This is in sharp contrast to the result of Shelah that if λ is a singular strong limit cardinal of uncountable cofinality, then $L(\mathcal{P}(\lambda))$ satisfies the Axiom of Choice ([25]). Since Axiom of Choice fails in $L(V_{\lambda+1})$, there can be no well-order of $\mathcal{P}(\lambda)$ which is second order definable on λ . This raises the following question:

Open Problem 4. Is every finite complete second order theory with a model of cardinality of an I_0 -cardinal categorical (or, at least categorical among all models of that cardinality)?

References

- [1] Miklós Ajtai. Isomorphism and higher order equivalence. Annals of Mathematical Logic, 1979.
- [2] David Asperó and Sy-David Friedman. Large cardinals and locally defined well-orders of the universe. *Ann. Pure Appl. Logic*, 157(1):1–15, 2009.
- [3] David Asperó and Sy-David Friedman. Definable well-orders of $H(\omega_2)$ and GCH. J. Symbolic Logic, 77(4):1101–1121, 2012.
- [4] David Asperó and Ralf Schindler. Martin's Maximum⁺⁺ implies Woodin's axiom (*). Ann. of Math. (2), 193(3):793–835, 2021.
- [5] Steve Awodey and Erich H. Reck. Completeness and categoricity. I. Nineteenth-century axiomatics to twentieth-century metalogic. *Hist. Philos. Logic*, 23(1):1–30, 2002.

- [6] Rudolf Carnap. *Untersuchungen zur allgemeinen Axiomatik*. Wissenschaftliche Buchgesellschaft, Darmstadt, 2000. Edited and with a foreword by Thomas Bonk and Jesus Mosterin.
- [7] M. Foreman and A. Kanamori. *Handbook of Set Theory*. Springer Netherlands, 2009.
- [8] Abraham Fraenkel. Einleitung in die Mengenlehre. 3. Aufl., volume 9. Springer, Berlin, 1928.
- [9] Roland Fraïssé. Sur les types de polyrelations et sur une hypothèse d'origine logistique. C. R. Acad. Sci. Paris, 230:1557–1559, 1950.
- [10] Roland Fraïssé. Sur la signification d'une hypothèse de la théorie des relations, du point de vue du calcul logique. C. R. Acad. Sci. Paris, 232:1793–1795, 1951.
- [11] William P Hanf and Dana Scott. Classifying inaccessible cardinals. *Notices of the American mathematical Society*, 8:445, 1961.
- [12] Akihiro Kanamori. The higher infinite: large cardinals in set theory from their beginnings. Springer monographs in mathematics. Springer, Berlin; 2nd ed. edition, 2003.
- [13] Itay Kaplan and Saharon Shelah. Forcing a countable structure to belong to the ground model. *MLQ Math. Log. Q.*, 62(6):530–546, 2016.
- [14] Eugene M. Kleinberg. Infinitary combinatorics and the axiom of determinateness, volume Vol. 612 of Lecture Notes in Mathematics. Springer-Verlag, Berlin-New York, 1977.
- [15] Kenneth Kunen. Elementary embeddings and infinitary combinatorics. J. Symbolic Logic, 36:407–413, 1971.
- [16] M. C. Laskowski and S. Shelah. On the existence of atomic models. *J. Symbolic Logic*, 58(4):1189–1194, 1993.
- [17] Alain Louveau. Borel sets and the analytical hierarchy. In *Proceedings of the Herbrand symposium (Marseilles, 1981)*, volume 107 of *Stud. Logic Found. Math.*, pages 209–215. North-Holland, Amsterdam, 1982.

- [18] Wiktor Marek. Consistance d'une hypothèse de Fraïssé sur la définissabilité dans un langage du second ordre. C. R. Acad. Sci. Paris Sér. A-B, 276:A1147–A1150, 1973.
- [19] Wiktor Marek. Sur la consistance d'une hypothèse de Fraïssé sur la définissabilité dans un langage du second ordre. C. R. Acad. Sci. Paris Sér. A-B, 276:A1169–A1172, 1973.
- [20] Donald A. Martin. The axiom of determinateness and reduction principles in the analytical hierarchy. Bull. Amer. Math. Soc., 74:687–689, 1968.
- [21] Telis K. Menas. Consistency results concerning supercompactness. Trans. Amer. Math. Soc., 223:61–91, 1976.
- [22] Yiannis N. Moschovakis. *Descriptive set theory*, volume 155 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, second edition, 2009.
- [23] Michael O. Rabin. A simple method for undecidability proofs and some applications. In *Logic*, *Methodology and Philos*. *Sci.* (*Proc.* 1964 Internat. Congr.), pages 58–68. North-Holland, Amsterdam, 1965.
- [24] Mitch Rudominer. The mouse set theorem just past projective. Journal of Mathematical Logic, 0(0):2450014, 0.
- [25] Saharon Shelah. Set theory without choice: not everything on cofinality is possible. Arch. Math. Logic, 36(2):81–125, 1997.
- [26] Robert Solovay. FOM posting, 2006. http://cs.nyu.edu/pipermail/fom/2006-May/010561.html.
- [27] John R. Steel. Scales in $L(\mathbf{R})$. In Cabal seminar 79–81, volume 1019 of Lecture Notes in Math., pages 107–156. Springer, Berlin, 1983.
- [28] W. Hugh Woodin. The axiom of determinacy, forcing axioms, and the nonstationary ideal, volume 1 of De Gruyter Series in Logic and its Applications. Walter de Gruyter GmbH & Co. KG, Berlin, revised edition, 2010.
- [29] W. Hugh Woodin. Suitable extender models II: beyond ω -huge. J. Math. Log., 11(2):115–436, 2011.

[30] W. Hugh Woodin. In search of Ultimate-L: the 19th Midrasha Mathematicae Lectures. Bull. Symb. Log., 23(1):1–109, 2017.