A note on the minimax risk of sparse linear regression

Yilin Guo¹, Shubhangi Ghosh¹, Haolei Weng³, and Arian Maleki¹

¹Columbia University ²Michigan State University **Abstract**

Sparse linear regression is one of the classical and extensively studied problems in high-dimensional statistics and compressed sensing. Despite the substantial body of literature dedicated to this problem, the precise determination of its minimax risk remains elusive. This paper aims to fill this gap by deriving asymptotically constant-sharp characterization for the minimax risk of sparse linear regression. More specifically, the paper focuses on scenarios where the sparsity level, denoted as k, satisfies the condition $(k \log p)/n \to 0$, with p and p representing the number of features and observations respectively. We establish that the minimax risk under isotropic Gaussian random design is asymptotically equal to $2\sigma^2 k/n \log(p/k)$, where σ denotes the standard deviation of the noise. In addition to this result, we will summarize the existing results in the literature, and mention some of the fundamental problems that have still remained open.

1 Sparse linear regression and minimaxity

Consider the linear regression model

$$y_i = x_i^T \beta + z_i, \qquad i = 1, \dots, n, \tag{1}$$

in which $y_i \in \mathbb{R}$ denotes the response, $x_i \in \mathbb{R}^p$ represents the feature or covariate vector, $\beta \in \mathbb{R}^p$ is the unknown signal vector to be estimated, and finally $z_1, \ldots, z_n \stackrel{i.i.d.}{\sim} \mathcal{N}(0, \sigma^2)$ are normal errors. We are interested in studying this problem for a broad range of p, considering p comparable with n, or even much larger than n. To ease one of the major concerns that linear regression procedures remain inconsistent unless $p/n \to 0$, following the rich literature of sparse linear regression [17, 6, 16, 33, 13], we consider the sparsity structure of the signal in this paper. Specifically, we assume that the true regression coefficients are k-sparse:

$$\beta \in \Theta_k := \{ \beta \in \mathbb{R}^p : \|\beta\|_0 \le k \}, \tag{2}$$

where $\|\beta\|_0$ denotes the number of non-zero components of β . In evaluating the performance of estimators, the minimax framework has been one of the most popular approaches, aiming to obtain an optimal estimator which has the best worst-case performance among all estimators. In order to define the minimax risk, the first step is to consider a model for the design matrix $X \in \mathbb{R}^{n \times p}$. Two models have been considered in the literature for matrix X:

 \bullet Fixed design: In this model, matrix X is regarded as a fixed entity, and no probabilistic assumptions are imposed upon it. Under this framework, the minimax risk is defined as

$$R_F(X, \Theta_k, \sigma) := \inf_{\hat{\beta}} \sup_{\beta \in \Theta_k} \mathbb{E}_{\beta} \|\hat{\beta} - \beta\|_2^2, \tag{3}$$

where $\|\cdot\|_2$ is the Euclidean norm. Note that in the above expression the expectation is with respect to the noise vector $z=(z_1,\ldots,z_n)$ only.

ullet Random design: In this model, X is presumed to be generated via a known probabilistic mechanism. One common random design model posits that the rows of matrix X are independent and identically distributed from a Gaussian distribution. In this case, the minimax risk is defined as

$$R_R(\Theta_k, \sigma) := \inf_{\hat{\beta}} \sup_{\beta \in \Theta_k} \mathbb{E}_{\beta} ||\hat{\beta} - \beta||_2^2, \tag{4}$$

where the expectation is with respect to both X and z.

Many researchers have considered characterizing the above two minimax risks for the sparse linear regression problem. However, obtaining the exact minimax risk is mathematically challenging and has remained open. Hence, researchers have explored approaches that aim to approximate the minimax risk. Below we summarize the existing approaches in the literature. On a related note, the minimax risk under other types of loss functions such as ℓ_q -norm loss $(q \ge 1)$ [11, 34, 1], prediction loss [24, 32, 10] and Hamming loss [19, 7, 22], has been also studied in the literature. The current paper is focused on the minimax risk with the squared loss as defined in (3) and (4).

2 Approximation of minimax risk

In this section, we would like to discuss three major approaches that researchers have explored for approximating the minimax risk.

2.1 Rate-optimal results under fixed design

Given the complexity involved in precisely calculating the minimax risk, numerous studies have tried to obtain the relationship between $R_F(X, \Theta_k, \sigma)$ and parameters such as k, n, and p. In this approach, the aim is often to obtain a function of k, n, and p, call it $f_F(n, p, k)$, for which there exist two constants c and C such that

$$cf_F(n, p, k)\sigma^2 \le R_F(X, \Theta_k, \sigma) \le Cf_F(n, p, k)\sigma^2.$$
 (5)

As evident, the constant in the lower and upper bounds may be different. Under this setting, once $f_F(n, p, k)$ is characterized, an estimator $\hat{\beta}$ is called minimax rate-optimal if

$$\sup_{\beta \in \Theta_k} \mathbb{E}_{\beta} \|\hat{\beta} - \beta\|_2^2 \le \tilde{C} f_F(n, p, k) \sigma^2.$$

Again, \tilde{C} can be different from C that appeared in the upper bound of the minimax risk. One important line of research on sparse linear regression, has been devoted to characterizing $f_F(n, p, k)$ and designing minimax rate-optimal estimators.

The minimax risk $R_F(X, \Theta_k, \sigma)$ generally depends on the design matrix X. For instance, if a few columns of X are linearly dependent, estimating the true β becomes impossible and $R_F(X, \Theta_k, \sigma)$ equals infinite. Hence, certain aspects of X need to appear in the upper bound (and lower bound). Some important conditions on X have been proposed in the literature, including restricted isometry property [9], compatibility condition [14] and restricted eigenvalue (RE) condition [3]. We adopt a slightly stronger version of the RE condition from [1] for later discussion.

Definition 1. The matrix X is said to satisfy the $SRE(k, c_0)$ condition for a given $c_0 > 0$ if its jth column is normalized $||X_j||_2 \le \sqrt{n}$ for $j = 1, \ldots, p$, and

$$\theta(k, c_0) := \min_{\delta \in \mathcal{C}_{SRE}(k, c_0) \setminus \{0\}} \frac{\|X\delta\|_2}{\sqrt{n} \|\delta\|_2} > 0, \tag{6}$$

where
$$C_{SRE}(k, c_0) := \{ \delta \in \mathbb{R}^p : \|\delta\|_1 \le (1 + c_0) \sqrt{k} \|\delta\|_2 \}$$
 is a cone in \mathbb{R}^p

According to this definition, if any k columns of the matrix X are linearly dependent (consequently the linear model is not identifiable on Θ_k), then $\theta(k, c_0) = 0$. Hence, the condition $\theta(k, c_0) > 0$ ensures model identifiability. Moreover, it essentially requires that the Hessian for quadratic cost function has a positive curvature in directions over the cone $C_{SRE}(k, c_0)$, which has implications for well-controlled estimation errors. To see how such a condition can be used to develop an upper bound for the minimax risk, consider the Lasso estimator [28]:

$$\hat{\beta}^L \in \underset{b \in \mathbb{R}^p}{\operatorname{arg\,min}} \, \frac{1}{2n} \|y - Xb\|_2^2 + \lambda \|b\|_1,$$

where $y = (y_1, ..., y_n)$ is the response vector and $\|\cdot\|_1$ is the ℓ_1 -norm. The following theorem taken from [1] obtains an upper bound for $\mathbb{E}_{\beta} \|\hat{\beta}^L - \beta\|_2^2$ based on the SRE condition.

Theorem 1 (Corollary 4.4 in [1]). Assume that X satisfies the SRE(k,7) condition. Let $\hat{\beta}^L$ denote the Lasso estimator with λ satisfying

$$\lambda \geq (8 + 2\sqrt{2})\sigma\sqrt{\frac{\log(2ep/k)}{n}}.$$

Then,

$$R_F(X, \Theta_k, \sigma) \le \sup_{\beta \in \Theta_k} \mathbb{E}_\beta \|\hat{\beta}^L - \beta\|_2^2 \le \frac{2401k\lambda^2}{64} \left(\frac{1}{\theta^4(k, 7)} + \frac{1}{(\log(2ep))^2} \right). \tag{7}$$

Based on our earlier discussion of $\theta(k,7)$ as a curvature-type condition, it is natural to see that $\theta(k,7)$ appears in the denominator of the upper bound. Following [1], we mention two significant features of this upper bound:

- Choosing $\lambda = (8+2\sqrt{2})\sigma\sqrt{\frac{\log(2ep)}{n}}$, the upper bound can be simplified to $C_{\theta}\frac{\sigma^2k\log p}{n}$ for some constant $C_{\theta} > 0$ depending on $\theta(k,7)$ whenever $p \geq 2$. Although upper bounds of this form are known from past work for $\|\hat{\beta}^L \beta\|_2^2$ (with high probability) [3, 34, 23], Theorem 1 is the first result to obtain such bounds in expectation, i.e. for $\mathbb{E}_{\beta}\|\hat{\beta}^L \beta\|_2^2$. The previous work provided only bounds in probability with the confidence level tied to the tuning parameter λ , thus not allowing for control of the moments of $\|\hat{\beta}^L \beta\|_2^2$. Similar bounds in probability have been obtained for several other estimators such as Dantzig selector [3] and square-root Lasso [2].
- Choosing $\lambda = (8+2\sqrt{2})\sigma\sqrt{\frac{\log(2ep/k)}{n}}$, the upper bound is improved to $\tilde{C}_{\theta}\frac{\sigma^2k\log(p/k)}{n}$ (with a different constant \tilde{C}_{θ}). In light of a minimax lower bound to be shown shortly, the rate $\frac{k\log(p/k)}{n}$ is the minimax optimal rate. The fact that Lasso can achieve the optimal rate (not just the suboptimal rate $\frac{k\log p}{n}$) was not known before the work $[1]^1$.

To evaluate the tightness of the upper bound in Theorem 1, we present one lower bound for the minimax risk taken from [32].

¹The optimal rate was known to be attained by ℓ_0 -constrained least squares under a sparse eigenvalue condition [24], although the bound was derived in probability instead of in expectation.

Theorem 2 (Proposition 6.2 in [32]). Suppose each column of X has been normalized to \sqrt{n} , i.e. $\|X_j\|_2 = \sqrt{n}$ for all $j = 1, \ldots, p$. For any $k \leq (n/4) \wedge (p/2)$, it holds that

$$R_F(X, \Theta_k, \sigma) \ge C_1 \sigma^2 \max\left(\frac{k \log(ep/k)}{n}, \frac{\exp\{C_2 k/n \log(p/k)\}}{n}\right),$$
 (8)

where $C_1, C_2 > 0$ are two universal constants.

The type of lower bound $C_1\sigma^2k/n\log(p/k)$ has been derived in several papers under different conditions: [24] assumes boundedness for both $k/n\log(p/k)$ and the maximum 2k-sparse eigenvalue²; [8] only requires $k \leq n$, and [1] has the minimal condition $k \leq p/2$. An interesting aspect of Theorem 2 is that in the ultra-high dimensional regime where $k/n\log(p/k) \gg \log n$, the term $C_1\sigma^2\frac{1}{n}\exp\{C_2\frac{k}{n}\log(\frac{p}{k})\}$ becomes dominating in the lower bound. This does not contradict with the upper bound $\tilde{C}_\theta\sigma^2k/n\log(p/k)$ from Theorem 1, because the SRE condition will not hold in such an ultra-high dimensional setting. In contrast, when $k/n\log(p/k)$ is bounded, the SRE condition holds with high probability if the rows of X are independent realizations of a large class of distributions [26, 21, 1].

In light of the aforementioned results, let us focus on the regime where $k/n \log(p/k) \leq C_3$ and $p \geq 2k$. Then, combining Theorems 1 and 2 we can reach the following conclusion about the minimax risk $R_F(X, \Theta_k, \sigma)$:

$$\frac{c\sigma^2 k \log(p/k)}{n} \le R_F(X, \Theta_k, \sigma) \le \frac{C\sigma^2 k \log(p/k)}{n},\tag{9}$$

as long as X satisfies the SRE(k,7) condition (which is true for a large set of matrices). Moreover, the Lasso estimator is minimax rate-optimal. This approximation approach offers very general non-asymptotic bounds such as the ones that have appeared in (9) to approximate the minimax risk, which is order-wise accurate. However, the constants that appear in the upper and lower bounds are not sharp. The looseness of constants is caused by the currently employed techniques (e.g. control of empirical process; information-theoretic inequalities).

By the nature of the constants (e.g. C depends on $\theta(k,7)$), the difference between c and C can be very large. For instance, as we describe later, the constant in the minimax risk under orthogonal designs should be close to 2 when p/k is large. However, the constant that appears in the upper bound of (7) is loose by a factor of $2401 \times (8 + 2\sqrt{2})^2/(2 \times 64) \approx 2199$ (even if we ignore the term $\frac{1}{\theta^4(k,7)}$). The constants obtained in lower bounds are often expected to be loose by a large factor as well. Therefore, the ratios between the upper and lower bounds can reach the magnitude of thousands, if not millions. Given that a constant-factor difference is already significant in most applications, the practical significance of minimax rate-optimal methods is sometimes questionable. This has led researchers to study the asymptotic minimax analysis that will be discussed in Section 2.3.

2.2 Rate-optimal results under the random design

In numerous scenarios, a more appropriate assumption regarding the data is that X is generated via a random mechanism. Consequently, assessing the risk solely over the specific dataset observed thus far is not desirable. Instead, we want our estimation procedure to generalize well to future

²The lower bound of [24] is in probability, but it implies the lower bound in expectation via Markov's inequality.

samples it encounters. In such cases, the random-design minimax risk defined below is a better criterion:

$$R_R(\Theta_k, \sigma) := \inf_{\hat{\beta}} \sup_{\beta \in \Theta_k} \mathbb{E}_{\beta} ||\hat{\beta} - \beta||_2^2.$$
 (10)

Compared to $R_F(X, \Theta_k, \sigma)$ discussed in the last section, the expectation in (10) is now with respect to both the noise vector z and the design matrix X, assuming independence between z and X. This seemingly straightforward change may further complicate the task of approximating the minimax risk. To quickly understand the issue, suppose $\max_j ||X_j||_2 \leq \sqrt{n}$. Then the expectation of the upper bound in (7) gives an upper bound for $R_R(\Theta_k, \sigma)$. This requires calculating the expectation of $\frac{1}{\theta^4(k,7)}$. However, calculating this expectation and proving that it is finite, is not straightforward. [32] managed to obtain a tight upper bound for $R_R(\Theta_k, \sigma)$ by analyzing the risk of a model selection procedure, and a matching lower bound was also derived. We summarize the results in the following theorem:

Theorem 3 (Proposition 6.4 of [32]). Suppose $\{x_i\}_{i=1}^n \stackrel{i.i.d}{\sim} \mathcal{N}(0,\Sigma)$ and Σ has ones on the diagonal. In the regime where $k/n\log(p/k) \leq C_1$ and $p \geq 2k$, it holds that

$$C_2 \frac{\sigma^2 k \log(p/k)}{n} \le R_R(\Theta_k, \sigma) \le C_3 \frac{\sigma^2 k \log(p/k)}{n\bar{\theta}_{2k}},$$

where $C_1, C_2, C_3 > 0$ are universal constants, and $\bar{\theta}_{2k} = \min_{\delta \in \Theta_{2k} \setminus \{0\}} \frac{\delta' \Sigma \delta}{\delta' \delta}$ is the minimal 2k-sparse eigenvalue of Σ .

As long as the sparse eigenvalue $\bar{\theta}_{2k}$ is bounded away from zero (which holds for rather general Σ 's [25]), the minimax risk $R_R(\Theta_k, \sigma)$ satisfies the same type of upper and lower bounds as in (9). Similar to the case of the fixed design, there can be a substantial gap between the constants that appear in the upper and lower bounds, which raises questions about the practical relevance of the rate-optimal minimax results. This major issue has led researchers to explore another approach that we will describe next.

2.3 Asymptotic approximation of minimax risk

The issue raised earlier regarding the loose constants in the upper and lower bounds of the minimax risk has been recognized and deliberated upon by many researchers. One proposed solution is to acquire a reliable approximation of the constants through asymptotic arguments. This approach was initially advocated by Donoho and Johnstone in the orthogonal design setting where $\frac{1}{n}X^TX = I_p$. In this simpler case of sparse linear regression, Donoho and Johnstone [12, 11, 20] demonstrated that

$$R_F(X, \Theta_k, \sigma) = \frac{(2 + o(1))\sigma^2 k \log(p/k)}{n},$$

as $p \to \infty$ and $k/p \to 0$.

Unfortunately, there has been limited exploration of this approach for the broader context of linear regression. A precise asymptotic approximation for $R_F(X,\Theta_k,\sigma)$ (under more general design) or $R_R(\Theta_k,\sigma)$ is still lacking. To our knowledge, the only existing result that has obtained sharp constants in a closely related context is the work of [27], which studies the Sorted L-One Penalized Estimator (SLOPE) introduced in [4]. For $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_p \geq 0$, the SLOPE estimator is defined as a solution of the minimization problem

$$\hat{\beta}_{\text{SLOPE}} \in \underset{b \in \mathbb{R}^p}{\text{arg min}} \frac{1}{2n} \|y - Xb\|_2^2 + \sum_{j=1}^p \lambda_j |b|_{(j)},$$

where $|b|_{(1)} \ge |b|_{(2)} \ge \cdots \ge |b|_{(p)}$ are the order statistics of $|b_1|, |b_2|, \ldots, |b_p|$. To show the optimality of SLOPE, [27] has proved the following results.

Theorem 4 (Theorem 1.2 & 1.3 in [27]). Assume model (1) with random Gaussian design $\{x_i\}_{i=1}^n \stackrel{i.i.d}{\sim} \mathcal{N}(0, I_p)$ and parameter space (2). Suppose $k/p \to 0$ and $(k \log p)/n \to 0$.

(i) For any $\epsilon > 0$, $\inf_{\hat{\beta}} \sup_{\beta \in \Theta_{L}} \mathbb{P} \bigg(\|\hat{\beta} - \beta\|_{2}^{2} > \frac{(2 - 2\epsilon)\sigma^{2}k \log(p/k)}{n} \bigg) \to 1.$

(ii) Fix 0 < q < 1 and set $\lambda_i = \sigma(1+\epsilon)n^{-1/2}\Phi^{-1}(1-iq/(2p))$, where Φ^{-1} is the quantile function of a standard normal and $\epsilon \in (0,1)$. Then, the SLOPE estimator satisfies

$$\sup_{\beta \in \Theta_k} \mathbb{P}\left(\|\hat{\beta}_{\text{SLOPE}} - \beta\|_2^2 > \frac{(2 + 6\epsilon)\sigma^2 k \log(p/k)}{n}\right) \to 0.$$

As is clear, the above theorem is under the random design setting. Intuitively speaking, Theorem 4 suggests the following for the minimax risk $R_R(\Theta_k, \sigma)$: as $k/p \to 0$ and $(k \log p)/n \to 0$, the minimax risk is approximately $2\sigma^2k/n\log(p/k)$, and that SLOPE asymptotically achieves the minimax risk. However, Theorem 4 does not exactly characterize the minimax risk. This is because the minimax risk $R_R(\Theta_k, \sigma)$ defined in (4) is based on the expected squared loss $\mathbb{E}\|\hat{\beta} - \beta\|_2^2$, while Theorem 4 characterizes the high-probability events for the squared loss $\|\hat{\beta} - \beta\|_2^2$. It is sometimes challenging to convert high-probability bounds on the squared loss to sharp (specially if we want sharp constants) bounds for the risk. To be more precise, Markov's inequality gives

$$\inf_{\hat{\beta}} \sup_{\beta \in \Theta_k} \mathbb{P} \bigg(\|\hat{\beta} - \beta\|_2^2 > \frac{(2 - 2\epsilon)\sigma^2 k \log(p/k)}{n} \bigg) \le \frac{\inf_{\hat{\beta}} \sup_{\beta \in \Theta_k} \mathbb{E}_{\beta} \|\hat{\beta} - \beta\|_2^2}{(2 - 2\epsilon)\sigma^2 k / n \log(p/k)}.$$

Based on Part (i) of Theorem 4, letting $n \to \infty$ and then $\epsilon \to 0+$ yields

$$\liminf_{n \to \infty} \frac{R_R(\Theta_k, \sigma)}{\sigma^2 k / n \log(p/k)} \ge 2.$$
(11)

We thus have converted the high-probability lower bound from Theorem 4 to a sharp lower bound for the minimax risk. However, the upper bound in Part (ii) of Theorem 4 is not directly transferable to a sharp upper bound for the minimax risk.

In summary, for both the fixed design (non-orthogonal) and random design cases, the asymptotic analysis of minimax risk that achieves sharp constants has remained an open problem. The main contribution of this paper is to show that for the isotropic Gaussian design,

$$R_R(\Theta_k, \sigma) = \frac{(2 + o(1))\sigma^2 k \log(p/k)}{n}.$$

As will be clarified later in the paper, in order to achieve this goal we have to address several technical challenges. We believe that our solutions can help create a viable path for evaluating the asymptotic approximations of $R_R(\Theta_k, \sigma)$ and $R_F(X, \Theta_k, \sigma)$ in more general settings, and even for problems beyond the sparse linear regression.

3 Our main contribution

As we described in the previous section, despite an extensive body of work on the sparse linear regression, the asymptotically exact characterization of the minimax risk has remained largely open. The main contribution of this paper is the following constant-sharp calculation for the minimax risk under Gaussian random designs.

Theorem 5. Assume model (1) with random Gaussian design $\{x_i\}_{i=1}^n \stackrel{i.i.d}{\sim} \mathcal{N}(0, I_p)$ and parameter space (2). As $k/p \to 0$ and $(k \log p)/n \to 0$, the minimax risk defined in (4) satisfies

$$R_R(\Theta_k, \sigma) = \frac{(2 + o(1))\sigma^2 k \log(p/k)}{n}.$$

In order to establish Theorem 5, we have used some of the techniques developed in [1] and [27]. However, proving Theorem 5 is not a straightforward application of the results presented in these two papers for the following reasons:

- For certain matrices, e.g. when a few columns of matrix X are linearly dependent, the minimax risk will be infinite. Let's call such matrices "infinite-risk matrices". Given our probabilistic assumption on matrix X, the probability of infinite-risk matrices is zero. However, the main concern in the minimax risk calculation is the matrices that are in the "vicinity" of infinite-risk matrices. For such matrices, the minimax risk is very large but still finite, and the closer they are to the infinite-risk matrices the risk is expected to be larger. On the other hand, the likelihood of being closer to infinite-risk matrices tends to be lower. Hence, in order to establish the sharp minimax risk characterization, one has to obtain sharp bounds on the probability of the vicinity of infinite-risk matrices and on the minimax risk for such matrices. As will be clarified in Section 4, this requires a delicate analysis.
- A key part of our minimax risk calculations is to obtain both a constant-sharp bound in probability and a rate-optimal bound in expectation for the Lasso estimator under a carefully chosen tuning parameter value. The former has been obtained in [27] for the SLOPE estimator instead of Lasso, and the latter has been derived for Lasso in [1] but with a larger value of tuning parameter (which does not yield sharp constants). Achieving both results requires a refined analysis of many arguments in [27] and [1]. We provide more detailed discussions in Section 4.

4 Roadmap of the proof of Theorem 5

We first introduce some notations used in this section. We use $\mathbbm{1}_{\mathcal{A}}$ to represent the indicator function of the set \mathcal{A} . For a given vector $v=(v_1,\ldots,v_p)\in\mathbb{R}^p, \|v\|_q=\left(\sum_{i=1}^p|v_i|^q\right)^{1/q}$ for $q\in(0,\infty)$, and $\mathrm{supp}(v)=\{1\leq i\leq p:v_i\neq 0\}$ denotes its support. We use $\{e_j\}_{j=1}^p$ to denote the natural basis in \mathbb{R}^p . For a matrix $X\in\mathbb{R}^{n\times p}$, X_j represents its jth column and $X_S\in\mathbb{R}^{n\times |S|}$ is the submatrix consisting of columns indexed by $S\subseteq\{1,2,\ldots,p\}$. For two non-zero real sequences $\{a_n\}_{n=1}^\infty$ and $\{b_n\}_{n=1}^\infty$, we use $a_n=o(b_n)$ to represent $|a_n/b_n|\to 0$ as $n\to\infty$, and use $a_n=O(b_n)$ for $\mathrm{sup}_n\,|a_n/b_n|<\infty$.

Given the lower bound result we have obtained in (11), to show $R_R(\Theta_k, \sigma) = (2+o(1))\sigma^2 k/n \log(p/k)$, it remains to prove the upper bound

$$\limsup_{n \to \infty} \frac{R_R(\Theta_k, \sigma)}{\sigma^2 k / n \log(p/k)} \le 2. \tag{12}$$

To prove the upper bound of the minimax risk, the main idea is to construct a good estimator and obtain a sharp upper bound for its maximum risk. Towards this goal, consider the following two estimators:

• Lasso:

$$\hat{\beta}^{L} \in \underset{b \in \mathbb{R}^{p}}{\min} \frac{1}{2n} \|y - Xb\|_{2}^{2} + \lambda \|b\|_{1}. \tag{13}$$

• Maximum likelihood estimator (MLE):

$$\hat{\beta}^M \in \underset{b \in \Theta_b}{\operatorname{arg\,min}} \|y - Xb\|_2^2. \tag{14}$$

Both estimators are known to achieve the minimax optimal rate (in probability or in expectation). To obtain constant-sharp upper bounds, we construct an aggregated estimator that combines Lasso and MLE, taking the form

$$\hat{\hat{\beta}} := \hat{\beta}^L \mathbb{1}_{\mathcal{A}} + \hat{\beta}^M \mathbb{1}_{\mathcal{A}^c}. \tag{15}$$

Here, the event $\mathcal{A} := \mathcal{A}(\delta_0, c_0, k)$ is defined as

$$\mathcal{A}(\delta_0, c_0, k) := \left\{ X \in \mathbb{R}^{n \times p} : \max_{j=1,\dots,p} \|X_j\|_2 \le (1 + \delta_0) \sqrt{n}, \ \theta(k, c_0) \ge 1 - \delta_0 \right\}, \tag{16}$$

where $\theta(k, c_0)$ was introduced in (6) and $\delta_0, c_0 > 0$ are constants that will be specified shortly. When the design matrix X is "well conditioned" (in the sense of \mathcal{A} , holding with high probability), our estimator $\hat{\beta}$ uses Lasso which will be shown to attain the sharp constant. Otherwise, our estimator resorts to the MLE that only induces a negligible error on rare events. The Lasso estimator depends on the tuning parameter $\lambda > 0$, and our choice of λ in the proof will be

$$\lambda_{\varepsilon} := (1 + \varepsilon)\sigma\sqrt{\frac{2\log(p/k)}{n}},\tag{17}$$

where $\varepsilon > 0$ is an arbitrarily small constant. We should emphasize two points here:

- 1. We do not expect Lasso to achieve constant-sharp results if we use larger values of λ that have been adopted in most existing work (e.g. [3, 1]).
- 2. As will be shown throughout the proof, it is much more challenging to obtain the optimal constant with the choice λ_{ε} , than obtaining the optimal rate using larger values of λ .

For the given tuning parameter λ_{ε} in (17), we set the corresponding constants δ_0 , c_0 of (16):

$$\delta_0 = \left(1 + \frac{\varepsilon}{2}\right)^{\frac{1}{3}} - 1, \quad c_0 = 8\sqrt{2}\varepsilon^{-1}\left(1 + \frac{\varepsilon}{2}\right)^{\frac{2}{3}} + 2\varepsilon^{-1} + 2. \tag{18}$$

Our goal is to prove that for any fixed $\varepsilon \in (0,1)$, our estimator $\hat{\beta}$ constructed in (15) satisfies

$$\sup_{\beta \in \Theta_k} \mathbb{E} \|\hat{\hat{\beta}} - \beta\|_2^2 \le \frac{(2 + o(1))(1 + \varepsilon)^2 \sigma^2 k \log(p/k)}{n}.$$
 (19)

The above result implies the upper bound for the minimax risk,

$$\limsup_{n \to \infty} \frac{R_R(\Theta_k, \sigma)}{\sigma^2 k / n \log(p/k)} \le 2(1 + \varepsilon)^2, \quad \forall \varepsilon \in (0, 1).$$

Further letting $\varepsilon \to 0+$ yields (12), and this will finish the proof.

To prove (19), from the construction of $\hat{\beta}$ we have

$$\mathbb{E}\|\hat{\hat{\beta}} - \beta\|_2^2 = \mathbb{E}\left(\|\hat{\beta}^L - \beta\|_2^2\mathbb{1}_{\mathcal{A}}\right) + \mathbb{E}\left(\|\hat{\beta}^M - \beta\|_2^2\mathbb{1}_{\mathcal{A}^c}\right),$$

and we aim to show

$$\sup_{\beta \in \Theta_k} \mathbb{E}\left(\|\hat{\beta}^M - \beta\|_2^2 \mathbb{1}_{\mathcal{A}^c}\right) = o\left(\sigma^2 k / n \log(p/k)\right),\tag{20}$$

$$\sup_{\beta \in \Theta_k} \mathbb{E}\left(\|\hat{\beta}^L - \beta\|_2^2 \mathbb{1}_{\mathcal{A}}\right) \le (2 + o(1))(1 + \varepsilon)^2 \sigma^2 k / n \log(p/k). \tag{21}$$

4.1 Proof of (20)

First note that from Hölder's inequality we have

$$\mathbb{E}\left(\|\hat{\beta}^M - \beta\|_2^2 \mathbb{1}_{\mathcal{A}^c}\right) \le \left(\mathbb{E}\|\hat{\beta}^M - \beta\|_2^m\right)^{\frac{2}{m}} \cdot \left(\mathbb{P}(\mathcal{A}^c)\right)^{\frac{m-2}{m}}, \quad m \in (2, \infty).$$
(22)

Hence, we will prove that $\sup_{\beta \in \Theta_k} \left(\mathbb{E} \| \hat{\beta}^M - \beta \|_2^m \right)^{\frac{2}{m}} = O(\sigma^2 k / n \log(p/k))$ and that $\mathbb{P}(\mathcal{A}^c) = o(1)$. Let us start with the simpler one, i.e. $\mathbb{P}(\mathcal{A}^c) = o(1)$. The following lemma proves this claim.

Lemma 1. Assume the design matrix $X \in \mathbb{R}^{n \times p}$ has i.i.d. $\mathcal{N}(0,1)$ entries and $p \geq 2$. For any constants $c_0 > 0, \delta_0 \in (0,1)$ and $k \in \{1,\ldots,p\}$, there exist absolute constants C, C' > 0 such that if

$$n \ge C\delta_0^{-2}(2+c_0)^2 k \log p,\tag{23}$$

then with probability at least $1 - 2\exp(-C'k\log p)$ we have

$$\max_{j=1,\dots,n} ||X_j||_2 \le (1+\delta_0)\sqrt{n}, \quad \theta(k,c_0) \ge 1-\delta_0.$$
(24)

Proof. Similar results exist in the literature of high-dimensional statistics. For completeness, we provide a proof based on uniform matrix deviation inequality from Lemma 7. Define the set

$$T := \Big\{ \delta \in \mathbb{R}^p : \|\delta\|_1 \le (1 + c_0)\sqrt{k}, \|\delta\|_2 = 1 \Big\},\,$$

and apply Lemma 7 to obtain: $\forall u > 0$, with probability at least $1 - 2e^{-u^2}$ it holds that

$$\sup_{\delta \in T} \left| \|X\delta\|_2 - \sqrt{n} \right| \le C_1 \left((1 + c_0) \sqrt{k} \cdot \mathbb{E} \|h\|_{\infty} + u \right), \quad h \sim \mathcal{N}(0, I_p),$$

$$\le C_1 \left((1 + c_0) C_2 \sqrt{k \log p} + u \right),$$

where the second inequality is due to Lemma 6, and $C_1, C_2 > 0$ are absolute constants. Choosing $u = C_2 \sqrt{k \log p}$ yields

$$\mathbb{P}\left(1 - \delta_0 \le \|X\delta\|_2 / \sqrt{n} \le 1 + \delta_0, \forall \delta \in T\right) \ge 1 - 2e^{-C_2^2 k \log p},$$

as long as $n \ge \delta_0^{-2} C_1^2 C_2^2 (2 + c_0)^2 k \log p$. The proof is thus completed because

$$\theta(k, c_0) = \inf_{\delta \in T} \|X\delta\|_2 / \sqrt{n}, \quad \sup_{1 \le i \le n} \|Xe_i\|_2 / \sqrt{n} \le \sup_{\delta \in T} \|X\delta\|_2 / \sqrt{n}.$$

As is clear from Lemma 1, under the conditions $k/p \to \infty$, $(k \log p)/n \to 0$, the event $\mathcal{A} = \mathcal{A}(\delta_0, c_0, k)$ with constants δ_0, c_0 in (18) satisfies $\mathbb{P}(\mathcal{A}^c) \leq 2e^{-C'k \log p} = o(1)$.

The next step is to obtain an upper bound for $\left(\mathbb{E}\|\hat{\beta}^M - \beta\|_2^m\right)^{\frac{2}{m}}$. This is done in the next proposition.

Proposition 1. Assume model (1) with isotropic Gaussian design. Suppose $k/p \to 0$ and $\frac{k \log(p/k)}{n} \to 0$. Then,

$$\sup_{\beta \in \Theta_h} \left(\mathbb{E} \| \hat{\beta}^M - \beta \|_2^m \right)^{\frac{2}{m}} = O(\sigma^2 k / n \log(p/k)), \quad \forall m \in [1, \infty).$$

The proof of this proposition is long, hence is deferred to Section B.1.

4.2 Proof of (21)

In establishing this bound, the main challenge is that we would like to obtain an upper bound with the sharp constant 2. Inspired by [27], we resort to the following oracle estimator:

$$\tilde{\beta}^O := \eta_\lambda (\beta + X^T z/n), \tag{25}$$

where $\eta_{\lambda}(u) := \arg\min_{b \in \mathbb{R}^p} \frac{1}{2} ||u - b||_2^2 + \lambda ||b||_1$ is known as the soft thresholding function. For each coordinate, the soft thresholding function can be characterized in closed-form as follows.

$$[\eta_{\lambda}(u)]_i = \arg\min_{b \in \mathbb{R}} \frac{1}{2} (u_i - b)^2 + \lambda |b| = \operatorname{sign}(u_i)(u_i - \lambda)_+$$

Note that this oracle estimator is using the same value of λ as the one for the Lasso estimator $\hat{\beta}^L$. The main idea is to show that the squared distance between $\hat{\beta}^L$ and $\tilde{\beta}^O$ is negligible, and hence the risk of Lasso $\hat{\beta}^L$ is asymptotically the same as the risk of this oracle estimator. Then the problem can be reduced to obtaining the upper bound for the risk of $\tilde{\beta}^O$. This is a simpler problem, because $\tilde{\beta}^O$ has a closed-form expression and thus its risk $\mathbb{E}\|\tilde{\beta}^O - \beta\|_2^2$ can be tightly upper bounded, as shown in the following lemma.

Lemma 2. Assume $X \in \mathbb{R}^{n \times p}$ has i.i.d. $\mathcal{N}(0,1)$ entries, $z \in \mathbb{R}^n$ has i.i.d. $\mathcal{N}(0,\sigma^2)$ entries, and X is independent of z. As $\frac{k}{p} \to 0$ and $\frac{\log(p/k)}{n} \to 0$, the oracle estimator $\tilde{\beta}^O$ with tuning parameter $\lambda = \lambda_{\varepsilon}$ in (17) satisfies

$$\sup_{\beta \in \Theta_k} \mathbb{E} \|\tilde{\beta}^O - \beta\|_2^2 \le (2 + o(1))(1 + \varepsilon)^2 \sigma^2 k / n \log(p/k), \quad \text{for any fixed } \epsilon > 0.$$

When $\varepsilon = 0$, $||z||_2^2 = n\sigma^2$, the sharp asymptotic risk of the oracle estimator (i.e. soft thresholding estimator) has been obtained in the study of sparse Gaussian sequence model [11, 12, 20, 15]. Lemma 2 can be viewed as a generalization to the scenario where $\varepsilon > 0$ and $||z||_2^2$ fluctuates around its mean $n\sigma^2$.

Proof. For a given $\tau \geq 0$, define the one-dimensional risk function:

$$r(\mu; \tau) = \mathbb{E}(\eta_{\tau}(\mu + w) - \mu)^2, \quad \mu \in \mathbb{R}, \ w \sim \mathcal{N}(0, 1).$$

By first conditioning on z and using the scale invariance property of the soft thresholding function, we can obtain that $\forall \beta \in \Theta_k$,

$$\mathbb{E}\|\tilde{\beta}^O - \beta\|_2^2 \le p \cdot \mathbb{E}\left(\frac{\|z\|_2^2}{n^2} \cdot r(0; \lambda_{\varepsilon} n/\|z\|_2)\right) + k \cdot \mathbb{E}\left(\frac{\|z\|_2^2}{n^2} \cdot \sup_{\mu \in \mathbb{R}} r(\mu; \lambda_{\varepsilon} n/\|z\|_2)\right)$$

$$\leq p \cdot \mathbb{E}\left(\frac{\|z\|_{2}^{2}}{n^{2}} \cdot \exp\left\{-\frac{\lambda_{\varepsilon}^{2} n^{2}}{2\|z\|_{2}^{2}}\right\}\right) + k \cdot \mathbb{E}\left(\frac{\|z\|_{2}^{2}}{n^{2}} \cdot \left(1 + \frac{\lambda_{\varepsilon}^{2} n^{2}}{\|z\|_{2}^{2}}\right)\right),$$

$$\leq p \cdot \mathbb{E}\left(\frac{\|z\|_{2}^{2}}{n^{2}} \cdot \exp\left\{-\frac{\lambda_{\varepsilon}^{2} n^{2}}{2\|z\|_{2}^{2}}\right\}\right) + \frac{k\sigma^{2}}{n} + k\lambda_{\varepsilon}^{2},$$
(26)

where the second inequality is due to the fact that $r(0;\tau) \leq e^{-\tau^2/2}$ and $r(\mu;\tau) \leq 1+\tau^2$ (see Chapter 8.2 in [20]). To bound the remaining expectation above, define the event

$$\mathcal{B} = \left\{ \left| \frac{\|z\|_2}{\sqrt{n}} - \sigma \right| \le \varepsilon \sigma \right\}.$$

According to the concentration of the norm of isotropic Gaussian (Theorem 3.1.1 in [30]), we obtain that $\mathbb{P}(\mathcal{B}^c) \leq 2e^{-Cn\varepsilon^2}$ for some absolute constant C > 0. We can then proceed to have

$$\mathbb{E}\left(\frac{\|z\|_{2}^{2}}{n^{2}}\cdot\exp\left\{-\frac{\lambda_{\varepsilon}^{2}n^{2}}{2\|z\|_{2}^{2}}\right\}\right)$$

$$=\mathbb{E}\left(\frac{\|z\|_{2}^{2}}{n^{2}}\cdot\exp\left\{-\frac{\lambda_{\varepsilon}^{2}n^{2}}{2\|z\|_{2}^{2}}\right\}\cdot\mathbb{1}_{\mathcal{B}}\right)+\mathbb{E}\left(\frac{\|z\|_{2}^{2}}{n^{2}}\cdot\exp\left\{-\frac{\lambda_{\varepsilon}^{2}n^{2}}{2\|z\|_{2}^{2}}\right\}\cdot\mathbb{1}_{\mathcal{B}^{c}}\right)$$

$$\leq\mathbb{E}\left(\frac{\|z\|_{2}^{2}}{n^{2}}\cdot\frac{k}{p}\cdot\mathbb{1}_{\mathcal{B}}\right)+\mathbb{E}\left(\frac{\|z\|_{2}^{2}}{n^{2}}\cdot\mathbb{1}_{\mathcal{B}^{c}}\right)\leq\frac{k\sigma^{2}}{np}+\sqrt{\mathbb{E}\left(\frac{\|z\|_{2}^{4}}{n^{4}}\right)}\cdot\sqrt{\mathbb{P}(\mathcal{B}^{c})}$$

$$\leq\frac{k\sigma^{2}}{np}+\sigma^{2}\sqrt{\frac{(2n+4)e^{-Cn\varepsilon^{2}}}{n^{3}}},$$
(27)

where in the last two inequalities we have used Cauchy–Schwarz inequality and the second moment of chi-squared distribution. Combining (26) and (27) gives us

$$\sup_{\beta \in \Theta_k} \mathbb{E} \|\tilde{\beta}^O - \beta\|_2^2 \le \frac{2(1+\epsilon)^2 \sigma^2 k \log(p/k)}{n} + \frac{2k\sigma^2}{n} + \sigma^2 p \sqrt{\frac{(2n+4)e^{-Cn\epsilon^2}}{n^3}},
= \frac{(2+o(1))(1+\epsilon)^2 \sigma^2 k \log(p/k)}{n},$$

where the last equality holds under the scaling condition $\frac{k}{p} \to 0, \frac{\log(p/k)}{n} \to 0.$

As we pointed out at the beginning of the section, to transfer the risk bound of $\tilde{\beta}^O$ to $\hat{\beta}^L$, we need to show that $\|\hat{\beta}^L - \tilde{\beta}^O\|_2^2$ is asymptotically negligible compared to $\|\tilde{\beta}^O - \beta\|_2^2$. The next lemma paves the way.

Lemma 3. For a given $\Delta \in (0,1)$, define the following event,

$$\mathcal{B}_{\Delta} = \left\{ \exists S_* \subseteq \{1, \dots, p\} \text{ such that } \operatorname{supp}(\hat{\beta}^L) \cup \operatorname{supp}(\tilde{\beta}^O) \cup \operatorname{supp}(\beta) \subseteq S_* \right.$$
and all the eigenvalues of $X_{S_*}^T X_{S_*} / n$ lie within $[1 - \Delta, 1 + \Delta] \right\}.$ (28)

Assume model (1). Then on the event \mathcal{B}_{Δ} ,

$$\|\tilde{\beta}^O - \hat{\beta}^L\|_2 \le \frac{\Delta}{1 - \Delta} \|\tilde{\beta}^O - \beta\|_2.$$

Proof. Recall the definition of $\hat{\beta}^L$ in (13) and $\tilde{\beta}^O$ in (25). Based on the KKT conditions for the Lasso problem, it is straightforward to verify that $\hat{\beta}^L$ satisfies the fixed point equation,

$$\hat{\beta}^L = \eta_{\lambda} \Big(\hat{\beta}^L + X^T (y - X \hat{\beta}^L) / n \Big) = \eta_{\lambda} \Big(\beta + (I - X^T X / n) (\hat{\beta}^L - \beta) + X^T z / n \Big),$$

where in the second equality we have used $y = X\beta + z$. Now utilizing the definition of S_* , we can proceed to obtain

$$\|\hat{\beta}^{L} - \tilde{\beta}^{O}\|_{2} = \|\eta_{\lambda} \Big(\beta_{S_{*}} + (I - X_{S_{*}}^{T} X_{S_{*}}/n)(\hat{\beta}_{S_{*}}^{L} - \beta_{S_{*}}) + X_{S_{*}}^{T} z/n\Big) - \eta_{\lambda} \Big(\beta_{S_{*}} + X_{S_{*}}^{T} z/n\Big)\|_{2}$$

$$\stackrel{(a)}{\leq} \|(I - X_{S_{*}}^{T} X_{S_{*}}/n)(\hat{\beta}_{S_{*}}^{L} - \beta_{S_{*}})\|_{2} \leq \|I - X_{S_{*}}^{T} X_{S_{*}}/n\|_{2} \cdot \|\hat{\beta}_{S_{*}}^{L} - \beta_{S_{*}}\|_{2}$$

$$\stackrel{(b)}{\leq} \Delta \|\hat{\beta}^{L} - \beta\|_{2} \leq \Delta (\|\hat{\beta}^{L} - \tilde{\beta}^{O}\|_{2} + \|\tilde{\beta}^{O} - \beta\|_{2}).$$

Here, (a) is due to the nonexpansiveness of the soft thresholding operator $\eta_{\lambda}(\cdot)$, and (b) holds because all the eigenvalues of $X_{S_*}^T X_{S_*}/n$ lie within $[1-\Delta, 1+\Delta]$. Solving the above inequality for $\|\tilde{\beta}^O - \hat{\beta}^L\|_2$ completes the proof.

Lemma 3 is similar to Lemma 4.2 in [27] that establishes a similar result for the SLOPE estimator. Here, we provided a more direct proof that yields a better dependence on Δ .

In light of Lemmas 2 and 3, to prove (21), we bound $\mathbb{E}(\|\hat{\beta}^L - \beta\|_2^2 \mathbb{1}_A)$ as follows:

$$\begin{split} \mathbb{E} \| \hat{\beta}^L - \beta \|_2^2 \mathbb{1}_{\mathcal{A}} &= \ \mathbb{E} \| \tilde{\beta}^O - \beta \|_2^2 \mathbb{1}_{\mathcal{A} \cap \mathcal{B}_{\Delta}} + 2 \mathbb{E} (\hat{\beta}^L - \tilde{\beta}^O)^T (\tilde{\beta}^O - \beta) \mathbb{1}_{\mathcal{A} \cap \mathcal{B}_{\Delta}} \\ &+ \mathbb{E} \| \hat{\beta}^L - \tilde{\beta}^O \|_2^2 \mathbb{1}_{\mathcal{A} \cap \mathcal{B}_{\Delta}} + \mathbb{E} \| \hat{\beta}^L - \beta \|_2^2 \mathbb{1}_{\mathcal{A} \cap \mathcal{B}_{\Delta}^c} \\ &\leq \ \mathbb{E} \| \tilde{\beta}^O - \beta \|_2^2 \mathbb{1}_{\mathcal{A} \cap \mathcal{B}_{\Delta}} + 2 \sqrt{\mathbb{E} \| \hat{\beta}^L - \tilde{\beta}^O \|_2^2 \mathbb{1}_{\mathcal{A} \cap \mathcal{B}_{\Delta}}} \cdot \sqrt{\mathbb{E} \| \tilde{\beta}^O - \beta \|_2^2 \mathbb{1}_{\mathcal{A} \cap \mathcal{B}_{\Delta}}} \\ &+ \mathbb{E} \| \hat{\beta}^L - \tilde{\beta}^O \|_2^2 \mathbb{1}_{\mathcal{A} \cap \mathcal{B}_{\Delta}} + \left(\mathbb{E} \| \hat{\beta}^L - \beta \|_2^q \mathbb{1}_{\mathcal{A}} \right)^{\frac{2}{q}} (\mathbb{P}(\mathcal{B}_{\Delta}^c))^{1 - \frac{2}{q}} \\ &\leq \ \mathbb{E} \| \tilde{\beta}^O - \beta \|_2^2 + 2 \sqrt{\mathbb{E} \| \hat{\beta}^L - \tilde{\beta}^O \|_2^2 \mathbb{1}_{\mathcal{B}_{\Delta}}} \cdot \sqrt{\mathbb{E} \| \tilde{\beta}^O - \beta \|_2^2} \\ &+ \mathbb{E} \| \hat{\beta}^L - \tilde{\beta}^O \|_2^2 \mathbb{1}_{\mathcal{B}_{\Delta}} + \left(\mathbb{E} \| \hat{\beta}^L - \beta \|_2^q \mathbb{1}_{\mathcal{A}} \right)^{\frac{2}{q}} (\mathbb{P}(\mathcal{B}_{\Delta}^c))^{1 - \frac{2}{q}} \end{split}$$

where in the first inequality we have used Hölder's inequality with a power factor q > 2. Plugging the bounds of Lemmas 2 and 3 into the above inequality gives us

$$\sup_{\beta \in \Theta_{k}} \mathbb{E}\left(\|\hat{\beta}^{L} - \beta\|_{2}^{2} \mathbb{1}_{\mathcal{A}}\right) \leq \frac{(2 + o(1))(1 + \epsilon)^{2} \sigma^{2} k / n \log(p/k)}{(1 - \Delta)^{2}} + \left(\sup_{\beta \in \Theta_{k}} \mathbb{E}\|\hat{\beta}^{L} - \beta\|_{2}^{q} \mathbb{1}_{\mathcal{A}}\right)^{\frac{2}{q}} \cdot \left(\sup_{\beta \in \Theta_{k}} \mathbb{P}(\mathcal{B}_{\Delta}^{c})\right)^{1 - \frac{2}{q}}.$$
(29)

Having (29), we will further show the following two results to eventually prove (21):

(i) We can pick $\Delta \to 0+$ such that $\sup_{\beta \in \Theta_k} \mathbb{P}(\mathcal{B}^c_{\Delta}) = o(1)$.

$$\text{(ii) } \Big(\sup_{\beta \in \Theta_k} \mathbb{E} \| \hat{\beta}^L - \beta \|_2^q \mathbb{1}_{\mathcal{A}} \Big)^{\frac{2}{q}} = O \Big(\sigma^2 k / n \log(p/k) \Big).$$

For the proof of result (i), we borrow the notion of resolvent set from [27].

Definition 2 (Resolvent set of Lasso). Fix $S = \text{supp}(\beta)$ of cardinality at most k, and an integer k^* obeying $k < k^* < p$. The set $S_* = S_*(S, k^*)$ is said to be a resolvent set if it is the union of S and the $k^* - k$ indices with the largest values of $|X_i^T z|$ among all $i \in \{1, ..., p\} \setminus S$.

It turns out that with a proper choice of k^* , the resolvent set satisfies both conditions required in \mathcal{B}_{Δ} . The next proposition aims to prove this point.

Proposition 2. Assume model (1) with isotropic Gaussian design. Let $S_* = S_*(S, k^*)$ be the resolvent set and $\Delta = 10\sqrt{\frac{k^*\log(p/k^*)}{n}}$. Suppose $k^* \geq 2k$, $k^*/p \to 0$ and $(k^*\log p)/n \to 0$. Then, S_* satisfies the conditions required in \mathcal{B}_{Δ} with probability converging to one. As a result,

$$\sup_{\beta \in \Theta_k} \mathbb{P}(\mathcal{B}^c_{\Delta}) = o(1).$$

The proof of this proposition is presented in Section B.2. Note that under the conditions $\frac{k}{p} \to 0$, $\frac{k \log p}{n} \to 0$ provided in Theorem 5, there are many choices of k^* that satisfy the required conditions of Proposition 2. For instance, we may simply pick $k^* = 2k$.

It remains to prove result (ii). The past work [1] has proved that the supremum risk of Lasso (on the event \mathcal{A}) can be bounded by $O(\sigma^2 k/n \log(p/k))$, which is what we aim to prove as well. However, in order to prove this upper bound, [1] uses $\lambda \geq C\sigma\sqrt{\frac{2\log(p/k)}{n}}$ for a constant $C>2\sqrt{2}+1$. As we discussed before, this larger choice of λ does not deliver the sharp constant for the (asymptotic) minimax risk. For our purpose, we refine the proof in [1] to obtain the same rate-optimal results under the smaller tuning parameter $\lambda_{\varepsilon}=(1+\varepsilon)\sigma\sqrt{\frac{2\log(p/k)}{n}}$ that is used for our estimator.

Proposition 3. Assume model (1) with isotropic Gaussian design. As $k/p \to 0$, the Lasso estimator $\hat{\beta}^L$ in (13) with regularization parameter λ_{ε} satisfies $\forall q \in [2, \infty)$,

$$\left(\sup_{\beta \in \Theta_k} \mathbb{E}\Big(\|\hat{\beta}^L - \beta\|_2^q \, \mathbb{1}_{\mathcal{A}}\Big)\right)^{\frac{2}{q}} = O\Big(\frac{\sigma^2 k \log(p/k)}{n}\Big).$$

Here, the event $\mathcal{A} = \mathcal{A}(\delta_0, c_0, k)$ is defined in (16) and the constants δ_0, c_0 are specified in (18). The proof of this proposition can be found in Section B.3.

5 Conclusion

In this paper, we study the minimax risk of the sparse linear regression problem. Despite the considerable volume of research dedicated to this area, as discussed in the paper, the constant-sharp analysis of minimax risk is still rather limited in the literature. To contribute along this line, we explored the asymptotic scenario where $(k \log p)/n \to 0$ and derived the sharp asymptotic minimax risk $2\sigma^2 k/n \log(p/k)$ under isotropic Gaussian design. Along the way, we provided a summary of existing literature results and highlighted some of the fundamental issues that have remained unresolved.

Proofs of technical results

We collect the notations used throughout the proof sections for convenience. For an integer n, let $[n] = \{1, 2, \ldots, n\}$. We use $\mathbbm{1}_{\mathcal{A}}$ to represent the indicator function of the set \mathcal{A} . For a given vector $v = (v_1, \ldots, v_p) \in \mathbb{R}^p$, $\|v\|_0 = \#\{i : v_i \neq 0\}$, $\|v\|_\infty = \max_i |v_i|$, $\|v\|_q = \left(\sum_{i=1}^p |v_i|^q\right)^{1/q}$ for $q \in (0, \infty)$, supp $(v) = \{1 \leq i \leq p : v_i \neq 0\}$ denotes its support, and $v_S \in \mathbb{R}^{|S|}$ denotes the subvector consisting of coordinates in $S \subseteq [p]$. The inner product of two vectors a, b is written as $\langle a, b \rangle$. We use $\{e_j\}_{j=1}^p$ to denote the natural basis in \mathbb{R}^p . For a matrix $X \in \mathbb{R}^{n \times p}$, $\sigma_{\min}(X)$ denotes its smallest singular value and $\sigma_{\max}(X)$ (or $\|X\|_2$) denotes its largest singular value; X_j represents its jth column and $X_S \in \mathbb{R}^{n \times |S|}$ is the submatrix consisting of columns indexed by $S \subseteq [p]$. The $p \times p$ identity matrix is denoted by I_p . For two real numbers a and b, $a \vee b$ and $a \wedge b$ represent $\max(a, b)$ and $\min(a, b)$, respectively. For two non-zero real sequences $\{a_n\}_{n=1}^\infty$ and $\{b_n\}_{n=1}^\infty$, we use $a_n = o(b_n)$ to represent $|a_n/b_n| \to 0$ as $n \to \infty$, and use $a_n = O(b_n)$ for $\sup_i |a_n/b_n| < \infty$. The notation $x \stackrel{d}{=} y$ means that the random variables x and y have the same distribution. For a random vector x, the notation $\|x\|_{\psi_2}$ denotes its sub-Gaussian norm. Finally, we reserve the notations $\Phi(y)$ and $\Phi^{-1}(y)$ for CDF of $\mathcal{N}(0,1)$ and its inverse function, respectively.

A Preliminaries

Lemma 4 (Binomial coefficient, Exercise 0.0.5 in [30]). For a given positive integer p,

$$\binom{p}{s} \le \left(\frac{ep}{s}\right)^s,$$

holds for all integers $s \in [1, p]$.

Lemma 5 (Covering number of unit sphere, Corollary 4.2.13 in [30]). The covering numbers $\mathcal{N}(S^{n-1}, \epsilon)$ of the unit Euclidean sphere $S^{n-1} := \{v \in \mathbb{R}^n : ||v||_2 = 1\}$ satisfy

$$\mathcal{N}(S^{n-1}, \epsilon) \le \left(\frac{3}{\epsilon}\right)^n, \quad \forall \epsilon \in (0, 1].$$

Lemma 6 (Maximum of sub-gaussians, Exercise 2.5.10 in [30]). Let w_1, w_2, \ldots, w_N be sub-gaussian random variables which are not necessarily independent. Let $K = \max_i ||w_i||_{\psi_2}$ be the maximum sub-gaussian norm. Then for every $N \geq 2$ we have

$$\mathbb{E}\max_{i\leq N}|w_i|\leq CK\sqrt{\log N},$$

where C > 0 is a universal constant.

Lemma 7 (Matrix deviation inequality, Exercise 9.1.8 in [30]). Let A be an $n \times p$ matrix whose rows $e_i^T A$ are independent, isotropic and sub-Gaussian random vectors in \mathbb{R}^p . Then for any given subset $T \subset \mathbb{R}^p$, the event

$$\sup_{x \in T} \left| ||Ax||_2 - \sqrt{n} ||x||_2 \right| \le CK^2 \left(\gamma(T) + u \cdot \operatorname{rad}(T) \right)$$

holds with probability at least $1 - 2e^{-u^2}$. Here, $\gamma(T) = \mathbb{E}\sup_{x \in T} |h^T x|, h \sim \mathcal{N}(0, I_p); \operatorname{rad}(T) = \sup_{x \in T} ||x||_2; K = \max_i ||e_i^T A||_{\psi_2}; C > 0$ is a universal constant and $u \geq 0$ is any constant.

Lemma 8 (χ^2 -concentration, Lemma 2 of [18]). Let $g_1, \ldots, g_d \stackrel{i.i.d.}{\sim} \mathcal{N}(0,1)$. Then,

$$\mathbb{P}\Big(\sum_{i=1}^{d} g_i^2 < d(1-\tau)\Big) \le e^{\frac{d}{2}\left(\tau + \log(1-\tau)\right)}, \quad \forall \tau \in (0,1).$$

Lemma 9 (Singular values of Gaussian matrices, Corollary 5.35 in [31]). Let A be an $N \times n$ $(N \ge n)$ matrix whose entries are independent standard normal random variables. Then,

$$\mathbb{P}\left(\sqrt{N} - \sqrt{n} - t \le \sigma_{\min}(A) \le \sigma_{\max}(A) \le \sqrt{N} + \sqrt{n} + t\right) \ge 1 - 2e^{-\frac{t^2}{2}}, \quad \forall t > 0.$$

Lemma 10 (Concentration of the maximum of Gaussians). Let $g_1, \ldots, g_p \stackrel{i.i.d}{\sim} \mathcal{N}(0,1)$ with $p \geq 2$. For all u > 0, we have

$$\mathbb{P}\left(\max_{1 \le i \le p} |g_i| \ge \sqrt{2\log p} + u\right) \le e^{-\frac{u^2}{2}}.$$

Proof. Using union bound and Gaussian tail bound, for $a \ge 1$,

$$\mathbb{P}\big(\max_{1 \le i \le p} |g_i| \ge a\big) \le \sum_{i=1}^p \mathbb{P}\big(|g_i| \ge a\big) \le p \cdot \frac{2}{a} \frac{1}{\sqrt{2\pi}} e^{-\frac{a^2}{2}} \le \frac{2p}{\sqrt{2\pi}} e^{-\frac{a^2}{2}} \le p e^{-\frac{a^2}{2}}.$$

For any u > 0, setting $a = \sqrt{2 \log p} + u \ge 1$ (since $p \ge 2$) in the above bound yields

$$\mathbb{P}\Big(\max_{1 \le i \le p} |g_i| \ge \sqrt{2\log p} + u\Big) \le e^{-\frac{u^2}{2} - u\sqrt{2\log p}} \le e^{-\frac{u^2}{2}}.$$

Lemma 11. Let $g_1, \ldots, g_p \stackrel{i.i.d}{\sim} \mathcal{N}(0,1)$, and $|g|_{(1)} \geq |g|_{(2)} \geq \cdots \geq |g|_{(p)}$ denote the order statistics of $(|g_1|, \ldots, |g_p|)$.

(i) For all $2 \le k \le p$, $\mathbb{E}|g|_{(k)} \le \sqrt{2\log(\frac{2p}{k-1})}$.

(ii) For all
$$1 \le k \le p$$
, $\mathbb{P}(|g|_{(k)} - \mathbb{E}|g|_{(k)} \ge u) \le e^{-\frac{u^2}{2}}, \ \forall u > 0$.

Proof. Prove (i). Let $U_1,\ldots,U_p\stackrel{i.i.d.}{\sim} \mathrm{Unif}(0,1)$ with order statistics $U_{(1)} \leq U_{(2)} \leq \cdots \leq U_{(p)}$, and $Y_1,\ldots,Y_p\stackrel{i.i.d.}{\sim} \mathrm{Exp}(1)$ with order statistics $Y_{(1)} \geq Y_{(2)} \geq \ldots \geq Y_{(p)}$. We will use the following well known distributional results: for $k=1,2,\ldots,p$,

$$|g|_{(k)} \stackrel{d}{=} \Phi^{-1}(1 - U_{(k)}/2), \quad Y_{(k)} \stackrel{d}{=} \log \frac{1}{U_{(k)}}, \quad Y_{(k)} - Y_{(k+1)} \stackrel{d}{=} \operatorname{Exp}(k),$$

where $\Phi^{-1}(\cdot)$ is the inverse function of CDF of standard normal and $Y_{(p+1)}:=0$. Then,

$$(\mathbb{E}|g|_{(k)})^{2} \leq \mathbb{E}|g|_{(k)}^{2} = \mathbb{E}\left(\Phi^{-1}(1 - U_{(k)}/2)\right)^{2} \stackrel{(a)}{\leq} \mathbb{E}\left[2\log\frac{2}{U_{(k)}}\right]$$

$$= 2\log 2 + 2\sum_{j=k}^{p} \mathbb{E}(Y_{(j)} - Y_{(j+1)}) = 2\log 2 + 2\sum_{j=k}^{p} \frac{1}{j} \stackrel{(b)}{\leq} 2\log\frac{2p}{k-1}.$$

Here, (a) is due to the Gaussian tail bound $1 - \Phi(t) \leq e^{-\frac{t^2}{2}}, \forall t > 0$; (b) holds because $\sum_{j=k}^p \frac{1}{j} \leq \sum_{j=k}^p \int_{j-1}^j \frac{1}{x} dx = \log p - \log(k-1)$ for $k \geq 2$.

Prove (ii). Given that each order statistic is a 1-Lipschitz function (e.g. Example 2.29 in [33]), we have

$$|g|_{(k)} - |\tilde{g}|_{(k)}| \le \sqrt{\sum_{i=1}^{p} (|g_i| - |\tilde{g}_i|)^2} \le \sqrt{\sum_{i=1}^{p} (g_i - \tilde{g}_i)^2},$$

where in the second inequality we have used $||a| - |b|| \le |a - b|, \forall a, b \in \mathbb{R}$. Hence, $|g|_{(k)}$, as a function of (g_1, \ldots, g_p) , is 1-Lipschitz as well. Applying standard Gaussian concentration inequality (Theorem 5.6 in [5]) completes the proof.

Lemma 12 (Proposition E.1 in [1]). Let $g_1, \ldots, g_p \stackrel{i.i.d}{\sim} \mathcal{N}(0,1)$, and $|g|_{(1)} \geq |g|_{(2)} \geq \cdots \geq |g|_{(p)}$ denote the order statistics of $(|g_1|, \ldots, |g_p|)$. Then for any $s \in \{1, \ldots, p\}$ and all t > 0, we have

$$\mathbb{P}\bigg(\frac{1}{s}\sum_{j=1}^{s}|g|_{(j)}^{2}>t\log(2p/s)\bigg)\leq (2p/s)^{1-\frac{3t}{8}}.$$

Lemma 13. Under the assumptions of Lemma 12, then for any fixed $\delta_1 > 0$,

$$\mathbb{P}\left(\left(\max_{1 \le j \le k} \frac{|g|_{(j)}}{4\sqrt{\log(2p/j)}}\right) \vee \frac{|g|_{(k+1)}}{(1+\delta_1)\sqrt{2\log(p/k)}} \le 1\right) \\
\ge 1 - \frac{k}{2p} - \exp\left\{-\frac{1}{2}\left((1+\delta_1)\sqrt{2\log(p/k)} - \sqrt{2\log(2p/k)}\right)^2\right\} > \frac{1}{2},$$

when p/k is large enough.

Proof. The proof follows that of Proposition E.2 in [1].

Lemma 12 with t = 16/3 and the inequality $|g|_{(j)}^2 \leq \frac{1}{j} \sum_{l=1}^{j} |g|_{(l)}^2$ imply

$$P(|g|_{(j)}^2 \le \frac{16}{3}\log(2p/j)) \ge 1 - \frac{j}{2p}, \quad j = 1, \dots, p.$$
 (30)

Let $q \ge 0$ be an integer such that $2^q \le k < 2^{q+1}$. Applying (30) to $j = 2^l$ for $l = 0, \dots, q-1$ and using the union bound, we obtain that the event

$$\Omega_0 := \left\{ \max_{l=0,\dots,q-1} \frac{|g|_{(2^l)} \sqrt{3}}{4\sqrt{\log(2p/2^l)}} \le 1 \right\}$$

satisfies $\mathbb{P}(\Omega_0) \ge 1 - \sum_{l=0}^{q-1} \frac{2^l}{2p} = 1 - \frac{2^q-1}{2p} \ge 1 - \frac{k}{2p}$. For any $j < 2^q$, there exists $l \in \{0, \dots, q-1\}$ such that $2^l \le j < 2^{l+1}$. On the event Ω_0 ,

$$|g|_{(j)} \le |g|_{(2^l)} \le \frac{4}{\sqrt{3}} \sqrt{\log \frac{2p}{2^l}} \le \frac{4}{\sqrt{3}} \sqrt{\log \frac{4p}{j}} \le 4 \sqrt{\log \frac{2p}{j}}, \quad \forall j < 2^q.$$

And for $2^q \leq j \leq k$,

$$|g|_{(j)} \le |g|_{(2^{q-1})} \le \frac{4}{\sqrt{3}} \sqrt{\log \frac{2p}{2^{q-1}}} < \frac{4}{\sqrt{3}} \sqrt{\log \frac{8p}{j}} \le 4\sqrt{\log \frac{2p}{j}}.$$

Thus, on the event Ω_0 we have $|g|_{(j)} \leq 4\sqrt{\log(2p/j)}$ for all $j=1,\ldots,k$. In addition, using Lemma 11, we have

$$\mathbb{P}\left(\frac{|g|_{(k+1)}}{(1+\delta_1)\sqrt{2\log\frac{p}{k}}} \ge 1\right) \le e^{-\frac{1}{2}\left((1+\delta_1)\sqrt{2\log(p/k)} - \sqrt{2\log(2p/k)}\right)^2}.$$

B Proof of Propositions 1-3

B.1 Proof of Proposition 1

Recalling the definition of $\hat{\beta}^M$ in (14), we start with the basic inequality

$$||y - X\hat{\beta}^M||_2^2 \le ||y - X\beta||_2^2.$$

With $y = X\beta + z$, this implies

$$\frac{1}{n} \|X(\hat{\beta}^M - \beta)\|_2^2 \le \frac{2}{n} z^T X(\hat{\beta}^M - \beta). \tag{31}$$

For a given $s \in \{1, \ldots, p\}$, define

$$V_s := \inf_{\Delta \in T_s} \frac{1}{n} \|X\Delta\|_2^2, \quad T_s := \left\{ \Delta \in \mathbb{R}^p : \|\Delta\|_2 = 1, \|\Delta\|_0 \le s \right\}. \tag{32}$$

Both $\hat{\beta}^M$ and β are in Θ_k , hence $\|\hat{\beta}^M - \beta\|_0 \le 2k$. We then continue from (31) to obtain

$$V_{2k} \cdot \|\hat{\beta}^{M} - \beta\|_{2}^{2} \leq \frac{1}{n} \|X(\hat{\beta}^{M} - \beta)\|_{2}^{2} \leq \frac{2}{n} z^{T} X(\hat{\beta}^{M} - \beta)$$
$$\leq \frac{2}{\sqrt{n}} \|\hat{\beta}^{M} - \beta\|_{2} \cdot \sup_{u \in T_{2k}} z^{T} Xu / \sqrt{n}.$$

Therefore,

$$\|\hat{\beta}^M - \beta\|_2 \le \frac{2}{\sqrt{n}V_{2k}} \cdot \sup_{u \in T_{2k}} z^T X u / \sqrt{n}.$$

Then, using Cauchy-Schwarz inequality, we have

$$\sup_{\beta \in \Theta_k} \mathbb{E} \|\hat{\beta}^M - \beta\|_2^m \le \frac{2^m}{n^{\frac{m}{2}}} \cdot \left(\mathbb{E} \frac{1}{V_{2k}^{2m}} \right)^{\frac{1}{2}} \cdot \left(\mathbb{E} \left(\sup_{u \in T_{2k}} z^T X u / \sqrt{n} \right)^{2m} \right)^{\frac{1}{2}}. \tag{33}$$

Hence, we need to bound the two terms on the right-hand side of (33). This is done in Lemmas 14 and 15. Combining (33) with Lemmas 14 and 15 completes the proof of Proposition 1.

Lemma 14. Suppose the matrix $X \in \mathbb{R}^{n \times p}$ has i.i.d $\mathcal{N}(0,1)$ entries. For $s \in \{1,\ldots,p\}$, let V_s be defined as in (32). If $(s \log(ep/s))/n \to 0$, then, for every fixed r > 0, we have

$$\mathbb{E}\frac{1}{V_s^r} = O(1).$$

Lemma 15. Suppose the matrix $X \in \mathbb{R}^{n \times p}$ has i.i.d $\mathcal{N}(0,1)$ entries, and is independent of $z \sim \mathcal{N}(0,\sigma^2 I_n)$. For $s \in \{1,\ldots,p\}$, let T_s be defined as in (32). Then,

$$\mathbb{E}\Big(\sup_{u \in T_*} z^T X u / \sqrt{n}\Big)^q \le c_q \Big(\sigma \sqrt{s \log(ep/s)}\Big)^q, \quad \forall q \in [1, \infty),$$

for some constant $c_q > 0$ that only depends on q.

Proof of Lemma 14. Throughout the proof, we fix $s \in \{1, ..., p\}$ and let $V := V_s$ for notational simplicity. We have

$$\mathbb{E}\frac{1}{V^r} = \mathbb{E}\left(\frac{1}{V^r}\mathbb{1}_{(V \le x)}\right) + \mathbb{E}\left(\frac{1}{V^r}\mathbb{1}_{(V > x)}\right),\tag{34}$$

where we set³ $x = e^{-8}$. It is clear that

$$\mathbb{E}\left(\frac{1}{V^r}\mathbb{1}_{(V>x)}\right) < \frac{1}{x^r}.\tag{35}$$

Hence, in the rest of the proof, we aim to obtain an upper bound for $\mathbb{E}\left(\frac{1}{V^r}\mathbb{1}_{(V\leq x)}\right)$. Towards this goal, we first bound $\mathbb{P}(V\leq 1-t)$. For $\forall t\in(0,1)$, using the union bound, we have

$$\mathbb{P}(V \le 1 - t) = \mathbb{P}\left(\min_{S \subseteq [p]: |S| = s} \inf_{\|\Delta\|_2 = 1} \frac{1}{n} \|X_S \Delta\|_2^2 \le 1 - t\right) \\
\le \binom{p}{s} \cdot \max_{S \subseteq [p]: |S| = s} \mathbb{P}\left(\inf_{\|\Delta\|_2 = 1} \frac{1}{n} \|X_S \Delta\|_2^2 \le 1 - t\right).$$
(36)

We focus on $\mathbb{P}\left(\inf_{\|\Delta\|_2=1} \frac{1}{n} \|X_S\Delta\|_2^2 \le 1-t\right)$ for now. This bound needs to be sharp for small values of 1-t to help us bound $\mathbb{E}(\frac{1}{V^r}\mathbb{1}(V \le x))$.⁴ Define the set

$$S^{s-1} := \{ \Delta \in \mathbb{R}^s : ||\Delta||_2 = 1 \}.$$

We discretize the set \mathcal{S}^{s-1} using an ε -net and write the union bound over the net in the following way. Let $\mathcal{N}(\varepsilon)$ denote the ε -net of \mathcal{S}^{s-1} . Then for $\forall \Delta \in \mathcal{S}^{s-1}$, there exists a $\Delta' \in \mathcal{N}(\varepsilon)$ such that $\|\Delta - \Delta'\|_2 \leq \varepsilon$ and

$$||X_{S}\Delta||_{2}^{2} = ||X_{S}\Delta'||_{2}^{2} + \langle X_{S}(\Delta - \Delta'), X_{S}(\Delta + \Delta')\rangle$$

$$\geq \inf_{\Delta \in \mathcal{N}(\varepsilon)} ||X_{S}\Delta||_{2}^{2} - ||\Delta - \Delta'||_{2}||\Delta + \Delta'||_{2} \langle X_{S} \frac{(\Delta - \Delta')}{||\Delta - \Delta'||_{2}}, X_{S} \frac{(\Delta + \Delta')}{||\Delta + \Delta'||_{2}} \rangle$$

$$\geq \inf_{\Delta \in \mathcal{N}(\varepsilon)} ||X_{S}\Delta||_{2}^{2} - 2\varepsilon\sigma_{\max}^{2}(X_{S}), \tag{37}$$

where to obtain the last inequality, we have used the fact that $\|\Delta - \Delta'\|_2 \le \varepsilon$, $\|\Delta + \Delta'\|_2 \le 2$ and the Cauchy-Schwartz inequality. Define the event

$$\mathcal{D} := \left\{ \frac{1}{n} \sigma_{\max}^2(X_S) \le \frac{1}{1-t} \right\},\,$$

and let $2\varepsilon = (1-t)^2$. We use (37) to have

$$\mathbb{P}\left(\inf_{\Delta \in \mathcal{S}^{s-1}} \frac{1}{n} \| X_S \Delta \|_2^2 \le 1 - t\right)
\le \mathbb{P}\left(\inf_{\Delta \in \mathcal{N}(\varepsilon)} \frac{1}{n} \| X_S \Delta \|_2^2 \le 2(1 - t), \ \mathcal{D}\right) + \mathbb{P}(\mathcal{D}^c)
\le \frac{6^s}{(1 - t)^{2s}} \cdot \max_{\Delta \in \in \mathcal{N}(\varepsilon)} \mathbb{P}\left(\frac{1}{n} \| X_S \Delta \|_2^2 \le 2(1 - t)\right) + \mathbb{P}(\mathcal{D}^c),$$
(38)

where the last inequality uses the union bound and the result $|\mathcal{N}(\varepsilon)| \leq (3/\varepsilon)^s$ from Lemma 5. Our next step is to bound the following two quantities from (38):

 $^{^{3}}$ Many other choices of x will work as well. We do not aim to optimize the constant.

⁴Note that standard concentration bounds for the singular values of Gaussian matrices are not sharp enough.

• $\mathbb{P}\left(\frac{1}{n}||X_S\Delta||_2^2 \leq 2(1-t)\right)$: Since S is a fixed set and Δ is a fixed unit-norm vector, we know $||X_S\Delta||_2^2 \sim \chi_n^2$. Applying Lemma 8 gives $\forall t \in (1/2,1)$,

$$\mathbb{P}\left(\frac{1}{n}\|X_S\Delta\|_2^2 \le 2 - 2t\right) \le \exp\left[\frac{n}{2}\left(2t - 1 + \log(2 - 2t)\right)\right]. \tag{39}$$

• $\mathbb{P}(\mathcal{D}^c)$: A direct use of Lemma 9 yields: $\forall t \in (1 - (\sqrt{s/n} + 1)^{-2}, 1),$

$$\mathbb{P}(\mathcal{D}^c) = \mathbb{P}\left(\sigma_{\max}(X_S) \ge \sqrt{\frac{n}{1-t}}\right) \le 2\exp\left(-\frac{1}{2}\left(((1-t)^{-1/2} - 1)\sqrt{n} - \sqrt{s}\right)^2\right). \tag{40}$$

We now use the bounds (36), (38)-(40) to obtain an upper bound for $\mathbb{E}\left(\frac{1}{V^r}\mathbb{1}_{(V\leq x)}\right)$. First, the rth moment of $\frac{1}{V}\mathbb{1}_{(V\leq x)}$ can be obtained via its tails:

$$\mathbb{E}\left(\frac{1}{V^r}\mathbb{1}_{(V \le x)}\right) = \int_0^\infty ru^{r-1} \mathbb{P}\left(\frac{1}{V}\mathbb{1}_{(V \le x)} > u\right) du$$

$$= \int_0^{1/x} ru^{r-1} \mathbb{P}\left(\frac{1}{V}\mathbb{1}_{(V \le x)} > u\right) du + \int_{1/x}^\infty ru^{r-1} \mathbb{P}\left(\frac{1}{V}\mathbb{1}_{(V \le x)} > u\right) du$$

$$= x^{-r} \mathbb{P}(V \le x) + \int_{1-x}^1 r(1-t)^{-r-1} \mathbb{P}(V < 1-t) dt,$$

where we use a change of variable $t = 1 - u^{-1}$ in the last equality. Plugging the bounds of (36), (38)-(40) into the above integral, we have

$$\mathbb{E}\left(\frac{1}{V^{r}}\mathbb{1}_{(V \leq x)}\right) \leq x^{-r}\mathbb{P}\left(V \leq x\right) + r\binom{p}{s}6^{s} \int_{1-x}^{1} (1-t)^{-r-1-2s} e^{\frac{n}{2}\left(2t-1+\log(2-2t)\right)} dt
+ 2r\binom{p}{s} \int_{1-x}^{1} (1-t)^{-r-1} e^{-\frac{1}{2}\left(((1-t)^{-1/2}-1)\sqrt{n}-\sqrt{s}\right)^{2}} dt
:= x^{-r}\mathbb{P}\left(V \leq x\right) + I_{1} + I_{2}.$$
(41)

Note that (39) and (40) can be applied here, because $x < \frac{1}{2} \wedge (\sqrt{s/n} + 1)^{-2}$ for $x = e^{-8}$ and $s \le n$. For the term I_1 , we can bound as follows:

$$I_{1} = r \binom{p}{s} 6^{s} \int_{1-x}^{1} (1-t)^{\frac{n}{2}-r-1-2s} e^{\frac{n}{2}(2t-1+\log 2)} dt$$

$$\leq r \binom{p}{s} 6^{s} e^{\frac{n}{2}(1+\log 2)} \int_{1-x}^{1} (1-t)^{\frac{n}{2}-r-1-2s} dt$$

$$= r \binom{p}{s} 6^{s} e^{\frac{n}{2}(1+\log 2)} \frac{x^{\frac{n}{2}-r-2s}}{\frac{n}{2}-r-2s}$$

$$\leq \frac{r}{\frac{n}{2}-r-2s} \exp\left(s \log(6ep/s) - (3.5-0.5\log 2)n + 8r + 16s\right) = o(1), \tag{42}$$

where we used Lemma 4 in the last inequality, and the last equality can be easily verified under the scaling condition $(s \log(ep/s))/n \to 0$.

Regarding the term I_2 , we have

$$I_{2} = 2r \binom{p}{s} \int_{1-x}^{1} \exp\left((r+1)\log\frac{1}{1-t} - \frac{1}{2}\left(((1-t)^{-1/2} - 1)\sqrt{n} - \sqrt{s}\right)^{2}\right) dt$$

$$\leq 2r \binom{p}{s} \int_{1-x}^{1} \exp\left(\frac{-n/8 + r + 1}{1-t}\right) dt \leq 2r \binom{p}{s} e^{-e^{8}(n/8 - r - 1) - 8}$$

$$\leq 2r \exp\left(s\log(ep/s) - e^{8}(n/8 - r - 1) - 8\right) = o(1). \tag{43}$$

Here, the first inequality uses the fact that when $n \ge s$ and $0 \le 1 - t \le x = e^{-8}$, it holds that $\left(((1-t)^{-1/2}-1)\sqrt{n}-\sqrt{s}\right)^2 \ge \frac{n}{4(1-t)}$ and $\log\frac{1}{1-t} \le \frac{1}{1-t}$; the second inequality holds by replacing t with 1-x in the integrand; the third inequality uses Lemma 4; and the last equality is seen under the scaling condition $(s\log(ep/s))/n \to 0$.

Putting together (41)-(43) completes the proof.

Proof of Lemma 15. Given that X and z/σ have independent $\mathcal{N}(0,1)$ entries, with a conditioning (on z) argument, we can obtain

$$\mathbb{E}\left(\sup_{u\in T_s} z^T X u / \sqrt{n}\right)^q = \mathbb{E}\left(\frac{\|z\|_2}{\sqrt{n}}\right)^q \cdot \mathbb{E}\left(\sup_{u\in T_s} \langle g, u \rangle\right)^q, \quad g \in \mathcal{N}(0, I_p). \tag{44}$$

We first construct an upper bound for $\mathbb{E}(\sup_{u\in T_s}\langle g,u\rangle)^q$. Using Minkowski's inequality,

$$\left[\mathbb{E}\left(\sup_{u\in T_s} \langle g, u \rangle\right)^q\right]^{1/q} \le \left(\mathbb{E}\left|\sup_{u\in T_s} \langle g, u \rangle - \mathbb{E}\sup_{u\in T_s} \langle g, u \rangle\right|^q\right)^{1/q} + \mathbb{E}\sup_{u\in T_s} \langle g, u \rangle. \tag{45}$$

The second term above is the Gaussian complexity of T_s , and it has a sharp upper bound (e.g. Exercise 5.7 in [33]),

$$\mathbb{E}\sup_{u\in T_s}\langle g, u\rangle \le C\sqrt{s\log(ep/s)},\tag{46}$$

for some absolute constant C>0. To bound the first term in (45), let $F(g):=\sup_{u\in T_s}\langle g,u\rangle$. Then, it is clear that $F(\cdot)$ is a 1-Lipschitz function. Using the concentration of Lipschitz function of Gaussians (e.g., Theorem 2.26 in [33]), we obtain

$$\mathbb{E} \left| \sup_{u \in T(s)} \langle g, u \rangle - \mathbb{E} \sup_{u \in T(s)} \langle g, u \rangle \right|^{q} = \int_{0}^{\infty} q t^{q-1} P(|F(g) - \mathbb{E}F(g)| > t) dt$$

$$\leq \int_{0}^{\infty} 2q t^{q-1} e^{-\frac{t^{2}}{2}} dt = 2^{\frac{q}{2}} q \Gamma(\frac{q}{2}), \tag{47}$$

where $\Gamma(\cdot)$ is the Gamma function. Putting together (45)-(47) gives us

$$\mathbb{E}\Big(\sup_{u \in T_0} \langle g, u \rangle\Big)^q \le C_q \cdot (\sqrt{s \log(ep/s)})^q, \tag{48}$$

for some constant $C_q > 0$ only depending on q. Finally, note that $\frac{\|z\|_2}{\sqrt{n}}$, as a function of the standard Gaussian z/σ , is a $(\frac{\sigma}{\sqrt{n}})$ -Lipschitz function. Hence, we can use similar arguments to derive the bound for $\mathbb{E}\left(\frac{\|z\|_2}{\sqrt{n}}\right)^q$: there exists some constant $\tilde{C}_q > 0$ only depending on q such that

$$\mathbb{E}\left(\frac{\|z\|_2}{\sqrt{n}}\right)^q \le \tilde{C}_q \sigma^q. \tag{49}$$

Combining (44), (48) and (49) finishes the proof.

B.2 Proof of Proposition 2

Note that the event \mathcal{B}_{Δ} introduced in (28), has two conditions:

- $\operatorname{supp}(\hat{\beta}^L) \cup \operatorname{supp}(\tilde{\beta}^O) \cup \operatorname{supp}(\beta) \subseteq S_*$
- All the eigenvalues $X_{S_*}^T X_{S_*}/n$ lie within $[1-\Delta, 1+\Delta]$

Let us start with the second condition and prove that it happens with high probability. In other words, we would like to prove that

$$1 - \Delta \le \sigma_{\min}(X_{S_*}^T X_{S_*}/n) < \sigma_{\max}(X_{S_*}^T X_{S_*}/n) \le 1 + \Delta.$$
 (50)

This result seems to be a simple conclusion of standard results on the minimum and maximum eigenvalues of Wishart matrices such as Lemma 9. However, there is a subtle issue that needs to be addressed before such lemmas are useful: the choice of S_* depends on X. Hence, the entries of the matrix X_{S_*} are not i.i.d. Gaussian. However, [27] has proposed a clever trick to circumvent this issue. For the sake of brevity, we do not mention their trick and only use the following lemma proved in [27].

Lemma 16 (Lemma A.11 in [27]). Let $k < k^* < \min\{n, p\}$ be any (deterministic) integer. Then for any t > 0,

$$\sigma_{\min}(X_{S_*}/\sqrt{n}) > \sqrt{1 - 1/n} - \sqrt{k^*/n} - t$$

holds with probability at least $1 - e^{-nt^2/2}$. Furthermore,

$$\sigma_{\max}(X_{S_*}/\sqrt{n}) < \sqrt{1 - 1/n} + \sqrt{k^*/n} + \sqrt{8k^* \log(p/k^*)/n} + t$$

holds with probability at least $1 - e^{-nt^2/2} - (\sqrt{2}ek^*/p)^{k^*}$.

By setting $t = \sqrt{k^* \log(p/k^*)/n}$ in Lemma 16, we have as $k^*/p \to 0$ and $k^* \log(p/k^*)/n \to 0$,

$$1 - 4\sqrt{k^* \log(p/k^*)/n} \le \sigma_{\min}(X_{S_*}/\sqrt{n}) \le \sigma_{\max}(X_{S_*}/\sqrt{n}) \le 1 + 4\sqrt{k^* \log(p/k^*)/n}$$

holds with probability larger than $1 - e^{-Ck^* \log(p/k^*)}$ for some absolute constant C > 0. This further implies that under the same scaling condition,

$$1 - 10\sqrt{k^* \log(p/k^*)/n} \le \sigma_{\min}(X_{S_*}^T X_{S_*}/n) \le \sigma_{\max}(X_{S_*}^T X_{S_*}/n) \le 1 + 10\sqrt{k^* \log(p/k^*)/n}, \quad (51)$$

holds with the same high probability. This means (50) holds with probability converging to one for $\Delta = 10\sqrt{k^*\log(p/k^*)/n}$.

The following lemma aims to prove the first condition required in \mathcal{B}_{Δ} .

Lemma 17. Suppose $k^* \geq 2k$. As $k^*/p \to 0$ and $(k^* \log p)/n \to 0$,

$$\inf_{\beta \in \Theta_k} \mathbb{P}\Big(\operatorname{supp}(\hat{\beta}^L) \cup \operatorname{supp}(\tilde{\beta}^O) \cup \operatorname{supp}(\beta) \subseteq S_*\Big) \ge 1 - (k^*/p)^{\delta_{\varepsilon}},$$

for some constant $\delta_{\varepsilon} > 0$ that only depends on ε .

We should emphasize that by combining (51) with Lemma 17, we conclude the statement of Proposition 2. Hence, the last remaining step is to prove Lemma 17, as we show below.

Proof of Lemma 17. By construction, $\operatorname{supp}(\beta) \subseteq S_*$. We thus only need to show that $\operatorname{supp}(\hat{\beta}^L) \subseteq S_*$ and $\operatorname{supp}(\hat{\beta}^C) \subseteq S_*$ hold with high probability. We first prove $\operatorname{supp}(\hat{\beta}^L) \subseteq S_*$. Define \hat{b}_{S_*} as the solution of the following reduced Lasso problem:

$$\hat{b}_{S_*} \in \arg\min_{b \in \mathbb{R}^{|S_*|}} \frac{1}{2n} \|y - X_{S_*}b\|_2^2 + \lambda_{\varepsilon} \|b\|_1.$$
 (52)

In other words, if we assume that $\operatorname{supp}(\hat{\beta}^L) \subseteq S_*$ holds, then the Lasso solution $\hat{\beta}^L$ has to be the same as \hat{b}_{S_*} on S_* . Hence, as our next step we aim to show that $\hat{\beta}^L_{S_*} = \hat{b}_{S_*}$ and $\hat{\beta}^L_{S_*^c} = \mathbf{0}$.

From the KKT conditions of the Lasso problem, we conclude that if

$$||X_{S_{\varepsilon}}^{T}(y - X_{S_{\varepsilon}}\hat{b}_{S_{\varepsilon}})||_{\infty} \le n\lambda_{\varepsilon}, \tag{53}$$

then $\hat{\beta}_{S_*}^L = \hat{b}_{S_*}$ and $\hat{\beta}_{S_*^c}^L = \mathbf{0}$. To show (53), it is sufficient to prove the following two results:

$$||X_{S_*}^T X_{S_*} (\beta_{S_*} - \hat{b}_{S_*})||_{\infty} \le \frac{\varepsilon}{2} \sigma \sqrt{2n \log(p/k)},$$
 (54)

and

$$||X_{S_*^c}^T z||_{\infty} \le \left(1 + \frac{\varepsilon}{2}\right) \sigma \sqrt{2n \log(p/k)}.$$
 (55)

Before analyzing (54) and (55), we first illustrate a property of the resolvent set using a clever trick from [27]. Let $Q \in \mathbb{R}^{n \times n}$ be an orthogonal matrix (measurable with respect to z) such that

$$Qz = (||z||_2, 0, \dots, 0).$$

Let $\tilde{\omega} \in \mathbb{R}^{1 \times p}$, $\tilde{W} \in \mathbb{R}^{(n-1) \times p}$ and

$$W = \begin{bmatrix} \tilde{\omega} \\ \tilde{W} \end{bmatrix} := QX.$$

The independence between Q and X implies that W is still a Gaussian random matrix with i.i.d. $\mathcal{N}(0,1)$ entries. Furthermore,

$$X_i^T z = (QX_i)^T (Qz) = ||z||_2 (QX_i)_1 = ||z||_2 \tilde{\omega}_i.$$
(56)

This indicates that S_* is the union of S and k^*-k indices in $\{1,\ldots,p\}\setminus S$ of the largest $|\tilde{\omega}_i|$. Since $\tilde{\omega}$ and \tilde{W} are independent, \tilde{W} and S_* are also independent. Thus, $\tilde{W}_{S_*^c}\in\mathbb{R}^{(n-1)\times(p-k^*)}$ and $\tilde{W}_{S_*}\in\mathbb{R}^{(n-1)\times k^*}$ are both Gaussian random matrices with i.i.d. $\mathcal{N}(0,1)$ entries. Now we aim to use this trick to prove (54) and (55).

• Proving $||X_{S_c}^T X_{S_*}(\beta_{S_*} - \hat{b}_{S_*})||_{\infty} \leq \frac{\varepsilon}{2} \sigma \sqrt{2n \log(p/k)}$. We have

$$X_{S_{*}^{c}}^{T}X_{S_{*}}(\beta_{S_{*}} - \hat{b}_{S_{*}}) = X_{S_{*}^{c}}^{T}X_{S_{*}}(X_{S_{*}}^{T}X_{S_{*}})^{-1}(X_{S_{*}}^{T}(y - X_{S_{*}}\hat{b}_{S_{*}}) - X_{S_{*}}^{T}z)$$

$$= X_{S_{*}^{c}}^{T}Q^{T}\underbrace{QX_{S_{*}}(X_{S_{*}}^{T}X_{S_{*}})^{-1}(X_{S_{*}}^{T}(y - X_{S_{*}}\hat{b}_{S_{*}}) - X_{S_{*}}^{T}z)}_{:=\xi}.$$
(57)

We first derive the bound for $\|\xi\|_2$. We have

$$\|\xi\|_{2} \leq \|X_{S_{*}}(X_{S_{*}}^{T}X_{S_{*}})^{-1}\|_{2} \cdot \|X_{S_{*}}^{T}(y - X_{S_{*}}\hat{b}_{S_{*}}) - X_{S_{*}}^{T}z\|_{2}.$$

Hence, our next goal is to bound the three terms $||X_{S_*}(X_{S_*}^TX_{S_*})^{-1}||_2$, $||X_{S_*}^T(y-X_{S_*}\hat{b}_{S_*})||_2$, and $||X_{S_*}^Tz||_2$. Since \hat{b}_{S_*} is the solution to the reduced Lasso problem (52), its KKT condition implies that

$$||X_{S_*}^T(y - X_{S_*}\hat{b}_{S_*})||_{\infty} \le n\lambda_{\varepsilon} = (1 + \varepsilon)\sigma\sqrt{2n\log(p/k)}.$$
 (58)

Furthermore, if we set t = 1/2 in Lemma 16 we obtain

$$\|\sqrt{n}X_{S_*}(X_{S_*}^TX_{S_*})^{-1}\|_2 \le \left(\sqrt{1-1/n} - \sqrt{k^*/n} - 1/2\right)^{-1} < 2.01$$
 (59)

with probability at least $1 - e^{-n/8}$ for sufficiently large n, where in the last step we have used $k^*/n \to 0$. Finally, we use the following lemma to bound $\|X_{S_*}^T z\|_2$.

Lemma 18 (Lemma A.7 in [27]). Let $1 \le k^* < p$ be any (deterministic) integer, then

$$\sup_{|T|=k^*} \|X_T^T z\|_2 \le \sigma \sqrt{32nk^* \log(p/k^*)}$$

with probability at least $1 - e^{-n/2} - (\sqrt{2}ek^*/p)^{k^*}$. Above, the supremum is taken over all the subsets of $\{1, \ldots, p\}$ with cardinality k^* .

Combining (58), (59) and Lemma 18, we have

$$\|\xi\|_{2} \leq \|X_{S_{*}}(X_{S_{*}}^{T}X_{S_{*}})^{-1}\|_{2} \cdot (\|X_{S_{*}}^{T}(y - X_{S_{*}}\hat{b}_{S_{*}})\|_{2} + \|X_{S_{*}}^{T}z\|_{2})$$

$$\leq 2.01n^{-1/2} \left(\sqrt{k^{*}}n\lambda_{\varepsilon} + 4\sigma\sqrt{2nk^{*}\log(p/k^{*})}\right)$$

$$\leq 2.01 \left((1+\varepsilon)\sqrt{2} + 4\sqrt{2}\right)\sigma\sqrt{k^{*}\log(p/k)}$$
(60)

with probability at least $1 - e^{-n/2} - (\sqrt{2}ek^*/p)^{k^*} - e^{-n/8} := 1 - P_{\xi}$ for large enough n; where in the third line we have used $k^* \ge k$.

We continue from (57) to obtain

$$X_{S^c}^T X_{S_*} (\beta_{S_*} - \hat{b}_{S_*}) = (\tilde{\omega}_{S^c}^T, 0) \xi + (0, \tilde{W}_{S^c}^T) \xi.$$
(61)

For the first term, we have

$$\|(\tilde{\omega}_{S_{\varepsilon}}^{T}, 0)\xi\|_{\infty} = |\xi_{1}| \cdot \|\tilde{\omega}_{S_{\varepsilon}}^{T}\|_{\infty} \le \|\xi\|_{2} \cdot \|\tilde{\omega}_{S_{\varepsilon}}^{T}\|_{\infty}.$$
(62)

Let $\zeta_1,\ldots,\zeta_{p-k}$ be i.i.d. $\mathcal{N}(0,1)$, and $|\zeta|_{(1)} \geq \ldots \geq |\zeta|_{(p-k)}$ denote the order statistics of $(|\zeta_1|,\ldots,|\zeta_{p-k}|)$. Based on (56), we recognize that $\tilde{\omega}_{S_s^c} \in \mathbb{R}^{p-k^*}$ is a subvector of a standard (p-k)-dimensional Gaussian by excluding the k^*-k largest (in absolute value) coordinates. Hence, to bound the term $\|(\tilde{\omega}_{S_s^c}^T,0)\xi\|_{\infty}$, we need to find an upper bound for $|\zeta|_{(k^*-k+1)}$. For this purpose we can use standard results on Gaussian order statistics such as Lemma 11. Applying Lemma 11 with $u=\frac{\varepsilon}{12+3\varepsilon}\sqrt{2\log\frac{2p}{k}}$, we obtain that

$$\|\tilde{\omega}_{S_*^c}\|_{\infty} \stackrel{d}{=} |\zeta|_{(k^*-k+1)} \le \sqrt{2\log\frac{2(p-k)}{k^*-k}} + \frac{\epsilon}{12+3\epsilon} \sqrt{2\log\frac{2p}{k}}$$

$$\le \frac{12+4\epsilon}{12+3\epsilon} \sqrt{2\log\frac{2p}{k}},$$
(63)

with probability at least $1-(\frac{k}{2p})^{\frac{\epsilon^2}{(12+3\epsilon)^2}}$; where in the last inequality we used the condition $k^* \geq 2k$. Thus, from (60), (62) and (63), we reach that with probability at least $1-P_{\xi}-(\frac{k}{2p})^{\frac{\epsilon^2}{(12+3\epsilon)^2}}$,

$$\|(\tilde{\omega}_{S_c^c}^T, 0)\xi\|_{\infty} \le C_{\epsilon}\sigma\sqrt{k^*}\log(2p/k),\tag{64}$$

for some constant $C_{\epsilon} > 0$ only depending on ϵ .

Our next goal is to bound the second term in (61), i.e. $\|(0, \tilde{W}_{S_*}^T)\xi\|_{\infty}$. From the discussion about (56), we conclude that $\tilde{W}_{S_*^c}$ is independent of ξ since $\tilde{W}_{S_*^c}$ is independent of (W_{S_*}, z, S_*) . Furthermore, conditional on ξ , the vector $(0, \tilde{W}_{S_*^c}^T)\xi$ has independent Gaussian coordinates with mean zero and variance $\xi_2^2 + \cdots + \xi_n^2$. Hence, we have

$$\|(0, \tilde{W}_{S_*^c}^T)\xi\|_{\infty} \stackrel{d}{=} \sqrt{\xi_2^2 + \dots + \xi_n^2} \cdot \max_{1 \le i \le (p-k^*)} |\zeta_i| \le \|\xi\|_2 \cdot \max_{1 \le i \le (p-k^*)} |\zeta_i|,$$

where $\zeta_1, \ldots, \zeta_{p-k^*} \stackrel{i.i.d.}{\sim} \mathcal{N}(0,1)$, independent of ξ . Lemma 10 implies that $\max_{1 \leq i \leq (p-k^*)} |\zeta_i| \leq 2\sqrt{2\log p}$, holds with probability at least $1-p^{-1}$. This together with (60) gives

$$\mathbb{P}\left(\|(0, \tilde{W}_{S_{\epsilon}^{*}}^{T})\xi\|_{\infty} \leq \tilde{C}_{\epsilon}\sigma\sqrt{k^{*}\log(p/k)}\sqrt{\log p}\right) \geq 1 - P_{\xi} - p^{-1},\tag{65}$$

for some constant $\tilde{C}_{\epsilon} > 0$ only depending on ϵ . Putting together (61), (64) and (65) yields

$$||X_{S_{\varepsilon}}^T X_{S_*} (\beta_{S_*} - \hat{b}_{S_*})||_{\infty} \le \bar{C}_{\varepsilon} \sigma \sqrt{k^* \log(2p/k)} \sqrt{\log(2p)}, \tag{66}$$

with probability at least $1 - P_{\xi} - (\frac{k}{2p})^{\frac{\varepsilon^2}{(12+3\varepsilon)^2}} - \frac{1}{p}$ for some constant $\bar{C}_{\varepsilon} > 0$ only depending on ε .

• Proving $||X_{S_*^c}^T z||_{\infty} \leq \left(1 + \frac{\varepsilon}{2}\right) \sigma \sqrt{2n \log(p/k)}$. First, using the concentration of the Gaussian norm (Theorem 3.1.1 in [30]), we have $\mathbb{P}\left(||z||_2 \leq (1 + \varepsilon/4)\sigma\sqrt{n}\right) \geq 1 - 2e^{-C\varepsilon^2 n}$ with some absolute constant C > 0. Combining this with (56) and (63), we have

$$||X_{S^c}^T z||_{\infty} = ||z||_2 \cdot ||\tilde{\omega}_{S_c^c}||_{\infty} \le (1 + \varepsilon/3)\sigma\sqrt{2n\log(2p/k)}$$

$$\tag{67}$$

holds with probability at least $1 - 2e^{-C\varepsilon^2 n} - (\frac{k}{2p})^{\frac{\varepsilon^2}{(12+3\varepsilon)^2}}$.

Now, under the given condition $k^* \geq 2k$, $\frac{k^*}{p} \to 0$, $\frac{k^* \log p}{n} \to 0$, it is straightforward to confirm: when n is sufficiently large, (66) and (67) imply that the two results (54) and (55) hold with probability at least $1 - (k^*/p)^{c_{\epsilon}}$ for some constant $c_{\epsilon} > 0$ only depending on ϵ . As a result,

$$\inf_{\beta \in \Theta_k} \mathbb{P}\left(\operatorname{supp}(\hat{\beta}^L) \subseteq S_*\right) \ge 1 - (k^*/p)^{c_{\epsilon}}.$$

The last remaining step of the proof is to show that $\operatorname{supp}(\tilde{\beta}^O) \subseteq S_*$ with high probability. Recalling $\tilde{\beta}^O$ defined in (25), the property of the resolvent set S_* and soft thresholding function $\eta_{\lambda_{\epsilon}}(\cdot)$ implies

$$\inf_{\beta \in \Theta_k} \mathbb{P}\left(\operatorname{supp}(\tilde{\beta}^O) \subseteq S_*\right) \ge \inf_{\beta \in \Theta_k} \mathbb{P}\left(\|X_{S_*^c}^T z\|_{\infty} \le n\lambda_{\varepsilon}\right) \ge 1 - (k^*/p)^{\tilde{c}_{\epsilon}},$$

where the last inequality was implied by (67) for large enough n.

B.3 Proof of Proposition 3

As pointed out in Section 4.2, we will follow the same proof strategy in [1] and do a refined analysis in some of the steps, to be able to obtain the optimal rates for Lasso with a smaller tuning parameter value $\lambda_{\varepsilon} = (1+\varepsilon)\sigma\sqrt{\frac{2\log(p/k)}{n}}$.

Inspired by [1], throughout the proof, we will use the following notation: for a given $u = (u_1, \ldots, u_p) \in \mathbb{R}^p$, let $|u|_{(1)} \ge |u|_{(2)} \ge \cdots \ge |u|_{(p)}$ denote the order statistics of $(|u_1|, \ldots, |u_p|)$, and define

$$H(u) := \sigma(1+\delta_2) \left(\sum_{j=1}^k |u|_{(j)} 4\sqrt{\frac{\log(2p/j)}{n}} + (1+\delta_1) \sum_{j=k+1}^p |u|_{(j)} \sqrt{\frac{2\log(p/k)}{n}} \right), \tag{68}$$

$$G(u) := \sigma(1 + \delta_2)\delta_2^{-1} \frac{\sqrt{2\log(1/\delta_3)}}{n(1 + \delta_0)} \|Xu\|_2, \tag{69}$$

where $\delta_0, \delta_1, \delta_2, \delta_3 \in (0, 1)$ are some constants. The following lemma shows the importance of the two terms defined above.

Lemma 19 (Bound on the stochastic error). Let $z \sim \mathcal{N}(0, \sigma^2 I_n)$ and $X \in \mathbb{R}^{n \times p}$ be a fixed matrix such that $\max_{j \in [p]} ||X_j||_2 \le (1 + \delta_0) \sqrt{n}$. For any $\delta_0, \delta_1, \delta_2, \delta_3 \in (0, 1)$, there exist a constant $C_{\delta_1} > 0$ only depending on δ_1 such that the event

$$\left\{ \frac{1}{n} z^T X u \le (1 + \delta_0) \cdot \max \left(H(u), G(u) \right), \ \forall u \in \mathbb{R}^p \right\}$$
 (70)

holds with probability at least $1 - \delta_3$, as long as $p/k \geq C_{\delta_1}$.

Proof. The lemma is a modified version of Theorem 4.1 in [1], tailored for the smaller tuning parameter value λ_{ε} . The proof is similar, hence we do not repeat all the details and only mention the major difference in bounding $\frac{1}{n}z^TXu$. Let $g_j = \frac{z^TX_j}{\sigma\sqrt{n}(1+\delta_0)}, j=1,2,\ldots,p$. We have

$$\frac{1}{n}z^{T}Xu \leq (1+\delta_{0})\sigma \sum_{j=1}^{p} \frac{1}{\sqrt{n}}|g|_{(j)}|u|_{(j)}
\leq (1+\delta_{0})\sigma \sum_{j=1}^{k} \left(|u|_{(j)}4\sqrt{\frac{\log(2p/j)}{n}}\right) \cdot \left(\frac{|g|_{(j)}}{4\sqrt{\log(2p/j)}}\right)
+ (1+\delta_{0})\sigma \sum_{j=k+1}^{p} \left((1+\delta_{1})|u|_{(j)}\sqrt{\frac{2\log(p/k)}{n}}\right) \cdot \left(\frac{|g|_{(j)}}{(1+\delta_{1})\sqrt{2\log(p/k)}}\right)
\leq \frac{(1+\delta_{0})H(u)}{1+\delta_{2}} \cdot \left(\max_{1\leq j\leq k} \frac{|g|_{(j)}}{4\sqrt{\log(2p/j)}}\right) \vee \frac{|g|_{(k+1)}}{(1+\delta_{1})\sqrt{2\log(p/k)}}.$$
(71)

Define

$$\mathcal{T} = \left\{ u \in \mathbb{R}^p : \max \left(H(u), G(u) \right) \le \frac{1 + \delta_2}{1 + \delta_0} \right\}.$$

The rest is to bound $\max_{u \in \mathcal{T}} \frac{1}{n} z^T X u$, using concentration of Gaussian measure around the median and using (71) together with Lemma 13 to bound the median. The detail is similar to that in [1] and is hence skipped.

Lemma 19 enables us to use similar proof steps as the ones presented in Theorem 4.2 of [1], to obtain optimal rates for Lasso under the smaller regularization $\lambda_{\varepsilon} = (1+\varepsilon)\sigma\sqrt{\frac{2\log(p/k)}{n}}$. It is this regularization λ_{ε} that eventually yields the sharp constant in the asymptotic minimax risk (see discussions in Section 4.2).

Lemma 20. Assume $p \ge 2k$ and $(1 + \varepsilon) - (1 + \delta_0)(1 + \delta_1)(1 + \delta_2) > 0$, and define

$$c_0 = \frac{4\sqrt{2}(1+\delta_0)(1+\delta_2) + 1 + \epsilon}{(1+\varepsilon) - (1+\delta_0)(1+\delta_1)(1+\delta_2)}.$$
 (72)

Consider model (1) with any fixed design $X \in \mathcal{A}(\delta_0, c_0, k)$ in (16). Then, on the event (70), the Lasso estimator $\hat{\beta}^L$ in (13) with tuning parameter λ_{ε} satisfies

$$\|\hat{\beta}^L - \beta\|_2 \le C_1(\delta_0, \delta_2, \epsilon)\sigma\sqrt{\frac{k\log(p/k)}{n}} + C_2(\delta_0, \delta_2, \epsilon)\sigma\frac{\log(1/\delta_3)}{\sqrt{nk\log(p/k)}},$$

for all $\beta \in \Theta_k$. Here,

$$C_1(\delta_0, \delta_2, \epsilon) := \frac{8(1 + \delta_0)(1 + \delta_2) + \sqrt{2}(1 + \epsilon)}{(1 - \delta_0)^2},$$

$$C_2(\delta_0, \delta_2, \epsilon) := \frac{4\sqrt{2}(1 + \delta_0)(1 + \delta_2) + 1 + \epsilon}{16\sqrt{2}(1 + \delta_0)^2 \delta_2^2}.$$

Proof of Lemma 20. We start with the two point inequality that has often been used to establish sharp oracle inequalities for Lasso (e.g. Lemma A.2 in [1]; Chapter 2.4 in [29]):

$$\frac{1}{n} \|X(\hat{\beta}^L - \beta)\|_2^2 \le \frac{1}{n} z^T X(\hat{\beta}^L - \beta) + \lambda_{\varepsilon} \|\beta\|_1 - \lambda_{\varepsilon} \|\hat{\beta}^L\|_1.$$
 (73)

Denoting $u = \hat{\beta}^L - \beta$, $S = \text{supp}(\beta)$ and using the following standard calculations

$$\|\beta\|_1 - \|\hat{\beta}^L\|_1 = \|\beta\|_1 - \|\beta + u\|_1 = \|\beta_S\|_1 - \|\beta_S + u_S\|_1 - \|u_{S^c}\|_1 \le \|u_S\|_1 - \|u_{S^c}\|_1,$$

we can continue from (73) to obtain that on the event (70),

$$\frac{1}{n} \|Xu\|_2^2 \le (1 + \delta_0) \max(H(u), G(u)) + \lambda_{\varepsilon} (\|u_S\|_1 - \|u_{S^c}\|_1)$$

To have an upper bound for H(u), define

$$\tilde{H}(u) := \sigma(1 + \delta_2) \left(8\|u\|_2 \sqrt{\frac{k \log(p/k)}{n}} + (1 + \delta_1) \sqrt{\frac{2 \log(p/k)}{n}} \|u_{S^c}\|_1 \right). \tag{74}$$

Using the Cauchy-Schwarz inequality,

$$\begin{split} H(u) &\leq \sigma(1+\delta_2) \bigg(4\|u\|_2 \sqrt{\sum_{j=1}^k \frac{\log(2p/j)}{n}} + (1+\delta_1) \sqrt{\frac{2\log(p/k)}{n}} \sum_{j=k+1}^p |u|_{(j)} \bigg) \\ &\leq \sigma(1+\delta_2) \bigg(4\|u\|_2 \sqrt{\frac{k\log(2ep/k)}{n}} + (1+\delta_1) \sqrt{\frac{2\log(p/k)}{n}} \sum_{j=k+1}^p |u|_{(j)} \bigg) \leq \tilde{H}(u), \end{split}$$

where the second inequality holds because $\sum_{j=1}^k \log(2p/j) = k \log(2p) - \log(k!) \le k \log(2ep/k)$ by Stirling's formula, and the third one is due to the fact that $\log(2ep/k) \le 4 \log(p/k)$ (as $p \ge 2k$) and $\sum_{j=k+1}^p |u|_{(j)} \le ||u_{S^c}||_1$. Hence, we can further proceed to have that on the event (70),

$$\frac{1}{n} \|Xu\|_{2}^{2} \le (1 + \delta_{0}) \max(\tilde{H}(u), G(u)) + \lambda_{\varepsilon}(\|u_{S}\|_{1} - \|u_{S^{c}}\|_{1}). \tag{75}$$

Based on (75), we will derive the bound for $||u||_2$ in two different cases:

• If $G(u) > \tilde{H}(u)$, then by comparing the two expressions given in (69) and (74) we have

$$||u||_2 \le \frac{\sqrt{2\log(1/\delta_3)}}{8(1+\delta_0)\delta_2\sqrt{nk\log(p/k)}}||Xu||_2.$$
(76)

This together with (75) gives us

$$\frac{1}{n} \|Xu\|_{2}^{2} \leq (1+\delta_{0})G(u) + \sqrt{k}\lambda_{\varepsilon} \|u\|_{2}
\leq \sigma(1+\delta_{2}) \frac{\sqrt{2\log(1/\delta_{3})}}{n\delta_{2}} \|Xu\|_{2} + \sigma(1+\varepsilon)\sqrt{\frac{2k\log(p/k)}{n}} \|u\|_{2}
\leq \left(1+\delta_{2} + \frac{1+\varepsilon}{4\sqrt{2}(1+\delta_{0})}\right) \cdot \frac{\sigma\sqrt{2\log(1/\delta_{3})}}{n\delta_{2}} \|Xu\|_{2}.$$

Solving the above for $||Xu||_2$ and then plugging it into (76), we obtain

$$||u||_2 \le C_2(\delta_0, \delta_2, \varepsilon) \frac{\sigma \log(1/\delta_3)}{\sqrt{nk \log(p/k)}}.$$

• If $G(u) \leq \tilde{H}(u)$, using $||u_S||_1 \leq \sqrt{k}||u||_2$ in (75) yields

$$0 \leq \frac{1}{n} \|Xu\|_{2}^{2} \leq \left(8(1+\delta_{0})(1+\delta_{2}) + \sqrt{2}(1+\varepsilon)\right) \sigma \sqrt{\frac{k \log(p/k)}{n}} \|u\|_{2} - \left((1+\varepsilon) - (1+\delta_{0})(1+\delta_{1})(1+\delta_{2})\right) \sigma \sqrt{\frac{2 \log(p/k)}{n}} \|u_{S^{c}}\|_{1},$$
(77)

which implies $||u_{S^c}||_1 \leq c_0 \sqrt{k} ||u||_2$ with c_0 defined in (72). This further shows that $||u||_1 = ||u_S||_1 + ||u_{S^c}||_1 \leq (1+c_0)\sqrt{k}||u||_2$. Therefore, applying $\theta(k,c_0)$ in (6) and the condition $X \in \mathcal{A}(\delta_0,c_0,k)$, we obtain

$$(1 - \delta_0)^2 ||u||_2^2 \le \frac{1}{n} ||Xu||_2^2.$$

Combining this result with (77) gives us

$$(1 - \delta_0)^2 ||u||_2^2 \le \left(8(1 + \delta_0)(1 + \delta_2) + \sqrt{2}(1 + \varepsilon)\right) \sigma \sqrt{\frac{k \log(p/k)}{n}} ||u||_2,$$

leading to the bound

$$||u||_2 \le C_1(\delta_0, \delta_2, \epsilon) \sigma \sqrt{\frac{k \log(p/k)}{n}}.$$

Putting together Lemmas 19 and 20, we can integrate the tail bounds to obtain bounds for q-moments $\mathbb{E}(\|\hat{\beta}^L - \beta\|_2^q \mathbb{1}_{\mathcal{A}})$.

Lemma 21. For any given constants ε , δ_0 , δ_1 , $\delta_2 \in (0,1)$, assume $(1+\varepsilon) > (1+\delta_0)(1+\delta_1)(1+\delta_2)$. Consider model (1) with isotropic Gaussian design. Then, there exists a constant $c_{\delta_1} > 0$ only depending on δ_1 such that as long as $p/k > c_{\delta_1}$, the Lasso estimator $\hat{\beta}^L$ in (13) with tuning parameter λ_{ε} satisfies $\forall q \in [2, \infty)$,

$$\sup_{\beta \in \Theta_k} \mathbb{E} \left(\| \hat{\beta}^L - \beta \|_2^q \, \mathbb{1}_{\mathcal{A}} \right) \le c_q \sigma^q \cdot \left[\left(C_1(\delta_0, \delta_2, \epsilon) \sqrt{\frac{k \log(p/k)}{n}} \right)^q + \left(\frac{C_2(\delta_0, \delta_2, \epsilon)}{\sqrt{nk \log(p/k)}} \right)^q \right]. \tag{78}$$

Here, $c_q > 0$ is a constant that only depends on q, and the constants $C_1(\delta_0, \delta_2, \epsilon), C_2(\delta_0, \delta_2, \epsilon), c_0$ and the event $\mathcal{A} = \mathcal{A}(\delta_0, c_0, k)$ are the same from Lemma 20.

Proof. Based on Lemmas 19 and 20, a conditioning (on X) argument shows

$$\mathbb{P}\left(\|\hat{\beta}^{L} - \beta\|_{2}\mathbb{1}_{\mathcal{A}} \ge C_{1}(\delta_{0}, \delta_{2}, \epsilon)\sigma\sqrt{\frac{k\log(p/k)}{n}} + \frac{C_{2}(\delta_{0}, \delta_{2}, \epsilon)\sigma}{\sqrt{nk\log(p/k)}}t\right) \le e^{-t}, \quad \forall t > 0.$$
 (79)

Denote $a_n := C_1(\delta_0, \delta_2, \epsilon) \sigma \sqrt{\frac{k \log(p/k)}{n}}, b_n := \frac{C_2(\delta_0, \delta_2, \epsilon) \sigma}{\sqrt{nk \log(p/k)}}$. We can compute the qth moment of $\|\hat{\beta}^L - \beta\|_2 \mathbb{1}_{\mathcal{A}}$ by integrating its tails in the following way:

$$\mathbb{E}\left(\|\hat{\beta}^{L} - \beta\|_{2}^{q} \mathbb{1}_{\mathcal{A}}\right) = \int_{0}^{\infty} q s^{q-1} \mathbb{P}\left(\|\hat{\beta}^{L} - \beta\|_{2} \mathbb{1}_{\mathcal{A}} > s\right) ds$$

$$= \int_{0}^{a_{n}} q s^{q-1} \mathbb{P}\left(\|\hat{\beta}^{L} - \beta\|_{2} \mathbb{1}_{\mathcal{A}} > s\right) ds + \int_{a_{n}}^{\infty} q s^{q-1} \mathbb{P}\left(\|\hat{\beta}^{L} - \beta\|_{2} \mathbb{1}_{\mathcal{A}} > s\right) ds$$

The first integral can be easily bounded as

$$\int_0^{a_n} q s^{q-1} \mathbb{P} \Big(\| \hat{\beta}^L - \beta \|_2 \mathbb{1}_{\mathcal{A}} > s \Big) ds \le \int_0^{a_n} q s^{q-1} ds = a_n^q.$$

For the second one, by a change of variable $s = a_n + b_n t$, we have

$$\int_{a_n}^{\infty} q s^{q-1} \mathbb{P} \Big(\| \hat{\beta}^L - \beta \|_2 \mathbb{1}_{\mathcal{A}} > s \Big) ds
= \int_{0}^{\infty} q b_n (a_n + b_n t)^{q-1} \mathbb{P} \Big(\| \hat{\beta}^L - \beta \|_2 \mathbb{1}_{\mathcal{A}} > a_n + b_n t \Big) dt
\leq \int_{0}^{\infty} q b_n (a_n + b_n t)^{q-1} e^{-t} dt \leq \int_{0}^{\infty} q b_n 2^{q-2} (a_n^{q-1} + b_n^{q-1} t^{q-1}) e^{-t} dt
= q 2^{q-2} b_n a_n^{q-1} + q 2^{q-2} \Gamma(q) b_n^q \leq q 2^{q-2} (1 + \Gamma(q)) (a_n^q + b_n^q).$$

Here, the first inequality is due to (79); the second inequality is based on the basic inequality $(c+d)^m \leq 2^{m-1}(c^m+d^m), \forall m \geq 1, c, d \geq 0$; and the third inequality holds because $b_n a_n^{q-1} \leq a_n^q + b_n^q$.

Based on Lemma 21, we can choose

$$\delta_0 = \delta_1 = \delta_2 = \left(1 + \frac{\varepsilon}{2}\right)^{\frac{1}{3}} - 1.$$

As a result, $(1 + \delta_0)(1 + \delta_1)(1 + \delta_2) = 1 + \frac{\varepsilon}{2} < 1 + \varepsilon$, and the constant c_0 of (72) becomes exactly the one specified in (18) (so is δ_0). Then, applying Lemma 21 and noticing that the first term in (78) is the dominating term as $p/k \to \infty$, completes the proof of Proposition 3.

References

- [1] Pierre C Bellec, Guillaume Lecué, and Alexandre B Tsybakov. "Slope meets lasso: improved oracle bounds and optimality". In: *The Annals of Statistics* 46.6B (2018), pp. 3603–3642.
- [2] Alexandre Belloni, Victor Chernozhukov, and Lie Wang. "Square-root lasso: pivotal recovery of sparse signals via conic programming". In: *Biometrika* 98.4 (2011), pp. 791–806.
- [3] P. J. Bickel, Y. Ritov, and A. Tsybakov. "Simultaneous analysis of LASSO and Dantzig selector". In: *The Annals of Statistics* 37.4 (2009), pp. 1705–1732.
- [4] Małgorzata Bogdan et al. "SLOPE—adaptive variable selection via convex optimization". In: *The annals of applied statistics* 9.3 (2015), p. 1103.
- [5] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. Concentration Inequalities: A Nonasymptotic Theory of Independence. Oxford University Press, 2013.
- [6] Peter Bühlmann and Sara Van De Geer. Statistics for high-dimensional data: methods, theory and applications. Springer Science & Business Media, 2011.
- [7] Cristina Butucea et al. "Variable selection with Hamming loss". In: *The Annals of Statistics* 46.5 (2018), pp. 1837–1875.
- [8] Emmanuel J Candes and Mark A Davenport. "How well can we estimate a sparse vector?" In: Applied and Computational Harmonic Analysis 34.2 (2013), pp. 317–323.
- [9] Emmanuel J Candes and Terence Tao. "Decoding by linear programming". In: *IEEE transactions on information theory* 51.12 (2005), pp. 4203–4215.
- [10] Arnak S Dalayan, Mohamed Hebiri, and Johannes Lederer. "On the prediction performance of the Lasso". In: *Bernoulli* 23.1 (2017), pp. 552–581.
- [11] David L Donoho and Iain M Johnstone. "Minimax risk over ℓ_p -balls for ℓ_q -error". In: *Probability theory and related fields* 99 (1994), pp. 277–303.
- [12] David L Donoho et al. "Maximum entropy and the nearly black object". In: Journal of the Royal Statistical Society: Series B (Methodological) 54.1 (1992), pp. 41–67.
- [13] Jianqing Fan et al. Statistical foundations of data science. CRC press, 2020.
- [14] Sara A van de Geer and Peter Bühlmann. "On the conditions used to prove oracle results for the Lasso". In: *Electronic Journal of Statistics* 3 (2009), pp. 1360–1392.
- [15] Yilin Guo, Haolei Weng, and Arian Maleki. "Signal-to-noise ratio aware minimaxity and higher-order asymptotics". In: arXiv preprint arXiv:2211.05954 (2022).
- [16] Trevor Hastie, Robert Tibshirani, and Martin Wainwright. Statistical learning with sparsity: the lasso and generalizations. CRC press, 2015.
- [17] Trevor Hastie et al. The elements of statistical learning: data mining, inference, and prediction. Vol. 2. Springer, 2009.
- [18] Shirin Jalali, Arian Maleki, and Richard Baraniuk. "Minimum complexity pursuit: Stability analysis". In: 2012 IEEE International Symposium on Information Theory Proceedings. 2012, pp. 1857–1861. DOI: 10.1109/ISIT.2012.6283602.
- [19] Pengsheng Ji and Jiashun Jin. "Optimal phase diagram in high-dimensional variable selection". In: *The Annals of Statistics* 40.1 (2012), pp. 73–103.
- [20] Iain M. Johnstone. Gaussian estimation: Sequence and wavelet models. 2019.
- [21] Guillaume Lecué and Shahar Mendelson. "Sparse recovery under weak moment assumptions". In: Journal of the European Mathematical Society 19.3 (2017), pp. 881–904.
- [22] Mohamed Ndaoud and Alexandre B Tsybakov. "Optimal variable selection and adaptive noisy compressed sensing". In: *IEEE Transactions on Information Theory* 66.4 (2020), pp. 2517– 2532.

- [23] Sahand N Negahban et al. "A Unified Framework for High-Dimensional Analysis of M-Estimators with Decomposable Regularizers". In: *Statistical Science* 27.4 (2012), pp. 538–557.
- [24] Garvesh Raskutti, Martin J Wainwright, and Bin Yu. "Minimax rates of estimation for high-dimensional linear regression over ℓ_q -balls". In: *IEEE transactions on information theory* 57.10 (2011), pp. 6976–6994.
- [25] Garvesh Raskutti, Martin J Wainwright, and Bin Yu. "Restricted eigenvalue properties for correlated Gaussian designs". In: The Journal of Machine Learning Research 11 (2010), pp. 2241–2259.
- [26] Mark Rudelson and Shuheng Zhou. "Reconstruction From Anisotropic Random Measurements". In: *IEEE Transactions on Information Theory* 59.6 (2013), pp. 3434–3447.
- [27] Weijie Su and Emmanuel Candes. "SLOPE is adaptive to unknown sparsity and asymptotically minimax". In: (2016).
- [28] Robert Tibshirani. "Regression shrinkage and selection via the lasso". In: Journal of the Royal Statistical Society Series B: Statistical Methodology 58.1 (1996), pp. 267–288.
- [29] Sara A Van de Geer et al. Estimation and testing under sparsity. Springer, 2016.
- [30] Roman Vershynin. High-dimensional probability: An introduction with applications in data science. Vol. 47. Cambridge university press, 2018.
- [31] Roman Vershynin. "Introduction to the non-asymptotic analysis of random matrices". In: arXiv preprint arXiv:1011.3027 (2010).
- [32] Nicolas Verzelen. "Minimax risks for sparse regressions: Ultra-high dimensional phenomenons". In: (2012).
- [33] Martin J Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*. Vol. 48. Cambridge university press, 2019.
- [34] Fei Ye and Cun-Hui Zhang. "Rate minimaxity of the Lasso and Dantzig selector for the ℓ_q loss in ℓ_r balls". In: The Journal of Machine Learning Research 11 (2010), pp. 3519–3540.