ON THE CONSTRUCTION OF A FAMILY OF WELL -POSED APPROXIMATE FORMULATIONS FOR THE STATIONARY STOKES PROBLEM USING AN EXTENDED SYSTEM

CĂTĂLIN - LIVIU BICHIR

ABSTRACT. We introduce an exact parameterized extended system such that, under adequate data, between the components of its solution, there is the solution of the weak formulation of the homogeneous Dirichlet problem for the stationary Stokes equations. In the extended system, we introduce the momentum equation together with two other forms of this one. This allows us to reformulate, for the stationary case, the consistent pressure Poisson equation of Sani, Shen, Pironneau, Gresho [Int. J. Numer. Meth. Fluids, 50 (2006), pp. 673-682], from the unsteady case. In this way, we can retain the information we need for the approximate pressure on the boundary. We obtain a parameterized perturbed pressure Poisson equation for the stationary Stokes problem. We prove that to solve the stationary Stokes problem is equivalent to solve a problem for the momentum equation, the parameterized equation and the equation that defines the Laplace operator acting on velocity. The approximation of this last problem give a family of well - posed approximate formulations. The solution of each element of this family approximates the solution of the stationary Stokes problem. Some necessary variants of existing results on general Banach spaces are also developed. These concern the density of subspaces related to isomorphisms and the connection between the exact and the approximate problems. First, the well - posedness of the exact extended system is proved in some dense subspaces. The well posed approximate problem does not necessitate the discrete inf-sup condition. The paper is also related to the existing discussions on the boundary conditions for the pressure and on the imposing of the incompressibility constraint on the boundary.

1. INTRODUCTION

Using the information from the momentum equation for the approximate pressure on the boundary and avoiding the discrete inf-sup condition, we obtain a parameterized family of approximate problems of the weak formulation of the homogeneous Dirichlet problem for the stationary Stokes equations. In the case of the finite element method, under some hypotheses,

²⁰²⁰ Mathematics Subject Classification. 35Q35, 47A58, 35A35, 65N30, 76M10.

Key words and phrases. pressure Poisson equation, stationary Stokes problem, extended system, density, isomorphism, connection between the exact and the approximate problems, approximation of the pressure on the boundary, approximate equation, finite element method, boundary condition for the pressure, incompressibility constraint on the boundary.

the corresponding matrix form is

(1.1)
$$\begin{bmatrix} A_h & B_h^T \\ G_h(\alpha) & D_h \end{bmatrix} \begin{bmatrix} \bar{\mathbf{u}}_h \\ \bar{p}_h \end{bmatrix} = \begin{bmatrix} \bar{\mathbf{f}} \\ \bar{\psi} \end{bmatrix}, \quad \alpha \neq 1,$$

with two block Poisson preconditioners A_h and D_h and a parameter α , instead of the well known matrix problem

(1.2)
$$\begin{bmatrix} A_h & B_h^T \\ B_h & 0 \end{bmatrix} \begin{bmatrix} \bar{\mathbf{u}}_h \\ \bar{p}_h \end{bmatrix} = \begin{bmatrix} \bar{\mathbf{f}} \\ \bar{g} \end{bmatrix}$$

The Stokes problem is well posed and one of the hypotheses that is satisfied is the inf-sup (or the LBB) condition. The approximate problem also needs this condition in the discrete formulation [2, 3, 6, 10, 12, 14, 20, 28, 31, 34, 39, 46]. The associated matrix problem can be solved by direct methods [30, 44]. Usually, the problem is reformulated using penalty methods or augmented Lagrangian variational methods or with methods that lead to well posed approximate problems without the need of the inf-sup condition such as stabilization methods and least-squares finite element methods ([4] - [10], [13, 14, 18, 24], [28] - [30], [34, 35, 39, 45, 46, 54, 55] and the references cited therein).

Another method is to replace the Stokes problem (where the equations are the momentum equation and the incompressibility constraint) by a problem for the momentum equation and the Poisson equation for the pressure [6, 28, 34, 35, 36, 38, 39, 46, 47, 56]. This method is accompanied by discussions on the boundary conditions for the pressure and on the imposing of the incompressibility constraint on the boundary.

In this paper, we first formulate a parameterized family of exact extended systems (related to the problem we study, on infinite dimensional spaces). Under adequate data, the solution (\mathbf{u}, p) of the weak formulation of the homogeneous Dirichlet problem for the stationary Stokes equations is between the components of the solution of each element of this family. To solve this problem for the stationary Stokes equations becomes equivalent to solve a such exact extended system. Each extended system is approximated by a well posed formulation without the use of the discrete inf-sup (or the LBB) condition. There results a solution set of approximate solutions of Stokes problem. In the case of the finite element method, under some hypotheses, this solution set is given by a family of well - posed problems (1.1). The results are obtained for $(\mathbf{u}, p) \in \mathbf{H}_0^1(\Omega) \times (H^1(\Omega) \cap L_0^2(\Omega))$.

A component of the exact extended system is the momentum equation. Some other components are constructed on the basis of the Glowinski -Pironneau decomposition for the pressure ([28, 34, 35, 36, 56]). The extended system also includes a reformulation, for the stationary case, of the consistent pressure Poisson equation of Sani, Shen, Pironneau, Gresho, [47], from the unsteady case. We use this one for the approximation of the pressure on the boundary. This equation is based on the second writing of the momentum equation. We construct the (discrete) Laplace operator acting on pressure by connecting this last equation and the rest of the equations. The third writing of the momentum equation leads to the connection between the velocity and the Laplace operator acting on velocity. All these considerations allow us to prove that to solve the weak formulation of the homogeneous Dirichlet problem for the stationary Stokes equations is equivalent to solve a problem for the momentum equation, the parameterized perturbed pressure Poisson equation and the equation that defines the Laplace operator acting on velocity. The conditions for the approximation of the Laplace operator acting on velocity on the boundary become essential since they determine the approximation of the pressure on the boundary. For instance, from the paper [5] of Bochev and Gunzburger, we retain a modality to evaluate the discrete Laplace operator acting on velocity in the case of finite element method.

A density result for isomorphisms, from the functional analysis book by Siretchi [51], allows us to provide proofs on dense subspaces. In order to prove the well - posedness of the exact extended system (where the parameter is fixed), we restrict the problem to some dense subspaces. In this way, we can combine, in the proof, the utilization of Glowinski - Pironneau of a potential related to velocity ([28, 34, 35, 36, 56]) and a method used by Ciarlet ([20]) for the variational formulation of the biharmonic problem. We do not impose the homogeneous Neumann condition for the potential, which is usually imposed in literature. This condition arises naturally related to the dense subspaces in a stage of the proof. Finally, the well posedness is obtained on the initial (extended) spaces using the result from [51]. No condition for the pressure on the boundary is necessary. No incompressibility constraint on the boundary is necessary. This proof and the modality we consider the approximate pressure on the boundary relate the paper to the above mentioned discussions on these conditions on the boundary.

In the exact extended system, some of the equations are kept in the exact form, others are approximated. In order to study the approximate extended system that results, we formulate a general theorem on the well-posedness of a such approximate problem. This theorem is a variant of a theorem of Kantorovich and Akilov [41], modified by some formulations of Girault and Raviart [34] that we use, and by our formulations. Resulting from the conditions for the Galerkin method and the conclusion of Céa's lemma, [2, 6, 10, 20, 26, 31, 40, 46, 54, 58, 59], some practical conditions are given. The approximate part of the problem is related to the corresponding exact one as in the case of nonlinear problems from [15, 23, 34], some elliptic operators allowing the transformations. The study of the approximate problem is reduced to the existing results for some approximate problems for second order elliptic partial differential equations.

Finally, we treat an application with equal order finite element spaces.

To the best of our knowledge, the approach in the paper and the results that we prove are new.

The paper is organized as follows. In Section 2, we review a summary of notations, some results from functional analysis and some reformulations of Stokes problem. In Sections 3 and 4, we develop some formulations on the well-posedness of general exact and approximate problems of the type of those used in the paper. Sections 5, 6 and 7 are devoted to the construction of the exact extended system. The approximate extended system is studied in Section 8. An analysis using finite element method is in Section 9.

2. Preliminaries and the Stokes problem

2.1. Notations. Let Ω be a bounded and connected open subset of \mathbb{R}^N (N = 2, 3) with a Lipschitz - continuous boundary $\partial \Omega$.

Let $L^2(\Omega)$ be space of all square integrable functions defined on Ω with the usual inner product (\cdot, \cdot) . Let $H^m(\Omega)$ be the Sobolev space of all functions having the derivatives up to order m as elements of $L^2(\Omega)$, with the standard norm $\|\cdot\|_m$ and seminorm $|\cdot|_m$. Let $\mathcal{D}(\Omega)$ be the linear space of infinitely differentiable functions with compact support in Ω , topologized as it is described in [51, 57]. $H_0^1(\Omega)$ denotes the closure of $\mathcal{D}(\Omega)$ with respect to the norm $\|\cdot\|_1$. $L_0^2(\Omega) = \{p \in L^2(\Omega) \mid \int_{\Omega} p \ dx = 0\}.$

 $H^{-1}(\Omega)$ denotes the dual space of $H^1_0(\Omega)$ and V' denotes the dual space of the linear space V. $\langle \cdot, \cdot \rangle$ is the duality pairing. $\mathcal{D}'(\Omega)$ is the space of distributions on Ω . $\|\cdot\|_{-1}$ is the norm of $H^{-1}(\Omega)$. On the product space $Y = \prod_{j=1}^M Y_j$, we use the norm $\|\cdot\|_{(1)}, \|\kappa\|_{(1)}$

On the product space $Y = \prod_{j=1}^{M} Y_j$, we use the norm $\|\cdot\|_{(1)}, \|\kappa\|_{(1)}$ = $\|\kappa\|_Y = \sum_{j=1}^{M} \|\kappa_j\|_{Y_j}$, where Y_j is a Banach space with norm $\|\cdot\|_{Y_j}$, for $j = 1, \ldots, M$, and $\kappa = (\kappa_1, \ldots, \kappa_M) \in Y$.

 $(v,w)_V = \sum_{i=1}^M (v_i, w_i)_{V_i}$ is the inner product on the product space V of the Hilbert spaces V_i with the inner product $(\cdot, \cdot)_{V_i}$, $v = (v_1, \ldots, v_M)$, $v_i \in V_i$. The corresponding norm is $||v||_V = (\sum_{i=1}^M ||v_i||_{V_i}^2)^{1/2}$. (grad v, grad w) $= \sum_{i,j=1}^N (\frac{\partial v_i}{\partial x_j}, \frac{\partial w_i}{\partial x_j})$ for $v, w \in H^1(\Omega)^N$, $(grad p, grad q) = \sum_{j=1}^N (\frac{\partial p}{\partial x_j}, \frac{\partial q}{\partial x_j})$ for $p, q \in H^1(\Omega)$. If $N \ge 2$, the norm and the seminorm on $H^m(\Omega)^N$ are simply denoted as in the case N = 1.

For vector-valued functions and their spaces, we use the boldface font notation. For instance, $\mathbf{H}_0^1(\Omega) = H_0^1(\Omega)^N$ and we write $\mathbf{u} \in \mathbf{H}_0^1(\Omega)$.

For the inner product on $L^2(\Omega)$ and $\mathbf{L}^2(\Omega)$, we use the same notation (\cdot, \cdot) .

2.2. Some results of functional analysis. We specify that, given two normed spaces E and F, an isomorphism of E onto F is a linear, continuous and bijective mapping $A : E \to F$ whose inverse A^{-1} is continuous [6, 7, 23, 32, 34, 51, 52]. We then say that problem Au = f is well-posed. L(E, F)denotes the space of all the continuous linear mappings (operators) from Eto F.

Lemma 2.1. ([2], Theorem 2.4.1; [42], Theorem 2.7-11; [51], Theorem 5.3.2, Corollary 1; [58], Proposition 18.29 (a)) Let E be a normed space, let $E_0 \subseteq E$ be a linear subspace dense in E, and let F be a Banach space. Assume $A_0 \in L(E_0, F)$. Then, A_0 has an unique extension $A \in L(E, F)$ with $Au = A_0u$, $\forall u \in E_0$, and $||A||_{L(E,F)} = ||A_0||_{L(E_0,F)}$.

Lemma 2.2. ([51], Theorem 5.3.2, Corollary 5) Let E, F be two Banach spaces. There exists an isomorphism of E onto F if and only if there exist two linear subspaces E_0 and F_0 of E and F respectively, E_0 dense in E, F_0 dense in F, such that there exists an isomorphism of E_0 onto F_0 .

Corollary 2.3. In Lemma 2.2, the isomorphism of E onto F is a unique extension from E_0 to E of the isomorphism of E_0 onto F_0 .

Proof. The proof (from [51]) of Lemma 2.2 uses the result from Lemma 2.1. \Box

Corollary 2.4. Consider the spaces E, F, E_0 and F_0 from Lemma 2.2. Let $\Phi_1 \in L(E, F)$ be given and let Φ_0 be the restriction of Φ_1 to E_0 . If Φ_0 is an isomorphism of E_0 onto F_0 , then Φ_1 is an isomorphism of E onto F.

Proof. From Lemma 2.2, it follows that there exists an isomorphism Φ of E onto F. By Corollary 2.3, Φ is a unique extension of Φ_0 to E. So $\Phi = \Phi_1$.

Corollary 2.5. If Φ is the isomorphism of E onto F and Φ_0 is the isomorphism of E_0 onto F_0 from Lemma 2.2, then $\Phi_0 = \Phi|_{E_0}$ and $\Phi_0^{-1} = \Phi^{-1}|_{F_0}$.

Proof. The proof follows from the proof (from [51]) of Lemma 2.2.

Lemma 2.6. Let E, F be two normed spaces. Assume that there exists an isomorphism A of E onto F. Then, for a linear subspace E_0 dense in E, we have that $A_0 = A|_{E_0}$ is an isomorphism of E_0 onto $F_0 = A(E_0)$ and F_0 is dense in F.

Proof. We mention only that since A is continuous and surjective, it follows that $A(E_0)$ is dense in F. (We use a result from [50]).

Lemma 2.7. Assume that the hypotheses of Lemma 2.6 are satisfied. Let i be the injection $E_0 \subset E$ and let j be the injection $F_0 \subset F$. If i is compact, then j is compact.

Proof. We have $A \circ i = j \circ A_0$, so $j = A \circ i \circ A_0^{-1}$. Since *i* is compact, it results that *j* is compact.

2.3. The Stokes problem. In the sequel, **u** is the velocity, p is the kinematic pressure, ν is the kinematic viscosity and **f** is a density of body forces per unit mass.

Consider the following homogeneous Dirichlet problem for the stationary Stokes equations:

(2.1)
$$-\nu \triangle \mathbf{u} + grad \, p = \mathbf{f} \text{ in } \Omega,$$

$$(2.2) div \mathbf{u} = 0 \text{ in } \Omega$$

(2.3)
$$\mathbf{u} = 0 \text{ on } \partial \Omega$$
,

under the condition

(2.4)
$$\int_{\Omega} p \, dx = 0 \, .$$

As usual [5, 34, 36], we take $\nu = 1$. For $\nu \neq 1$, the pressure is νp and the free term is $\nu \mathbf{f}$.

In the literature, since $div(\Delta \mathbf{u}) = \Delta(div \mathbf{u}) = 0$, the pressure Poisson equation is obtained,

(2.5)
$$-\triangle p = -div \mathbf{f} \text{ in } \Omega.$$

Problem (2.1), (2.5), (2.3) is approximated in [36], where it is introduced the following problem derived from (2.1) - (2.3): find $(\mathbf{u}, \zeta, p) \in \mathbf{H}_0^1(\Omega) \times$ $H_0^1(\Omega) \times (H^1(\Omega)/\mathbb{R})$ such that

- (2.6) $(\operatorname{grad} p, \operatorname{grad} \tilde{\mu}) = (\mathbf{f}, \operatorname{grad} \tilde{\mu}), \ \forall \, \tilde{\mu} \in H^1_0(\Omega),$
- (2.7) $(grad \mathbf{u}, grad \mathbf{w}) (p, div \mathbf{w}) = (\mathbf{f}, \mathbf{w}), \ \forall \mathbf{w} \in \mathbf{H}_0^1(\Omega),$
- (2.8) $(\operatorname{grad} \zeta, \operatorname{grad} \bar{\mu}) (\operatorname{div} \mathbf{u}, \bar{\mu}) = 0, \ \forall \, \bar{\mu} \in H^1(\Omega),$

under the condition that the boundary $\partial\Omega$ is smooth (such that from $\phi \in H_0^1(\Omega)$ it can be deduced that $\phi \in H_0^1(\Omega) \cap H^2(\Omega)$) and $\mathbf{f} \in \mathbf{L}^2(\Omega)$. $(\mathbf{u}, \zeta, p) \in \mathbf{H}_0^1(\Omega) \times H_0^1(\Omega) \times (H^1(\Omega)/\mathbb{R})$ is also the solution of the problem [36]

(2.9)
$$-\triangle p = -div \mathbf{f} \text{ in } \Omega.$$

(2.10)
$$-\Delta \mathbf{u} + \operatorname{grad} p = \mathbf{f} \text{ in } \Omega,$$

- (2.11) $\mathbf{u} = 0 \text{ on } \partial\Omega,$
- (2.12) $-\triangle \zeta = div \, \mathbf{u} \text{ in } \Omega,$
- (2.13) $\zeta = 0 \text{ on } \partial\Omega,$
- (2.14) $(\mathbf{n} \cdot \nabla \zeta)|_{\partial \Omega} = 0 \text{ on } \partial \Omega.$

Problem (2.9) - (2.14) is decomposed, equivalently, in some variational problems in [34, 35]. We retain only the equations for pressure: $p \in L^2(\Omega)$, $q \in H^1_0(\Omega)$, $\hat{p} \in H^1(\Omega)$,

(2.15)
$$p = q + \hat{p},$$

(2.16)
$$(\operatorname{grad} q, \operatorname{grad} \overline{\lambda}) = (\mathbf{f}, \operatorname{grad} \overline{\lambda}), \ \forall \, \overline{\lambda} \in H^1_0(\Omega),$$

- (2.17) $(\operatorname{grad} \hat{p}, \operatorname{grad} \tilde{\lambda}) = 0, \ \forall \, \tilde{\lambda} \in H_0^1(\Omega),$
- (2.18) $\hat{p} = \lambda_w \text{ on } \partial\Omega,$

where $\lambda_w \in H^{-1/2}(\partial\Omega)$ is determined by the methods from [34, 35].

In [47], for the unsteady case, for the time dependent incompressible Stokes problem with Cauchy and Dirichlet data in Ω over the time interval (0,T), under the assumption $-\nu \Delta \mathbf{u} + \operatorname{grad} p \in \mathbf{L}^2(\Omega)$, the following equation is obtained

(2.19)
$$(-\nu \triangle \mathbf{u} + \operatorname{grad} p, \operatorname{grad} \tilde{\mu}) = (\mathbf{f}, \operatorname{grad} \tilde{\mu}), \ \forall \, \tilde{\mu} \in H^1(\Omega) \,,$$

It is the weak form of the equation ([47])

(2.20)
$$-\nu \operatorname{div}(\Delta \mathbf{u}) + \Delta p = \operatorname{div} \mathbf{f} \operatorname{in} \Omega,$$

which is one of the equations of the consistent pressure Poisson equation formulation of the mentioned unsteady Stokes problem [38, 47].

According to [28, 38, 47], the problem (2.1), (2.2), (2.3), (2.21) is equivalent to the problem (2.1), (2.5), (2.3), (2.21), where

$$(2.21) div \mathbf{u} = 0 \text{ on } \partial\Omega.$$

Condition (2.14) is equivalent to (2.21) [55, 56].

Let us also retain the Neumann boundary condition for the pressure

(2.22)
$$\frac{\partial p}{\partial n}|_{\partial\Omega} = ((\nu \bigtriangleup \mathbf{u} + \mathbf{f}) \cdot \mathbf{n})|_{\partial\Omega},$$

where **n** is the unit outward normal to $\partial \Omega$ [13, 38, 39, 45, 46, 47].

Remark 2.8. We do not use (2.21) or (2.22) or a Dirichlet boundary condition for the pressure in the paper. We do not use a boundary unknown λ_w (as in (2.18)) or a variational problem related to $\partial\Omega$. We do not impose condition (2.14). This one arises naturally, in a stage of the proof, related to some dense subspaces. (See (6.32) and Remark 6.10).

3. A LEMMA THAT REDUCES THE PROOFS OF WELL-POSEDNESS TO DENSE SUBSPACES

Consider the equation

(3.1)
$$\widehat{\Phi}(\widehat{U}) = \widehat{\mathcal{F}}.$$

Using Lemma 2.2, we want to reduce the study of (3.1) to the study of the equation

(3.2)
$$\widehat{\Phi}_0(\widehat{U}) = \widehat{\mathcal{F}}_0,$$

where $\widehat{\Phi}_0$ is a restriction of $\widehat{\Phi}$.

Lemma 3.1. (proof of the open mapping theorem [11, 19, 25, 42, 51, 57]) Let E and F be two Banach spaces and let $f: E \to F$ be a linear, continuous and surjective mapping. Then, there exist $\varepsilon > 0$ and $\delta > 0$ such that $f(B_E(0,\varepsilon)) \supset B_F(0,\delta)$.

Lemma 3.2. (Lemma 5.4, page 149, [19]) A continuous linear mapping f from a normed space E into a normed space F is open if and only if there exists an open ball B of E such that f(B) is a subset of F with nonempty interior.

Lemma 3.3. Let $\widehat{\Gamma}$, $\widehat{\Sigma}$ be two Banach spaces. Let τ , τ' be the (norm) topologies on $\widehat{\Gamma}$ and $\widehat{\Sigma}$ respectively. Consider a continuous linear mapping $\widehat{\Phi}:\widehat{\Gamma}\to\widehat{\Sigma}$.

Let $\widehat{\Gamma}_0 \subset \widehat{\Gamma}$ be a linear subspace dense in $\widehat{\Gamma}$ and let $\widehat{\Sigma}_0 \subset \widehat{\Sigma}$ be a linear subspace dense in $\widehat{\Sigma}$ such that $\widehat{\Phi}(\widehat{\Gamma}_0) \subseteq \widehat{\Sigma}_0$. We take the induced topology $\tau \cap \widehat{\Gamma}_0$ on $\widehat{\Gamma}_0$ and the induced topology $\tau' \cap \widehat{\Sigma}_0$ on $\widehat{\Sigma}_0$.

Let $\widehat{\Phi}_0 : \widehat{\Gamma}_0 \to \widehat{\Sigma}_0, \ \widehat{\Phi}_0(\widehat{U}_0) = \widehat{\Phi}(\widehat{U}_0), \ \forall \ \widehat{U}_0 \in \widehat{\Gamma}_0.$

Then, $\widehat{\Phi}_0$ is a continuous linear mapping.

Moreover, assume that $\widehat{\Phi}$ is surjective and $\widehat{\Phi}_0$ is an bijection of $\widehat{\Gamma}_0$ onto $\widehat{\Sigma}_0$. Then, $\widehat{\Phi}_0$ is an isomorphism of $\widehat{\Gamma}_0$ onto $\widehat{\Sigma}_0$ and $\widehat{\Phi}$ is an isomorphism of $\widehat{\Gamma}$ onto $\widehat{\Sigma}$.

Proof. Let us first prove that $\widehat{\Phi}_0$ is a continuous mapping.

Let ι_0 be the injection $\widehat{\Gamma}_0 \subset \widehat{\Gamma}$ and let \jmath_0 be the injection $\widehat{\Sigma}_0 \subset \widehat{\Sigma}$.

We have the induced topology $\tau \cap \widehat{\Gamma}_0$ on $\widehat{\Gamma}_0$ and the induced topology $\tau' \cap \widehat{\Sigma}_0$ on $\widehat{\Sigma}_0$, so injections i_0 and j_0 are continuous.

Define $\widehat{\Phi}_1: \widehat{\Gamma}_0 \to \widehat{\Sigma}, \ \widehat{\Phi}_1(\widehat{U}_0) = \widehat{\Phi}_0(\widehat{U}_0) = \widehat{\Phi}(\widehat{U}_0), \ \forall \ \widehat{U}_0 \in \widehat{\Gamma}_0.$

We obtain: $\widehat{\Phi}_1 = \widehat{\Phi} \circ \imath_{\widehat{\Gamma}_0}$, so $\widehat{\Phi}_1$ is continuous. $\widehat{\Phi}_1 = \jmath_0 \circ \widehat{\Phi}_0$, so $\widehat{\Phi}_0$ is continuous.

In order to prove that $\widehat{\Phi}_0$ is open, we use Lemma 3.2.

 Φ is defined on Banach spaces and it is linear, continuous and surjective, so, by the open mapping theorem ([11, 19, 25, 42, 51, 57]), $\hat{\Phi}$ is open. From Lemma 3.1, we have that there exist $\varepsilon > 0$ and $\delta > 0$ such that

 $\widehat{\Phi}(B_{\widehat{\Gamma}}(0,\varepsilon)) \supset B_{\widehat{\Sigma}}(0,\delta).$

 $B_{\widehat{\Sigma}}(0,\delta) \cap \widehat{\Sigma}_{0} = \widehat{\Phi}_{0}(\widehat{\Phi}_{0}^{-1}(B_{\widehat{\Sigma}}(0,\delta) \cap \widehat{\Sigma}_{0})) = \widehat{\Phi}(\widehat{\Phi}_{0}^{-1}(B_{\widehat{\Sigma}}(0,\delta) \cap \widehat{\Sigma}_{0}) \cap \widehat{\Gamma}_{0}) \subset \widehat{\Phi}(\widehat{\Phi}^{-1}(B_{\widehat{\Sigma}}(0,\delta) \cap \widehat{\Sigma}_{0}) \cap \widehat{\Gamma}_{0}) = \widehat{\Phi}(\widehat{\Phi}^{-1}(B_{\widehat{\Sigma}}(0,\delta)) \cap \widehat{\Phi}^{-1}(\widehat{\Sigma}_{0}) \cap \widehat{\Gamma}_{0}) = \widehat{\Phi}(\widehat{\Phi}^{-1}(B_{\widehat{\Sigma}}(0,\delta)) \cap \widehat{\Gamma}_{0}) = \widehat{\Phi}(\widehat{\Phi}^{-1}(B_{\widehat{\Sigma}(0,\delta)) \cap \widehat{\Gamma}_{0}) = \widehat{\Phi$

$$\begin{split} &\widehat{\Phi}_0(\widehat{\Phi}^{-1}(B_{\widehat{\Sigma}}(0,\delta)) \cap \widehat{\Gamma}_0) = \widehat{\Phi}(\widehat{\Phi}^{-1}(B_{\widehat{\Sigma}}(0,\delta)) \cap \widehat{\Gamma}_0) \supset B_{\widehat{\Sigma}}(0,\delta) \cap \widehat{\Sigma}_0 \neq \emptyset. \\ &\widehat{\Phi} \text{ is continuous, so } \widehat{\Phi}^{-1}(B_{\widehat{\Sigma}}(0,\delta)) \text{ is an open subset of } \widehat{\Gamma}. \end{split}$$

 $\widehat{B}_0 = \widehat{\Phi}^{-1}(B_{\widehat{\Sigma}}(0,\delta)) \cap \widehat{\Gamma}_0 \text{ is an open subset of } \widehat{\Gamma}_0. \ \widehat{\Phi}_0(\widehat{B}_0) \supset B_{\widehat{\Sigma}}(0,\delta) \cap \widehat{\Sigma}_0 \neq \emptyset. \ B_{\widehat{\Sigma}}(0,\delta) \cap \widehat{\Sigma}_0 \text{ is a nonempty open subset of } \widehat{\Sigma}_0.$

From Lemma 3.2, we obtain that $\widehat{\Phi}_0$ is open.

 $\widehat{\Phi}_0$ is linear, continuous and bijective. $\widehat{\Phi}_0$ is open, so $\widehat{\Phi}_0^{-1}$ is continuous. So $\widehat{\Phi}_0$ is an isomorphism of $\widehat{\Gamma}_0$ onto $\widehat{\Sigma}_0$.

Applying Lemma 2.2, in the form of Corollary 2.4, it results that $\widehat{\Phi}$ is an isomorphism of $\widehat{\Gamma}$ onto $\widehat{\Sigma}$.

4. A THEOREM ON THE WELL-POSEDNESS OF A CLASS OF APPROXIMATE LINEAR PROBLEMS

An equation equivalent to a form of (3.1) is introduced. A theorem on the well-posedness of its approximation is formulated. Some practical conditions are given in two corollaries.

The general framework from this section is applied in the following sections. Let Ω , \mathcal{X} , \mathcal{Y} , \mathcal{Z} be four Banach spaces. Define

 $\widehat{\Gamma} = \mathfrak{Q} \times \mathfrak{X}, \, \widehat{\Sigma} = \mathfrak{Z} \times \mathfrak{Y}, \, \widehat{\Delta} = \mathfrak{Z} \times \mathfrak{X}.$

Assume that \mathfrak{T} is an isomorphism of \mathfrak{Y} onto \mathfrak{X} . $\widehat{\mathfrak{T}} = [\mathfrak{J}_{\mathfrak{Z}}, \mathfrak{T}]^T$ is an isomorphism of $\widehat{\Sigma}$ onto $\widehat{\Delta}$ and $\widehat{\Delta} = \mathfrak{Z} \times \mathfrak{T}(\mathfrak{Y}) = \widehat{\mathfrak{T}}(\widehat{\Sigma})$.

 $\mathcal{J}_{\mathcal{I}}$ is the identity operator on \mathcal{I} , $\mathcal{I}_{\mathcal{X}}$ is the identity operator on \mathcal{X} . $\mathcal{J}_{\mathcal{I}}$ and $\mathcal{I}_{\mathcal{X}}$ are automorphisms. (The domain and the range are endowed with the same norm in each case).

Assume that $\widehat{\Phi}$ is an isomorphism of $\widehat{\Gamma}$ onto $\widehat{\Sigma}$ and $\widehat{\Phi}$ has a formulation which is deducible from the definition of the operator $\widehat{\mathcal{A}}$ below. In the present context, we denote $\widehat{U} = (x, U) \in \widehat{\Gamma}$ and we consider equation (3.1) in the following formulation

(4.1)
$$\widehat{\Phi}(x,U) = \widehat{\mathcal{F}},$$

Let us introduce the operators $\mathcal{G} \in L(\mathbb{Q} \times \mathfrak{X}, \mathfrak{Y})$ and $\mathcal{B} \in L(\mathbb{Q} \times \mathfrak{X}, \mathfrak{Z})$ and let $\widehat{\mathcal{A}} \in L(\widehat{\Gamma}, \widehat{\Delta})$ be defined by $\widehat{\mathcal{A}} = \widehat{\mathcal{T}}\widehat{\Phi} = [\mathcal{J}_{\mathcal{Z}}, \mathfrak{T}]^T [\mathcal{B}, \Phi]^T = [\mathcal{J}_{\mathcal{Z}} \mathfrak{B}, \mathfrak{T}\Phi]^T = [\mathcal{B}, \mathfrak{I}_{\mathcal{X}} - \mathfrak{T}\mathcal{G}]^T = [\mathcal{B}, \mathcal{A}]^T$ or $\widehat{\mathcal{A}}(x, U) = [\mathcal{B}(x, U), U - \mathfrak{T}\mathcal{G}(x, U)]^T = [\mathcal{B}(x, U), \mathcal{A}(x, U)]^T$. It results that $\widehat{\mathcal{A}}$ is an isomorphism of $\widehat{\Gamma}$ onto $\widehat{\Delta}$. $\widehat{\Phi}(\widehat{\Gamma}) = \widehat{\Sigma}, \ \widehat{\mathcal{A}}(\widehat{\Gamma}) = \widehat{\mathfrak{T}}\widehat{\Phi}(\widehat{\Gamma}) = \widehat{\mathfrak{T}}(\widehat{\Sigma}) = \widehat{\Delta} = \mathfrak{Z} \times \mathfrak{T}(\mathfrak{Y}).$ $\widehat{\Gamma} = \widehat{\mathcal{A}}^{-1}\widehat{\mathfrak{T}}(\widehat{\Sigma}) = \widehat{\mathcal{A}}^{-1}\widehat{\Delta} = \widehat{\mathcal{A}}^{-1}(\mathcal{Z} \times \mathfrak{T}(\mathfrak{Y})) = \widehat{\mathcal{A}}^{-1}(\mathcal{Z}, \mathfrak{T}(\mathfrak{Y})).$ Let \mathfrak{X}_h be a closed subspace of \mathfrak{X} . Define

$$\begin{split} \widehat{\Gamma}_h &= \mathfrak{Q} \times \mathfrak{X}_h \subset \mathfrak{Q} \times \mathfrak{X} = \widehat{\Gamma}, \, \widehat{\Delta}_h = \mathfrak{Z} \times \mathfrak{X}_h \subset \mathfrak{Z} \times \mathfrak{X} = \widehat{\Delta}.\\ \text{Let } \mathfrak{T}_h &\in L(\mathfrak{Y}, \mathfrak{X}_h), \, \mathfrak{X}_h = \mathfrak{T}_h(\mathfrak{Y}), \, \widehat{\mathfrak{T}}_h \in L(\widehat{\Sigma}, \mathfrak{Z} \times \mathfrak{X}_h), \, \widehat{\mathfrak{T}}_h = [\mathfrak{J}_{\mathfrak{Z}}, \mathfrak{T}_h]^T. \end{split}$$

Let $\widehat{\mathcal{K}} \in L(\widehat{\Gamma}, \widehat{\Delta}), \widehat{\mathcal{K}} = [\mathcal{B}, \mathfrak{I}_{\mathcal{X}} - \mathfrak{T}_h \mathfrak{G}]^T = [\mathcal{B}, \mathcal{K}]^T$ or $\widehat{\mathcal{K}}(x, U) = [\mathcal{B}(x, U), U - \mathfrak{T}_h \mathfrak{G}(x, U)]^T = [\mathcal{B}(x, U), \mathcal{K}(x, U)]^T$.

Observe that we have $Range(\mathcal{K}) \subseteq Range(\mathcal{A}) = \mathfrak{X} = \mathfrak{T}(\mathfrak{Y}).$ We denote $\widehat{\mathfrak{F}} = [\mathfrak{F}_{\mathfrak{B}}, \mathfrak{F}]^T \in \widehat{\Sigma}.$

Define $\widehat{\mathcal{A}}_h : \widehat{\Gamma}_h \to \widehat{\Delta}_h, \widehat{\mathcal{A}}_h \in L(\widehat{\Gamma}_h, \widehat{\Delta}_h), \widehat{\mathcal{A}}_h = [\mathfrak{B}, \mathcal{A}_h]^T \text{ or } \widehat{\mathcal{A}}_h(x_h, U_h) = [\mathfrak{B}(x_h, U_h), U_h - \mathfrak{T}_h \mathfrak{G}(x_h, U_h)]^T = [\mathfrak{B}(x_h, U_h), \mathcal{A}_h(x_h, U_h)]^T, \mathcal{A}_h \in L(\widehat{\Gamma}_h, \mathfrak{X}_h), \mathcal{A}_h(x_h, U_h) = U_h - \mathfrak{T}_h \mathfrak{G}(x_h, U_h),$

Equation (4.1) is equivalent to the following equation

(4.2)
$$\begin{array}{c} \mathcal{B}(x,U) = \mathcal{J}_{\mathcal{Z}} \mathcal{F}_{\mathcal{B}}, \\ U - \mathfrak{I} \mathfrak{I}(x,U) = \mathfrak{I} \mathfrak{F}, \end{array} \quad \text{or} \quad \widehat{\mathcal{A}}(x,U) = \widehat{\mathfrak{I}} \widehat{\mathfrak{F}}, \end{array}$$

Given the conforming space \mathfrak{X}_h , approximate (4.2) by

(4.3)
$$\begin{array}{c} \mathfrak{B}(x_h, U_h) = \mathfrak{J}_{\mathfrak{Z}} \mathfrak{F}_{\mathfrak{B}}, \\ U_h - \mathfrak{T}_h \mathfrak{G}(x_h, U_h) = \mathfrak{T}_h \mathfrak{F}, \end{array} \quad \text{or} \quad \widehat{\mathcal{A}}_h(x_h, U_h) = \widehat{\mathfrak{T}}_h \widehat{\mathfrak{F}}, \end{array}$$

Lemma 4.1. ([52]) Let *E*, *F* be two Banach spaces and let $T, S \in L(E, F)$. If the operator *T* is bijective and $||T^{-1}||_{L(F,E)}||T - S||_{L(E,F)} < 1$, then the operator *S* is bijective and $||S^{-1}||_{L(F,E)} \leq (1-q)^{-1} ||T^{-1}||_{L(F,E)}, \forall q \in \mathbb{R}$ that satisfies $||T^{-1}||_{L(F,E)}||T - S||_{L(E,F)} \leq q < 1$.

In order to use Theorem XIV.1.1, [41], and the proof of Theorem IV.3.8, [34], to relate equations (4.2) and (4.3), we must reformulate these theorems into the following Theorem 4.2. The proof is an adaptation, to our conditions, of the proofs of these mentioned theorems from [41] and [34].

Theorem 4.2. Consider the framework introduced above and the isomorphism $\widehat{\mathcal{A}}$ of $\widehat{\Gamma}$ onto $\widehat{\Delta}$. If

(4.4)
$$q_* = \|(\mathfrak{T} - \mathfrak{T}_h)\mathfrak{G}\|_{L(\widehat{\Gamma},\mathfrak{X})} \|\widehat{\mathcal{A}}^{-1}\|_{L(\widehat{\Delta},\widehat{\Gamma})} < 1,$$

then there exists $\widehat{\Gamma}_*$, $\widehat{\Gamma}_* \subseteq \widehat{\Gamma}_h$, $\widehat{\Gamma}_* \neq \emptyset$, $\widehat{\Gamma}_*$ closed in $\widehat{\Gamma}_h$, such that the restriction $\widehat{\mathcal{A}}_*$ of $\widehat{\mathcal{A}}_h$ to $\widehat{\Gamma}_*$, $\widehat{\mathcal{A}}_* = \widehat{\mathcal{A}}_h|_{\widehat{\Gamma}_*}$, is an isomorphism of $\widehat{\Gamma}_*$ onto $\widehat{\Delta}_h$. We have $\widehat{\Gamma}_* = \widehat{\mathcal{K}}^{-1}(\mathbb{Z} \times \mathfrak{T}_h(\mathfrak{Y})) = \widehat{\mathcal{K}}^{-1}(\mathbb{Z} \times \mathfrak{X}_h) = \widehat{\mathcal{K}}^{-1}(\widehat{\Delta}_h)$ and

(4.5)
$$\|\widehat{\mathcal{A}}_*^{-1}\|_{L(\widehat{\Delta}_h,\widehat{\Gamma}_h)} \le (1-q_*)^{-1} \|\widehat{\mathcal{A}}^{-1}\|_{L(\widehat{\Delta},\widehat{\Gamma})},$$

Moreover, the following results hold. Let $\widehat{\mathcal{F}} \in \widehat{\Sigma}$. Then, there exists an unique solution (x_h, U_h) of (4.3) that approximates the unique solution (x, U) of (4.2). We have

(4.6)
$$\left\| \begin{bmatrix} x \\ U \end{bmatrix} - \begin{bmatrix} x_h \\ U_h \end{bmatrix} \right\|_{\widehat{\Gamma}} \le (1 - q_*)^{-1} \|\widehat{\mathcal{A}}^{-1}\|_{L(\widehat{\Delta},\widehat{\Gamma})} \| (\mathcal{T} - \mathcal{T}_h) \mathcal{T}^{-1} U\|_{\mathfrak{X}},$$

Proof. We have $\|\widehat{\mathcal{A}} - \widehat{\mathcal{K}}\|_{L(\widehat{\Gamma},\widehat{\Delta})} \|\widehat{\mathcal{A}}^{-1}\|_{L(\widehat{\Delta},\widehat{\Gamma})} = q_* < 1$. Applying Lemma 4.1, since $\widehat{\mathcal{A}}$ is an isomorphism of $\widehat{\Gamma}$ onto $\widehat{\Delta}$, it results that $\widehat{\mathcal{K}}$ is an isomorphism of $\widehat{\Gamma}$ onto $\widehat{\Delta}$ and

(4.7)
$$\|\widehat{\mathcal{K}}^{-1}\|_{L(\widehat{\Delta},\widehat{\Gamma})} \le (1-q_*)^{-1} \|\widehat{\mathcal{A}}^{-1}\|_{L(\widehat{\Delta},\widehat{\Gamma})},$$

In the space $L(\widehat{\Gamma}_h, \widehat{\Delta}_h)$, let us consider the operator $\widehat{\mathcal{A}}_h$. $(\widehat{\mathcal{A}}_h \in L(\widehat{\Gamma}_h, \widehat{\Delta}_h))$. Observe that we have: $\forall (x_h, U_h) \in \widehat{\Gamma}_h, \widehat{\mathcal{A}}_h(x_h, U_h) = \widehat{\mathcal{K}}(x_h, U_h)$. Let $\widehat{\Gamma}_* = \widehat{\mathcal{K}}^{-1}(\mathcal{Z} \times \mathfrak{T}_h(\mathcal{Y})) = \widehat{\mathcal{K}}^{-1}(\mathcal{Z} \times \mathfrak{X}_h) = \widehat{\mathcal{K}}^{-1}(\widehat{\Delta}_h)$. If $\eta_{\mathcal{B}} \in \mathcal{Z}, \eta_h \in \mathcal{T}_h(\mathcal{Y})$ and if $(V', V_h) = \widehat{\mathcal{K}}^{-1}(\eta_{\mathcal{B}}, \eta_h) \in \widehat{\Gamma} = \mathcal{Q} \times \mathfrak{X}$, then $\widehat{\mathcal{K}}(V', V_h) = (\eta_{\mathcal{B}}, \eta_h)$ or $\mathcal{B}(V', V_h) = \eta_{\mathcal{B}}, V_h - \mathfrak{T}_h \mathcal{G} V_h = \eta_h$ or $\mathcal{B}(V', V_h) = \eta_{\mathcal{B}}, V_h = \mathcal{T}_h \mathcal{G} V_h + \eta_h$, so $V_h \in \mathfrak{X}_h$. It follows that $V_h \in \mathfrak{X}_h$ and, since $V' \in \mathcal{Q}$, we obtain $(V', V_h) \in \widehat{\Gamma}_h = \mathcal{Q} \times \mathfrak{X}_h$ or $\widehat{\mathcal{K}}^{-1}(\eta_{\mathcal{B}}, \eta_h) \in \widehat{\Gamma}_h = \mathcal{Q} \times \mathfrak{X}_h$. It follows that $\widehat{\Gamma}_* \subseteq \widehat{\Gamma}_h = \mathcal{Q} \times \mathfrak{X}_h$.

 $\widehat{\mathcal{K}}$ is an isomorphism, so $\widehat{\Gamma}_*$ is closed in $\widehat{\Gamma}_h = \mathcal{Q} \times \mathfrak{X}_h$.

Since $\forall (x_h, U_h) \in \widehat{\Gamma}_h$, we have $\widehat{\mathcal{A}}_h(x_h, U_h) = \widehat{\mathcal{K}}(x_h, U_h)$, it follows that the restriction $\widehat{\mathcal{A}}_*$ of $\widehat{\mathcal{A}}_h$ to $\widehat{\Gamma}_*$, $\widehat{\mathcal{A}}_* = \widehat{\mathcal{A}}_h|_{\widehat{\Gamma}_*}$, is an isomorphism of $\widehat{\Gamma}_*$ onto $\widehat{\Delta}_h$.

Hence it follows that the operator $\widehat{\mathcal{A}}_*$ has a continuous inverse that coincides with $\widehat{\mathcal{K}}^{-1}$ on $\widehat{\Delta}_h$ and

(4.8)
$$\|\widehat{\mathcal{A}}_{*}^{-1}\|_{L(\widehat{\Delta}_{h},\widehat{\Gamma}_{h})} \leq \|\widehat{\mathcal{K}}^{-1}\|_{L(\widehat{\Delta},\widehat{\Gamma})}.$$

Relation (4.5) results from (4.8) and (4.7).

Solutions (x, U) and (x_h, U_h) are given by the isomorphisms $\widehat{\mathcal{A}}$ and $\widehat{\mathcal{A}}_*$ respectively. Given $\widehat{\mathcal{F}}$, (x_h, U_h) approximates (x, U).

Viewing the identity operator and the projection operator from the proof of an identity from Theorem 12.1.2, page 479, [2], as \mathcal{T} and \mathcal{T}_h from our case, we easy obtain (4.6): We have

$$\begin{split} & \mathbb{T}^{-1}U - \mathbb{G}(x,U) = \mathfrak{F}, \\ & \mathbb{T}_{h}\mathbb{T}^{-1}U - \mathbb{T}_{h}\mathbb{G}(x,U) = \mathbb{T}_{h}\mathfrak{F}, \\ & U_{h} = \mathbb{T}_{h}\mathbb{G}(x_{h},U_{h}) + \mathbb{T}_{h}\mathfrak{F} = \mathbb{T}_{h}\mathbb{G}(x_{h},U_{h}) + \mathbb{T}_{h}\mathbb{T}^{-1}U - \mathbb{T}_{h}\mathbb{G}(x,U) \\ & \mathbb{B}(x,U) - \mathbb{B}(x_{h},U_{h}) = \mathbb{B}(x,U) - \mathfrak{F}_{\mathbb{B}} = 0 \\ & \mathbb{K}((x,U) - (x_{h},U_{h})) = U - \mathbb{T}_{h}\mathbb{G}(x,U) - (U_{h} - \mathbb{T}_{h}\mathbb{G}(x_{h},U_{h})) = U - \mathbb{T}_{h}\mathbb{T}^{-1}U \\ & = (\mathbb{T} - \mathbb{T}_{h})\mathbb{T}^{-1}U \\ & \left[\begin{array}{c} \mathbb{B}(x,U) - \mathbb{B}(x_{h},U_{h}) \\ \mathbb{K}(x,U) - \mathbb{K}(x_{h},U_{h}) \end{array} \right] = \left[\begin{array}{c} 0 \\ (\mathbb{T} - \mathbb{T}_{h})\mathbb{T}^{-1}U \end{array} \right] . \\ & \text{Hence} \\ & \left[\begin{array}{c} x \\ U \\ \end{array} \right] - \left[\begin{array}{c} x_{h} \\ U_{h} \end{array} \right] = \widehat{\mathcal{K}}^{-1} \left[\begin{array}{c} 0 \\ (\mathbb{T} - \mathbb{T}_{h})\mathbb{T}^{-1}U \end{array} \right] . \\ & \text{From this identity, using (4.7), we deduce (4.6).} \\ \end{array} \right]$$

Corollary 4.3. Let $\widehat{\Gamma}_r$ be a linear subspace dense in $\widehat{\Gamma}$. Then, (4.4) has the form

(4.9)
$$q_* = \|(\mathfrak{T} - \mathfrak{T}_h)\mathcal{G}\|_{L(\widehat{\Gamma}_r, \mathfrak{X})} \|\widehat{\mathcal{A}}^{-1}\|_{L(\widehat{\Delta}, \widehat{\Gamma})} < 1,$$

Thinking to \mathcal{T} as related to the solution operators of an elliptic problems, let us introduce some additional hypotheses that result from the approximation of elliptic problems (conditions for the Galerkin method and the conclusion of Céa's lemma [2, 20, 46]). In this way, the convergence of Galerkin method is sufficient.

Corollary 4.4. Assume that the hypotheses of Theorem 4.2 are satisfied. Let X_d be a linear subspace dense in X. Assume that the embedding $X_d \subseteq X$ is continuous and the embedding operator i_0 is compact. Assume that

(4.10)
$$\lim_{h \to 0} \| (\mathfrak{T} - \mathfrak{T}_h) \Psi \|_{\mathfrak{X}} = 0, \ \forall \Psi \in \mathfrak{Y},$$

Then, $\exists h_0$ such that $\forall h < h_0, q_* < 1$, that is, (4.4). Furthermore,

(4.11)
$$\lim_{h \to 0} \|U - U_h\|_{\mathfrak{X}} = 0, \ \lim_{h \to 0} \|x - x_h\|_{\mathfrak{Q}} = 0,$$

Proof. \mathcal{T}^{-1} is an isomorphism of \mathfrak{X} onto \mathfrak{Y} .

We apply 2.6 with $E = \mathfrak{X}$, $F = \mathfrak{Y}$, $A = \mathfrak{T}^{-1}$, $E_0 = \mathfrak{X}_d$. It results $A_0 = \mathfrak{T}_{d,isom}^{-1} = (\mathfrak{T}^{-1})|_{\mathfrak{X}_d}$ is an isomorphism of E_0 onto $F_0 = \mathfrak{Y}_d = \mathfrak{T}^{-1}(\mathfrak{X}_d)$ and \mathfrak{Y}_d is dense in \mathfrak{Y} . $\mathfrak{T}_{d,isom} \in L(\mathfrak{Y}_d, \mathfrak{X}_d), \mathfrak{T}_{d,isom}^{-1}(U) = (\mathfrak{T}^{-1})|_{\mathfrak{X}_d}(U), \forall U \in \mathfrak{X}_d$.

 $\mathcal{Y}_{d,isom} = (\mathfrak{T}^{-1})_{\mathcal{X}_{d}} = \mathcal{T}^{-1}$ \mathcal{Y}_{d} is dense in \mathcal{Y} . $\mathcal{T}_{d,isom} \in L(\mathcal{Y}_{d}, \mathcal{X}_{d}), \mathcal{T}_{d,isom}^{-1}(U) = (\mathfrak{T}^{-1})|_{\mathcal{X}_{d}}(U), \forall U \in \mathcal{X}_{d}.$ Let j_{0} be the injection $\mathcal{Y}_{d} \subset \mathcal{Y}$. We now apply Lemma 2.7 with $E = \mathcal{X},$ $F = \mathcal{Y}, A = \mathfrak{T}^{-1}, E_{0} = \mathcal{X}_{d}, F_{0} = \mathcal{Y}_{d}, A_{0} = \mathfrak{T}_{d,isom}^{-1}, i = i_{0}, j = j_{0}.$ So j_{0} is compact. j_{0} is also continuous. (This follows from the proof of Lemma 2.7).

Let \mathfrak{T}_d be the restriction of \mathfrak{T} to \mathfrak{Y}_d , $\mathfrak{T}_d \in L(\mathfrak{Y}_d, \mathfrak{X})$. We have $\mathfrak{T}_{d,isom}(\mathfrak{F}) = \mathfrak{T}_d(\mathfrak{F}), \forall \mathfrak{F} \in \mathfrak{Y}_d$. Let $\mathfrak{T}_{d,h}$ be the restriction of \mathfrak{T}_h to \mathfrak{Y}_d . $\mathfrak{T}_{d,h} \in L(\mathfrak{Y}_d, \mathfrak{X}_h)$. From Lemma 2.1, it follows that

(4.12)
$$\|\mathfrak{T} - \mathfrak{T}_h\|_{L(\mathfrak{Y},\mathfrak{X})} = \|\mathfrak{T}_d - \mathfrak{T}_{d,h}\|_{L(\mathfrak{Y}_d,\mathfrak{X})},$$

where $\mathfrak{T} - \mathfrak{T}_h$ is the unique extension of $\mathfrak{T}_d - \mathfrak{T}_{d,h}$. We have

(4.13)
$$\mathfrak{T}_d - \mathfrak{T}_{d,h} = (\mathfrak{T} - \mathfrak{T}_h)|_{\mathfrak{Y}_d} = (\mathfrak{T} - \mathfrak{T}_h) \circ \mathfrak{z}_0,$$

We can now apply the end of Remark IV.3.4, page 306, [34], and also page 33, [43]. This and (4.10) give (4.14), where

(4.14)
$$\lim_{h \to 0} \|\mathcal{T}_d - \mathcal{T}_{d,h}\|_{L(\mathcal{Y}_d, \mathfrak{X})} = 0,$$

Taking into account (4.12), we deduce where

(4.15)
$$\lim_{h \to 0} \|\mathfrak{T} - \mathfrak{T}_h\|_{L(\mathfrak{Y},\mathfrak{X})} = 0 \,,$$

and $\|(\mathfrak{T}-\mathfrak{T}_h)\mathfrak{G}\|_{L(\widehat{\Gamma},\mathfrak{X})} \leq \|\mathfrak{T}-\mathfrak{T}_h\|_{L(\mathfrak{Y},\mathfrak{X})}\|\mathfrak{G}\|_{L(\widehat{\Gamma},\mathfrak{Y})}.$

Using (4.15), we obtain that $\exists h_0$ such that $\forall h < h_0, q_* < 1$, that is, (4.4). We have $\|(\mathcal{T} - \mathcal{T}_h)\mathcal{T}^{-1}U\|_{\mathcal{X}} \leq \|\mathcal{T} - \mathcal{T}_h\|_{L(\mathcal{Y},\mathcal{X})}\|\mathcal{T}^{-1}U\|_{\mathcal{Y}}$.

We use (4.6). Finally, (4.15) leads to (4.11).

Corollary 4.5. Assume that \mathfrak{X} , \mathfrak{Q} and \mathfrak{Z} are Hilbert spaces. Assume that the hypotheses of Theorem 4.2 are satisfied. Let h be a parameter. Let $\{\mathfrak{X}_h; h > 0\}$ be a family of finite dimensional subspaces of \mathfrak{X} . Let \mathfrak{S} be a set dense in \mathfrak{X} and assume that

(4.16)
$$\lim_{h \to 0} \inf_{V_h \in \mathfrak{X}_h} \|V - V_h\|_{\mathfrak{X}} = 0, \ \forall V \in \mathfrak{S},$$

Assume that there exists $\gamma_0 > 0$ such that for all $\Psi \in \mathcal{Y}$ and $W = \mathcal{T}\Psi$, $W_h = \mathcal{T}_h \Psi$, we have

(4.17)
$$||W - W_h||_{\mathfrak{X}} \le \gamma_0 \inf_{V_h \in \mathfrak{X}_h} ||W - V_h||_{\mathfrak{X}},$$

Then, (4.10) holds.

Proof. Let us fix $\Psi, \Psi \in \mathcal{Y}$.

(4.18)
$$\|(\mathcal{T} - \mathcal{T}_h)\Psi\|_{\mathfrak{X}} = \|W - W_h\|_{\mathfrak{X}},$$

CĂTĂLIN - LIVIU BICHIR

The arguments from the proof of the convergence of Galerkin method for the elliptic problems ([2, 20, 46]) lead to (4.10). W corresponds to the solution of exact variational equation, W_h corresponds to the solution of approximate variational equation and (4.17) corresponds to the Céa's lemma.

5. Preliminary remarks for the formulation of the problem we study

5.1. Some remarks on some operators and subspaces. Related to the Stokes problem, we introduce an extended system in Section 6. This extended system has the forms (3.1) and (4.1). In order to apply Lemma 3.3 to the study, we define now some adequate spaces $\widehat{\Gamma}$, $\widehat{\Sigma}$, $\widehat{\Gamma}_0$ and $\widehat{\Sigma}_0$. The results are obtained for $(u, p) \in \mathbf{H}_0^1(\Omega) \times (H^1(\Omega) \cap L_0^2(\Omega))$ and these spaces are components of some spaces $\widehat{\Gamma}_1$ and $\widehat{\Sigma}_1$ which are also defined below.

Some remarks on the Stokes problem

Let Ω be a bounded and connected open subset of \mathbb{R}^N (N = 2, 3) with a Lipschitz - continuous boundary $\partial \Omega$.

Remark 5.1. The conditions for Ω and $\partial \Omega$ are the ones from Theorem I.5.1, page 80, [34]. This theorem establishes the existence and the uniqueness of the solution of the nonhomogeneous Dirichlet problem for the stationary Stokes equations.

Let us consider the variational formulation associated to (2.1) - (2.3) [27, 34]: given $\mathbf{f} \in \mathbf{H}^{-1}(\Omega), g \in L^2_0(\Omega)' \simeq L^2_0(\Omega)$, find $(\mathbf{u}, p) \in \mathbf{H}^1_0(\Omega) \times L^2_0(\Omega)$ such that

(5.1)
$$(\operatorname{grad} \mathbf{u}, \operatorname{grad} \mathbf{w}) - (p, \operatorname{div} \mathbf{w}) = \langle \mathbf{f}, \mathbf{w} \rangle, \ \forall \, \mathbf{w} \in \mathbf{H}_0^1(\Omega),$$

(5.2)
$$(\operatorname{div} \mathbf{u}, \mu) = (g, \mu), \ \forall \, \mu \in L^2_0(\Omega)$$

The compatibility condition (page 828, [27]) is satisfied in this case,

(5.3)
$$\int_{\Omega} g \, dx = \int_{\partial\Omega} (0 \cdot \mathbf{n})|_{\partial\Omega} \, ds \, .$$

Let Ψ_S be the operator defined by the left hand sides of problem (5.1) - (5.2), $\Psi_S \in L(\mathbf{H}_0^1(\Omega) \times L_0^2(\Omega); \mathbf{H}^{-1}(\Omega) \times L_0^2(\Omega))$. Then, problem (5.1) - (5.2) has the form

(5.4)
$$\Psi_S(\mathbf{u}, p) = (\mathbf{f}, g),$$

Theorem I.5.1, page 80, [34], together with Theorem I.4.1, page 59, [34], and Theorem XIX.9, page 828, [27], together with Remark XIX.6, page 828, [27], provide that problem (5.1) - (5.2) is well - posed. Hence Ψ_S is an isomorphism of $\mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$ onto $\mathbf{H}^{-1}(\Omega) \times L_0^2(\Omega)$.

Since we need $p \in H^1(\Omega)$, we take $(u, p) \in \mathbf{H}_0^1(\Omega) \times (H^1(\Omega) \cap L_0^2(\Omega))$ and $\Upsilon_1 \times M = \Psi_S(\mathbf{H}_0^1(\Omega) \times (H^1(\Omega) \cap L_0^2(\Omega)))$, Υ_1 is a dense subspace of $\mathbf{H}^{-1}(\Omega)$ and $M = L_0^2(\Omega)$. This follows from Lemma 2.6 and from the fact that $H^1(\Omega) \cap L_0^2(\Omega)$ is dense in $L_0^2(\Omega)$. These data correspond to the spaces $\widehat{\Gamma}_1$ si $\widehat{\Sigma}_1$ defined below.

12

To obtain (5.7) below for the steady case, we take $p \in H^1(\Omega) \cap L^2_0(\Omega)$ and we adapt the technique used in [47] to deduce (2.19) in the unsteady case.

From (5.1), we obtain

(5.5)
$$-\bigtriangleup \mathbf{u} + \operatorname{grad} p - \mathbf{f} = 0, \text{ in } (\mathcal{D}'(\Omega))^N,$$

that is, (6.21) below. In other words,

(5.6)
$$\operatorname{grad} p = \Delta \mathbf{u} + \mathbf{f}, \text{ in } (\mathcal{D}'(\Omega))^N,$$

 $grad p \in \mathbf{L}^{2}(\Omega)$, hence $\Delta \mathbf{u} + \mathbf{f} \in \mathbf{L}^{2}(\Omega)$, so we have

(5.7)
$$(-\bigtriangleup \mathbf{u} - \mathbf{f}, \operatorname{grad} \bar{\mu}) + (\operatorname{grad} p, \operatorname{grad} \bar{\mu}) = 0, \ \forall \, \bar{\mu} \in H^1(\Omega),$$

Let $\mathbf{z} = -\Delta \mathbf{u}$. Then, (5.7) is written

(5.8)
$$(\mathbf{z} - \mathbf{f}, grad\,\bar{\mu}) + (grad\,p, grad\,\bar{\mu}) = 0, \ \forall \,\bar{\mu} \in H^1(\Omega),$$

We observe that, instead (2.5), we can use

(5.9)
$$\alpha \nu \operatorname{div} (\Delta \mathbf{u}) - \Delta p = -\operatorname{div} \mathbf{f} \operatorname{in} \Omega,$$

with a fixed real $\alpha \neq 1$ ($\nu = 1$ as we established). We quickly deduce that $div \mathbf{u} = 0$. Indeed, from (2.1) and (5.9), we obtain $(1 - \alpha) \triangle (div \mathbf{u}) = 0$. Using (2.12), we deduce $\triangle \triangle \zeta = 0$, Using (2.13), (2.14), it results $\zeta = 0$, so $div \mathbf{u} = 0$, that is, (2.2). For each $\alpha \neq 1$, we can obtain an approximate solution of the given Stokes problem.

Definition of the subspace \mathcal{W}

Let us retain, from [20, 35, 36, 37], the formulations of the Dirichlet problem for the biharmonic operator and the space \mathcal{V} ,

(5.10)
$$\mathcal{V} = \{(v, \psi) \in H_0^1(\Omega) \times L^2(\Omega); \forall \mu \in H^1(\Omega), b((v, \psi), \mu) = 0\},\$$

where $b((v, \psi), \mu) = (\operatorname{grad} v, \operatorname{grad} \mu) - (\psi, \mu)$. $(\psi = \operatorname{div} \mathbf{u}, \operatorname{with} u \in H^1(\Omega)^N, u = u_w \text{ on } \partial\Omega, \operatorname{in} [35, 36])$. \mathcal{V} is defined for Ω convex or with $\partial\Omega$ sufficiently smooth and, in this case, $v \in H^2(\Omega)$. Theorem 7.1.1, page 384, [20], tells us that $\mathcal{V} = \mathcal{V}_1$, where

(5.11)
$$\mathcal{V}_1 = \{(v,\psi) \in H^2_0(\Omega) \times L^2(\Omega); -\Delta v = \psi\}.$$

Since we consider a Lipschitz - continuous boundary $\partial\Omega$, we take directly $v \in H_0^1(\Omega) \cap H^2(\Omega)$ and consider the space

(5.12)
$$\mathcal{W} = \{(v,\psi) \in (H^1_0(\Omega) \cap H^2(\Omega)) \times L^2(\Omega); (v,\psi) \in \mathcal{V}\}.$$

The proof of Theorem 7.1.1, page 384, [20], remains valid if \mathcal{V} is replaced by \mathcal{W} , so $\mathcal{W} = \mathcal{V}_1$ and we can use the formulations of the Dirichlet problem for the biharmonic operator, from [20, 35, 36, 37], using the space \mathcal{W} instead of the space \mathcal{V} .

A remark on the density of a subspace

Let $T_{PD} \in L(H^{-1}(\Omega), H_0^1(\Omega))$ be the solution operator for the (scalar) Poisson equation with homogeneous Dirichlet boundary conditions. Details are given in Section 7. Following page 370, [25], where we take $V = H_0^1(\Omega)$, $H = L^2(\Omega)$ and $A = T_{PD}^{-1}$, we have

(5.13)
$$D = D(T_{PD}^{-1}) = T_{PD}(L^2(\Omega)) = \{ v \in L^2(\Omega), T_{PD}^{-1}v \in L^2(\Omega) \},\$$

and $T_{PD}^{-1}|_{T_{PD}(L^2(\Omega))}$ is an isomorphism of $D(T_{PD}^{-1})$, considered with the norm of the graph, onto $L^2(\Omega)$ [25]. $D(T_{PD}^{-1})$ is dense in $L^2(\Omega)$ [25]. $D(T_{PD}^{-1})$ is dense in $H_0^1(\Omega)$ [25].

Let

(5.14)
$$D_0 = T_{PD}(L_0^2(\Omega)) = \{ v \in L^2(\Omega), T_{PD}^{-1}v \in L_0^2(\Omega) \},\$$

 $L_0^2(\Omega)$ is closed in $L^2(\Omega)$, so $D_0 = T_{PD}(L_0^2(\Omega))$ is closed in $D(T_{PD}^{-1}) = T_{PD}(L^2(\Omega))$ (in the topology of norm of the graph). $T_{PD}^{-1}(D_0) = L_0^2(\Omega)$.

Lemma 5.2. $D_0 \cap H^2_0(\Omega)$ is dense in D_0 in the topology of the norm of the graph.

Proof. If $v \in H^2(\Omega)$, then $-\Delta v \in L^2(\Omega)$. So $T_{PD}^{-1}(H_0^2(\Omega)) \subset L^2(\Omega)$ and $H_0^2(\Omega) \subset D$.

Let us prove that $H_0^2(\Omega)$ is dense D.

We denote $\mathbb{B}_{\varepsilon}(v) = \{ \tilde{v} \in L^2(\Omega), \text{ such that } (5.15) \text{ holds.} \}$, where

(5.15)
$$\|v - \tilde{v}\|_{L^{2}(\Omega)} + \|T_{PD}^{-1}|_{T_{PD}(L^{2}(\Omega))}(v - \tilde{v})\|_{L^{2}(\Omega)} < \varepsilon.$$

Let $v \in D$.

Assume that $\exists \varepsilon > 0$ such that $\forall \tilde{v} \in H_0^2(\Omega)$ we have

(5.16)
$$\|v - \tilde{v}\|_{L^{2}(\Omega)} + \|T_{PD}^{-1}|_{T_{PD}(L^{2}(\Omega))}(v - \tilde{v})\|_{L^{2}(\Omega)} \ge \varepsilon.$$

or $\mathbb{B}_{\varepsilon}(v) \cap H_0^2(\Omega) = \emptyset$ or $\forall \ \tilde{v} \in \mathbb{B}_{\varepsilon}(v)$, we have $\tilde{v} \notin H_0^2(\Omega)$ and (5.15).

Hence $\forall \ \tilde{v} \in \mathbb{B}_{\varepsilon}(v)$, we have $\tilde{v} \notin H_0^2(\Omega)$ and $||v - \tilde{v}||_{L^2(\Omega)} < \varepsilon$. This contradicts the density of $H_0^2(\Omega)$ in $L^2(\Omega)$.

Hence, for all $v \in D$ and for all $\varepsilon > 0$, there exists $\tilde{v} \in H_0^2(\Omega)$ such that (5.15) holds, so $H_0^2(\Omega)$ is dense D.

Let $v \in int D_0$.

There exists $\varepsilon > 0$ such that $\mathbb{B}_{\varepsilon}(v) \subset int D_0$

Assume that for a such ε we have $\mathbb{B}_{\varepsilon}(v) \cap H_0^2(\Omega) = \emptyset$.

Proceeding as before, we obtain that for $v \in int D_0$, for all $\varepsilon > 0$, there exists $\tilde{v} \in D_0 \cap H^2_0(\Omega)$ such that (5.15) holds.

Hence, $int D_0 \cap H_0^2(\Omega) \neq \emptyset$ and, as a consequence, $int L_0^2(\Omega) \cap T_{PD}^{-1}(H_0^2(\Omega)) \neq \emptyset$. *int* D_0 is in the induced topology of D on D_0 , $int L_0^2(\Omega)$ is in the induced topology of $L^2(\Omega)$ on $L_0^2(\Omega)$.

The space \mathcal{V}_1 is a closed subspace of $H^1_0(\Omega) \times L^2(\Omega)$ [20]. Instead of \mathcal{V}_1 , we work with

(5.17)
$$\widetilde{\mathcal{V}}_1 = \{(\psi, v) \in L^2(\Omega) \times H^2_0(\Omega); T_{PD}\psi = v\},\$$

The space $\widetilde{\mathcal{V}}_1$ is a closed subspace of $L^2(\Omega) \times H^2_0(\Omega)$.

It results that the restriction $T_2 \in L(D(T_2), H_0^2(\Omega))$ of T_{PD} to $D(T_2) \subset L^2(\Omega)$ is a closed operator. $D(T_2)$ is complete for the norm of the graph. If $v \in H^2(\Omega)$, then $-\Delta v \in L^2(\Omega)$. So $T_{PD}^{-1}(H_0^2(\Omega)) \subset L^2(\Omega)$.

 $T_{PD}^{-1}(H_0^2(\Omega)) = D(T_2)$ as sets (or as subspaces of $L^2(\Omega)$ (in the topology of $L^2(\Omega)$)).

Hence, $D(T_2) \cap int \ L^2_0(\Omega) \neq \emptyset$ (in the topology of $L^2(\Omega)$).

Let us prove that the restriction \hat{L} of $L_0^2(\Omega)$ to $D(T_2)$ is closed in the topology of the norm of the graph.

Consider a sequence $\{l_n\} \subset L$ such that $||l_n - l||_{D(T_2)} \to 0$ in $D(T_2)$ as $n \to \infty$. $l \in D(T_2)$ since $D(T_2)$ is closed. $||l_n - l||_{L^2(\Omega)} \leq ||l_n - l||_{D(T_2)}$, so $||l_n - l||_{L^2(\Omega)} \to 0$ in $L^2(\Omega)$ with $\{l_n\} \subset L^2_0(\Omega)$. $L^2_0(\Omega)$ is closed in $L^2(\Omega)$, hence $l \in L^2_0(\Omega)$. We obtain $l \in \widetilde{L}$ and \widetilde{L} is closed in $D(T_2)$.

Inspired by some density proofs from [48], where some projection operators are used, let us work with the operators P_2 and P, where

- $P_2 = I_2$ on $D(T_2) \ominus (D(T_2) \cap L_0^2(\Omega)) = D(T_2) \ominus L$,
- $P_2 = 0$ on $D(T_2) \cap L_0^2(\Omega) = L$,
- P = I on $L^2(\Omega) \ominus L^2_0(\Omega)$,
- P = 0 on $L_0^2(\Omega)$.

Here, I_2 is the identity operator on $D(T_2)$, I is the identity operator on $L^2(\Omega)$, P_2 is the projection operator of $D(T_2)$ onto $D(T_2) \ominus L$, P_2 $\in L(D(T_2), D(T_2)), P$ is the projection operator of $L^2(\Omega)$ onto $L^2(\Omega) \ominus$ $L_0^2(\Omega), P \in L(L^2(\Omega), L^2(\Omega)).$

We have $P_2(y) = P(y)$ and $I_2(y) = I(y)$ for $y \in D(T_2)$.

We write T_{PD} , but the operator is the restriction $T_{PD}|_{L^2(\Omega)}$ of T_{PD} to $L^2(\Omega)$. $T_{PD}|_{L^2(\Omega)}$ is the inverse of $T_{PD}^{-1}|_{T_{PD}(L^2(\Omega))}$ from the definition of D. Let $u \in D_0$. There exists $y \in L_0^2(\Omega)$ such that $u = T_{PD}y$. $u \in D$.

 $H_0^2(\Omega)$ is dense in D, so there exists a sequence $\{u_n\} \subset H_0^2(\Omega)$ such that $||u - u_n||_D \to 0 \text{ as } n \to \infty$.

There exists $y_n \in D(T_2) \subset L^2(\Omega)$ such that $u_n = T_2 y_n = T_{PD} y_n$. It results $||y - y_n||_{L^2(\Omega)} \to 0$ as $n \to \infty$. Let $v_n = T_2(I_2 - P_2)y_n = T_{PD}(I - P)y_n$.

 $y_n \in D(T_2)$, so $(I_2 - P_2)y_n \in \widetilde{L}$. Hence, $v_n \in D_0$ and $v_n \in H^2_0(\Omega)$, so v_n $\in D_0 \cap H^2_0(\Omega).$

We have $||y - y_n||_{L^2(\Omega)} \to 0$ as $n \to \infty$.

It results $v_n = T_{PD}(I - P)y_n \to T_{PD}(I - P)y = u$ in the topology of D (since $(I - P)y_n \to (I - P)y$ in the topology of $L^2(\Omega)$).

Hence, $D_0 \cap H_0^2(\Omega)$ is dense in D_0 in the topology of the norm of the graph.

Definition of the subspaces $\widetilde{V^{\perp}}$ and Υ_0 It results that $T_{PD}^{-1}(D_0 \cap H_0^2(\Omega))$ is dense in $T_{PD}^{-1}(D_0) = L_0^2(\Omega)$ in the topology of $L^2(\Omega)$.

div is an isomorphism of V^{\perp} onto $L^2_0(\Omega)$, where $V = \{ \mathbf{v} \in \mathbf{H}^1_0(\Omega) | div \, \mathbf{v} = \{ \mathbf{v} \in \mathbf{H}^1_0(\Omega) | div \, \mathbf{v} = \{ \mathbf{v} \in \mathbf{H}^1_0(\Omega) | div \, \mathbf{v} = \{ \mathbf{v} \in \mathbf{H}^1_0(\Omega) | div \, \mathbf{v} \in \mathbf{H}^1_0(\Omega) \}$ 0} and $\mathbf{H}_{0}^{1}(\Omega) = V \oplus V^{\perp}$ [34].

Let us define $\widetilde{V^{\perp}} = (div|_{V^{\perp}})^{-1}(T_{PD}^{-1}(D_0 \cap H_0^2(\Omega))). T_{PD}^{-1}(D_0 \cap H_0^2(\Omega))$ is dense in $T_{PD}^{-1}(D_0) = L_0^2(\Omega)$, in the topology of $L_0^2(\Omega)$ (or $L^2(\Omega)$), so $\widetilde{V^{\perp}}$ is dense in $V^{\perp} = (div|_{V^{\perp}})^{-1}(L_0^2(\Omega)).$

It results that $V \oplus \widetilde{V^{\perp}}$ is dense in $\mathbf{H}_0^1(\Omega) = V \oplus V^{\perp}$ (since $(V \oplus \widetilde{V^{\perp}})^{\perp} =$ $\{0\}$).

Let us define Υ_0 by $\Upsilon_0 \times T_{PD}^{-1}(D_0 \cap H_0^2(\Omega)) = \Psi_S(V \oplus \widetilde{V^{\perp}}; H^1(\Omega) \cap U^{\perp}(\Omega))$ $L^2_0(\Omega)$). If we take $(\mathbf{u},p) \in V \oplus \widetilde{V^{\perp}} \times (H^1(\Omega) \cap L^2_0(\Omega))$, we have $(\mathbf{f},g) \in V$ $\Upsilon_0 \times T_{PD}^{-1}(D_0 \cap H_0^2(\Omega)).$

It results that $\Upsilon_0 \times T_{PD}^{-1}(D_0 \cap H_0^2(\Omega))$ is dense in $\mathbf{H}^{-1}(\Omega) \times L_0^2(\Omega)$, so Υ_0 is a dense subspace of $\mathbf{H}^{-1}(\Omega)$.

Let $\mathbf{F} \in \mathbf{H}^{-1}(\Omega)$. Let $S_{\mathbf{F}} \in \mathcal{D}'(\Omega)$ be defined by

(5.18)
$$\langle S_{\mathbf{F}}, \varphi \rangle = -\langle \mathbf{F}, grad \, \varphi \rangle, \ \forall \, \varphi \in \mathcal{D}(\Omega) \,,$$

Let $\widetilde{S}_{\mathbf{F}}$ be the unique continuous extension of $S_{\mathbf{F}}$ to $H_0^1(\Omega)$ (using page 49, [1], and page 291, [2]). $\widetilde{S}_{\mathbf{F}} \in H^{-1}(\Omega)$.

5.2. Some definitions introduced in order to apply Lemma 3.3 and Theorem 4.2. In order to gather some mathematical entities that we use in the sequel, we consider some definitions from Sections 7 and 8 below. Let us now define:

 $\widehat{\Gamma} = \mathfrak{Q} \times \mathfrak{X}, \, \widehat{\Sigma} = \mathfrak{Z} \times \mathfrak{Y}, \, \widehat{\Gamma}_h = \mathfrak{Q} \times \mathfrak{X}_h,$ $\begin{aligned} &\mathcal{Q} = H^1(\Omega) \times L^2_0(\Omega)^2, \\ &\mathcal{X} = \mathbf{H}^1_0(\Omega) \times H^1_0(\Omega)^2 \times \mathbf{L}^2(\Omega) \times H^1(\Omega) \cap L^2_0(\Omega) \times H^1(\Omega) \times \mathbf{L}^2(\Omega), \end{aligned}$ $\begin{aligned} \mathcal{Z} &= H^{1}(\Omega) \times L^{2}_{0}(\Omega)^{2}, \\ \mathcal{Y} &= \mathbf{H}^{-1}(\Omega) \times H^{-1}(\Omega)^{2} \times \mathbf{L}^{2}(\Omega) \times B^{-1}_{1}(H^{1}(\Omega) \cap L^{2}_{0}(\Omega)) \times (H^{1}(\Omega))' \end{aligned}$ $\times \mathbf{L}^2(\Omega),$ $\mathfrak{X}_{h} = \mathbf{X}_{0h} \times Y_{0h}^{2} \times \mathbf{X}_{h} \times M_{h} \times Y_{h} \times \mathbf{X}_{h},$
$$\begin{split} \widehat{\Gamma}_0 &= \mathfrak{Q}_0 \times \mathfrak{X}_0, \ \widehat{\Sigma}_0 &= \mathfrak{Z}_0 \times \mathfrak{Y}_0, \\ \mathfrak{Q}_0 &= H^1(\Omega) \times H^1(\Omega) \cap L^2_0(\Omega) \times T^{-1}_{PD}(D_0 \cap H^2_0(\Omega)), \end{split}$$
 $\mathfrak{X}_{0} = V \oplus \widetilde{V^{\perp}} \times H^{1}_{0}(\Omega)^{2} \times \mathbf{L}^{2}(\Omega) \times H^{1}(\Omega) \cap L^{2}_{0}(\Omega) \times H^{1}(\Omega) \times \mathbf{L}^{2}(\Omega),$ $\begin{aligned} \mathcal{Z}_0 &= H^1(\Omega) \times H^1(\Omega) \cap L^2_0(\Omega) \times (T^{-1}_{PD}(D_0 \cap H^2_0(\Omega))), \\ \mathcal{Y}_0 &= \Upsilon_0 \times H^{-1}(\Omega)^2 \times \mathbf{L}^2(\Omega) \times B_1^{-1}(H^1(\Omega) \cap L^2_0(\Omega)) \times (H^1(\Omega))' \times \end{aligned}$ $\mathbf{L}^{2}(\Omega),$
$$\begin{split} \widehat{\Gamma}_1 &= \mathfrak{Q}_1 \times \mathfrak{X}_1, \, \widehat{\Sigma}_1 = \mathfrak{Z}_1 \times \mathfrak{Y}_1, \\ \mathfrak{Q}_1 &= H^1(\Omega) \times H^1(\Omega) \cap L^2_0(\Omega) \times L^2_0(\Omega), \end{split}$$
 $\mathfrak{X}_1 = \mathfrak{X},$ $\begin{aligned} \boldsymbol{\mathcal{Z}}_1 &= \boldsymbol{H}^1(\Omega) \times \boldsymbol{H}^1(\Omega) \cap \boldsymbol{L}_0^2(\Omega) \times \boldsymbol{L}_0^2(\Omega), \\ \boldsymbol{\mathcal{Y}}_1 &= \boldsymbol{\Upsilon}_1 \times \boldsymbol{H}^{-1}(\Omega)^2 \times \mathbf{L}^2(\Omega) \times \boldsymbol{B}_1^{-1}(\boldsymbol{H}^1(\Omega) \cap \boldsymbol{L}_0^2(\Omega)) \times (\boldsymbol{H}^1(\Omega))' \times \end{aligned}$ $\mathbf{L}^{2}(\Omega),$ $x = (\hat{p}, p_S, y),$ $U = (\mathbf{u}, q, \hat{q}, \mathbf{z}, p, r, \mathbf{t}),$ $x_h = (\hat{p}_h, p_{S,h}, y_h),$ $U_h = (\mathbf{u}_h, q_h, \hat{q}_h, \mathbf{z}_h, p_h, r_h, \mathbf{t}_h),$ $\mathfrak{T} \in L(\mathfrak{Y}, \mathfrak{X}), \, \mathfrak{T} = (T_{VPD}, T_{PD}, T_{PD}, \pi, B_1, B_1, \pi),$ $\mathfrak{G} \in L(\mathfrak{Q} \times \mathfrak{X}, \mathfrak{Y}), \, \mathfrak{G}(y, U) = (G_{\mathbf{u}}(p_S), G_q(\mathbf{t}, \mathbf{z}), G_{\hat{q}}(p), G_{\mathbf{z}}(p), G_p(p, \hat{p}, \hat{q}, \mathbf{z}, r),$ $G_r(q, \hat{q}, r), G_t(\mathbf{u})),$ $\mathcal{B} \in L(\mathbb{Q} \times \mathfrak{X}, \mathfrak{Z}),$ $\widehat{\mathcal{F}} = [\mathcal{F}_{\mathcal{B}}, \mathcal{F}]^T, \, \mathcal{F}_{\mathcal{B}} = (\hat{g}, \, \ell_S, \, g), \, \mathcal{F} = (\mathbf{f}, \, \psi, \, \hat{\psi}, \, \mathbf{f}_{\triangle}, \, \varrho, \, \hat{\varrho}, \, \hat{\mathbf{f}}),$ $\widehat{\Upsilon}\widehat{\mathscr{F}} = [\mathscr{J}_{\mathcal{I}}\mathscr{F}_{\mathcal{B}}, \Upsilon \mathscr{F}]^T, \, \mathscr{J}_{\mathcal{I}}\mathscr{F}_{\mathcal{B}} = \mathscr{F}_{\mathcal{B}}, \, \Upsilon \mathscr{F} = (T_{VPD}\mathbf{f}, \, T_{PD}\psi, \, T_{PD}\hat{\psi}, \, \pi\mathbf{f}_{\wedge}, \, B_1\varrho,$ $B_1\hat{\rho}, \,\pi\hat{\mathbf{f}}),$ $\mathfrak{T}_h \in L(\mathfrak{Y}, \mathfrak{X}_h), \, \mathfrak{T}_h = (T_{VPD,h}, T_{PD,h}, T_{PD,h}, \pi_h, B_{1,h}, B_{1,h}, \pi_h),$ $\widehat{\mathfrak{T}}_h\widehat{\mathfrak{F}} = [\mathfrak{J}_{\mathbb{Z}}\mathfrak{F}_{\mathbb{B}},\mathfrak{T}_h\mathfrak{F}]^T, \ \mathfrak{J}_{\mathbb{Z}}\mathfrak{F}_{\mathbb{B}} = \mathfrak{F}_{\mathbb{B}}, \ \mathfrak{T}_h\mathfrak{F} = (T_{VPD,h}\mathbf{f}, \ T_{PD,h}\psi, \ T_{PD,h}\hat{\psi}, \ T_{PD,h}\hat{\psi}$ $\pi_h \mathbf{f}_{\Delta}, B_{1,h} \varrho, B_{1,h} \hat{\varrho}, \pi_h \mathbf{f}),$ $\mathfrak{X}_{d} = \mathbf{H}_{0}^{2}(\Omega) \times H_{0}^{2}(\Omega)^{2} \times \mathbf{H}_{0}^{1}(\Omega) \times H^{m+1}(\Omega) \cap L_{0}^{2}(\Omega) \times H^{m+1}(\Omega) \times$ $\mathbf{H}_{0}^{1}(\Omega),$

$$\begin{split} & \mathcal{S} = (C^{\infty}(\overline{\Omega}) \cap H^1_0(\Omega))^{N+2} \times C^{\infty}_0(\Omega)^N \times C^{\infty}(\overline{\Omega}) \cap L^2_0(\Omega) \times C^{\infty}(\overline{\Omega}) \times \\ & C^{\infty}_0(\Omega)^N. \\ & \text{The correspondences between components are shown in the following } \end{split}$$

tables, where we skip " (Ω) " in the notations.

	()			â		â		â	â	
	(x, U)	Ĵ		$\frac{1}{u^1}$		$\frac{\Gamma_0}{\mu_1}$		$\frac{1}{1}$	Γ_h	
	p			L^2_0		$H^1 \cap L^2_0$	H^1	$\cap L^2_0$	L^2_0	
	y		T_P^-	$L_D^{-0}(D_0) = L_0^2$	T_P^-	${}^{1}_{D}(D_0 \cap H_0^2)$		L_0^2	L_0^{-0}	
	u	1	11	\mathbf{H}_{0}^{1}	11	$V \oplus \widetilde{V^{\perp}}$]	\mathbf{H}_{0}^{1}	\mathbf{X}_{0h}	
	q	2	$H_0^{\check{1}}$		H_0^1		H_0^1		Y_{0h}	
	\hat{q}	3	H_0^1		H_0^1		H_0^1 Y		Y_{0h}	
	\mathbf{Z}	4	L^2		L^2		L^2		\mathbf{X}_h	
	$p \\ r$	5 6	$H^1 \cap L^2_0$ H^1		$H^1 \sqcup L_0^-$ H^1		$H^1 \sqcup L_0^-$ H^1		M_h V_l	
	\mathbf{t}	$\frac{1}{7}$	\mathbf{L}^2			\mathbf{L}^2		\mathbf{L}^2		
	(y, U)	j	$\widehat{\mathcal{F}}$	$\widehat{\Sigma}$		$\widehat{\Sigma}_0$			$\widehat{\Sigma}_1$	
	\hat{p}		\hat{g} H^1			H^1		H^1		
	p_S		$\ell_S = L_0^2$		τ2	$\begin{bmatrix} H^{1} \cap L_{0}^{2} \\ T^{-1} (D \cap H^{2}) \end{bmatrix}$		$H^{1} \cap L_{0}^{2}$		
	$\frac{y}{y}$	1	$\begin{array}{c c}g\\f\end{array} & I_{PD}(D_0) = I\\ \mathbf{H}^{-1}\end{array}$		L_0	$\Gamma_{PD}(D_0 + H_0)$ Υ_0		$\Gamma_0 \\ \Upsilon_1$		
	q	2	ψ	ψ H^{-1}		H^{-1}		H^{-1}		
	\hat{q}	3	$\hat{\psi}$	$\hat{\psi}$ H^{-1}		H^{-1}		H^{-1}		
	\mathbf{Z}	4	\mathbf{f}_{\triangle} \mathbf{L}^2			\mathbf{L}^2		\mathbf{L}^2		
	p	5	$\varrho \mid B_1^{-1}(H^1 \cap L)$		$L_0^2)$	$B_1^{-1}(H^1 \cap L_0^2)$		$B_1^{-1}(H^1 \cap L_0^2)$		
	r	6 7	$ \begin{array}{c} \varrho \\ \hat{\mathbf{f}} \\ \end{array} $ $(H^{\perp})'$ $(H^{\perp})'$			$(H^{\perp})^{\prime}$		$(H^{T})^{r}$ T^{2}		
	t	1	I			L			L	
	(u, U)	i	For $\widehat{\Gamma}$, topology of		of	\mathfrak{X}_d		S		
	\hat{p}	J	H^1			-		-		
	p_S			L_{0}^{2}		-		-		
	y	1		$L_0^2 (\text{or } L^2)$		- TT2		-		
	u	1		\mathbf{H}_{0}^{1}	\mathbf{H}_{0}^{2} ((C^{α})	$(C^{\infty}(\Omega) \cap H_0^1)^N = C^{\infty}(\overline{\Omega}) \cap H_0^1$		
ļ	\hat{q}	⊿ 3		H_{1}^{1}	$\begin{array}{c c} & \Pi_{\overline{0}} \\ & H_{2}^{2} \end{array}$		C°	$C^{\infty}(\underline{\Omega}) \cap H^{1}_{2}$		
ļ	ч Z	4		\mathbf{L}^2		\mathbf{H}_{0}^{n}		$C_0^{\infty}(\Omega)^N$		
ļ	p	5	H^1			$H^{m+1} \cap L^2_0$		$C^{\infty}(\overline{\Omega}) \cap L^2_0$		
	r	6		H^1		H^{m+1}		$C^{\infty}(\overline{\Omega})$		
	\mathbf{t}	7	\mathbf{L}^2			\mathbf{H}_{0}^{1}		$C_0^\infty(\Omega)^N$		

Lemma 5.3. $\widehat{\Gamma}_0$ and $\widehat{\Gamma}_1$ are dense subspaces of $\widehat{\Gamma}$. $\widehat{\Sigma}_0$ and $\widehat{\Sigma}_1$ are dense subspaces of $\widehat{\Sigma}$. The embedding $\chi_d \subseteq \chi$ is continuous and the embedding operator i_0 is compact. S is a set dense in \mathfrak{X} .

CĂTĂLIN - LIVIU BICHIR

Proof. The elements from the column of $\widehat{\Gamma}_0$ are dense in the corresponding elements, on a line, from the column of $\widehat{\Gamma}$ and so on. We use the references [1, 2, 6, 7, 10, 11, 20, 21, 34, 46, 58].

6. An extended system based on the Stokes problem

6.1. The formulation of the extended system. Let us introduce the following exact extended system, related to the Stokes problem, using the notations from Section 5: given a fixed $\alpha \neq 1$ and $\widehat{\mathcal{F}} \in \widehat{\Sigma}$, find $(x, U) \in \widehat{\Gamma}$ such that

- (6.1) $p (q + \hat{p}) = \hat{g},$
- $(6.2) \qquad p_S p = \ell_S \,,$
- (6.3) $(\operatorname{div} \mathbf{u} y, \hat{\mu}) = (g, \hat{\mu}), \ \forall \, \hat{\mu} \in L^2_0(\Omega),$

(6.4)
$$(\operatorname{grad} \mathbf{u}, \operatorname{grad} \mathbf{w}) - (p_S, \operatorname{div} \mathbf{w}) = \langle \mathbf{f}, \mathbf{w} \rangle, \ \forall \, \mathbf{w} \in \mathbf{H}_0^1(\Omega),$$

- $(6.5) \quad -\alpha \langle \widetilde{S}_{\mathbf{z}}, \overline{\lambda} \rangle + \langle \widetilde{S}_{\mathbf{t}}, \overline{\lambda} \rangle + (\operatorname{grad} q, \operatorname{grad} \overline{\lambda}) = \langle \psi, \overline{\lambda} \rangle, \ \forall \, \overline{\lambda} \in H^1_0(\Omega) \,,$
- (6.6) $(\operatorname{grad} \hat{q}, \operatorname{grad} \tilde{\lambda}) + (\operatorname{grad} p, \operatorname{grad} \tilde{\lambda}) = \langle \hat{\psi}, \tilde{\lambda} \rangle, \ \forall \, \tilde{\lambda} \in H_0^1(\Omega),$

(6.7)
$$(\mathbf{z}, \tilde{\mathbf{w}}) - (p, div \, \tilde{\mathbf{w}}) = (\mathbf{f}_{\triangle}, \tilde{\mathbf{w}}), \, \forall \, \tilde{\mathbf{w}} \in \mathbf{H}_0^1(\Omega)$$

(6.8)
$$(\mathbf{z} - \mathbf{t} + grad \, (\hat{p} - \hat{q}), grad \, \varphi) + (grad \, r, grad \, \varphi) = \langle \varrho, \varphi \rangle, \, \forall \, \varphi \, \in \, H^1(\Omega) \, ,$$

(6.9)
$$(r,\tilde{\varphi}) + (grad(q+\hat{q}), grad\tilde{\varphi}) = \langle \hat{\varrho}, \tilde{\varphi} \rangle, \ \forall \tilde{\varphi} \in H^1(\Omega),$$

(6.10) $(\mathbf{t}, \hat{\mathbf{w}}) = (\hat{\mathbf{f}}, \hat{\mathbf{w}}), \forall \hat{\mathbf{w}} \in \mathbf{L}^2(\Omega),$

Remark 6.1. Since \mathbf{z} , grad p, $\mathbf{f}_{\Delta} \in \mathbf{L}^{2}(\Omega)$ and $H_{0}^{1}(\Omega)$ is dense in $L^{2}(\Omega)$, (6.7) is equivalent to the following formulations

(6.11)
$$(\mathbf{z}, \tilde{\mathbf{w}}) + (\operatorname{grad} p, \tilde{\mathbf{w}}) = (\mathbf{f}_{\triangle}, \tilde{\mathbf{w}}), \ \forall \, \tilde{\mathbf{w}} \in \mathbf{L}^2(\Omega),$$

$$\mathbf{z} + grad\, p = \mathbf{f}_{\triangle}$$

In the sequel, we work with (6.11) instead of (6.7). We use \mathcal{M}_5 and $\mathcal{M}_{h,5}$ below.

Remark 6.2. Assume that problem (6.1) - (6.10) has a solution. Let us introduce p_S from (6.2) into (6.4) and then replace $(p, div \mathbf{w})$ using (6.7). We obtain

(6.13)
$$(\operatorname{grad} \mathbf{u}, \operatorname{grad} \mathbf{w}) - (\mathbf{z}, \mathbf{w}) = (\ell_S, \operatorname{div} \mathbf{w}) + \langle \mathbf{f}, \mathbf{w} \rangle - (\mathbf{f}_{\triangle}, \mathbf{w}), \ \forall \, \mathbf{w} \in \mathbf{H}_0^1(\Omega) \,.$$

Remark 6.3. Equation (6.4) is the weak formulation (5.1) of the fist equation of the Stokes system (that is, of the momentum equation). Equation (6.3) corresponds to (5.2).

Equation (6.1) corresponds to (2.15).

We introduce equation (6.2) to relate $p_S \in L^2_0(\Omega)$ and $p \in H^1(\Omega) \cap L^2_0(\Omega)$ in order to use Lemma 3.3. $p_S = p$ for $\ell_S = 0$.

(6.5) corresponds to (2.16) and introduces (5.9).

(6.6) corresponds to (2.17) rewritten as an homogeneous Dirichlet prob-

Dirichlet boundary condition). The connection between p, \hat{p} and \hat{q} is in (6.8).

Equations (6.7) and (6.8) are two forms of the momentum equation. Equation (6.7) defines \mathbf{z} to be $-\Delta \mathbf{u}$.

Equation (6.8) is (5.7) and it is like (2.19).

Equation (6.8) gives an equation, on the boundary, for the pressure p in the approximate case. We regain the equation, containing the pressure, that we lose when the velocity boundary condition is implemented, for instance, in the case of nodal finite element method.

The unknown r and equation (6.9) are introduced in order to avoid a compatibility condition between some components of the right hand term.

Equation (6.10) is intended to introduce the free term \mathbf{f} between the unknowns.

We do not use a boundary unknown λ_w , as in (2.18), that is, as in [34, 35]. We do not use a variational problem related to $\partial\Omega$.

Remark 6.4. The following equation, which corresponds to (5.9),

(6.14)
$$-\alpha \langle S_{\mathbf{z}}, \lambda \rangle + \langle S_{\mathbf{t}}, \lambda \rangle + (\operatorname{grad} p, \operatorname{grad} \lambda) = \langle \psi, \lambda \rangle, \ \forall \lambda \in H_0^1(\Omega),$$

is obtained from (6.24).

Remark 6.5. Replacing (6.1), (6.10) and (6.12) in (6.8), it results

(6.15)
$$(\mathbf{f}_{\bigtriangleup} - \mathbf{f} - \operatorname{grad} \hat{g}, \operatorname{grad} \varphi)$$

 $-(grad (q + \hat{q}), grad \varphi) + (grad r, grad \varphi) = \langle \varrho, \varphi \rangle, \ \forall \varphi \in H^1(\Omega).$

Let $\tilde{\psi}_r \in (H^1(\Omega))'$ be defined by

(6.16)
$$\langle \tilde{\psi}_r, \varphi \rangle = (-\mathbf{f}_{\triangle} + \hat{\mathbf{f}} + \operatorname{grad} \hat{g}, \operatorname{grad} \varphi) + \langle \varrho, \varphi \rangle + \langle \hat{\varrho}, \varphi \rangle, \ \forall \varphi \in H^1(\Omega) ,$$

Summing (6.15) and (6.9), we obtain that r is the solution of the equation

(6.17)
$$((r,\varphi))_1 = \langle \tilde{\psi}_r, \varphi \rangle, \ \forall \varphi \in H^1(\Omega),$$

so $r = B_1 \tilde{\psi}_r$. Then, in (6.8), $(\operatorname{grad} r, \operatorname{grad} \varphi)$ is $(\operatorname{grad} B_1 \tilde{\psi}_r, \operatorname{grad} \varphi)$. Equation (6.9) is written

(6.18)
$$((r,\tilde{\varphi}))_1 + (grad (q + \hat{q}), grad \tilde{\varphi}) - (grad r, grad \tilde{\varphi}) = \langle \hat{\varrho}, \tilde{\varphi} \rangle, \ \forall \, \tilde{\varphi} \in H^1(\Omega) \,,$$

 \mathbf{SO}

(6.19)
$$r - B_1 G_r(q, \hat{q}, r) = B_1 \hat{\varrho},$$

From (6.10) and (6.12), we deduce

(6.20)
$$(\mathbf{z} - \mathbf{t} + \operatorname{grad} p, \operatorname{grad} \varphi) = (\mathbf{f}_{\bigtriangleup} - \hat{\mathbf{f}}, \operatorname{grad} \varphi), \ \forall \varphi \in H^1(\Omega),$$

The same is obtained if we replace (6.1), (6.10) in (6.8) and then sum (6.9) to (6.8), where we use (6.17). In both cases, we have a form of equation (5.8).

6.2. The well-posedness of the exact extended system. Let $\widehat{\Phi}$ be the operator defined by the left hand side of problem (6.1) - (6.10). For this problem, the spaces $\widehat{\Gamma}$, $\widehat{\Sigma}$, $\widehat{\Gamma}_0$, $\widehat{\Sigma}_0$ are defined in Section 5. $\widehat{\Phi} \in L(\widehat{\Gamma}, \widehat{\Sigma})$. Then, problem (6.1) - (6.10) has the forms (3.1) and (4.1). Let us use Lemma 3.3 in order to prove that $\widehat{\Phi}$ is an isomorphism of $\widehat{\Gamma}$ onto $\widehat{\Sigma}$. Let $\widehat{\Phi}_0$ be the restriction of $\widehat{\Phi}$ to $\widehat{\Gamma}_0$. Consider equation (3.2) in this particular case.

Lemma 6.6. $\widehat{\Phi}_0$ is a bijection of $\widehat{\Gamma}_0$ onto $\widehat{\Sigma}_0$.

Proof. We constructed $\widehat{\Gamma}_0$ and $\widehat{\Sigma}_0$ such that $\widehat{\Phi}(\widehat{\Gamma}_0) \subseteq \widehat{\Sigma}_0$. We have $\widehat{\Phi}_0(x, U)$ $= \widehat{\Phi}(x, U), \forall (x, U) \in \widehat{\Gamma}_0.$

Fix $\widehat{\mathcal{F}}_0 = (\hat{g}, \ell_S, g, \mathbf{f}, \psi, \hat{\psi}, \mathbf{f}_{\Delta}, \varrho, \hat{\varrho}, \hat{\mathbf{f}}) \in \widehat{\Sigma}_0.$ By the density of $\mathcal{D}(\Omega)^N$ in $\mathbf{H}_0^1(\Omega)$, from (6.4), we obtain equation (2.1) in the sense of distributions,

(6.21)
$$\langle -\Delta \mathbf{u} + grad \, p_S - \mathbf{f}, \hat{\varphi} \rangle = 0, \forall \, \hat{\varphi} \in \mathcal{D}(\Omega)^N$$

Let $\varphi \in \mathcal{D}(\Omega)$. Then, $\langle \bigtriangleup \mathbf{u}, grad \, \varphi \rangle = \langle \mathbf{u}, \bigtriangleup (grad \, \varphi) \rangle = \langle \mathbf{u}, grad \, (\bigtriangleup \varphi) \rangle$ $= \langle div \mathbf{u}, -\Delta \varphi \rangle$. Using (6.21) with $\hat{\varphi} = grad \varphi$, we deduce

(6.22)
$$\langle div \mathbf{u}, -\Delta \varphi \rangle = \langle grad p_S, grad \varphi \rangle - \langle \mathbf{f}, grad \varphi \rangle, \ \forall \varphi \in \mathcal{D}(\Omega).$$

Using (6.2), we obtain

(6.23)
$$\langle div \, \mathbf{u}, -\Delta \varphi \rangle = \langle grad \, (p + \ell_S), grad \, \varphi \rangle - \langle \mathbf{f}, grad \, \varphi \rangle, \ \forall \, \varphi \in \mathcal{D}(\Omega) \,.$$

Taking the test functions $\varphi \in H_0^1(\Omega)$ in (6.8), replacing \hat{p} from (6.1) in (6.8), summing (6.5), (6.6) to (6.8), using (6.12) and $r = B_1 \tilde{\psi}_r$ from (6.17), we obtain

(6.24)
$$(\operatorname{grad} p, \operatorname{grad} \tilde{\lambda}) = (\operatorname{grad} \hat{g}, \operatorname{grad} \tilde{\lambda}) + \alpha \langle \tilde{S}_{\mathbf{z}}, \tilde{\lambda} \rangle - \langle \tilde{S}_{\mathbf{t}}, \bar{\lambda} \rangle$$

 $+ \langle \psi, \tilde{\lambda} \rangle + \langle \hat{\psi}, \tilde{\lambda} \rangle - \langle \mathbf{f}_{\Delta}, \operatorname{grad} \tilde{\lambda} \rangle + \langle \varrho, \tilde{\lambda} \rangle - (\operatorname{grad} (B_1 \tilde{\psi}_r), \operatorname{grad} \tilde{\lambda})$
 $+ \langle \hat{\mathbf{f}}, \operatorname{grad} \tilde{\lambda} \rangle, \ \forall \ \tilde{\lambda} \in \mathcal{D}(\Omega),$

Using (6.12) and (6.10), relation (6.24) becomes:

(6.25)
$$(1-\alpha)(\operatorname{grad} p, \operatorname{grad} \varphi) = (\operatorname{grad} \hat{g}, \operatorname{grad} \varphi) + \langle \psi, \varphi \rangle + \langle \tilde{\psi}, \varphi \rangle$$
$$-(1+\alpha)\langle \mathbf{f}_{\triangle}, \operatorname{grad} \varphi \rangle + \langle \varrho, \varphi \rangle - (\operatorname{grad} (B_1 \tilde{\psi}_r), \operatorname{grad} \varphi)$$
$$+(1+\gamma)\langle \hat{\mathbf{f}}, \operatorname{grad} \varphi \rangle, \ \forall \varphi \in \mathcal{D}(\Omega),$$

Relation (6.23) and (6.25) give

(6.26)
$$(\operatorname{div} \mathbf{u}, -\Delta \varphi) = \langle \psi_{\operatorname{div}}, \varphi \rangle, \ \forall \varphi \in \mathcal{D}(\Omega).$$

where $\psi_{div} \in H^{-1}(\Omega)$ is

(6.27)
$$\langle \psi_{div}, \varphi \rangle = (1-\alpha)^{-1} (grad \,\hat{g}, grad \,\varphi) + \langle grad \,\ell_S, grad \,\varphi \rangle$$

 $+ \langle \widetilde{S}_{\mathbf{f}}, \varphi \rangle + (1-\alpha)^{-1} (\langle \psi, \varphi \rangle + \langle \hat{\psi}, \varphi \rangle + (1+\alpha) \langle \widetilde{S}_{\mathbf{f}_{\Delta}}, \varphi \rangle$
 $+ \langle \varrho, \varphi \rangle - (grad \,(B_1 \tilde{\psi}_r), grad \,\varphi) - (1+\gamma) \langle \widetilde{S}_{\mathbf{f}}, \varphi \rangle), \ \forall \varphi \in H_0^1(\Omega),$

Let ψ_g be given by the sum of some distributional derivatives of g by

(6.28)
$$(g, \bigtriangleup \varphi) = \langle \psi_g, \varphi \rangle, \ \forall \varphi \in \mathcal{D}(\Omega).$$

Equation (6.3) gives $div \mathbf{u} = y + g$ and with (6.26) and (6.28), we have

(6.29)
$$(y, -\Delta\varphi) = \langle \psi_g + \psi_{div}, \varphi \rangle, \ \forall \varphi \in \mathcal{D}(\Omega)$$

The density of $\mathcal{D}(\Omega)$ in $H^2_0(\Omega)$ leads to

(6.30)
$$(y, -\Delta \varphi) = \langle \psi_g + \psi_{div}, \varphi \rangle, \ \forall \varphi \in H^2_0(\Omega).$$

or, using the space \mathcal{W} as the space \mathcal{V} is used in Theorem 7.1.1, page 384, [20] ($\Delta \varphi \in L^2(\Omega)$ since $\varphi \in H^2_0(\Omega)$),

(6.31)
$$(y, -\Delta \varphi) = \langle \psi_g + \psi_{div}, \varphi \rangle, \ \forall (\varphi, -\Delta \varphi) \in \mathcal{W}.$$

Since $y \in T_{PD}^{-1}(D_0 \cap H_0^2)$, there exists $\phi \in H_0^2(\Omega)$ such that

$$(6.32) \qquad \qquad (\phi, y) \in \mathcal{W},$$

Following the technique from the proof of Theorem 7.1.2, page 385, [20], we infer that (6.32) and (6.31) are equivalent to (6.32) and (6.33), where

(6.33)
$$(\triangle \phi, \triangle \varphi) = \langle \psi_g + \psi_{div}, \varphi \rangle, \ \forall \varphi \in H^2_0(\Omega).$$

Equation (6.33) satisfies the conditions of Proposition I.1.3, page 16, [34] (where we take $u_0 = 0$), for a bounded and connected open subset of \mathbb{R}^N (N = 2, 3) with a Lipschitz - continuous boundary $\partial\Omega$, so this equation has an unique solution $\phi \in H_0^2(\Omega)$.

y is uniquely determined by (6.32).

 (\mathbf{u}, p_S) is the solution of the Stokes problem given by (6.4), (6.3).

 $p, \mathbf{z}, \mathbf{t}, q, \hat{p}, \hat{q}, r$ are obtained from (6.2), (6.7), (6.10), (6.5), (6.1), (6.6), (6.17) respectively. So $\widehat{\Phi}_0$ is surjective.

If $\widehat{\mathcal{F}}_0 = (\widehat{g}, \ell_S, g, \mathbf{f}, \psi, \widehat{\psi}, \mathbf{f}_{\triangle}, \varrho, \widehat{\varrho}, \widehat{\mathbf{f}}) = 0$, we deduce $\widehat{U}_0 = (\widehat{p}, p_S, y, \mathbf{u}, q, \widehat{q}, \mathbf{z}, p, r, \mathbf{t})$ = 0 uniquely determined. $Ker(\widehat{\Phi}_0) = \{0\}$ and $\widehat{\Phi}_0$ is injective. So $\widehat{\Phi}_0$ is a bijection of $\widehat{\Gamma}_0$ onto $\widehat{\Sigma}_0$.

Lemma 6.7. $\widehat{\Phi}$ is a linear, continuous and surjective mapping.

Proof. Fix $\widehat{\mathcal{F}}$ and $\widehat{\mathcal{F}}_0 \in \widehat{\Sigma}$.

 $\widehat{\mathcal{F}} = (\widehat{g}, \, \ell_S, \, g, \, \mathbf{f}, \, \widehat{\psi}, \, \widehat{\psi}, \, \mathbf{f}_{\triangle}, \, \varrho, \, \widehat{\varrho}, \, \widehat{\mathbf{f}}).$

 $\widehat{\mathcal{F}}_0 = (\widehat{g}, (\ell_S)_0, g_0, \mathbf{f}_0, \psi, \widehat{\psi}, \mathbf{f}_{\triangle}, \varrho, \widehat{\varrho}, \widehat{\mathbf{f}})$, where the index "0" is written only for the elements that do not coincide with the corresponding element from $\widehat{\mathcal{F}}$.

 $\widehat{\Phi}_0: \widehat{\Gamma}_0 \to \widehat{\Sigma}_0$ is bijective. Let $\widehat{U}_0 = (x_0, U_0) = (\widehat{p}_0, (p_S)_0, y_0, \mathbf{u}_0, q_0, \widehat{q}_0, \mathbf{z}_0, p_0, r_0, \mathbf{t}_0)$ be the solution of

(6.34)
$$\widehat{\Phi}_0(\widehat{U}) = \widehat{\mathcal{F}}_0,$$

We have $\widehat{\mathcal{F}} - \widehat{\mathcal{F}}_0 = (0, \ell_S - (\ell_S)_0, g - g_0, \mathbf{f} - \mathbf{f}_0, 0, 0, 0, 0, 0, 0)$. Consider the equation

(6.35)
$$\widehat{\Phi}(\widehat{V}) = \widehat{\mathcal{F}} - \widehat{\mathcal{F}}_0$$

Let us determine a solution $\widehat{U}' = (x', U') = (\hat{p}', p'_S, y', \mathbf{u}', q', \hat{q}', \mathbf{z}', p', r', \mathbf{t}')$ of (6.35).

If we take $\hat{p}' = 0$, q' = 0, $\hat{q}' = 0$, $\mathbf{z}' = 0$, p' = 0, r' = 0, $\mathbf{t}' = 0$, then, from the equations (6.1) - (6.10), it remains

 $(6.36) \qquad p'_S = \ell_S - (\ell_S)_0 \,,$

(6.37) $(div \mathbf{u}' - y', \hat{\mu}) = (g - g_0, \hat{\mu}), \ \forall \, \hat{\mu} \in L^2_0(\Omega),$

(6.38) $(\operatorname{grad} \mathbf{u}', \operatorname{grad} \mathbf{w}) - (p'_S, \operatorname{div} \mathbf{w}) = \langle \mathbf{f} - \mathbf{f}_0, w \rangle, \ \forall \, \mathbf{w} \in \mathbf{H}_0^1(\Omega),$

We obtain $p'_S = \ell_S - (\ell_S)_0$ from (6.36), **u'** from (6.38) and y' from (6.37), so \widehat{U}' is a solution of the equation (6.35). We have

(6.39)
$$\widehat{\Phi}(\widehat{U}_0 + \widehat{U}') = \widehat{\mathcal{F}},$$

Hence $\widehat{\Phi}$ is a surjective mapping.

Remark 6.8. \widehat{U}' and $\widehat{\mathcal{F}} - \widehat{\mathcal{F}}_0$ also satisfy (6.13). $\forall \mathbf{w} \in \mathbf{H}_0^1(\Omega)$, we have

$$(grad \mathbf{u}', grad \mathbf{w}) - (\ell_S - (\ell_S)_0, div \mathbf{w}) - \langle \mathbf{f} - \mathbf{f}_0, w \rangle = (\mathbf{z}', \mathbf{w}).$$

Theorem 6.9. $\widehat{\Phi}$ is an isomorphism of $\widehat{\Gamma}$ onto $\widehat{\Sigma}$.

Proof. $\widehat{\Gamma}$, $\widehat{\Sigma}$, $\widehat{\Gamma}_0$, $\widehat{\Sigma}_0$, $\widehat{\Phi}$, $\widehat{\Phi}_0$ satisfy the hypotheses of Lemma 3.3.

Remark 6.10. We do not impose condition (2.14). This one arises for $y \in T_{PD}^{-1}(D_0 \cap H_0^2(\Omega))$ in the form $\phi \in D_0 \cap H_0^2(\Omega)$, related to some dense subspaces. See (6.32). In this situation, we also have $\mathbf{u} \in V \oplus \widetilde{V^{\perp}}$.

Corollary 6.11. Let $\mathbf{f} \in \Upsilon_1 \cap \mathbf{L}^2(\Omega)$ be given. If $\hat{g} = 0$, $\ell_S = 0$, g = 0, $\psi = 0$, $\hat{\psi} = 0$, $\mathbf{f}_{\triangle} = \mathbf{f}$, $\varrho = 0$, $\hat{\varrho} = 0$, $\hat{\mathbf{f}} = \mathbf{f}$ in problem (6.1) - (6.10), then $(\mathbf{u}, p) \in \mathbf{H}_0^1(\Omega) \times (H^1(\Omega) \cap L_0^2(\Omega))$ is the unique solution of problem (5.1) - (5.2) with g = 0.

Proof. We obtain $\psi_{div} = 0$, $\psi_g = 0$, so $\phi = 0$, and so on.

Corollary 6.12. Let $\mathbf{f} \in \Upsilon_1 \cap \mathbf{L}^2(\Omega)$ be given. To solve the problem (5.1) - (5.2) with g = 0 is equivalent to solve the problem (6.1) - (6.10) with $\hat{g} = 0, \ell_S = 0, g = 0, \psi = 0, \hat{\psi} = 0, \mathbf{f}_{\triangle} = \mathbf{f}, \varrho = 0, \hat{\varrho} = 0, \hat{\mathbf{f}} = \mathbf{f}.$

Corollary 6.13. Under the hypotheses of Corollaries 6.11 and 6.12, for $\alpha_1 \neq \alpha_2$, we have $(x(\alpha_1), U(\alpha_1)) = (x(\alpha_2), U(\alpha_2))$, where $(x(\alpha_i), U(\alpha_i))$ is the solution of (6.1) - (6.10) for α_i .

Proof. Under the hypotheses of the Corollary, $\psi_{div} = 0, \forall \alpha \neq 1$.

Corollary 6.14. Under the hypotheses of Corollaries 6.11 and 6.12, $(\mathbf{z}, \operatorname{grad} \varphi) = 0, \forall \varphi \in H_0^1(\Omega).$

Proof. Using (6.7), we deduce

(6.40)
$$(\mathbf{z}, \operatorname{grad} \varphi) + (\operatorname{grad} p, \operatorname{grad} \varphi) = (\mathbf{f}_{\triangle}, \operatorname{grad} \varphi), \forall \varphi \in \mathcal{D}(\Omega),$$

Subtracting (6.40) from (6.24), we obtain

(6.41)
$$(1-\alpha)(\mathbf{z}, \operatorname{grad} \varphi) = 0, \forall \varphi \in \mathcal{D}(\Omega),$$

Hence, $(\mathbf{z}, grad \varphi) = 0, \forall \varphi \in H_0^1(\Omega).$

22

Corollary 6.15. We have

(6.42)
$$(\alpha \boldsymbol{z} - \boldsymbol{t}, \operatorname{grad} \bar{\lambda}) + (\operatorname{grad} p, \operatorname{grad} \bar{\lambda}) = (r, \bar{\lambda}) + \langle \psi, \bar{\lambda} \rangle + \langle \hat{\psi}, \bar{\lambda} \rangle - \langle \hat{\varrho}, \bar{\lambda} \rangle, \ \forall \bar{\lambda} \in H_0^1(\Omega),$$

where we use (6.17), and

$$(6.43)(\boldsymbol{z} - \boldsymbol{t} + \operatorname{grad} p, \operatorname{grad} \varphi) = (\boldsymbol{f}_{\bigtriangleup} - \boldsymbol{f}, \operatorname{grad} \varphi), \ \forall \, \varphi \, \in \, H^1(\Omega) \ominus H^1_0(\Omega) \,,$$

Proof. If we add (6.5) and (6.6) and subtract (6.9), where the test functions $\varphi \in H_0^1(\Omega)$, we have (6.42) where we use (6.17).

(6.20) gives (6.43).

The parameterized perturbed pressure Poisson equation is given by (6.42) and (6.43). In the following corollary, we formulate the announced result that to solve the stationary Stokes problem is equivalent to solve a problem for the momentum equation, the parameterized perturbed pressure Poisson equation and the equation that defines the Laplace operator acting on velocity.

Corollary 6.16. Let $\mathbf{f} \in \Upsilon_1 \cap \mathbf{L}^2(\Omega)$ be given. To solve the problem (5.1) - (5.2) with g = 0 is equivalent to solve the problem (6.4), (6.13) (6.42), (6.43) with $\hat{g} = 0$, $\ell_S = 0$, g = 0, $\psi = 0$, $\hat{\psi} = 0$, $\mathbf{f}_{\triangle} = \mathbf{f}$, $\varrho = 0$, $\hat{\varrho} = 0$, $\hat{\mathbf{f}} = \mathbf{f}$.

Proof. The proof follows from Corollary 6.11 and the proof of Lemma 6.6 using the density.

7. Construction of another formulation of the isomorphism for the exact problem

In the sequel, we write the problem (6.1) - (6.10) in the form of equation (4.2).

Let us consider the generic form of a variational equation: find $v \in E$ such that

(7.1)
$$a(v,\lambda) = \langle \psi, \lambda \rangle, \ \forall \lambda \in E$$

[2, 6, 10, 20, 26, 31, 40, 46, 54, 58]. E is a Hilbert space, \mathcal{E} is a dense subset in E. $a(\cdot, \cdot)$ is a continuous, (strongly) coercive bilinear form on $E \times E$ and $\langle \psi, \cdot \rangle$ is a linear continuous functional on E. $T \in L(E', E)$, T is the solution operator defined by: given $\psi \in E'$, $v = T\psi$ if and only if $v \in E$ is the unique solution of (7.1) (it exists and it is unique according to Lax -Milgram lemma). T is an isomorphism from E' onto E.

Let us associate a set $\mathcal{M}_i = \{i, (7.1), E, E', \mathcal{E}_i, v, \psi, T\}$ to equation (7.1), where the first element is an index *i* to be fixed for an particular equation, the second element is the number of the equation. Let us retain the following particular cases $\mathcal{M}_1, \ldots, \mathcal{M}_5$ of \mathcal{M}_i (In all the cases, the bilinear form $a(\cdot, \cdot)$ is an inner product):

 $\mathcal{M}_{1} = \{1, (7.2), \mathbf{H}_{0}^{1}(\Omega), \mathbf{H}^{-1}(\Omega), \mathcal{E}_{1}, \mathbf{u}, \mathbf{f}, T_{VPD}\} \text{ with}$ $(7.2) \qquad (grad \mathbf{u}, grad \mathbf{w}) = \langle \mathbf{f}, \mathbf{w} \rangle, \ \forall \mathbf{w} \in \mathbf{H}_{0}^{1}(\Omega).$

CĂTĂLIN - LIVIU BICHIR

 $\mathcal{M}_2 = \{2, (7.3), H_0^1(\Omega), H^{-1}(\Omega), \mathcal{E}_2, q, \psi, T_{PD}\}$ with

(7.3)
$$(\operatorname{grad} q, \operatorname{grad} \lambda) = \langle \psi, \lambda \rangle, \ \forall \lambda \in H^1_0(\Omega).$$

 $\mathbb{M}_3=\{3,\,(7.4),\,H^1(\Omega),\,H^1(\Omega)',\,\mathbb{E}_3,\,\phi,\,\psi_1,\,B_1\}$ with

(7.4)
$$(\operatorname{grad}\phi, \operatorname{grad}\mu) + (\phi, \mu) = \langle \psi_1, \mu \rangle, \ \forall \mu \in H^1(\Omega).$$

 $\mathcal{M}_4 = \{4, (7.4), H^1(\Omega) \cap L^2_0(\Omega), B_1^{-1}(H^1(\Omega) \cap L^2_0(\Omega)), \mathcal{E}_4, \phi, \psi_1, B_1\}.$ Let π be the identity operator on $\mathbf{L}^2(\Omega)$. We have: $\mathbf{z}, \mathbf{f}_* \in \mathbf{L}^2(\Omega), \pi \mathbf{f}_* = \mathbf{z}$ if and only if

(7.5)
$$(\mathbf{z}, \mathbf{w}) = (\mathbf{f}_*, \mathbf{w}), \ \forall \mathbf{w} \in \mathbf{L}^2(\Omega),$$

 $\mathfrak{M}_5 = \{5, (7.5), \, \mathbf{L}^2(\Omega), \, \mathbf{L}^2(\Omega), \, \mathfrak{E}_5, \, \mathbf{z}, \, \mathbf{f}_*, \, \pi\}.$

Let us define the following linear and continuous mappings:

$$\begin{split} & \mathcal{B}(y,U) = \begin{bmatrix} p - (q + \hat{p}) \\ p_{S} - p \\ div \,\mathbf{u} - y \end{bmatrix}, \\ & G_{\mathbf{u}} : L_{0}^{2}(\Omega) \rightarrow \mathbf{H}^{-1}(\Omega), \\ & \langle G_{\mathbf{u}}(p_{S}), \mathbf{w} \rangle = (p_{S}, div \,\mathbf{w}), \ \forall \,\mathbf{w} \in \mathbf{H}_{0}^{1}(\Omega), \\ & G_{q} : \mathbf{L}^{2}(\Omega)^{2} \rightarrow H^{-1}(\Omega), \ \langle G_{q}(\mathbf{t}, \mathbf{z}), \bar{\lambda} \rangle = \alpha \langle \tilde{S}_{\mathbf{z}}, \bar{\lambda} \rangle - \langle \tilde{S}_{\mathbf{t}}, \bar{\lambda} \rangle, \ \forall \bar{\lambda} \in H_{0}^{1}(\Omega), \\ & G_{\hat{q}} : H^{1}(\Omega) \cap L_{0}^{2}(\Omega) \rightarrow H^{-1}(\Omega), \\ & \langle G_{\hat{q}}(p), \hat{\lambda} \rangle = -(grad \, p, grad \, \hat{\lambda}), \ \forall \, \hat{\lambda} \in H_{0}^{1}(\Omega), \\ & G_{\mathbf{z}} : H^{1}(\Omega) \cap L_{0}^{2}(\Omega) \rightarrow \mathbf{L}^{2}(\Omega), \ G_{\mathbf{z}}(p) = -grad \, p, \\ & G_{p} : (H^{1}(\Omega) \cap L_{0}^{2}(\Omega)) \times H^{1}(\Omega) \times H_{0}^{1}(\Omega) \times \mathbf{L}^{2}(\Omega)^{2} \\ & \times H^{1}(\Omega) \rightarrow B_{1}^{-1}(H^{1}(\Omega) \cap L_{0}^{2}(\Omega)), \\ & \langle G_{p}(p, \hat{p}, \hat{q}, \mathbf{z}, \mathbf{t}, r), \varphi \rangle = -(\mathbf{z} - \mathbf{t} + grad \, (\hat{p} - \hat{q} + r), grad \, \varphi) \\ & + ((p, \varphi))_{1}, \ \forall \, \varphi \in H^{1}(\Omega), \\ & G_{r} : H_{0}^{1}(\Omega)^{2} \times H^{1}(\Omega) \rightarrow H^{1}(\Omega)', \\ & \langle G_{r}(q, \hat{q}, r), \varphi \rangle = -(grad \, (q + \hat{q} - r), grad \, \tilde{\varphi}), \ \forall \, \varphi \in H^{1}(\Omega), \\ & G_{\mathbf{t}} : \mathbf{H}_{0}^{1}(\Omega) \rightarrow \mathbf{L}^{2}(\Omega), \ G_{\mathbf{t}}(\mathbf{u}) = 0, \end{split}$$

where the index u indicates, in $G_{\mathbf{u}}$, that this function is used to construct the equation obtained by the perturbation of \mathbf{u} and so on.

Problem (6.1) - (6.10) becomes: find $(x, U) \in \widehat{\Gamma}$ such that

$$\begin{array}{ll} (7.6) & p - (q + \hat{p}) = \hat{g} \,, \\ (7.7) & p_S - p = \ell_S \,, \\ (7.8) & div \, \mathbf{u} - y = g \,, \\ (7.9) & \mathbf{u} - T_{VPD}G_{\mathbf{u}}(p_S) = T_{VPD}\mathbf{f} \,, \\ (7.10) & q - T_{PD}G_{q}(\mathbf{t}, \mathbf{z}) = T_{PD}\psi \,, \\ (7.11) & \hat{q} - T_{PD}G_{\hat{q}}(p) = T_{PD}\hat{\psi} \,, \\ (7.12) & \mathbf{z} - \pi G_{\mathbf{z}}(p) = \pi \mathbf{f}_{\Delta} \,, \\ (7.13) & p - B_1G_p(p, \hat{p}, \hat{q}, \mathbf{z}, \mathbf{t}, r) = B_1\varrho \,, \\ (7.14) & \mathbf{t} - \pi G_{\mathbf{t}}(\mathbf{u}) = \pi \hat{\mathbf{f}} \,, \end{array}$$

Lemma 7.1. Problem (7.6) - (7.15) has the form of equation (4.2).

Proof. Consider the spaces Q, X, Y, Z defined in Section 5. The operator $\widehat{\mathcal{A}}$ defined by the left hand side of (7.6) - (7.15) has the form of the operator $\widehat{\mathcal{A}}$ defined in Section 4.

Theorem 7.2. Assume the hypotheses of Theorem 6.9. Then, to solve (6.1) - (6.10) is equivalent to solve (7.6) - (7.15). In other words, (4.1) is equivalent to (4.2) in the particular case considered.

Theorem 7.3. \mathfrak{T} is an isomorphism of \mathfrak{Y} onto \mathfrak{X} . $\mathfrak{T}^{-1} = (T_{VPD}^{-1}, T_{PD}^{-1}, T_{PD}^{-1}, T_{PD}^{-1}, \pi^{-1}, B_1^{-1}, B_1^{-1}, \pi^{-1}), \mathfrak{T}^{-1}U = (T_{VPD}^{-1}\boldsymbol{u}, T_{PD}^{-1}\boldsymbol{q}, T_{PD}^{-1}\hat{\boldsymbol{q}}, \pi^{-1}\boldsymbol{z}, B_1^{-1}\boldsymbol{p}, B_1^{-1}\boldsymbol{r}, \pi^{-1}\boldsymbol{t}).$

8. An abstract approximation. Construction of the isomorphism for the approximate problem

8.1. The approximate problem. In the sequel, we write the approximate problem of the problem (6.1) - (6.10) in a general framework. We then formulate this approximate problem as equation (4.3). Particular approximations can be obtained by finite element method, finite differences method, finite volume method, spectral methods, wavelets and so on. In Section 9, we use finite element method.

We use the general approximation context formulated in [6, 34, 41]. Let X_h and Y_h be two closed subspaces of $H^1(\Omega)$. Following [20, 22, 34], we approximate $H^1(\Omega)$ and $L^2(\Omega)$ with the same space.

In order to approximate (6.1) - (6.10), we introduce the spaces $X_{0h} = X_h \cap H_0^1(\Omega) = \{x_h \in X_h; x_h = 0 \text{ on } \partial\Omega\},$ $M_h = Y_h \cap L_0^2(\Omega) = \{y_h \in Y_h; \int_{\Omega} y_h dx = 0\},$ $Y_{0h} = Y_h \cap H_0^1(\Omega) = \{y_h \in Y_h; y_h = 0 \text{ on } \partial\Omega\},$ $\mathbf{X}_{0h} = \mathbf{X}_h \cap \mathbf{H}_0^1(\Omega).$

Let us consider the approximate equation: find $v_h \in E_h$ such that

(8.1)
$$a(v_h, \lambda_h) = \langle \psi, \lambda_h \rangle, \ \forall \lambda_h \in E_h,$$

corresponding to (7.1) [2, 6, 10, 20, 26, 31, 40, 46, 54, 59]. E_h is a closed subspace of E. The hypotheses of Céa's lemma are satisfied for this problem. So (8.1) has an unique solution $v_h \in E_h$ and, for the solutions v, v_h of (7.1), (8.1), there exists $\gamma_{E,a} > 0$ (that depends on the continuity and ellipticity of $a(\cdot, \cdot)$ on E) such that

(8.2)
$$\|v - v_h\|_E \le \gamma_{E,a} \inf_{e_h \in E_h} \|v - e_h\|_E ,$$

Define the operator $T_h : E' \to E_h$ by: given $\psi \in E'$, $v_h = T_h \psi$ if and only if $v_h \in E_h$ is the unique solution of problem (8.1).

Let us associate the set $\mathcal{M}_{h,i} = \{i, \mathcal{M}_i, (7.1), (8.1), E_h, v_h, \gamma_{E,a}, T_h\}$ to the set \mathcal{M}_i . Let us retain the following particular cases $\mathcal{M}_{h,1}, ..., \mathcal{M}_{h,5}$ of $\mathcal{M}_{h,i}$:

$$\mathcal{M}_{h,1} = \{1, \mathcal{M}_1, (7.2), (8.3), \mathbf{X}_{0h}, \mathbf{u}_h, \gamma_1, T_{VPD,h}\}$$
 with

(8.3)
$$(\operatorname{grad} \mathbf{u}_h, \operatorname{grad} \mathbf{w}_h) = \langle \mathbf{f}, \mathbf{w}_h \rangle, \ \forall w_h \in \mathbf{X}_{0h}$$

 $\mathcal{M}_{h,2} = \{2, \mathcal{M}_2, (7.3), (8.4), Y_{0h}, q_h, \gamma_2, T_{PD,h}\}$ with

(8.4)
$$(\operatorname{grad} q_h, \operatorname{grad} \lambda_h) = \langle \psi, \lambda_h \rangle, \ \forall \lambda_h \in Y_{0h}.$$

 $\mathcal{M}_{h,3} = \{3, \mathcal{M}_3, (7.4), (8.5), Y_h, \phi_h, \gamma_3, B_{1,h}\}$ with

(8.5)
$$(\operatorname{grad} \phi_h, \operatorname{grad} \mu_h) + (\phi_h, \mu_h) = \langle \psi_1, \mu_h \rangle, \ \forall \mu_h \in Y_h.$$

 $\mathcal{M}_{h,4} = \{4, \mathcal{M}_4, (7.4), (8.5), M_h, \phi_h, \gamma_4, B_{1,h}\}.$

Let π_h be the L^2 projection operator onto \mathbf{X}_h defined by $\pi_h \in L(\mathbf{L}^2(\Omega), \mathbf{X}_h)$ such that $\mathbf{z}_h \in \mathbf{X}_h$, $\mathbf{f}_* \in \mathbf{L}^2(\Omega)$, $\pi_h \mathbf{f}_* = \mathbf{z}_h$ if and only if

(8.6)
$$(\mathbf{z}_h, \mathbf{w}_h)_0 = (\mathbf{f}_*, \mathbf{w}_h)_0, \ \forall \, \mathbf{w}_h \in \mathbf{X}_h$$

 $\mathcal{M}_{h,5} = \{5, \mathcal{M}_5, (7.5), (8.6), \mathbf{X}_h, \mathbf{z}_h, \gamma_5, \pi_h\}.$

Approximate (6.1) - (6.10), where (6.7) is considered with the formulation (6.11), by the approximate extended system: find $(x_h, U_h) \in \widehat{\Gamma}_h$ such that

$$(8.7) p_{h} - (q_{h} + \hat{p}_{h}) = \hat{g},$$

$$(8.8) p_{S,h} - p_{h} = \ell_{S},$$

$$(8.9) (div \mathbf{u}_{h} - y_{h}, \hat{\mu}) = (g, \hat{\mu}), \forall \hat{\mu} \in L_{0}^{2}(\Omega),$$

$$(8.10)(grad \mathbf{u}_{h}, grad \mathbf{w}_{h}) - (p_{S,h}, div \mathbf{w}_{h}) = \langle \mathbf{f}, \mathbf{w}_{h} \rangle, \forall \mathbf{w}_{h} \in \mathbf{X}_{0h},$$

$$(8.11) - \alpha \langle \widetilde{S}_{\mathbf{z}_{h}}, \overline{\lambda}_{h} \rangle + \langle \widetilde{S}_{\mathbf{t}_{h}}, \overline{\lambda}_{h} \rangle + (grad q_{h}, grad \overline{\lambda}_{h}) = \langle \psi, \overline{\lambda}_{h} \rangle, \forall \overline{\lambda}_{h} \in Y_{0h},$$

$$(8.12)(grad \hat{q}_{h}, grad \tilde{\lambda}_{h}) + (grad p_{h}, grad \overline{\lambda}_{h})$$

$$= \langle \hat{\psi}, \widetilde{\lambda}_{h} \rangle, \forall \widetilde{\lambda}_{h} \in Y_{0h},$$

$$(8.13)(\mathbf{z}_{h}, \mathbf{\tilde{w}}_{h}) + (grad p_{h}, \mathbf{\tilde{w}}_{h}) = (\mathbf{f}_{\triangle}, \mathbf{\tilde{w}}_{h}), \forall \mathbf{\tilde{w}}_{h} \in \mathbf{X}_{h},$$

$$(8.14)(\mathbf{z}_{h} - \mathbf{t}_{h} + grad (\hat{p}_{h} - \hat{q}_{h}), grad \varphi_{h})$$

$$+ (grad r_{h}, grad \varphi_{h}) = \langle \varrho, \varphi_{h} \rangle, \forall \varphi_{h} \in Y_{h},$$

$$(8.15)(r_{h}, \tilde{\varphi}_{h}) + (grad (q_{h} + \hat{q}_{h}), grad \tilde{\varphi}_{h}) = \langle \hat{\varrho}, \tilde{\varphi}_{h} \rangle, \forall \tilde{\varphi}_{h} \in Y_{h},$$

$$(8.16)(\mathbf{t}_{h}, \mathbf{\hat{w}}_{h}) = (\mathbf{\hat{f}}, \mathbf{\hat{w}}_{h}), \forall \mathbf{\hat{w}}_{h} \in \mathbf{X}_{h},$$

Remark 8.1. Equation (8.13) approximates equation (6.11).

Let $\mathbf{f}_{\Delta,h}$ and $\mathbf{f}_h \in \mathbf{X}_h$ be defined by

$$(8.17)(\mathbf{f}_{\triangle,h},\tilde{\mathbf{w}}_h) = (\mathbf{f}_{\triangle},\tilde{\mathbf{w}}_h), \ \forall \,\tilde{\mathbf{w}}_h \in \mathbf{X}_h, \ (\hat{\mathbf{f}}_h,\tilde{\mathbf{w}}_h) = (\hat{\mathbf{f}},\hat{\mathbf{w}}_h), \ \forall \,\hat{\mathbf{w}}_h \in \mathbf{X}_h.$$

Using $\mathbf{f}_{\Delta,h}$, we work with (8.13) as we work with (6.12).

Using (8.17), equations (8.13) and (8.16) are equivalent, respectively, to

(8.18)
$$\mathbf{z}_h + \operatorname{grad} p_h = \mathbf{f}_{\triangle,h}, \ \mathbf{t}_h = \mathbf{f}_h$$

Replacing (8.7) and (8.18) in (8.14), it results

(8.19)
$$(\mathbf{f}_{\Delta,h} - \mathbf{f}_h - \operatorname{grad} \hat{g}, \operatorname{grad} \varphi_h) - (\operatorname{grad} (q_h + \hat{q}_h), \operatorname{grad} \varphi_h) + (\operatorname{grad} r_h, \operatorname{grad} \varphi_h) = \langle \varrho, \varphi_h \rangle, \ \forall \varphi_h \in Y_h,$$

Let $\tilde{\psi}_{r,h} \in Y'_h$ be defined by

(8.20)
$$\langle \hat{\psi}_{r,h}, \varphi_h \rangle = (-\mathbf{f}_{\triangle,h} + \hat{\mathbf{f}}_h + \operatorname{grad} \hat{g}, \operatorname{grad} \varphi_h) + \langle \varrho, \varphi_h \rangle + \langle \hat{\varrho}, \varphi_h \rangle, \ \forall \varphi_h \in Y_h,$$

Summing (8.19) and (8.15), we obtain that r_h is the solution of the equation

(8.21)
$$((r_h, \varphi_h))_1 = \langle \bar{\psi}_{r,h}, \varphi_h \rangle, \ \forall \varphi_h \in Y_h ,$$

From (8.18), we deduce

(8.22)
$$(\mathbf{z}_h - \mathbf{t}_h + \operatorname{grad} p_h, \operatorname{grad} \varphi_h) = (\mathbf{f}_{\Delta,h} - \hat{\mathbf{f}}_h, \operatorname{grad} \varphi_h), \ \forall \varphi_h \in Y_h,$$

Equation (8.22) approximates (6.20). The same is obtained if we replace (8.7), (8.16) in (8.14) and then sum (8.15) to (8.14), where we use (8.21).

Corollary 8.2. We have

(8.23)
$$(\alpha \boldsymbol{z}_{h} - \boldsymbol{t}_{h}, \operatorname{grad} \bar{\lambda}_{h}) + (\operatorname{grad} p_{h}, \operatorname{grad} \bar{\lambda}_{h}) = (r_{h}, \bar{\lambda}_{h}) + \langle \psi, \bar{\lambda}_{h} \rangle + \langle \hat{\psi}, \bar{\lambda}_{h} \rangle - \langle \hat{\varrho}, \bar{\lambda}_{h} \rangle, \ \forall \, \bar{\lambda}_{h} \in Y_{0h},$$

where we use (8.21). This equation approximates (6.42).

(8.24)
$$(\boldsymbol{z}_h - \boldsymbol{t}_h + \operatorname{grad} p_h, \operatorname{grad} \varphi_h)$$
$$= (\boldsymbol{f}_{\triangle,h} - \hat{\boldsymbol{f}}_h, \operatorname{grad} \varphi_h), \ \forall \, \varphi_h \,\in \, Y_h \ominus Y_{0h} \,,$$

This equation approximates (6.43).

Proof. If we add (8.11) and (8.12) and subtract (8.15), where the test functions $\varphi_h \in Y_{0h}$, we have (8.23), where we use (8.21). (8.22) gives (8.24).

Remark 8.3. As in the exact case, in (6.13), we obtain

(8.25)
$$(\operatorname{grad} \mathbf{u}_h, \operatorname{grad} \mathbf{w}_h) - (\mathbf{z}_h, \mathbf{w}_h) = (\ell_S, \operatorname{div} \mathbf{w}_h) + \langle \mathbf{f}, \mathbf{w}_h \rangle - (\mathbf{f}_{\triangle}, \mathbf{w}_h), \ \forall \, \mathbf{w}_h \in \mathbf{X}_{0h},$$

Approximate (7.6) - (7.15) by the problem: find $(x_h, U_h) \in \widehat{\Gamma}_h$ such that

 $p_h - (q_h + \hat{p}_h) = \hat{g} \,,$ (8.26) $p_{S,h} - p_h = \ell_S \,,$ (8.27) $div\,\mathbf{u}_h - y_h = g\,,$ (8.28) $\mathbf{u}_h - T_{VPD,h} G_{\mathbf{u}}(p_{S,h}) = T_{VPD,h} \mathbf{f},$ (8.29) $q_h - T_{PD,h}G_q(\mathbf{t}_h, \mathbf{z}_h) = T_{PD,h}\psi,$ (8.30) $\hat{q}_h - T_{PD,h} G_{\hat{q}}(p_h) = T_{PD,h} \hat{\psi} \,,$ (8.31) $\mathbf{z}_h - \pi_h G_{\mathbf{z}}(p_h) = \pi_h \mathbf{f}_{\Delta} \,,$ (8.32) $p_h - B_{1,h}G_p(p_h, \hat{p}_h, \hat{q}_h, \mathbf{z}_h, \mathbf{t}_h, r_h) = B_{1,h}\varrho,$ (8.33) $r_h - B_{1,h}G_r(q_h, \hat{q}_h, r_h) = B_{1,h}\hat{\varrho},$ (8.34) $\mathbf{t}_h - \pi_h G_{\mathbf{t}}(\mathbf{u}_h) = \pi_h \hat{\mathbf{f}},$ (8.35)

Lemma 8.4. Problem (8.26) - (8.35) is equivalent to problem (8.7) - (8.16).

Proof. The conclusion follows from the definition of operators $T_{VPD,h}$, $T_{PD,h}$, π_h , $B_{1,h}$.

Lemma 8.5. Problem (8.26) - (8.35) has the form of equation (4.3), so Theorem 4.2, Corollary 4.4, Corollary 4.5 can be used in order to analyze (8.26) - (8.35).

Proof. Consider the spaces Ω , \mathfrak{X} , \mathfrak{Y} , \mathfrak{Z} and \mathfrak{X}_h defined in Section 5. The operator $\widehat{\mathcal{A}}_h$ defined by the left hand side of (8.26) - (8.35) has the form of the operator $\widehat{\mathcal{A}}_h$ defined in Section 4.

Theorem 8.6. Assume the hypotheses of Theorem 4.2. Taking $\alpha \in \mathbb{R}$, $\alpha \neq 1$, we obtain a family of well - posed approximate formulations (8.7) - (8.16).

Proof. The results follow from Lemma 8.5, Theorem 4.2 and Lemma 8.4. \Box

Theorem 8.7. Assume the hypotheses of Corollaries 6.11 and 6.12. Assume the hypotheses of Theorem 4.2. For each $\alpha \neq 1$, the solution (\boldsymbol{u}, p) of problem (5.1) - (5.2) is approximated by the components (\boldsymbol{u}_h, p_h) of the solution of (8.7) - (8.16).

Proof. This follows from Theorem 8.6.

8.2. Approximation by the finite element method. In order to define the following operator π_{0h} and the discrete Laplace operator $-\triangle^h$, assume that the approximate spaces are finite element spaces and the following inverse inequality [5, 6, 20, 34] is satisfied

(8.36)
$$\|\mathbf{u}_h\|_1 \leq C_{FEM} h^{-1} \|\mathbf{u}_h\|_0, \ \forall \, \mathbf{u}_h \in \mathbf{X}_h,$$

We refer to [5]. Assume (8.36). Let π_{0h} be the L^2 projection operator onto \mathbf{X}_{0h} defined by $\pi_{0h} : \mathbf{L}^2(\Omega) \to \mathbf{X}_{0h}$ such that $\mathbf{z}_h \in \mathbf{X}_{0h}$, $\mathbf{f}_* \in \mathbf{L}^2(\Omega)$, $\pi_{0h}\mathbf{f}_* = \mathbf{z}_h$ if and only if

(8.37)
$$(\mathbf{z}_h, \mathbf{w}_h) = (\mathbf{f}_*, \mathbf{w}_h), \ \forall \, \mathbf{w}_h \in \mathbf{X}_{0h},$$

 $\mathcal{M}_{h,6} = \{6, \mathcal{M}_5, (7.5), (8.37), \mathbf{X}_{0h}, \mathbf{z}_h, \gamma_6, \pi_{0h}\}.$

We refer to [5]. Assume (8.36). The discrete Laplace operator $-\Delta^h$ is defined to be the mapping $-\Delta^h : \mathbf{H}_0^1(\Omega) \to \mathbf{X}_{0h}$ be such that $-\Delta^h \mathbf{u} = \xi_h$ if and only if

(8.38)
$$(\xi_h, \mathbf{w}_h) = (grad \, \mathbf{u}, grad \, \mathbf{w}_h), \ \forall \, \mathbf{w}_h \in \mathbf{X}_{0h},$$

Remark 8.8. Assume that Ω is a polygon in \mathbb{R}^2 . N = 2. Let us consider a regular triangulation Θ_h of Ω by triangles K that are triangles with three nodal points that coincide with the three vertices.

(8.39)
$$X_h = \{v_h \in C^0(\overline{\Omega}), v_h | _K \in P_1(K), \forall K \in \Theta_h\}.$$

We take $X_h = Y_h$. A basis functions of the space $P_1(K)$ is defined by the three nodal points of K (e.g. subsection 10.2.1, page 394, [2]). The inverse inequality (8.36) holds ([5, 6, 20, 34]). Then, in equation (8.24), we use $Y_h \oplus Y_{0h} = \Xi_h$. The space Ξ_h denotes the complementary space M_h from the subsection 5.7.2.2, page 268, [35]. From this reference, we also retain that $\mu_h \in \Xi_h$ is determined by its values at the boundary nodes.

Corollary 8.9. The approximation p_h of the pressure p on the boundary depends on the approximation \mathbf{z}_h of the \mathbf{z} on the boundary. In order to obtain the approximate solution (\mathbf{u}_h, p_h) as a component of the solution of (8.10), (8.23), (8.24), where we use $(8.16), (8.21), \mathbf{z}_h$ can be determined from equation (8.25) under some additional conditions which are decided in every studied particular case of approximation. One such condition is $\mathbf{z}_h \in \mathbf{X}_{0h}$ in the case of the finite element method, under the hypothesis (8.36), that is, $\mathbf{z}_h = \xi_h$ from (8.38), where we use the approximation $-\Delta^h$: $\mathbf{H}_0^1(\Omega) \to \mathbf{X}_{0h}$ of the Laplace operator from [5].

Proof. The first affirmation follows from (8.24). $\psi_{r,h} = 0$ so $r_h = 0$ from (8.21). If we determine u_h , p_h , \mathbf{z}_h , \mathbf{t}_h as it is indicated in the Corollary 8.9, the rest of the components of the solution (x_h, U_h) are obtained from the rest of the equations of (8.7) - (8.16). u_h , p_h , \mathbf{z}_h , \mathbf{t}_h exist as components of the solution (x_h, U_h) of (8.7) - (8.16) or (8.26) - (8.35). If we assume that they are not unique, then we contradict the uniqueness of (x_h, U_h) .

Corollary 8.10. Let us continue the discussion from Corollary 8.9 in the case of the finite element method, under the hypothesis (8.36).

We refer as to "the basic case" to the exact and approximated problems studied till now, (6.1) - (6.10) (where we work with (6.11) instead of (6.7)), (7.6) - (7.15), (8.7) - (8.16), (8.26) - (8.35).

I. A modality to introduce the condition $z_h \in X_{0h}$ is obtained by taking two steps:

1. The basic case. For the study, we use Theorem 9.5 (or similar to this).

2. The approximate problem (8.7) - (8.16) from the basic case is approximated by a problem of the same form, with the same data, with a single modification: $\mathbf{z}_h \in \mathbf{X}_h$ from the basic case is approximated by the new \mathbf{z}_h $\in \mathbf{X}_{0h}$; we replace (8.13), (8.32) by (8.40), (8.41), respectively, where

(8.40) $(\boldsymbol{z}_h, \tilde{\boldsymbol{w}}_h) - (p_h, div \, \tilde{\boldsymbol{w}}_h) = (\boldsymbol{f}_{\triangle}, \tilde{\boldsymbol{w}}_h), \, \forall \, \tilde{\boldsymbol{w}}_h \in \boldsymbol{X}_{0h},$

For this second step of the study, for a fixed h, we use only Theorem 4.2, without the convergence $h \rightarrow 0$. In order to verify the estimate (4.4), it is necessary only the use of (9.11) (or similar to this) in order to compare (8.32), from the basic case, with (8.32) from this second step.

Finally, for these two steps, the estimate (4.4) must be satisfied comparing (7.6) - (7.15) with (8.26) - (8.35), where (8.13), (8.32) are replaced by (8.40), (8.41) (and where $\mathbf{z}_h \in \mathbf{X}_{0h}$). The convergence $h \to 0$ from the first step and (9.11) (or similar to this) in the second step lead us to the existence of an index h_2 such that $\forall h < h_2, q_* < 1$, that is, (4.4).

II. Another modality to introduce the condition $\mathbf{z}_h \in \mathbf{X}_{0h}$ is obtained in the following manner. In the basic case, we approximate (6.7) by (8.40) that replaces (8.13) in (8.7) - (8.16), and, in (8.26) - (8.35), we replace (8.32) by (8.41). For the study, it can be used Theorem 4.2 and estimates such those from Lemmas 9.2 and 9.3 (or similar to these), accompanied to laborious correlations between spaces, norms and the components of \mathcal{G} .

Corollary 8.11. Assume the hypotheses of Corollary 8.10. Then, the problem has the following matrix form

(8.42)
$$\begin{bmatrix} A_h & 0 & B_h^T \\ A_h & K_h & 0 \\ 0 & \begin{bmatrix} \alpha C_h \\ C_{h,0} \end{bmatrix} & D_h \end{bmatrix} \begin{bmatrix} \bar{\boldsymbol{u}}_h \\ \bar{\boldsymbol{z}}_h \\ \bar{p}_h \end{bmatrix} = \begin{bmatrix} \bar{\boldsymbol{f}} \\ \bar{\boldsymbol{0}} \\ \bar{\boldsymbol{\psi}} \end{bmatrix} ,$$

with three block Poisson preconditioners A_h , K_h and D_h . $(\bar{u}_h, \bar{z}_h, \bar{p}_h)$ is the coefficient vector corresponding to the finite element solution (u_h, z_h, p_h) .

Proof. We use (8.10), (8.23), (8.24), (8.25) and some bases of X_h and Y_h .

Corollary 8.12. Assume that $\bar{\mathbf{z}}_h$ is evaluated in terms of $\bar{\mathbf{u}}_h$, $\bar{\mathbf{z}}_h = -K_h^{-1}A\bar{\mathbf{u}}_h$. We denote $G_h(\alpha) = -\begin{bmatrix} \alpha C_h \\ C_{h,0} \end{bmatrix} K_h^{-1}A_h$. Then, the problem has the formulation (1.1) with two block Poisson preconditioners A_h and D_h .

Proof. This comes from (8.42).

Corollary 8.13. Assume the hypotheses of Corollaries 6.11, 6.12 and 8.10. It is expected that $(\mathbf{z}_h, \operatorname{grad} \overline{\lambda}_h) \neq 0, \forall \overline{\lambda}_h \in Y_{0h}$. Then $(\mathbf{u}_{h,1}, \mathbf{z}_{h,1}, p_{h,1}) \neq (\mathbf{u}_{h,2}, \mathbf{z}_{h,2}, p_{h,2})$, for $\alpha_1 \neq \alpha_2$, where $(\mathbf{u}_{h,i}, \mathbf{z}_{h,i}, p_{h,i})$ is the solution of (8.10), (8.25), (8.23), (8.24) for $\alpha = \alpha_i$.

30

Proof. We obtain

 $(grad (\mathbf{u}_{h,1} - \mathbf{u}_{h,2}), grad \mathbf{w}_h) - ((p_{h,1} - p_{h,2}), div \mathbf{w}_h) = 0, \ \forall \mathbf{w}_h \in \mathbf{X}_{0h},$ $(grad (\mathbf{u}_{h,1} - \mathbf{u}_{h,2}), grad \widetilde{\mathbf{w}}_h) - ((\mathbf{z}_{h,1} - \mathbf{z}_{h,2}), \widetilde{\mathbf{w}}_h) = 0, \ \forall \widetilde{\mathbf{w}}_h \in \mathbf{X}_{0h},$ $\alpha_1(\mathbf{z}_{h,1} - \mathbf{z}_{h,2}, grad \overline{\lambda}_h) + (grad (p_{h,1} - p_{h,2}), grad \overline{\lambda}_h)$ $= (\alpha_2 - \alpha_1)(\mathbf{z}_{h,2}, grad \overline{\lambda}_h), \ \forall \overline{\lambda}_h \in Y_{0h},$ $((\mathbf{z}_{h,1} - \mathbf{z}_{h,2}) + grad (p_{h,1} - p_{h,2}), grad \varphi_h) = 0, \ \forall \varphi_h \in Y_h \ominus Y_{0h},$

From Theorem 8.6, we deduce that $(\mathbf{u}_{h,1}, \mathbf{z}_{h,1}, p_{h,1}) = (\mathbf{u}_{h,2}, \mathbf{z}_{h,2}, p_{h,2})$ if and only if $(\mathbf{z}_{h,2}, grad \bar{\lambda}_h) = 0, \forall \bar{\lambda}_h \in Y_{0h}$.

In the situation considered in Corollary 8.10, it is expected that $(\mathbf{z}_h, grad \,\overline{\lambda}_h) = -(div \, \mathbf{z}_h, \overline{\lambda}_h) \neq 0.$

Remark 8.14. Assume the hypotheses of Corollary 8.10. We can simplify the problem by replacing $(grad \mathbf{u}_h, grad \mathbf{w}_h)$ from (8.25) by $(\mathbf{z}_h, \mathbf{w}_h)$ in (8.10). In other words, we replace (8.10), (8.25) by (8.13). The system is given by (8.13), (8.16), (8.23), (8.24) with the unknowns \mathbf{z}_h , p_h , \mathbf{t}_h . \mathbf{u}_h results from (8.25).

9. An analysis using the finite element method

Assume that Ω is a polygon in \mathbb{R}^2 or a polyhedron in \mathbb{R}^3 . Let X_h be a finite dimensional subspace of $H^1(\Omega)$. h is the mesh parameter.

We take $X_h = Y_h$. In the case of Stokes problem, if the approximation space of the velocity is the same as the approximation space, then the discrete inf-sup (or the LBB) condition is not satisfied (for instance, [4, 5, 6]).

Taking a regular triangulation Θ_h of Ω by triangles K, define

(9.1)
$$X_h = \{ v_h \in C^0(\overline{\Omega}), v_h | _K \in \mathbb{P}_{k_Q}(K), \forall K \in \Theta_h \}, \quad k_Q \ge 1 ,$$

where $\mathbb{P}_k(K)$ is the space of polynomials in the variables x_1, \ldots, x_N , of degree less than or equal to k, defined on $K, k \geq 1$. Let us assume the following hypotheses: there exist some operators $I_1 \in L(H^1(\Omega), X_h)$, $I_0 \in L(L^2(\Omega), X_h)$, some integers j_1, j_0 , and some positive constants δ_1, δ_0 , independent of h, such that

(9.2)
$$\|\mu - I_1\mu\|_1 \le \delta_1 \cdot h^m \cdot |\mu|_{m+1}, \ \forall \mu \in H^{m+1}(\Omega), \ 1 \le m \le j_1.$$

(9.3)
$$\|\mu - I_0\mu\|_0 \le \delta_0 \cdot h^{m+1} \cdot |\mu|_{m+1}, \ \forall \mu \in H^{m+1}(\Omega), \ 1 \le m \le j_0.$$

Lemma 9.1. Let us consider $\mathcal{M}_{h,i}$, $i = 1, \ldots, 5$. Assume Hypotheses (9.2) - (9.3). Let $\mathcal{E}_1 = (C^{\infty}(\overline{\Omega}) \cap H_0^1)^N$, $\mathcal{E}_2 = C^{\infty}(\overline{\Omega}) \cap H_0^1$, $\mathcal{E}_3 = C^{\infty}(\overline{\Omega})$, $\mathcal{E}_4 = C^{\infty}(\overline{\Omega}) \cap L_0^2(\Omega)$, $\mathcal{E}_5 = C_0^{\infty}(\Omega)^N$. Then, i) \mathcal{E}_i is dense in E_i and

(9.4)
$$\lim_{h \to 0} \inf_{v_h \in E_{h,i}} \|v - v_h\|_{E_i} = 0, \ \forall v \in \mathcal{E}_i,$$

ii)

(9.5)
$$||v_i - v_{h,i}||_{E_i} \to 0 \text{ as } h \to 0$$

Proof. The proof follows from [2, 10, 20, 26, 31, 46].

Lemma 9.2. Assume that the hypotheses of Lemma 9.1. Let $m \ge 1$. If $\mathbf{u} \in \mathbf{H}_0^1(\Omega) \cap \mathbf{H}^{m+1}(\Omega)$, $q \in H_0^1(\Omega) \cap H^{m+1}(\Omega)$, $\phi \in H^{m+1}(\Omega)$, $\mathbf{z} \in \mathbf{H}^{m+1}(\Omega)$, then there exist some constants $C_i > 0$, $i = 1, \ldots, 4$, such that

(9.6)
$$|(T_{VPD} - T_{VPD,h})T_{VPD}^{-1}\boldsymbol{u}|_1 \le C_1 \cdot h^m \cdot |\boldsymbol{u}|_{m+1}.$$

(9.7)
$$|(T_{PD} - T_{PD,h})T_{PD}^{-1}q|_1 \le C_2 \cdot h^m \cdot |q|_{m+1}.$$

- (9.8) $\| (B_1 B_{1,h}) B_1^{-1} \phi \|_1 \le C_3 \cdot h^m \cdot \| \phi \|_{m+1} .$
- (9.9) $\|(\pi \pi_h)\pi^{-1} \mathbf{z}\|_0 \le C_4 \cdot h^{m+1} \cdot \|\mathbf{z}\|_{m+1}.$

Proof. These are obtained by standard methods ([2, 10, 20, 26, 31, 46]). We also use $\mathbf{f} = T_{VPD}^{-1}\mathbf{u}, \ \psi = T_{PD}^{-1}q, \ \psi_1 = B_1^{-1}\phi, \ \mathbf{f}_* = \pi^{-1}\mathbf{z}$ and $\pi_h\pi = \pi_h$.

Lemma 9.3. ([5], Theorem 5.1, k = 1) For π_{0h} from (8.37) and for any $u \in L^2(\Omega)$, we retain for k = 1 in Theorem 5.1, [5], the following facts:

(9.10)
$$\|(I - \pi_{0h})\boldsymbol{u}\|_{-1} = \sup_{\phi \in \boldsymbol{H}_{0}^{1}(\Omega)} \frac{((I - \pi_{0h})\boldsymbol{u}, \phi)}{\|\phi\|_{1}}$$

(9.11)
$$||(I - \pi_{0h})\boldsymbol{u}||_{-1} \leq Ch ||\boldsymbol{u}||_{0}.$$

In order to verify (4.16) and (4.17), let us remember:

Lemma 9.4. ([49]) Consider two sets $A, B \subseteq \mathbb{R}$. Then,

(9.12)
$$\inf(A+B) = \inf A + \inf B,$$

Theorem 9.5. i) Consider the spaces Q, X, Y, Z and X_h defined in Section 5. In the definition of X_h , the spaces X_h and Y_h are those from the present Section. Let \widehat{A}_h be the operator defined by (8.26) - (8.35).

Then, the conclusions of Theorem 4.2 and of Corollary 4.4 hold for (8.26) - (8.35).

ii) Assume that Ω is a bounded convex polygon in \mathbb{R}^2 or a polyhedron in \mathbb{R}^3 . Let $m \geq 1$ and $\mathbf{u} \in \mathbf{H}_0^1(\Omega) \cap \mathbf{H}^{m+1}(\Omega)$, $q, \hat{q} \in H_0^1(\Omega) \cap \mathbf{H}^{m+1}(\Omega)$, \mathbf{z} , $\mathbf{t} \in \mathbf{H}^{m+1}(\Omega)$, $p, r \in H^{m+1}(\Omega)$. Let $q_{*,1}$ be q_* for some fixed $h = h_1 \leq h_0$ such that $q_* \leq q_{*,1}, \forall h < h_1$. Then, there exists a constant $C_0 > 0$ such that we have the estimate

(9.13)
$$\left\| \begin{bmatrix} x \\ U \end{bmatrix} - \begin{bmatrix} x_h \\ U_h \end{bmatrix} \right\|_{\widehat{\Gamma}} \le (1 - q_{*,1})^{-1} \|\widehat{\mathcal{A}}^{-1}\|_{L(\widehat{\Delta},\widehat{\Gamma})} C_0 \cdot h^m \cdot |||U|||,$$

where $|||U||| = |\mathbf{u}|_{m+1} + |q|_{m+1} + |\hat{q}|_{m+1} + ||\mathbf{z}||_{m+1} + ||p||_{m+1} + ||r||_{m+1} + ||\mathbf{t}||_{m+1}$.

Proof. We use Theorem 4.2, Corollary 4.4, Corollary 4.5.

i) $S = \mathcal{E}_1 \times \mathcal{E}_2 \times \mathcal{E}_2 \times \mathcal{E}_4 \times \mathcal{E}_5 \times \mathcal{E}_3 \times \mathcal{E}_4$ and it is also given in Section 5.

Using an enumeration different from that for \mathcal{E}_k and for $\mathcal{M}_{h,k}$, let us write $V = (V_1, \ldots, V_7) \in \mathfrak{X} = \mathfrak{X}_1 \times \ldots \times \mathfrak{X}_7$, $\mathfrak{S} = \mathfrak{S}_1 \times \ldots \times \mathfrak{S}_7$, $\mathfrak{X}_h = \mathfrak{X}_{h,1} \times \ldots \times \mathfrak{X}_{h,7}$, $\mathfrak{T} = (\mathfrak{T}_1, \ldots, \mathfrak{T}_7)$, $\mathfrak{T}_h = (\mathfrak{T}_{h,1}, \ldots, \mathfrak{T}_{h,7})$, and so on.

 $||V||_{\mathcal{X}} = ||V_1||_{\mathcal{X}_1} + \ldots + ||V_7||_{\mathcal{X}_7}.$

Let us define the following sets of real numbers: $A_j = \{ \|V_j - V_{h,j}\|_{\mathfrak{X}_j} | V_{h,j} \in \mathfrak{X}_{h,j} \}, A = \{ \|V - V_h\|_{\mathfrak{X}} | V_h \in \mathfrak{X}_h \}.$

We have $A = \{ \|V_1 - V_{h,1}\|_{X_1} + \ldots + \|V_n - V_{h,7}\|_{X_7} | V_{h,1} \in \mathcal{X}_{h,1}, \ldots, V_{h,7} \in \mathcal{X}_{h,7} \}$, so $A = A_1 + \ldots + A_5$ and $\inf(A_1 + \ldots + A_5) = \inf A_1 + \ldots + \inf A_5$ (we use (9.12)), that is, $\inf_{V_h \in \mathcal{X}_h} \|V - V_h\|_{X} = \sum_{j=1}^7 \inf_{V_{h,j} \in \mathcal{X}_{h,j}} \|V_j - V_{h,j}\|_{X_j}$. Let us verify (4.16). $\forall V \in \mathcal{S}$, we have: $\lim_{h \to 0} \inf_{V_h \in \mathcal{X}_h} \|V - V_h\|_{X} = \sum_{j=1}^7 \lim_{h \to 0} \inf_{V_{h,j} \in \mathcal{X}_{h,j}} \|V_j - V_{h,j}\|_{X_j}$ Using Lemma 9.1 i), it results (4.16). Let us verify (4.17). We denote $\gamma_0 = \sup_{i \in \{1,\ldots,7\}} \gamma_i$. Let us use (8.2). $\|W - W_h\|_{X} = \sum_{j=1}^7 \|W_j - W_{h,j}\|_{X_j} \le \sum_{j=1}^7 \gamma_j \inf_{V_{h,j} \in \mathcal{X}_{h,j}} \|W_j - V_{h,j}\|_{X_j} \le \gamma_0 \sum_{j=1}^7 \inf_{V_{h,j} \in \mathcal{X}_{h,j}} \|W_j - V_{h,j}\|_{X_j} = \gamma_0 \inf_{V_h \in \mathcal{X}_h} \|W - V_h\|_{X}$, so (4.17). ii) We have (4.10). So we can take h_1 and $q_{*,1}$. $q_* < q_{*,1}$ implies $(1-q_*)^{-1} < \psi h < h_1$, let us use (4.6) and the estimates from Lemma 9.2. We denote

 $C_0 = \sup\{C_1, C_2, C_3, C_4h_1\}.$

We have $\|(\mathcal{T}-\mathcal{T}_{h})\mathcal{T}^{-1}U\|_{\mathcal{X}} = \sum_{j=1}^{7} \|(\mathcal{T}_{j}-\mathcal{T}_{h,j})\mathcal{T}_{j}^{-1}U_{j}\|_{\mathcal{X}_{j}} \leq C_{1}\cdot h^{m}\cdot |\mathbf{u}|_{m+1} + C_{2}\cdot h^{m}\cdot |q|_{m+1} + C_{2}\cdot h^{m}\cdot |\hat{q}|_{m+1} + C_{4}\cdot h^{m+1}\cdot \|\mathbf{z}\|_{m+1} + C_{3}\cdot h^{m}\cdot \|p\|_{m+1} + C_{3}\cdot h^{m}\cdot \|r\|_{m+1} + C_{4}\cdot h^{m+1}\cdot \|\mathbf{t}\|_{m+1} < C_{0}\cdot h^{m}\cdot (|\mathbf{u}|_{m+1} + |q|_{m+1} + |\hat{q}|_{m+1} + \|\hat{q}|_{m+1} + \|\mathbf{z}\|_{m+1} + \|\mathbf{z}\|_{m+1}$

Acknowledgments

The author owes gratitude to Mr. Costache Bichir and Mrs. Georgeta Bichir for their moral and financial support without which this paper would not have existed.

Many thanks to Mrs. Georgeta Stoenescu for her advice with the English language.

References

- [1] R. A. Adams, Sobolev Spaces, Academic Press, London, 1978.
- [2] K. Atkinson, W. Han, *Theoretical Numerical Analysis*, Springer, Dordrecht, Heidelberg, London, New York, 2009.
- [3] I. Babuška, Error bounds for finite element method, Numer. Math., 16 (1971), 322-333.
- [4] T. Barth, P. Bochev, M. Gunzburger, J. Shadid, A taxonomy of consistently stabilized finite element methods for the Stokes problem, SIAM J. Sci. Comput., 25 (5) (2004), 1585-1607.
- [5] P. Bochev, M. Gunzburger An absolutely stable pressure-Poisson stabilized finite element method for the Stokes equations, SIAM J. Numer. Anal., 42(3) (2004), 1189-1207.
- [6] P. B. Bochev, M. D. Gunzburger, *Least-Squares Finite Element Methods*, Springer, New York, 2009.
- [7] D. Braess, Finite Elements. Theory, Fast Solvers, and Applications in Elasticity Theory, Cambridge University Press, 2007.
- [8] J. Bramble, R. Lazarov, and J. Pasciak, A Least Squares Approach Based on a Discrete Minus One Inner Product for First Order Systems, Technical Report 94-32, Math. Sci. Institute, Cornell University, Ithaca, NY, 1994.
- [9] J. H. Bramble and J. E. Pasciak, Least-squares methods for Stokes equations based on a discrete minus one inner product, J. Comput. Appl. Math., 74 (1996), 155-173.
- [10] S. C. Brenner, L. R. Scott, The Mathematical Theory of Finite Element Methods, Springer, New York, 2008.

CĂTĂLIN - LIVIU BICHIR

- [11] H. Brezis, Functional Analysis, Sobolev Spaces and Partial Differential Equations, Springer, New York, 2010.
- [12] F. Brezzi, On the existence, uniqueness and approximation of saddle-point problems arising from lagrangian multipliers, R.A.I.R.O. Anal. Numer., 8 (1974), 129-151.
- [13] F. Brezzi, J. Douglas, Jr., Stabilized mixed methods for the Stokes problem, Numer. Math., 53 (1988), 225-235.
- [14] F. Brezzi, M. Fortin, Mixed and Hybrid Finite Element Methods, Springer, New York, 1991.
- [15] F. Brezzi, J. Rappaz, P. A. Raviart, Finite-dimensional approximation of nonlinear problems, I. Branches of nonsingular solutions, Numer. Math., 36 (1980), 1-25.
- [16] F. Brezzi, J. Rappaz, P. A. Raviart, Finite-dimensional approximation of nonlinear problems, II. Limit points, Numer. Math., 37 (1981), 1-28.
- [17] F. Brezzi, J. Rappaz, P. A. Raviart, Finite-dimensional approximation of nonlinear problems, III. Simple bifurcation points, Numer. Math., 38 (1981), 1-30.
- [18] F. Brezzi, J. Pitkäranta, On the stabilization of finite element approximations of the Stokes equations, in Efficient Solutions of Elliptic Systems, W. Hackbusch ed., Friedr. Vieweg & Sohn, Braunschweig, pp. 11-19, 1984. Paper cited in [46].
- [19] G. Christol, A. Cot, C.-M. Marle, *Topologie*, Ellipses, Paris, 1997.
- [20] P. G. Ciarlet, The Finite Element Method for Elliptic Problems, SIAM, Philadelphia, 2002.
- [21] P. G. Ciarlet, J. L. Lions (Eds.), Handbook of Numerical Analysis, vol. II, part 1, North-Holland, Amsterdam, 1997.
- [22] P. G. Ciarlet, P.-A. Raviart, A mixed finite element method for the biharmonic equation, in Mathematical Aspects of Finite Elements in Partial Differential Equations, C. de Boor ed., Academic Press, New York, pp. 125-145, 1974. Paper cited in [20, 34].
- [23] M. Crouzeix, J. Rappaz, On Numerical Approximation in Bifurcation Theory, Masson-Paris, Springer-Berlin, 1990.
- [24] C. Cuvelier, A. Segal, A.A. van Steenhoven, *Finite Element Methods and Navier-Stokes Equations*, Reidel, Amsterdam, 1986.
- [25] R.Dautray, J.-L.Lions, Mathematical Analysis and Numerical Methods for Science and Technology, 2, Functional and Variational Methods, Springer, Berlin, 1988.
- [26] R.Dautray, J.-L.Lions, Analyse mathématique et calcul numérique pour les sciences et les techniques, 6, Méthodes intégrales et numériques, Masson, Paris, 1988.
- [27] R.Dautray, J.-L.Lions, Analyse mathématique et calcul numérique pour les sciences et les techniques, 8, Évolution: semi-groupe, variationnel, Masson, Paris, 1988.
- [28] M. O. Deville, P. F. Fischer, E. H. Mund, High-Order Methods for Incompressible Fluid Flow, Cambridge University Press, 2004.
- [29] J. Douglas, J. Wang, An absolutely stabilized finite element method for the Stokes problem, Math. Comp., 52 (1989), 495-508.
- [30] H. C. Elman, D. J. Silvester, A. J. Wathen, *Finite Elements and Fast Iterative Solvers: with Applications in Incompressible Fluid Dynamics*, Oxford University Press, 2005.
- [31] A. Ern, Aide-mémoire. Éléments finis, DUNOD, Paris, 2005.
- [32] M. Fabian, P. Habala, P. Hájek, V. Montesinos, V. Zizler, Banach Space Theory. The Basis for Linear and Nonlinear Analysis, Springer, New York, 2011.
- [33] A. Georgescu, Hydrodynamic Stability Theory, Kluwer, Dordrecht, 1985.
- [34] V. Girault, P.-A. Raviart, Finite Element Methods for Navier-Stokes Equations. Theory and Algorithms, Springer, Berlin, 1986.
- [35] R. Glowinski, Numerical Methods for Nonlinear Variational Problems, Springer, New York, 1984.
- [36] R. Glowinski, O. Pironneau, On a mixed finite element approximation of the Stokes problem. I. Convergence of the approximate solution, Numer. Math., 33 (1979), 397-424.
- [37] R. Glowinski, O. Pironneau, Numerical methods for the first biharmonic equation and for the two-dimensional Stokes problem, SIAM Review, 21 (1979), 167-212.
- [38] P. M. Gresho, R. L. Sani, On the pressure boundary conditions for the incompressible Navier-Stokes equations, Int. J. Numer. Meth. Fluids, 7 (1987), 1111-1145.

- [39] M. D. Gunzburger, Finite Element Methods for Viscous Incompressible Flows, Academic, Boston, 1989.
- [40] C.Ilioi, Optimization Problems and Algorithms for Approximate Solutions, Editura Academiei Române, Bucharest, 1980 (in Romanian).
- [41] L.V.Kantorovich, G.P.Akilov, Functional Analysis, Ed. Ştiinţifică şi Enciclopedică, Bucharest, 1986 (Romanian translation from the Second Russian Edition, NAUKA, Moscow, 1977).
- [42] E. Kreyszig, Introductory functional analysis with applications, John Wiley & Sons, New York. 1978.
- [43] M. T. Nair, Linear Operator Equations. Approximation and regularization, World Scientific, Singapore, 2009.
- [44] J. Nocedal, S. J. Wright, Numerical Optimization, Springer, New York, 2006.
- [45] R.Peyret, T.D.Taylor, *Computational Methods for Fluid Flow*, Springer, Berlin, 1983.
- [46] A. Quarteroni, A. Valli, Numerical Approximation of Partial Differential Equations, Springer, Berlin, 2008.
- [47] R. L. Sani, J. Shen, O. Pironneau, P. M. Gresho, Pressure boundary condition for the time-dependent incompressible Navier - Stokes equations, Int. J. Numer. Meth. Fluids, 50 (2006), 673-682.
- [48] M. Schechter, Principles of Functional Analysis, Graduate Studies in Mathematics, 36, American Mathematical Society, Providence, Rhode Island, 2002.
- [49] Gh. Sireţchi, Mathematical Analysis, 1, Multiplication Centre of Bucharest University, 1978 (in Romanian).
- [50] Gh. Sireţchi, General Topology, Multiplication Centre of Bucharest University, 1983 (in Romanian).
- [51] Gh. Sireţchi, Functional Analysis, Multiplication Centre of Bucharest University, 1982 (in Romanian).
- [52] Gh. Sireţchi, *Concrete Spaces in Functional Analysis*, Multiplication Centre of Bucharest University, 1982 (in Romanian).
- [53] C. Taylor, T.G. Hughes, Finite Element Programming of the Navier-Stokes Equations, Pineridge Press, Swansea, U.K., 1981.
- [54] R.Temam, Navier-Stokes equations. Theory and numerical analysis, North-Holland, Amsterdam, 1979.
- [55] F. Thomasset, Implementation of Finite Element Methods for Navier-Stokes Equations, Springer, New York, 1981.
- [56] L. K. Waters, G. J. Fix and C. L. Cox, The Method of Glowinski and Pironneau for the Unsteady Stokes Problem, Comput. Math. Appl., 48 (2004), 1191-1211.
- [57] K. Yosida, Functional Analysis, Springer, Berlin, 1980.
- [58] E. Zeidler, Nonlinear Functional Analysis and its Applications II/A: Linear Monotone Operators, Springer, New York, 1990.
- [59] E. Zeidler, Nonlinear Functional Analysis and its Applications II/B: Nonlinear Monotone Operators, Springer, New York, 1990.

INDEPENDENT RESEARCHER, 74 NICOLAE BĂLCESCU STREET, FLAT V1, APARTMENT 5, GALAȚI, 800001, ROMANIA

Email address: catalinliviubichir@yahoo.com