The Sunburst Arc with JWST: III. An Abundance of Direct Chemical Abundances

BRIAN WELCH (D, 1, 2, 3) T. EMIL RIVERA-THORSEN (D, 4) JANE R. RIGBY (D, 2) TAYLOR A. HUTCHISON (D, 2)GRACE M. OLIVIER (D, 5) DANIELLE A. BERG (D, 6) KEREN SHARON (D, 7) HÅKON DAHLE (D, 8) M. RILEY OWENS (D, 9)MATTHEW B. BAYLISS (D, 9) GOURAV KHULLAR (D, 10) JOHN CHISHOLM (D, 6) MATTHEW HAYES (D, 4) AND KEUNHO J. KIM (D^{11})

¹Department of Astronomy, University of Maryland, College Park, MD 20742, USA

²Astrophysics Science Division, Code 660, NASA Goddard Space Flight Center, 8800 Greenbelt Rd., Greenbelt, MD 20771, USA

³Center for Research and Exploration in Space Science and Technology, NASA/GSFC, Greenbelt, MD 20771

⁴ The Oskar Klein Centre, Department of Astronomy, Stockholm University, AlbaNova 10691, Stockholm, Sweden

⁵Department of Physics and Astronomy and George P. and Cynthia Woods Mitchell Institute for Fundamental Physics and Astronomy, Texas A&M University, 4242 TAMU, College Station, TX 77843-4242, USA

⁶Department of Astronomy, The University of Texas at Austin, 2515 Speedway, Stop C1400, Austin, TX 78712, USA

⁷ Department of Astronomy, University of Michigan, 1085 S. University Ave, Ann Arbor, MI 48109, USA

⁸Institute of Theoretical Astrophysics, University of Oslo, P.O. Box 1029, Blindern, NO-0315 Oslo, Norway

⁹Department of Physics, University of Cincinnati, Cincinnati, OH 45221, USA

¹⁰Department of Physics and Astronomy and PITT PACC, University of Pittsburgh, Pittsburgh, PA 15260, USA ¹¹IPAC, California Institute of Technology, 1200 E. California Boulevard, Pasadena, CA 91125, USA

ABSTRACT

We measure the gas-phase abundances of the elements He, N, O, Ne, S, Ar, and Fe in the Lyman-continuum emitting region of the Sunburst Arc, a highly magnified galaxy at redshift z = 2.37. We detect the temperature-sensitive auroral lines [SII] $\lambda\lambda$ 4069, 4076, [OIII] λ 4363, [SIII] λ 6312, [OII] $\lambda\lambda$ 7320, 7330, and [NeIII] λ 3343 in a stacked spectrum of 5 multiple images of the Lyman-continuum emitter (LCE), from which we directly measure the electron temperature in the low, intermediate, and high ionization zones. We also detect the density-sensitive doublets of [OII] $\lambda\lambda$ 3727, 3729, [SII] $\lambda\lambda$ 6717, 6731, and [ArIV] $\lambda\lambda$ 4713, 4741, which constrain the density in both the low- and high-ionization gas. With these temperature and density measurements, we measure gas-phase abundances with similar rigor as studies of local galaxies. We measure a gas-phase metallicity for the LCE of $12 + \log(O/H) = 7.97 \pm 0.05$, and find an enhanced nitrogen abundance $\log(N/O) = -0.65^{+0.16}_{-0.25}$. This nitrogen abundance is consistent with enrichment from a population of Wolf-Rayet stars, additional signatures of which are reported in a companion paper. Abundances of sulfur, argon, neon, and iron are consistent with local low-metallicity HII regions and low-redshift galaxies. This study represents the most complete chemical abundance analysis of a galaxy at Cosmic Noon to date, which enables direct comparisons between local HII regions and those in the distant universe.

Keywords: Chemical abundances, Gravitational lensing, galaxies

1. INTRODUCTION

As stars shed mass in their late evolutionary stages and explode as supernovae, they recycle newly synthesized elements into the surrounding gas, enriching the nebula. Analysis of the chemical abundance patterns in

Corresponding author: Brian Welch brian.d.welch@nasa.gov

nebulae thus give insight into the evolution of stars and the star formation histories of galaxies.

Recent measurements of high redshift galaxies have found unusual abundance patterns, including anomalously high nitrogen abundances (N/O, Cameron et al. 2023; Senchyna et al. 2023; Marques-Chaves et al. 2024; Castellano et al. 2024) and carbon abundances (C/O, D'Eugenio et al. 2023). Meanwhile similar carbon abundance measurements at slightly lower redshifts (z = 4 - 6) find much lower abundances (Citro et al. 2023; Jones et al. 2023). The anomalous abundance patterns in the highest redshift galaxies may be indicative of chemical evolution via e.g. Wolf-Rayet (WR) stars or extremely metal poor stars, as the standard production pathway for nitrogen and carbon via intermediate-mass AGB stars would not have time to develop.

One of the most robust methods to determine chemical abundance patterns in galaxies is the so-called "direct" method, which directly measures electron temperature (T_e) and density from the strengths of auroral emission lines relative to bright collisionally-excited lines. These temperatures and densities are then used in conjunction with the strengths of the bright lines to calculate the abundance of each ion within the nebula (Dinerstein 1990).

The auroral lines are intrinsically faint, making this measurement difficult. Direct T_e abundances have been well studied in nearby galaxies (e.g., van Zee & Haynes 2006; Berg et al. 2015; Croxall et al. 2015, 2016; Berg et al. 2020; Rogers et al. 2021, 2022). In these local galaxies, multiple auroral lines are detected, allowing direct measurements of the electron temperature in multiple ionization zones within the nebular gas. This eliminates the need for empirical temperature relations to infer the physical conditions of different zones within the gas, thereby making the final abundance determinations more accurate. Additionally, including muliple ionization zones minimizes the impact of empiricallycalibrated ionization corrections, which account for unobserved ionization states of a given element (e.g., Izotov et al. 2006; Amayo et al. 2021). In the best case, all relevant ionization states of each element are observed, resulting in the most accurate determinations of elemental abundances.

Only a handful of auroral line detections were available for galaxies at higher redshifts $(z \sim 1-3)$ prior to the launch of JWST (Christensen et al. 2012; James et al. 2014; Bayliss et al. 2014; Ly et al. 2015; Kojima et al. 2017; Sanders et al. 2020, 2023b; Citro et al. 2023). The sensitivity and wavelength coverage of JWST (Gardner et al. 2023; Rigby et al. 2023a) have changed the landscape of direct T_e chemical abundance measurements in distant galaxies. The auroral $[OIII]\lambda 4363$ line was detected out to $z \sim 8$ for the first time in the first data release from the Early Release Observation (ERO, Pontoppidan et al. 2022) of the lensing cluster SMACS0723 (Arellano-Córdova et al. 2022; Schaerer et al. 2022; Taylor et al. 2022; Brinchmann 2023; Curti et al. 2023; Katz et al. 2023; Rhoads et al. 2023; Trump et al. 2023; Trussler et al. 2023). This line has since been clearly detected out to $z \sim 8.7$ in a field galaxy (Sanders et al. 2023a), and at z = 10.17 in a gravitationally lensed galaxy (Hsiao et al. 2024).

Each of the aforementioned studies have relied on detection of a single auroral line for their abundance Several early JWST observing promeasurements. grams have detected multiple auroral lines in galaxies at Cosmic Noon, enabling more robust studies of abundance patterns in distant galaxies. The CECILIA program (PID 2593, PI Strom, Co-PI Rudie) targeted the $[NII]\lambda 5755$, $[SIII]\lambda 6312$, and $[OII]\lambda \lambda 7320, 7330$ lines in a sample of galaxies at $z \sim 2-3$, aiming to measure multiple ionization states and thus obtain more accurate abundances. This program has so far yielded an analysis of the stacked sample (Strom et al. 2023), and measurements of all three lines in a single galaxy at $z \sim 3$ with a notably sub-solar S and Ar abundance (Rogers et al. 2024). Meanwhile the TEMPLATES ERS program (PID 1355, PI Rigby, Co-PI Viera) has detected $[OIII]\lambda 4363$, $[SIII]\lambda 6312$, and $[OII]\lambda \lambda 7320$, 7330 in a lensed galaxy at $z \sim 1.3$, finding a slightly enhanced nitrogen abundance (Welch et al. 2024). These studies measuring multiple ionization states at $z \sim 1-3$ represent a positive step towards applying the same detailed methods used in local galaxies to high redshift. However none have managed to reach the complete standard set by studies of local galaxies.

This paper presents the analysis of direct T_e abundances in the Sunburst Arc (Dahle et al. 2016), a highly magnified gravitationally lensed galaxy at redshift z =2.37 (Sharon et al. 2022; Diego et al. 2022; Pignataro et al. 2021). We focus on a single physical region of the galaxy that is leaking ionizing photons (Rivera-Thorsen et al. 2019), which we refer to as the Lyman-continuum emitter (LCE). This region is a compact star cluster (Vanzella et al. 2022) hosting a young stellar population with a light-weighted age of ~ 3.6 Myr (Chisholm et al. 2019; Rivera-Thorsen et al. 2024) and a steep UV slope $\beta \simeq -3$ (Kim et al. 2023). Several authors have found evidence of very massive stars in the LCE (Chisholm et al. 2019; Meštrić et al. 2023; Pascale et al. 2023). A companion paper presents evidence that the LCE hosts a population of Wolf-Rayet stars (Rivera-Thorsen et al. 2024).

In report the this paper, we detection of $[SII]\lambda\lambda4069, 4076,$ $[OIII]\lambda 4363,$ $[SIII]\lambda 6312,$ the $[OII]\lambda\lambda7320,7330$, and $[NeIII]\lambda3343$ auroral lines, from which we measure T_e directly in the low, intermediate, and high ionization zones of the nebula. We also measure the electron temperature in ionized hydrogen gas from the Balmer jump. We measure abundances for seven elements (He, N, O, Ne, S, Ar, Fe). For three of these (O, S, Ar) we measure multiple ionization states, which minimizes the impact of ionization corrections in our abundance determinations. With this

set of diagnostics, for the first time we apply the same level of rigor used in local galaxies to an object at Cosmic Noon, bridging the gap between local abundance measurements and those at high redshift.

This paper is organized as follows. The data are presented in Section 2. Our emission line measurements and reddening corrections are described in Section 3. We describe our electron temperature, density, and chemical abundance measurements and results in Section 4, and discuss the context of these results in Section 5. We summarize our conclusions in Section 6. We assume flat Λ CDM cosmology with $\Omega_{\Lambda} = 0.7$, $\Omega_m = 0.3$, and $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

2. DATA

The Sunburst Arc was observed by JWST in Cycle 1 using NIRCam (Rieke et al. 2023) and the Integral Field Spectroscopy (IFS) mode of NIRSpec (Böker et al. 2023), taken in program GO-02555 (PI: Rivera-Thorsen). NIRCam data were taken in six broadband filters (F115W, F150W, F200W, F277W, F356W, and F444W). NIRSpec data were taken with two grating settings (G140H/F100LP, G235H/F170LP). The IFS data were taken in three pointings, shown in Figure 1 of Rivera-Thorsen et al. (2024). The observations and data reduction are described in greater detail in Rivera-Thorsen et al., in prep; we briefly summarize the relevant steps here. The NIRSpec IFS data were reduced following the methods described in Rigby et al. (2023b), using the TEMPLATES NIRSpec data reduction notebook (Rigby et al. 2024). We used the JWST data reduction pipeline version 1.11.4 (Bushouse et al. 2023), along with calibration reference files from pmap_1105. After running the main JWST pipeline, we use the baryon-sweep¹ code to remove any remaining outliers from the final data cubes (Hutchison et al. 2023, 2024).

In this work, we analyze a spatially-integrated spectrum that consists of 5 multiple images of the LCE covered by the three IFU pointings. We extract spectra of the LCE images from 5×5 spaxel apertures, using a flux-weighted sum. The spectra are normalized, no magnification correction is applied, however all results presented herein rely exclusively on flux ratios which are unaffected by magnification or normalization. The full extraction method is documented in greater detail in Rivera-Thorsen et al. (2024).

3. METHODS

3.1. Continuum Subtraction

Prior to fitting emission lines, we first remove the continuum. We mask known emission lines and sigma-clip the remaining continuum to remove any artifacts that passed our previous artifact removal steps described in Section 2. The remaining continuum is then smoothed with a running median to produce a smooth continuum model. The smooth model is subtracted from the original spectrum prior to fitting any emission lines. This method of continuum subtraction will not account for features such as Balmer absorption. However, we do not expect significant Balmer absorption in the young stellar population seen in the LCE (≤ 4 Myr, Chisholm et al. 2019; Rivera-Thorsen et al. 2024), and visual inspection of the original spectrum shows no sign of Balmer absorption features.

3.2. Emission Line Measurements

Emission lines are fit using Gaussian profiles, utilizing the publicly-available tools developed by the TEM-PLATES team and available on GitHub.² We initially attempt to fit each line with a single Gaussian component, and iteratively add up to two additional components if visual inspection determines the fit to be poor. We find that the brightest lines (e.g. $[OIII]\lambda\lambda4959, 5007,$ $H\alpha$) require three components - one narrow (unresolved) component, one medium component with a full-width at half maximum (FWHM) of $\sim 200-300 \text{ km s}^{-1}$, and one broad component with a FWHM of $\sim 600 - 700 \text{ km s}^{-1}$. As line strengths decrease, the broadest component is the first to become lost in the noise. Thus for many lines, only the unresolved and the medium width components are used in fitting. For the faintest lines (including all of the auroral lines), only a single unresolved component is sufficient to obtain a good fit. When fitting both singleand multi-component lines, we allow the centroids of each component to vary within ± 2 Å in the observed frame to allow for possible velocity offsets from our assumed systemic redshift ($z = 2.371062 \pm 6 \times 10^{-6}$ Rivera-Thorsen et al. 2024). For bright multi-component lines, the amplitude and width of the Gaussians are left unconstrained. For the faintest emission features, we choose to fix the line widths to the instrumental resolution at the observed wavelength, as is observed in the narrow components of the brighter emission lines. This prevents noise around the edge of the lines from artificially broadening the Gaussian fits. Final line fluxes are obtained by integrating the Gaussian fits. For line fits with two components, we sum the Gaussian integrals to obtain the total line flux. For three compo-

 2 github.com/JWST-Templates




Figure 1. Stacked spectrum of 5 multiple images of the Lyman-continuum emitter in the Sunburst Arc, zoomed in to highlight detected auroral emission lines. The extracted spectrum is shown in blue, and the continuum fit is shown in orange. Flux errors are shown in grey, offset by the continuum fit to show the SNR of the continuum-subtracted lines. The [ArIV] $\lambda\lambda$ 4713, 4741 lines abut the blue Wolf-Rayet bump (Rivera-Thorsen et al. 2024). Our continuum fit treats this broad feature similar to the continuum, so it is removed prior to fitting the argon emission lines.

nent fits, we choose to not include the broadest component in our total flux measurements, as the velocities ($\sim 600 - 700 \text{ km s}^{-1}$) are such that this gas has likely traveled far from the main nebula. This only affects the H α and [OII] $\lambda\lambda$ 3727, 3729 lines. Uncertainties in measured line fluxes are propagated through from fitting uncertainties on the Gaussian amplitude and width.

Blended emission lines (e.g., $H\alpha$ and $[NII]\lambda\lambda 6548, 6584$) are fit simultaneously, using the same priors and uncertainty calculations described above. One notable exception in our fitting process is the $[OII]\lambda\lambda 3727, 3729$ doublet. These lines are blended together, and because of the high signal-to-noise ratio (SNR) of this spectrum, the H13 and H14 lines of hydrogen are also blended with the $[OII]\lambda\lambda3727, 3729$ doublet. We find that the $[OII]\lambda\lambda3727, 3729$ lines require two components each to fit properly. The combination of six blended Gaussian profiles leads to a large degeneracy in model parameters, particularly within the $[OII]\lambda\lambda 3727, 3729$ doublet. We choose to fix the parameters of the H13 and H14 lines to improve the accuracy of the $[OII]\lambda\lambda 3727, 3729$ fit. The fluxes of H13 and H14 are calculated from our measured fluxes of the H11 and H12 lines, which are well separated from other nearby lines. We use a simple mean of the line fluxes predicted from H11 and H12 as our estimates for the H13 and H14 line flux. We fix the line widths of the H13 and H14 lines to be unresolved, and fix the observed wavelengths using the systemic redshift of the system. This process leads to a reliable $[OII]\lambda\lambda3727,3729$ fit, however the degeneracy between parameters in these blended lines still leads to unreasonably large flux uncertainties. To estimate the true uncertainty on the fluxes of these two lines, we run a MCMC ensemble using the python package emcee (Foreman-Mackey et al. 2013). We calculate the total flux of each line from each step of the MCMC, and use the standard deviation of the resulting distribution as our estimate of the flux uncertainty.

3.3. Extinction Correction

Emission line fluxes are first corrected for Milky Way dust reddening using the dust law of Cardelli et al. (1989). We query the Milky Way dust map of Schlafly & Finkbeiner (2011) via the NASA/IPAC Infrared Science Archive³ to obtain the ISM reddening correction. We find the Galactic $E(B - V) = 0.0812 \pm 0.003$.

We next correct for dust within the LCE itself. This correction is done using an iterative process using the $H\alpha$, $H\beta$, $H\gamma$, and $H\delta$ line strengths. We start by calculating the expected Balmer line ratios, assuming an initial electron temperature of 10^4 K, and an initial electron density of 10^3 cm⁻³. The measured line ratios are then compared to the predicted values to calculate E(B - V), assuming the dust law of Cardelli et al. (1989). We then apply the error-weighted average dust correction, weighted by the uncertainty on the Balmer line ratios, to relevant emission lines, and calculate the electron temperatures using the $[OIII]\lambda 4363$, $[OII]\lambda\lambda7320,7330$, and $[SIII]\lambda6312$ lines. The electron density is calculated using the $[SII]\lambda\lambda6717, 6731$ doublet ratio, as that is the best-measured of the densitysensitive doublets used in this study. We update the electron temperature using an error-weighted average of the three calculated temperatures and repeat the dust correction measurement. This process is continued iteratively until the change in error-weighted electron temperature is less than 15 K. The reddening that we calculated from this iterative process is $E(B-V) = 0.04 \pm 0.1$. The reddening-corrected line intensities are reported in Table 1.

As a consistency check, we also calculated the reddening correction assuming the dust law of Gordon et al. (2003), calculated for the Large Magellanic Cloud, to see if low metallicity alters the reddening result. We find $E(B-V) = 0.06 \pm 0.1$ in this case, consistent with the value calculated using Cardelli et al. (1989).

³ https://irsa.ipac.caltech.edu/applications/DUST/

Table 1. Emission Line Fluxes

$\begin{tabular}{ c c c c c c c c c c c c c c c c c c c$
[Ne III] 3969 3968.59 0.17 ± 0.02 0.18 ± 0.03 H ϵ 3971.20 0.14 ± 0.03 0.15 ± 0.03
He 3971.20 0.14 ± 0.03 0.15 ± 0.03
[S II] 4060 4060.75 0.006 ± 0.001 0.006 ± 0.001
S II] 4076 4075.79 0.002 ± 0.001 0.003 ± 0.001
Hδ 4102.89 0.24 ± 0.02 0.25 ± 0.02
H γ 4341.68 0.52 ± 0.07 0.53 ± 0.07
[Fe II] 4360 4359.59 0.010 ± 0.002 0.010 ± 0.002
[O III] 4363 4364.44 0.135 ± 0.006 0.138 ± 0.006
He I 4473 4472.70 0.04 ± 0.01 0.04 ± 0.01
[Fe III] 4658 4659.35 0.008 ± 0.002 0.008 ± 0.002
[Ar IV] 4713 4712.69 0.010 ± 0.003 0.010 ± 0.003
[Ar IV] 4741 4741.45 0.010 ± 0.002 0.010 ± 0.002
Hβ 4862.68 1.0 ± 0.1 1.0 ± 0.1
[O III] 4960 4960.30 2.2 ± 0.3 2.2 ± 0.3
[O III] 5008 5008.24 6.8 ± 0.4 6.8 ± 0.4
[N II] 5756 5756.24 $< 0.0004 < 0.0004$
He I 5878 5877.59 0.142 ± 0.007 0.139 ± 0.007
[O I] 6302 6302.05 0.026 ± 0.004 0.025 ± 0.004
[S III] 6312 6313.80 $0.010 \pm 0.001 0.010 \pm 0.001$
[N II] 6550 6549.85 $0.040 \pm 0.004 0.039 \pm 0.004$
Hα 6564.61 3.0 ± 0.3 2.9 ± 0.3
[N II] 6585 6585.28 0.113 ± 0.005 0.109 ± 0.005
He I 6679 6680.00 0.04 ± 0.02 0.04 ± 0.02
[S II] 6718 6718.29 0.040 ± 0.003 0.039 ± 0.003
[S II] 6733 6732.67 0.044 ± 0.003 0.043 ± 0.003
He I 7064.21 0.08 ± 0.01 0.08 ± 0.01
[Ar III] 7138 7137.80 0.062 ± 0.007 0.059 ± 0.006
[O II] 7322 7322.01 0.031 ± 0.002 0.030 ± 0.002
[O II] 7332 7331.68 0.023 ± 0.001 0.022 ± 0.001
[Ar III] 7753 7753.20 0.015 ± 0.001 0.014 ± 0.001
[S III] 9071 9071.10 0.151 ± 0.007 0.139 ± 0.007

NOTE—Emission line fluxes and dereddened intensities relative to H β . Columns are: (1) Line identification, (2) Vacuum wavelength, in Å, (3) Measured flux relative to H β , (4) Dereddened intensity relative to H β . Line fluxes were measured from the stacked spectrum of multiple images of the LCE, which has arbitrary units; to enable conversion to physical units, we report the measured and the demagnified flux of Hbeta for the image of the LCE that appears in image 4 of the lensed galaxy: the measured flux is $1.01 \pm 0.03 \times 10^{-17} \text{erg s}^{-1} \text{cm}^{-1}$, and the demagnified flux is $6.6 \pm 0.2 \times 10^{-19} \text{erg s}^{-1} \text{cm}^{-1}$, assuming

the calculated magnification of $\mu = 15.3$ from Sharon et al. (2022). Note this demagnified flux does not include

magnification uncertainties.

4. DIRECT ELECTRON TEMPERATURES AND CHEMICAL ABUNDANCES

4.1. Electron Temperature and Density

The broad wavelength coverage (rest-frame $\sim 2900 -$ 9400Å) and high SNR of the stacked LCE spectrum enables calculations of direct electron temperatures and densities from multiple ions. We detect the auroral emission lines of $[OIII]\lambda 4363$, $[OII]\lambda \lambda 7320$, 7330, $[SIII]\lambda 6312$, and $[SII]\lambda\lambda4069, 4076$, which are shown in Figure 1. These let us use the temperature-sensitive auroral-tonebular line ratios to calculate the electron temperatures for O^{++} , O^{+} , S^{++} , and S^{+} . Our spectra do not go red enough to cover the $[SIII]\lambda 9532$ line, so the S⁺⁺ temperature is measured from $[SIII]\lambda 6312/\lambda 9069$. While we detect both $[SII]\lambda\lambda4069, 4076$ lines, the redder line is barely detected ($< 3\sigma$). We therefore choose to use the ratio $[SII]\lambda 4069/\lambda\lambda 6717, 6731$ to calculate the temperature for this ion. We do not detect the $[NII]\lambda 5755$ line, and instead set an upper limit on its strength. This sets an upper limit on the ratio $[NII]\lambda 5755/\lambda 6584$, and thus an upper limit on the N^+ temperature. We also detect the density-sensitive lines $[SII]\lambda\lambda 6717, 6731$, $[OII]\lambda\lambda3727, 3729$, and $[ArIV]\lambda\lambda4713, 4741$. We calculate the electron temperatures and densities from these lines using the getTemDen function from the python package PYNEB (Luridiana et al. 2015). The resulting measurements are presented in Table 3.

Uncertainties on temperatures and densities are calculated following the same method as Welch et al. (2024). We draw 300 points from a Gaussian distribution centered on the measured line ratio, with standard deviation equal to the uncertainty on the line ratio. We calculate the temperature or density for each point, then take the standard deviation of the resulting temperature/density distribution as our uncertainty. In cases where the final distribution is highly asymmetric, we use the 16th and 84th percentiles as our 1σ -low and 1σ -high uncertainties.

We calculate the electron temperatures assuming a fixed density. We use the best fit value from the $[SII]\lambda\lambda6717,6731$ ratio $(n_e = 1200 \text{ cm}^{-3})$ as our fiducial density for temperature calculations. Similarly, we use a fixed temperature for our density calculations. We adopt the O⁺⁺ temperature ($T_e = 14000 \text{ K}$) as our fiducial value for density calculations. The temperatures and densities are not strongly dependent on each other. We find that any reasonable change in density (within $\sim 1\sigma$ of the S⁺ density) does not change our temperature estimates beyond the quoted uncertainty range, and any reasonable change in temperature (within $\sim 1\sigma$




Figure 2. The spectrum of the Sunburst LCE shows a clear Balmer jump at 3646 Å (marked with the vertical dashed line). The strength of this jump is sensitive to the electron temperature in the ionized hydrogen gas. The jump is measured by fitting the continuum on either side of the discontinuity, as described in Section 4.1.1. The wavelength region used to fit the continuum blueward of the jump is shown in blue, while the region used to fit the continuum redward of the jump is shown in red. Linear fits to each continuum region are shown in the corresponding colors.

of the O^{++} temperature) does not change the density estimate beyond the quoted uncertainty range.

4.1.1. Balmer Jump Temperature

Our spectrum covers the Balmer jump at λ 3646Å, and the high SNR of the stacked spectrum allow us to detect this continuum feature. The strength of the jump is sensitive to the electron temperature of the ionized hydrogen in the nebula. We measure the jump following the procedure of Liu et al. (2001). The continuum redward and blueward of the jump is measured with a linear fit (shown in Figure 2). On the red side of the jump, we mask emission lines prior to fitting. We also mask the region between the jump at λ 3646Å and the H12 line, where blended high-order Balmer lines will bias the continuum level. We evaluate the blue linear fit at 3643Å, and the red linear fit at 3681Å.

Following Equation 3 of Liu et al. (2001), the Balmer jump temperature depends on the ratio of the jump strength to the H11 line flux, and the helium abundances He^+/H^+ and He^{++}/H^+ . Our helium abundance measurements are described in greater detail in the following section. We do not detect nebular HeII λ 4686; while the line is present, we only see evidence of a broad component originating in the WR stars (see Rivera-Thorsen et al. 2024). We therefore assume the He⁺⁺/H⁺ = 0. The dependence on helium abundance is weak (see also Fig. 9 of Liu et al. 2001). As a test of this weak dependence, if we assume the He⁺⁺ abundance is 50% of the He⁺, the calculated temperature does not change beyond our quoted 1σ uncertainties.

With this method, we measure a Balmer jump temperature of $T_e(\text{Bal}) = 12000 \pm 300 \text{ K}$. This temperature is largely consistent with our other measured electron temperatures.

4.1.2. Comparison of Electron Temperatures & Densities

We have measured electron temperatures and densities from multiple ionization zones within the Sunburst LCE nebula. The densities we measure are generally consistent within 1σ uncertainties, however there is some evidence that the high-ionization gas density is higher than the lower ionization states. While the low ionization estimates from S⁺ ($n_e = 1200^{+500}_{-300} \text{cm}^{-3}$) and O⁺ $(n_e = 1600^{+1500}_{-200} \text{ cm}^{-3})$ are very consistent, the density measured from $[ArIV]\lambda\lambda4713, 4741(n_e) =$ 4500^{+10000}_{-2500} cm⁻³) is somewhat higher. However, the $[ArIV]\lambda\lambda4713,4741$ lines are much fainter than either $[SII]\lambda\lambda 6717, 6731$ or $[OII]\lambda\lambda 3727, 3729$, leading to a much larger uncertainty on the resulting den-The high density reported from sity estimate. $[ArIV]\lambda\lambda4713,4741$ could thus be evidence that the high ionization gas is denser than the surrounding lowionization gas, however the large uncertainty means that we cannot rule out these two ionization states having equal density.

Electron temperature measurements show greater variation across ionization zones and across diagnostics. The lower-ionization gas probed by $T_e(S^+)$ (10.36–23.34 eV) and $T_e(S^{++})$ (23.34–34.86 eV) favors slightly lower electron temperatures (10000 ± 1000 and 10300 ± 600 K, respectively). Meanwhile, the higher ionization states observed with $T_e(O^{++})$ (35.12–54.94 eV) and $T_e(Ne^{++})$ (40.96–63.42 eV) show slightly higher temperatures (15100 ± 500 and 15400 ± 1500 K, respectively). These differences between ionization zones highlight the value of obtaining T_e measurements from ions at different ionization energies, as empirical temperature relations do not fully capture such variations.

One notable exception to this trend is $T_e(O^+) = 28000\pm3000$ K, which is an outlier in our measurements. At high densities $(n_e > 10^3 \text{ cm}^{-3})$, dielectronic recombination contributes substantially to the [OII] emission (Liu et al. 2001); this artificially increases the inferred O^+ electron temperature. All of our densities are above this threshold, thus we conclude that the O^+ temperature is not reliable for this object.

We also have a measurement of the Balmer jump, which is sensitive to the electron temperature in the hydrogen gas across the whole nebula. We find that the




Figure 3. Comparison of electron temperatures measured from auroral lines of different ions in the Sunburst Arc (black point), plotted alongside local HII regions from CHAOS (blue points, Berg et al. 2020). The black dashed line represents equal temperatures. The electron temperature for O^+ is biased high due to significant contribution to the $[OII]\lambda\lambda7320,7330$ flux from dielectronic recombination. We therefore do not consider the O^+ temperature reliable. The upper limit placed on the N⁺ temperature does not offer a strong constraint when compared to expectations based on other well-measured temperatures.

Balmer temperature is closer to the measurements from low ionization gas, though the larger uncertainty on the Balmer temperature means that it is still consistent with the high ionization temperatures within 1σ uncertainty.

4.2. Abundances and Ionization Corrections

Ionic abundances relative to hydrogen are calculated using

$$\frac{N(X^i)}{N(H^+)} = \frac{I_{\lambda(i)}}{I_{H\beta}} \frac{j_{H\beta}}{j_{\lambda(i)}}.$$
(1)

The emissivity coeffecients $j_{\lambda(i)}$ are temperature dependent. We calculate these coeffecients using the getIonAbundance function from PYNEB (Luridiana et al.

2015). We use the temperature from the nearest ionization zones for each ionic abundance calculation, as described below.

Atomic data used in our calculations are tabulated in Table 2.

4.2.1. Oxygen Abundance

We calculate the total oxygen abundance O/H = $O^+/H^+ + O^{++}/H^+$. We use the electron temperature $T_e(O^{++})$ determined from [OIII] λ 4363 to calculate the O^{++}/H^+ ionic abundance.

For O⁺, the electron temperature measured from the $[OII]\lambda\lambda7320,7330$ lines is not reliable at the densities measured, as discussed in the Section 4.1.2. We there-

Table 2. Atomic Data Sources

Ion	Transition Probabilities	Collision Strengths
O^+	Froese Fischer & Tachiev (2004)	Kisielius et al. (2009)
O^{++}	Froese Fischer & Tachiev (2004)	Aggarwal & Keenan (1999)
N^+	Froese Fischer & Tachiev (2004)	Tayal (2011)
S^+	Rynkun et al. (2019)	Tayal & Zatsarinny (2010)
S^{++}	Tayal et al. (2019)	Grieve et al. (2014)
Ar^{++}	Mendoza & Zeippen (1983)	Munoz Burgos et al. (2009)
Ar^{+3}	Rynkun et al. (2019)	Ramsbottom et al. (1997)
Ne^{++}	Froese Fischer & Tachiev (2004)	McLaughlin & Bell (2000)

fore use the S⁺ temperature as our low-density electron temperature, as the [SII] $\lambda\lambda$ 4069, 4076 lines are not similarly inflated. Calculating the O⁺/H⁺ abundance using the high O⁺ temperature leads to a ~ 10× reduction in the inferred ionic abundance.

The final oxygen ion abundances are reported in Table 3. We find a total metallicity for this object of $12 + \log(O/H) = 7.97 \pm 0.05$.

4.2.2. Nitrogen Abundance

We make the assumption that $N/O \simeq N^+/O^+$, based on the similar ionization zones of the N^+ and O^+ ions (Peimbert 1967). This assumption has been found to be valid within ~ 10% (Nava et al. 2006; Amayo et al. 2021).

Because we do not detect the $[NII]\lambda 5755$ line, we use the temperature from the $[SII]\lambda 4069/\lambda\lambda 6717, 6731$ ratio as our low-ionization temperature when calculating the nitrogen abundance. One potential caveat when using the S⁺ temperature is that the lower bound of the S⁺ ionization energy (10.36 eV) is lower than the hydrogen ionization energy (13.6 eV). Thus S⁺ can form outside the H⁺ boundary. We note however that the Balmer jump, sensitive to the electron temperature of H⁺, yields a temperature that is consistent within uncertainties of the S⁺ temperature. Thus we would not expect a significant change in our measured N⁺ abundance from this effect.

4.2.3. Sulfur Abundance

Our spectrum contains the temperature sensitive lines for both S^+ and S^{++} , which we use to calculate the abundance of the lower ionization states of sulfur. We do not have a measurement of a higher ionization state of sulfur, so we must correct for the unobserved high ionization sulfur. There may be a small contribution from S^{+3} (34.86 – 47.22 eV ionization energy). We use the ionization correction factor (ICF) of Izotov et al. (2006), specifically the "low-Z" component of their Equation 20, to make this correction. We find that the ICF is consistent with 1 within uncertainties, indicating that we have measured the primary states of sulfur within the nebula. We use the direct S^+ and S^{++} temperatures to measure the abundances of these sulfur ions.

4.2.4. Argon Abundance

We detect both $[ArIII]\lambda7138$ and $[ArIV]\lambda\lambda4713, 4741$, allowing us to calculate the ionic abundances of both Ar^{++} and Ar^{+3} . We correct for unobserved higher ionization states of argon using the ICF in Equation 23 of Izotov et al. (2006), specifically using the "low-Z" component based on our measured oxygen abundance. Most of the gas-phase argon is in the two ionization states which we have measured, resulting in an ICF of 1 ± 0.2 and a decreased systematic uncertainty on the total argon abundance.

The ionization energy range of Ar⁺⁺ does not neatly align with any ion for which we have a temperature measurement; it lies between the S⁺⁺ and O⁺⁺ zones. The overlap with S⁺⁺ is slightly greater, so we choose to use $T_e(S^{++})$ when calculating Ar⁺⁺/H⁺. We did test the effect of using the O⁺⁺ temperature, and found that the resulting Ar⁺⁺/H⁺ ionic abundance is lower by a factor ~ 2 . Ar⁺³ better aligns with O⁺⁺, so we use $T_e(O^{++})$ when calculating Ar³⁺/H⁺.

4.2.5. Neon Abundance

We only detect neon in the Ne⁺⁺ state, requiring a correction for unobserved ionization states. We again use the ICF of Izotov et al. (2006) for this correction. The neon abundance and ICF are reported in Table 3. We use the electron temperature from O⁺⁺ to calculate the neon ionic abundance. While we have a direct measurement of $T_e(Ne^{++})$, this temperature has a relatively large uncertainty (owing to the faintness of the [NeIII] λ 3343 line). Additionally, it is consistent within 1 σ uncertainties of $T_e(O^{++})$. Thus using the doubly ionized oxygen temperature reduces the statistical uncertainty on the neon abundance measurement without introducing any additional systematic uncertainty, as the two temperatures have been found to be consistent.

We find that the ionization correction for neon is 1.1 ± 0.1 , indicating that the majority of the neon is contained within the measured Ne⁺⁺ state.

4.2.6. Iron Abundance

We calculate the gas-phase iron abundance based on the strength of the nebular [FeIII] λ 4658 line. To account for the unobserved ionization states of iron, we use the ICF from Equation 24 of Izotov et al. (2006). The ionization energy of Fe⁺⁺ (16.2–30.7 eV) overlaps both the low and intermediate ionization zones. We choose to use the low-ionization temperature from S⁺ for the iron abundance calculation, though the final result would not change significantly if we instead used the \mathbf{S}^{++} temperature.

The iron abundance here relies on only a single ionization state, and as such the ICF (6 ± 2) is large. This abundance is thus subject to additional systematic uncertainties. While we detect [FeII] λ 4360, this line is strongly affected by flourescence and is thus not suitable for abundance determinations. The [FeII] λ 4288 line appears to be partially detected, however it straddles the detector gap in the NIRSpec IFS data so we cannot measure a reliable flux. We do not detect any higherionization iron lines, though we may expect some contribution from Fe⁺³, which has similar ionization energy to O⁺⁺.

4.3. Helium Abundance

We measure a variety of He I lines that allow us to calculate the helium abundance, which is needed for the Balmer jump temperature calculation. While we detect HeII λ 4686 emission, we only see evidence of the broad stellar component, without significant contribution from a narrow nebular component. We therefore assume that the He⁺⁺/H⁺ abundance is small relative to He⁺/H⁺, and thus He/H = He⁺/H⁺.

The He I ionization energy is most similar to that of O^{++} , so we use $T_e(O^{++})$ when calculating the helium abundance. We have a number of He I lines available to calculate the ionic abundance. We find that the various lines give similar abundance results (reported in Table 3. For our final value, we use an error-weighted average of the abundance measurements from each line.

5. DISCUSSION

5.1. High-Quality Abundances of Multiple Species at z = 2.37

We have detected the auroral lines $[SII]\lambda\lambda4069, 4076$, $[OIII]\lambda4363$, $[SIII]\lambda6312$, and $[OII]\lambda\lambda7320, 7330$, allowing us to measure the electron temperature in the low, intermediate, and high ionization zones of the nebula. The resulting temperatures are reported in Table 3, and plotted in Figure 3 alongside temperature measurements from local HII regions from the CHAOS project (Berg et al. 2015; Croxall et al. 2015, 2016; Berg et al. 2020). We do not detect [NII] λ 5755, so we only set an upper limit on the N⁺ temperature.

These electron temperatures allow us to calculate abundances of multiple species via the direct method. The resulting ionic abundance, ICFs, and abundances relative to oxygen are reported in Table 3. Our spectrum includes emission lines from the dominant ionization states of oxygen, nitrogen, sulfur, argon, and neon,

 Table 3. Temperature, Density, and Abundance Measurements from the Ly-C Leaking Cluster

Property	Measurement
$T_e({\rm O}^{++})~(10^3{\rm K})$	15.1 ± 0.5
$T_e({\rm O}^+)~(10^3{\rm K})$	28 ± 3
$T_e({ m S}^{++})~(10^3{ m K})$	10.3 ± 0.6
$T_e({ m S}^+)~(10^3{ m K})$	10 ± 1
$T_e({\rm N^+})~(10^3{\rm K})$	< 26.3
$T_e({\rm Ne}^{++})~(10^3{\rm K})$	15.4 ± 1.5
$T_e(\text{Balmer}) (10^3 \text{K})$	12 ± 3
$n_e({\rm O}^+)~({\rm cm}^{-3})$	1600^{+1500}_{-200}
$n_e({ m S}^+)~({ m cm}^{-3})$	1200^{+500}_{-400}
$n_e({\rm Ar}^{+3})~({\rm cm}^{-3})$	4500^{+10000}_{-2500}
$O^+/H^+ (\times 10^5)$	$2.0^{+1.1}_{-0.7}$
$O^{++}/H^+ (\times 10^5)$	7.3 ± 1.2
$12 + \log(O/H)$	7.97 ± 0.05
$S^+/H^+ (\times 10^7)$	2.3 ± 0.7
$S^{++}/H^+ (\times 10^7)$	21 ± 4
S ICF	1.3 ± 0.3
$12 + \log(S/H)$	6.48 ± 0.05
$\log(S/O)$	-1.49 ± 0.14
$N^+/H^+ (\times 10^7)$	46 ± 7
$\log(N/O)$	$-0.65^{+0.16}_{-0.25}$
$Ar^{++}/H^+ (\times 10^7)$	4.5 ± 0.9
${\rm Ar^{+3}/H^{+}}~(\times 10^7)$	0.7 ± 0.2
Ar ICF	1.0 ± 0.2
$12 + \log(Ar/H)$	5.72 ± 0.04
$\log(Ar/O)$	-2.24 ± 0.14
$Ne^{++}/H^+ (\times 10^5)$	1.5 ± 0.3
Ne ICF	1.1 ± 0.1
$12 + \log({\rm Ne/H})$	7.21 ± 0.04
$\log({\rm Ne/O})$	-0.66 ± 0.11
${\rm Fe^{++}/H^+}~(\times 10^7)$	3.1 ± 0.1
Fe ICF	6 ± 2
$12 + \log(\mathrm{Fe/H})$	6.28 ± 0.06
$\log({\rm Fe}/{\rm O})$	-1.7 ± 0.2
$\mathrm{He^{+}/H^{+}(\lambda4471)}$	0.08 ± 0.01
$\mathrm{He^{+}/H^{+}}(\lambda5876)$	0.09 ± 0.01
$\mathrm{He^+/H^+}(\lambda6678)$	0.09 ± 0.01
$\rm Mean~He^+/H^+$	0.08 ± 0.02
$12 + \log(\mathrm{He/H})$	10.93 ± 0.09

minimizing the uncertainties propagated to our abundances from ionization correction factors.

This combination of directly measured electron temperatures for each of the low, intermediate, and high ionization zones, plus detection of the relevant emission lines for the primary states of multiple elements, enables the same quality of abundance measurements as has been used in nearby galaxies. We can therefore directly compare the gas-phase abundances of this young star cluster at z = 2.37 to abundances measured in nearby HII regions. This is the first time chemical abundance patterns have been measured to this level of precision in a galaxy beyond the local universe.

We plot the abundances of nitrogen, sulfur, argon, neon, and iron relative to oxygen in Figure 4. We compare our abundances to measurements from local HII regions from CHAOS (Berg et al. 2015; Croxall et al. 2015, 2016; Berg et al. 2020), low-redshift galaxies from SDSS (Izotov et al. 2006), and other galaxies measured at cosmic noon (Sanders et al. 2023b; Rogers et al. 2024; Welch et al. 2024). Solar abundances from Asplund et al. (2021) are also marked. We find that our measured abundances of sulfur, argon, and neon are all consistent with solar abundance patterns. Additionally, these elements are consistent with local HII regions, indicating the enrichment of alpha elements in the LCE is following standard patterns, in contrast with the low alpha abundances for a $z \sim 3$ galaxy reported in Rogers et al. (2024). Our nitrogen abundance is high compared to local HII regions, similar to the $z \sim 2$ galaxies reported in Sanders et al. (2023b). We discuss this enhanced nitrogen abundance in the following section. We find an iron abundance that is slightly below the solar value, however it matches the Fe/O abundances of low-redshift galaxies with similar oxygen abundances (Izotov et al. 2006). Our measured Fe/O is still higher than what has been observed in the Magellanic Clouds (Domínguez-Guzmán et al. 2022), however our measured dust E(B-V) is also lower, indicating that there is less dust available onto which the iron can condense.

5.2. Nitrogen Enrichment from Wolf-Rayet Stars

The oxygen abundance of the LCE places it just below the point at which measurements of local HII regions have begun to find significant contribution from secondary nitrogen production. We might therefore expect it to only be affected by primary nitrogen production, placing it along the N/O plateau for low-metallicity galaxies and HII regions (e.g. within uncertainties of the photoionization model prediction from Nicholls et al. (2017)). Additionally, the young stellar age that has been previously reported for this star cluster (< 4 Myr, Chisholm et al. (2019)) would typically be indicative of only primary nitrogen production driving the N/O abundance. The elevated N/O measured for the LCE is in tension with this interpretation, indicating another source of nitrogen enrichment is driving the abundance.

Elevated nitrogen abundances have been observed previously in low-redshift galaxies (e.g., Brinchmann et al. 2008; Amorín et al. 2010; Berg et al. 2011), galaxies at cosmic noon (e.g., Masters et al. 2014; Sanders et al. 2023b), and more recently in high redshift galaxies (Cameron et al. 2023; Senchyna et al. 2023; Marques-Chaves et al. 2024; Castellano et al. 2024). The elevated nitrogen has often been attributed to nitrogen enrichment from Wolf-Rayet stars (Henry et al. 2000; Kobayashi & Ferrara 2024). However it has also been suggested that inflows of low-metallicity gas can dilute the oxygen abundance, driving increases in the N/O ratio without requiring enrichment from Wolf-Rayet stars (e.g., Köppen & Hensler 2005; Amorín et al. 2010; Andrews & Martini 2013).

A companion paper utilizing the same stacked spectrum determined that the LCE contains spectral signatures of Wolf-Rayet stars (Rivera-Thorsen et al. 2024). Previous analyses of galaxies showing WR signatures at low redshift have found that abundances of alpha elements (Ar & Ne) relative to oxygen were not elevated (Brinchmann et al. 2008). Recent theoretical analysis of WR abundance patterns from Kobayashi & Ferrara (2024) find a similar result, that only the nitrogen abundance will change significantly with enrichment from WR stars. We see a similar abundance pattern here, with only N/O elevated relative to similarly low-O/H galaxies and HII regions. The similar abundance pattern could be indicative of WR stars driving the elevated N/O in the Sunburst LCE. However earlier studies have suggested that the WR stellar winds driving the enhanced nitrogen abundance would also carry significant helium (Kobulnicky et al. 1997). We find a helium abundance $12 + \log(\text{He/H}) = 10.93 \pm 0.09$ that is consistent with solar $(12 + \log(\text{He/H})_{\odot} = 10.914 \pm 0.013,$ Asplund et al. 2021). Brinchmann et al. (2008) also find no evidence for elevated helium abundance in their WR galaxy sample, yet still conclude that the elevated N/O is likely due to WR stellar winds. We thus make the same conclusion here, that WR stellar winds are driving increased N/O abundance in the Sunburst LCE. A recent analysis of several high-redshift $(z \sim 6)$ galaxies with elevated N/O also found evidence for elevated He/H (Yanagisawa et al. 2024). These authors suggest that elevated densities $(n_e \sim 10^3 - 10^4)$ could be related to the high helium abundance. We measure a similar density $(n_e \sim 1200 - 4500)$ but see no increase in helium abundance.

The type of WR stars present can also influence the measured chemical abundances. Analyses of wind abundances in WN and WC stars found that WNs exhibit fairly typical neon abundance, while WC stars show significantly elevated levels of neon (Smith & Houck 2005; Ignace et al. 2007). We see no evidence of enhanced neon abundance here, which could be indicative of a




Figure 4. Abundances of the LyC leaking clump, compared to local HII regions (Berg et al. 2020), low-redshift galaxies (Izotov et al. 2006), and other galaxies at Cosmic Noon (Rogers et al. 2024; Sanders et al. 2023b; Welch et al. 2024). Solar abundances from Asplund et al. (2021) are shown as black dashed lines. We find an enhanced nitrogen abundance, while other elements are consistent with local HII regions and low-redshift star-forming galaxies.

majority of the WR stars present being WN-type. In exploratory fitting by Rivera-Thorsen et al. (2024), the best-fitting BPASS model had only 13% of the WR stars of WC-type, offering some support to this hypothesis. However, more detailed analysis would be required to constrain the relative quantities of WC and WN stars.

A previous analysis of the LCE using rest-frame UV spectra also measured a high N/O abundance (Pascale et al. 2023). However, their measurement $(\log(N/O) =$ -0.21 ± 0.1) is higher than our direct measurement by ~ 0.3 dex, though the two are still consistent within 2σ . Pascale et al. (2023) hypothesize that the nitrogen enrichment is most enhanced in the high-density, highionization regions of the cluster. Our measurement using N^+ being slightly lower than their measurement based on N⁺⁺ would be consistent with this picture. However we also note that the [NII] transitions used here have lower critical densities than the higher-ionization lines, so collisional de-excitation could be lowering the measured flux. Given the high measured densities here, it is possible that our measured N/O is slightly low, owing to the effects of collisional de-excitation.

6. CONCLUSIONS

We have measured direct gas-phase chemical abundances in the stacked spectrum of the Lyman-continuum emitting star cluster in the Sunburst Arc. We detect the $[SII]\lambda\lambda4069, 4076, [OIII]\lambda4363, [SIII]\lambda6312$, and $[OII]\lambda\lambda7320, 7330$ lines, allowing measurement of the electron temperature in the low, intermediate, and high ionization zones of the nebular gas. With these multiple temperature measurements, we measure direct abundances of oxygen, nitrogen, sulfur, argon, neon, and iron, following best practices established in nearby galaxies from the CHAOS program. These measurements enable a direct comparison between well-studied nearby HII regions and an object at Cosmic Noon (z = 2.37).

These high-quality abundance measurements enable detailed studies of the origins of abundance patterns in distant galaxies. In the Sunburst LCE, the combination of these abundance measurements and the detection of Wolf-Rayet features (Rivera-Thorsen et al. 2024) allow us to conclude that nitrogen-rich Wolf-Rayet stars are responsible for increasing the nitrogen abundance.

Further high-quality abundance measurements of galaxies and galactic substructures at Cosmic Noon and beyond will continue to drive our understanding of how galaxies built up the chemical elements.

The authors wish to dedicate this paper to the memory of Prof. Liese Van Zee, a pioneer of chemical abundance measurements in galaxies. This work was based on observations taken with the NASA/ESA/CSA JWST. JWST is operated by the Space Telescope Science Institute under the management of the Association of Universities for Research in Astronomy, Inc., under NASA contract no. NAS 5-03127. These observations were taken as part of JWST GO program 2555, support for which was provided by NASA through a grant from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-03127. BW is supported by NASA under award number 80GSFC21M0002.

Facilities: JWST

Software: Astropy (Astropy Collaboration et al. 2013, 2018, 2022), Matplotlib (Hunter 2007), Scipy (Virtanen et al. 2020), emcee (Foreman-Mackey et al. 2013), PyNeb (Luridiana et al. 2015)

REFERENCES

- Aggarwal, K. M., & Keenan, F. P. 1999, ApJS, 123, 311, doi: 10.1086/313232
- Amayo, A., Delgado-Inglada, G., & Stasińska, G. 2021, MNRAS, 505, 2361, doi: 10.1093/mnras/stab1467
- Amorín, R. O., Pérez-Montero, E., & Vílchez, J. M. 2010, ApJL, 715, L128, doi: 10.1088/2041-8205/715/2/L128
- Andrews, B. H., & Martini, P. 2013, ApJ, 765, 140, doi: 10.1088/0004-637X/765/2/140
- Arellano-Córdova, K. Z., Berg, D. A., Chisholm, J., et al. 2022, ApJL, 940, L23, doi: 10.3847/2041-8213/ac9ab2
- Asplund, M., Amarsi, A. M., & Grevesse, N. 2021, A&A, 653, A141, doi: 10.1051/0004-6361/202140445

Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, A&A, 558, A33,

doi: 10.1051/0004-6361/201322068

- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, AJ, 156, 123, doi: 10.3847/1538-3881/aabc4f
- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, ApJ, 935, 167, doi: 10.3847/1538-4357/ac7c74
- Bayliss, M. B., Rigby, J. R., Sharon, K., et al. 2014, ApJ, 790, 144, doi: 10.1088/0004-637X/790/2/144
- Berg, D. A., Pogge, R. W., Skillman, E. D., et al. 2020, ApJ, 893, 96, doi: 10.3847/1538-4357/ab7eab

- Berg, D. A., Skillman, E. D., Croxall, K. V., et al. 2015, ApJ, 806, 16, doi: 10.1088/0004-637X/806/1/16
- Berg, D. A., Skillman, E. D., & Marble, A. R. 2011, ApJ, 738, 2, doi: 10.1088/0004-637X/738/1/2
- Böker, T., Beck, T. L., Birkmann, S. M., et al. 2023, PASP, 135, 038001, doi: 10.1088/1538-3873/acb846
- Brinchmann, J. 2023, MNRAS, 525, 2087, doi: 10.1093/mnras/stad1704
- Brinchmann, J., Kunth, D., & Durret, F. 2008, A&A, 485, 657, doi: 10.1051/0004-6361:200809783
- Bushouse, H., Eisenhamer, J., Dencheva, N., et al. 2023, JWST Calibration Pipeline, 1.11.4, Zenodo, doi: 10.5281/zenodo.8247246
- Cameron, A. J., Katz, H., Rey, M. P., & Saxena, A. 2023, MNRAS, 523, 3516, doi: 10.1093/mnras/stad1579
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, ApJ, 345, 245, doi: 10.1086/167900
- Castellano, M., Napolitano, L., Fontana, A., et al. 2024, arXiv e-prints, arXiv:2403.10238, doi: 10.48550/arXiv.2403.10238
- Chisholm, J., Rigby, J. R., Bayliss, M., et al. 2019, ApJ, 882, 182, doi: 10.3847/1538-4357/ab3104
- Christensen, L., Laursen, P., Richard, J., et al. 2012, MNRAS, 427, 1973, doi: 10.1111/j.1365-2966.2012.22007.x
- Citro, A., Berg, D. A., Erb, D. K., et al. 2023, arXiv
- e-prints, arXiv:2305.14414,
- doi: 10.48550/arXiv.2305.14414
- Croxall, K. V., Pogge, R. W., Berg, D. A., Skillman, E. D., & Moustakas, J. 2015, ApJ, 808, 42, doi: 10.1088/0004-637X/808/1/42
- —. 2016, ApJ, 830, 4, doi: 10.3847/0004-637X/830/1/4
- Curti, M., D'Eugenio, F., Carniani, S., et al. 2023, MNRAS, 518, 425, doi: 10.1093/mnras/stac2737
- Dahle, H., Aghanim, N., Guennou, L., et al. 2016, A&A, 590, L4, doi: 10.1051/0004-6361/201628297
- D'Eugenio, F., Maiolino, R., Carniani, S., et al. 2023, arXiv e-prints, arXiv:2311.09908, doi: 10.48550/arXiv.2311.09908
- Diego, J. M., Pascale, M., Kavanagh, B. J., et al. 2022, A&A, 665, A134, doi: 10.1051/0004-6361/202243605
- Dinerstein, H. L. 1990, in Astrophysics and Space Science Library, Vol. 161, The Interstellar Medium in Galaxies, ed. J. Thronson, Harley A. & J. M. Shull, 257–285, doi: 10.1007/978-94-009-0595-5_10
- Domínguez-Guzmán, G., Rodríguez, M., García-Rojas, J., Esteban, C., & Toribio San Cipriano, L. 2022, MNRAS, 517, 4497, doi: 10.1093/mnras/stac2974
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, PASP, 125, 306, doi: 10.1086/670067

- Froese Fischer, C., & Tachiev, G. 2004, Atomic Data and Nuclear Data Tables, 87, 1, doi: https://doi.org/10.1016/j.adt.2004.02.001
 Gardner, J. P., Mather, J. C., Abbott, R., et al. 2023, PASP, 135, 068001, doi: 10.1088/1538-3873/acd1b5
 Gordon, K. D., Clayton, G. C., Misselt, K. A., Landolt,
- A. U., & Wolff, M. J. 2003, ApJ, 594, 279,
 doi: 10.1086/376774
- Grieve, M. F. R., Ramsbottom, C. A., Hudson, C. E., & Keenan, F. P. 2014, ApJ, 780, 110, doi: 10.1088/0004-637X/780/1/110
- Henry, R. B. C., Edmunds, M. G., & Köppen, J. 2000, ApJ, 541, 660, doi: 10.1086/309471
- Hsiao, T. Y.-Y., Álvarez-Márquez, J., Coe, D., et al. 2024, arXiv e-prints, arXiv:2404.16200, doi: 10.48550/arXiv.2404.16200
- Hunter, J. D. 2007, Computing in Science & Engineering, 9, 90, doi: 10.1109/MCSE.2007.55
- Hutchison, T. A., Welch, B. D., Rigby, J. R., et al. 2023, arXiv e-prints, arXiv:2312.12518, doi: 10.48550/arXiv.2312.12518
- —. 2024, baryon-sweep: Outlier rejection algorithm for JWST/NIRSpec IFS data, Astrophysics Source Code Library, record ascl:2401.012
- Ignace, R., Cassinelli, J. P., Tracy, G., Churchwell, E., & Lamers, H. J. G. L. M. 2007, ApJ, 669, 600, doi: 10.1086/521718
- Izotov, Y. I., Stasińska, G., Meynet, G., Guseva, N. G., & Thuan, T. X. 2006, A&A, 448, 955, doi: 10.1051/0004-6361:20053763
- James, B. L., Pettini, M., Christensen, L., et al. 2014, MNRAS, 440, 1794, doi: 10.1093/mnras/stu287
- Jones, T., Sanders, R., Chen, Y., et al. 2023, ApJL, 951, L17, doi: 10.3847/2041-8213/acd938
- Katz, H., Saxena, A., Cameron, A. J., et al. 2023, MNRAS, 518, 592, doi: 10.1093/mnras/stac2657
- Kim, K. J., Bayliss, M. B., Rigby, J. R., et al. 2023, ApJL, 955, L17, doi: 10.3847/2041-8213/acf0c5
- Kisielius, R., Storey, P. J., Ferland, G. J., & Keenan, F. P. 2009, MNRAS, 397, 903,
- doi: 10.1111/j.1365-2966.2009.14989.x
- Kobayashi, C., & Ferrara, A. 2024, ApJL, 962, L6, doi: 10.3847/2041-8213/ad1de1
- Kobulnicky, H. A., Skillman, E. D., Roy, J.-R., Walsh,
 J. R., & Rosa, M. R. 1997, ApJ, 477, 679,
 doi: 10.1086/303742
- Kojima, T., Ouchi, M., Nakajima, K., et al. 2017, PASJ, 69, 44, doi: 10.1093/pasj/psx017
- Köppen, J., & Hensler, G. 2005, A&A, 434, 531, doi: 10.1051/0004-6361:20042266

- Liu, X. W., Luo, S. G., Barlow, M. J., Danziger, I. J., & Storey, P. J. 2001, MNRAS, 327, 141, doi: 10.1046/j.1365-8711.2001.04676.x
- Luridiana, V., Morisset, C., & Shaw, R. A. 2015, A&A, 573, A42, doi: 10.1051/0004-6361/201323152
- Ly, C., Rigby, J. R., Cooper, M., & Yan, R. 2015, ApJ, 805, 45, doi: 10.1088/0004-637X/805/1/45
- Marques-Chaves, R., Schaerer, D., Kuruvanthodi, A., et al. 2024, A&A, 681, A30, doi: 10.1051/0004-6361/202347411
- Masters, D., McCarthy, P., Siana, B., et al. 2014, ApJ, 785, 153, doi: 10.1088/0004-637X/785/2/153
- McLaughlin, B. M., & Bell, K. L. 2000, Journal of Physics
 B: Atomic, Molecular and Optical Physics, 33, 597, doi: 10.1088/0953-4075/33/4/301
- Mendoza, C., & Zeippen, C. J. 1983, MNRAS, 202, 981, doi: 10.1093/mnras/202.4.981
- Meštrić, U., Vanzella, E., Upadhyaya, A., et al. 2023, A&A, 673, A50, doi: 10.1051/0004-6361/202345895
- Munoz Burgos, J. M., Loch, S. D., Ballance, C. P., & Boivin, R. F. 2009, A&A, 500, 1253, doi: 10.1051/0004-6361/200911743
- Nava, A., Casebeer, D., Henry, R. B. C., & Jevremovic, D. 2006, ApJ, 645, 1076, doi: 10.1086/504416
- Nicholls, D. C., Sutherland, R. S., Dopita, M. A., Kewley, L. J., & Groves, B. A. 2017, MNRAS, 466, 4403, doi: 10.1093/mnras/stw3235
- Pascale, M., Dai, L., McKee, C. F., & Tsang, B. T. H. 2023, ApJ, 957, 77, doi: 10.3847/1538-4357/acf75c
- Peimbert, M. 1967, ApJ, 150, 825, doi: 10.1086/149385
- Pignataro, G. V., Bergamini, P., Meneghetti, M., et al. 2021, A&A, 655, A81, doi: 10.1051/0004-6361/202141586
- Pontoppidan, K. M., Barrientes, J., Blome, C., et al. 2022, ApJL, 936, L14, doi: 10.3847/2041-8213/ac8a4e
- Ramsbottom, C. A., Bell, K. L., & Keenan, F. P. 1997, MNRAS, 284, 754, doi: 10.1093/mnras/284.3.754
- Rhoads, J. E., Wold, I. G. B., Harish, S., et al. 2023, ApJL, 942, L14, doi: 10.3847/2041-8213/acaaaf
- Rieke, M. J., Kelly, D. M., Misselt, K., et al. 2023, PASP, 135, 028001, doi: 10.1088/1538-3873/acac53
- Rigby, J., Hutchison, T. A., Rivera-Thorsen, T. E., et al. 2024, JWST-Templates/Notebooks: minor update, added DOI tracker, v1.0.2, Zenodo, doi: 10.5281/zenodo.10933642
- Rigby, J., Perrin, M., McElwain, M., et al. 2023a, PASP, 135, 048001, doi: 10.1088/1538-3873/acb293
- Rigby, J. R., Vieira, J. D., Phadke, K. A., et al. 2023b, arXiv e-prints, arXiv:2312.10465, doi: 10.48550/arXiv.2312.10465
- Rivera-Thorsen, T. E., Dahle, H., Chisholm, J., et al. 2019, Science, 366, 738, doi: 10.1126/science.aaw0978

- Rivera-Thorsen, T. E., Chisholm, J., Welch, B., et al. 2024, arXiv e-prints, arXiv:2404.08884, doi: 10.48550/arXiv.2404.08884
- Rogers, N. S. J., Skillman, E. D., Pogge, R. W., et al. 2022, ApJ, 939, 44, doi: 10.3847/1538-4357/ac947d
- —. 2021, ApJ, 915, 21, doi: 10.3847/1538-4357/abf8b9
- Rogers, N. S. J., Strom, A. L., Rudie, G. C., et al. 2024, ApJL, 964, L12, doi: 10.3847/2041-8213/ad2f37
- Rynkun, P., Gaigalas, G., & Jönsson, P. 2019, A&A, 623, A155, doi: 10.1051/0004-6361/201834931
- Sanders, R. L., Shapley, A. E., Topping, M. W., Reddy, N. A., & Brammer, G. B. 2023a, arXiv e-prints, arXiv:2303.08149, doi: 10.48550/arXiv.2303.08149
- Sanders, R. L., Shapley, A. E., Reddy, N. A., et al. 2020, MNRAS, 491, 1427, doi: 10.1093/mnras/stz3032
- Sanders, R. L., Shapley, A. E., Clarke, L., et al. 2023b, ApJ, 943, 75, doi: 10.3847/1538-4357/aca9cc
- Schaerer, D., Marques-Chaves, R., Barrufet, L., et al. 2022, A&A, 665, L4, doi: 10.1051/0004-6361/202244556
- Schlafly, E. F., & Finkbeiner, D. P. 2011, ApJ, 737, 103, doi: 10.1088/0004-637X/737/2/103
- Senchyna, P., Plat, A., Stark, D. P., & Rudie, G. C. 2023, arXiv e-prints, arXiv:2303.04179, doi: 10.48550/arXiv.2303.04179
- Sharon, K., Mahler, G., Rivera-Thorsen, T. E., et al. 2022, ApJ, 941, 203, doi: 10.3847/1538-4357/ac927a
- Smith, J.-D. T., & Houck, J. R. 2005, ApJ, 622, 1044, doi: 10.1086/427979
- Strom, A. L., Rudie, G. C., Trainor, R. F., et al. 2023, arXiv e-prints, arXiv:2308.13508, doi: 10.48550/arXiv.2308.13508
- Tayal, S. S. 2011, ApJS, 195, 12, doi: 10.1088/0067-0049/195/2/12
- Tayal, S. S., & Zatsarinny, O. 2010, ApJS, 188, 32, doi: 10.1088/0067-0049/188/1/32
- Tayal, S. S., Zatsarinny, O., & Sossah, A. M. 2019, ApJS, 242, 9, doi: 10.3847/1538-4365/ab17e1
- Taylor, A. J., Barger, A. J., & Cowie, L. L. 2022, ApJL, 939, L3, doi: 10.3847/2041-8213/ac959d
- Trump, J. R., Arrabal Haro, P., Simons, R. C., et al. 2023, ApJ, 945, 35, doi: 10.3847/1538-4357/acba8a
- Trussler, J. A. A., Adams, N. J., Conselice, C. J., et al. 2023, MNRAS, 523, 3423, doi: 10.1093/mnras/stad1629
- van Zee, L., & Haynes, M. P. 2006, ApJ, 636, 214, doi: 10.1086/498017
- Vanzella, E., Castellano, M., Bergamini, P., et al. 2022, A&A, 659, A2, doi: 10.1051/0004-6361/202141590
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, Nature Methods, 17, 261, doi: 10.1038/s41592-019-0686-2

Welch, B., Olivier, G. M., Hutchison, T. A., et al. 2024, arXiv e-prints, arXiv:2401.13046, doi: 10.48550/arXiv.2401.13046 Yanagisawa, H., Ouchi, M., Watanabe, K., et al. 2024, arXiv e-prints, arXiv:2405.01823, doi: 10.48550/arXiv.2405.01823