

Assorted remarks on bending measures and energies for plates and shells, and their invariance properties

E. Vitral*

*Department of Mechanical Engineering, Rose-Hulman Institute of Technology
5500 Wabash Ave., Terre Haute, IN 47803, U.S.A.*

J. A. Hanna†

*Department of Mechanical Engineering, University of Nevada,
1664 N. Virginia St. (0312), Reno, NV 89557-0312, U.S.A.
(Dated: May 13, 2024)*

In this note, we address several issues, including some raised in recent works and commentary, related to bending measures and energies for plates and shells, and certain of their invariance properties. We discuss the distinction between definitions and results in our and others' approaches, correct an error and citation oversights in our work, and provide additional brief observations regarding the relative size of energetic terms and the symmetrization of bending measures. Particular points of emphasis are a reiteration of some of the early history of dilation-invariant measures, the similarities between all such measures, and the non-dilation-invariance of our recently introduced bending measure for shells and curved rods. In the course of this discussion, we provide a simpler presentation of the elementary, but much overlooked, fact that the additional tangential stretch of material near the mid-surface of a thin body is the product of the mid-surface stretch and the change in curvature.

This note serves as an extended comment on our own and others' papers, particularly [1–6]. Therefore, some conceptual background is presumed on the part of the reader, and we forego introductory or concluding discussion.

I. REMARKS ON DIFFERENT DEFINITIONS OF STRETCHING, AND RESULTING INVARIANT BENDING MEASURES

In the first part of our work [1], we introduced and employed a structure-dependent definition of stretching, one that treats the normal direction through a thin body differently than the tangential directions. We called “pure stretching” of such a body a deformation featuring through-thickness uniformity of stretch of tangentially-oriented fibers. By extension of this idea, an appropriate bending measure would be unchanged by such deformations. This approach gave rise to certain bending measures and associated bending energies with a simple constitutive response, such that a pure moment is both linear in the bending measure and induces no stretching on a neutral surface. The location of this preserved neutral surface depends on the referential curvature, but not the magnitude of the applied moment; for a flat plate, it is the mid-surface. An independent and equally important path to these same bending measures was taken in the second part of our work [2], in which we dimensionally reduced a 3D isotropic elastic energy quadratic in Biot-like strains, which are linear in the stretch tensor. Regardless of the choice of energy, typical thin shell kinematics provides the form of the tangential stretch off of the mid-surface which leads to our bending measures, a point we will re-emphasize below.

* vitralfr@rose-hulman.edu

† jhanna@unr.edu

In our work, we overlooked an earlier paper by Acharya [3] in which the topic of bending measures was addressed by a different approach. Acharya's approach employs a definition of "pure stretch" which, at its core, restricts the rotation field in the polar decomposition of the deformation gradient to be uniform. This definition is universal, remaining the same whether applied to a bulk solid or a thin body or a 2D surface of any shape; it does not distinguish between the normal direction and the tangential directions of a plate or shell (further considerations of drilling rotations that build on this core definition in the context of surfaces are reserved for brief remarks in the following Section II). Again, one can construct bending measures that are unchanged by such deformations.

One requirement is through-thickness uniformity of the stretch, in which different elements along a referentially-curved body may experience different rotations (while through-thickness uniformity of rotation is part of the presumed kinematics of the thin body). The other requirement is uniformity of the rotation field in every direction, in which the stretch may be nonuniform through the thickness of a referentially-curved body. These invariance requirements are not the same, but coincide for plates. Using either requirement leads to the identification of dilations (also referred to as scaling) of plate mid-surfaces as pure stretching. Note that the dilation is of the surface only—the material on either side must follow kinematics appropriate to a thin body, and this is what dictates its stretch, as detailed in Figure 2 of [1]. In a dilation, these kinematics result in variations in stretch through a body with a curved reference shape.

A simple bending measure that is dilation-invariant is the referential gradient of the (present) normal, a (rotated) product of the stretch and the present curvature. This object can be further modified by transposition, or rotation of one or more legs to/from the reference/present configuration, operations that are irrelevant to the construction of the isotropic invariants that will appear in associated bending energies. Therefore, to avoid excessive descriptors and modifiers in subsequent discussion, we will informally refer to tensors related by such operations as "equivalent", or refer to such equivalent tensors interchangeably, as for the present purposes they encode the same physical information of interest. An early example of such a tensor was given by Atluri [7], who did not consider its invariance properties. Such invariance was recognized as important by Ghiba and co-workers [5], and (a few months later) by the present authors, who considered a symmetrization of equivalent referential and present tensors [1, 2]. In planar bending, we can use the scalar mid-surface stretch λ and curvature κ to write the corresponding dilation-invariant bending measure $\lambda\kappa$ or, equivalently, the arc length derivative of the tangential angle. This measure and its invariance properties were introduced and recognized even earlier by Antman [8].

In the case of curved shells or rods, the requirements of through-thickness uniformity of stretch, or of a globally uniform rotation field, do not correspond. Ours is a structure-dependent definition of "pure stretching" of a particular surface, which does not correspond to a "pure stretch" of a continuum except in the special case of flat plates or straight rods. The use of the same term, "stretching", for both cases is thus an unfortunate distraction from the underlying concepts. Our coinage arose from the way the term "stretching" is informally used in studies of thin sheets in soft matter physics.

In the bending measures for shells that arise from one or the other of these requirements, the difference lies in what is being subtracted from the plate term, the product of stretch and present curvature. We subtract the product of stretch and referential curvature—for details, including the relations between various tensors mentioned above and below, see [1, Section 2] and [2, Section 5.2]. In planar bending, the measure would be written $\lambda(\kappa - \bar{\kappa})$. For a curved reference state ($\bar{\kappa} \neq 0$), these measures are not dilation-invariant. To our knowledge, such shell or curved rod measures did not appear in any prior work. They may be contrasted with a dilation-invariant measure that subtracts the referential curvature, which also appears in Antman [8] as a scalar and Atluri [7] as a tensor. In planar bending, this measure would be $\lambda\kappa - \bar{\kappa}$. This type of measure can be found in several other near-contemporary and subsequent works, including Acharya [3]. Further history may

be found in the references cited in [1, 2], noting however that this discussion missed not only [3] but another work [4], to be discussed shortly below. The two types of measures agree for flat plates and straight rods. However, a deformation that takes a body in a curved reference state and extends it at the same curvature, so that the stretch is uniform through the thickness, is considered a pure stretching in our definition, but a combination of stretching and bending in the other definition, as this process changes the normals attached to material points. Meanwhile, dilation of the mid-surface of a referentially-curved body is considered a combination of stretching and bending in our definition, but a pure stretching in the other definition.

The quantity in our measures, namely the product of the mid-surface stretch and the change in mid-surface curvature, also arises naturally as the additional (tangential) stretch of material located just off of the mid-surface. The manner in which this enters a reduction of a 3D quadratic-Biot-type bulk energy, *via* the through-thickness derivative of the deformation gradient, can be seen in Equations 27-31 of [2], noting that the coefficient α_1 appearing in these expressions is unity at lowest order in stretch. While a simple planar or axisymmetric version of this type of derivation was performed in several places for flat bodies [9–11], we are not aware of such a simple derivation for curved rods in the literature. Because of this apparent lacuna, we provide a sketch of this calculation below.

We note that nearly every introduction of Antman/Atluri-type curved rod or shell measures was a direct prescription, not derived through a reduction. However, in our discussion we overlooked a work of Ghiba and co-workers [4], who perform a reduction of a 3D quadratic-Biot-type energy but with additional degrees of freedom, and obtain shell measures akin to Atluri's (where a referential curvature is subtracted) rather than ours (where the product of stretch and referential curvature is subtracted). Their approach starts with a more complex material and kinematics, employs different assumptions, and follows different procedures, and we are unable to see how this bending measure appears in the midst of their derivation, leading to the disagreement. We note that some of the text within Figure 3 of [5] implies that they have not yet performed the reduction directly from a 3D Biot-type energy to a 2D Biot-type energy.

The appearance of our measure can be understood from the following informal and incomplete planar derivation in the style of [9, 11], or implicitly through computation of through-thickness-uniform stretches in [1], or explicitly and in full tensorial form in [2], a detailed and more careful approach built on [12, 13] that leads to the same results. Consider a thin body in the plane, with uniform thickness h , parameterized by material coordinates x and z in the tangential and normal directions, respectively. Let x be the arc length of the referential mid-line. The deformed mid-line has stretch λ , tangential angle θ , and scalar curvature $\kappa = d_x\theta/\lambda$. The corresponding referential quantities are $\bar{\theta}$ and $\bar{\kappa} = d_x\bar{\theta}$. The angular description is not necessary, but we include it for completeness.

The tangential stretch is the ratio of lengths of deformed and referential line elements in the x direction. Considering shear-free deformations for simplicity, this stretch varies with distance from the mid-surface approximately as

$$\frac{l(z)}{\bar{l}(z)} \approx \frac{l(0)[1 - zd_x\theta/\lambda]}{\bar{l}(0)[1 - zd_x\bar{\theta}]} = \lambda \frac{1 - z\kappa}{1 - z\bar{\kappa}} \approx \lambda [1 - z(\kappa - \bar{\kappa})]. \quad (1)$$

The first approximation corresponds to neglect of the Poisson effect in the z direction, which forms part of the classical Kirchhoff-Love kinematic assumptions and remains in their more consistent generalization [12], while the second approximation corresponds to small $z\kappa$ and $z\bar{\kappa}$. Thus, we already see that, to a first approximation, *the additional tangential stretch near the mid-surface is the product of the mid-surface stretch and the change in curvature*. This quantity will then appear in strains and energies of any order of stretch. Further kinematic assumptions allow the transverse

stretch to be related to the tangential stretch, so the energy can be written in terms of the latter only. Merely as an example, we may consider an energy quadratic in Biot strains, which are obtained by subtracting unity from (1); on the mid-line we define $\Delta = \lambda - 1$. This energy would be of the form

$$\mathcal{E} \sim \int dx \int_{-h/2}^{h/2} dz (1 - z\bar{\kappa}) [\Delta - z\lambda(\kappa - \bar{\kappa})]^2 , \quad (2)$$

whose leading-order terms for small strain and thickness are of the form $h\Delta^2$ and $h^3\lambda^2(\kappa - \bar{\kappa})^2$. Again, we cannot account for the discrepancy with the results of the reduction of Ghiba and co-workers [4]. The additional question of whether a symmetric or asymmetric tensorial measure might arise in a general derivation is addressed later in Section IV.

We believe that if such simple considerations had been taken into account, with or without additional consideration of through-thickness stretch uniformity, the bending measure $\lambda(\kappa - \bar{\kappa})$ might have appeared and been adopted earlier. Before we had performed either the stretch uniformity [1] or reduction [2] calculations, we were also ready to extend the property of dilation invariance to shells, misguided by its elegance, instead of allowing the geometry of the shell to break this symmetry.

II. FURTHER COMPARISONS AND COMMENTS

We have focused on the isotropic invariant content of certain bending measures, while others have introduced more stringent requirements that distinguish between tensors we refer to as “equivalent”, and have further delved into the significance of drilling rotations or other issues. We refer the reader to these works for details [3, 5, 6, 14, 15], and here only briefly summarize a few points.

Acharya introduced a restrictive requirement and several bending measures, one of which (Equation 9 of [3]) is equivalent to that of Atluri [7], where the plate measure appears in Equation 9.10 and the shell measure implicitly in a 2D Biot-like energy in Equation 9.25. Ghiba and co-workers [5] argue for relaxed and modified requirements, noting that Acharya’s requirements may be too stringent to include even his own measures, due to considerations related to drilling rotations in the reference configuration. One “novel invariance requirement” they suggest is dilation invariance for plates, but this concept, illustrated recently in Figure 6 of [5] and Figure 1 of [1], was actually introduced alongside Antman’s original measure [8]— see also Antman [16, Chapter 4] for further discussion of inflation of a ring and preservation of this bending measure within the context of a planar Cosserat rod. Ghiba and co-workers then suggest a “new bending tensor” for shells (Equation 5.9 or 6.2 of [5]), which is equivalent to Atluri’s and Acharya’s, although re-expressed using an additional “fictitious planar configuration”. This tensor is taken from the aforementioned reduction paper (Equation 5.5 of [4]). They compare this measure with Acharya’s work, but overlook the appropriate tensor for comparison (Equation 9 of [3]) and instead compare to other measures of Acharya. A short note by Acharya [14] points out this oversight. These Antman/Atluri-type measures satisfy the requirement of dilation-invariance, being equivalent to the referential gradient of the normal, or the product of stretch and curvature, minus the referential curvature. Acharya [14] writes that dilation invariance is “natural for shells as well”, surely because his core definition is equally applicable to bulk solids or any structure; a similar viewpoint seems present in a recent draft by Ghiba and co-workers [15], where they rename their extended Antman/Atluri-type bending measure the “change of curvature” tensor (which rather subtracts referential curvature from the product of stretch and curvature, not from the curvature). As outlined above, we instead consider a different non-dilation-invariant bending measure, one that features the product of stretch and the

change in curvature, arising from the form of the stretch near the mid-surface, and corresponding to a different, structure-dependent, invariance condition, to be “natural” for shells, although this measure takes the same dilation-invariant form for the special case of plates.

Virga [6] confines his treatment of bending measures to plates. He introduces a requirement that shares some of the restrictions of Acharya that Ghiba and co-workers relax; further comment can be found in [15]. Its restrictions distinguish between referential and present tensorial measures with the same invariants, although the invariants themselves are considered acceptable as scalar measures. He introduces a plate measure that satisfies his requirement; it is one of two possible “squares” of the Atluri-type dilation-invariant referential gradient of the normal $\bar{\nabla}\mathbf{n}$. While $(\bar{\nabla}\mathbf{n})^\top \cdot \bar{\nabla}\mathbf{n}$ satisfies his requirements, $\bar{\nabla}\mathbf{n} \cdot (\bar{\nabla}\mathbf{n})^\top$ does not, although they have the same invariants; their trace provides the continuum limit of the discrete Seung-Nelson energy. Working with either squared quantity, one misses additional invariant content at quadratic order in stretch that can be obtained by squaring the trace of an Atluri-type measure. However, this and its parent Atluri-type tensors do not satisfy certain of Virga’s requirements related to drilling rotations of a *deformed* plate [17]. For details of these, and similar quadratic-stretch invariants arising from reduction of a neo-Hookean energy or other quadratic-stretch energies, see Section 5.2.2 of [2]. While Virga is concerned with purity of bending, our focus has been on purity of stretching. We note that some mid-surface stretching operations will be detected by both our and Virga’s bending measures, and drilling operations (shear stretching) by our bending measures, but all only in the presence of a change in curvature, and this last is consistent with the idea that the bending content associated with changes in curvature may depend on tangential strains of a thin body. Extending or drilling at the reference curvature corresponds to through-thickness-uniform stretches, whereas extending or drilling at a different curvature does not, so the former will not contribute to our bending measure, but the latter will. In a body that has been deformed away from its reference curvature, particular additional combinations of extension and curvature change will not constitute bending—for a plate, this is dilation. Presumably, there are also particular combinations of drilling and curvature change that are “not bending”.

Our perspective considers the internal kinematics of the thin bodies, which are neither bulk continua nor surfaces, and thereby differs from the other authors’ use of a universal definition of a uniform rotation field augmented with a purely two-dimensional view of drilling rotations.

III. CORRECTION, WITH A COMMENT ON THE SIZE OF RELEVANT TERMS

In the papers [1, 2], we discuss a plate bending measure $\mathbf{J}\mathbf{I} = \text{sym}(\mathbf{V} \cdot \mathbf{b})$ formed by symmetrizing the asymmetric product of two symmetric tensors, the left stretch \mathbf{V} and curvature \mathbf{b} . In both papers, a similar incorrect statement is made in the discussion of the results, to the effect that the determinant of $\mathbf{J}\mathbf{I}$ is the product of determinants of \mathbf{V} and \mathbf{b} , which are respectively the Jacobian determinant J and Gaussian curvature K . This elementary mistake in linear algebra was never employed in any of the derivations, but contributed to incorrect conclusions about the geometric nature of certain terms in a bending energy, as expressed in two places in our texts.

Let \mathbf{A} and \mathbf{B} be second-order tensors. While it is true that $\text{Det}(\mathbf{A} \cdot \mathbf{B}) = \text{Det}(\mathbf{B} \cdot \mathbf{A}) = \text{Det } \mathbf{A} \text{ Det } \mathbf{B}$, this is not equal to $\text{Det}(\text{sym}(\mathbf{A} \cdot \mathbf{B}))$ unless \mathbf{A} and \mathbf{B} commute. For tensors of dimension two, the determinant of the sum and difference of tensors \mathbf{A} and \mathbf{B} can be expressed using (see [18,

Section 9] for several more general relevant expressions):

$$\begin{aligned}
\text{Det}(\mathbf{A} \pm \mathbf{B}) &= \text{Det } \mathbf{A} \text{ Det}(\mathbf{I} \pm \mathbf{A}^{-1} \cdot \mathbf{B}) \\
&= \text{Det } \mathbf{A} (1 \pm \text{Tr}(\mathbf{A}^{-1} \cdot \mathbf{B}) + \text{Det}(\mathbf{A}^{-1} \cdot \mathbf{B})) \\
&= \text{Det } \mathbf{A} \pm \text{Tr}(\text{Det}(\mathbf{A}) \mathbf{A}^{-1} \cdot \mathbf{B}) + \text{Det } \mathbf{B}.
\end{aligned} \tag{3}$$

Alternately, we may use the relations [17]

$$\begin{aligned}
2\text{Det}(\mathbf{A} \pm \mathbf{B}) &= (\text{Tr}(\mathbf{A} \pm \mathbf{B}))^2 - \text{Tr}((\mathbf{A} \pm \mathbf{B}) \cdot (\mathbf{A} \pm \mathbf{B})) \\
&= (\text{Tr } \mathbf{A})^2 \pm 2\text{Tr } \mathbf{A} \text{ Tr } \mathbf{B} + (\text{Tr } \mathbf{B})^2 - \text{Tr}(\mathbf{A} \cdot \mathbf{A}) \mp 2\text{Tr}(\mathbf{A} \cdot \mathbf{B}) - \text{Tr}(\mathbf{B} \cdot \mathbf{B}) \\
&= 2\text{Det } \mathbf{A} + 2\text{Det } \mathbf{B} \pm 2\text{Tr } \mathbf{A} \text{ Tr } \mathbf{B} \mp 2\text{Tr}(\mathbf{A} \cdot \mathbf{B}).
\end{aligned} \tag{4}$$

Combining either pair of expressions, we find that

$$\text{Det}(\mathbf{A} + \mathbf{B}) + \text{Det}(\mathbf{A} - \mathbf{B}) = 2(\text{Det } \mathbf{A} + \text{Det } \mathbf{B}), \tag{5}$$

and, therefore, Det commutes with the decomposition into symmetric and antisymmetric parts,

$$\text{Det}(\text{sym}(\mathbf{A} \cdot \mathbf{B})) + \text{Det}(\text{skew}(\mathbf{A} \cdot \mathbf{B})) = 2(\text{Det}_{\frac{1}{2}}(\mathbf{A} \cdot \mathbf{B}) + \text{Det}_{\frac{1}{2}}(\mathbf{B} \cdot \mathbf{A})) = \text{Det}(\mathbf{A} \cdot \mathbf{B}) = \text{Det } \mathbf{A} \text{ Det } \mathbf{B}. \tag{6}$$

For the particular case at hand, employing the definition [19] of the Bell strain $\mathbf{E}_{\text{Bell}} = \mathbf{V} - \mathbf{I}$, we have

$$\begin{aligned}
\text{Det } \mathbf{J}\mathbf{I} &= \text{Det } \mathbf{V} \text{ Det } \mathbf{b} - \text{Det}(\text{skew}(\mathbf{V} \cdot \mathbf{b})) \\
&= JK - \text{Det}(\text{skew}(\mathbf{E}_{\text{Bell}} \cdot \mathbf{b}))
\end{aligned} \tag{7}$$

The second term only vanishes if the principal axes of stretch and curvature align (if these two symmetric tensors commute, they are coaxial).

The relevant errors are those referring to the determinant of the plate bending measure $\mathbf{J}\mathbf{I}$ and its referential-area-weighted integral: in [1], Section 2, the paragraph following the paragraph containing Equation 4, and in [2], Section 5.2.1, the final paragraph.

The determinant of any symmetric plate bending measure linear in curvature will contain K , which is related to the derivatives of the metric. The referential \bar{K} of a plate is zero. Changes in metric, and thus K , scale as stretch squared minus one, and are thus linear in Δ , where Δ is an eigenvalue of the Bell strain (or equivalently the more common Biot strain that employs the right stretch). The first term in (7) is $O(\Delta)$ while the second is $O(\Delta^2)$. This determinant of the antisymmetric piece can be compared with the terms coming from the square of the trace of the symmetric bending measure, which will be $O(1)$ with $O(\Delta)$ corrections. These terms can be considered in the context of an energy density that is an expansion in small thickness h and small strains Δ , with a stretching term of $O(h\Delta^2)$ and a bending term of $O(h^3B^2)$ (where B is some suitable eigenvalue of the bending measure), in which we are already dropping terms coming from Poisson coupling to bending of $O(h^3\Delta B^2)$, among terms of various other orders. In shells, which provide additional complications, we also often ignore terms of the same order coming from the change in area off the mid-surface, of the form $O(h^3\Delta\bar{H}B)$ (with \bar{H} the referential mean curvature).

All of this is to say that arguments might be made to ignore some terms for small strain theories, should this be desirable for other reasons. It is worth remembering that the distinctions being made between bending measures are small in magnitude for small-strain theories, their importance being primarily in keeping certain natural objects intact, and in producing a certain qualitative response.

IV. REMARKS ON ANISYMMETRIC CONTRIBUTIONS TO THE ENERGY

In our work [1, 2], we construct the most general isotropic quadratic energy for symmetric tensors such as the plate measure $\mathbf{J}\mathbf{I} = \text{sym}(\mathbf{V} \cdot \mathbf{b})$, which includes $(\text{Tr } \mathbf{J}\mathbf{I})^2$ and $\text{Tr}(\mathbf{J}\mathbf{I} \cdot \mathbf{J}\mathbf{I})$, or equivalently $(\text{Tr } \mathbf{J}\mathbf{I})^2$ and $\text{Det } \mathbf{J}\mathbf{I}$. We also derive such an energy by reduction from a symmetric 3D strain [2]. However, the latter process requires care to preserve the symmetry of the measure. The symmetric stretch can be equivalently expressed in two different ways, using the rotation and deformation gradient or their transposes. We take derivatives of this quantity through the thickness of the body, and neglect as higher-order the derivative of the rotation (see Equation 5 and the text in the short paragraph spanning pages 587-588; to see how this strain measure enters the energy, the relevant Equations are 18-20, 23-24). This process can potentially result in the spurious introduction of asymmetry into this quantity, unless it is expressed symmetrically, as is done in Equation 31 of [2].

If we, instead, directly begin with an asymmetric tensor, the most general isotropic quadratic energy has three parameters (here λ, μ, α), as the tensor can be “squared” with itself or with its transpose. The asymmetric tensor $\mathbf{V} \cdot \mathbf{b}$, the rotated referential gradient of the normal that appears in some manner in all of the plate bending tensors discussed in this note, has some additional structure, as it is formed from two symmetric tensors \mathbf{V} and \mathbf{b} (so that $(\mathbf{V} \cdot \mathbf{b})^\top = \mathbf{b} \cdot \mathbf{V}$). In this case, the most general form of an isotropic quadratic energy is, after arranging to isolate the antisymmetric piece, and further rewriting compactly using $\mathbf{J}\mathbf{I} = \text{sym}(\mathbf{V} \cdot \mathbf{b})$ and its counterpart $\mathbf{Z} = \text{skew}(\mathbf{V} \cdot \mathbf{b})$,

$$\lambda(\text{Tr}(\mathbf{V} \cdot \mathbf{b}))^2 + \mu(\text{Tr}(\mathbf{V} \cdot \mathbf{b} \cdot \mathbf{b} \cdot \mathbf{V}) + \text{Tr}(\mathbf{V} \cdot \mathbf{b} \cdot \mathbf{V} \cdot \mathbf{b})) + \alpha(\text{Tr}(\mathbf{V} \cdot \mathbf{b} \cdot \mathbf{b} \cdot \mathbf{V}) - \text{Tr}(\mathbf{V} \cdot \mathbf{b} \cdot \mathbf{V} \cdot \mathbf{b})) \quad (8)$$

$$= \lambda(\text{Tr } \mathbf{J}\mathbf{I})^2 + 2\mu\text{Tr}(\mathbf{J}\mathbf{I} \cdot \mathbf{J}\mathbf{I}) - 2\alpha\text{Tr}(\mathbf{Z} \cdot \mathbf{Z}), \quad (9)$$

noting further the two-dimensional relations $\text{Tr}(\mathbf{J}\mathbf{I} \cdot \mathbf{J}\mathbf{I}) = (\text{Tr } \mathbf{J}\mathbf{I})^2 - 2\text{Det } \mathbf{J}\mathbf{I}$ and $-\text{Tr}(\mathbf{Z} \cdot \mathbf{Z}) = \text{Tr}(\mathbf{Z} \cdot \mathbf{Z}^\top) = 2\text{Det } \mathbf{Z}$. We remark that the original tensor $\mathbf{V} \cdot \mathbf{b}$ has two invariants, and it is possible to re-express the three terms in (9) in terms of these using (6), that is, $\text{Det } \mathbf{J}\mathbf{I} + \text{Det } \mathbf{Z} = \text{Det}(\mathbf{V} \cdot \mathbf{b})$.

What information is contained in the antisymmetric term? The determinant only provides a small, positive term of $O(\Delta^2)$; expressed in terms of Cartesian components it is $((V_{11} - V_{22})b_{12} - (b_{11} - b_{22})V_{12})^2$. This couples the deviatoric parts (difference of eigenvalues) of the stretch and curvature. Consider equal stretching and compression of a square element, resulting in pure shear along a diagonal, followed by bending into a saddle with equal-magnitude principal curvatures along the diagonals. The principal bending is along the unstretched (sheared) directions, and the principal stretching is along the unbent directions of the saddle. A positive α corresponds to a positive “interaction energy” between the two deformations. We reserve consideration of the physical meaning, origins, and permissibility of such terms, and their counterparts in the more complex case of shells, for another time.

ACKNOWLEDGMENTS

We are grateful to E. G. Virga for alerting us to the error corrected in Section III and for helpful input on this note, including suggesting the simple derivation (4). We also thank A. Acharya for discussion and clarification of his definitions.

- [1] E. Vitral and J. A. Hanna. Dilation-invariant bending of elastic plates, and broken symmetry in shells. *Journal of Elasticity*, 153(4-5):571–579, 2023.
- [2] E. Vitral and J. A. Hanna. Energies for elastic plates and shells from quadratic-stretch elasticity. *Journal of Elasticity*, 153(4-5):581–598, 2023.
- [3] A. Acharya. A nonlinear generalization of the Koiter–Sanders–Budiansky bending strain measure. *International Journal of Solids and Structures*, 37:5517–5528, 2000.
- [4] I.-D. Ghiba, M. Bîrsan, P. Lewintan, and P. Neff. The isotropic Cosserat shell model including terms up to $O(h^5)$. Part I: Derivation in matrix notation. *Journal of Elasticity*, 142:201–262, 2020.
- [5] I.-D. Ghiba, M. Bîrsan, P. Lewintan, and P. Neff. A constrained Cosserat shell model up to order $O(h^5)$: Modelling, existence of minimizers, relations to classical shell models and scaling invariance of the bending tensor. *Journal of Elasticity*, 146:83–141, 2021.
- [6] E. G. Virga. Pure measures of bending for soft plates. *Soft Matter*, 20:144, 2024.
- [7] S. N. Atluri. Alternate stress and conjugate strain measures, and mixed variational formulations involving rigid rotations, for computational analyses of finitely deformed solids, with application to plates and shells—I: Theory. *Computers & Structures*, 18(1):93–116, 1984.
- [8] S. Antman. General solutions for plane extensible elasticae having nonlinear stress-strain laws. *Quarterly of Applied Mathematics*, 26(1):35–47, 1968.
- [9] H. Irschik and J. Gerstmayr. A continuum mechanics based derivation of Reissner’s large-displacement finite-strain beam theory: the case of plane deformations of originally straight Bernoulli-Euler beams. *Acta Mechanica*, 206:1–21, 2009.
- [10] O. Oshri and H. Diamant. Strain tensor selection and the elastic theory of incompatible thin sheets. *Physical Review E*, 95:053003, 2017.
- [11] H. G. Wood and J. A. Hanna. Contrasting bending energies from bulk elastic theories. *Soft Matter*, 15:2411–2417, 2019.
- [12] O. Ozenda and E. G. Virga. On the Kirchhoff-Love hypothesis (revised and vindicated). *Journal of Elasticity*, 143:359–384, 2021.
- [13] D. J. Steigmann. Koiter’s shell theory from the perspective of three-dimensional nonlinear elasticity. *Journal of Elasticity*, 111(1):91–107, 2013.
- [14] A. Acharya. Mid-surface scaling invariance of some bending strain measures. *Journal of Elasticity*, doi.org/10.1007/s10659-024-10066-9, 2024.
- [15] I.-D. Ghiba, P. Lewintan, A. Sky, and P. Neff. An essay on deformation measures in isotropic thin shell theories. Bending versus curvature. [arXiv:2312.10928].
- [16] S. S. Antman. *Nonlinear Problems of Elasticity*. Springer, New York, second edition, 2005.
- [17] E. G. Virga. Personal communication, 2024.
- [18] C. Truesdell and W. Noll. *The Non-linear Field Theories of Mechanics*. Springer, second edition, 1992.
- [19] E. Vitral and J. A. Hanna. Quadratic-stretch elasticity. *Mathematics and Mechanics of Solids*, 27(3):462–473, 2022.