

Integrable sigma model with generalized \mathcal{F} structure, Yang-Baxter sigma model with generalized complex structure and multi-Yang-Baxter sigma model

A. Rezaei-Aghdam* and A. Taghavi†

*Department of Physics, Faculty of Basic science, Azarbaijan Shahid Madani University
53714-161, Tabriz, Iran*

May 14, 2024

Abstract

We construct an integrable sigma model with a generalized \mathcal{F} structure, which involves a generalized Nijenhuis structure \mathcal{J} satisfying $\mathcal{J}^3 = -\mathcal{J}$. Utilizing the expression of the generalized complex structure on the metric Lie group manifold G in terms of operator relations on its Lie algebra \mathfrak{g} , we formulate a Yang-Baxter sigma model with a generalized complex structure. Additionally, we present multi-Yang-Baxter sigma models featuring two and three compatible Nijenhuis structures. Examples for each of these models are provided.

*e-mail: rezaei-a@azaruniv.ac.ir

†e-mail: amanetaghavi@gmail.com

1 Introduction

Two-dimensional integrable σ models and their deformations have garnered considerable attention over the past 45 years [1–3]. The integrable deformation of the principal chiral model on $SU(2)$ was first presented in [4–7]. Additionally, the Yang-Baxter deformation of the chiral model was introduced by Klimcik [8–10]. The Yang-Baxter sigma model relies on R-operators that satisfy the (modified) classical Yang-Baxter equation ((m)CYBE) [11–13]. The integrable sigma model on a Lie group with a complex structure was also studied in [14] as a special case of the Yang-Baxter sigma model [8]. Recently, Mohammadi proposed a deformation of WZW models using two invertible linear operators [15]. Utilizing the general method outlined in another work by Mohammadi [16], we have constructed an integrable sigma model on a general manifold, particularly on a Lie group with a complex structure, in our previous work [17]. Furthermore, employing this method, we have explored an integrable sigma model on Lie groups equipped with a generalized complex structure [17].

Following the methodology outlined in [15] and [16], and building upon our previous work [17], we aim to construct integrable sigma models on a Lie group with generalized \mathcal{F} structure [18] (a generalized Nijenhuis structure \mathcal{J} [19, 20] with $\mathcal{J}^3 = -\mathcal{J}$). By utilizing the expression of the generalized complex structure on a metric Lie group G in terms of operator relations on its Lie algebra \mathfrak{g} (as discussed in [17]), we will further develop the Yang-Baxter sigma model with generalized complex structure. Consequently, we will present the multi-Yang-Baxter sigma model. The plane of the paper is outlined as follows:

In Section 2, we will review the methods outlined in [15] and [16], as well as the concepts related to generalized complex structure [19, 20]. Then, in Section 3, we will present the integrable sigma model on a Lie group with generalized \mathcal{F} structure. We will illustrate this with two examples: one on the Heisenberg Lie group $H_4 \equiv A_{4,8}$ [21], and another on the Lie group $GL(2, R)$. In Section 4, by employing the expression of the generalized complex structure on a metric Lie group G in terms of operator relations on its Lie algebra (as discussed in our previous work [17]), we will construct the Yang-Baxter sigma model with generalized complex structure. An example on the Nappi-Witten Lie group ($\mathbf{A}_{4,10}$) [22] will be presented. Finally, in Section 5, we will introduce the multi-Yang-Baxter sigma model with two and three compatible Nijenhuis structures and provide corresponding examples.

2 Some Review on integrable sigma model and generalized complex structure

Before delving into our main discussions, let's provide a brief overview of the two methods proposed by Mohammadi for constructing integrable sigma models [15, 16]. Additionally, we'll briefly review the fundamental concepts and relations concerning generalized complex structure, as discussed in the literature [19, 20].

2.1 First Method

Consider the following sigma model action

$$S = \int_{\Sigma} dz d\bar{z} (G_{\mu\nu}(x) + B_{\mu\nu}(x)) \partial x^{\mu} \bar{\partial} x^{\nu}, \quad (1)$$

where $x^{\mu}(z, \bar{z})$ ($\mu = 1, 2, \dots, d$) are coordinates of d dimensional manifold M , such that $G_{\mu\nu}$ and $B_{\mu\nu}$ are invertible metric and anti-symmetric tensor fields on M ; the (z, \bar{z}) are coordinates of the world-sheet Σ . The equations of motion for this model have the following form [16]

$$\bar{\partial} \partial x^{\lambda} + \Omega^{\lambda}_{\mu\nu} \partial x^{\mu} \bar{\partial} x^{\nu} = 0, \quad (2)$$

where

$$\Omega^{\lambda}_{\mu\nu} = \Gamma^{\lambda}_{\mu\nu} - H^{\lambda}_{\mu\nu}, \quad (3)$$

such that $\Gamma^{\lambda}_{\mu\nu}$ is the Christoffel coefficients and components of the torsion are given by

$$H^{\lambda}_{\mu\nu} = \frac{1}{2} G^{\lambda\eta} (\partial_{\eta} B_{\mu\nu} + \partial_{\nu} B_{\eta\mu} + \partial_{\mu} B_{\nu\eta}). \quad (4)$$

As in [16] one can construct a linear system whose consistency conditions are equivalence to the equations of motion (2) as follows:

$$[\partial + \partial x^\mu \alpha_\mu(x)]\psi = 0, \quad (5)$$

$$[\bar{\partial} + \bar{\partial} x^\mu \beta_\mu(x)]\psi = 0, \quad (6)$$

where the matrices α_μ and β_μ are functions of coordinates x^μ . The compatibility condition of this linear system yields the equations of motion, provided that the matrices $\alpha_\mu(x)$ and $\beta_\mu(x)$ must satisfies the following relations [16]

$$\beta_\mu - \alpha_\mu = \mu_\mu, \quad (7)$$

$$\partial_\mu \beta_\nu - \partial_\nu \alpha_\mu + [\alpha_\mu, \beta_\nu] = \Omega^\lambda_{\mu\nu} \mu_\lambda, \quad (8)$$

such that the second equation of the above set can then be rewritten as

$$F_{\mu\nu} = -(\nabla_\mu \mu_\nu - \Omega^\lambda_{\mu\nu} \mu_\lambda), \quad (9)$$

where the field strength $F_{\mu\nu}$ and covariant derivative corresponding to the matrices α_μ are given as follows:

$$F_{\mu\nu} = \partial_\mu \alpha_\nu - \partial_\nu \alpha_\mu + [\alpha_\mu, \alpha_\nu], \quad \nabla_\mu X = \partial_\mu X + [\alpha_\mu, X]. \quad (10)$$

Then by splitting symmetric and anti-symmetric parts of (9), we have [16]

$$0 = \nabla_\mu \mu_\nu + \nabla_\nu \mu_\mu - 2\Gamma^\lambda_{\mu\nu} \mu_\lambda, \quad (11)$$

$$F_{\mu\nu} = -\frac{1}{2}(\nabla_\mu \mu_\nu - \nabla_\nu \mu_\mu) - H^\lambda_{\mu\nu} \mu_\lambda. \quad (12)$$

In this manner the integrability condition of the sigma model (1) is equivalent to finding the matrices α_μ and μ_μ such that they satisfy in (9) (or (11) and (12)).

2.2 Second Method

In [15] Mohammadi give a general integrable deformation of chiral and WZW models. Here we review his results as the following proposition:

Proposition [15]: Consider invertible linear operators \mathcal{P}' and \mathcal{Q}' (such that $\mathcal{P}' + \mathcal{Q}'$ is also invertible) on a metric Lie algebra \mathfrak{g} (with ad invariant invertible metric¹ $\langle \cdot, \cdot \rangle_{\mathfrak{g}}$) with Lie group G , such that the operators satisfy the following relations [15]:

$$\forall X, Y \in \mathfrak{g}: \quad \langle \mathcal{P}'X, Y \rangle_{\mathfrak{g}} = \langle X, \mathcal{Q}'Y \rangle_{\mathfrak{g}} - 2\lambda \langle \mathcal{P}'X, \mathcal{Q}'Y \rangle_{\mathfrak{g}}, \quad (13)$$

$$[\mathcal{P}'X, \mathcal{Q}'Y] - \mathcal{P}'[X, \mathcal{Q}'Y] - \mathcal{Q}'[\mathcal{P}'X, Y] = \frac{\varepsilon}{2}(\mathcal{P}' + \mathcal{Q}')[X, Y], \quad (14)$$

with $\mathcal{P}'^{-1} = \mathcal{Q}'^{-1} + 2\lambda I$, where I is identity operator; λ and ε are parameters. Then the following two-dimensional sigma model is classically integrable [15]:

$$\begin{aligned} S(g) = & \lambda \int_{\partial B} dz d\bar{z} \langle g^{-1} \partial g, g^{-1} \bar{\partial} g \rangle_{\mathfrak{g}} + \frac{\lambda}{6} \int_B d^3 x \epsilon^{\mu\nu\rho} \langle g^{-1} \partial_\mu g, [g^{-1} \partial_\nu, g^{-1} \partial_\rho] \rangle_{\mathfrak{g}} \\ & + \int_{\partial B} dz d\bar{z} \langle g^{-1} \partial g, \mathcal{Q}'^{-1}(g^{-1} \bar{\partial} g) \rangle_{\mathfrak{g}}, \end{aligned} \quad (15)$$

where B is a three-dimensional manifold with boundary ∂B . Indeed the equations of motion of the above model are given as conservation of the following currents:

$$J = g(\mathcal{P}'^{-1}(g^{-1} \partial g))g^{-1},$$

¹The ad invariant metric is obtained as [22]

$$f^\gamma_{\alpha\beta} g_{\gamma\delta} + f^\gamma_{\alpha\delta} g_{\gamma\beta} = 0,$$

where the $f^\gamma_{\alpha\beta}$ are the structure constants of the Lie algebra \mathfrak{g} with bases $\{T_\alpha\}$, such that $[T_\alpha, T_\beta] = f^\gamma_{\alpha\beta} T_\gamma$.

$$\bar{J} = g(\mathcal{Q}'^{-1}(g^{-1}\bar{\partial}g))g^{-1}, \quad (16)$$

i.e.

$$\partial\bar{J} + \bar{\partial}J = 0. \quad (17)$$

Such that these currents satisfy the zero curvature condition

$$\partial\bar{J} - \bar{\partial}J + \varepsilon[J, \bar{J}] = 0, \quad (18)$$

if \mathcal{P}' and \mathcal{Q}' satisfy relation (14) [15].

2.3 Review on generalized complex structure

The generalized complex structure [19, 20] \mathcal{J} on a manifold M with even dimension is an endomorphism from $\mathcal{J} : TM \oplus T^*M \rightarrow TM \oplus T^*M$, such that this structure is invariant with respect to inner product \langle , \rangle on $TM \oplus T^*M$:

$$\forall X, Y \in TM, \xi, \eta \in T^*M : \quad \langle \mathcal{J}(X + \xi), \mathcal{J}(Y + \eta) \rangle = \langle X + \xi, Y + \eta \rangle, \quad (19)$$

and

$$\mathcal{J}^2 = -1, \quad (20)$$

also the generalized Nijenhuis tensor with Courant bracket [23]

$$[X + \xi, Y + \eta]_C = [X, Y] + L_X\eta - L_Y\xi - \frac{1}{2}d_M[i_X\eta - i_Y\xi], \quad (21)$$

is zero [19, 20] i.e.

$$N_{\mathcal{J}}(X + \xi, Y + \eta) = [X + \xi, Y + \eta]_C + \mathcal{J}[X + \xi, \mathcal{J}(Y + \eta)]_C + \mathcal{J}[\mathcal{J}(X + \xi), (Y + \eta)]_C - [\mathcal{J}(X + \xi), \mathcal{J}(Y + \eta)]_C = 0. \quad (22)$$

Utilizing the generalized complex structure \mathcal{J} in the following block form [23]

$$\mathcal{J} = \begin{pmatrix} J & P \\ Q & -J^* \end{pmatrix}, \quad (23)$$

where $J \in C^\infty(TM \otimes T^*M)$, $P \in C^\infty(\wedge^2 TM)$, $Q \in C^\infty(\wedge^2 T^*M)$; then in the coordinate representation $\{x^\mu\}$ of the manifold M we have $J = J^\mu{}_\nu \partial_\mu \otimes dx^\nu$, $P = P^{\mu\nu} \partial_\mu \wedge \partial_\nu$ and $Q = Q_{\mu\nu} dx^\mu \wedge dx^\nu$ such that by applying the above block form (23) in (20), we have the following relations for tensors J , P and Q :

$$P^{\nu k} + P^{k\nu} = 0, \quad Q_{\nu k} + Q_{k\nu} = 0, \quad (24)$$

$$J^\nu{}_\mu J^\mu{}_k + P^{\nu\mu} Q_{\mu k} + \delta^\nu{}_k = 0, \quad (25)$$

$$J^\nu{}_\mu P^{\mu k} + J^k{}_\mu P^{\mu\nu} = 0, \quad (26)$$

$$Q_{\nu\mu} J^\mu{}_k + Q_{k\mu} J^\mu{}_\nu = 0. \quad (27)$$

Furthermore, relation (22) (zero of the generalized Nijenhuis tensor) leads to the following relations [24]

$$\mathcal{A}^{\nu k\mu} = P^{\nu\lambda} \partial_\lambda P^{k\mu} + P^{k\lambda} \partial_\lambda P^{\mu\nu} + P^{\mu\lambda} \partial_\lambda P^{\nu k} = 0, \quad (28)$$

$$\mathcal{B}^{k\mu}{}_\nu = J^\lambda{}_\nu \partial_\lambda P^{k\mu} + P^{k\lambda} (\partial_\nu J^\mu{}_\lambda - \partial_\lambda J^\mu{}_\nu) - P^{\mu\lambda} (\partial_\nu J^k{}_\lambda - \partial_\lambda J^k{}_\nu) - \partial_\nu (J^k{}_\lambda P^{\lambda\mu}) = 0, \quad (29)$$

$$\mathcal{C}_{\nu k}{}^\mu = J^\lambda{}_\nu \partial_\lambda J^\mu{}_k - J^\lambda{}_k \partial_\lambda J^\mu{}_\nu - J^\mu{}_\lambda \partial_\nu J^\lambda{}_k + J^\mu{}_\lambda \partial_k J^\lambda{}_\nu + P^{\mu\lambda} (\partial_\lambda Q_{\nu k} + \partial_\nu Q_{k\lambda} + \partial_k Q_{\lambda\nu}) = 0, \quad (30)$$

$$\begin{aligned} \mathcal{D}_{\nu k\mu} = & J^\lambda{}_\nu (\partial_\lambda Q_{k\mu} + \partial_k Q_{\mu\lambda} + \partial_\mu Q_{\lambda k}) + J^\lambda{}_k (\partial_\lambda Q_{\mu\nu} + \partial_\mu Q_{\nu\lambda} + \partial_\nu Q_{\lambda\mu}) + J^\lambda{}_\mu (\partial_\lambda Q_{\nu k} + \partial_\nu Q_{k\lambda} + \partial_k Q_{\lambda\nu}) \\ & - \partial_\nu (Q_{k\lambda} J^\lambda{}_\mu) - \partial_k (Q_{\mu\lambda} J^\lambda{}_\nu) - \partial_\mu (Q_{\nu\lambda} J^\lambda{}_k) = 0. \end{aligned} \quad (31)$$

So the components of a generalized complex structure (23) on a manifold M must be satisfy relations (24-31).

3 Integrable sigma model with generalized \mathcal{F} structure

In our previous work [17], we constructed integrable sigma models with generalized complex structures, i.e., structures that satisfy relations (24-31). Here, we will consider a generalized \mathcal{F} structure, i.e., instead of condition (20) (or (24-27)), we consider the \mathcal{F} structure [18]

$$\mathcal{J}^3 + \mathcal{J} = 0. \quad (32)$$

So by using of block form (23) in the coordinate representation, we have the following conditions (instead of (24-27)):

$$P^{\mu\lambda}Q_{\lambda\rho}J^\rho{}_\nu + J^\mu{}_\lambda P^{\lambda\rho}Q_{\rho\nu} - P^{\mu\lambda}J^\rho{}_\lambda Q_{\rho\nu} + J^\mu{}_\lambda J^\lambda{}_\rho J^\rho{}_\nu + J^\mu{}_\nu = 0, \quad (33)$$

$$J^\mu{}_\lambda J^\lambda{}_\rho P^{\rho\nu} + P^{\mu\lambda}Q_{\lambda\rho}P^{\rho\nu} + P^{\mu\lambda}J^\rho{}_\lambda J^\nu{}_\rho - J^\mu{}_\lambda P^{\lambda\rho}J^\nu{}_\rho + P^{\mu\nu} = 0, \quad (34)$$

$$Q_{\mu\lambda}J^\lambda{}_\rho J^\rho{}_\nu - J^\lambda{}_\mu Q_{\lambda\rho}J^\rho{}_\nu + Q_{\mu\lambda}P^{\lambda\rho}Q_{\rho\nu} + J^\lambda{}_\mu J^\rho{}_\lambda Q_{\rho\nu} + Q_{\mu\nu} = 0. \quad (35)$$

Now for the case that M is a Lie group G (with ad invariant metric) by using the veilbein formalism $\forall g \in G, g^{-1}\partial g = (g^{-1}\partial g)^\alpha T_\alpha = e^\alpha{}_\mu \partial x^\mu T_\alpha$ with inverse $e_\beta{}^\mu e^\alpha{}_\mu = \delta^\alpha{}_\beta$ [25] and

$$J^\mu{}_\nu = e_\alpha{}^\mu J^\alpha{}_\beta e^\beta{}_\nu, \quad P^{\mu\nu} = e_\alpha{}^\mu P^{\alpha\beta} e_\beta{}^\nu, \quad Q_{\mu\nu} = e^\alpha{}_\mu Q_{\alpha\beta} e^\beta{}_\nu, \quad (36)$$

the above relations can be rewritten as follows:

$$P^{\alpha\delta}Q_{\delta\sigma}J^\sigma{}_\beta + J^\alpha{}_\delta P^{\delta\sigma}Q_{\sigma\beta} - P^{\alpha\delta}J^\sigma{}_\delta Q_{\sigma\beta} + J^\alpha{}_\delta J^\delta{}_\sigma J^\sigma{}_\beta + J^\alpha{}_\beta = 0, \quad (37)$$

$$J^\alpha{}_\delta J^\delta{}_\sigma P^{\sigma\beta} + P^{\alpha\delta}Q_{\delta\sigma}P^{\sigma\beta} + P^{\alpha\delta}J^\sigma{}_\delta J^\beta{}_\sigma - J^\alpha{}_\delta P^{\delta\sigma}J^\beta{}_\sigma + P^{\alpha\beta} = 0, \quad (38)$$

$$Q_{\alpha\delta}J^\delta{}_\sigma J^\sigma{}_\beta - J^\delta{}_\alpha Q_{\delta\sigma}J^\sigma{}_\beta + Q_{\alpha\delta}P^{\delta\sigma}Q_{\sigma\beta} + J^\delta{}_\alpha J^\sigma{}_\delta Q_{\sigma\beta} + Q_{\alpha\beta} = 0, \quad (39)$$

Furthermore, for ease of computation, we assume that J , P , and Q satisfy relations (26) and (27). Therefore, we will consider (26) and (27) in the following algebraic forms [17]

$$J^\alpha{}_\delta P^{\delta\beta} + J^\beta{}_\delta P^{\delta\alpha} = 0, \quad (40)$$

$$Q_{\alpha\delta}J^\delta{}_\beta + Q_{\beta\delta}J^\delta{}_\alpha = 0. \quad (41)$$

In the same way the relations (28-31) can be rewritten as follows [17]

$$A^{\alpha\beta\gamma} = f^\alpha{}_{\delta\sigma}P^{\beta\sigma}P^{\gamma\delta} + f^\gamma{}_{\delta\sigma}P^{\beta\delta}P^{\alpha\sigma} + f^\beta{}_{\delta\sigma}P^{\alpha\delta}P^{\gamma\sigma} = 0, \quad (42)$$

$$B^{\beta\gamma}{}_\alpha = f^\delta{}_{\sigma\alpha}P^{\beta\sigma}J^\gamma{}_\delta + f^\delta{}_{\alpha\sigma}P^{\gamma\sigma}J^\beta{}_\delta + f^\gamma{}_{\sigma\delta}P^{\beta\delta}J^\sigma{}_\alpha + f^\beta{}_{\sigma\delta}P^{\gamma\sigma}J^\delta{}_\alpha = 0, \quad (43)$$

$$\begin{aligned} C^{\alpha}{}_{\beta\gamma} = & -f^\delta{}_{\beta\gamma}J^\alpha{}_\sigma J^\sigma{}_\delta - f^\delta{}_{\beta\gamma}P^{\alpha\sigma}Q_{\sigma\delta} - f^\alpha{}_{\delta\sigma}J^\delta{}_\beta J^\sigma{}_\gamma - f^\delta{}_{\gamma\sigma}J^\alpha{}_\delta J^\sigma{}_\beta + f^\delta{}_{\beta\sigma}J^\alpha{}_\delta J^\sigma{}_\gamma \\ & + f^\delta{}_{\sigma\beta}P^{\alpha\sigma}Q_{\delta\gamma} + f^\delta{}_{\gamma\sigma}P^{\alpha\sigma}Q_{\delta\beta} = 0, \end{aligned} \quad (44)$$

$$D_{\alpha\beta\gamma} = f^\delta{}_{\alpha\sigma}J^\sigma{}_\beta Q_{\delta\gamma} + f^\delta{}_{\gamma\sigma}J^\sigma{}_\beta Q_{\alpha\delta} + f^\delta{}_{\gamma\sigma}J^\sigma{}_\alpha Q_{\delta\beta} + f^\delta{}_{\beta\sigma}J^\sigma{}_\alpha Q_{\gamma\delta} + f^\delta{}_{\beta\sigma}J^\sigma{}_\gamma Q_{\delta\alpha} + f^\delta{}_{\alpha\sigma}J^\sigma{}_\gamma Q_{\beta\delta} = 0, \quad (45)$$

Note that for obtaining relations (42-45) we have used the Maurer-Cartan relation [25]

$$f^\gamma{}_{\alpha\beta} = e^\gamma{}_\nu (e_\alpha{}^\mu \partial_\mu e_\beta{}^\nu - e_\beta{}^\mu \partial_\mu e_\alpha{}^\nu). \quad (46)$$

3.1 The model

Now, we will attempt to consider an integrable sigma model on the Lie group G with the aforementioned generalized \mathcal{F} structure, subject to conditions (37-45). We propose the action of the sigma model with generalized \mathcal{F} structure as follows:

$$S = \int (g^{-1}\partial g)^\alpha \{g_{\alpha\beta} + kg_{\alpha\delta}J_\beta^\delta + k'Q_{\alpha\beta} + k''g_{\alpha\delta}P^{\delta\sigma}g_{\sigma\beta} + k_1g_{\alpha\delta}J_\sigma^\delta J_\beta^\sigma + k_2g_{\alpha\delta}P^{\delta\sigma}Q_{\sigma\beta} \\ + k_3g_{\alpha\delta}J_\sigma^\delta P^{\sigma\gamma}g_{\gamma\beta} + k_4Q_{\alpha\delta}J_\beta^\delta\} (g^{-1}\partial g)^\beta dzd\bar{z}, \quad (47)$$

where $k, k', k'', k_1, \dots, k_4$ are real coupling constants. To obtain the condition under which the above model is integrable, one can utilize the first method mentioned in subsection 2.1 by introducing the matrices α_μ and μ_μ as follows:

$$\alpha_\mu = (\lambda_1 J^\alpha_\gamma + \lambda_2 P^{\alpha\delta}g_{\delta\gamma} + \lambda_3 g^{\alpha\delta}Q_{\delta\gamma} + \lambda_4 J^\alpha_\delta J^\delta_\gamma + \lambda_5 P^{\alpha\delta}Q_{\delta\gamma} + \lambda_6 J^\alpha_\delta P^{\delta\sigma}g_{\sigma\gamma} + \lambda_7 g^{\alpha\sigma}Q_{\sigma\delta}J^\delta_\gamma)e^\gamma_\mu T_\alpha, \\ \mu_\mu = (\lambda'_1 J^\alpha_\gamma + \lambda'_2 P^{\alpha\delta}g_{\delta\gamma} + \lambda'_3 g^{\alpha\delta}Q_{\delta\gamma} + \lambda'_4 J^\alpha_\delta J^\delta_\gamma + \lambda'_5 P^{\alpha\delta}Q_{\delta\gamma} + \lambda'_6 J^\alpha_\delta P^{\delta\sigma}g_{\sigma\gamma} + \lambda'_7 g^{\alpha\sigma}Q_{\sigma\delta}J^\delta_\gamma)e^\gamma_\mu T_\alpha, \quad (48)$$

here $(\lambda_1, \dots, \lambda_7, \lambda'_1, \dots, \lambda'_7)$ are constant parameters and T_α are the matrix representation of the bases of Lie algebra. Since this process is quite intricate, it's preferable to perform it for examples individually. Here, we will consider the model (47) for the Heisenberg Lie group $\mathbf{H}_4 \equiv \mathbf{A}_{4,8}$ and the Lie group $\mathbf{GL}(2, \mathbf{R})$.

3.2 Examples

a) Here, we construct the integrable sigma model with generalized \mathcal{F} structure (47) on the Heisenberg Lie group $\mathbf{H}_4 \equiv \mathbf{A}_{4,8}$ with the following commutation relations for its Lie algebra $A_{4,8}$ [21]:

$$[P_2, T] = P_2, \quad [P_2, J] = P_1, \quad [T, J] = J. \quad (49)$$

To calculate the vierbein we parameterize the corresponding Lie group \mathbf{H}_4 with coordinates $x^\mu = \{x, y, u, v\}$ by the g elements as follows

$$g = e^{xT_1} e^{yT_2} e^{vT_4} e^{uT_3}, \quad (50)$$

where we use the generators $T_\alpha = \{P_1, P_2, J, T\}$. Thus the vierbein and ad-invariant metric have the following form [12, 26]

$$(e^\alpha_\mu) = \begin{pmatrix} 1 & ue^v & 0 & 0 \\ 0 & e^v & 0 & 0 \\ 0 & 0 & 1 & u \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (g_{\alpha\beta}) = \begin{pmatrix} 0 & 0 & 0 & -k_0 \\ 0 & 0 & k_0 & 0 \\ 0 & k_0 & 0 & 0 \\ -k_0 & 0 & 0 & 0 \end{pmatrix}, \quad (51)$$

where $k_0 \in \mathbb{R} - \{0\}$ is an arbitrary real constant. Now, by utilizing the conditions for the generalized \mathcal{F} structure (37-39) and (40-41), along with its integrability conditions (42-45), one can obtain the following results:

$$J^\alpha_\beta = \begin{pmatrix} 0 & a & \frac{b(b-d)}{c} & \frac{ad}{c} \\ 0 & b-d & 0 & \frac{d(b-d)}{c} \\ 0 & 0 & b & 0 \\ 0 & c & 0 & d \end{pmatrix}, \quad Q_{\alpha\beta} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & e & 0 \\ 0 & -e & 0 & -\frac{de}{c} \\ 0 & 0 & \frac{de}{c} & 0 \end{pmatrix} \\ P^{\alpha\beta} = \frac{(b^2+1)(b-d)}{be} \begin{pmatrix} 0 & -\frac{(b-d)}{c} & \frac{a}{b-d} & -1 \\ \frac{(b-d)}{c} & 0 & 1 & 0 \\ -\frac{a}{b-d} & -1 & 0 & -\frac{c}{b-d} \\ 1 & 0 & \frac{c}{b-d} & 0 \end{pmatrix}, \quad (52)$$

where $a, d \in \mathbb{R}$ and $b, c, e \in \mathbb{R} - \{0\}$. Here, for simplicity, we will choose $a = 0$ and $d = 0$. Then, the action (47) of this model, after omitting surface terms, can be written as:²

$$S = \int dzd\bar{z} \{ -k_0(\partial v \bar{\partial} x + \partial x \bar{\partial} v) - \frac{k_0 c e^v}{2b} (bk + b^2 k_1 - k_2(b^2 + 1))(\partial y \bar{\partial} x + \partial x \bar{\partial} y) - \frac{k_0 c e^{2v}}{b} (bk + b^2 k_1 - k_2(b^2 + 1))u \partial y \bar{\partial} y \}$$

²Note that here and in the following examples, we have omitted the surface terms.

$$\begin{aligned}
& +k_0 e^v (1 + bk + b^2 k_1 - k_2(b^2 + 1))(\partial y \bar{\partial} u + \partial u \bar{\partial} y) + k_0 u e^v (bk + b^2 k_1 - k_2(b^2 + 1))(\partial y \bar{\partial} v + \partial v \bar{\partial} y) \\
& - \frac{k_0 b}{2c} (bk + b^2 k_1 - k_2(b^2 + 1))(\partial u \bar{\partial} v + \partial v \bar{\partial} u) - \frac{k_0 b u}{c} (bk + b^2 k_1 - k_2(b^2 + 1)) \partial v \bar{\partial} v \\
& + \frac{e^v}{e} (-k' e^2 + k_0^2 k''(1 + b^2) + k_0^2 k_3 b(1 + b^2) - k_4 b e^2)(\partial u \bar{\partial} y - \partial y \bar{\partial} u) + e(k' + k_4 b) u e^v (\partial y \bar{\partial} v - \partial v \bar{\partial} y) \\
& - \frac{k_0 c e^v}{2be} (kbe + 2k'' k_0(1 + b^2) + k_1 b^2 e - k_2 e(1 + b^2) + 2k_3 k_0 b(1 + b^2))(\partial x \bar{\partial} y - \partial y \bar{\partial} x). \tag{53}
\end{aligned}$$

The model (53) satisfy the integrability condition (9) if we set

$$\begin{aligned}
\lambda_1 &= -\frac{\lambda_4 b^2 - (1 + b^2) \lambda_5}{b}, \lambda_2 = -b \lambda_6 + k'' + k_3 b, \lambda_3 = -b \lambda_7, \\
\lambda'_1 &= -\frac{\lambda'_4 b^2 - (1 + b^2) \lambda'_5}{b}, \lambda'_2 = -b \lambda'_6 - 2(k'' + k_3 b), \lambda'_3 = -b \lambda'_7, \tag{54}
\end{aligned}$$

$$e = -k_0(1 + b^2)(k'' + b k_3), k_1 = \frac{-kb + k_2(1 + b^2)}{b^2}, \tag{55}$$

with $\lambda_4, \lambda_5, \lambda_6, \lambda_7, \lambda'_4, \lambda'_5, \lambda'_6, \lambda'_7$ are arbitrary constants and one of them can be consider as spectral parameters. Now by considering the WZW (Witten-Zumino-Witten) model on the Lie group $\mathbf{A}_{4,8}$ [26]:

$$S_{WZW_{A_{48}}} = k_0 \int dz d\bar{z} \{ -(\partial x \bar{\partial} v + \partial v \bar{\partial} x) + e^v (\partial y \bar{\partial} u + \partial u \bar{\partial} y) + \frac{e^v}{2} ((\partial u \bar{\partial} y - \partial y \bar{\partial} u) + u(\partial y \bar{\partial} v - \partial v \bar{\partial} y)) \}, \tag{56}$$

and assuming $bk + b^2 k_1 - k_2(b^2 + 1) = 0$, $\frac{1}{e}(k_0^2 k''(1 + b^2) + k_0^2 k_3 b(1 + b^2)) = k_0$ and $e(k' + k_4 b) = \frac{k_0}{2}$; then the action (53) can be considered as a following integrable deformation of WZW action

$$S = S_{WZW_{A_{48}}} + \int dz d\bar{z} \{ -\frac{3k_0 c e^v}{2b} (\partial x \bar{\partial} y - \partial y \bar{\partial} x) \}. \tag{57}$$

b) As second example, we construct integrable sigma model with generalised \mathcal{F} structure on the $\mathbf{GL}(2, \mathbf{R})$ Lie group. By considering the following commutation relations between the generators $\{T_1, T_2, T_3, T_4\}$ [13, 22]

$$[T_1, T_2] = 2T_2, \quad [T_1, T_3] = -2T_3, \quad [T_2, T_3] = T_1, \quad [T_4, \dots] = 0, \tag{58}$$

where T_4 is a central generator, and parameterize the corresponding Lie group $\mathbf{GL}(2, \mathbf{R})$ with coordinates $x^\mu = \{x, y, u, v\}$ by the g elements as

$$g = e^{xT_1} e^{yT_2} e^{uT_3} e^{vT_4}, \tag{59}$$

then the veirbein and ad-invariant metric are given by [13]

$$(e^\alpha{}_\mu) = \begin{pmatrix} 1 & ue^{-2x} & 0 & 0 \\ 0 & e^{-2x} & 0 & 0 \\ -2u & -u^2 e^{-2x} & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (g_{\alpha\beta}) = \begin{pmatrix} 2k_0 & 0 & 0 & 0 \\ 0 & 0 & k_0 & 0 \\ 0 & k_0 & 0 & 0 \\ 0 & 0 & 0 & d \end{pmatrix}, \tag{60}$$

where $k_0, d \in \mathbb{R} - \{0\}$. By using the conditions for the integrability of the generalized complex structure ((42)-(45)) and the relations (37-41) then the following components of generalized \mathcal{F} structure are obtained

$$\begin{aligned}
J^\alpha{}_\beta &= \begin{pmatrix} a & 0 & b & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad Q_{\alpha\beta} = \frac{(1 + a^2)}{ac} \begin{pmatrix} 0 & a & 0 & 0 \\ -a & 0 & -b & 0 \\ 0 & b & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\
P^{\alpha\beta} &= \begin{pmatrix} 0 & c & 0 & 0 \\ -c & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \tag{61}
\end{aligned}$$

where $a, c \in \mathbb{R} - \{0\}$, $b \in \mathbb{R}$. Here for simplicity we will choose $b = 0$; then one can construct sigma model (47), after omitting surfaces terms, as follows:

$$\begin{aligned}
S = \int dz d\bar{z} \{ & 2k_0(1 + ka + k_1a^2 - k_2(1 + a^2))\partial x \bar{\partial} x + e^{-2x}k_0(ka + k_1a^2 - k_2(1 + a^2))u(\partial x \bar{\partial} y + \partial y \bar{\partial} x) \\
& + k_0u^2e^{-4x}(ka + k_1a^2 - k_2(1 + a^2))\partial y \bar{\partial} y + \frac{k_0}{2}e^{-2x}(2 + ka + k_1a^2 - k_2(1 + a^2))(\partial u \bar{\partial} y + \partial y \bar{\partial} u) + d\partial v \bar{\partial} v \\
& + e^{-2x}(2k_0^2c(k'' + k_3a)u^2 - k_0(ka + k_1a^2 - k_2(1 + a^2))u)(\partial x \bar{\partial} y - \partial y \bar{\partial} x) \\
& + \frac{k_0}{2}e^{-2x}(-4uk_0c(k'' + k_3a) + ka + k_1a^2 - k_2(1 + a^2))(\partial u \bar{\partial} y - \partial y \bar{\partial} u)\}. \tag{62}
\end{aligned}$$

To investigate the integrability of this model, such that it satisfies the relation (9) (or (11) and (12)), we assume that α_μ and μ_μ have similar forms as the relation (48). Then, by setting:

$$\lambda_1 = -\frac{a^2\lambda_4 - (1 + a^2)\lambda_5}{a}, \lambda'_1 = -\frac{a^2\lambda'_4 - (1 + a^2)\lambda'_5}{a}, \lambda_2 = -a\lambda_6, \lambda_3 = -a\lambda_7, \lambda'_2 = -a\lambda'_6, \lambda'_3 = -a\lambda'_7, \tag{63}$$

such that $\lambda_4, \lambda_5, \lambda_6, \lambda_7, \lambda'_4, \lambda'_5, \lambda'_6, \lambda'_7$ are arbitrary constants (and one of them can be considered as spectral parameter), then this model satisfy the integrability condition (9).

4 Yang-Baxter sigma model with generalized complex structure

As mentioned in our previous work [17], the generalized complex structure relations on a metric Lie group (24-27) and (42-45) can be expressed as operator relations by considering the operators $J, \mathcal{P}, \mathcal{Q} : \mathfrak{g} \rightarrow \mathfrak{g}$ that act on the generators T_α of the Lie algebra \mathfrak{g} as follows:

$$J(T_\alpha) = J^\beta{}_\alpha T_\beta, \quad \mathcal{P}(T_\alpha) = \mathcal{P}^\beta{}_\alpha T_\beta, \quad \mathcal{Q}(T_\alpha) = \mathcal{Q}^\beta{}_\alpha T_\beta, \tag{64}$$

such that we define

$$\mathcal{P}^\beta{}_\alpha = g_{\alpha\gamma} P^{\gamma\beta}, \quad \mathcal{Q}^\beta{}_\alpha = Q_{\alpha\gamma} g^{\gamma\beta}, \tag{65}$$

where $g_{\alpha\beta} = \langle T_\alpha, T_\beta \rangle$ is an ad-invariant invertible metric on Lie algebra \mathfrak{g} . Then conditions (24-31) (on a metric Lie algebra \mathfrak{g}) can be rewritten as the following operator relations on \mathfrak{g} [17]:

$$\mathcal{P}^t = -\mathcal{P}, \quad \mathcal{Q}^t = -\mathcal{Q}, \tag{66}$$

$$J^2 + \mathcal{P}\mathcal{Q} + I = 0, \tag{67}$$

$$(J\mathcal{P})^t = -(J\mathcal{P}), \tag{68}$$

$$(J\mathcal{Q})^t = -(J\mathcal{Q}). \tag{69}$$

Furthermore relations (42-45) can be rewritten on the following operator relations [17]:

$\forall X, Y \in \mathfrak{g} :$

$$[\mathcal{P}(X), \mathcal{P}(Y)] - \mathcal{P}[X, \mathcal{P}(Y)] - \mathcal{P}[\mathcal{P}(X), Y] = 0, \tag{70}$$

$$[\mathcal{P}(X), J(Y)] + [J(X), \mathcal{P}(Y)] = -J^t([\mathcal{P}(X), Y] + [X, \mathcal{P}(Y)]), \tag{71}$$

$$[X, Y] - [J(X), J(Y)] + J[J(X), Y] + J[X, J(Y)] + \mathcal{P}[X, \mathcal{Q}(Y)] + \mathcal{P}[\mathcal{Q}(X), Y] = 0, \tag{72}$$

$$[J(X), \mathcal{Q}(Y)] + [\mathcal{Q}(X), J(Y)] + J^t([\mathcal{Q}(X), Y] + [X, \mathcal{Q}(Y)]) - \mathcal{Q}([J(X), Y] + [X, J(Y)]) = 0, \tag{73}$$

where transpose of an operator O is defined as

$$\forall X, Y \in \mathfrak{g}, \quad \langle OX, Y \rangle = \langle X, O^t Y \rangle. \tag{74}$$

In the following, for constructing the action (15), we assume that $\lambda = 0$, and \mathcal{P}' and \mathcal{Q}' are given with the following relations:³

$$\mathcal{Q}' = kI + \xi_1 J^t - \xi_2 \mathcal{Q} - \xi_3 \mathcal{P},$$

³Note that in the previous work [17], we have considered the model (15) with $\mathcal{Q}'^{-1} = kI + \xi_1 J^t - \xi_2 \mathcal{Q} - \xi_3 \mathcal{P}$ and $\lambda = 0$.

$$\mathcal{P}' = kI + \xi_1 J + \xi_2 \mathcal{Q} + \xi_3 \mathcal{P}, \quad (75)$$

where $\mathcal{P}' = \mathcal{Q}'^t$. Then by inserting (75) in the relation (14) and using (66-73) we must have

$$\begin{aligned} & \xi_2^2(\mathcal{Q}[X, \mathcal{Q}(Y)] + \mathcal{Q}[\mathcal{Q}(X), Y] - [\mathcal{Q}(X), \mathcal{Q}(Y)]) \\ & + \xi_2 \xi_3(\mathcal{Q}[X, \mathcal{P}(Y)] + \mathcal{Q}[\mathcal{P}(X), Y] - [\mathcal{Q}(X), \mathcal{P}(Y)] - [\mathcal{P}(X), \mathcal{Q}(Y)]) \\ & - \xi_1 \xi_3(\mathcal{P}[J(X), Y] + \mathcal{P}[X, J(Y)]) = 0. \end{aligned} \quad (76)$$

Here, we assume that $k\varepsilon - k^2 = \xi_1^2 = \xi_2 \xi_3$ and $J = -J^t$. Indeed, from (14), the relation (76) represents the integrability condition for the following Yang-Baxter sigma model [9] with a generalized complex structure (with three deformation parameters)

$$S = \int dz d\bar{z} \langle g^{-1} \partial g, \frac{1}{kI - \xi_1 J - \xi_2 \mathcal{Q} - \xi_3 \mathcal{P}}, g^{-1} \bar{\partial} g \rangle. \quad (77)$$

In the following we investigate an example for the model (77).

4.1 Example

c) As an example we construct the sigma model (77) on Nappi-Witten Lie group $\mathbf{A}_{4,10}$, with the following Lie algebra commutators [21, 22]

$$[J, P_i] = \varepsilon_{ij} P_j, \quad [P_i, P_j] = \varepsilon_{ij} T, \quad [T, J] = [T, P_i] = 0. \quad (78)$$

We parameterize the corresponding Lie group $\mathbf{A}_{4,10}$ with coordinates $x^\mu = \{v, a_1, a_2, u\}$ and consider the following parametrization for the group element [21]

$$g = e^{\Sigma a_i P_i} e^{uJ+vT}, \quad (79)$$

with the generators $T_\alpha = \{T, P_1, P_2, J\}$. The veirbein and ad-invariant metric is given by [22]:

$$(e^\alpha{}_\mu) = \begin{pmatrix} 1 & \frac{1}{2}a_2 & -\frac{1}{2}a_1 & 0 \\ 0 & \cos(u) & \sin(u) & 0 \\ 0 & -\sin(u) & \cos(u) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (g_{\alpha\beta}) = \begin{pmatrix} 0 & 0 & 0 & k_0 \\ 0 & k_0 & 0 & 0 \\ 0 & 0 & k_0 & 0 \\ k_0 & 0 & 0 & 0 \end{pmatrix}, \quad (80)$$

where $k_0 \in \mathbb{R} - \{0\}$. The corresponding components of the generalized complex structure (which satisfies the algebraic form of the relations (24-27) and (42-45)) are

$$(J^\alpha{}_\beta) = \begin{pmatrix} a & -ab & b & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & a \end{pmatrix}, \quad (Q_{\alpha\beta}) = \begin{pmatrix} 0 & 0 & 0 & 1+a^2 \\ 0 & 0 & 0 & -ba^2-b \\ 0 & 0 & 0 & 0 \\ -1-a^2 & ba^2+b & 0 & 0 \end{pmatrix}, \quad (P^{\alpha\beta}) = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}, \quad (81)$$

where a and b are real constants. We set $a = b = 0$ for satisfying $J = -J^t$, so the action (77) for this model is given by

$$\begin{aligned} S(g) = \int dz d\bar{z} \{ & \frac{kk_0^3}{(k^2 k_0^2 - (\xi_2 - \xi_3 k_0^2)^2)} ((\partial v \bar{\partial} u + \partial u \bar{\partial} v) + \frac{a_2}{2} (\partial a_1 \bar{\partial} u + \partial u \bar{\partial} a_1)) \\ & - \frac{a_1}{2} (\partial a_2 \bar{\partial} u + \partial u \bar{\partial} a_2)) + \frac{kk_0}{k^2 + \xi_1^2} (\partial a_2 \bar{\partial} a_2 + \partial a_1 \bar{\partial} a_1) + \\ & \frac{k_0^2 (\xi_2 - k_0^2 \xi_3)}{(k^2 k_0^2 - (\xi_2 - \xi_3 k_0^2)^2)} (\frac{a_2}{2} (\partial u \bar{\partial} a_1 - \partial a_1 \bar{\partial} u) + \frac{a_1}{2} (\partial a_2 \bar{\partial} u - \partial u \bar{\partial} a_2)) \}, \end{aligned} \quad (82)$$

where we omit the surfaces terms. This model satisfy the integrability condition (76).

5 Multi-Yang-Baxter sigma models

The model (77) and its integrability conditions (76) are motivated to consider Yang-Baxter sigma model with two and three (or multiple) compatible Nijenhuis structures⁴. If we consider \mathcal{Q}' and \mathcal{P}' in the following forms i.e. a two, three or in general a multi-parametric deformation of principal chiral model

$$\begin{aligned}\mathcal{Q}' &= kI + \xi_1 J_1 + \xi_2 J_2 + \xi_3 J_3, \\ \mathcal{P}' &= kI + \xi_1 J_1^t + \xi_2 J_2^t + \xi_3 J_3^t,\end{aligned}\tag{83}$$

then from (15) (with $\lambda = 0$) for the following multi-Yang-Baxter sigma model

$$S = \int dz d\bar{z} \langle g^{-1} \partial g, \frac{1}{kI + \xi_1 J_1 + \xi_2 J_2 + \xi_3 J_3}, g^{-1} \bar{\partial} g \rangle,\tag{84}$$

the condition (13) (by assuming $J_i^t = -J_i$ $i = 1, 2, 3$) can be written as

$$\forall X, Y \in \mathfrak{g} : \quad \langle (kI - \xi_1 J_1 - \xi_2 J_2 - \xi_3 J_3)X, Y \rangle_{\mathfrak{g}} = \langle X, (kI + \xi_1 J_1 + \xi_2 J_2 + \xi_3 J_3)Y \rangle_{\mathfrak{g}},\tag{85}$$

and integrability condition (14) result as the following relations

$$\omega_i[X, Y] - [J_i(X), J_i(Y)] + J_i[J_i(X), Y] + J_i[X, J_i(Y)] = 0, \quad \omega_i = 0, \pm 1, \quad i = 1, 2, 3,\tag{86}$$

$$\begin{aligned}(\omega_i + \omega_j)[X, Y] - [J_i(X), J_j(Y)] - [J_j(X), J_i(Y)] + J_i[J_j(X), Y] + J_j[J_i(X), Y] + \\ J_i[X, J_j(Y)] + J_j[X, J_i(Y)] = 0, \quad i, j = 1, 2, 3, \quad i \neq j,\end{aligned}\tag{87}$$

by assuming that

$$k(k + \varepsilon) + \omega_1 \xi_1^2 + \omega_2 \xi_2^2 + \omega_3 \xi_3^2 + (\omega_1 + \omega_2) \xi_1 \xi_2 + (\omega_1 + \omega_3) \xi_1 \xi_3 + (\omega_2 + \omega_3) \xi_2 \xi_3 = 0,\tag{88}$$

i.e. one can construct integrable multi-Yan-Baxter sigma model (84) with three compatible Nijenhuis structures (87) that satisfy in mCYBE (86). One can also write the multi-Yang-Baxter sigma model (84) with two J_i such that the conditions (86) and (87) are satisfied with $i, j = 1, 2$ and the conditions (88) is replaced by

$$k(k + \varepsilon) + \omega_1 \xi_1^2 + \omega_2 \xi_2^2 + (\omega_1 + \omega_2) \xi_1 \xi_2 = 0.\tag{89}$$

Indeed the relations (86)-(87) demonstrate that the two or three Nijenhuis structures are compatible with zero concomitant [27]. In this manner one can construct a multi-Yang-Baxter sigma model as multi-parametric Poisson-Lie deformation of principal sigma model with dual Lie algebra $\tilde{\mathfrak{g}} = \mathfrak{g}_{\xi_1 J_1 + \xi_2 J_2 + \xi_3 J_3}$ [9]; such that $\forall X, Y \in \mathfrak{g}$, $[X, Y]_{\xi_1 J_1 + \xi_2 J_2 + \xi_3 J_3} = \xi_1 [X, Y]_{J_1} + \xi_2 [X, Y]_{J_2} + \xi_3 [X, Y]_{J_3}$ where $[X, Y]_{J_i} = [J_i(X), Y] + [X, J_i(Y)]$. Relations (86-88) can also obtained from satisfying the combination of Nijenhuis structures $\xi_1 J_1 + \xi_2 J_2 + \xi_3 J_3$ in Yang-Baxter equation. Note that if we have $J_i^2 = -1$ (i.e. complex structure J_i) and have Hermitian structure then we have $N = 2$ structure and two of J_i are independent [27]. In the following we investigate two examples on $\mathbf{IX} + \mathbf{R}$ and $\mathbf{G}_{6,23}$ Lie groups.

5.1 Examples

d) As first example we consider the compatible Nijenhuis structures J_1, J_2 over $\mathbf{IX} + \mathbf{R}$ Lie group [28] with the following commutation relations for their Lie algebra

$$[T_2, T_3] = T_1, \quad [T_1, T_3] = -T_2, \quad [T_1, T_2] = T_3,\tag{90}$$

we parameterize the corresponding Lie group $\mathbf{IX} + \mathbf{R}$ with coordinates $x = \{x_1, x_2, x_3, x_4\}$ and consider the the group element $g = e^{x_1 T_1} e^{x_2 T_2} e^{x_3 T_3} e^{x_4 T_4}$. Then the veirbein and ad-invariant metric are given by:

$$(e^\alpha{}_\mu) = \begin{pmatrix} \cos x_3 \cos x_2 & \sin x_3 & 0 & 0 \\ -\sin x_3 \cos x_2 & \cos x_3 & 0 & 0 \\ \sin x_2 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (g_{\alpha\beta}) = \begin{pmatrix} k_0 & 0 & 0 & 0 \\ 0 & k_0 & 0 & 0 \\ 0 & 0 & k_0 & 0 \\ 0 & 0 & 0 & k_0 \end{pmatrix},\tag{91}$$

⁴Note that our model is different from bi-Yang-Baxter sigma model where presented in [9] with Nijenhuis structures J and $J_g = Ad_{g^{-1}} J Ad_g$ with $g \in G$.

where $k_0 \in \mathbb{R} - \{0\}$. The compatible Nijenhuis structures J_1, J_2 that satisfies the conditions (86) and (87) with $\omega_1 = \omega_2 = 1$ are given as follows

$$(J_1^\alpha)_\beta = \begin{pmatrix} 0 & 0 & 0 & a \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -a & 0 & 0 & 0 \end{pmatrix}, \quad (J_2^\alpha)_\beta = \begin{pmatrix} 0 & 0 & 0 & b \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -b & 0 & 0 & 0 \end{pmatrix}, \quad (92)$$

where $a \neq b$; then the action (84) can be obtained as

$$\begin{aligned} S(g) = \int dz d\bar{z} \{ & \frac{kk_0}{k^2 + (\xi_1 + \xi_2)^2} ((1 - \cos^2 x_2 \cos^2 x_3) \partial x_1 \bar{\partial} x_1 + \cos^2 x_3 \partial x_2 \bar{\partial} x_2 + \partial x_3 \bar{\partial} x_3 + \partial x_4 \bar{\partial} x_4 \\ & - \sin x_3 \cos x_2 \cos x_3 (\partial x_1 \bar{\partial} x_2 + \partial x_2 \bar{\partial} x_1) + \sin x_2 (\partial x_1 \bar{\partial} x_3 + \partial x_3 \bar{\partial} x_1)) \\ & + \frac{kk_0}{k^2 + (a\xi_1 + b\xi_2)^2} (\cos^2 x_2 \cos^2 x_3 \partial x_1 \bar{\partial} x_1 + \sin^2 x_3 \partial x_2 \bar{\partial} x_2 + \partial x_4 \bar{\partial} x_4 + \sin x_3 \cos x_2 \cos x_3 (\partial x_1 \bar{\partial} x_2 + \partial x_2 \bar{\partial} x_1)) \\ & - \frac{k_0(\xi_1 + \xi_2)}{k^2 + (\xi_1 + \xi_2)^2} (\sin x_3 \cos x_2 (\partial x_3 \bar{\partial} x_1 - \partial x_1 \bar{\partial} x_3) + \sin x_2 \cos x_3 (\partial x_2 \bar{\partial} x_1 - \partial x_1 \bar{\partial} x_2)) \\ & + \frac{k_0(a\xi_1 + b\xi_2)}{k^2 + (a\xi_1 + b\xi_2)^2} (\cos x_2 \cos x_3 (\partial x_4 \bar{\partial} x_1 - \partial x_1 \bar{\partial} x_4) + \sin x_3 (\partial x_4 \bar{\partial} x_2 - \partial x_2 \bar{\partial} x_4)) \}. \end{aligned} \quad (93)$$

e) As another example, we construct (84) model on six-dimensional Lie group $\mathbf{G}_{6,23}$. The $g_{6,23}$ Lie algebra have the following commutations [29, 30]:

$$[T_2, T_3] = T_1, \quad [T_2, T_6] = T_3, \quad [T_3, T_6] = T_4. \quad (94)$$

Now using $l = e^{x_1 T_1} e^{x_2 T_2} e^{x_3 T_3} e^{x_4 T_4} e^{x_5 T_5} e^{x_6 T_6}$ as a Lie group element with coordinates $x^\mu = \{x_1, x_2, x_3, x_4, x_5, x_6\}$ and the generators $\{T_1, T_2, T_3, T_4, T_5, T_6\}$, the algebraic metric and veilbeins related to this Lie algebra are given as [30]:

$$(e^\alpha)_\mu = \begin{pmatrix} 1 & x_3 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & x_6 & 1 & 0 & 0 & 0 \\ 0 & \frac{1}{2}x_6^2 & x_6 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad (g_{\alpha\beta}) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & m_2 \\ 0 & m_1 & 0 & m_2 & m_3 & m_4 \\ 0 & 0 & -m_2 & 0 & 0 & 0 \\ 0 & m_2 & 0 & 0 & 0 & 0 \\ 0 & m_3 & 0 & 0 & m_5 & m_6 \\ m_2 & m_4 & 0 & 0 & m_6 & m_7 \end{pmatrix}, \quad (95)$$

where m_1, \dots, m_7 are arbitrary real parameters. By using equations (86) and (87) with $\omega_1 = \omega_2 = \omega_3 = 1$ three compatible complex structure J_1, J_2, J_3 obtain as follows:

$$\begin{aligned} (J_1^\alpha)_\beta &= \begin{pmatrix} 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & a & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -a & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad (J_2^\alpha)_\beta = \begin{pmatrix} 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & b & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -b & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \end{pmatrix}, \\ (J_3^\alpha)_\beta &= \begin{pmatrix} 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & c & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -c & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \end{pmatrix}, \end{aligned} \quad (96)$$

where $a \neq b \neq c$ are arbitrary real parameters. For simplicity we assume that $m_1 = m_3 = m_4 = m_6 = m_7 = 0$, thus our multi-Yang-Baxter integrable sigma model (84) can be expressed as:

$$\begin{aligned}
S(g) = \int dz d\bar{z} \{ & \frac{km_2}{k^2 + (\xi_1 + \xi_2 + \xi_3)^2} ((\partial x_1 \bar{\partial} x_6 + \partial x_6 \bar{\partial} x_1) + x_3 (\partial x_2 \bar{\partial} x_6 + \partial x_6 \bar{\partial} x_2) + x_6^2 \partial x_2 \bar{\partial} x_2 \\
& + x_6 (\partial x_3 \bar{\partial} x_2 + \partial x_2 \bar{\partial} x_3) + (\partial x_4 \bar{\partial} x_2 + \partial x_2 \bar{\partial} x_4)) - \frac{1}{k^2 + (a\xi_1 + b\xi_2 + c\xi_3)^2} (km_2 x_6 \partial x_2 \bar{\partial} x_2 \\
& + km_2 (\partial x_3 \bar{\partial} x_2 + \partial x_2 \bar{\partial} x_3) + x_6 (a\xi_1 + b\xi_2 + c\xi_3) (m_2 + m_5) (\partial x_5 \bar{\partial} x_2 + \partial x_2 \bar{\partial} x_5) \\
& + (a\xi_1 + b\xi_2 + c\xi_3) (m_2 + m_5) (\partial x_5 \bar{\partial} x_3 + \partial x_3 \bar{\partial} x_5)) + \frac{(a\xi_1 + b\xi_2 + c\xi_3) (m_2 - m_5)}{k^2 + (a\xi_1 + b\xi_2 + c\xi_3)^2} x_6 (\partial x_2 \bar{\partial} x_5 - \partial x_5 \bar{\partial} x_2) \}. \quad (97)
\end{aligned}$$

6 Conclusion

Following the methodology outlined in the previous work [17], we construct a new integrable sigma model with generalized \mathcal{F} structure. Additionally, utilizing the expression of the generalized complex structure on the metric Lie group in terms of operator relations on its Lie algebra, we construct the Yang-Baxter sigma model with generalized complex structure. Furthermore, we present the multi-Yang-Baxter sigma model with two or three compatible Nijenhuis structures. Investigating the Poisson-Lie symmetry of these models remains an open problem, as does their conformality up to one loop.

Acknowledgements

This work has been supported by the research vice chancellor of Azarbaijan Shahid Madani University under research fund No. 1402/231.

References

- [1] K. Pohlmeyer, *Integrable Hamiltonian systems and interactions through quadratic constraints*, Commun. Math. Phys. 46 (1976) 207.
- [2] V. G. Zakharov and A. V. Mikhailov, *Relativistically invariant two-dimensional models of field theory which are integrable by means of the inverse scattering problem method*, Sov. Phys. JETP 47 (1978) 1017.
- [3] H. Eichenherr and M. Forger, *On the dual symmetry of the non-linear sigma models*, Nucl. Phys. B 155 (1979) 381; H. Eichenherr, M. Forger, *More about non-linear sigma models on symmetric spaces*, Nucl. Phys. B 164 (1980) 528.
- [4] I. V. Cherednik, *Relativistically invariant quasiclassical limits of integrable two-dimensional quantum models*, Theo. Math. phys. 47 (1981) 422.
- [5] J. Balog, P. Forgacs, Z. Horvath and L. Palla, *A new family of $SU(2)$ symmetric integrable sigma models*, Phys. Lett. B 324 (1994) 403, [arXiv:hep-th/9307030](#).
- [6] V. A. Fateev, *The sigma model (dual) representation for a two-parameter family of integrable quantum field theories*, Nucl. Phys. B 473 (1996) 509.
- [7] N. Sochen, *Integrable generalized principal chiral models*, Phys. Lett. B 391 (1997) 374, [arXiv:hep-th/9607009](#).
- [8] C. Klimcik, *Yang-Baxter σ -models and dS/AdS T-duality*, JHEP 0212 (2003) 051, [arXiv:hep-th/0210095](#).
- [9] C. Klimcik, *On integrability of the Yang-Baxter σ -model*, J. Math. Phys. 50 (2009) 043508, [arXiv:0802.3518 \[hep-th\]](#).
- [10] C. Klimcik, *Integrability of the bi-Yang-Baxter sigma-model*, Lett. Math. Phys. 104 (2014) 1095, [arXiv:1402.2105 \[math-ph\]](#).

- [11] H. Kyono and K. Yoshida, *Yang–Baxter invariance of the Nappi–Witten model*, Nucl. Phys. B 905 (2016) 242.
- [12] A. Eghbali, T. Parvizi, A. Rezaei-Aghdam, *Yang-Baxter deformations of WZW model on the Heisenberg Lie group*, Nucl. Phys. B 967 (2021) 115423, [arXiv:2103.01646 \[hep-th\]](#).
- [13] A. Eghbali, T. Parvizi and A. Rezaei-Aghdam, *Yang-Baxter deformations of the $GL(2, R)$ WZW model and non-Abelian T-duality*, E. P. J. C 83 (2023) 917, [arXiv:2305.12187 \[hep-th\]](#).
- [14] D. Bykov, *Complex structures and zero-curvature equations for sigma-models*, Phys. Lett. B 760 (2016) 341, [arXiv:1605.01093 \[hep-th\]](#).
- [15] N. Mohammadi, *On some integrable deformations of the Wess-Zumino-Witten model*, Phys. Rev. D 104 (2021) 126028, [arXiv:2012.09753 \[hep-th\]](#).
- [16] N. Mohammadi, *On the geometry of classically integrable two-dimensional non-linear sigma models*, Nucl. Phys. B 839 (2010) 420, [arXiv:0806.0550 \[hep-th\]](#).
- [17] A. Rezaei-Aghdam and A. Taghavi, *Integrable sigma models with complex and generalized complex structures*, arXiv preprint (2023) [arXiv:2309.13413](#).
- [18] M. Kon and K. Yano, *Structures on manifolds*, Vol. 3, World scientific, 1985.
- [19] N. Hitchin, *Generalized Calabi–Yau manifolds*, Quart. J. Math. 54 (2003) 281, [arXiv:math/0209099 \[math.DG\]](#).
- [20] M. Gualtieri, *Generalized complex geometry*, Ann. Math. (2011) 75, [arXiv:math/0401221 \[math.DG\]](#).
- [21] J. Patera and P. Winternitz, *Subalgebras of real three and four-dimensional Lie algebras*, J. Math. Phys. 18 (1977) 1449.
- [22] C. R. Nappi and E. Witten, *A WZW model based on a non-semi-simple group*, Phys. Rev. Lett. 71 (1993) 3751, [arXiv:hep-th/9310112](#).
- [23] T. J. Courant, *Dirac manifolds*, Trans. Am. Math. Soc. 319 (1990) 631.
- [24] R. Zucchini, *A sigma model field theoretic realization of Hitchin’s generalized complex geometry*, JHEP 11 (2004) 045, [arXiv:hep-th/0409181](#).
- [25] M. Nakahara, *Geometry, topology and physics*, CRC press, 2018.
- [26] A. Eghbali and A. Rezaei-Aghdam, *Poisson Lie symmetry and D-branes in WZW model on the Heisenberg Lie group H_4* , Nucl. Phys. B 899 (2015) 165, [arXiv:1506.06233 \[hep-th\]](#).
- [27] P. S. Howe and G. Papadopoulos, *Further remarks on the geometry of two-dimensional non-linear σ -models*, Classic. Quant. Grav. 12 (1988) 1647.
- [28] M. Aali-Javanangrouh and A. Rezaei-Aghdam, *Algebraic Structures of $N=(4, 4)$ and $N=(8, 8)$ SUSY Sigma Models on Lie groups and SUSY WZW Models*, arXiv preprint (2014) [arXiv:1402.5600](#).
- [29] G. M. Mubarakzyanov, *Classification of solvable Lie algebras of sixth order with a non-nilpotent basis element*, Izv. Vyssh. Uchebn. Zaved. Mat. 4 (1963) 104.
- [30] A. Rezaei-Aghdam, M. Sephid and S. Fallahpour, *Automorphism group and ad-invariant metric on all six dimensional solvable real Lie algebras*, (2010), [arXiv:1009.0816 \[math-ph\]](#).