On the distance problem over finite p-adic rings

Thang Pham Boqing Xue

May 14, 2024

Abstract

In this paper, we study the distance problem in the setting of finite p-adic rings. In odd dimensions, our results are essentially sharp. In even dimensions, we clarify the conjecture and provide examples to support it. Surprisingly, compared to the finite field case, in this setting, we are able to provide a large family of sets such that the distance conjecture holds. By developing new restriction type estimates associated to circles and orbits, with a group theoretic argument, we will prove the 4/3-parallel result in the two dimensions. This answers a question raised by Alex Iosevich. In a more general scenario, the existence/distribution of geometric/graph configurations will be also considered in this paper. Our results present improvements and extensions of recent results due to Ben Lichtin (2019, 2023). In comparison with Lichtin's method, our approach is much simpler and flexible, which is also one of the novelties in this paper.

Contents

1	Introduction	2
	1.1 Results on the generalized-distance sets	3
	1.2 The conjecture in even dimensions	5
	1.3 Results on geometric/graph configurations	6
2	Notations and Lemmas	7
3	Proof of Theorems 1.4, 1.6, and Corollary 1.5	14
4	Extension estimates associated to circles and orbits	19
	4.1 Extension theorems	20
	4.1.1 Extension theorems associated to circles	20
	4.1.2 Extension theorems associated to orbits	20
	4.2 Preliminary lemmas	21
	4.3 Proof of extension theorems	26
5	Proof of Theorem 1.8 and 1.9	27
6	Proof of Theorems 1.11, 1.12, 1.13, and 1.14	31
7	Sharpness examples	33
8	Acknowledgements	34

1 Introduction

Let p be a prime, r be a positive integer, and $\mathbb{Z}/p^r\mathbb{Z}$ be a finite p-adic ring. For a postive integer n, the "distance" between two points $\mathbf{x} = (x_1, \ldots, x_n)$ and $\mathbf{y} = (y_1, \ldots, y_n)$ in $(\mathbb{Z}/p^r\mathbb{Z})^n$, denoted by ||x - y||, is defined by

$$||x - y|| = (x_1 - y_1)^2 + \dots + (x_n - y_n)^2 \pmod{p^r}.$$

Given subsets $E_1, E_2 \subset (\mathbb{Z}/p^r\mathbb{Z})^n$, the distance set determined by points in $E_1 \times E_2$ is denoted by $\Delta_{n,r}(E_1, E_2)$, i.e.,

$$\Delta_{n,r}(E_1, E_2) = \{ \| \mathbf{x} - \mathbf{y} \| : \, \mathbf{x} \in E_1, \, \mathbf{y} \in E_2 \}.$$

For simplicity, we write $\Delta_{n,r}(E) = \Delta_{n,r}(E, E)$. We also denote the density by $\delta_{E_1,E_2} := \frac{\sqrt{|E_1||E_2|}}{p^{rn}}$ and $\delta_E := \frac{|E|}{p^{rn}}$.

In the setting of finite *p*-adic rings, the Erdős-Falconer distance problem is stated as follows.

Question 1.1. What is the smallest density threshold $\delta \in (0,1)$ independent of r such that $|\Delta_{n,r}(E)| \gg p^r$ whenever E is a subset of $(\mathbb{Z}/p^r\mathbb{Z})^n$ with $\delta_E \geq \delta$?

This problem was initially studied in the case r = 1, i.e. over finite fields, due to Iosevich and Rudnev [7]. More precisely, they proved that if $|E| \ge Cp^{\frac{n+1}{2}}$ for some sufficient large constant C, then the distance set $\Delta_{n,1}(E)$ covers the whole field. Hart, Iosevich, Koh, and Rudnev [4] indicated that the exponent $\frac{n+1}{2}$ is sharp in odd dimensions. In even dimensions, it is conjectured that the right exponent should be n/2. In two dimensions, Chapman, Erdogan, Hart, Iosevich, and Koh [2] proved the exponent 4/3 by using an extension theorem associated to circles in the plane. This result is recently improved to 5/4 by Murphy, Petridis, the first listed author, Rudnev, and Stevens in [20] by using algebraic methods and results from incidence geometry.

When r > 1, by extending the techniques from finite fields, Covert, Iosevich and Pakianathan [1] proved that $U_r \subset \Delta_{n,r}(E)$ whenever $|E| \gg r(r+1)p^{rn}p^{-\frac{n-1}{2}}$, where $U_r := (\mathbb{Z}/p^r\mathbb{Z})^*$ is the set of units. This result is only non-trivial when r is bounded, and does not offer a uniform density independent of r, to Question 1.1. By using a different and sophisticated approach, namely, a combination of p-adic analysis and estimates for a class of exponential sums mod p, Lichtin [15] proved that $U_r \subset \Delta_{n,r}(E)$ if $\delta_E \gg p^{-\frac{n-1}{2}}$. As mentioned in his paper, the main advantage of his approach is that the argument detects nontrivial cancellations within certain exponential sums mod p^r which were not used in the work of Covert et al. [1].

In this paper, the first purpose is to study the distance problem in a general setting when the distance function is replaced by a diagonal polynomial. In the case of the usual distance function, one of the novelties of this paper is to present a much simpler and flexible approach than that of Lichtin. Our results are essentially sharp in odd dimensions. The second purpose of this paper is to clarify the conjecture in even dimensions and provide examples to support it. In two dimensions, by developing new restriction type estimates associated to circles and orbits, with a group theoretic argument, we will prove the 4/3-parallel result. Surprisingly, in comparison with the finite field case, in this setting, we are able to provide a large family of sets such that the distance conjecture holds. The third purpose is to study the existence/distribution of geometric/graph configurations.

To be precise, we are interested in the following questions.

Question 1.2. Let H be a given graph, E be any subset of $(\mathbb{Z}/p^r\mathbb{Z})^n$ and j be any element of U_r . What is the smallest density threshold $\delta \in (0,1)$ independent of r such that E contains a copy of H at distance j whenever $\delta_E \gg \delta$?

Question 1.3. Let C be a given geometric configuration and E be any set in $(\mathbb{Z}/p^r\mathbb{Z})^n$. What is the smallest density threshold $\delta \in (0,1)$ not depending on r such that E contains a copy of Cwhenever $\delta_E \gg \delta$?

This paper addresses the graphs of being cycles, chains, and trees, and a geometric configuration of rectangles. In contrast to initial methods/results over finite fields, the main challenge one has to deal with in this setting is to find an effective approach to obtain a uniform density which does not depend on r. Lichtin's method in [15] is based on a combination of p-adic analysis and estimates for a class of exponential sums mod p. To find cancellations in exponential sums, he investigated paths of points (descending family of neighbourhoods), and his arguments are heavily relied on computing Hessian matrix of all levels. The approach we introduce in this paper is much simpler, which involves polynomial congruences and Fourier analysis in finite rings. Our arguments only rely on the first order derivative ∇F , and avoid the second derivative Hessian matrix.

1.1 Results on the generalized-distance sets

For a polynomial $F(\mathbf{x}) \in \mathbb{Z}[\mathbf{x}]$ in *n* variables. We denote $F_i(\mathbf{x}) = \frac{\partial F}{\partial x_i}(\mathbf{x})$ for all $1 \leq i \leq n$, and denote $(\nabla F)(\mathbf{x}) = (F_1(\mathbf{x}), F_2(\mathbf{x}), \dots, F_n(\mathbf{x}))$. Our first result reads as follows.

Theorem 1.4. Let $F(\mathbf{x})$ be a given polynomial in $\mathbb{Z}[\mathbf{x}]$ in $n \ge 2$ variables. Suppose that the following conditions hold for some positive constants c_1, c_2, c_3 with $c_1 < 1$, for some prime p, and for some $j \in \mathbb{Z}$ with (j, p) = 1:

(i) $(\nabla F)(\mathbf{x}) \not\equiv \mathbf{0} \pmod{p}$ when $F(\mathbf{x}) \equiv j \pmod{p}$;

(*ii*) $|\#\{\mathbf{x} (mod \ p) : F(\mathbf{x}) \equiv j (mod \ p)\} - p^{n-1}| \le c_1 p^{n-1};$

(iii) When $\mathbf{m} \not\equiv \mathbf{0} \pmod{p}$,

$$\Big|\sum_{\substack{\mathbf{x} \pmod{p}\\F(\mathbf{x}) \equiv j \pmod{p}}} e_p \left(-\mathbf{m} \cdot \mathbf{x}\right)\Big| \le c_2 p^{\frac{n-1}{2}};$$

(iv) When $\mathbf{m} \not\equiv \mathbf{0} \pmod{p}$,

$$\begin{split} \# \{ \mathbf{x} (\operatorname{mod} p) : F(\mathbf{x}) &\equiv j \ (\operatorname{mod} p), \ \exists \ 1 \leq t \leq n, \ s.t., F_t(\mathbf{x}) \not\equiv 0 \ (\operatorname{mod} p), \\ m_t \not\equiv 0 \ (\operatorname{mod} p), m_i F_t(\mathbf{x}) \equiv m_t F_i(\mathbf{x}) \ (\operatorname{mod} p) \ (1 \leq i \leq n) \} \leq c_3 p^{\frac{n-1}{2}}. \end{split}$$

Let r be any positive integer and E_1 , E_2 be subsets of $(\mathbb{Z}/p^r\mathbb{Z})^n$. Assume that

$$\delta_{E_1,E_2} > \frac{C}{p^{\frac{n-1}{2}}},$$

where $C = (1-c)^{-1} \max\{c_2, c_3\}$. Then

$$\#\{(\mathbf{x}, \mathbf{y}) \in E_1 \times E_2 : F(\mathbf{x} - \mathbf{y}) \equiv j \pmod{p^r}\} > 0.$$

Remark 1.1. In applications, the constants c_1, c_2, c_3 depend only on the polynomial F (degree, coefficients, number of variables), and the prime p needs to be sufficiently large compared to the degree and the coefficients of F. So the result is uniform in all natural numbers $r \ge 1$ and all integers j with (j,p) = 1. Here and throughout, by depending on F we mean in terms of the degree, coefficients, and number of variables.

Corollary 1.5. Let $F(\mathbf{x}) = \sum_{i=1}^{n} a_i x_i^k$ be a polynomial in $\mathbb{Z}[\mathbf{x}]$ with $n \ge 2$, $k \ge 2$, and $a_i \ne 0$ for all $1 \le i \le n$. Then, for any sufficiently large prime p, the conclusion of Theorem 1.4 holds for all integers r, j with $r \ge 1$ and (j, p) = 1, and with the constant C depending only on F.

It also worth noting that when k = 3 and r = 1, Corollary 1.5 recovers a result established by Iosevich and Koh in [6]. Moreover, Example 7.1 in the last section will show that this distance result is sharp in odd dimensions.

We mention that condition (iii) in Theorem 1.4 is equivalent to

$$\Big|\sum_{\mathbf{x} \,(\mathrm{mod}\ p)}\sum_{s \not\equiv 0 \,(\mathrm{mod}\ p)}\ e_p \left(-\mathbf{m} \cdot \mathbf{x} + sF(\mathbf{x}) - sj\right)\Big| \ll p^{\frac{n+1}{2}}$$

for $\mathbf{m} \neq \mathbf{0} \pmod{p}$. However, the above bound is not applicable to polynomials of the form $F(\mathbf{x}) = \sum_{j=1}^{n} a_j x_j^{k_j}$ with distinct exponents. In [11], Koh and Shen worked with a weaker exponential sum

$$\Big|\sum_{\mathbf{x} \,(\mathrm{mod}\ p)} e_p \left(-\mathbf{m} \cdot \mathbf{x} + sF(\mathbf{x})\right)\Big| \ll p^{\frac{n}{2}}$$

for $\mathbf{m} \neq \mathbf{0} \pmod{p}$ and $s \neq 0 \pmod{p}$. And they proved that $|\Delta_{n,1}(E_1, E_2)| \gg p$ whenever E_1, E_2 are subsets of \mathbb{F}_p satisfying

$$\delta_{E_1,E_2} \gg p^{-(n-1)/2}.$$
 (1)

Merely using the approach in [11] can not lead to a result that is uniform in r. With the method developed in this paper, we are able to provide a generalization of the above result in the setting of finite p-adic rings.

Theorem 1.6. Let $F(\mathbf{x}) = \sum_{i=1}^{n} a_i x_i^{k_i}$ be a polynomial in $\mathbb{Z}[\mathbf{x}]$ with $n \ge 2$, $k_i \ge 2$ and $a_i \ne 0$ for all $1 \le i \le n$. Denote $k_* = \min_{1 \le i \le n} k_i$. Then, for any sufficiently large prime p, for any natural number $r \ge 1$, and any subsets $E_1, E_2 \subseteq \mathbb{Z}/p^r\mathbb{Z}$, we have

$$|\Delta_{n,r}(E_1, E_2)| \gg \min\left\{p^r, \frac{|E_1||E_2|}{p^{r(2n-2/k_*)-n+1}}\right\}.$$

In particular, one has $|\Delta_{n,r}(E_1, E_2)| \gg p^r$ if

$$\delta_{E_1,E_2} \gg p^{r\left(\frac{1}{2} - \frac{1}{k_*}\right) - \frac{n-1}{2}}.$$
 (2)

Here, the implied constants depend only on F.

If $k_* = 2$, then we can see that the lower bound of the density is independent of r. However, if $k_* \ge 3$, then the formula (2) is meaningful only when $r \le \frac{k_*(n-1)}{k_*-2}$.

1.2 The conjecture in even dimensions

In even dimensions, one might ask about the conjectured densities to guarantee that $U_r \subset \Delta_{n,r}(E)$ or $|\Delta_{n,r}(E)| \gg p^r$. We propose the following conjecture.

Conjecture 1.7. Let $F(\mathbf{x}) = \sum_{i=1}^{n} a_i x_i^k$ be a polynomial in $\mathbb{Z}[\mathbf{x}]$ with $n \ge 2$ even, $k \ge 2$, and $a_i \ne 0$ for all $1 \le i \le n$. Then, for any sufficiently large prime p, if the density $\delta_{E_1,E_2} \gg p^{-\frac{n}{2}}$, then we have $|\Delta_{n,r}(E_1,E_2)| \gg p^r$.

Example 7.2 in the last section will support this conjecture for the case k = 2.

In the setting of finite fields, i.e. r = 1, as mentioned above, Chapman, Erdogan, Hart, Iosevich, and Koh [2] proved the exponent 4/3 by using an extension theorem associated to circles in the plane. Another proof by using geometric properties of rigid-motions in the plane can be found in [5]. Alex Iosevich asked in several conferences/workshops if the 4/3-parallel result exists in the setting of finite p-adic rings. In this paper, we give an affirmative answer to his question.

Theorem 1.8. Let $r \ge 1$ be an integer. Let $E \subset (\mathbb{Z}/p^r\mathbb{Z})^2$ with $|E| \gg p^{2r-\frac{2}{3}}$. Then $|\Delta_{2,r}(E)| \gg p^r$. Here the implied constant is independent of r.

Compared to Theorem 1.4, the exponent $2r - \frac{2}{3}$ gives a smaller threshold density, namely, of $p^{-\frac{2}{3}}$. Regarding the proof of Theorem 1.8, it is challenging if we just want to follow the methods in [2] or in [5]. On the one hand, in a recent exposition, Liao showed that the method in [5] implies too many degenerate cases, even with r = 2, which are very hard to deal with. On the other hand, a direct computation shows that the same happens with the approach developed in [2], namely, the main difficulty arises when finding the explicit form of the sum $\sum_{j \in \mathbb{Z}/p^r\mathbb{Z}} \widehat{1_{C_{r,j}}}(\mathbf{m}) \widehat{1_{C_{r,j}}}(\mathbf{m}')$ for \mathbf{m} and \mathbf{m}' in $(\mathbb{Z}/p^r\mathbb{Z})^n$, where $C_{r,j}$ is the circle centered at the origin of radius j.

In this paper, to prove Theorem 1.8, we first use a group theoretic argument, which we learned from [19, Appendix], to reduce the theorem to an extension type question for circles, then the rest is devoted to study such type estimates in the p-adic setting. The extension theorems are of independent interest and are expected to have many other applications.

Unlike the finite field case, in finite p-adic rings, we are able to provide a large family of sets E such that Conjecture 1.7 is true, namely, the family of sets E satisfying the property that the density of the fibre of the natural projection over each element in $(\mathbb{Z}/p\mathbb{Z})^2$ is not too large in E. The precise statement reads as follows.

Theorem 1.9. Let $r \geq 2$ be an integer. Let $E \subset (\mathbb{Z}/p^r\mathbb{Z})^2$ with $|E| \gg p^{2r-1}$. Assume that

$$\#\{(\mathbf{y}_1, \mathbf{y}_2) \in E^2 : \mathbf{y}_1 \equiv \mathbf{y}_2 \pmod{p}\} \ll p^{2r - \frac{7}{3}} |E|.$$

Then $|\Delta_{2,r}(E)| \gg p^r$.

Corollary 1.10. Let $r \geq 2$ be an integer. Let $E \subset (\mathbb{Z}/p^r\mathbb{Z})^2$ with $|E| \gg p^{2r-1}$. Assume that

$$\#\{\mathbf{x}' \in E \colon \mathbf{x}' \equiv \mathbf{x} \,(\textit{mod } p)\} \ll p^{2r - \frac{7}{3}}$$

for each $\mathbf{x} \in (\mathbb{Z}/p\mathbb{Z})^2$. Then $|\Delta_{2,r}(E)| \gg p^r$.

The implied constants in above two corollaries are independent of r.

1.3 Results on geometric/graph configurations

Let $F(\mathbf{x})$ be a given polynomial in $\mathbb{Z}[\mathbf{x}]$ in $n \geq 2$ variables. We call $F(\mathbf{x})$ good if it satisfies the conditions of Theorem 1.4. In the following, we obtain a number of extensions of the distance result in the setting of geometric/graph configurations in which the polynomials $F(\mathbf{x})$ are assumed to be good.

The first result is on the existence of rectangles of given side-length $j \in U_r$, which is an extension of an earlier result due to Lyall and Magyar in [17].

Theorem 1.11. For given $0 < \delta' < \delta < 1$ and an integer j with (j, p) = 1, there exists $\epsilon > 0$ such that the following holds. If $E \subset (\mathbb{Z}/p^r\mathbb{Z})^{2n}$, with $p^{-\frac{2n-1}{2}} \leq \epsilon$, satisfies $|E| \geq \delta p^{2rn}$, then E contains at least $\delta' p^{4rn-2}$ tuples

$$(\mathbf{u}_1, \mathbf{v}_1), \ (\mathbf{u}_1, \mathbf{v}_2), \ (\mathbf{u}_2, \mathbf{v}_1), \ (\mathbf{u}_2, \mathbf{v}_2)$$

that form rectangles of side-length j, i.e. $F(\mathbf{u}_1 - \mathbf{u}_2) = F(\mathbf{v}_1 - \mathbf{v}_2) \equiv j \pmod{p}$.

If we only want to count the number of quadruples $(\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3, \mathbf{u}_4)$ in E that form a cycle, i.e., $F(\mathbf{u}_1 - \mathbf{u}_2) = F(\mathbf{u}_2 - \mathbf{u}_3) = F(\mathbf{u}_3 - \mathbf{u}_4) = F(\mathbf{u}_4 - \mathbf{u}_1) \equiv j \pmod{p}$, then a weaker condition is sufficient.

Theorem 1.12. Let $E \subset (\mathbb{Z}/p^r\mathbb{Z})^n$. If $\delta_E \gg p^{-\frac{n-1}{2}}$, then E contains cycles of length 4 with distinct vertices and of side-length j.

The next result is on the existence of k-chains, i.e., a graph of k+1 vertices $\mathbf{u}_1, \ldots, \mathbf{u}_{k+1}$ such that $F(\mathbf{u}_i - \mathbf{u}_{i+1}) \equiv j \pmod{p}$ for all $1 \leq i \leq k-1$.

Theorem 1.13. Let $E \subset (\mathbb{Z}/p^r\mathbb{Z})^n$ and $k \geq 1$ be an integer. If $\delta_E \gg p^{-\frac{n-1}{2}}$, then E contains k-chains of side-length j.

We note that this theorem improves the density of $p^{-\frac{n-1}{3}}$ due to Lichtin in a recent paper [16].

Our last result is on the distribution of pinned trees. We first need to introduce some notations.

Let T be an arbitrary tree with k+1 vertices and k edges. Assume that $V(T) = {\mathbf{v}_1, \ldots, \mathbf{v}_{k+1}} \subset E$, then the edge set of T can be ordered as follows:

$$\mathcal{E}(T) = \{ (\mathbf{v}_{i_1}, \mathbf{v}_{i_2}), (\mathbf{v}_{i_3}, \mathbf{v}_{i_4}), \dots, (\mathbf{v}_{i_{2k-1}}, \mathbf{v}_{i_{2k}}) \},\$$

where $i_1 \le i_3 \le \dots \le i_{2k-1}$, and $i_{2s} < i_{2t}$ if s < t and $i_{2s-1} = i_{2t-1}$.

For such a tree T, the vector

$$(|\mathbf{v}_{i_1} - \mathbf{v}_{i_2}|, |\mathbf{v}_{i_3} - \mathbf{v}_{i_4}|, \dots, |\mathbf{v}_{i_{2k-1}} - \mathbf{v}_{i_{2k}}|) \in (\mathbb{Z}/p^r\mathbb{Z})^k$$

is called the edge-length vector of T. Two trees T and T' are called distinct if the corresponding edge-length vectors are not the same. We also recall that two trees T and T' are isotropic if there

exists a bijective map φ from V(T) to V(T') such that the edges are preserved. For $\mathbf{x} \in V(T')$ and $\mathbf{v} \in V(T)$, we say (T', \mathbf{x}) is isomorphic to (T, \mathbf{v}) if T is isomorphic to T' under φ and $\varphi(\mathbf{x}) = \mathbf{v}$.

Theorem 1.14. Let $E \subset (\mathbb{Z}/p^r\mathbb{Z})^n$, $k \geq 1$ be an integer, and (T, \mathbf{v}) be a given tree with k + 1 vertices and k edges. If $\delta_E \gg_k p^{-\frac{n-1}{2}}$, then there exists $\mathbf{x} \in E$ such that the number of distinct trees (T', \mathbf{x}) with vertices in E and isotropic to (T, \mathbf{v}) is at least $\gg p^{rk}$.

In the rest of the paper, in some places, we may denote $q = p^r$ for simplicity if it causes no harm to the argument.

2 Notations and Lemmas

In this paper, the letter p always denote a given prime p > 2. The letter n denotes the dimension of the space under consideration, and r is always a positive integer. The cardinality of a finite set S is denoted by either |S| or #S.

When we write $\mathbf{z} \pmod{p^r}$ for an *n*-dimensional vector \mathbf{z} , it always means that \mathbf{z} is considered as an element in $(\mathbb{Z}/p^r\mathbb{Z})^n$. The expression $\operatorname{ord}_p(z) = u$ means that $z \equiv 0 \pmod{p^u}$ and $z \not\equiv 0 \pmod{p^{u+1}}$. We also denote $\operatorname{ord}_p(0) = r$ for $0 \in \mathbb{Z}/p^r\mathbb{Z}$. When $\mathbf{z} = (z_1, z_2, \ldots, z_n)$, we write $v_{\mathbf{z}} = \min_{1 \leq i \leq n} \operatorname{ord}_p(z_i)$. Then the vector \mathbf{z} can be expressed as $\mathbf{z} = p^{v_{\mathbf{z}}} \tilde{\mathbf{z}}$, where $\tilde{\mathbf{z}}$ is a vector in $(\mathbb{Z}/p^{r-v_{\mathbf{z}}}\mathbb{Z})$ such that $v_{\tilde{\mathbf{z}}} = 0$.

For any $j \in \mathbb{Z}/p^r\mathbb{Z}$, we use $C_{n,r,j}$ to denote the sphere in $(\mathbb{Z}/p^r\mathbb{Z})^n$ centered at origin with radius j, i.e.,

$$C_{n,r,j} = \{ \mathbf{z} \in (\mathbb{Z}/p^r \mathbb{Z})^n : \|\mathbf{z}\| = j \}.$$

Similarly, for a given polynomial F(x) in n variables with coefficients in \mathbb{Z} , we define

$$S_{n,r,j} = \{ \mathbf{z} \in (\mathbb{Z}/p^r \mathbb{Z})^n : F(\mathbf{z}) = j \},$$

When it makes no confusion, we abbreviate $C_{n,r,j}$ (or $S_{n,r,j}$) as $C_{r,j}$ (or $S_{r,j}$, respectively).

The additive character modulo p^r is denoted by $e_{p^r}(x) = e^{\frac{2\pi i x}{p^r}}$, $(x \mod p^r)$. For a function f: $(\mathbb{Z}/p^r\mathbb{Z})^n \to \mathbb{C}$, the Fourier transformation is defined by

$$\widehat{f}(\mathbf{m}) = \frac{1}{p^{rn}} \sum_{\mathbf{m} \pmod{p^r}} f(\mathbf{x}) e_{p^r}(-\mathbf{m} \cdot \mathbf{x}), \quad \left(\mathbf{m} \in (\mathbb{Z}/p^r \mathbb{Z})^n\right),$$

and the Fourier inverse is given by

$$f(\mathbf{x}) = \sum_{\mathbf{m} \, (\mathrm{mod} \ p^r)} \widehat{f}(\mathbf{m}) e_{p^r}(\mathbf{m} \cdot \mathbf{x}), \quad \big(\mathbf{x} \in (\mathbb{Z}/p^r\mathbb{Z})^n).$$

The convolution of two functions f_1 and f_2 is defined by

$$(f_1 * f_2)(\mathbf{x}) = \frac{1}{p^{rn}} \sum_{\mathbf{y} \pmod{p^r}} f_1(\mathbf{x} - \mathbf{y}) f_2(\mathbf{y}).$$

We have the property that $\widehat{f_1 * f_2} = \widehat{f_1}\widehat{f_2}$. The Parseval's identity is

$$\frac{1}{p^{rn}}\sum_{\mathbf{x} \pmod{p^r}} f_1(\mathbf{x})\overline{f_2(\mathbf{x})} = \sum_{\mathbf{m} \pmod{p^r}} \widehat{f_1}(\mathbf{m})\overline{\widehat{f_2}(\mathbf{m})}.$$

The following is one version of Hensel's lemma that we will regularly applied in this paper.

Lemma 2.1 (Hensel's lemma). Let

$$\mathbf{G}(\mathbf{x}) = (G_1(x_1, \dots, x_n), \dots, G_m(x_1, \dots, x_n))$$

be a map from \mathbb{Z}^n to \mathbb{Z}^m , with G_i polynomials with integer coefficients. Let l be a positive integer and $\mathbf{y} \in \mathbb{Z}^n$. Suppose that $\mathbf{G}(\mathbf{y}) \equiv \mathbf{0} \pmod{p^l}$. Let R be the rank of $J(\mathbf{G})|_{\mathbf{y}}$ modulo p, where $J(\mathbf{G})|_{\mathbf{y}}$ is the Jocobi matrix

$$J(\mathbf{G})|_{\mathbf{y}} = \begin{bmatrix} \frac{\partial G_1}{\partial x_1}(\mathbf{y}) & \frac{\partial G_1}{\partial x_2}(\mathbf{y}) & \dots & \frac{\partial G_1}{\partial x_n}(\mathbf{y}) \\ \frac{\partial G_2}{\partial x_1}(\mathbf{y}) & \frac{\partial G_2}{\partial x_2}(\mathbf{y}) & \dots & \frac{\partial G_2}{\partial x_n}(\mathbf{y}) \\ \dots & \dots & \dots & \dots \\ \frac{\partial G_m}{\partial x_1}(\mathbf{y}) & \frac{\partial G_m}{\partial x_2}(\mathbf{y}) & \dots & \frac{\partial G_m}{\partial x_n}(\mathbf{y}) \end{bmatrix}$$

Then

$$# \{ \mathbf{z} \,(\operatorname{mod} \, p^k) : \, \mathbf{G}(\mathbf{y} + p^l \mathbf{z}) \equiv \mathbf{0} \,(\operatorname{mod} \, p^{l+k}) \} \le p^{k(n-R)}$$
(3)

for any integer $k \ge 1$. When R = m, the " \le " can be replaced by "=".

Proof. Firstly, let us consider the case k = 1. Let $\mathbf{G}(\mathbf{y}) = p^l \mathbf{b}$, where $\mathbf{b} = (b_1, b_2, \dots, b_m) \in \mathbb{Z}^m$. Note that

$$G_i(\mathbf{y} + p^l \mathbf{z}) \equiv G_i(\mathbf{y}) + p^l(\nabla G_i)(\mathbf{y}) \cdot \mathbf{z} \pmod{p^{l+1}}$$

for any i = 1, 2, ..., m. Then $G_i(\mathbf{y} + p^l \mathbf{z}) \equiv 0 \pmod{p^{l+1}}$ (i = 1, 2, ..., m) if and only if $b_i + (\nabla G_i)(\mathbf{y}) \cdot \mathbf{z} \equiv 0 \pmod{p}$ (i = 1, 2, ..., m), if and only if $J_{\mathbf{G}}(\mathbf{y})\mathbf{z} \equiv -\mathbf{b} \pmod{p}$. The number of solutions to such a system of linear equations does not exceed p^{n-R} . Thus,

$$\# \big\{ \mathbf{z} \, (\mathsf{mod} \ p) : \ \mathbf{G}(\mathbf{y} + p^l \mathbf{z}) \equiv \mathbf{0} \, (\mathsf{mod} \ p^{l+1}) \big\} \leq p^{n-R}$$

Moreover, the system is consistent when R = m, and has exactly p^{n-m} solutions.

Secondly, let us consider the case k = 2. For any $\mathbf{z}_1 \pmod{p}$ with $\mathbf{G}(\mathbf{y} + p^l \mathbf{z}_1) \equiv \mathbf{0} \pmod{p^{l+1}}$, we write $\mathbf{y}_1 = \mathbf{y} + p^l \mathbf{z}_1$. Noting that $J(\mathbf{G})|_{\mathbf{y}_1} \equiv J(\mathbf{G})|_{\mathbf{y}} \pmod{p}$, the rank of $J(\mathbf{G})|_{\mathbf{y}_1}$ modulo p is also R. Hence

$$\# \{ \mathbf{z}_2 \,(\mathsf{mod} \, p) : \, \mathbf{G}(\mathbf{y}_1 + p^{l+1}\mathbf{z}_2) \equiv \mathbf{0} \,(\mathsf{mod} \, p^{l+2}) \} \le p^{n-R}.$$

It follows that

$$\#\{(\mathbf{z}_1, \mathbf{z}_2) \,(\text{mod } p): \, \mathbf{G}\big(\mathbf{y} + p^l(\mathbf{z}_1 + p\mathbf{z}_2)\big) \equiv \mathbf{0} \,(\text{mod } p^{l+2})\} \le p^{2(n-R)}.$$

Finally, the conclusion follows by an induction on k.

Lemma 2.2 (Weil's theorem). Let $f(x) \in \mathbb{F}_p[x]$ be a polynomial of degree $k \ge 1$ with (k, p) = 1. Then

$$\Big|\sum_{x\in\mathbb{F}_p} e_p(f(x))\Big| \le (k-1)p^{1/2}.$$

Lemma 2.3 (Theorem 2.3, [11]). Let $F(\mathbf{x}) = \sum_{i=1}^{n} a_i x_i^{k_i}$ be a polynomial in $\mathbb{F}_p[\mathbf{x}]$ such that $n \ge 2$, $k_i \ge 1$, and $a_i \ne 0$ for all $1 \le i \le n$. Let $j \in \mathbb{F}_p^*$. Then

$$\#\{\mathbf{x} \in \mathbb{F}_p : F(\mathbf{x}) = j\} = (1 + o_F(1))p^{n-1}$$

as $p \to \infty$.

Lemma 2.4 (Lemma 2.1, [11]). Let $F(\mathbf{x}) = \sum_{i=1}^{n} a_i x_i^k$ be a polynomial in $\mathbb{F}_p[\mathbf{x}]$ with $n \ge 1$, $k \ge 2$, and $a_i \ne 0$ for all $1 \le i \le n$. Let $j \in \mathbb{F}_p^*$. Then, for any $\mathbf{m} \in \mathbb{F}_p^n \setminus \{\mathbf{0}\}$, one has

$$\sum_{\substack{\mathbf{x}\in\mathbb{F}_p^n\\F(\mathbf{x})=j}} e_p(-\mathbf{m}\cdot\mathbf{x}) \ll_{n,k} p^{\frac{n-1}{2}}$$

when p is sufficiently large.

Lemma 2.5. Let p be a prime and j be an integer with (j,p) = 1. Assume that $F(\mathbf{x})$ is a polynomial in $\mathbb{Z}[\mathbf{x}]$ in $n \ge 1$ variables satisfying the conditions (i) and (iii) in Theorem 1.4, and the following:

(*iv*'): When $\mathbf{m} \not\equiv \mathbf{0} \pmod{p}$,

$$\# \{ \mathbf{y}(\operatorname{mod} p) : F(\mathbf{y}) \equiv j(\operatorname{mod} p), \exists 1 \le t \le n, \ s.t., F_t(\mathbf{y}) \not\equiv 0(\operatorname{mod} p), \\ m_t \not\equiv 0(\operatorname{mod} p), m_i F_t(\mathbf{y}) \equiv m_t F_i(\mathbf{y}) (\operatorname{mod} p) \text{ for all } 1 \le i \le n \} \le c_3 p^{\kappa}$$

for some $c_3 > 0$ and $0 \le \kappa \le \frac{n-1}{2}$.

Then, for any $r \geq 1$, we have

$$|\widehat{\mathbf{1}_{S_{r,j}}}(\mathbf{m}')| \leq \begin{cases} c_2 \, p^{-r - \frac{n-1}{2}}, & \text{if } \mathbf{m}' \equiv \mathbf{0} \, (\text{mod } p^{r-1}) \, \text{ but } \mathbf{m}' \not\equiv \mathbf{0} \, (\text{mod } p^r), \\ c_3 \, p^{-r - n + 1 + \kappa}, & \text{if } \mathbf{m}' \not\equiv \mathbf{0} \, (\text{mod } p^{r-1}). \end{cases}$$

Proof. Recall that $S_{r,j} = \{ \mathbf{x} \in (\mathbb{Z}/p^r \mathbb{Z})^n : F(\mathbf{x}) \equiv j \pmod{p^r} \}$. For simplicity, we denote $v_{\mathbf{m}'} = \nu$. Then $\mathbf{m}' = p^{\nu}\mathbf{m}$ with $\mathbf{m} \in (\mathbb{Z}/p^{r-\nu}\mathbb{Z})^n$ and $v_{\mathbf{m}} = 0$.

When r = 1, the conclusion follows from the condition (iii). In the following, we assume that $r \ge 2$.

For $1 \le \nu \le r - 1$, by a change of variables $\mathbf{x} = \mathbf{y} + p^{r-\nu} \mathbf{z}$, we have that

$$\widehat{\mathbf{1}_{S_{r,j}}}(\mathbf{m}') = \frac{1}{p^{rn}} \sum_{\substack{\mathbf{x}(\text{mod } p^{r}) \\ F(\mathbf{x}) \equiv j(\text{mod } p^{r})}} e_{p^{r}}(-\mathbf{m}' \cdot \mathbf{x})$$

$$= \frac{1}{p^{rn}} \sum_{\substack{\mathbf{y}(\text{mod } p^{r-\nu}) \\ F(\mathbf{y}) \equiv j(\text{mod } p^{r-\nu})}} \sum_{\substack{\mathbf{z}(\text{mod } p^{\nu}) \\ F(\mathbf{y}+p^{r-\nu}\mathbf{z}) \equiv j(\text{mod } p^{r})}} e_{p^{r-\nu}}\left(-\mathbf{m} \cdot \mathbf{y}\right) \sum_{\substack{\mathbf{z}(\text{mod } p^{\nu}) \\ F(\mathbf{y}) \equiv j(\text{mod } p^{r-\nu})}} 1.$$

Since $F(\mathbf{y}) \equiv j \pmod{p}$, one has $(\nabla F)(\mathbf{y}) \not\equiv \mathbf{0} \pmod{p}$ by the condition (i). By Hensel's lemma, we have

$$\sum_{\substack{\mathbf{z}(\bmod p^{\nu})\\ (\mathbf{y}+p^{r-\nu}\mathbf{z})\equiv j(\bmod p^r)}} 1 = p^{\nu(n-1)},$$

for each given $\mathbf{y} \pmod{p^{r-\nu}}$. Therefore,

$$\widehat{\mathbf{1}_{S_{r,j}}}(\mathbf{m}') = p^{-rn+\nu(n-1)} \sum_{\substack{\mathbf{y} \pmod{p^r - \nu} \\ F(\mathbf{y}) \equiv j \pmod{p^r - \nu}}} e_{p^{r-\nu}} \left(-\mathbf{m} \cdot \mathbf{y}\right), \tag{4}$$

for $1 \le \nu \le r - 1$. Moreover, when $\nu = 0$, the expression (4) trivially holds.

F

Let us substitute the parameters \mathbf{x} , γ for \mathbf{y} , $r - \nu$, respectively. Then, it is sufficient to show that

$$T_{\gamma}(\mathbf{m}) := \sum_{\substack{\mathbf{x} (\text{mod } p^{\gamma}) \\ F(\mathbf{x}) \equiv j (\text{mod } p^{\gamma})}} e_{p^{\gamma}} (-\mathbf{m} \cdot \mathbf{x}) \le \begin{cases} c_2 \, p^{(\gamma - 1/2)(n-1)}, & \text{if } \nu = r-1, \\ c_3 \, p^{(\gamma - 1)(n-1) + \kappa}, & \text{if } 0 \le \nu \le r-2. \end{cases}$$

Here $\mathbf{m} \not\equiv \mathbf{0} \pmod{p}$.

When $\gamma = 1$, the bound follows from the condition (iii). In the following, we consider the situation that $\gamma \geq 2$. By a change of variables $\mathbf{x} = \mathbf{y} + p^{\gamma - 1}\mathbf{z}$,

$$\begin{split} T_{\gamma}(\mathbf{m}) &= \sum_{\substack{\mathbf{y}(\text{mod } p^{\gamma-1})\\F(\mathbf{y}) \equiv j(\text{mod } p^{\gamma-1})\\F(\mathbf{y}) \equiv j(\text{mod } p^{\gamma-1})}} \sum_{\substack{F(\mathbf{y} + p^{\gamma-1}\mathbf{z}) \equiv j(\text{mod } p^{\gamma})\\F(\mathbf{y}) \equiv j(\text{mod } p^{\gamma-1})}} e_{p^{\gamma}} \Big(- \mathbf{m} \cdot \mathbf{y} \Big) \sum_{\substack{\mathbf{z}(\text{mod } p)\\F(\mathbf{y} + p^{\gamma-1}\mathbf{z}) \equiv j(\text{mod } p^{\gamma-1})}} e_{p^{\gamma}} \Big(- \mathbf{m} \cdot \mathbf{z} \Big) \end{split}$$

Let us denote

$$T_{\gamma}(\mathbf{m};\mathbf{y}) := \sum_{\substack{\mathbf{z} \pmod{p} \\ F(\mathbf{y} + p^{\gamma-1}\mathbf{z}) \equiv j \pmod{p^{\gamma}}}} e_p(-\mathbf{m} \cdot \mathbf{z}).$$

The condition $F(\mathbf{y} + p^{\gamma-1}\mathbf{z}) \equiv j \pmod{p^{\gamma}}$ is equivalent to

$$F(\mathbf{y}) + p^{\gamma-1}(\nabla F)(\mathbf{y}) \cdot \mathbf{z} \equiv j \pmod{p^{\gamma}}.$$

Suppose that $F(\mathbf{y}) \equiv j + p^{\gamma - 1} h_{\gamma}(\mathbf{y}) \pmod{p^{\gamma}}$ for some $h_{\gamma}(\mathbf{y}) \pmod{p}$. Recall that $F_i(\mathbf{y}) = \frac{\partial F}{\partial x_i}(\mathbf{y})$.

Also recall the condition (i) that $(\nabla F)(\mathbf{y}) \neq \mathbf{0} \pmod{p}$. Without loss of generality, let us proceed with $F_1(\mathbf{y}) \neq 0 \pmod{p}$. Then

$$z_1 \equiv -F_1^{-1}(\mathbf{y})(h_\gamma(\mathbf{y}) + F_2(\mathbf{y})z_2 + \ldots + F_n(\mathbf{y})z_n) \pmod{p}$$

Here $F_1^{-1}(\mathbf{y})$ is the inverse of $F_1(\mathbf{y}) \pmod{p}$. So

$$T_{\gamma}(\mathbf{m};\mathbf{y}) = e_p(m_1 F_1^{-1}(\mathbf{y}) h_{\gamma}(\mathbf{y})) \prod_{i=2}^n \sum_{z_i \pmod{p}} e_p((-m_i + m_1 F_1^{-1}(\mathbf{y}) F_i(\mathbf{y})) z_i).$$

It follows that

$$|T_{\gamma}(\mathbf{m};\mathbf{y})| = \begin{cases} p^{n-1}, & \text{if } m_i F_1(\mathbf{y}) \equiv m_1 F_i(\mathbf{y}) \pmod{p} \text{ for } 1 \leq i \leq n, \\ 0, & \text{otherwise.} \end{cases}$$

(Note that, for i = 1, the formula $m_i F_1(\mathbf{y}) \equiv m_1 F_i(\mathbf{y}) \pmod{p}$ trivially holds.)

Moreover, if $m_1 \equiv 0 \pmod{p}$, then the assumption $\mathbf{m} \neq \mathbf{0} \pmod{p}$ shows that $m_{i_0} \neq 0 \pmod{p}$ for some $2 \leq i_0 \leq n$. Then $m_{i_0}F_1(\mathbf{y}) \neq m_1F_{i_0}(\mathbf{y}) \pmod{p}$, which leads to $T_{\gamma}(\mathbf{m}; \mathbf{y}) = 0$. As a result, if $T(\mathbf{m}; \mathbf{y}) \neq 0$ and $F_t(\mathbf{y}) \neq 0 \pmod{p}$ for some $1 \leq t \leq n$, then $m_t \neq 0 \pmod{p}$.

Now we conclude that

$$|T_{\gamma}(\mathbf{m})| \le p^{n-1} |V_{\gamma-1}(\mathbf{m})|,$$

where

$$V_{\mu}(\mathbf{m}) := \{ \mathbf{y}(\mathsf{mod}\ p^{\mu}) :\ F(\mathbf{y}) \equiv j(\mathsf{mod}\ p^{\mu}), \ \exists \ 1 \le t \le n, \ s.t., F_t(\mathbf{y}) \not\equiv 0(\mathsf{mod}\ p), \\ m_t \not\equiv 0(\mathsf{mod}\ p), m_i F_t(\mathbf{y}) \equiv m_t F_i(\mathbf{y})(\mathsf{mod}\ p) \text{ for all } 1 \le i \le n \}.$$

When $\gamma \geq 3$, we further have

$$|V_{\gamma-1}(\mathbf{m})| \leq \sum_{\mathbf{w} \in V_1(\mathbf{m})} \#\{\mathbf{y} (\text{mod } p^{\gamma-1}): F(\mathbf{y}) \equiv j (\text{mod } p^{\gamma-1}), \, \mathbf{y} \equiv \mathbf{w} \, (\text{mod } p)\}.$$

Note that $F(\mathbf{w}) \equiv j \pmod{p}$ for any $\mathbf{w} \in V_1$. So $(\nabla F)(\mathbf{w}) \not\equiv 0 \pmod{p}$ by the condition (i). Applying Hensel's lemma again, we have

$$\begin{aligned} &\#\{\mathbf{y}(\text{mod } p^{\gamma-1}): F(\mathbf{y}) \equiv j(\text{mod } p^{\gamma-1}), \, \mathbf{y} \equiv \mathbf{w} \,(\text{mod } p)\} \\ &= \#\{\mathbf{z} \,(\text{mod } p^{\gamma-2}): F(\mathbf{w} + p\mathbf{z}) \equiv j \,(\text{mod } p^{\gamma-1})\} = p^{(\gamma-2)(n-1)} \end{aligned}$$

for each given **w**. It follows that, for any $\gamma \geq 2$,

$$|T_{\gamma}(\mathbf{m})| \le p^{(\gamma-1)(n-1)} |V_1(\mathbf{m})|.$$

Recalling that $\mathbf{m} \neq \mathbf{0} \pmod{p}$, one can apply the condition (iv') to get $|V_1(\mathbf{m})| \leq c_3 p^{\kappa}$. Thus, we conclude that $|T_{\gamma}(\mathbf{m})| \leq c_3 p^{(\gamma-1)(n-1)+\kappa}$. The proof is completed.

Remark: When n = 1, the cardinality of the set in condition (iv') becomes

$$\#\{y \,(\mathsf{mod} \,\, p): \, F(y) \equiv j \,(\mathsf{mod} \,\, p)\},\$$

which is no larger than the degree of F(y). So one may take $\kappa = 0$ in this case.

Lemma 2.6. Let κ be a number with $0 \leq \kappa \leq n/2$ and $c_1, c_2 > 0$ be constants. Let $F(\mathbf{x}) \in \mathbb{Z}[\mathbf{x}]$ be a polynomial in $n \geq 1$ variables of degree ≥ 2 such that the upper bounds

$$\sum_{\mathbf{x} \pmod{p}} e_p \left(sF(\mathbf{x}) + \mathbf{m} \cdot \mathbf{x} \right) \le c_1 p^{n/2}$$
(5)

and

$$\#\{\mathbf{x} (mod \ p): \ s (\nabla F)(\mathbf{x}) + \mathbf{m} \equiv \mathbf{0} (mod \ p)\} \le c_2 p^{\kappa}$$
(6)

both hold when $(\mathbf{m}, s) \not\equiv (\mathbf{0}, 0) \pmod{p}$. Then, for any $r \ge 1$, we have

$$\begin{split} &\sum_{\mathbf{x} \ (\text{mod } p^{r})} e_{p^{r}} \left(s'F(\mathbf{x}) + \mathbf{m}' \cdot \mathbf{x} \right) \\ &\leq \begin{cases} c_{1}p^{(r-1/2)n}, & \text{if } (\mathbf{m}', \ s') \equiv (\mathbf{0}, 0) \ (\text{mod } p^{r-1}) \ but \ (\mathbf{m}', \ s') \not\equiv (\mathbf{0}, 0) \ (\text{mod } p^{r}), \\ c_{2}p^{(r-1)n+\kappa}, & \text{if } (\mathbf{m}', \ s') \not\equiv (\mathbf{0}, 0) \ (\text{mod } p^{r-1}). \end{cases} \end{split}$$

Proof. For simplicity, let us write $v_{(\mathbf{m}',s')} = \nu$. Since $(\mathbf{m}', s') \not\equiv (\mathbf{0}, 0) \pmod{p^r}$, we have $0 \leq \nu \leq r-1$. Write $\mathbf{m}' = p^{\nu}\mathbf{m}$ and $s' = p^{\nu}s$. Then $(\mathbf{m}, s) \not\equiv (\mathbf{0}, 0) \pmod{p}$.

When $1 \le \nu \le r - 1$, we obtain by a change of variables $\mathbf{x} = \mathbf{y} + p^{r-\nu}\mathbf{z}$ that

$$\begin{split} G_r(\mathbf{m}',s') &:= \sum_{\mathbf{x} \pmod{p^r}} e_{p^r} \left(s'F(\mathbf{x}) + \mathbf{m}' \cdot \mathbf{x} \right) \\ &= \sum_{\mathbf{y} \pmod{p^{r-\nu}}} \sum_{\mathbf{z} \pmod{p^{\nu}}} e_{p^r} \left(p^{\nu} sF(\mathbf{y}) + p^{\nu} s(\nabla F)(\mathbf{y}) \cdot (p^{r-\nu}\mathbf{z}) + p^{\nu}\mathbf{m} \cdot (\mathbf{y} + p^{r-\nu}\mathbf{z}) \right) \\ &= \sum_{\mathbf{y} \pmod{p^{r-\nu}}} e_{p^{r-\nu}} \left(sF(\mathbf{y}) + \mathbf{m} \cdot \mathbf{y} \right) \sum_{\mathbf{z} \pmod{p^{\nu}}} 1 = p^{\nu n} \cdot G_{r-\nu}(\mathbf{m}, s). \end{split}$$

When $\nu = 0$, the above equality trivially holds. By writing $\gamma = r - \nu$, it is sufficient to prove that

$$G_{\gamma}(\mathbf{m}, s) \le \begin{cases} c_1 p^{(\gamma - 1/2)n}, & \text{if } \nu = r - 1, \\ c_2 p^{(\gamma - 1)n + \kappa}, & \text{if } 0 \le \nu \le r - 2, \end{cases}$$

where $(\mathbf{m}, s) \not\equiv (\mathbf{0}, 0) \pmod{p}$.

When $\gamma = 1$, the conclusion follows from (5). In the following, let us assume that $\gamma \geq 2$. By a

change of variables $\mathbf{x} = \mathbf{y} + p^{\gamma - 1} \mathbf{z}$, we obtain that

$$\begin{split} G_{\gamma}(\mathbf{m},s) &= \sum_{\mathbf{x} \pmod{p^{\gamma}}} e_{p^{\gamma}} \left(sF(\mathbf{x}) + \mathbf{m} \cdot \mathbf{x} \right) \\ &= \sum_{\mathbf{y} \pmod{p^{\gamma-1}}} e_{p^{\gamma}} \left(sF(\mathbf{y}) + \mathbf{m} \cdot \mathbf{y} \right) \sum_{\mathbf{z} \pmod{p}} e_{p} \left(\left(s\left(\nabla F\right)(\mathbf{y}) + \mathbf{m}\right) \cdot \mathbf{z} \right) \\ &= p^{n} \sum_{\substack{\mathbf{y} \pmod{p^{\gamma-1}}\\ s(\nabla F)(\mathbf{y}) + \mathbf{m} \equiv \mathbf{0} \pmod{p}}} e_{p^{\gamma}} \left(sF(\mathbf{y}) + \mathbf{m} \cdot \mathbf{y} \right) \\ &\leq p^{n} \cdot \#\{\mathbf{y} \pmod{p^{\gamma-1}}: \ s\left(\nabla F\right)(\mathbf{y}) + \mathbf{m} \equiv \mathbf{0} \pmod{p}\}. \end{split}$$

When $\gamma = 2$, it follows from (6) that $G_2(\mathbf{m}, s) \leq c_2 p^{n+\kappa}$. When $\gamma \geq 3$, we further have

$$\begin{aligned} &\#\{\mathbf{y} \ (\text{mod } p^{\gamma-1}): \ s \ (\nabla F)(\mathbf{y}) + \mathbf{m} \equiv \mathbf{0} \ (\text{mod } p)\} \\ &= \#\{(\mathbf{w}, \mathbf{z}) \in (\mathbb{Z}/p\mathbb{Z}) \times (\mathbb{Z}/p\mathbb{Z})^{\gamma-2}: \ s \ (\nabla F)(\mathbf{w} + p\mathbf{z}) + \mathbf{m} \equiv \mathbf{0} \ (\text{mod } p)\} \\ &= p^{(\gamma-2)n} \cdot \#\{\mathbf{w} \ (\text{mod } p): \ s \ (\nabla F)(\mathbf{w}) + \mathbf{m} \equiv \mathbf{0} \ (\text{mod } p)\}. \end{aligned}$$

By (6) again, one deduces that

$$G_{\gamma}(\mathbf{m},s) \leq p^n \cdot p^{(\gamma-2)n} \cdot c_2 p^{\kappa} = c_2 p^{(\gamma-1)n+\kappa}.$$

The proof is completed.

Lemma 2.7. Let a be an integer with (a, p) = 1. Let $k \ge 2$ be a natural number with (k, p) = 1. Then for any $m \pmod{p}$,

$$\sum_{s \pmod{p}} \left| \sum_{x \pmod{p}} e_p(-mx + sax^k) \right|^2 \ll_k p^2.$$
(7)

Moreover, for any natural number $r \geq 2$, it satisfies that

$$\sum_{s \, (\text{mod } p^r)} \left| \sum_{x \, (\text{mod } p^r)} e_{p^r}(-mx + sax^k) \right|^2 \ll_k \begin{cases} p^{2r}, & \text{if } \nu(m) < r - \lceil \frac{r}{k} \rceil, \\ p^{(3-2/k)r}, & \text{if } \nu(m) \ge r - \lceil \frac{r}{k} \rceil, \end{cases}$$
(8)

for $m \in \mathbb{Z}/p^r\mathbb{Z}$ with $0 \leq \nu(m) \leq r$. Here $\nu(m) = r$ means $m \equiv 0 \pmod{p^r}$.

Proof. Note that

$$\sum_{s \,(\text{mod } p^r)} \left| \sum_{x \,(\text{mod } p^r)} e_{p^r}(-mx + sax^k) \right|^2 = \sum_{x,y \,(\text{mod } p^r)} \sum_{s \,(\text{mod } p^r)} e_{p^r}\left(-m(x-y) + sa(x^k - y^k)\right)$$
$$= p^r \sum_{\substack{x,y \,(\text{mod } q) \\ x^k \equiv y^k \,(\text{mod } p^r)}} \sum_{\substack{x,y \,(\text{mod } p^r) \\ x^k \equiv y^k \,(\text{mod } p^r)}} e_q\left(-m(x-y)\right) = p^r \sum_{j \,(\text{mod } p^r)} \left| \sum_{\substack{x \,(\text{mod } p^r) \\ x^k \equiv j \,(\text{mod } p^r)}} e_{p^r}(-mx) \right|^2.$$

When $j \not\equiv 0 \pmod{p}$, one has

$$\left|\sum_{\substack{x \pmod{p^r} \\ x^k \equiv j \pmod{p^r}}} e_{p^r}(-mx)\right| \le \#\{x \pmod{p^r} : x^k \equiv j \pmod{p^r}\}.$$
(9)

When r = 1, it is not hard to see that the above quantity is $\ll_k 1$. By Hensel's lemma, we conclude that (9) is $\ll_k 1$ for any $r \ge 1$. Hence

$$q\sum_{\substack{j\not\equiv 0\,(\mathrm{mod}\,\,p^r)\\x^k\equiv j\,(\mathrm{mod}\,\,p^r)}}\Big|\sum_{\substack{x\,(\mathrm{mod}\,\,p^r)\\x^k\equiv j\,(\mathrm{mod}\,\,p^r)}}e_{p^r}(-mx)\Big|^2\ll p^{2r}.$$

Next, we consider the case that $j \equiv 0 \pmod{p}$. When r = 1, it is easy to see that

$$p \cdot \Big| \sum_{\substack{x \pmod{p} \\ x^k \equiv 0 \pmod{p}}} e_p(-mx) \Big|^2 = p.$$

When $r \ge 2$, let us take $\mu = \lceil \frac{r}{k} \rceil$. Then $x^k \equiv 0 \pmod{p^r}$ if and only if $p^{\mu}|x$. By writing $x = p^{\mu}y$, we obtain that

$$\sum_{\substack{x \pmod{p^r} \\ x^k \equiv 0 \pmod{p^r}}} e_{p^r}(-mx) = \sum_{y \pmod{p^{r-\mu}}} e_{p^{r-\mu}}(-my) \ll p^{r-\mu} \ll p^{r(1-1/k)}.$$

Donote $\nu = \nu(m)$ and write $m = p^{\nu} \tilde{m}$. When $1 \leq \nu < r - \mu$, one deduces by a change of variables $y = w + p^{r-\mu-\nu}z$ that

$$\sum_{y \pmod{p^{r-\mu}}} e_{p^{r-\mu}}(-my) = \sum_{w \pmod{p^{r-\mu-\nu}}} e_{p^{r-\mu-\nu}}(-\tilde{m}w) \sum_{z \pmod{p^{\nu}}} 1 = 0.$$

When $\nu = 0$, the above result also holds. So the summand over $j \equiv 0 \pmod{p^r}$ is

$$p^{r} \cdot \Big| \sum_{\substack{x \pmod{p^{r}} \\ x^{k} \equiv 0 \pmod{p^{r}}}} e_{p^{r}}(-mx) \Big|^{2} \ll \begin{cases} 0, & \text{if } \nu(m) < r - \lceil \frac{r}{k} \rceil, \\ p^{r(3-2/k)}, & \text{if } \nu(m) \ge r - \lceil \frac{r}{k} \rceil. \end{cases}$$

1

The lemma then follows.

3 Proof of Theorems 1.4, 1.6, and Corollary 1.5

For simplicity, we write $q = p^r$ in this section.

Proof of Theorem 1.4. Recall that $S_{r,j} = \{ \mathbf{z} \in (\mathbb{Z}/q\mathbb{Z})^n \colon F(\mathbf{z}) \equiv j \pmod{q} \}$. First, we have

$$\begin{split} \mathcal{N}_{r,j} &:= \#\{(\mathbf{x}, \mathbf{y}) \in E_1 \times E_2 : \ F(\mathbf{x} - \mathbf{y}) \equiv j \ (\mathsf{mod} \ q)\} \\ &= \sum_{\mathbf{x}, \mathbf{y} \ (\mathsf{mod} \ q)} \mathbf{1}_{E_1}(\mathbf{x}) \mathbf{1}_{E_2}(\mathbf{y}) \mathbf{1}_{S_{r,j}}(\mathbf{x} - \mathbf{y}) = q^{2n} \cdot \frac{1}{q^n} \sum_{\mathbf{x} \ (\mathsf{mod} \ q)} \mathbf{1}_{E_1}(\mathbf{x}) (\mathbf{1}_{E_2} * \mathbf{1}_{S_{r,j}})(\mathbf{x}) \\ &= q^{2n} \sum_{\mathbf{m} \ (\mathsf{mod} \ q)} \overline{\widehat{\mathbf{1}_{E_1}(\mathbf{m})} \mathbf{1}_{\widehat{E_1} * \widehat{\mathbf{1}_{S_{r,j}}}(\mathbf{m})} = q^{2n} \sum_{\mathbf{m} \ (\mathsf{mod} \ q)} \overline{\widehat{\mathbf{1}_{E_1}(\mathbf{m})} \widehat{\mathbf{1}_{E_1}}(\mathbf{m})} \widehat{\mathbf{1}_{E_1}(\mathbf{m})} (\mathbf{m}) = \mathcal{M} + \mathcal{E}, \end{split}$$

where

$$\mathcal{M} = q^{-n} |E_1| |E_2| |S_{r,j}| \text{ and } \mathcal{E} = q^{2n} \sum_{\mathbf{m} \not\equiv \mathbf{0} \pmod{q}} \overline{\widehat{\mathbf{1}_{E_1}}(\mathbf{m})} \widehat{\mathbf{1}_{E_1}}(\mathbf{m}) \widehat{\mathbf{1}_{S_{r,j}}}(\mathbf{m}).$$

By the condition (ii), one has

$$|\#S_{1,j} - p^{n-1}| \le c_1 p^{n-1}.$$

With the condition (i), we may apply Hensel's lemma to get

$$|\#S_{r,j} - q^{n-1}| \le c_1 q^{n-1}$$

for all r, j with $r \ge 1$ and (j, p) = 1.

Applying Lemma 2.5, one obtains that

$$\left|\widehat{1_{S_{r,j}}}(\mathbf{m})\right| \le \max\{c_2, c_3\} \cdot q^{-1} p^{-\frac{n-1}{2}}.$$

Hence,

$$\begin{aligned} |\mathcal{E}| &\leq q^{2n} \cdot \sup_{\mathbf{m} \neq \mathbf{0} \pmod{q}} \left| \widehat{\mathbf{1}_{S_{r,j}}}(\mathbf{m}) \right| \cdot \left(\sum_{\mathbf{m} \pmod{q}} \left| \overline{\mathbf{1}_{E_1}}(\mathbf{m}) \right|^2 \right)^{1/2} \left(\sum_{\mathbf{m} \pmod{q}} \left| \overline{\mathbf{1}_{E_2}}(\mathbf{m}) \right|^2 \right)^{1/2} \\ &= q^n |E_1|^{1/2} |E_2|^{1/2} \cdot \sup_{\mathbf{m} \neq \mathbf{0} \pmod{q}} \left| \widehat{\mathbf{1}_{S_{r,j}}}(\mathbf{m}) \right| \leq \max\{c_2, c_3\} \cdot \sqrt{|E_1||E_2|} \cdot q^{n-1} p^{-(n-1)/2}. \end{aligned}$$

Combining all bounds, we conclude that

$$\left|\mathcal{N}_{r,j} - \frac{|E_1||E_2|}{q}\right| \le \frac{c_1|E_1||E_2|}{q} + \max\{c_2, c_3\} \cdot \sqrt{|E_1||E_2|} \cdot q^{n-1} p^{-(n-1)/2}$$

So $\mathcal{N}_{r,j} > 0$ if

$$\frac{\sqrt{|E_1||E_2|}}{q^n} > \frac{\max\{c_2, c_3\}}{(1-c_1)p^{\frac{n-1}{2}}}.$$

The proof is completed.

Proof of Corollary 1.5. Let $p > \max\{k(k-1), |a_1|, \dots, |a_n|\}$ be sufficiently large. Then (k, p) = 1 and $(a_i, p) = 1$ for all $1 \le i \le n$.

Let us verify the four conditions in Theorem 1.4 for all $j \in \mathbb{Z}$ with (j, p) = 1. We have

$$(\nabla F)(\mathbf{x}) = (ka_1x_1^{k-1}, \dots, ka_nx_n^{k-1}) \not\equiv \mathbf{0} \pmod{p}$$

when $\mathbf{x} \neq \mathbf{0} \pmod{p}$. Then the condition (i) holds. Moreover, the conditions (ii) and (iii) follow from Lemmas 2.3 and 2.4, respectively. In the following, let us verify condition (iv).

For $\mathbf{m} \neq \mathbf{0} \pmod{p}$ and (j, p) = 1, the cardinality of the set in the condition (iv), denoted by \mathfrak{N} , satisfies that

$$\begin{split} \mathfrak{N} &\leq \sum_{\substack{1 \leq t \leq n \\ m_t \not\equiv 0 \;(\text{mod } p)}} \# \left\{ \mathbf{x} \;(\text{mod } p) : \; \sum_{i=1}^n a_i x_i^k \equiv j \;(\text{mod } p), \\ x_t \not\equiv 0 \;(\text{mod } p), \; m_i a_t x_t^{k-1} \equiv m_t a_i x_i^{k-1} \;(\text{mod } p) \text{ for all } 1 \leq i \leq n \right\}. \end{split}$$

Thus, for the **x** taken into account, we have $x_i^{k-1} \equiv (m_t a_i)^{-1} m_i a_t x_t^{k-1} \pmod{p}$. When k = 2, it follows that

$$\mathfrak{N} \le \sum_{\substack{1 \le t \le n \\ m_t \not\equiv 0 \; (\text{mod } p)}} \# \left\{ x_t \; (\text{mod } p) : \; \Big(\sum_{i=1}^n \left((m_t a_i)^{-1} m_i a_t \right)^2 a_i \Big) x_t^2 \equiv j \; (\text{mod } p) \right\} \le 2n \ll 1.$$

When k = 3, we further have

$$x_i^3 \equiv x_i^{3+p} \equiv \left((m_t a_i)^{-1} m_i a_t x_t^2 \right)^{\frac{3+p}{2}} \equiv \left((m_t a_i)^{-1} m_i a_t \right)^{\frac{3+p}{2}} x_t^3 \,(\text{mod } p)$$

 So

$$\mathfrak{N} \leq \sum_{\substack{1 \leq t \leq n \\ m_t \not\equiv 0 \, (\text{mod } p)}} \# \left\{ x_t \, (\text{mod } p) : \, \left(\sum_{i=1}^n \left((m_t a_i)^{-1} m_i a_t \right)^{\frac{3+p}{2}} a_i \right) x_t^3 \equiv j \, (\text{mod } p) \right\} \leq 3n \ll 1.$$

When $k \ge 4$, we have

$$\mathfrak{N} \leq \sum_{\substack{1 \leq t \leq n \\ m_t \not\equiv 0 \pmod{p}}} \# \left\{ \mathbf{x} \pmod{p} : x_i^{k-1} \equiv (m_t a_i)^{-1} m_i a_t x_t^{k-1} \pmod{p} \text{ for all } 1 \leq i \leq n \right\}.$$

The variable x_t can be chosen mod p, and each x_i $(i \neq t)$ has at most k-1 choices. So

$$\mathfrak{N} \le n \cdot p \cdot k^{n-1} \ll p \ll p^{\frac{n-1}{2}}$$

when $n \geq 3$.

Finally, when n = 2, let us deal with t = 1 without loss of generality. In this situation, one has $m_1, x_1 \neq 0 \pmod{p}$. Note that

$$j \equiv a_1 x_1^k + a_2 x_2^k \equiv a_1 x_1^k + a_2 (m_1 a_2)^{-1} m_2 a_1 x_1^{k-1} x_2 \equiv a_1 x_1^{k-1} (x_1 + m_1^{-1} m_2 x_2) \pmod{p}.$$

If $m_2 \equiv 0 \pmod{p}$, then the above equivalence has at most k-1 solutions in $x_1 \pmod{p}$. Otherwise we have $x_2 \equiv m_1 m_2^{-1} (ja_1^{-1}x_1^{-(k-1)} - x_1) \pmod{p}$. Then

$$(m_1a_2)^{-1}m_2a_1x_1^{k-1} \equiv x_2^{k-1} \equiv \left(m_1m_2^{-1}(ja_1^{-1}x_1^{-(k-1)} - x_1)\right)^{k-1} \pmod{p},$$

i.e.,

$$\begin{split} \big(-(m_1a_2)^{-1}m_2a_1 + (-1)^{k-1}\big)x_1^{k(k-1)} + (-1)^{k-2}m_1^{k-1}m_2^{-(k-1)}(k-1)(ja_1^{-1})x_1^{k(k-2)} \\ + \sum_{i=2}^{k-1}(-1)^{k-1-i}\binom{k-1}{i}(ja_1^{-1})^ix_1^{k(k-1-i)} \equiv 0 \,(\text{mod } p). \end{split}$$

This involves a non-zero polynomial of degree no larger than k(k-1). So there are at most k(k-1) solutions in $x_1 \pmod{p}$. Now we conclude that $\mathfrak{N} \leq 2k(k-1)^2 \ll 1$.

Corollary 1.5 then follows from Theorem 1.4.

Proof of Theorem 1.6. For sufficiently large p, we have

$$(\nabla F)(x) = (k_1 a_1 x_1^{k_1 - 1}, \dots, k_n a_n x_n^{k_n - 1}) \not\equiv \mathbf{0} \pmod{p}$$

when $\mathbf{x} \neq \mathbf{0} \pmod{p}$. Hensel's lemma can be applied on the roots of $F(\mathbf{x}) \equiv j \pmod{p^l}$ $(l \geq 1)$. Combining Lemma 2.3, one can obtain that $|S_{r,j}| \ll q^{n-1}$. Then we deduce similarly as in the proof of Theorem 1.4 that

$$\mathcal{N}_{r,j} \ll_F q^{-1}|E_1||E_2| + q^{2n} \cdot \sum_{\mathbf{m} \not\equiv \mathbf{0} \pmod{q}} \widehat{\widehat{\mathbf{1}_{E_1}}(\mathbf{m})} \widehat{\mathbf{1}_{E_2}}(\mathbf{m}) \widehat{\mathbf{1}_{S_{r,j}}}(\mathbf{m}).$$

Now

$$\sum_{j \pmod{q}} |\mathcal{N}_{r,j}|^2 \ll \Sigma_1 + \Sigma_2,$$

where

$$\Sigma_1 = \sum_{j \pmod{q}} \left(q^{-1} |E_1| |E_2| \right)^2 = q^{-1} |E_1|^2 |E_2|^2,$$

and

$$\Sigma_2 = q^{4n} \sum_{\mathbf{m}, \mathbf{l} \not\equiv \mathbf{0} \, (\mathsf{mod} \, q)} \overline{\widehat{\mathbf{1}_{E_1}}(\mathbf{m})} \widehat{\mathbf{1}_{E_2}}(\mathbf{m}) \widehat{\mathbf{1}_{E_1}}(\mathbf{l}) \overline{\widehat{\mathbf{1}_{E_2}}(\mathbf{l})} \sum_{j \, (\mathsf{mod} \, q)} \widehat{\mathbf{1}_{S_{r,j}}}(\mathbf{m}) \overline{\widehat{\mathbf{1}_{S_{r,j}}}(\mathbf{l})}.$$

Note that

$$\begin{split} \Sigma_{3} &:= \sum_{j \text{ (mod } q)} \widehat{\mathbf{1}_{S_{r,j}}}(\mathbf{m}) \overline{\widehat{\mathbf{1}_{S_{r,j}}}(\mathbf{l})} \\ &= \frac{1}{q^{2n}} \sum_{j \text{ (mod } q)} \sum_{\substack{\mathbf{x} \text{ (mod } q) \\ F(\mathbf{x}) \equiv j \text{ (mod } q)}} \sum_{\substack{\mathbf{y} \text{ (mod } q) \\ F(\mathbf{y}) \equiv j \text{ (mod } q)}} \sum_{\substack{\mathbf{y} \text{ (mod } q) \\ F(\mathbf{y}) \equiv j \text{ (mod } q)}} e_{q}(-\mathbf{m} \cdot \mathbf{x} + \mathbf{l} \cdot \mathbf{y}) \\ &= \frac{1}{q^{2n+1}} \sum_{\substack{\mathbf{x}, \mathbf{y} \text{ (mod } q) \\ F(\mathbf{x}) \equiv F(\mathbf{y}) \text{ (mod } q)}} \sum_{\substack{\mathbf{x}, \mathbf{y} \text{ (mod } q) \\ q \text{ (mod } q)}} e_{q}(-\mathbf{m} \cdot \mathbf{x} + \mathbf{l} \cdot \mathbf{y} + sF(\mathbf{x}) - sF(\mathbf{y})) \\ &= \frac{1}{q^{2n+1}} \sum_{\substack{s \text{ (mod } q) \\ \mathbf{x}, \mathbf{y} \text{ (mod } q)}} \sum_{\substack{\mathbf{x}, \mathbf{y} \text{ (mod } q)}} e_{q}(-\mathbf{m} \cdot \mathbf{x} + \mathbf{l} \cdot \mathbf{y} + sF(\mathbf{x}) - sF(\mathbf{y})), \end{split}$$

where the last two steps follows from the orthogonality and the fact that $\mathbf{m}, \mathbf{l} \neq \mathbf{0} \pmod{q}$. Noting that the inner summation is a product of exponential sums in one variable, one has

$$|\Sigma_3| \leq \frac{1}{q^{2n+1}} \cdot \sum_{s \not\equiv 0 \pmod{q}} \left| \prod_{i=1}^n \sum_{x_i \pmod{q}} e_q \left(-m_i x_i + sa_i x_i^{k_i} \right) \cdot \prod_{i=1}^n \sum_{y_i \pmod{q}} e_q \left(l_i y_i - sa_i y_i^{k_i} \right) \right|.$$

For $(m_i, s), (l_i, s) \neq (0, 0) \pmod{p}$, the condition (5) in Lemma 2.6 follows from Lemma 2.2, and the condition (6) in Lemma 2.6 is confirmed by observing that

$$\#\{x_i (\text{mod } p): -m_i + sa_ik_ix_i^{k_i - 1} \equiv 0 (\text{mod } p)\}, \ \#\{y_i (\text{mod } p): -l_i + sa_ik_iy_i^{k_i - 1} \equiv 0 (\text{mod } p)\} \le k_i - 1.$$

For $s \not\equiv 0 \pmod{q}$, we deduce from Lemma 2.6 that

$$\left| \sum_{x_i \, (\text{mod } q)} e_q \left(-m_i x_i + sa_i x_i^{k_i} \right) \right|, \, \left| \sum_{y_i \, (\text{mod } q)} e_q \left(l_i y_i - sa_i y_i^{k_i} \right) \right| \, \ll_k \, q p^{-1/2} \tag{10}$$

for any $m_i, l_i \pmod{q}$ with $1 \le i \le n$. Moreover, let us assume without loss of generality that $k_1 = k_* = \min_{1 \le i \le n} k_i$. Then

$$\begin{aligned} |\Sigma_{3}| &\ll q^{-2n-1} \cdot \left(qp^{-1/2}\right)^{2n-2} \cdot \sum_{\substack{s \neq 0 \pmod{q}}} \left| \sum_{\substack{x_{1} \pmod{q}}} e_{q} \left(-m_{1}x_{1} + sa_{1}x_{1}^{k_{1}} \right) \cdot \sum_{\substack{y_{1} \pmod{q}}} e_{q} \left(l_{1}y_{1} - sa_{1}y_{1}^{k_{1}} \right) \right| \\ &\leq q^{-3}p^{-n+1} \cdot \left(\sum_{\substack{s \pmod{q}}} \left| \sum_{\substack{x_{1} \pmod{q}}} e_{q} \left(-m_{1}x_{i} + sa_{1}x_{1}^{k_{1}} \right) \right|^{2} \right)^{\frac{1}{2}} \left(\sum_{\substack{s \pmod{q}}} \left| \sum_{\substack{y_{1} \pmod{q}}} e_{q} \left(-l_{1}x_{1} + sa_{1}y_{1}^{k_{1}} \right) \right|^{2} \right)^{\frac{1}{2}} \\ &\leq q^{-3}p^{-n+1} \cdot q^{3-2/k_{*}} = q^{-2/k_{*}}p^{-n+1}, \end{aligned}$$
(11)

where we have applied (10), the Cauchy-Schwartz inequality and Lemma 2.7.

It follows that

$$\begin{split} \Sigma_2 &\ll q^{4n} \cdot q^{-2/k_*} p^{-n+1} \cdot \prod_{i=1,2} \left(\sum_{\mathbf{m} \pmod{q}} |\widehat{\mathbf{1}_{E_i}}(\mathbf{m})|^2 \right)^{1/2} \cdot \prod_{i=1,2} \left(\sum_{\mathbf{l} \pmod{q}} |\widehat{\mathbf{1}_{E_i}}(\mathbf{l})|^2 \right)^{1/2} \\ &\ll q^{4n} \cdot q^{-2/k_*} p^{-n+1} \cdot q^{-2n} |E_1| |E_2| = q^{2n-2/k_*} p^{-n+1} |E_1| |E_2|. \end{split}$$

As a result, we have

$$\sum_{j \pmod{q}} \mathcal{N}_{r,j}^2 \ll q^{-1} |E_1|^2 |E_2|^2 + q^{2n-2/k_*} p^{-n+1} |E_1| |E_2|.$$

Moreover, since

$$|E_1|^2 |E_2|^2 = \left(\sum_{j \in \Delta_{n,r}(E_1, E_2)} \mathcal{N}_{r,j}\right)^2 \le |\Delta_{n,r}(E_1, E_2)| \cdot \sum_{j \ (\text{mod } q)} \mathcal{N}_{r,j}^2$$

we conclude that

$$|\Delta_{n,r}(E_1,E_2)| \geq \frac{|E_1|^2 |E_2|^2}{\sum\limits_{j \ (\text{mod } q)} \mathcal{N}_{r,j}^2} \gg \min\{q, \ q^{-2n+2/k_*} p^{n-1} |E_1| |E_2|\}.$$

In particular, one has $|\Delta_{n,r}(E_1, E_2)| \gg q$ if

$$\frac{\sqrt{|E_1||E_2|}}{q^n} \gg q^{\frac{1}{2} - \frac{1}{k_*}} p^{-\frac{n-1}{2}}.$$

This completes the proof.

Remark 3.1. For the case r = 1, we may use (7) instead of (8) in the above proof, and obtain $\ll p^{-n}$ at the end of (11). As a consequence, the same result (1) as in [11] can be obtained.

4 Extension estimates associated to circles and orbits

Let V be a variety over \mathbb{F}_p^n . The $L^u \to L^{u'}$ Fourier extension problem for V asks us to determine all exponents $1 \le u, u' \le \infty$ such that the following inequality

$$||(fd\sigma)^{\vee}||_{L^{u'}(\mathbb{F}^n_n,dm)} \le C||f||_{L^u(V,d\sigma)},\tag{12}$$

holds for some constant C > 0 and all complex valued functions f on V. Here dm is the counting measure on \mathbb{F}_p^n , $d\sigma$ is the normalized surface measure on V, and

$$(fd\sigma)^{\vee}(m) = \frac{1}{|V|} \sum_{x \in V} f(x) e_p(m \cdot x)$$
$$||f||_{L^u(V,d\sigma)} = \frac{1}{|V|} \sum_{x \in V} |f(x)|^u,$$

and

$$||g||_{L^{u'}(\mathbb{F}_p^n,dm)} = \sum_{m \in \mathbb{F}_p^n} |g(m)|^{u'}$$

for any functions f and g. The $L^u \to L^{u'}$ Fourier extension problem has been studied intensively in the literature for paraboloids, spheres, cones, and homogeneous varieties with applications in several areas of Mathematics including *Discrete Geometry* and *Combinatorial Number Theory*. We refer the interested reader to a series of papers [8, 9, 10, 12, 13, 14] for more discussions.

When V is a circle in the plane \mathbb{F}_p^2 , it has been proved in [2] that

$$||(fd\sigma)^{\vee}||_{L^4(\mathbb{F}^2_p,dm)} \ll ||f||_{L^2(V,d\sigma)}$$

In this paper, to improve Corollary 1.5 in two dimensions, we need to study the finite p-adic ring analog of this estimate. The extension estimates are of independent interest and are expected to have more applications in other topics.

4.1 Extension theorems

In the rest of this section, we always assume that n = 2. Recall that

$$C_{r,j} = \{ \mathbf{x} \in (\mathbb{Z}/p^r \mathbb{Z})^2 : \|\mathbf{x}\| \equiv j \pmod{p^r} \}.$$

We denote by $d\sigma_{r,j}$ the normalised surface measure on $C_{r,j}$. Our results will be presented separately for two cases $p \equiv 3 \pmod{4}$ and $p \equiv 1 \pmod{4}$.

4.1.1 Extension theorems associated to circles

Theorem 4.1. Let $p \equiv 3 \pmod{4}$ and $r \ge 1$ be an integer. Let $j \in \mathbb{Z}/p^r\mathbb{Z}$. Then

$$\Big(\sum_{m \in (\mathbb{Z}/p^r \mathbb{Z})^2} |(f d\sigma_{r,j})^{\vee}(m)|^4\Big)^{1/2} \ll p^{\kappa} \sum_{x \in C_{r,j}} |f(x)|^2,$$

where

$$\kappa = \begin{cases} -\frac{r+1}{2}, & \text{if } j \not\equiv 0 \ (\textit{mod } p), \\ -1, & \text{if } j \equiv 0 \ (\textit{mod } p) \ and \ j \not\equiv 0 \ (\textit{mod } p^r), \\ (-1)^{r-1}, & \text{if } j \equiv 0 \ (\textit{mod } p^r). \end{cases}$$

Theorem 4.2. Let $p \equiv 1 \pmod{4}$ and $r \ge 1$ be an integer. Let $j \in \mathbb{Z}/p^r\mathbb{Z}$.

$$\left(\sum_{m \in (\mathbb{Z}/p^r \mathbb{Z})^2} |(f d\sigma_{r,j})^{\vee}(m)|^4\right)^{1/2} \ll K \sum_{x \in C_{r,j}} |f(x)|^2,$$

where

$$K = \begin{cases} p^{-\frac{r}{2}}, & \text{if } j \not\equiv 0 \ (\textit{mod } p), \\ p^{-1/2} (\textit{ord}_p(j))^{-2}, & \text{if } j \equiv 0 \ (\textit{mod } p). \end{cases}$$

4.1.2 Extension theorems associated to orbits

We now define a more general extension problem. Denote

$$G_r := SO_2(\mathbb{Z}/p^r\mathbb{Z}) = \left\{ \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \pmod{p^r} : a^2 + b^2 \equiv 1 \pmod{p^r} \right\}$$

for $r \geq 1$. The group G_r acts on $(\mathbb{Z}/p^r\mathbb{Z})^2$ naturally by $\mathbf{x} \mapsto \theta \mathbf{x}$, where $\theta \in G_r$ and $\mathbf{x} \in (\mathbb{Z}/p^r\mathbb{Z})^2$. Write the orbit of \mathbf{x} by

$$\operatorname{orb}_r(\mathbf{x}) = \{ \theta \mathbf{x} : \theta \in G_r \},\$$

and the stabilizer of ${\bf x}$ is

$$\mathsf{stab}_r(\mathbf{x}) = \{ \theta \in G_r : \ \theta \mathbf{x} \equiv \mathbf{x} \ (\mathsf{mod} \ p^r) \}$$

Theorem 4.3. Let $p \equiv 3 \pmod{4}$ and $r \geq 1$ be an integer. Let $\mathbf{m} \in (\mathbb{Z}/p^r\mathbb{Z})^2$ be such that $\mathbf{m} \neq \mathbf{0}$. Denote by $d\sigma$ the normalized surface measure on $orb_r(\mathbf{m})$. Then

$$\left(\sum_{x \in (\mathbb{Z}/p^r\mathbb{Z})^2} |(fd\sigma)^{\vee}(x)|^4\right)^{1/2} \ll p^{-\frac{r-3v_{\mathbf{m}}+1}{2}} \sum_{x \in orb_r(\mathbf{m})} |f(x)|^2.$$

Theorem 4.4. Let $p \equiv 1 \pmod{4}$ and $r \geq 1$ be an integer. Let $\mathbf{m} \in (\mathbb{Z}/p^r\mathbb{Z})^2$ be such that $\mathbf{m} \neq \mathbf{0}$. Denote by $d\sigma$ the normalized surface measure on $\operatorname{orb}_r(\mathbf{m})$. Write $\mathbf{m} = p^{v_m} \tilde{\mathbf{m}}$ with $\tilde{\mathbf{m}} \in (\mathbb{Z}/p^{r-v_m}\mathbb{Z})^2$ and $v_{\tilde{\mathbf{m}}} = 0$. Then

$$\Big(\sum_{x\in (\mathbb{Z}/p^r\mathbb{Z})^2} |(fd\sigma)^{\vee}(x)|^4\Big)^{1/2} \ll p^{-\frac{r-3v_{\mathbf{m}}+1-\delta_{\tilde{\mathbf{m}}}}{2}} \sum_{x\in \mathit{orb}_r(\mathbf{m})} |f(x)|^2,$$

where $\delta_{\tilde{\mathbf{m}}} = 1$ if $\|\tilde{\mathbf{m}}\| \equiv 0 \pmod{p}$, and $\delta_{\tilde{\mathbf{m}}} = 0$ if $\|\tilde{\mathbf{m}}\| \neq 0 \pmod{p}$.

4.2 Preliminary lemmas

To prove these extension theorems, we first need to collect and prove a number of preliminary results. All detailed proofs in this subsection are written under the assumption that $p \equiv 3 \pmod{4}$. In the case $p \equiv 1 \pmod{4}$, the identical argument works and will be omitted.

Let p be an prime with $p \equiv 3 \pmod{4}$, $r \geq 1$ be an integer. For any $l \in \mathbb{Z}$, one has (l, p) = 1if and only if $(l, p^r) = 1$. For any $\theta \in G_r$ and $\mathbf{x} \in (\mathbb{Z}/p^r\mathbb{Z})^2$, it is not hard to verify that $\|\theta\mathbf{x}\| \equiv \|\mathbf{x}\| \pmod{p^r}$. Moreover, we have $v_{\theta\mathbf{x}} \geq v_{\mathbf{x}}$. Since θ is invertible, one obtains that $v_{\theta\mathbf{x}} = v_{\mathbf{x}}$.

It is well-known (see [5] for example) that

$$|C_{1,j}| = |G_1| = p + 1, \quad (j \in (\mathbb{Z}/p\mathbb{Z})^*), \qquad |C_{1,0}| = 1$$

when $p \equiv 3 \pmod{4}$. In particular, $C_{1,0} = \{\mathbf{0}\}$. For any $\mathbf{x} \in \mathbb{Z}/p^r\mathbb{Z}$, it follows that $\|\mathbf{x}\| \equiv 0 \pmod{p}$, if and only if $\mathbf{x} \equiv \mathbf{0} \pmod{p}$, if and only if $v_{\mathbf{x}} > 0$. Or equivalently, we have $\mathbf{x} \in (\mathbb{Z}/p^r\mathbb{Z})^*$ if and only if $\mathbf{x} \not\equiv \mathbf{0} \pmod{p}$, if and only if $\|\mathbf{x}\| \not\equiv 0 \pmod{p}$, if and only if $\|\mathbf{x}\| \not\equiv 0 \pmod{p}$, if and only if $v_{\mathbf{x}} = 0$.

Lemma 4.5. We have $|G_r| = p^r (1 + 1/p)$.

Proof. For r = 1, we have $|G_1| = p + 1$. Now consider the circumstances that $r \ge 2$. For any $\theta = \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \in G_r$, there is some $\theta_0 = \begin{bmatrix} a_0 & -b_0 \\ b_0 & a_0 \end{bmatrix} \in G_1$ such that $\theta \equiv \theta_0 \pmod{p}$. Applying Lemma 2.1 to (a_0, b_0) and the polynomial $F(x, y) = x^2 + y^2 - 1$. Then $(\nabla F)(a_0, b_0) = (2a_0, 2b_0) \neq \mathbf{0} \pmod{p}$, since $a_0^2 + b_0^2 \equiv 1 \pmod{p}$. Thus,

$$\#\left\{(z_1, z_2) \,(\mathsf{mod} \ p^{r-1}): \,(a_0 + pz_1)^2 + (b_0 + pz_2)^2 \equiv 1 \,(\mathsf{mod} \ p^r)\right\} = p^{r-1}.$$

It follows that

$$|G_r| = p^{r-1}|G_1| = p^r(1+1/p).$$

Lemma 4.6. For any $\mathbf{0} \neq \mathbf{x} \in (\mathbb{Z}/p^r\mathbb{Z})^2$, let $\mathbf{x} = p^{v_{\mathbf{x}}}\tilde{\mathbf{x}}$ for some $\tilde{\mathbf{x}} \in (\mathbb{Z}/p^{r-v_{\mathbf{x}}}\mathbb{Z})^2$ with $v_{\tilde{\mathbf{x}}} = 0$. Then

$$\textit{orb}_r(\mathbf{x}) = \left\{ p^{v_{\mathbf{x}}} \theta_0 \tilde{\mathbf{x}} : \, \theta_0 \in G_{r-v_{\mathbf{x}}} \right\} = p^{v_{\mathbf{x}}} C_{r-v_{\mathbf{x}}, \|\tilde{\mathbf{x}}\|}.$$

And

$$|stab_r(\mathbf{x})| = p^{v_{\mathbf{x}}}, \quad |orb_r(\mathbf{x})| = p^{r-v_{\mathbf{x}}}(1+1/p)$$

Proof. Write $v = v_{\mathbf{x}}$ for simplicity. Here $0 \le v \le r - 1$. Let $\tilde{\mathbf{x}} = (\tilde{x_1}, \tilde{x_2})$. The equation $\theta \mathbf{x} \equiv \mathbf{x} \pmod{p^r}$ is equivalent to

$$\begin{bmatrix} a-1 & -b \\ b & a-1 \end{bmatrix} \begin{bmatrix} p^v \tilde{x_1} \\ p^v \tilde{x_2} \end{bmatrix} \equiv \begin{bmatrix} 0 \\ 0 \end{bmatrix} \pmod{p^r},$$
(13)

or equivalently,

$$\begin{bmatrix} \tilde{x_1} & -\tilde{x_2} \\ \tilde{x_2} & \tilde{x_1} \end{bmatrix} \begin{bmatrix} a-1 \\ b \end{bmatrix} \equiv \begin{bmatrix} 0 \\ 0 \end{bmatrix} \pmod{p^{r-v}}.$$
(14)

Since $v_{\tilde{\mathbf{x}}} = 0$, one has $\tilde{x_1}^2 + \tilde{x_2}^2 \not\equiv 0 \pmod{p}$ (recalling that $p \equiv 3 \pmod{4}$). The coefficient matrix is invertible and

$$\begin{bmatrix} a \\ b \end{bmatrix} \equiv \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \pmod{p^{r-v}}.$$

By Lemma 2.1, the number of $(a, b) \in (\mathbb{Z}/p^r\mathbb{Z})^2$ satisfying the above equivalence and $a^2 + b^2 \equiv 1 \pmod{p^r}$ is exactly p^v . So

$$|\mathsf{stab}_r(\mathbf{x})| = p^v.$$

It then follows that $|\operatorname{orb}_r(\mathbf{x})| = |G_r|/|\operatorname{stab}_r(\mathbf{x})| = p^{r-v}(1+1/p).$

Moreover, for any $\theta \in G_r$, there is some $\theta_0 \in G_{r-v}$ such that $\theta \equiv \theta_0 \pmod{p^{r-v}}$. It can be verified that $\theta \tilde{\mathbf{x}} \equiv \theta_0 \tilde{\mathbf{x}} \pmod{p^{r-v}}$. So

$$\mathsf{orb}_r(\mathbf{x}) = \left\{ \theta(p^v \tilde{\mathbf{x}}) : \, \theta \in G_r \right\} = \left\{ p^v \theta_0 \tilde{\mathbf{x}} : \, \theta_0 \in G_{r-v} \right\}.$$

Note that $|G_{r-v}| = p^{r-v}(1+1/p)$ by Lemma 4.5, the elements on the right-hand side of above formula give different members of the orbit. Furthermore, one has $\|\tilde{\mathbf{x}}\| \in (\mathbb{Z}/p^{r-v}\mathbb{Z})^*$, since $v_{\tilde{\mathbf{x}}} = 0$ and $p \equiv 3 \pmod{4}$. We have $\{\theta_0 \tilde{\mathbf{x}} : \theta_0 \in G_{r-v}\} = C_{r-v,\|\tilde{\mathbf{x}}\|}$. The proof is completed.

Lemma 4.7. Let $p \equiv 3 \pmod{4}$. Let j be an integer with $ord_p(j) = v$. Then

$$|C_{r,j}| = \begin{cases} p^r (1+1/p), & \text{if } 0 \le v < r \text{ and } v \text{ is even} \\ p^{2\lfloor r/2 \rfloor}, & \text{if } v = r, \\ 0, & \text{otherwise.} \end{cases}$$

More concretely, we have

$$C_{r,j} = \begin{cases} \left\{ p^{u} \mathbf{z} + p^{r-u} \mathbf{w} : \, \mathbf{z} \in C_{r-2u,\tilde{j}}, \, \mathbf{w} \in \mathbb{Z}/p^{u}\mathbb{Z} \right\}, & \text{if } j = p^{2u}\tilde{j}, \, 0 \le u \le \lfloor \frac{r-1}{2} \rfloor, \, \tilde{j} \in (\mathbb{Z}/p^{r-2u}\mathbb{Z})^{*} \\ \left\{ p^{\lceil r/2 \rceil} \mathbf{w} : \, \mathbf{w} \in \mathbb{Z}/p^{\lfloor r/2 \rfloor}\mathbb{Z} \right\}, & \text{if } j = 0, \\ \emptyset, & \text{otherwise.} \end{cases}$$

Proof. We first consider the case $j \in (\mathbb{Z}/p^r\mathbb{Z})^*$. By applying Lemma 2.1, similar arguments as previous show that

$$|C_{r,j}| = p^{r-1}|C_{1,j}| = p^r(1+1/p).$$

Next, consider the circle $C_{r,0}$ $(r \ge 2)$. Since $C_{1,0} = \{0\}$, we can write

$$C_{r,0} = \{ p\mathbf{y} : \mathbf{y} \in C'_{r,0} \}, \quad C'_{r,0} = \big\{ \mathbf{y} \, (\text{mod } p^{r-1}) : \, \|p\mathbf{y}\| \equiv 0 \, (\text{mod } p^r) \big\}.$$

When r = 2, one sees that $C_{2,0} = \{p\mathbf{y} : \mathbf{y} \in (\mathbb{Z}/pZ)^2\}$ and $|C_{2,0}| = p^2$. When $r \ge 3$, we further write $\mathbf{y} = \mathbf{z} + p^{r-2}\mathbf{w}$ with $\mathbf{z} \in \mathbb{Z}/p^{r-2}\mathbb{Z}$ and $\mathbf{w} \in \mathbb{Z}/pZ$. The condition $\|p\mathbf{y}\| \equiv \mathbf{0} \pmod{p^r}$ holds if and only if $\|\mathbf{z} + p^{r-2}\mathbf{w}\| = \|\mathbf{y}\| \equiv \mathbf{0} \pmod{p^{r-2}}$, if and only if $\|\mathbf{z}\| \equiv \mathbf{0} \pmod{p^{r-2}}$. It follows that

$$C_{r,0} = \left\{ p(\mathbf{z} + p^{r-2}\mathbf{w}) : \mathbf{z} \in C_{r-2,0}, \, \mathbf{w} \in (\mathbb{Z}/p\mathbb{Z})^2 \right\},\$$

and $|C_{r,0}| = p^2 |C_{r-2,0}|.$

By induction, we conclude that

$$C_{r,0} = \left\{ p^{\lceil r/2 \rceil} \mathbf{w} : \mathbf{w} \in \mathbb{Z}/p^{\lfloor r/2 \rfloor} \mathbb{Z} \right\}$$

and $|C_{r,0}| = p^{2\lfloor r/2 \rfloor}$.

Now, let us consider the case $j = p^v \tilde{j}$, where $1 \le v \le r-1$ and $\tilde{j} \in (\mathbb{Z}/p^{r-v}\mathbb{Z})^*$. For any $\mathbf{x} \in C_{r,j}$, it satisfies that $\|\mathbf{x}\| \equiv 0 \pmod{p}$. Recalling that $C_{1,0} = \{\mathbf{0}\}$, one has $\mathbf{x} \equiv \mathbf{0} \pmod{p}$. So

$$C_{r,j} = \{ p\mathbf{y} : \, \mathbf{y} \in C'_{r,j} \}, \quad C'_{r,j} = \big\{ \mathbf{y} \, (\text{mod } p^{r-1}) : \, \|p\mathbf{y}\| \equiv p^v \tilde{j} \, (\text{mod } p^r) \big\}.$$

When v = 1, it is easy to see that $C'_{r,j} = \emptyset$. When $v \ge 2$, which means $r \ge 3$, similar arguments as previous show that

$$C_{r,p^{v}\tilde{j}} = \left\{ p(\mathbf{z} + p^{r-2}\mathbf{w}) : \, \mathbf{z} \in C_{r-2,p^{v-2}\tilde{j}}, \, \mathbf{w} \in (\mathbb{Z}/p\mathbb{Z})^2 \right\},$$

and $|C_{r,p^v\tilde{j}}|=p^2\cdot |C_{r-2,p^{v-2}\tilde{j}}|.$ By induction, it follows that

$$|C_{r,p^v\tilde{j}}| = \begin{cases} p^r(1+1/p), & \text{if } v \text{ is even}, \\ 0, & \text{if } v \text{ is odd} \end{cases}$$

when $1 \le v \le r - 1$. Indeed,

$$C_{r,p^{2u}\tilde{j}} = \left\{ p^{u} \mathbf{z} + p^{r-u} \mathbf{w} : \, \mathbf{z} \in C_{r-2u,\tilde{j}}, \, \mathbf{w} \in (\mathbb{Z}/p^{u}\mathbb{Z})^{2} \right\}$$
(15)

when $r \geq 3$ and $1 \leq u \leq (r-1)/2$.

Remark: When $p \equiv 3 \pmod{4}$, we have the disjoint union

$$(\mathbb{Z}/p^{r}\mathbb{Z})^{2} = \left(\bigcup_{u=0}^{\lfloor (r-1)/2 \rfloor} \bigcup_{\tilde{j} \in (\mathbb{Z}/p^{r-2u}\mathbb{Z})^{*}} C_{r,p^{2u}\tilde{j}}\right) \cup C_{r,0},$$

whose cardinalities give

$$p^{2r} = \sum_{u=0}^{\lfloor (r-1)/2 \rfloor} p^r (1+1/p) \cdot p^{r-2u} (1-1/p) + p^{2\lfloor r/2 \rfloor}$$
$$= \sum_{u=0}^{\lfloor (r-1)/2 \rfloor} (p^{2r-2u} - p^{2r-2u-2}) + p^{2\lfloor r/2 \rfloor}.$$

Lemma 4.8. Let $j \in \mathbb{Z}/p^r\mathbb{Z}$. For any $\mathbf{0} \neq \mathbf{z} \in (\mathbb{Z}/p^r\mathbb{Z})^2$, we have

$$\#\big\{(\mathbf{x},\mathbf{y})\in C^2_{r,j}: \mathbf{x}-\mathbf{y}\equiv \mathbf{z} \,(\operatorname{mod} \, p^r)\big\} \,\leq\, \begin{cases} 2p^{r-1}, & \text{if} \, j\not\equiv 0 \,(\operatorname{mod} \, p), \\ 2p^{2(r-1)}, & \text{if} \, j\equiv 0 \,(\operatorname{mod} \, p). \end{cases}$$

Remark: The set on the left-hand side is empty if $\operatorname{ord}_p(j) > 2v_z$.

Proof. The set on the left-hand side involves system of congruences

$$\begin{aligned}
x_1^2 + x_2^2 &\equiv j, \\
y_1^2 + y_2^2 &\equiv j, \\
x_1 - y_1 &\equiv z_1, \\
x_2 - y_2 &\equiv z_2.
\end{aligned}$$
(16)

We first consider the system (16) modulo p. One sees that

$$2y_1z_1 + 2y_2z_2 + z_1^2 + z_2^2 \equiv 0 \pmod{p}.$$

When $\mathbf{z} \neq \mathbf{0} \pmod{p}$, we assume without loss of generality that $z_1 \neq 0 \pmod{p}$. Inserting $y_1 \equiv -(2z_1)^{-1}(2y_2z_2 + z_1^2 + z_2^2) \pmod{p}$ into $y_1^2 + y_2^2 \equiv j \pmod{p}$, one obtains

$$z_1^{-2}(z_1^2+z_2^2)y_2^2+z_1^{-2}z_2(z_1^2+z_2^2)y_2+4^{-1}z_1^{-2}(z_1^2+z_2^2)^2\equiv j \pmod{p}$$

When $p \equiv 3 \pmod{4}$, $\mathbf{z} \not\equiv \mathbf{0} \pmod{p}$ implies that $z_1^2 + z_2^2 \not\equiv 0 \pmod{p}$. The left-hand side of above congruence has degree 2. So there are at most two solutions in y_2 . Moreover, each choice of y_2 exactly determines the choices of x_1 , y_1 and x_2 .

When $\mathbf{z} \equiv \mathbf{0} \pmod{p}$, the number of the system (16) modulo p is $|C_{1,j}|$, which equals p+1 when $j \not\equiv 0 \pmod{p}$ and equals one otherwise.

In the following, we will apply Hensel's lemma. The Jacobi matrix, in (x_1, y_1, x_2, y_2) , is given by

$$\begin{bmatrix} 2x_1 & 0 & 2x_2 & 0 \\ 0 & 2y_1 & 0 & 2y_2 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 \end{bmatrix}$$

By elementary operations, we obtain

$$\begin{bmatrix} 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & x_1 & x_2 \\ 0 & 0 & y_1 & y_2 \end{bmatrix}$$

The rank of above Jacobi matrix, modulo p, is at least 2. Moreover, when $j \neq 0$, (mod p), it satisfies that $\mathbf{x}, \mathbf{y} \neq \mathbf{0} \pmod{p}$ and the rank is at least 3. By Lemma 2.1, one concludes that

$$\#\big\{(\mathbf{x},\mathbf{y})\in C_{r,j}^2:\,\mathbf{x}-\mathbf{y}\equiv\mathbf{z}\,(\mathrm{mod}\,\,p^r)\big\} \leq \begin{cases} 2p^{r-1}, & \text{if}\,\,j\not\equiv 0\,(\mathrm{mod}\,\,p), \text{ and}\,\,\mathbf{z}\not\equiv \mathbf{0}\,(\mathrm{mod}\,\,p),\\ 2p^{2(r-1)}, & \text{if}\,\,j\equiv 0\,(\mathrm{mod}\,\,p) \text{ and}\,\,\mathbf{z}\not\equiv \mathbf{0}\,(\mathrm{mod}\,\,p),\\ p^{2(r-1)}, & \text{if}\,\,j\equiv 0\,(\mathrm{mod}\,\,p) \text{ and}\,\,\mathbf{z}\equiv \mathbf{0}\,(\mathrm{mod}\,\,p). \end{cases}$$

Finally, let us deal with the case $j \not\equiv 0 \pmod{p}$ and $\mathbf{z} \equiv \mathbf{0} \pmod{p}$. Assume that $\mathbf{z} = p^k \tilde{\mathbf{z}}$ with $k = v_{\mathbf{z}}$ and $\tilde{\mathbf{z}} \in (\mathbb{Z}/p^{r-k}\mathbb{Z})^2$. Here $1 \leq v < r$, since $\mathbf{z} \neq \mathbf{0}$. Note that, for any solution (\mathbf{x}, \mathbf{y}) to the system (16) modulo p^r , it also satisfies (16) modulo p^k or p^{k+1} . For the former situation, one deduces that $\mathbf{x} \equiv \mathbf{y} \pmod{p^k}$ since $\mathbf{z} \equiv \mathbf{0} \pmod{p^k}$. For the latter situation, similar arguments as previous shows that

$$(y_1 + p^k \tilde{z_1})^2 + (y_2 + p^k \tilde{z_2})^2 \equiv j \equiv y_1^2 + y_2^2, \, (\mathrm{mod} \ p^{k+1}),$$

i.e., $2y_1\tilde{z_1} + 2y_2\tilde{z_2} \equiv 0 \pmod{p}$. Noting that $\tilde{z_1} \not\equiv 0 \pmod{p}$. We have

$$x_1 \equiv y_1 \equiv y_2 \tilde{z_2} \tilde{z_1}^{-1} \equiv x_2 \tilde{z_2} \tilde{z_1}^{-1}, \pmod{p}$$

Now any solution (\mathbf{x}, \mathbf{y}) to the system (16) modulo p^r satisfies that

$$\begin{cases} x_1^2 + x_2^2 \equiv j, \\ x_1 \equiv x_2 \tilde{z_2} \tilde{z_1}^{-1}, \\ x_1 \equiv y_1, \\ x_2 \equiv y_2, \end{cases} \pmod{p}.$$

It is not hard to see that there are at most 2 solutions modulo p, since $\tilde{z_1}^2 + \tilde{z_2}^2 \neq 0 \pmod{p}$. Now we lift these solutions modulo p to solutions to (16) modulo p^r . By applying Hensel's lemma with the rank of Jacobian matrix being 3, we conclude that

$$\#\{(\mathbf{x}, \mathbf{y}) \in C^2_{r, j} : \mathbf{x} - \mathbf{y} \equiv \mathbf{z} \,(\mathsf{mod} \, p^r)\} \le 2p^{r-1}.$$

Lemma 4.9. Let $\mathbf{m}, \mathbf{z} \in (\mathbb{Z}/p^r\mathbb{Z})^2$ be such that $\mathbf{m}, \mathbf{z} \neq \mathbf{0}$. Then

$$\#\big\{(\mathbf{x},\mathbf{y})\in(\textit{orb}_r(\mathbf{m}))^2:\,\mathbf{x}-\mathbf{y}\equiv\mathbf{z}\,(\textit{mod}\,\,p^r)\big\}\,\leq\,2p^{r-v_{\mathbf{m}}-1}.$$

Proof. Write $v = v_{\mathbf{m}}$ and $\mathbf{m} = p^{v}\tilde{\mathbf{m}}$. Here $0 \leq v \leq r-1$, $\tilde{\mathbf{m}} \in (\mathbb{Z}/p^{r-v_{\mathbf{m}}}\mathbb{Z})^{2}$ and $v_{\tilde{\mathbf{m}}} = 0$. Write $\|\tilde{\mathbf{m}}\| = \tilde{j}$ for simplicity. Then $\tilde{j} \in (\mathbb{Z}/p^{r-v}\mathbb{Z})^{*}$. By Lemma 4.6, we have $\mathsf{orb}(\mathbf{m}) = p^{v}C_{\tilde{j},r-v}$. Write $\mathbf{x} = p^{v}\tilde{\mathbf{x}}$ and $\mathbf{y} = p^{v}\tilde{\mathbf{y}}$ with $\tilde{\mathbf{x}}, \tilde{\mathbf{y}} \in C_{\tilde{j},r-v}$. When $\mathbf{z} \not\equiv \mathbf{0} \pmod{p^{v}}$, it is obvious that

$$\#\{(\mathbf{x},\mathbf{y})\in (\operatorname{orb}_r(\mathbf{m}))^2: \, \mathbf{x}-\mathbf{y}\equiv \mathbf{z}\,(\operatorname{mod}\,p^r)\}=0.$$

When $\mathbf{z} \equiv \mathbf{0} \pmod{p^v}$, we write $\mathbf{z} = p^v \tilde{\mathbf{z}}$ for some $\mathbf{0} \neq \tilde{\mathbf{z}} \in (\mathbb{Z}/p^{r-v}\mathbb{Z})^2$. It follows that

$$\#\big\{(\mathbf{x},\mathbf{y})\in(\operatorname{orb}_{r}(\mathbf{m}))^{2}:\,\mathbf{x}-\mathbf{y}\equiv\mathbf{z}\,(\operatorname{mod}\,p^{r})\big\}=\#\big\{(\tilde{\mathbf{x}},\tilde{\mathbf{y}})\in(C_{\tilde{j},r-v})^{2}:\,\tilde{\mathbf{x}}-\tilde{\mathbf{y}}\equiv\tilde{\mathbf{z}}\,(\operatorname{mod}\,p^{r-v})\big\}.$$

By Lemma 4.8, the above quantity can be bounded by $2p^{r-v-1}$.

With the same argument, we obtain similar results for the case $p \equiv 1 \pmod{4}$.

Lemma 4.10. For any $0 \neq \mathbf{x} \in (\mathbb{Z}/p^r\mathbb{Z})^2$, let $\mathbf{x} = p^{v_{\mathbf{x}}}\tilde{\mathbf{x}}$ for some $\tilde{\mathbf{x}} \in (\mathbb{Z}/p^{r-v_{\mathbf{x}}}\mathbb{Z})^2$ with $v_{\tilde{\mathbf{x}}} = 0$. Then

$$|stab_r(\mathbf{x})| = p^{v_{\mathbf{x}}}, \quad |orb_r(\mathbf{x})| = p^{r-v_{\mathbf{x}}}(1-1/p).$$

And $orb_r(\mathbf{x}) = p^{v_{\mathbf{x}}} orb_{r-v}(\tilde{\mathbf{x}}).$

Lemma 4.11. Let $p \equiv 1 \pmod{4}$. Let j be an integer with $ord_p(j) = v$. Then

$$|C_{r,j}| = \begin{cases} (v+1)(p-1)p^{r-1}, & \text{if } 0 \le v < r, \\ (rp+p-r)p^{r-1}, & \text{if } v = r \end{cases}$$

Lemma 4.12. For $p \equiv 1 \pmod{4}$ be a prime, $r \geq 1$ be an integer. Let $j \in \mathbb{Z}/p^r\mathbb{Z}$. For any $\mathbf{0} \neq \mathbf{z} \in (\mathbb{Z}/p^r\mathbb{Z})^2$, we have

$$\#\{(\mathbf{x},\mathbf{y})\in C^2_{r,j}\colon \mathbf{x}-\mathbf{y}\equiv \mathbf{z} \,(\textit{mod } p^r)\}\leq 2p^{(1+\delta_j)(r-1)+\delta_{\mathbf{z}}},$$

where

$$\delta_j = \begin{cases} 1, & j \equiv 0 \pmod{p}, \\ 0, & j \not\equiv 0 \pmod{p}, \end{cases}, \quad \delta_{\mathbf{z}} = \begin{cases} 1, & \|\mathbf{z}\| \equiv 0 \pmod{p}, \\ 0, & \|\mathbf{z}\| \not\equiv 0 \pmod{p}. \end{cases}$$

Lemma 4.13. For $p \equiv 1 \pmod{4}$ be a prime, $r \geq 1$ be an integer and $\mathbf{m} = p^{v_{\mathbf{m}}} \tilde{\mathbf{m}} \in (\mathbb{Z}/p^r \mathbb{Z})^2$ with $v_{\tilde{\mathbf{m}}} = 0$, let $\mathbf{z} \in (\mathbb{Z}/p^r \mathbb{Z})^2$ we have

$$\#\{(\mathbf{x},\mathbf{y})\in(\textit{orb}_r(\mathbf{m}))^2\colon\mathbf{x}-\mathbf{y}\equiv\mathbf{z}(\textit{mod }p^r)\}\ll\begin{cases}p^{r-\upsilon_{\mathbf{m}}-1}, & \textit{if }\|\tilde{\mathbf{m}}\|\neq0\,(\textit{mod }p),\\p^{r-\upsilon_{\mathbf{m}}}, & \textit{if }\|\tilde{\mathbf{m}}\|\equiv0\,(\textit{mod }p).\end{cases}$$

Note that the second bound is trivial since it is almost the same as the size of $orb_r(\mathbf{m})$.

4.3 **Proof of extension theorems**

Theorems 4.1 and 4.3 are proved in the same way. Let V be a variety with $V \subseteq (\mathbb{Z}/p^r\mathbb{Z})^2$, which can be $C_{r,j}$ or $\operatorname{orb}_r(\mathbf{m})$, on which the normalized surface measure is denoted by $d\sigma$. For simplicity, we denote $q = p^r$ in this subsection.

We have

$$\sum_{\mathbf{m} \pmod{q}} |(fd\sigma)^{\vee}(\mathbf{m})|^4 = \sum_{\mathbf{m} \pmod{q}} \left| \frac{1}{|V|} \sum_{\mathbf{x} \in V} f(\mathbf{x}) e_q(\mathbf{m} \cdot \mathbf{x}) \right|^4$$
$$= \frac{q^2}{|V|^4} \sum_{\substack{\xi, \xi', \eta, \eta' \in V\\ \xi - \eta \equiv \xi' - \eta' \pmod{q}}} f(\xi) f(\xi') \overline{f(\eta)} \overline{f(\eta')} = \frac{q^2}{|V|^4} \sum_{\substack{\zeta \pmod{q}}} \left| \sum_{\substack{\xi, \eta \in V\\ \xi - \eta \equiv \zeta \pmod{q}}} f(\xi) \overline{f(\eta)} \right|^2.$$

For $\zeta \equiv \mathbf{0} \pmod{q}$, we have

$$\Big|\sum_{\substack{\xi,\eta\in V\\\xi\equiv\eta \pmod{q}}} f(\xi)\overline{f(\eta)}\Big|^2 = \Big(\sum_{\xi\in V} |f(\xi)|^2\Big)^2.$$

For $\zeta \not\equiv \mathbf{0} \pmod{p^r}$, the Cauchy-Schwarz inequality implies

$$\sum_{\zeta \not\equiv \mathbf{0} \, (\mathrm{mod} \, q)} \Big| \sum_{\substack{\xi, \eta \in V \\ \xi - \eta \equiv \zeta \, (\mathrm{mod} \, q)}} f(\xi) \overline{f(\eta)} \Big|^2 \leq \sum_{\zeta \not\equiv \mathbf{0} \, (\mathrm{mod} \, p^r)} \Big(\sum_{\substack{\xi, \eta \in V \\ \xi - \eta \equiv \zeta \, (\mathrm{mod} \, q)}} 1\Big) \Big(\sum_{\substack{\xi, \eta \in V \\ \xi - \eta \equiv \zeta \, (\mathrm{mod} \, q)}} |f(\xi)|^2 |f(\eta)|^2 \Big).$$

Assuming that $\sum_{\xi,\eta\in V\,\xi-\eta\equiv\zeta\,(\mathsf{mod}\ q)} 1 \ll U$, then

$$\sum_{\zeta \not\equiv \mathbf{0} \pmod{q}} \Big| \sum_{\substack{\xi, \eta \in V \\ \xi - \eta \equiv \zeta \pmod{q}}} f(\xi) \overline{f(\eta)} \Big|^2 \ll U \sum_{\substack{\zeta \pmod{p^r}}} \Big(\sum_{\substack{\xi, \eta \in V \\ \xi - \eta \equiv \zeta \pmod{q}}} |f(\xi)|^2 |f(\eta)|^2 \Big) = U \left(\sum_{\xi \in V} |f(\xi)|^2 \right)^2.$$

It follows that

$$\left(\sum_{m \, (\text{mod } q)} |(f d\sigma_{r,j})^{\vee}(m)|^4\right)^{1/2} \ll \frac{q U^{1/2}}{|V|^2} \sum_{x \in V} |f(x)|^2.$$

On the one hand, to get the bound U, we use Lemmas 4.8 and 4.9. On the other hand, Lemma 4.7 gives estimates on the size of V. Hence, the theorems follow.

5 Proof of Theorem 1.8 and 1.9

These two theorems are proved by the same argument.

By Lemma 4.7, it is clear that the number of pairs $(\mathbf{x}, \mathbf{y}) \in E \times E$ such that $||\mathbf{x}-\mathbf{y}|| \notin (\mathbb{Z}/p^r \mathbb{Z})^*$ is much smaller than $|E|^2$, since $|E| \gg p^{2r-1}$. By the Cauchy-Schwarz inequality, we have

$$|E|^{4} \ll \Big(\sum_{\substack{\mathbf{x},\mathbf{y}\in E\\ \|\mathbf{x}-\mathbf{y}\|\in (\mathbb{Z}/p^{r}\mathbb{Z})^{*}}} 1\Big)^{2} = \Big(\sum_{\substack{j\in\Delta_{2,r}(E)\cap (\mathbb{Z}/p^{r}\mathbb{Z})^{*} \ \mathbf{x}\in E}} \sum_{\substack{\mathbf{y}\in E\\ \|\mathbf{x}-\mathbf{y}\|\equiv j \pmod{p^{r}}}} 1\Big)^{2}$$
$$\leq |\Delta_{2,r}(E)| \cdot |E| \cdot \sum_{\substack{j\neq 0 \pmod{p} \ \mathbf{x}\in E}} \sum_{\substack{\mathbf{y}\in E\\ \|\mathbf{x}-\mathbf{y}\|\equiv j \pmod{p^{r}}}} 1\Big)^{2}.$$

$$|\Delta_{2,r}(E)| \gg \frac{|E|^3}{\mathcal{N}},$$

where

$$\mathcal{N} := \sum_{j \not\equiv 0 \pmod{p}} \sum_{\mathbf{x} \in E} \Big(\sum_{\substack{\mathbf{y} \in E \\ \|\mathbf{x} - \mathbf{y}\| \equiv j \pmod{p^r}}} 1 \Big)^2 = \sum_{x \in E} \sum_{j \not\equiv 0 \pmod{p}} \Big(\sum_{\mathbf{z} \in C_{r,j}} 1_E(\mathbf{x} - \mathbf{z}) \Big)^2.$$

For any given $\mathbf{x} \in E$, we have

$$\sum_{\substack{j \neq 0 \pmod{p}}} \left(\sum_{\mathbf{z} \in C_{r,j}} 1_E(\mathbf{x} - \mathbf{z})\right)^2 = \sum_{\substack{j \neq 0 \pmod{p} \ \mathbf{y}, \mathbf{z} \in C_{r,j}}} \sum_{\substack{1_E(\mathbf{x} - \mathbf{y}) 1_E(\mathbf{x} - \mathbf{z})} 1_E(\mathbf{x} - \mathbf{z})$$
$$\leq \sum_{\substack{j \neq 0 \pmod{p} \ \mathbf{z} \in C_{r,j}}} \sum_{\substack{1_E(\mathbf{x} - \mathbf{z}) \ \mathcal{D}_{\theta} \in G_r}} 1_E(\mathbf{x} - \theta \mathbf{z}) \leq \sum_{\substack{\theta \in G_r \ ||\mathbf{z}|| \neq 0 \pmod{p^r}}} 1_E(\mathbf{x} - \theta \mathbf{z}) 1_E(\mathbf{x} - \theta \mathbf{z}).$$

Here the terms with $\|\mathbf{z}\| \neq 0 \pmod{p^r}$ but $\|\mathbf{z}\| \equiv 0 \pmod{p}$ can also be involved, since the summands are non-negative. In the next step, we write

$$\begin{split} \sum_{\mathbf{z} \pmod{p^r}} \sum_{\theta \in G_r} \mathbf{1}_E(\mathbf{x} - \mathbf{z}) \mathbf{1}_E(\mathbf{x} - \theta \mathbf{z}) \\ &= \sum_{\mathbf{z} \pmod{p^r}} \sum_{\theta \in G_r} \sum_{\mathbf{m} \pmod{p^r}} \widehat{\mathbf{1}_E}(\mathbf{m}) e_{p^r}(\mathbf{m} \cdot (\mathbf{x} - \mathbf{z})) \overline{\sum_{\mathbf{m}' \pmod{p^r}} \widehat{\mathbf{1}_E}(\mathbf{m}') e_{p^r}(\mathbf{m}' \cdot (\mathbf{x} - \theta \mathbf{z}))} \\ &= \sum_{\theta \in G_r} \sum_{\mathbf{m} \pmod{p^r}} \sum_{\mathbf{m}' \pmod{p^r}} \widehat{\mathbf{1}_E}(\mathbf{m}) \overline{\widehat{\mathbf{1}_E}(\mathbf{m}')} e_{p^r} \left((\mathbf{m} - \mathbf{m}') \cdot \mathbf{x} \right) \sum_{\mathbf{z} \pmod{p^r}} e_{p^r} \left(- (\mathbf{m} \cdot \mathbf{z} - \mathbf{m}' \cdot (\theta \mathbf{z})) \right) . \end{split}$$

Note that

$$\sum_{\mathbf{z} \pmod{p^r}} e_{p^r} \left(-(\mathbf{m} \cdot \mathbf{z} - \mathbf{m}' \cdot (\theta \mathbf{z}) \right) = \sum_{\mathbf{z} \pmod{p^r}} e_{p^r} \left(-(\mathbf{m} - \theta^{-1} \mathbf{m}') \cdot \mathbf{z} \right) = \begin{cases} p^{2r}, & \text{if } \mathbf{m}' = \theta \mathbf{m}, \\ 0, & \text{otherwise.} \end{cases}$$

One deduces that

$$\begin{split} \sum_{\mathbf{z} \pmod{p^r}} \sum_{\theta \in G_r} \mathbf{1}_E(\mathbf{x} - \mathbf{z}) \mathbf{1}_E(\mathbf{x} - \theta \mathbf{z}) \\ &= p^{2r} \sum_{\mathbf{m} \pmod{p^r}} \sum_{\theta \in G_r} \widehat{\mathbf{1}_E}(\mathbf{m}) \overline{\widehat{\mathbf{1}_E}(\theta \mathbf{m})} e_{p^r}((\mathbf{m} - \theta \mathbf{m}) \cdot \mathbf{x}) \\ &\ll \frac{|E|^2}{p^r} + \mathcal{E}(\mathbf{x}), \end{split}$$

where

$$\mathcal{E}(\mathbf{x}) := p^{2r} \sum_{v=0}^{r-1} \mathcal{E}_v(\mathbf{x}), \qquad \mathcal{E}_v(\mathbf{x}) = \sum_{\substack{\mathbf{m} \pmod{p^r} \\ v_{\mathbf{m}=v}}} \sum_{\theta \in G_r} \widehat{\mathbf{1}_E}(\mathbf{m}) \overline{\widehat{\mathbf{1}_E}(\theta \mathbf{m})} e_{p^r}((\mathbf{m} - \theta \mathbf{m}) \cdot \mathbf{x}).$$

Here, the first term $|E|^2/p^r$ comes from the summands with $\mathbf{m} \equiv \mathbf{0} \pmod{p^r}$, in view of $|\widehat{\mathbf{1}_E}(\mathbf{0})| = |E|/p^{2r}$ and $|G_r| \ll p^r$.

Write $\mathbf{m} = p^v \tilde{\mathbf{m}}$ and $\tilde{\theta} \equiv \theta \pmod{p^{r-v}}$. Then $\tilde{\theta}\tilde{\mathbf{m}} \equiv \theta \mathbf{m} \pmod{p^{r-v}}$. Noting that $|G_r/G_{r-v}| = p^v$,

we have

$$\begin{aligned} \mathcal{E}_{v}(\mathbf{x}) &= p^{v} \sum_{\substack{\mathbf{\tilde{m}} \pmod{p^{r-v}}\\ v_{\mathbf{\tilde{m}}=0}}} \sum_{\tilde{\theta} \in G_{r-v}} \widehat{1_{E}}(p^{v}\tilde{\mathbf{m}}) \overline{1_{E}(p^{v}\tilde{\theta}\tilde{\mathbf{m}})} e_{p^{r-v}} \left((\tilde{\mathbf{m}} - \tilde{\theta}\tilde{\mathbf{m}}) \cdot \mathbf{x} \right) \right) \\ &= p^{v} \sum_{\tilde{j} \in (\mathbb{Z}/p^{r-v}\mathbb{Z})^{*}} \sum_{\substack{\mathbf{\tilde{m}} \in C_{r-v,\tilde{j}}}} \sum_{\mathbf{m} \in C_{r-v,\tilde{j}}} \widehat{1_{E}}(p^{v}\tilde{\mathbf{m}}) \overline{1_{E}(p^{v}\mathbf{m})} e_{p^{r-v}} \left((\tilde{\mathbf{m}} - \mathbf{m}) \cdot \mathbf{x} \right) \right) \\ &= p^{v} \sum_{\tilde{j} \in (\mathbb{Z}/p^{r-v}\mathbb{Z})^{*}} \left| \sum_{\substack{\tilde{m} \in C_{r-v,\tilde{j}}}} \widehat{1_{E}}(p^{v}\tilde{\mathbf{m}}) e_{p^{r-v}} (\tilde{\mathbf{m}} \cdot \mathbf{x}) \right|^{2}. \end{aligned}$$

Rewrite

$$\begin{split} \widehat{\mathbf{1}_E}(-p^v \tilde{\mathbf{m}}) &= \frac{1}{p^{2r}} \sum_{\mathbf{y} \pmod{p^r}} \mathbf{1}_E(\mathbf{y}) e_{p^r}(-p^v \tilde{\mathbf{m}} \cdot \mathbf{y}) \\ &= \frac{1}{p^{2r}} \sum_{\mathbf{y}_1 \pmod{p^{r-v}}} \sum_{\mathbf{y}_2 \pmod{p^v}} \mathbf{1}_E(\mathbf{y}_1 + p^{r-v} \mathbf{y}_2) e_{p^{r-v}}(-\tilde{\mathbf{m}} \cdot \mathbf{y}_1) := \widehat{g_{E,r-v}}(\tilde{\mathbf{m}}). \end{split}$$

Here, the Fourier transformation on the left-hand side is over $(\mathbb{Z}/p^r\mathbb{Z})^2$, while that on the righthand side is over $(\mathbb{Z}/p^{r-v}\mathbb{Z})^2$. And, for $\mathbf{y}_1 \pmod{p^{r-v}}$,

$$g_{E,r-v}(\mathbf{y}_1) = \frac{1}{p^{2v}} \sum_{\mathbf{y}_2 \pmod{p^v}} \mathbb{1}_E(\mathbf{y}_1 + p^{r-v}\mathbf{y}_2) = \frac{1}{p^{2v}} \#\{\mathbf{y} \in E : \mathbf{y} \equiv \mathbf{y}_1 \pmod{p^{r-v}}\}.$$

We note that for all $\mathbf{y}_1 \pmod{p^{r-v}}$, one has $g_{E,r-v}(\mathbf{y}_1) \leq 1$. Write $f = \widehat{g_{E,r-v}}|_{C}$. It follows that

write
$$J = g_{E,r-v}|_{C_{r-v,\tilde{j}}}$$
. It follows that

$$\begin{aligned} \mathcal{E}_{v}(\mathbf{x}) &= p^{v} \sum_{\tilde{j} \in (\mathbb{Z}/p^{r-v}\mathbb{Z})^{*}} \left| \sum_{\tilde{m} \in C_{r-v,\tilde{j}}} \widehat{g_{E,r-v}}(\tilde{\mathbf{m}}) e_{p^{r-v}}(\tilde{\mathbf{m}} \cdot \mathbf{x}) \right|^{2} \\ &\ll p^{2r-v} \sum_{\tilde{j} \in (\mathbb{Z}/p^{r-v}\mathbb{Z})^{*}} \left| (fd\sigma_{\tilde{j}})^{\vee}(\mathbf{x}) \right|^{2}. \end{aligned}$$

Moreover, we have $\mathcal{E}_v(\mathbf{x}) = \mathcal{E}_v(\tilde{\mathbf{x}})$ if $\tilde{\mathbf{x}} \equiv \mathbf{x} \pmod{p^{r-v}}$. Now we write

$$E_{[r-v]} := \{ \tilde{\mathbf{x}} \pmod{p^{r-v}} : \tilde{\mathbf{x}} \equiv \mathbf{x} \pmod{p^{r-v}} \text{ for some } \mathbf{x} \in E \}.$$

Then, by Cauchy-Schwartz inequality and Theorem 4.1, one deduces that

$$p^{2r} \sum_{\mathbf{x} \in E} \mathcal{E}_{v}(\mathbf{x}) \ll p^{4r-v} \sum_{\tilde{\mathbf{x}} \in A_{[r-v]}} p^{2v} g_{E,r-v}(\tilde{\mathbf{x}}) \sum_{\tilde{j} \in (\mathbb{Z}/p^{r-v}\mathbb{Z})^{*}} |(fd\sigma_{\tilde{j}})^{\vee}(\tilde{\mathbf{x}})|^{2} \\ \ll p^{4r+v} \sum_{\tilde{j} \in (\mathbb{Z}/p^{r-v}\mathbb{Z})^{*}} \left(\sum_{\tilde{\mathbf{x}} \in E_{[r-v]}} |g_{E,r-v}(\tilde{x})|^{2} \right)^{1/2} \left(\sum_{\tilde{\mathbf{x}} \in E_{[r-v]}} |(fd\sigma_{\tilde{j}})^{\vee}(\tilde{\mathbf{x}})|^{4} \right)^{1/2} \\ \ll p^{4r+v} \left(\sum_{\tilde{\mathbf{x}} \in E_{[r-v]}} |g_{E,r-v}(\tilde{\mathbf{x}})|^{2} \right)^{1/2} \cdot p^{-\frac{r-v+1}{2}} \sum_{\tilde{j} \in (\mathbb{Z}/p^{r-v}\mathbb{Z})^{*}} \sum_{\tilde{\mathbf{y}} \in C_{r-v,\tilde{j}}} |\widehat{g_{E,r-v}}(\tilde{\mathbf{y}})|^{2} \\ \ll p^{\frac{3r+7v-1}{2}} \left(\sum_{\tilde{\mathbf{y}} \pmod{p^{r-v}}} |g_{E,r-v}(\tilde{\mathbf{y}})|^{2} \right)^{3/2}.$$

The last sum can be bounded trivially from above

$$\sum_{\tilde{\mathbf{y}} \pmod{p^{r-v}}} |g_{E,r-v}(\tilde{\mathbf{y}})|^2 \leq \sum_{\tilde{\mathbf{y}} \pmod{p^{r-v}}} |g_{E,r-v}(\tilde{\mathbf{y}})| \leq \frac{|E|}{p^{2v}}.$$

In other words,

$$p^{2r} \sum_{\mathbf{x} \in E} \mathcal{E}_v(\mathbf{x}) \ll p^{\frac{3r+v-1}{2}} |E|^{3/2}.$$

With this bound in hand, one has

$$\mathcal{N} \ll \frac{|E|^3}{p^r} + p^{\frac{3r-1}{2}} |E|^{3/2} \sum_{v=0}^{r-1} p^{v/2} \ll \frac{|E|^3}{p^r} + p^{2r-1} |E|^{3/2} \ll \frac{|E|^3}{p^r},$$

whenever $|E| \gg p^{2r-\frac{2}{3}}$. This proves Theorem 1.8.

Indeed,

$$p^{2r} \sum_{\mathbf{x}\in E} \sum_{r=0}^{r-2} \mathcal{E}_v(\mathbf{x}) \ll p^{\frac{3r-1}{2}} |E|^{3/2} \sum_{v=0}^{r-2} p^{v/2} \ll p^{2r-\frac{3}{2}} |E|^{3/2} \ll \frac{|E|^3}{p^r}$$

whenever $|E| \gg p^{2r-1}$. In the statement of Theorem 1.9, we know that

$$\#\{(\mathbf{y}_1, \mathbf{y}_2) \in E^2 : \mathbf{y}_1 \equiv \mathbf{y}_2 \pmod{p} \ll p^{2r - \frac{7}{3}} |E|.$$

Then

$$\sum_{\tilde{\mathbf{y}} \pmod{p}} |g_{E,1}(\tilde{\mathbf{y}})|^2 = \frac{1}{p^{4r-4}} \#\{(\mathbf{y}_1, \mathbf{y}_2) \in E^2 : \mathbf{y}_1 \equiv \mathbf{y}_2 \,(\text{mod } p)\} \ll p^{-2r+\frac{5}{3}} |E|.$$

It follows that

$$p^{2r} \sum_{x \in E} \mathcal{E}_{r-1}(\mathbf{x}) \ll p^{5r-4} \cdot \left(p^{-2r+\frac{5}{3}} |E| \right)^{3/2} \ll p^{2r-\frac{3}{2}} |E|^{\frac{3}{2}} \ll \frac{|E|^3}{p^r},$$

and then $\mathcal{N} \ll \frac{|E|^3}{p^r}$. Plugging this bound to the above argument gives us Theorem 1.9.

6 Proof of Theorems 1.11, 1.12, 1.13, and 1.14

In this section, we write $q = p^r$ for simplicity. To prove Theorems 1.11 and 1.13, we use a general framework developed for pseudo-random graphs in which Lemma 2.5 plays a crucial role.

Proof of Theorem 1.11: Let G = (V, E) be a graph with the vertex set V and the edge set E. Let M be its adjacency matrix, i.e., $M_{ij} = 1$ if and only if there is an edge between i and j and zero otherwise. Let $\lambda_1 \geq \lambda_2 \geq \ldots \geq \lambda_{|V|}$ be its eigenvalues. Assume that |V| = N, the graph G is called a (N, d, λ) -graph if each vertex is of degree d and the second eigenvalue, determined by $\max{\lambda_2, -\lambda_N}$, is at most λ .

Let G and H be two graphs. The Cartesian product of G and H, denoted by $G\Box H$, is defined as follows. The vertex set $V(G\Box H) = V(G) \times V(H)$, and there is an edge between (u_1, v_1) and (u_2, v_2) if and only if either $u_1 = u_2$ and $(v_1, v_2) \in E(H)$ or $v_1 = v_2$ and $(u_1, u_2) \in V(G)$.

We recall the following result on the number of rectangles in a Cartesian product of graphs in [21].

Theorem 6.1. Let G_i be (N_i, d_i, λ_i) -graphs with $1 \leq i \leq 2$. Set $G = G_1 \square G_2$. For any $0 < \delta' < \delta < 1$, there exists $\epsilon > 0$ such that for any $A \subset V(G_1 \square G_2)$ with $|A| \geq \delta |V(G_1 \square G_2)|$, if $\max \left\{\frac{\lambda_1}{d_1}, \frac{\lambda_2}{d_2}\right\} < \epsilon$, then

$$N = \sum_{(u_1, u_2) \in E(G_1), (v_1, v_2) \in E(G_2)} A(u_1, v_1) A(u_1, v_2) A(u_2, v_1) A(u_2, v_2) > \delta'^4 N_1 N_2 d_1 d_2 A(u_1, v_2) A(u_2, v_2) = \delta'^4 N_1 N_2 d_1 d_2 A(u_1, v_2) A(u_2, v_2) A(u_2, v_2) = \delta'^4 N_1 N_2 d_1 d_2 A(u_1, v_2) A(u_2, v_2) A(u_2, v_2) = \delta'^4 N_1 N_2 d_1 d_2 A(u_1, v_2) A(u_2, v_2) A(u_2, v_2) = \delta'^4 N_1 N_2 d_1 d_2 A(u_1, v_2) A(u_2, v_2) A(u_2, v_2) = \delta'^4 N_1 N_2 d_1 d_2 A(u_1, v_2) A(u_2, v_$$

Given $j \in (\mathbb{Z}/p^r\mathbb{Z})^*$. To prove Theorem 1.11, we will apply the above theorem with appropriate graphs G and H. More precisely, define G = H with $V(G) = V(H) = (\mathbb{Z}/p^r\mathbb{Z})^n$, and there is an edge between two vertices \mathbf{x} and \mathbf{y} if and only if $F(\mathbf{x} - \mathbf{y}) \equiv j \pmod{q}$. So G and H have q^n vertices and are regular of degree $(1 + o(1))q^{n-1}$. On the other hand, since G and H are Cayley graphs with the generating set $S = \{\mathbf{x} \in (\mathbb{Z}/p^r\mathbb{Z})^n \colon F(\mathbf{x}) \equiv j \pmod{q}\}$. Thus, all non-trivial eigenvalues of G and H are bounded by $q^n \cdot \max_{\mathbf{m} \not\equiv \mathbf{0} \pmod{q}} |\widehat{1_S}(\mathbf{m})| \ll q^{n-1}p^{-\frac{n-1}{2}}$ by Lemma 2.5. In other words, G and H are $(q^n, (1 + o(1))q^{n-1}, q^{n-1}p^{-\frac{n-1}{2}})$ -graphs.

Thus, Theorem 1.11 follows directly from Theorem 6.1 with A = E.

Proof of Theorem 1.13: To prove Theorem 1.13, we follow the same approach as in the proof of Theorem 1.11. More precisely, we make use of the following result on (N, d, λ) -graphs, which is also taken from [21].

Theorem 6.2. Let G be a (N, d, λ) -graph, $\ell \geq 1$ an integer, and A be a vertex set with $|A| \gg_{\ell} \lambda \cdot \frac{N}{d}$. Let $P_{\ell}(U)$ denotes the number of paths of length ℓ in A. Then we have

$$P_{\ell}(E) \sim \frac{|A|^{\ell+1} d^{\ell}}{N^{\ell}}.$$

With the graph G defined as above, Theorem 1.13 follows from Theorem 6.2 immediately.

Proof of Theorem 1.12: We first recall the statement of Theorem 1.4: for $E \subset (\mathbb{Z}/p^r\mathbb{Z})^n$ if $|E| \gg q^n p^{-\frac{n-1}{2}}$, then for any $j \in (\mathbb{Z}/p^r\mathbb{Z})^*$, the number of pairs $(\mathbf{x}, \mathbf{y}) \in E \times E$ such that $F(\mathbf{x} - \mathbf{y}) \equiv j \pmod{q}$ is at least $\gg q^{-1} |E|^2$.

By the pigeon-hole principle, there exists $\mathbf{u} \in (\mathbb{Z}/p^r\mathbb{Z})^n$ such that $F(\mathbf{u}) \equiv j \pmod{q}$ and the number of pairs $(\mathbf{x}, \mathbf{y}) \in E \times E$ such that $\mathbf{x} = \mathbf{y} + \mathbf{u}$ is at least $\gg \frac{|E|^2}{q^n}$. Let $E_1 \subset E$ be the set of $\mathbf{y} \in E$ such that $\mathbf{y} + \mathbf{u} \in E$. Without loss of generality, we assume that $|E_1| = |E|^2/q^n$. By Theorem 1.4 again, the number of pairs $(\mathbf{y}_1, \mathbf{y}_2) \in E_1 \times E_2$ such that $F(\mathbf{y}_1 - \mathbf{y}_2) \equiv j \pmod{q}$ is at least $|E_1|^2/q = |E|^4/q^{2n+1}$. Among these pairs, the number of pairs with $\mathbf{y}_1 = \mathbf{y}_2 + \mathbf{u}$ is at most $|E_1|$, which is smaller than $|E_1|^2/q$. Therefore, by the pigeon-hole principle again, there exists $\mathbf{u}' \in (\mathbb{Z}/p^r\mathbb{Z})^n$ such that $\mathbf{u} \neq \mathbf{u}'$, $F(\mathbf{u}') \equiv j \pmod{q}$, and the number of pairs $(\mathbf{y}_1, \mathbf{y}_2) \in E_1 \times E_1$ such that $\mathbf{y}_1 = \mathbf{y}_2 + \mathbf{u}'$ is at least $\gg \frac{|E|^2}{q^n} > 0$.

With each pair $(\mathbf{y}_1, \mathbf{y}_2)$ satisfying the above property, we have a cycle of the from $(\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_2 + \mathbf{u}, \mathbf{y}_1 + \mathbf{u})$. This completes the proof.

Proof of Theorem 1.14: To prove Theorem 1.14, we need a stronger version of Theorem 1.4, namely, the pinned distance version.

Theorem 6.3. Let $E_1, E_2 \subset (\mathbb{Z}/p^r\mathbb{Z})^n$ with $|E_1|, |E_2| \gg q^n p^{-\frac{n-1}{2}}$, then there exists $E'_1 \subset E_1$ with $|E'_1| \gg |E_1|$ and $|\Delta_{n.r,\mathbf{x}}(E_2)| \gg q$ for all $\mathbf{x} \in E'_1$. Here

$$\Delta_{n.r,\mathbf{x}}(E_2) := \{ F(\mathbf{x} - \mathbf{y}) \colon \mathbf{y} \in E_2 \}.$$

To prove this strong version, we introduce an incidence problem between points and F-spheres in $(\mathbb{Z}/p^r\mathbb{Z})^n$.

A F-sphere centered at x of radius j is the set

$$\{y \in (\mathbb{Z}/p^r\mathbb{Z})^n \colon F(x-y) \equiv j \mod q\}.$$

Let \mathcal{P} be a set of points in $(\mathbb{Z}/p^r\mathbb{Z})^n$ and \mathcal{S} be a set of *F*-spheres. The number of incidences between \mathcal{P} and \mathcal{S} , denoted by $I(\mathcal{P}, \mathcal{S})$, is defined by

$$I(\mathcal{P},\mathcal{S}) = \#\{(p,s) \in \mathcal{P} \times \mathcal{S} \colon p \in s\}$$

Theorem 6.4. Let \mathcal{P} be a set of points in $(\mathbb{Z}/p^r\mathbb{Z})^n$ and \mathcal{S} be a set of F-spheres with radii in $(\mathbb{Z}/p^r\mathbb{Z})^*$. Then the number of incidences between \mathcal{P} and \mathcal{S} satisfies

$$I(\mathcal{P}, \mathcal{S}) \le \frac{|\mathcal{P}||\mathcal{S}|}{q} + Cq^{n-\frac{1}{2}}p^{-\frac{n-1}{2}}|\mathcal{P}|^{1/2}|cS|^{1/2}.$$

Proof. We partition S into S_i , $i \in (\mathbb{Z}/p^r\mathbb{Z})^*$, such that spheres in S_i are of radius i. By abuse of notation, we denote the set of centers in S_i by S_i . By Theorem 1.4, we have

$$I(\mathcal{P}, \mathcal{S}_i) \le \frac{|\mathcal{P}||\mathcal{S}_i|}{q} + q^{n-1}p^{-\frac{n-1}{2}}|\mathcal{P}|^{1/2}|\mathcal{S}_i|^{1/2}.$$

Taking the sum over all i and by using the Cauchy-Schwarz inequality, we have the conclusion. \Box

Proof of Theorem 6.3. For $\mathbf{x} \in E_1$, let $\Delta_{n,r,\mathbf{x}}^*(E_2)$ be the set of distances in $(\mathbb{Z}/p^r\mathbb{Z})^*$ between \mathbf{x} and points in E_2 . Let $\mathcal{S}(\mathbf{x})$ be the set of spheres centered at \mathbf{x} of radii in $\Delta_{n,r,\mathbf{x}}^*(E_2)$, and set $\mathcal{S} = \bigcup_{\mathbf{x} \in E_1} \mathcal{S}(\mathbf{x})$. We have $|\mathcal{S}| = \sum_{\mathbf{x} \in E_1} |\Delta_{n,r,\mathbf{x}}^*(E_2)|$. Under the conditions on the sizes of E_1 and E_2 , for any $j \in (\mathbb{Z}/p^r\mathbb{Z})^*$, it follows from Theorem 1.4 that the number of pairs $(\mathbf{x}, \mathbf{y}) \in E_1 \times E_2$ such that $F(\mathbf{x} - \mathbf{y}) \equiv j \pmod{q}$ is at least $|E_1||E_2|/2q$.

Thus, $I(E_2, S) \ge |E_1||E_2|/4$. Applying Theorem 6.4 gives

$$I(E_2, \mathcal{S}) \le \frac{|E_2| \sum_{\mathbf{x} \in E_1} |\Delta_{\mathbf{x}}(E_2)^*|}{q} + Cq^{n-\frac{1}{2}} p^{-\frac{n-1}{2}} |E_2|^{1/2} \left(\sum_{\mathbf{x} \in E_1} |\Delta_{n.r, \mathbf{x}}(E_2)^*| \right)^{1/2}$$

If $\sum_{\mathbf{x}\in E_1} |\Delta_{n.r,\mathbf{x}}(E_2)^*| \le |E_1|q/8$, then $|E_1||E_2| \le 8C^2q^{2n}p^{-(n-1)}$, so we reach a contradiction as long as $|E_1||E_2| > 8C^2q^{2n}p^{-(n-1)}$.

Thus, if $|E_1||E_2| > 8C^2q^{2n}p^{-(n-1)}$, then we have

$$\sum_{\mathbf{x}\in E_1} |\Delta_{n.r,\mathbf{x}}(E_2)^*| > \frac{|E_1|q}{8}.$$
(17)

Set $E'_1 := \{ \mathbf{x} \in E_1 : |\Delta_{\mathbf{x}}(E_2)^*| \ge q/32 \}$. We are going to show that $|E'_1| \ge |E_1|/32$. Indeed, otherwise, we have

$$\sum_{\mathbf{x}\in E_1} |\Delta_{\mathbf{x}}(E_2)^*| = \sum_{\mathbf{x}\in E_1'} + \sum_{\mathbf{x}\in E_1\setminus E_1'} \le q|E_1'| + \frac{q|E_1|}{32} \le \frac{q|E_1|}{16},$$

which contradicts (17). Therefore, we have proved that there exists $E'_1 \subset E_1$ with $|E'_1| \ge |E_1|/32$ and for all $\mathbf{x} \in E'_1$, we have $|\Delta_{n.r,\mathbf{x}}(E_2)^*| \ge q/32$. The proof is complete.

With Theorem 6.3 in hand, Theorem 1.14 now follows from a combinatorial argument which is identical to [22, Proof of Theorem 3.1] with Theorem 6.3 in place of [22, Lemma 2.2]. So we omit the details.

7 Sharpness examples

In this section, we provide some sharpness examples for results stated in the introduction.

Example 7.1. Let p be a large prime. Suppose that n is odd and -1 is a k-th power modulo p, i.e., $\xi^k \equiv -1 \pmod{p}$. We consider the polynomial $F(\mathbf{x}) = x_1^k + x_2^k + \ldots + x_n^k$.

For r = 1, let us take

$$\Pi_0 = \{ (u, \xi u) : u \in \mathbb{Z}/p\mathbb{Z} \}.$$

Then, for any $(u, \xi u) \in \Pi_0$, one has $||(u, \xi u)|| = u^k + (\xi u)^k \equiv 0 \pmod{p}$. This is a one-dimensional subspace of $(\mathbb{Z}/p\mathbb{Z})^2$. We have $||(u, \xi u) - (u', \xi u')|| = 0$ for any $(u, \xi u)$, $(u', \xi u') \in \Pi_0$. Let $1 \leq l < p/2$ be a parameter to be determined later. Take

$$E_1 = \{ (x_1, x_2, \dots, x_n) \in \mathbb{Z} / p\mathbb{Z} : x_1 \in \{1, 2, \dots, l\}, (x_2, x_3), \dots, (x_{n-1}, x_n) \in \Pi_0 \}.$$

Then $|E_1| = lp^{\frac{n-1}{2}}$, and

$$|\Delta_{n,1}(E_1)| = \#\left\{(-l+1)^k, (-l+2)^k, \dots, (l-1)^k\right\} < 2l.$$

For $r \geq 2$, let us take

$$E_r = \left\{ \mathbf{x} + p\mathbf{y} : \, \mathbf{x} \in E_1, \, \mathbf{y} \in (\mathbb{Z}/p^{r-1}\mathbb{Z})^n \right\}$$

as a subset of $(\mathbb{Z}/p^r\mathbb{Z})^n$. It is not hard to see that $|E_r| = lp^{nr - \frac{n+1}{2}}$. Note that, for any two elements $\mathbf{z} = \mathbf{x} + p\mathbf{y}, \mathbf{z}' = \mathbf{x}' + p\mathbf{y}' \in E_r$, we have $\|\mathbf{z} - \mathbf{z}'\| \equiv \|\mathbf{x} - \mathbf{x}'\| \pmod{p}$. So

$$|\Delta_{n,r}(E_r)| \le p^{r-1} \cdot |\Delta_{n,1}(E_1)| < 2lp^{r-1}.$$

To sum up, if we take $l = \lceil p \cdot w(p) \rceil$ with w(p) = o(1) $(p \to \infty)$ being a positive-valued function tending to 0 arbitrarily slowly, then there are sets E_r with density $p^{-\frac{n-1}{2}}w(p)$ such that $|\Delta_{n,r}(E_r)| < 2p^r w(p) = o(p^r)$. So Corollary 1.5 is optimal in general.

Example 7.2. Given a positive integer C, we first claim that there exists a set $E_1 \subset (\mathbb{Z}/p\mathbb{Z})^n$ of size $p^{n/2}$ such that $\Delta_{n,1}(E_1) = \{0\}$ or $|\Delta_{n,1}(E_1)| \leq (p+1)/C$.

If -1 is a square or $n \equiv 0 \pmod{4}$, as above, we can find such a set $E_1 \subset (\mathbb{Z}/p\mathbb{Z})^n$ of size $p^{n/2}$ such that $\Delta_{n,1}(E_1) = \{0\}$.

If -1 is a non-square and $n \equiv 2 \pmod{4}$, then let θ be a generator of the group $G_1 = SO_2(\mathbb{F}_p)$ of rotations, which is cyclic of order p+1, in $(\mathbb{Z}/p\mathbb{Z})^2$, and let $v_0 \in (\mathbb{Z}/p\mathbb{Z})^2$ such that $||v_0|| = 1$. By the Dirichlet's theorem on arithmetic progressions, one can choose p large enough such that C divides p+1. Let G be the subgroup spanned by θ^C . Then we have |G| = (p+1)/C. Set $X = \operatorname{orb}_1(\mathbf{v}_0)$. A direct computation shows that X satisfies the desired property, namely, $|\Delta_{2,1}(X)| \leq |X| = (p+1)/C$. Let $X' \subset (\mathbb{Z}/p^r\mathbb{Z})^{n-2}$ of size $p^{(n-2)/2}$ such that $\Delta_{n,1}(X') = \{0\}$. Set $E_1 = X \times X'$. Then we have $|E_1| = p^{\frac{n-2}{2}} \cdot \frac{p+1}{C}$. By the definition of E_1 , we obtain $|\Delta_{n,1}(E_1)| \leq (p+1)/C$.

For $r \geq 2$, we set

$$E_r = \{ \mathbf{x} + ((\mathbb{Z}/p^r \mathbb{Z}) \setminus U_r)^n \colon \mathbf{x} \in E_1 \}.$$

So, $|E_r| = p^{rn} p^{-\frac{n}{2}}$ and $|\Delta_{n,r}(E_r)| \le p^r/C$.

8 Acknowledgements

Thang Pham would like to thank to the VIASM for the hospitality and for the excellent working condition. Thang Pham was supported by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 101.99–2021.09.

We thank Le Quang Hung for checking all proofs in Section 4 for the case $p \equiv 1 \pmod{4}$.

References

[1] D. Covert, A. Iosevich, and J. Pakianathan, Geometric configurations in the ring of integers modulo p^{ℓ} , Indiana university mathematics journal, **61**(5) (2012), 1949-1969.

- [2] J. Chapman, M. B. Erdogan, D. Hart, A. Iosevich, and D. Koh, *Pinned distance sets*, ksimplices, Wolff's exponent in finite fields and sum-product estimates, Mathematische Zeitschrift, 271(1-2) (2012), 63-93.
- [3] X. Du, A. Iosevich, Y. Ou, H. Wang, and R. Zhang, An improved result for Falconer's distance set problem in even dimensions, Mathematische Annalen, 380 (2021), 1215–1231.
- [4] D. Hart, A. Iosevich, D. Koh, and M. Rudnev, Averages over hyperplanes, sum-product theory in vector spaces over finite fields and the Erdős-Falconer distance conjecture, Transactions of the American Mathematical Society, 363(6) (2007), 3255-3275.
- [5] B. Hanson, B. Lund, and O. Roche-Newton, On distinct perpendicular bisectors and pinned distances in finite fields, Finite Fields and Their Applications, 37 (2016), 240–264.
- [6] A. Iosevich and D. Koh, The Erdős-Falconer distance problem, exponential sums, and Fourier analytic approach to incidence theorems in vector spaces over finite fields, SIAM Journal on Discrete Mathematics, 23(1) (2009), 123-135.
- [7] A. Iosevich and M. Rudnev, Erdős-Falconer distance problem in vector spaces over finite fields, Transactions of the American Mathematical Society, 359(12) (2007), 6127-6142.
- [8] A. Iosevich and D. Koh, Extension theorems for paraboloids in the finite field setting, Mathematische Zeitschrift, 266(2) (2010), 471–487.
- [9] A. Iosevich and D. Koh, Extension theorems for spheres in the finite field setting, Forum Mathematicum, 22(3) (2010), 457–483.
- [10] A. Iosevich, D. Koh, and M. Lewko, Finite field restriction estimates for the paraboloid in high even dimensions, Journal of Functional Analysis, 278(11) (2020), 108450.
- [11] D. Koh and C-Y. Shen, The generalized Erdős-Falconer distance problems in vector spaces over finite fields, Journal of Number Theory, 132(11) (2012), 2455-2473.
- [12] M. Lewko, Finite field restriction estimates based on Kakeya maximal operator estimates, Journal of European Mathematics Society, 21(12) (2019), 3649–3707.
- [13] M. Lewko, New restriction estimates for the 3-d paraboloid over finite fields, Advances in Mathematics, 270(1) (2015), 457–479.
- [14] G. Mockenhaupt and T. Tao, Restriction and Kakeya phenomena for finite fields, Duke Mathematical Journal, 121(1) (2004), 35–74.
- [15] B. Lichtin, Distance and sum-product problems over finite p-adic rings, Proceedings of the London Mathematical Society, 118(6) (2019), 1450-1470.
- [16] B. Lichtin, k-chain configurations of points over p-adic rings, Proceedings of the American Mathematical Society, 151(10) (2023), 4113-4125.
- [17] N. Lyall and A. Magyar, Weak hypergraph regularity and applications to geometric Ramsey theory, Transactions of the American Mathematical Society, Series B, 9(5) (2022), 160-207.
- [18] S-Y. Liao, Exploration on Incidence Geometry and Sum-Product Phenomena, arXiv:2310.07964 (2023).

- [19] B. Murphy, M. Rudnev, and S. Stevens, Bisector energy and pinned distances in positive characteristic, arXiv:1908.04618 (version 1) (2019).
- [20] B. Murphy, G. Petridis, T. Pham, M. Rudnev, and S. Stevens, On the pinned distances problem in positive characteristic, Journal of the London Mathematical Society, 105(1) (2022), 469-499.
- [21] T. Pham, S. Senger, M. Tait, and V.T.H. Thu, Geometric structures in pseudo-random graphs, arXiv:2208.04399 (2022).
- [22] T. Pham, S. Senger, and T.D. Tran, Distribution of pinned distance trees in the plane \mathbb{F}_p^2 , Discrete Mathematics, **346**(12), p.113613.