The obstacle problem for linear scalar conservation laws with constant velocity

Paulo Amorim^{4,5}, Alexander Keimer¹, Lukas Pflug^{2,3}, Jakob Rodestock²

¹Department Mathematics, Friedrich-Alexander Universität Erlangen-Nürnberg (FAU), Cauerstr. 11, Erlangen, 91058, Germany. ²Competence Center for Scientific Computing (FAU CSC), Friedrich-Alexander Universität Erlangen-Nürnberg (FAU), Martensstr.

5a, Erlangen, 91058, Germany.

³Chair of Applied Mathematics, Continuous Optimization, Department Mathematics, Friedrich-Alexander Universität Erlangen-Nürnberg (FAU), Cauerstrasse 11, Erlangen, 91058, Germany.

⁴School of Applied Mathematics (FGV-EMAp), Fundação Getúlio Vargas, Praia de Botafogo 190, Rio de Janeiro, 22250-900, Brazil. ⁵Instituto de Matemática, Universidade Federal do Rio de Janeiro (UFRJ), Av. Athos da Silveira Ramos 149, Rio de Janeiro, C.P. 68530,

21941-909, Brazil.

Contributing authors: paulo@im.ufrj.br; alexander.keimer@fau.de; lukas.pflug@fau.de; jakob.w.rodestock@fau.de;

Abstract

In this contribution, we present a novel approach for solving the obstacle problem for (linear) conservation laws. Usually, given a conservation law with an initial datum, the solution is uniquely determined. How to incorporate obstacles, i.e., inequality constraints on the solution so that the resulting solution is still "physically reasonable" and obeys the obstacle, is unclear. The proposed approach involves scaling down the velocity of the conservation law when the solution approaches the obstacle. We demonstrate that this leads to a reasonable solution and show that, when scaling down is performed in a discontinuous fashion, we still obtain a suitable velocity - and the solution satisfying a discontinuous conservation law. We illustrate the developed solution concept using numerical approximations.

Keywords: obstacle problem for conservation laws, obstacle problem, viscosity approximation, discontinuous conservation laws, solution constrained PDE

MSC Classification: 35L65, 35L03, 35L81, 35N99

1 Introduction

1.1 The obstacle problem in the literature

Hyperbolic conservation laws involving constraints on the solution (usually called *obstacles*) arise naturally in many applications. For instance, in traffic modeling, it is natural to impose that vehicle density cannot exceed a certain threshold. Bounds on the velocity can be envisioned, given by speed limits, or on the vehicular flux in a certain region [1–11]. Elsewhere, in multi-phase flows modeled by hyperbolic equations, constraints may appear in the solutions, leading to obstacles. For related problems involving hyperbolic settings, [12–19] provide a few examples. In addition, pedestrian flows may involve congestion, so population density should be limited; see [20] for a recent survey.

More generally, in cases where the solution of a partial differential equation represents the density of a medium, e.g., a population, fluid, group of vehicles, or group of pedestrians, one may consider a variety of situations where a constraint on the density is imposed in a certain region of space or time. Therefore, since, in many cases, such dynamics are described by parabolic or hyperbolic evolution equations, a modeling mechanism capable of modifying an existing problem is useful to introduce density constraints in a meaningful way. This is especially interesting in cases where the solutions of the "free" problem (i.e., without constraint) typically violate the constraint; then, additional mechanisms that enforce the obstacle can be defined while remaining consistent with the underlying model.

The existing literature on obstacle problems is already substantial, especially concerning parabolic or elliptic problems. Since this is not the focus of our work, we cite only a few selected contributions, which are either foundational/surveys/monographs [21–27] or publications involving interesting applications [28–30]. Most approaches, especially in the parabolic setting, state the obstacle problem in a variational setting, where connecting the mathematical tools used to proper physical interpretations is not easy. Namely, a typical method to obtain a solution to an obstacle problem is by *penalization*. For instance, when a function u that must remain below some obstacle ϕ is sought, the evolution equation is supplemented by some source terms acting to decrease u on the set where $u > \phi$ in the setting of some approximation scheme. In general, though, such terms may have no ready interpretation in the model setting and often represent mathematical artifacts reminiscent of more abstract problems intended to find appropriate projections on convex sets [21, 25].

For hyperbolic conservation laws, in particular, the penalization approach has been applied by a few authors, e.g., [12, 17, 31]. One drawback is that a penalization term is,



Fig. 1 Solution q_A proposed in [31] with constant velocity of 1 for times t = 0, 0.5, 1, 2 (top from left to right) and the solution suggested in Eq. (1) for times t = 0, 0.5, 1, 2 (bottom from left to right) and $\varepsilon = \frac{1}{1024}$ with identical initial datum q_0 and obstacle o (black). The dashed line shows the function $q_0(\cdot - t)$.

in general, not consistent with the conservation (or evolution) of the total mass. However, a constraint on a point-wise density should not necessarily affect the evolution of the total mass. In view of this, in the setting of an obstacle problem for a conservation law preserving mass, authors in [31] introduced a Lagrange multiplier (effectively a source term) designed to counterbalance the effect of the penalization on the total mass. However, the resulting formulation relies on mass creation and destruction to enforce the obstacle constraint while preserving the total mass. Here, we propose a different approach, which we argue is more natural.

Indeed, when trying to enforce an obstacle-like condition, density variations should originate from rearrangement of individuals rather than the destruction/creation of mass, especially in the case of traffic or pedestrian models. Thus, we expect that changes in the local *velocity* of the vehicles or individuals can be enough to adjust the density to satisfy the constraint. For a comparison of the two approaches, see Fig. 1.

With these remarks in mind, we propose a new formulation for the obstacle problem for a one-dimensional hyperbolic conservation law. In this work, we consider the case of a linear flux only. As shown below, the linear case allows for a cleaner exposition, focusing on the main innovations of our approach.

1.2 Problem statement and outline

The obstacle problems for linear conservation laws with constant coefficients reads

$$\partial_t q + \partial_x q = 0$$
, $q(0, \cdot) \equiv q_0$ such that $q(t, x) \leq o(x) \ \forall (t, x) \in (0, T) \times \mathbb{R}$

for an obstacle $o : \mathbb{R} \to \mathbb{R}$. As evident, the proposed system is over-determined in the sense that the Cauchy problem admits a unique solution without the obstacle. As a result, how to incorporate the obstacle into the solution is not straightforward when the following holds:

• mass is conserved even when hitting the obstacle;

- the solution still satisfies a semi-group property in time;
- the solution behaves physically reasonably in the sense that mass is not instantaneously transported across space.

Having this in mind, one may need to adjust the velocity of the conservation law accordingly. Following this approach, we want the velocity of the conservation law to decrease when the solution approaches the obstacle. Thus, for a smoothed version of the Heaviside function $V_{\varepsilon}, \ \varepsilon \in \mathbb{R}_{>0}$, we consider the following nonlinear conservation law:

$$\partial_t q(t, x) + \partial_x \left(V_{\varepsilon}(o - q)q \right) = 0. \tag{1}$$

Here, the velocity becomes small when q approaches o, and whenever q is not close to the obstacle, the dynamics evolve with the constant speed of (almost) 1 as expected for the conservation law without the obstacle. However, the smoothness of V_{ε} remains problematic, as we indeed want the dynamics to hit the obstacle and only slow down when hitting it. Thus, the singular limit $\varepsilon \to 0$ of the solution and the velocity V_{ε} is of interest as well as what values V_{ε} attains at this limit point, resulting in a discontinuous (in the solution) conservation law (compare, in particular, Rem. 5),

$$\partial_t q(t, x) + \partial_x \left(H(o - q)q \right) = 0, \tag{2}$$

where H denotes the Heaviside function. All the raised points are addressed in this manuscript, starting in Sec. 2 with the assumptions on the conservation law, the obstacle, and more (see Asm. 2). Additionally, as we deal with the viscous approximation of Eq. (1), we look into its well-posedness, i.e., existence and uniqueness of solutions.

In Sec. 3, we then show that the solutions to the viscosity approximation and Eq. (1) satisfy the obstacle constraint strictly as long as the initial datum is "compatible" and the density remains non-negative in the viscosity limit (for non-negative initial datum), uniformly in $\varepsilon \in \mathbb{R}_{>0}$.

As we are interested in the mentioned convergence for $\varepsilon \to 0$, crefsec:Lipschitz presents one-sided Lipschitz (OSL) bounds for the solution and the velocity, uniformly in $\varepsilon \in \mathbb{R}_{>0}$, from which we can conclude certain compactness properties of the solution and the velocity, culminating in the convergence result in Thm. 13. In addition, we show that even in the limit $\varepsilon \to 0$, the tuple $(q_{\varepsilon}, V_{\varepsilon}(o - q_{\varepsilon}))$ satisfies a (discontinuous) conservation law, which is indeed of the form stated in Eq. (2).

Sec. 5 then characterizes the limit solution and velocity in specific, more restrictive setups, illustrating the reasonableness of the approach and providing insights into the limiting velocity. Surprisingly, the velocity does not become zero when the density approaches the obstacle. This phenomenon may seem counter-intuitive at first glance, but assuming that the velocity is zero, no mass is transported when the obstacle is active, i.e., a "full blocking" results once the obstacle is active. We also showcase how a backward shock front originating from the point of contact of the obstacle and the solution propagate.

Next, Sec. 6 continues to motivate the chosen approach for the obstacle problem, this time via optimization. On a formal level, we want to maximize the velocity of the

conservation law at each point in time so the obstacle is not violated when forward propagating in time. This leads to an optimization problem for all time $t \in [0, T]$. We again see how the velocity behaves if the solution touches the obstacle, coinciding with the result in Sec. 5.

In Sec. 7, we conclude the contribution with numerical approximations by means of a tailored Godunov scheme, demonstrating the reasonableness of the approach and showing the approximate solution for some specific cases for a better understanding and intuition.

Preliminaries 2

First, we define a sequence of functions that approximates the Heaviside function smoothly and later plays an important role in approximating a solution to the obstacle problem. These functions should have the following properties:

Assumption 1 (Approximation of Heaviside function). To approximate the Heaviside function, we consider a sequence $V_{\varepsilon}(x) := V\left(\frac{x}{\varepsilon}\right)$ for all $x \in [0,\infty), \varepsilon > 0$. Here, $V \in C^{\infty}([0,\infty))$ fulfills the following:

- $\lim_{x\to\infty} V(x) = 1;$
- V(0) = 0;
- V(0) = 0, $V|_{(0,\infty)} > 0$, $(-1)^{k+1}V^{(k)}|_{[0,\infty)} > 0$ for all $n \in \mathbb{N}$; $V^{(k)} \in L^{\infty}([0,\infty))$ for all $k \in \mathbb{N}$, $\|V\|_{L^{\infty}([0,\infty))} = 1$ and $\sup_{\varepsilon > 0} \|V'_{\varepsilon}\|_{L^{\infty}([\tilde{c},\infty))} < \infty$ and all $\tilde{c} > 0$;
- it holds that

$$-\infty < v_{2,1}^- \coloneqq \inf_{y \in \mathbb{R}_{\geq 0}} \frac{V^{\prime\prime}(y)}{V^{\prime}(y)} \le v_{2,1}^+ \coloneqq \sup_{y \in \mathbb{R}_{\geq 0}} \frac{V^{\prime\prime}(y)}{V^{\prime}(y)} < 0,$$

and

• $\lim_{x \to \pm \infty} x V_{\varepsilon}^{(k)}(x) = 0$ for all $k \in \mathbb{N}$.

From the first point, we see that $\lim_{\varepsilon \searrow 0} V_{\varepsilon} = \chi_{(0,\infty)}$ pointwise. The last point is not explicitly referenced again in this work. However, the assumption is not very restrictive and is used in the proof of Thm. 4.

To demonstrate that the above requirements are reasonable, the following is a simple example of such function V.

Example 1 (Example of approximation). We set $V : \mathbb{R} \to \mathbb{R}$, $V(x) := 1 - \exp(-x)$ for $x \in \mathbb{R}$, where V satisfies all the properties outlined in Asm. 1.

Now, as stated in the introduction, we want to investigate Eq. (1) and its limiting behavior for $\varepsilon \to 0$ and $T \in \mathbb{R}_{>0}$ with initial condition $q(0, \cdot) \equiv q_0$. To proceed, assumptions about the regularity of the data are necessary.

Assumption 2 (Regularity). When working with Eq. (1), we assume the following:

Velocity field: The velocity field V_{ε} adheres to Asm. 1. *Initial datum:* $q_0 \in BV(\mathbb{R}; \mathbb{R}_{>0})$ **Obstacle:** $o \in W^{6,\infty}(\mathbb{R})$ such that

•
$$o' \in W^{6,1}(\mathbb{R}),$$

- essinf o(x) q₀(x) > 0, and
 lim_{x→±∞} o(x) both exist in ℝ.

Here BV $\subseteq L^1$ denotes the space of functions with bounded variation and $|\cdot|_{\text{TV}}$ will denote the total variation (semi-norm).

Existence and uniqueness to Eq. (1) for $\varepsilon \in \mathbb{R}_{>0}$ can be found in the standard literature. We present the notion of solution (i.e., weak entropy solution) and a precise statement about existence and uniqueness.

Definition 1 (Entropy solution). Let q_0 be as in Asm. 2. We say that $q \in$ $C([0,T]; L^1_{loc}(\mathbb{R}))$ is an entropy solution to the Cauchy problem Eq. (1) with $q(0, \cdot) \equiv q_0$ if $q(0, \cdot) = q_0$ in $L^1_{loc}(\mathbb{R})$, and for any $(\kappa, \varphi) \in \mathbb{R} \times C^{\infty}_0((0, T) \times \mathbb{R}; \mathbb{R}_{\geq 0})$, the following holds:

$$\int_0^T \!\!\!\int_{\mathbb{R}} |q(t,x) - \kappa| \partial_t \varphi(t,x) + \operatorname{sgn}(q(t,x) - \kappa) \big(f(x,q(t,x)) - f(x,\kappa) \big) \partial_x \varphi(t,x) \, \mathrm{d}x \, \mathrm{d}t \\ - \int_0^T \!\!\!\!\int_{\mathbb{R}} \operatorname{sgn}(q(t,x) - \kappa) \partial_1 f(x,\kappa) \varphi(t,x) \, \mathrm{d}x \, \mathrm{d}t \ge 0.$$

where $f(x,q) \coloneqq V_{\varepsilon}(o(x)-q)q \ \forall (x,q) \in \{(y,z) \in \mathbb{R}^2 : o(y) \ge z\}.$

Given the definition of an entropy solution, we can state the existence and uniqueness of Eq. (1) as follows:

Theorem 3 (Existence and uniqueness to Eq. (1)). There is a unique entropy solution $q_{\varepsilon} \in C([0,T]; L^{1}_{loc}(\mathbb{R}))$ to Eq. (1) as defined in Defn. 1. Moreover, $0 \leq q_{\varepsilon} \leq o$ a.e. in $(0,T) \times \mathbb{R}.$

Proof. Existence follows from [32, §4]. In addition, [32, p.229, Theorem 3] implies directly that $0 \leq q_{\varepsilon} \leq o$ a.e., as 0 and o are both solutions to Eq. (1). Using these bounds, uniqueness can be obtained owing to the doubling of variables technique [32, §3].

To obtain quantitative estimates with regard to the ε parameter, we approximate Eq. (1) with a viscosity to work with smooth solutions, as detailed in the following **Definition 2** (Viscosity approximation). Let $\nu, \varepsilon \in \mathbb{R}_{>0}$ and let Asm. 2 hold. Moreover, we set $q_{0,\nu} = \varphi_{\nu} * q_0$, where * denotes the convolution [33, 4.13] and φ_{ν} is given by

$$\varphi_{\nu}(x) \coloneqq \frac{1}{\nu} \varphi\left(\frac{x}{\nu}\right), \ x \in \mathbb{R},$$

where φ is the standard mollifier [34, 4.2.1, Notation (ii)]. Then, we call the solution $q_{\nu,\varepsilon}: (0,T) \times \mathbb{R} \to \mathbb{R}$ to the initial value problem

$$\partial_t q(t,x) + \partial_x \big(V_{\varepsilon}(o(x) - q(t,x) \big) = \nu \partial_x^2 \big(q(t,x) - o(x) \big), \quad (t,x) \in (0,T) \times \mathbb{R}$$
(3)
$$q_{\nu,\varepsilon}(0,\cdot) \equiv q_{0,\nu}, \qquad on \mathbb{R}$$
(4)

the viscosity approximation of Eq. (1).

The viscosity approximation approach has been laid out by, e.g., Kružkov [32]. However, it is not canonical to have $\nu o''$ on the right-hand side of a viscosity approximation. In fact, this term ensures the obstacle condition $q_{\nu,\varepsilon} < o$ even for the viscosity approximation.

In the following, we state properties of the viscosity approximation and start with the well-posedness of Eqs. (3) to (4). For existence, we resort to parabolic theory.

Theorem 4 (Existence for the viscous equation Eq. (3)). If q_0 and o satisfy assumption Asm. 2, then there is a unique solution $q_{\nu,\varepsilon} \in H^5((0,T) \times \mathbb{R})$ to the problem Eq. (3)-(4).

Proof. To prove this statement, we follow the steps in [35, Chapter II, Section 2] while extending the arguments to a space-dependent flux (as is our case).

To apply the results in [35], we first extend V_{ε} to a function $\tilde{V}_{\varepsilon} \in C^{\infty}(\mathbb{R})$ such that $\|\tilde{V}_{\varepsilon}\|_{L^{\infty}(\mathbb{R})} = 1$ and $\tilde{V}_{\varepsilon}|_{(-\infty, -\ell]} = 0$ for some $\ell > 0$ and for all $0 < \varepsilon < 1$.

The result is \tilde{V}_{ε} , and after obtaining the maximum principle Lem. 7, which is proved for first the extension \tilde{V}_{ε} as well, we obtain the result for Eq. (3). This is because \tilde{V}_{ε} will then only be evaluated on $[0, \infty)$.

Thanks to classical embedding theorems, we also obtain the following

Remark 1 (Asymptotic behavior of $q_{\nu,\varepsilon}$). For $\nu, \varepsilon \in \mathbb{R}_{>0}$, the solution $q_{\nu,\varepsilon}$ to problem Eq. (3) belonging to $H^5((0,T) \times \mathbb{R})$ implies that $\lim_{x \to \pm \infty} \partial_{\alpha}g(t,x) = 0$ for every $t \in [0,T]$ and every multi-index α up to at least order 4. This can be found, e.g., in Brézis [36, Corollary 8.9]. We use this fact extensively in Sec. 3.

Now, by embedding theorems, we have a *classical* solution $q_{\nu,\varepsilon} \in C^3([0,T] \times \mathbb{R})$ of the viscous approximation. Thus, we can obtain bounds in L^1 as well as TV: **Theorem 5** (Estimates for $q_{\nu,\varepsilon}$). For $\varepsilon, \nu \in \mathbb{R}_{>0}$, the solution $q_{\nu,\varepsilon} \in H^5((0,T) \times \mathbb{R})$ to Eq. (3) satisfies

 L^1 bound: $\|q_{\nu,\varepsilon}\|_{L^{\infty}((0,T);L^1(\mathbb{R}))} \leq (T\nu\|o''\|_{L^1(\mathbb{R})} + \|q_0\|_{L^1(\mathbb{R})})$ Spatial TV-bound: For all $t \in [0,T]$, the following holds:

 $|q_{\nu,\varepsilon}(t,\cdot) - o|_{\mathrm{TV}(\mathbb{R})} \le \left(\|o'\|_{L^1(\mathbb{R})} + |q_0|_{\mathrm{TV}(\mathbb{R})} \right) T \|o''\|_{L^1(\mathbb{R})} e^{T} \|V'_{\varepsilon}\|_{L^{\infty}([0,\infty)]} \|o'\|_{L^{\infty}(\mathbb{R})}.$

TV-bound in time: There exists $\bar{\nu} > 0$ such that for all $t \in [0, T]$,

$$\int_{\mathbb{R}} |\partial_t q_{\nu,\varepsilon}(t,x)| \, \mathrm{d}x \le \sup_{\varepsilon > 0} \|V'_{\varepsilon}\|_{L^{\infty}\left([\hat{K},\infty)\right)} \|q_0\|_{L^{\infty}(\mathbb{R})} \left(\|o'\|_{L^1(\mathbb{R})} + |q_0|_{\mathrm{TV}(\mathbb{R})}\right) \\ + \left(1 + 2e^{-1}\right) |q_0|_{\mathrm{TV}(\mathbb{R})} + \nu \|o''\|_{L^1(\mathbb{R})}$$

for all $0 < \nu < \bar{\nu}$, where $\hat{K} \coloneqq \frac{\operatorname{essinf}_{y \in \mathbb{R}} o(y) - q_0(y)}{2}$

Even more, for monotonically decreasing o, we even obtain uniform TV bounds in $\varepsilon \in \mathbb{R}_{>0}$:

Remark 2 (Uniform TV bounds for monotone obstacles). Let o be decreasing and $q_{\nu,\varepsilon}$ be the solution as in Eq. (3). Then, in $\varepsilon, \nu \in \mathbb{R}_{>0}$ and $t \in (0,T)$ the following

holds uniformly:

$$\int_{\mathbb{R}} |\partial_x q_{\nu,\varepsilon}(t,x) - o'(x)| \, \mathrm{d}x \le \left(\|o'\|_{L^1(\mathbb{R})} + |q_0|_{\mathrm{TV}(\mathbb{R})} \right) T \|o''\|_{L^1(\mathbb{R})}.$$

The case of increasing obstacles is trivial as the obstacle never becomes active.

Now, we have collected the necessary tools to prove that $\lim_{\nu \to} q_{\nu,\varepsilon}$ is indeed an entropy solution to Eq. (1):

Theorem 6 $(q_{\nu,\varepsilon} \text{ converges toward the entropy solution for } \nu \to 0)$. Let $\varepsilon, \nu \in \mathbb{R}_{>0}$ and $q_{\varepsilon,\nu}$ the viscosity solution in Defn. 2. Then, there exists $q_{\varepsilon} \in C([0,T]; L^{1}_{loc}(\mathbb{R}))$ so that

$$\lim_{\nu \to 0} \|q_{\varepsilon,\nu} - q_{\varepsilon}\|_{C([0,T];L^1_{loc}(\mathbb{R}))} = 0$$

and q_{ε} is the unique entropy solution to Eq. (1) in the sense of Defn. 1.

Proof. The convergence of $q_{\nu,\varepsilon}$ for $\nu \to 0$ on subsequences follows from Thm. 5 and the Riesz–Kolmogorov-type Theorem for Bochner spaces [37, Theorem 3]. Showing that q_{ε} is an entropy solution and by the uniqueness of entropy solution in Thm. 3, we indeed obtain convergence on all subsequences. This a standard argument that can be found in [32, p.236]. For this, $||q_{\nu,\varepsilon}||_{L^{\infty}([0,T]\times\mathbb{R})}$ must be bounded uniformly in $0 < \nu < 1$. This is proven in Sec. 3.

3 Comparison principles

Thm. 3 shows that q_{ε} respects the obstacle. However, we can even prove that the obstacle is never hit. This prevents the "full blocking", as mentioned in the introduction.

Lemma 7 (Maximum principle/Obstacle is respected). Let $q_{\nu,\varepsilon}$ be a solution to Eq. (3) for $\nu, \varepsilon > 0$. Then,

$$q_{\nu,\varepsilon}(t,x) < o(x), \qquad \forall (t,x) \in [0,T] \times \mathbb{R}.$$
 (5)

Proof. First, extend V_{ε} to $\tilde{V}_{\varepsilon} \in C^{\infty}(\mathbb{R})$ as discussed in the proof of Thm. 4. Let $q_{\nu,\varepsilon}$ be the solution to

$$\partial_t q(t,x) + \partial_x \left(\tilde{V}_{\varepsilon}(o(x) - q(t,x)) = \nu \partial_x^2 \left(q(t,x) - o(x) \right), \quad (t,x) \in (0,T) \times \mathbb{R} \quad (6)$$

$$q_{\nu,\varepsilon}(0,\cdot) \equiv q_{0,\nu}, \qquad \text{on } \mathbb{R}. \quad (7)$$

Not only will we obtain the maximum principle for $q_{\nu,\varepsilon}$, but the maximum principle also implies that $q_{\nu,\varepsilon}$ solves Eqs. (3) and (4):

Fix $t \in (0,T]$. In the end, we want to use [38, Theorem 1]. Here, $(0,T) \mapsto \sup_{y \in \mathbb{R}} q_{\nu,\varepsilon}(t,y) - o(y)$ must be differentiable a.e. Fortunately, this is guaranteed by Lipschitz continuity of all involved functions. Moreover, $\lim_{y\to\pm\infty} q_{\nu,\varepsilon}(t,y) - o(y)$ has to exist, and $\lim_{y\to\pm\infty} \partial_t (q_{\nu,\varepsilon}(t,y) - o(y)) = 0$ has to hold, both of which can be seen

by invoking Asm. 2 and Rem. 1.

In the spirit of [38, Theorem 1], set

$$X(t) := \left\{ x \in \mathbb{R} \cup \{-\infty, \infty\} : q_{\nu, \varepsilon}(t, x) - o(x) = \sup_{y \in \mathbb{R}} q_{\nu, \varepsilon}(t, y) - o(x) \right\} \neq \emptyset$$

for $t \in [0, T]$. Set $m(t) \coloneqq \sup_{y \in \mathbb{R}} q_{\nu,\varepsilon}(t, y) - o(y)$ for $t \in [0, T]$ and choose any $x \in X(t)$. Then, as a first case, if $x \in \{-\infty, \infty\}$, we have

$$\lim_{y \to \infty} \partial_t \left(q_{\nu,\varepsilon}(t,y) - o(y) \right) = \lim_{y \to \infty} \partial_t q_{\nu,\varepsilon}(t,y) = 0 = \partial_t \left(q_{\nu,\varepsilon}(t,x) - o(x) \right).$$
(8)

In the **second case**, if $x \in \mathbb{R}$, then, according to standard optimality conditions [39, Chapter 2], we obtain

$$\partial_x q_{\nu,\varepsilon}(t,x) - o'(x) = 0 \qquad \wedge \qquad \partial_x^2 q_{\nu,\varepsilon}(t,x) - o''(x) \le 0 \tag{9}$$

Then, we can differentiate as follows:

$$\begin{aligned} \partial_t \big(q_{\nu,\varepsilon}(t,x) - o(x) \big) &= \partial_t q_{\nu,\varepsilon}(t,x) \\ \stackrel{(1)}{=} - \tilde{V}_{\varepsilon}'(o(x) - q_{\nu,\varepsilon}(t,x)) \underbrace{(o'(x) - \partial_x q_{\nu,\varepsilon}(t,x))}_{=0} q_{\nu,\varepsilon}(t,x) \\ - \tilde{V}_{\varepsilon} \underbrace{(o(x) - q_{\nu,\varepsilon}(t,x))}_{=-m(t)} \underbrace{\partial_x q_{\nu,\varepsilon}(t,x)}_{=o'(x)} + \nu \underbrace{(\partial_x^2 q_{\nu,\varepsilon}(t,x) - o''(x))}_{\leq 0} \end{aligned}$$

Using Eq. (9),

$$\leq \left| \tilde{V}_{\varepsilon} \big(- m(t) \big) \right| \| o' \|_{L^{\infty}(\mathbb{R})}$$

Now, $q_{\nu,\varepsilon} - o \in W^{2,\infty}([0,T] \times \mathbb{R})$, and $\lim_{x \to \pm\infty} q_{\nu,\varepsilon}(t,x) - o(x)$ exists for all $t \in [0,T]$ because of Asm. 2 and Rem. 1. Similarly,

$$\lim_{x \to \pm \infty} \partial_t \left(q_{\nu,\varepsilon}(t,x) - o(x) \right) = 0 \ \forall t \in [0,T].$$

Altogether, [38, Theorem 1] is applicable, and combining both cases for a.e. $t \in (0, T)$ and all $x \in X(t)$ yields

$$\frac{\mathrm{d}}{\mathrm{d}t} \sup_{y \in \mathbb{R}} q_{\nu,\varepsilon}(t,y) - o(y) = m'(t) \le \partial_t \left(q_{\nu,\varepsilon}(t,x) - o(x) \right) = \|o'\|_{L^{\infty}(\mathbb{R})} \left| \tilde{V}_{\varepsilon} \left(-m(t) \right) \right|$$

Now, consider the Cauchy problem for the corresponding differential equation (instead of the inequality)

$$\tilde{m}'(t) = \|o'\|_{L^{\infty}(\mathbb{R})} \left| \tilde{V}_{\varepsilon}(-\tilde{m}(t)) \right|, \quad \tilde{m}(0) = \sup_{y \in \mathbb{R}} q_{0,\nu}(y) - o(y)$$

on [0, T] that uniquely defines \tilde{m} on [0, T] because \tilde{V}_{ε} is globally Lipschitz (see Asm. 1). Since $\tilde{V}_{\varepsilon}(0) = 0$ and $-\tilde{m}(0) > 0$, $-\tilde{m}(t) > 0$ for all $t \in [0, T]$ because equilibria are not allowed to be crossed. We also note that \tilde{m} does not depend on ν ! Then, the ODE comparison principle (see, e.g., [40, Lemma 1.2, p.24]) implies for $(t, y) \in [0, T] \times \mathbb{R}$,

$$q_{\nu,\varepsilon}(t,y) - o(y) \le m(t) \le \tilde{m}(t) < 0$$

This yields the claim.

So, in fact, the obstacle is not even hit at all for $\varepsilon \in \mathbb{R}_{>0}$, which also implies that the mass is transported with a positive velocity.

The same argument also allows for the proof of a minimum principle:

Theorem 8 (Minimum principle). Let $q_{\nu,\varepsilon}$ be a solution to Eq. (3) for $\nu, \varepsilon > 0$. Then,

$$q_{\nu,\varepsilon}(t,x) \ge -\frac{\nu \|o''\|_{L^{\infty}(\mathbb{R})}}{\|o'\|_{L^{\infty}(\mathbb{R})}\tilde{K}} \left(-1 + \exp\left(\tilde{K}\|o'\|_{L^{\infty}(\mathbb{R})}t\right)\right)$$
(10)

for all $(t,x) \in [0,T] \times \mathbb{R}$, where $c_0 \coloneqq \inf_{x \in \mathbb{R}} o(x) > 0$, $\tilde{K} := \|V_{\varepsilon}'\|_{L^{\infty}([c_0,\infty))}$. We recall Asm. 1 to know that \tilde{K} is finite.

Proof. The proof is similar to the previous one that shows that the obstacle is respected. It is again based on [38, Theorem 1]. \Box

Of note, the solution can be negative for $\nu, \varepsilon \in \mathbb{R}_{>0}$; however, in the limit, it will satisfy the expected lower bound of 0.

This also allows us to state bounds for the entropy solution:

Remark 3 (Bounds on entropy solution). Let $\varepsilon > 0$. Because L^1_{loc} -convergence implies point-wise convergence a.e. of a subsequence, the entropy solution $q_{\varepsilon} \in L^1_{loc}(\mathbb{R})$ from Thm. 6 fulfills the bounds

$$0 \le q_{\varepsilon}(t,x) < o(x)$$
 for a.e. $(t,x) \in (0,T) \times \mathbb{R}$.

This can be obtained by letting $\nu \searrow 0$ in Eq. (10) and Eq. (5) point-wise a.e.. Here, we note that an upper bound \tilde{m} exists on $q_{\nu,\varepsilon} - o$ such that $\tilde{m} < 0$ and does not depend on ν , as stated in the proof of Lem. 7. This allows for the strict inequality $q_{\varepsilon} < o$ to hold pointwise a.e.

4 OSL bounds for $q_{\nu,\varepsilon}$ and $V_{\varepsilon}(o-q_{\nu,\varepsilon})$ uniformly in ν, ε

As stated earlier, the condition $o' \leq 0$ gives uniform TV-bounds of $q_{\nu,\varepsilon}$ in ν and ε . We now want to find OSL bounds that provide a more general compactness result in $\nu, \varepsilon > 0$. We need the following:

Lemma 9 (OSL bounds and compactness). Let $\Omega \subseteq \mathbb{R}$ be open and $(f_{\mu})_{\mu>0} \subseteq C^{1}(\Omega)$ be a sequence of functions such that

$$C_1 \coloneqq \sup_{\mu > 0} \|f_{\mu}\|_{L^{\infty}(\Omega)} < \infty \text{ and } C_2 \coloneqq \sup_{\mu > 0} \sup_{x \in \Omega} f'_{\mu}(x) < \infty.$$

Then, $(f_{\mu})_{\mu>0} \subseteq BV(V)$ for every compact $V \subseteq \Omega$, and there exist $f \in L^{1}_{loc}(\Omega)$ and a subsequence $(\mu_{k})_{k\in\mathbb{N}}$ fulfilling $\lim_{k\to\infty} \mu_{k} = 0$ such that

$$\lim_{k \to \infty} \|f_{\mu_k} - f\|_{L^1_{loc}(\Omega)} = 0.$$

Proof. First, we prove that $|f_{\mu}|_{\mathrm{TV}(V)}$ is finite and does not depend on μ for every compact $V \subseteq \overline{\Omega}$. Let $V \subseteq \Omega$ be compact, and a compact interval $K \supseteq V$. Set $f_{\mu}^{+} := \max\{f_{\mu}, 0\}$ and $f_{\mu}^{-} := -\{f_{\mu}, 0\}$ for any $\mu > 0$.

$$\begin{split} \int_{V} |f'_{\mu}(x)| \, \mathrm{d}x &\leq \int_{K} |f'_{\mu}(x)| \, \mathrm{d}x = \int_{K} f'_{\mu}(x)^{+} + f'_{\mu}(x)^{-} \, \mathrm{d}x \\ &= 2 \int_{K} f'_{\mu}(x)^{+} \, \mathrm{d}x - \int_{K} f_{\mu}(x)' \, \mathrm{d}x \\ &= 2 \int_{K} \underbrace{f'_{\mu}(x)^{+}}_{\leq C_{2}} \, \mathrm{d}x - f_{\mu}(\sup K) + f_{\mu}(\inf K) \leq 2|K|C_{2} + 2C_{1} \end{split}$$

Consequently, $|f_{\mu}|_{\text{TV}(V)} \leq 2|K|C_2 + 2C_1$, and per [41, Theorem 13.35], the classical compactness result for functions of bounded variation gives convergence to f in $L^1_{\text{loc}}(\Omega)$ along a subsequence.

Note that the result from Lem. 9 also holds if $\inf_{\mu>0} \inf_{x\in\Omega} f'_{\mu}(x) > -\infty$ instead of $\sup_{\mu>0} \sup_{x\in\Omega} f'_{\mu}(x) < \infty$. To obtain this, replace f with -f and use Lem. 9. Unfortunately, linear conservation laws have no intrinsic "smoothing" because of the lack of Oleinik-type estimates. However, as we later want to pass to the limit nevertheless, we impose an OSL condition on the initial datum, which is preserved over time (the same cannot easily be obtained with a TV argument).

Assumption 10 (Initial datum revisited). In addition to Asm. 2, we assume for the remainder of Sec. 4 that the initial datum is OSL continuous from below, i.e., for a $C \in \mathbb{R}$, the following holds:

$$\inf_{(x,y)\in\mathbb{R}^2} \frac{q_0(x) - q_0(y)}{x - y} \ge C.$$
(11)

In the following, we derive OSL bounds for $q_{\nu,\varepsilon}$ uniform in ν, ε : **Theorem 11** (OSL condition for $q_{\nu,\varepsilon}$ in space). There exists $\nu(\varepsilon) \in \mathbb{R}_{>0}$ with $\lim_{\varepsilon \searrow 0} \nu(\varepsilon) = 0$ such that $\forall \nu \in \mathbb{R}_{>0}$ with $0 \le \nu \le \nu(\varepsilon)$, and it holds $\forall (t, x) \in [0, T] \times \mathbb{R}$ that

$$\partial_x q_{\nu,\varepsilon}(t,x) - o'(x) > \min\left\{-2\|o'\|_{L^{\infty}(\mathbb{R})}, \underset{y \in \mathbb{R}}{\operatorname{ess\,inf}}\left(q'_0(y) - o'(y)\right)\right\} - T\|o''\|_{L^{\infty}(\mathbb{R})},$$

where the right-hand side term is bounded thanks to Asm. 10.

Consequently, with $o' \in L^{\infty}(\mathbb{R})$ —see Asm. 2— $\forall \nu \in \mathbb{R}_{>0}$ with $0 \leq \nu \leq \nu(\varepsilon)$ and $\forall (t,x) \in [0,T] \times \mathbb{R}$,

$$\partial_x q_{\nu,\varepsilon}(t,x) > \min\{-2\|o'\|_{L^{\infty}(\mathbb{R})}, \operatorname{ess\,inf}_{y \in \mathbb{R}} \left(q'_0(y) - o'(y)\right)\} - T\|o''\|_{L^{\infty}(\mathbb{R})} - \|o'\|_{L^{\infty}(\mathbb{R})}.$$

Here, $\operatorname{ess\,inf}_{y\in\mathbb{R}}\left(q_0'(y) - o'(y)\right)$ is interpreted as the lower OSL bound of $q_0 - o$. *Proof.* As before, we apply [38, Theorem 1]. Let

$$Z(t) \coloneqq \left\{ x \in \mathbb{R} \cup \{-\infty, \infty\} : \partial_x q_{\nu,\varepsilon}(t,x) - o'(x) = \inf_{y \in \mathbb{R}} \left(\partial_y q_{\nu,\varepsilon}(t,y) - o'(y) \right) \right\}$$

and $m(t) \coloneqq (\inf_{y \in \mathbb{R}} \partial_y q_{\nu,\varepsilon}(t,y) - o'(y))$ for all $t \in [0,T]$. Choose $x \in Z(t)$. We distinguish two cases:

The first case is $x \in \{-\infty, \infty\} \cap Z(t)$. Here,

$$\partial_t \left(\partial_x q_{\nu,\varepsilon}(t,x) - o'(x) \right) = 0$$

since Eq. (8) and $o' \in H^1(\mathbb{R})$ as in Asm. 2.

The second case is $x \in \mathbb{R} \cap Z(t)$. Once again, optimality conditions lead to

$$\partial_x^2 q_{\nu,\varepsilon}(t,x) - o''(x) = 0 \quad \text{and} \quad \partial_x^3 q_{\nu,\varepsilon}(t,x) - o'''(x) \ge 0, \tag{12}$$

and we then obtain

$$\partial_t \big(\partial_x q_{\nu,\varepsilon}(t,x) - o'(x) \big) \\= \partial_{xt}^2 q_{\nu,\varepsilon}(t,x).$$

Plugging in the strong form Eq. (3), we have

$$\begin{split} &= \partial_x \Big(-V_{\varepsilon}'(o(x) - q_{\nu,\varepsilon}(t,x))(o'(x) - \partial_x q_{\nu,\varepsilon}(t,x))q_{\nu,\varepsilon}(t,x) \\ &- V_{\varepsilon}(o(x) - q_{\nu,\varepsilon}(t,x))\partial_x q_{\nu,\varepsilon}(t,x) + \nu(\partial_x^2 q_{\nu,\varepsilon}(t,x) - o''(x))) \Big) \\ &= -V_{\varepsilon}''(o(x) - q_{\nu,\varepsilon}(t,x))\underbrace{(o'(x) - \partial_x q_{\nu,\varepsilon}(t,x))}_{=m(t)^2} q_{\nu,\varepsilon}(t,x) \\ &- V_{\varepsilon}'(o(x) - q_{\nu,\varepsilon}(t,x))\underbrace{(o''(x) - \partial_x^2 q_{\nu,\varepsilon}(t,x))}_{=0} q_{\nu,\varepsilon}(t,x) \\ &- 2V_{\varepsilon}'(o(x) - q_{\nu,\varepsilon}(t,x))\underbrace{(o'(x) - \partial_x q_{\nu,\varepsilon}(t,x))}_{=-m(t)} \underbrace{\partial_x q_{\nu,\varepsilon}(t,x)}_{m(t)+o'(x)} \\ &- V_{\varepsilon}(o(x) - q_{\nu,\varepsilon}(t,x))\underbrace{\partial_x^2 q_{\nu,\varepsilon}(t,x)}_{=o''(x)} + \underbrace{\nu(\partial_x^3 q_{\nu,\varepsilon}(t,x) - o'''(x))}_{\geq 0} \\ \\ &\stackrel{(12)}{\geq} -V_{\varepsilon}''(o(x) - q_{\nu,\varepsilon}(t,x))m(t)^2 q_{\nu,\varepsilon}(t,x) + V_{\varepsilon}'(o(x) - q_{\nu,\varepsilon}(t,x))\Big(2m(t)^2 + m(t)o'(x)) \end{split}$$

$$-V_{\varepsilon}(o(x)-q_{\nu,\varepsilon}(t,x))o''(x).$$

We write $q_{\nu,\varepsilon} = q_{\nu,\varepsilon}^+ - q_{\nu,\varepsilon}^-$ for the positive and negative parts, respectively, and incorporate $-V_{\varepsilon}''m^2q_{\nu,\varepsilon}^+ \ge 0$:

$$V_{\varepsilon}^{\prime\prime}(o(x) - q_{\nu,\varepsilon}(t,x))m(t)^2 q_{\nu,\varepsilon}(t,x)^- + V_{\varepsilon}^{\prime}(o(x) - q_{\nu,\varepsilon}(t,x))(2m(t)^2 + m(t)o^{\prime}(x)) - V_{\varepsilon}(o(x) - q_{\nu,\varepsilon}(t,x))o^{\prime\prime}(x)$$

Per Asm. 2, $V_{\varepsilon}''(x) = \frac{1}{\varepsilon^2} \frac{V''(\frac{x}{\varepsilon})}{V'(\frac{x}{\varepsilon})} V'\left(\frac{x}{\varepsilon}\right) \ge \frac{1}{\varepsilon} \underbrace{v_{2,1}^{-1}}_{<0} V_{\varepsilon}'(x)$ holds for $x \in \mathbb{R}$, and we estimate

as follows

$$\geq V_{\varepsilon}'(o(x) - q_{\nu,\varepsilon}(t,x)) \left(m(t)^2 \frac{v_{2,1}^- q_{\nu,\varepsilon}(t,x)}{\varepsilon} + 2m(t)^2 + m(t)o'(x) \right) \\ - \|V_{\varepsilon}\|_{L^{\infty}(\mathbb{R})} \|o''\|_{L^{\infty}(\mathbb{R})}.$$

According to Eq. (10), there exists $\nu(\varepsilon)$ s.t. $q_{\nu,\varepsilon}^- < \frac{\varepsilon}{v_{2,1}^-}$ for all $0 < \nu < \nu(\varepsilon)$. Because $\lim_{y \to \pm \infty} \partial_y q_{\nu,\varepsilon}(t,y) - o'(y) = 0$, $m(t) \leq 0$ holds, so we can continue the estimate as

$$\geq V_{\varepsilon}'(o(x) - q_{\nu,\varepsilon}(t,x)) \left(m(t)^2 + \|o'\|_{L^{\infty}(\mathbb{R})} m(t) \right) - \|o''\|_{L^{\infty}(\mathbb{R})} \|V_{\varepsilon}\|_{L^{\infty}(\mathbb{R})}.$$

Applying [38, Theorem 1] and combining both cases, we obtain, for a.e. $t \in (0, T)$,

$$m'(t) \ge \min \left\{ V_{\varepsilon}'(o(x) - q_{\nu,\varepsilon}(t,x)) \left(m(t)^2 + \|o'\|_{L^{\infty}(\mathbb{R})} m(t) \right) - \|o''\|_{L^{\infty}(\mathbb{R})} \|V_{\varepsilon}\|_{L^{\infty}(\mathbb{R})}, 0 \right\}$$

Then, fix any $t \in [0,T]$. Now, if $m(t) \geq -2 \|o'\|_{L^{\infty}(\mathbb{R})}$, we have a desirable lower bound on m.

However, if $m(t) < -2 \| o' \|_{L^{\infty}(\mathbb{R})}$, then $m(t)^2 + \| o' \|_{L^{\infty}(\mathbb{R})} m(t) \ge 0$. So,

$$m'(t) \ge -\|o''\|_{L^{\infty}(\mathbb{R})}\|V_{\varepsilon}\|_{L^{\infty}(\mathbb{R})} = -\|o''\|_{L^{\infty}(\mathbb{R})}$$

holds, which implies, after integration,

$$m(t) \ge \min\{m(0), -2\|o'\|_{L^{\infty}(\mathbb{R})}\} - t\|o''\|_{L^{\infty}(\mathbb{R})},$$

which is the claimed inequality. Here, we remind the reader that $m(0) > -\infty$ is assumed in Eq. (11).

A similar result can be derived for the velocity $V_{\varepsilon}(o-q_{\nu,\varepsilon})$. **Theorem 12.** (OSL condition for $V_{\varepsilon}(o - q_{\nu,\varepsilon})$ in space) Let $\varepsilon, \nu \in \mathbb{R}_{>0}$, $q_{\varepsilon,\nu}$ be a solution to Eq. (3). Then, for fixed ε , there exists $\nu(\varepsilon) > 0$ with $\lim_{\varepsilon \searrow 0} \nu(\varepsilon) = 0$ such that

$$\partial_x \left(V_{\varepsilon} \left(o(x) - q_{\nu,\varepsilon}(t,x) \right) \right)$$

$$\leq \max\left\{C_1, C_2 \sup_{\varepsilon > 0} \|V_{\varepsilon}'\|_{L^{\infty}([k_0,\infty))}, \operatorname{ess\,sup}_{y \in \mathbb{R}}(o'(y) - q'_0(y)) \sup_{\varepsilon > 0} \|V_{\varepsilon}'\|_{L^{\infty}([c_0,\infty))}\right\}$$

for all $0 < \nu \leq \nu(\varepsilon)$, $(t, x) \in (0, T) \times \mathbb{R}$, where

$$k_0 \coloneqq \frac{\inf_{y \in \mathbb{R}} o(y)}{2} \tag{13}$$

$$c_0 \coloneqq \underset{y \in \mathbb{R}}{\operatorname{ess\,inf}}(o(y) - q_0(y)) \tag{14}$$

$$v \coloneqq \left\| \frac{V''}{V'} \right\|_{L^{\infty}(\mathbb{R})} \tag{15}$$

$$C_{1} \coloneqq \sqrt{\frac{(v+2\|V'\|_{L^{\infty}([0,\infty)]})^{2}\|o'\|_{L^{\infty}(\mathbb{R})}^{2}}{\left(v_{2,1}^{+}\inf_{y\in\mathbb{R}}o(y)\right)^{2}} + \frac{2\|V'\|_{L^{\infty}([0,\infty))}(-\|o''\|_{L^{\infty}(\mathbb{R})}-1)}{v_{2,1}^{+}\inf_{y\in\mathbb{R}}o(y)}}$$
(16)

$$+\frac{2(v+2\|V'\|_{L^{\infty}([0,\infty))})\|o'\|_{L^{\infty}(\mathbb{R})}}{v_{2,1}^{+}\inf_{y\in\mathbb{R}}o(y)}\tag{17}$$

$$C_2 \coloneqq T \| o'' \|_{L^{\infty}(\mathbb{R})} - \min\left\{ -2 \| o' \|_{L^{\infty}(\mathbb{R})}, \operatorname{essinf}_{y \in \mathbb{R}} \left(q'_0(y) - o'(y) \right) \right\}$$
(18)

Proof. As before, we want to invoke [38, Theorem 1]. Fix $t \in (0, T)$ and $\nu, \varepsilon > 0$. Since

$$\partial_x \big(V_{\varepsilon}(o(x) - q_{\nu,\varepsilon}(t,x)) \big) = V'_{\varepsilon}(o(x) - q_{\nu,\varepsilon}(t,x))(o'(x) - \partial_t q_{\nu,\varepsilon}(t,x)),$$

for all $x \in \mathbb{R}$, from $\lim_{x \to \pm \infty} o'(x) - \partial_x q_{\nu,\varepsilon}(t,x) = 0$ and the boundedness of V'_{ε} , we obtain

$$\lim_{x \to \pm \infty} \partial_x \left(V_{\varepsilon}(o(x) - q_{\nu,\varepsilon}(t,x)) \right) = 0.$$

The same argument can be employed for derivatives thereof. Therefore, [38] is indeed applicable.

Now, choose $x \in X(t)$, where

$$X(t) \coloneqq \left\{ x \in \mathbb{R} \cup \{-\infty, \infty\} : \partial_x V_{\varepsilon} \big(o(x) - q_{\nu, \varepsilon}(t, x) \big) = \sup_{y \in \mathbb{R}} \partial_y V_{\varepsilon} \big(o(y) - q_{\nu, \varepsilon}(t, y) \big) \right\}.$$

We distinguish two cases.

Case 1: $(x \in \{-\infty, \infty\})$ Then,

$$\begin{split} \lim_{y \to \infty} \partial_t \partial_y \big(V_{\varepsilon}(o(y) - q_{\nu,\varepsilon}(t,y)) \big) \\ &= \lim_{y \to \infty} V_{\varepsilon}''(o(y) - q_{\nu,\varepsilon}(t,y)) (-\partial_t q_{\nu,\varepsilon}(t,y)) (o'(y) - \partial_y q_{\nu,\varepsilon}(t,y)) \\ &+ V_{\varepsilon}'(o(y) - q_{\nu,\varepsilon}(t,y)) (-\partial_t y q_{\nu,\varepsilon}(t,y)) = 0, \end{split}$$

which follows from Rem. 1.

Case 2: $(x \in \mathbb{R})$ For the sake of clarity, we (have to) introduce the *following* abbreviations (for $k \in \mathbb{N}_{\geq 0}$, multi-index α):

$$V_{\varepsilon}^{(k)} \coloneqq V_{\varepsilon}^{(k)}(o(x) - q_{\nu,\varepsilon}(t,x)), \ q_{\nu,\varepsilon} \coloneqq q_{\nu,\varepsilon}(t,x), \ o^{(k)} \coloneqq o^{(k)}(x), \ \partial_{\alpha}q_{\nu,\varepsilon} \coloneqq \partial_{\alpha}q_{\nu,\varepsilon}(t,x)$$

$$V^{(k)} \coloneqq V^{(k)} \left(\frac{o(x) - q_{\nu,\varepsilon}(t,x)}{\varepsilon}\right) = \varepsilon^k V_{\varepsilon}^{(k)}$$

Note that $V^{(k)}$ needs to be *sharply distinguished* from $V_{\varepsilon}^{(k)}$; see also Asm. 1. Moreover, the notation $V^{(k)}$ is a bit misleading, as it still depends on ε . For $V^{(k)}$, however, we have uniform L^{∞} -bounds in ε , which further motivates omitting the index.

In the following, we sometimes write $V'_{\varepsilon} \cdot (o'' - \partial_x^2 q_{\nu,\varepsilon})$ rather than, e.g., $V'_{\varepsilon}(o - q_{\nu,\varepsilon})(o'' - \partial_x^2 q_{\nu,\varepsilon})$, i.e., V_{ε} without its argument. Remember that X(t) is defined so that $\forall x \in X(t)$,

$$\partial_x V_{\varepsilon} \big(q_{\nu,\varepsilon}(t,x) - o(x) \big) = \sup_{y \in \mathbb{R}} \partial_y V_{\varepsilon} \big(q_{\nu,\varepsilon}(t,y) - o(y) \big).$$

So, based on standard optimality conditions, the following must hold:

$$\partial_x^2 V_{\varepsilon} = \partial_x^2 \big(V_{\varepsilon}(o(x) - q_{\nu,\varepsilon}(t,x)) \big) = 0 \quad \wedge \quad \partial_x^3 V_{\varepsilon} = \partial_x^3 \big(V_{\varepsilon}(o(x) - q_{\nu,\varepsilon}(t,x)) \big) \le 0.$$
(19)

Expanding the above expressions (in the abbreviated notation), we obtain

$$\partial_x^2 V_{\varepsilon} = V_{\varepsilon}'' \cdot (o' - \partial_x q_{\nu,\varepsilon})^2 + V_{\varepsilon}' \cdot (o'' - \partial_x^2 q_{\nu,\varepsilon})$$

$$\partial_x^3 V_{\varepsilon} = V_{\varepsilon}''' \cdot (o' - \partial_x q_{\nu,\varepsilon})^3 + 3V_{\varepsilon}'' \cdot (o' - \partial_x q_{\nu,\varepsilon})(o'' - \partial_x^2 q_{\nu,\varepsilon}) + V_{\varepsilon}' \cdot (o''' - \partial_x^3 q_{\nu,\varepsilon}),$$

and setting $m(t) \coloneqq \sup_{y \in \mathbb{R}} \partial_y V_{\varepsilon} (q_{\nu,\varepsilon}(t,y) - o(y)) = V_{\varepsilon} (q_{\nu,\varepsilon}(t,x) - o(x))$ yields

$$\partial_{t}\partial_{x}V_{\varepsilon} = V_{\varepsilon}'' \cdot (o' - \partial_{x}q_{\nu,\varepsilon})(-\partial_{t}q_{\nu,\varepsilon}) + V_{\varepsilon}' \cdot (-\partial_{tx}q_{\nu,\varepsilon})$$

$$\stackrel{(1)}{=} V_{\varepsilon}'' \cdot (o' - \partial_{x}q_{\nu,\varepsilon}) \left(V_{\varepsilon}' \cdot (o' - \partial_{x}q_{\nu,\varepsilon})q_{\nu,\varepsilon} + V_{\varepsilon} \cdot \partial_{x}q_{\nu,\varepsilon} - \nu(\partial_{x}^{2}q_{\nu,\varepsilon} - o'')\right)$$

$$+ V_{\varepsilon}' \cdot \partial_{x} \left(V_{\varepsilon}' \cdot (o' - \partial_{x}q_{\nu,\varepsilon})q_{\nu,\varepsilon} + V_{\varepsilon} \cdot \partial_{x}q_{\nu,\varepsilon} - \nu(\partial_{x}^{2}q_{\nu,\varepsilon} - o'')\right)$$

$$= V_{\varepsilon}'' \cdot (o' - \partial_{x}q_{\nu,\varepsilon}) \left(V_{\varepsilon}' \cdot (o' - \partial_{x}q_{\nu,\varepsilon})q_{\nu,\varepsilon} + V_{\varepsilon} \cdot \partial_{x}q_{\nu,\varepsilon} - \nu(\partial_{x}^{2}q_{\nu,\varepsilon} - o'')\right)$$

$$+ V_{\varepsilon}' \cdot \left(V_{\varepsilon}'' \cdot (o' - \partial_{x}q_{\nu,\varepsilon})^{2}q_{\nu,\varepsilon} + V_{\varepsilon}' \cdot (o'' - \partial_{x}^{2}q_{\nu,\varepsilon})q_{\nu,\varepsilon} + V_{\varepsilon} \cdot \partial_{x}^{2}q_{\nu,\varepsilon} - \nu(\partial_{x}^{2}q_{\nu,\varepsilon} - o''')\right). \tag{20}$$

We apply Eq. (19) to Eq. (20), use $\partial_x q_{\nu,\varepsilon} = -(o' - \partial_x q_{\nu,\varepsilon}) + o'$ twice, $\partial_{xx}^2 q_{\nu,\varepsilon} = -(o'' - \partial_{xx}^2 q_{\nu,\varepsilon}) + o''$, and expand to arrive at the estimate

$$\leq V_{\varepsilon}'' \cdot V_{\varepsilon}' \cdot (o' - \partial_x q_{\nu,\varepsilon})^2 q_{\nu,\varepsilon} - V_{\varepsilon}'' \cdot V_{\varepsilon} \cdot (o' - \partial_x q_{\nu,\varepsilon})^2 + V_{\varepsilon}'' \cdot V_{\varepsilon} \cdot (o' - \partial_x q_{\nu,\varepsilon})o' - \nu V_{\varepsilon}'' \cdot (o' - \partial_x q_{\nu,\varepsilon})(\partial_x^2 q_{\nu,\varepsilon} - o'') - 2 (V_{\varepsilon}' \cdot (o' - \partial_x q_{\nu,\varepsilon}))^2 + 2 (V_{\varepsilon}')^2 \cdot (o' - \partial_x q_{\nu,\varepsilon})o' - V_{\varepsilon} \cdot V_{\varepsilon}' \cdot (o'' - \partial_x^2 q_{\nu,\varepsilon}) + V_{\varepsilon}' \cdot V_{\varepsilon} \cdot o'' - \nu V_{\varepsilon}' \cdot (\partial_x^3 q_{\nu,\varepsilon} - o''').$$

$$(21)$$

Using Eq. (19) as well as the identities $\partial_x^2 q_{\nu,\varepsilon} - o'' = \frac{1}{V_{\varepsilon}'} \left(\partial_x^2 V_{\varepsilon} - V_{\varepsilon}'' \cdot (o' - \partial_x q_{\nu,\varepsilon})^2 \right)$ and $V_{\varepsilon}' \cdot (o'' - \partial_x^2 q_{\nu,\varepsilon}) = \partial_x^2 V_{\varepsilon} - V_{\varepsilon}'' \cdot (o' - \partial_x q_{\nu,\varepsilon})^2$, we obtain

$$= \frac{V_{\varepsilon}''}{V_{\varepsilon}'} \left(V_{\varepsilon}' \cdot (o' - \partial_x q_{\nu,\varepsilon}) \right)^2 q_{\nu,\varepsilon} - \frac{V_{\varepsilon}'' V_{\varepsilon}}{(V_{\varepsilon}')^2} \left(V_{\varepsilon}' \cdot (o' - \partial_x q_{\nu,\varepsilon}) \right)^2 - \frac{V_{\varepsilon}''}{V_{\varepsilon}'} V_{\varepsilon} V_{\varepsilon}' \cdot (o' - \partial_x q_{\nu,\varepsilon}) o' + \nu \frac{(V_{\varepsilon}'')^2}{(V_{\varepsilon}')^4} \left(V_{\varepsilon}' \cdot (o' - \partial_x q_{\nu,\varepsilon}) \right)^3 - 2 \left(V_{\varepsilon}' \cdot (o' - \partial_x q_{\nu,\varepsilon}) \right)^2 + 2 (V_{\varepsilon}')^2 \cdot (o' - \partial_x q_{\nu,\varepsilon}) o' + \frac{V_{\varepsilon} V_{\varepsilon}''}{(V_{\varepsilon}')^2} \left(V_{\varepsilon}' \cdot (o' - \partial_x q_{\nu,\varepsilon}) \right)^2 + V_{\varepsilon}' V_{\varepsilon} o'' - \nu \frac{V_{\varepsilon}'''}{(V_{\varepsilon}')^3} (V_{\varepsilon}' \cdot (o' - \partial_x q_{\nu,\varepsilon}))^3 - 3\nu V_{\varepsilon}'' \cdot (o' - \partial_x q_{\nu,\varepsilon}) (o'' - \partial_x^2 q_{\nu,\varepsilon}) dv_{\varepsilon} dv_{$$

and eventually replacing $V'_{\varepsilon} \cdot (o' - \partial_x q_{\nu,\varepsilon})$ with m(t) in most cases yields

$$\leq \frac{V_{\varepsilon}''}{V_{\varepsilon}'}m(t)^2 q_{\nu,\varepsilon} - \frac{V_{\varepsilon}''}{V_{\varepsilon}'}V_{\varepsilon}o'm(t) + \nu \frac{(V_{\varepsilon}'')^2}{(V_{\varepsilon})^4} \left(V_{\varepsilon}' \cdot (o' - \partial_x q_{\nu,\varepsilon})\right)^3 - 2m(t)^2 + 2V_{\varepsilon}'o'm(t) + V_{\varepsilon}'V_{\varepsilon}o'' - \nu \frac{V_{\varepsilon}'''}{(V_{\varepsilon}')^3}m(t)^3 + 3\nu \frac{(V_{\varepsilon}'')^2}{V_{\varepsilon}'} \cdot (o' - \partial_x q_{\nu,\varepsilon})^3.$$

We factor out V_{ε}' for several expressions. Additionally, for $0 \leq \nu \leq \nu(\varepsilon) \leq \varepsilon$ according to Thm. 11, we have C > 0 independent of $\nu, \varepsilon > 0$ (see Eq. (25)) such that $\partial_x q_{\nu,\varepsilon} - o' \geq C$ for all $(t, x) \in [0, T] \times \mathbb{R}$. Therefore, we can estimate $(o' - \partial_x q_{\nu,\varepsilon})^3 \leq C^3$ if ν is sufficiently small:

$$\leq -2m(t)^2 + V_{\varepsilon}' \left(\frac{V_{\varepsilon}''}{(V_{\varepsilon}')^2} m(t)^2 q_{\nu,\varepsilon} - \frac{V_{\varepsilon} V_{\varepsilon}''}{(V_{\varepsilon}')^2} o'm(t) + \nu \frac{(V_{\varepsilon}'')^2}{(V_{\varepsilon}')^2} C^3 + 2m(t)o' \right)$$
(22)

$$+ V_{\varepsilon}o'' - \nu \frac{V_{\varepsilon}'''}{(V_{\varepsilon}')^4} m(t)^3 + 3\nu \frac{(V_{\varepsilon}'')^2}{(V_{\varepsilon}')^2} C^3 \bigg)$$
(23)

Therefore, combining both cases, we show that by using [38, Theorem 1], for a.e. $t \in (0,T)$,

$$m'(t) \leq \sup_{x \in X(t)} \max\left\{ -2m(t)^{2} + V_{\varepsilon}' \left(\frac{V_{\varepsilon}''}{(V_{\varepsilon}')^{2}} m(t)^{2} q_{\nu,\varepsilon} - \frac{V_{\varepsilon} V_{\varepsilon}''}{(V_{\varepsilon}')^{2}} o'm(t) + \nu \frac{(V_{\varepsilon}'')^{2}}{(V_{\varepsilon}')^{2}} C^{3} + 2m(t)o' + V_{\varepsilon}o'' - \nu \frac{V_{\varepsilon}'''}{(V_{\varepsilon}')^{4}} m(t)^{3} + 3\nu \frac{(V_{\varepsilon}'')^{2}}{(V_{\varepsilon}')^{2}} C^{3} \right\}, 0 \right\}.$$

$$(24)$$

Again, two cases have to be distinguished. For this, fix $t \in (0,T)$ such that m is differentiable (this is possible a.e.).

• Case 2.1: $(\exists x \in X(t) : q_{\nu,\varepsilon}(t,x) \leq \frac{o(x)}{2})$ In this case, a bound on *m* can be obtained immediately without Eq. (24): Choose some $x \in X(t)$ such that $q_{\nu,\varepsilon}(t,x) \leq \frac{o(x)}{2}$. If we choose $\nu > 0$ as in Thm. 11, then from Thm. 11,

$$o'(x) - \partial_x q_{\nu,\varepsilon}(t,x) \le T \| o'' \|_{L^{\infty}(\mathbb{R})} - \min\{-2\| o' \|_{L^{\infty}(\mathbb{R})}, L_{-}\} =: C$$
(25)

holds. Here, $L_{-} \coloneqq \operatorname{ess\,inf}_{y \in \mathbb{R}} (q'_{0}(y) - o'(y))$. Then, we can estimate the following:

$$m(t) = \sup_{y \in \mathbb{R}} \partial_y \left(V_{\varepsilon}(o(y) - q_{\nu,\varepsilon}(t,y)) \right) = \partial_x \left(V_{\varepsilon}(o(x) - q_{\nu,\varepsilon}(t,x)) \right)$$
$$= V_{\varepsilon}'(o(x) - q_{\nu,\varepsilon}(t,x))(o'(x) - \partial_x q_{\nu,\varepsilon}(t,x))$$
$$\stackrel{(25)}{\leq} CV_{\varepsilon}'(o(x) - q_{\nu,\varepsilon}(t,x))$$

where $q_{\nu,\varepsilon}(t,x) \leq \frac{o(x)}{2}$ and that V_{ε} is decreasing.

$$\leq CV_{\varepsilon}'\left(\frac{o(x)}{2}\right) \leq CV_{\varepsilon}'\left(\frac{\inf_{y\in\mathbb{R}}o(y)}{2}\right)$$

Since the infimum of o is positive per Asm. 2, the last expression has a uniform bound in $\varepsilon > 0$. Namely,

$$\sup_{y \in \mathbb{R}} \partial_y \left(V_{\varepsilon}(o(y) - q_{\nu,\varepsilon}(t,y)) \right) = \partial_x \left(V_{\varepsilon}(o(x) - q_{\nu,\varepsilon}(t,x)) \right) \le C \sup_{\varepsilon > 0} \|V_{\varepsilon}'\|_{L^{\infty}([k_0,\infty))},$$
(26)

where $k_0 := \frac{\inf_{y \in \mathbb{R}} o(y)}{2}$. • Case 2.2: $(q_{\nu,\varepsilon}(t,x) > \frac{o(x)}{2} \text{ for all } x \in X(t))$: Here, we estimate Eqs. (22) and (23) (once again in the abbreviated notation) using $q_{\nu,\varepsilon}(t,x) > \frac{o(x)}{2}$ and $\frac{V_{\varepsilon}''}{(V_{\varepsilon}')^2} < 0$:

$$m'(t) \leq -2m(t)^{2} + V_{\varepsilon}' \left(\frac{V_{\varepsilon}''}{(V_{\varepsilon}')^{2}} m(t)^{2} \frac{o(x)}{2} - \frac{V_{\varepsilon}V_{\varepsilon}''}{(V_{\varepsilon}')^{2}} o'm(t) + \nu \frac{(V_{\varepsilon}'')^{2}}{(V_{\varepsilon}')^{2}} C^{3} + 2m(t)o' + V_{\varepsilon}o'' - \nu \frac{V_{\varepsilon}'''}{(V_{\varepsilon}')^{4}} m(t)^{3} + 3\nu \frac{(V_{\varepsilon}'')^{2}}{(V_{\varepsilon}')^{2}} C^{3} \right)$$

Recall $-\inf_{y\in\mathbb{R}} o(y) < 0$ and the estimates from the fifth point of Asm. 1 to control the quotients. We set $v := \left\| \frac{V''}{V'} \right\|_{L^{\infty}([0,\infty)])} < \infty$, factor out $(V')^{-1}$ and estimate the other terms roughly:

$$\leq -2m(t)^{2} + V_{\varepsilon}' \left(\frac{1}{V'} \left(\frac{1}{2} \underbrace{v_{2,1}^{+}}_{<0 \text{ by Asm. 1}} \inf_{y \in \mathbb{R}} o(y)m(t)^{2} + v \|o'\|_{L^{\infty}(\mathbb{R})} |m(t)| + 2V' \|o'\|_{L^{\infty}(\mathbb{R})} |m(t)| \right) + \|o''\|_{L^{\infty}(\mathbb{R})} + \frac{4\nu v^{2}}{\varepsilon^{2}}C^{3} - \nu \frac{V_{\varepsilon}'''}{(V_{\varepsilon}')^{4}}m(t)^{3} \right)$$

We additionally require $\nu \leq \frac{\varepsilon^2}{4v^2C^3}$ to get remove ε .

$$\leq -2m(t)^{2} + V_{\varepsilon}' \Big(\frac{1}{V'} \Big(\frac{1}{2} v_{2,1}^{+} \inf_{y \in \mathbb{R}} o(y) m(t)^{2} + (v + 2 \|V'\|_{L^{\infty}(\mathbb{R})} \Big) \|o'\|_{L^{\infty}(\mathbb{R})} |m(t)| \Big)$$

$$+ \|o''\|_{L^{\infty}(\mathbb{R})} + 1 - \nu \frac{V_{\varepsilon}'''}{(V_{\varepsilon}')^4} m(t)^3 \right)$$

Let C_1 be as in Eq. (17). If $m(t) \leq C_1$, a feasible bound is found. If $m(t) > C_1$, then one can compute $\frac{1}{2}v_{2,1}^+ \inf_{y \in \mathbb{R}} o(y)m(t)^2 + (v+2||V'||_{L^{\infty}(\mathbb{R})})||o'||_{L^{\infty}(\mathbb{R})}|m(t)| < ||V'||_{L^{\infty}([0,\infty))}(-||o''||_{L^{\infty}(\mathbb{R})}-1)$. In particular, $-\nu V_{\varepsilon}^{'''}(V_{\varepsilon}')^{-4} \leq 0$. So, we estimate for this case

$$\leq -2m(t)^2 + V_{\varepsilon}' \left(\frac{\|V'\|_{L^{\infty}([0,\infty))}}{V'} \left(-\|o''\|_{L^{\infty}(\mathbb{R})} - 1 \right) + \|o''\|_{L^{\infty}(\mathbb{R})} + 1 \right)$$

$$\leq -2m(t)^2 \leq 0$$

Thus, if a maximal interval I exists such that $m(t) > C_1$ with C_1 as in Eq. (17) and $q_{\nu,\varepsilon}(t,x) > \frac{o(x)}{2}$ for all $t \in I$, then Eq. (24) becomes $m'(t) \leq 0$ for a.e. $t \in (0,T)$, whence

$$m(t) \le \max\{m(0), C_1\}$$

If no such interval I exists, then, by continuity, m stays below the maximum C_1 and the bound computed in Eq. (26).

Finally, in **all cases**, we obtain that

$$m(t) \le \max\left\{C_1, m(0), C_2 \sup_{\varepsilon > 0} \|V_{\varepsilon}'\|_{L^{\infty}([k_0, \infty))}\right\},\$$

with C_2 defined as in Eq. (18). It remains to prove that m(0) is bounded uniformly in ν, ε , which is a consequence of

$$m(0) = \sup_{x \in \mathbb{R}} \partial_x \left(V_{\varepsilon} \left(o(x) - q_0(x) \right) \right) \stackrel{Asm. \ 1, e)}{\leq} \operatorname{ess\,sup}_{y \in \mathbb{R}} (o'(y) - q'_0(y)) \sup_{\varepsilon > 0} \|V'_{\varepsilon}\|_{L^{\infty}([c_0, \infty))},$$

where $c_0 \coloneqq \operatorname{ess\,inf}_{y \in \mathbb{R}}(o(y) - q_0(y))$, which completes the proof.

The obtained results allow us to pass to the limit, i.e., $\varepsilon \searrow 0$:

Theorem 13 (Convergence of q_{ε} and $V_{\varepsilon}(o-q_{\varepsilon})$). Let Asm. 2 and Asm. 10 hold and q_{ε} be the solution to Eq. (1) according to Thm. 6 for $\varepsilon \in \mathbb{R}_{>0}$. Then, there exists $q^* \in C([0,T]; L^1_{loc}(\mathbb{R}))$ s.t. along a subsequence $(\varepsilon_k)_{k \in \mathbb{N}}$, $\lim_{k \to \infty} \varepsilon_k = 0$, and the following holds:

$$q_{\varepsilon_k} \to q^{\star} \text{ in } C([0,T]; L^1_{loc}(\mathbb{R})),$$

and there exists $V^{\star} \in L^{\infty}((0,T) \times \mathbb{R})$ such that

$$V_{\varepsilon_k}(o-q_{\varepsilon_k}) \stackrel{*}{\rightharpoonup} V^* \text{ in } L^{\infty}((0,T) \times \mathbb{R}).$$

Proof. This follows from Theorems 5, 6 and 11 and Lem. 9.

Remark 4 (Missing time compactness of V_{ε} to obtain strong convergence). Note that thanks to Thm. 12, one obtains by Lem. 9 total variation estimates of $V_{\varepsilon}(o-q_{\varepsilon}(t,\cdot))$ in a space uniform in $\varepsilon \in \mathbb{R}_{>0}$. However, we could not obtain the required (compare [37, Theorem 3]) time compactness for strong convergence in $C([0,T]; L^p_{loc}(\mathbb{R})), p \in [1,\infty),$ but we are left with a weaker form of convergence: Either the weak-star convergence in Thm. 13 or, for every $t \in [0,T]$, there exists $V_t^* \in L^{\infty}(\mathbb{R})$ and a sequence $(\varepsilon_k)_{k \in \mathbb{N}}$ with $\lim_{k\to\infty} \varepsilon_k = 0$ such that

$$\lim_{k \to \infty} \|V_{\varepsilon_k}(o(\cdot) - q_{\varepsilon_k}(t, \cdot)) - V_t^*\|_{L^1_{loc}(\mathbb{R})} = 0.$$

Indeed, V_t^* is OSL-bounded from above (see Thm. 12).

One may wonder what the velocity $V_{\varepsilon}(o - q_{\varepsilon})$ and solution q_{ε} satisfy in the limit. This is characterized in the following:

Lemma 14 (Dynamics of the solution and velocity in the limit $\varepsilon \to 0$). The sequence q_{ε_k} , $V_{\varepsilon_k}(o - q_{\varepsilon_k})$ of Thm. 13 satisfies in the limit $\varepsilon \to 0$ the following weak form of a conservation law for $\varphi \in C_c^1((-42, T) \times \mathbb{R})$:

$$\int_0^T \int_{\mathbb{R}} q^*(t,x) \partial_t \varphi(t,x) + V^*(t,x) q^*(t,x) \partial_x \varphi(t,x) \,\mathrm{d}x \,\mathrm{d}t = -\int_{\mathbb{R}} q_0(x) \varphi(0,x) \,\mathrm{d}x \quad (27)$$

Proof. This is a direct consequence of Thm. 13 together with the fact that the product of a strongly and a weakly-* convergent sequence converges weak-*. \Box

Remark 5 (Discontinuous conservation laws/relation to Lem. 14). Looking back to the regularized obstacle problem in Eq. (1), this equation converges to a discontinuous (in the solution and spatial variable (!)) conservation law as the velocity converges to a discontinuous function, i.e.,

$$\lim_{\varepsilon \to 0} V_{\varepsilon}(o(x) - q_{\varepsilon}(t, x)) \in \begin{cases} \{1\} & \text{if } q^*(t, x) < o(x) \\ [0, 1] & \text{if } q^*(t, x) = o(x), \end{cases} \qquad (t, x) \in (0, T) \times \mathbb{R}$$

and thus, we obtain

$$\int_0^T \!\!\!\int_{\mathbb{R}} q^*(t,x) \partial_t \varphi(t,x) + H(q^*(t,x))q^*(t,x) \partial_x \varphi(t,x) \,\mathrm{d}x \,\mathrm{d}t = -\int_{\mathbb{R}} q_0(x)\varphi(0,x) \,\mathrm{d}x,$$
(28)

where φ is a test function as in Lem. 14 and H is the Heaviside function with a value at zero that is undetermined, i.e., can attain any value in [0,1]. One may wonder whether applying results on the uniqueness of these discontinuous conservation laws is possible [42–47]; however, the discontinuous conservation law considered here does not seem to fit into the required frameworks.

5 Characterization of the limit $\varepsilon \searrow 0$ in specific cases

This section presents characterizations of the solutions approached in Thm. 13 under additional assumptions on its smoothness and more. They serve to build an intuition on what happens in this limit as well as how the dynamics behave at the obstacle.

Suppose we have a solution $q := q^*$ as in Eq. (27) that is piecewise smooth in the sense that it is smooth outside of a finite number of curves in (t, x), across which q or $\partial_x q$ has jump discontinuities. We take a representative that is also piecewise smooth, such that q and $\partial_x q$ have well-defined traces at every point of discontinuity and the coincidence set $E \subset [0, T) \times \mathbb{R}$, $E = \{(t, x) \in [0, T) \times \mathbb{R} : q(t, x) = o(x)\}$ is of the form

$$E = \{(t, x) \in [0, T) \times \mathbb{R} : \gamma_L(t) \le x \le \gamma_R(t)\},\$$

for two continuous piecewise smooth curves $\gamma_L \leq \gamma_R$ on [0, T] with γ_R non-decreasing. Suppose there exists a well-defined first collision time. We can set it without loss of generality to be t = 0 (this is a contradiction to Asm. 2 as we postulated $q_0 - o < 0$ on \mathbb{R} ; however, thanks to the semi-group property of any conservation law, we can shift time accordingly).

Then, the following result relying on the Rankine–Hugoniot condition [48, 49] gives a characterization of the boundary curves of the coincidence region E in terms of the traces of q and $\partial_x q$ from the outside of E:

Theorem 15. Assume that q_0 and o adhere to Asm. 2, o is strictly decreasing on some interval $(-\infty, x_0)$, and the limits for $t \in [0, T]$,

$$q^{L}(t) \coloneqq \lim_{y \nearrow \gamma_{L}(t)} q(t, y), \quad q^{R}(t) \coloneqq \lim_{y \searrow \gamma_{R}(t)} q(t, y), \quad q^{R}_{x}(t) \coloneqq \lim_{y \searrow \gamma_{R}(t)} \partial_{x} q(t, y),$$

exist for a.e. point along the curves γ_L, γ_R . Suppose further that $|\partial_x^2 q|$ is uniformly bounded on the (open) set E^c . Under the previous assumptions, we have the following:

1. The velocity field $(t,x) \mapsto V^*(t,x)$ as in Lem. 14 (and thus the speed of the characteristics) is given by

$$V^{\star}(t,x) = \begin{cases} 1, & (t,x) \notin E, \\ \frac{o(\gamma_R(t))}{o(x)}, & (t,x) \in E. \end{cases}$$
(29)

2. $\forall t \in [0,T), \ o(\gamma_L(t)) - q^L(t) > 0$ holds, and

$$\gamma_L'(t) = \frac{o(\gamma_R(t)) - q^L}{o(\gamma_L(t)) - q^L}.$$
(30)

3. In intervals where $t \mapsto \gamma_R(t)$ is differentiable, either $\gamma'_R(t) = 0$, or if q is smooth at $x = \gamma_R(t)$,

$$\gamma_{R}'(t) = \frac{q_{x}^{R}(t)}{q_{x}^{R}(t) - o'(\gamma_{R}(t))} > 1,$$
(31)

which, if q has a jump discontinuity at $x = \gamma_R(t)$, results in

$$\gamma_R'(t) = 1. \tag{32}$$

In particular, q has a discontinuity along γ_L . When $\gamma'_L(t) \leq 0$, then $\gamma'_L(t)$ is the speed of a backward congestion shock originated by the collision with the obstacle.

Proof. Consider a sufficiently small open ball B centered on a point $\gamma_L(t)$ where γ_L is smooth. Let $\Gamma_L = \{(t, y) \in [0, T] \times \mathbb{R} : (t, y) = (t, \gamma_L(t))\}$ and set $B = B_E \cup (\Gamma_L \cap B) \cup B_{E^c}$ with obvious notation (see Fig. 2).



Fig. 2 Illustration of the sets $\Gamma_L = \{(t, \gamma_L(t))\}, B_{E^c}$ and B_E . E is the set "above" the green and orange curves. E^c denotes is its complement.

Take a test function φ supported in *B*. Since *q* is a distributional solution of the conservation law $q_t + (V^*q)_x = 0$ in $(0,T) \times \mathbb{R}$, we have (here, for the sake of brevity, we omit the arguments of the functions)

$$0 = \int_{B} q\partial_{t}\varphi + V^{*}q\partial_{x}\varphi \ d(t,x)$$

=
$$\int_{B_{E}} q\varphi_{t} + V^{*}q\partial_{x}\varphi \ d(t,x) + \int_{B_{E^{c}}} q\varphi_{t} + V^{*}q\partial_{x}\varphi \ d(t,x)$$

as $q \equiv o$ on $E, V^* \equiv 1$ on E^c , and we obtain

$$= \int_{B_E} o\partial_t \varphi + V^* o\partial_x \varphi \,\mathrm{d}(t, x) + \int_{B_{E^c}} q\partial_t \varphi + q\partial_x \varphi \,\mathrm{d}(t, x). \tag{33}$$

Let us look at each of the two integrals in turn. First, note that in the coincidence set E, the obstacle o is a distributional solution of

$$\partial_t o(x) + \partial_x (V^*(t, x) o(x)) = 0 \ \forall x \in E.$$
(34)

Since o does not depend on t, we conclude that $(t, x) \to \partial_x(V^*(t, x)o(x))$ is the zero distribution on E. Therefore, owing to distribution theory, the function $((t, x) \mapsto V^*(t, x)o(x)) \in L^{\infty}(E)$ is constant in x for almost all t. Thus, $V^*(t, \cdot)o(\cdot) \equiv c_t$ holds a.e. on $(\gamma_L(t), \gamma_R(t))$ for a.e. t.

To identify c_t , we recall that $V_{\varepsilon}(o-q_{\varepsilon})$ is OSL continuous from below in the spatial variable (see Thm. 12 and apply the limit) uniformly in ε . Let us see that, actually,

$$c_t = o(\gamma_R(t)).$$

Let L > 0 be this Lipschitz constant. Writing $x_1 = \gamma_R(t)$, observe that for a.e. t for which $\gamma_{L,R}(t)$ are defined, and $\varepsilon > 0$:

$$V_{\varepsilon}\left(o\left(\underbrace{x_1+\frac{1}{k}}_{\notin E}\right)-q_{\varepsilon}\left(t,x_1+\frac{1}{k}\right)\right)-\frac{L}{k}\leq V_{\varepsilon}\left(o(x_1)-q_{\varepsilon}(t,x_1)\right)\leq 1.$$

As $\varepsilon \searrow 0$, we obtain that the left-hand side converges to $1 - \frac{L}{k}$ since $x_1 + \frac{1}{k} \notin E$ (see the definition of V_{ε}), whereas $V_{\varepsilon}(o(x_1) - q_{\varepsilon}(t, x_1)) \to V^*(t, x_1)$ as $\varepsilon \searrow 0$. So, in total,

$$1 - \frac{L}{k} \le V^*(t, x_1) \le 1$$

for all $k \in \mathbb{N}$, whence $V^*(t, x_1) = 1$ for a.e. $t \in [t_0, t_1]$. Thus, plug $x = x_1$ into $V^*(t, x)o(x) = c_t$ to obtain $c_t = o(x_1) = o(\gamma_R(t))$. This proves the representation Eq. (29).

Going back to Eq. (33), we deduce for the first integral, using the divergence theorem (let σ denote the 1D surface measure),

$$\int_{B_E} o\partial_t \varphi + V^* o\partial_x \varphi \, \mathrm{d}(t, x)$$

$$= \int_{B_E} o\partial_t \varphi + (o \circ \gamma_R) \partial_x \varphi \, \mathrm{d}(t, x)$$

$$\overset{\text{div-Thm.}}{=} \int_{B_E} \operatorname{div}_{t,x} (o\varphi, (o \circ \gamma_R)\varphi)^\top \, \mathrm{d}(t, x)$$

$$= \int_{\Gamma_L \cap B} (o, o \circ \gamma_R)^\top \cdot (-\gamma'_L, 1)^\top \varphi \, \mathrm{d}\sigma$$

$$= \int_{\Gamma_L \cap B} [-o\gamma_L + o \circ \gamma_R] \varphi \, \mathrm{d}\sigma.$$

For the second integral in Eq. (33), we find again with the divergence theorem and the fact that on E^c , q satisfy the transport equation,

$$\begin{split} &\int_{B_{E^c}} q\partial_t \varphi + q\partial_x \varphi \, \mathrm{d}(t,x) \\ &= -\int_{B_{E^c}} (\partial_t q + \partial_x q) \varphi \, \mathrm{d}(t,x) + \int_{\Gamma_L \cap B} (q,q) \cdot (\gamma'_L,-1) \varphi \, \mathrm{d}\sigma \\ &= 0 + \int_{\Gamma_L \cap B} (q^L \gamma'_L - q^L) \varphi \, \mathrm{d}\sigma. \end{split}$$

Therefore, the following holds:

$$\int_{\Gamma_L \cap B} (-o\gamma'_L + (o \circ \gamma_R)\varphi \,\mathrm{d}\sigma = -\int_{\Gamma_L} (q^L\gamma'_L - q^L)\varphi \,\mathrm{d}\sigma.$$

Since φ is arbitrary, we conclude that

$$-o(\gamma_L(t))\gamma'_L(t) + o(\gamma_R(t)) = -q^L\gamma'_L(t) + q^L$$
(35)

for a.e. $t \in (0,T)$ exploiting the given structure of Γ_L and the fact that the center of B is arbitrary.

Now, suppose that $o(\gamma_L(t)) = q^L$ and $\gamma_L(t)$ is strictly smaller than $\gamma_R(t)$. Then, from the strict monotonicity of o and Eq. (35), we have $o(\gamma_L(t)) < o(\gamma_R(t)) = q^L$, which is absurd. Therefore, $o(\gamma_L(t)) - q^L \neq 0$, and Eq. (30) follows by rearranging.

Eqs. (31) and (32) remain to be proven. Suppose that q has a jump discontinuity at $x = \gamma_R(t)$. Then, by assumption, there exists the trace q^R , and, necessarily, $q^R < o(\gamma_R(t))$. The proof of Eq. (32) is entirely analogous to the proof of Eq. (30), but instead of Eq. (35), we obtain

$$-o(\gamma_R(t))\gamma_R'(t) + o(\gamma_R(t)) = -q^R \gamma_R'(t) + q^R$$

$$\implies \gamma_R'(t)(o(\gamma_R(t)) - q^R) = o(\gamma_R(t)) - q^R,$$

and thus, $\gamma'_R(t) = 1$ since $q^R - o(\gamma_R(t)) \neq 0$.

Suppose now that q is continuous at $x = \gamma_R(t)$, q_x has a trace from the right, q_x^R , and $\gamma_R'(t) > 0$. Then, $\frac{d}{dt}o(\gamma_R(t)) < 0$, which implies that $o'(\gamma_R(t)) < 0$. Observe that necessarily, $q_x^R < o'(\gamma_R(t)) < 0$. The formal calculation to show Eq. (31) is as follows. Compute the derivative of o along the curve γ_R to find $\frac{d}{dt}o(\gamma_R(t)) = o'(\gamma_R(t))\gamma_R'(t)$. But, along γ_R , o = q holds; therefore, using Eq. (34) for q on E^c ,

$$\frac{\mathrm{d}}{\mathrm{d}t}o(\gamma_R(t)) = \frac{\mathrm{d}}{\mathrm{d}t}q(t,\gamma_R(t)) = \partial_t q(t,\gamma_R(t)) + \gamma'_R(t)\partial_x(t,\gamma_R(t))
= -\partial_2 q(t,\gamma_R(t)) + \gamma'_R(t)\partial_2 q(t,\gamma_R(t))
= -q_x^R + \gamma'_R(t)q_x^R,$$
(36)

giving Eq. (31). However, while $t \mapsto q(t, \gamma_R(t))$ is differentiable, q is not differentiable at $(t, \gamma_R(t))$, and so, the calculation is not justified. To circumvent this problem, a more careful computation is needed, which we now provide.

Let $\lambda > 0$ and consider the quantity

$$S_{t,\lambda} \coloneqq \frac{q(t,\gamma_R(t)) - q(t-\lambda,\gamma_R(t-\lambda))}{\lambda}$$

On the one hand, we have

$$\lim_{\lambda \searrow 0} S_{t,\lambda} = \frac{\mathrm{d}}{\mathrm{d}t} q(t, \gamma_R(t)).$$
(37)

On the other hand, we find

$$S_{t,\lambda} = \frac{q(t,\gamma_R(t)) - q(t-\lambda,\gamma_R(t))}{\lambda} + \frac{q(t-\lambda,\gamma_R(t)) - q(t-\lambda,\gamma_R(t-\lambda))}{\lambda}.$$
 (38)

The first term gives

$$\frac{q(t,\gamma_R(t)) - q(t-\lambda,\gamma_R(t))}{\lambda} = -\frac{1}{\lambda} \int_0^\lambda \frac{\mathrm{d}}{\mathrm{d}s} (q(t-s,\gamma_R(t))) \,\mathrm{d}s$$
$$= \frac{1}{\lambda} \int_0^\lambda \partial_t q(t-s,\gamma_R(t)) \,\mathrm{d}s$$
$$= -\frac{1}{\lambda} \int_0^\lambda \partial_2 q(t-s,\gamma_R(t)) \,\mathrm{d}s \to -q_x^R$$

as $\lambda \to 0$, from our smoothness assumptions on q from outside of E. Note that we can use the transport equation for q since $(t - s, \gamma_R(t))$ is *outside* the coincidence region E. For the second term in Eq. (38), we find

$$\begin{split} \frac{q(t-\lambda,\gamma_R(t)) - q(t-\lambda,\gamma_R(t-\lambda))}{\lambda} &= -\frac{1}{\lambda} \int_0^\lambda \frac{\mathrm{d}}{\mathrm{d}s} \big(q(t-\lambda,\gamma_R(t-s)) \big) \,\mathrm{d}s \\ &= \frac{1}{\lambda} \int_0^\lambda \partial_2 q(t-\lambda,\gamma_R(t-s)) \gamma_R'(t-s) \,\mathrm{d}s \\ &= \frac{1}{\lambda} \int_0^\lambda \partial_2 q(t-\lambda,\gamma_R(t)) \gamma_R'(t-s) \,\mathrm{d}s \\ &+ \frac{1}{\lambda} \int_0^\lambda [\partial_2 q(t-\lambda,\gamma_R(t-s)) - \partial_2 q(t-\lambda,\gamma_R(t))] \gamma_R'(t-s) \,\mathrm{d}s \\ &= \partial_2 q(t-\lambda,\gamma_R(t)) \frac{1}{\lambda} \int_0^\lambda \gamma_R'(t-s) \,\mathrm{d}s \end{split}$$

Now, clearly the first term converges to $q_x^R \gamma'_R(t)$ when $\lambda \to 0$, while the second can be easily bounded by $\lambda \|\partial_x^2 q\|_{L^{\infty}(E^c)} \|\gamma'_R\|_{L^{\infty}([0,T])}$, which also vanishes in the limit $\lambda \to 0$. Therefore, $\lim_{\lambda \to 0} S_{t,\lambda} = -q_x^R + \gamma'_R(t)q_x^R$, which together with Eq. (37) gives the

rigorous analogue of Eq. (36). Thus, we find that

$$o'(\gamma_R(t))\gamma'_R(t) = -q_x^R + \gamma'_R(t)q_x^R \iff \gamma'_R(t)(o'(\gamma_R(t)) - q_x^R) = -q_x^R.$$

Finally, recalling that $q_x^R < o'(\gamma_R(t)) < 0$, we obtain Eq. (31) by rearranging. This completes the proof.

6 Motivation of velocity for $\varepsilon \searrow 0$ by optimization

The result from Thm. 15 can also be *formally* motivated from an optimization perspective. We can, in fact, demonstrate that the velocity V^* , found in the previous chapter, is maximal in an L^1 sense for each time.

To this end, let us first assume that the density q moves with a space- and timedependent velocity $v: (0,T) \times \mathbb{R} \to [0,\infty)$ (the choice of this dependency becomes clear later) while obeying the scalar conservation law

$$\partial_t q(t,x) + \partial_x (v(t,x)q(t,x)) = 0, \quad q(0,x) = q_0(x) \text{ on } (0,T) \times \mathbb{R}.$$
 (39)

The conservation laws' characteristics emanating from $(t, x) \in (0, T) \times \mathbb{R}$, i.e., $\xi_v[t, x] : [0, T] \to \mathbb{R}$ are given as a solution of the ODE (see, e.g., [50, Sec. 2]):

$$\partial_s \xi_v[t, x](s) = v \left(s, \xi_v[t, x](s) \right), \quad \xi_v[t, x](t) = x, \ s \in [0, T]$$
(40)

We assume that v is OSL from below in the spatial variable and essentially bounded in $L^{\infty}((0,T) \times \mathbb{R})$. According to [50, 51], the solution to Eq. (39) satisfies

$$q(t,x) \coloneqq q(t_0,\xi_v[t,x](t_0))\partial_2\xi_v[t,x](t_0)$$

for all $(t, x) \in [0, T] \times \mathbb{R}$ and $t_0 \in [0, T]$. The previously mentioned solution formula makes the semi-group property (in time) of the dynamics visible.

We then define the L^2 -obstacle violation by introducing the function $V : [0, T] \to \mathbb{R}$ for all $t \in [0, T]$ and choose a $t_0 \in (0, T)$:

$$V(t) \coloneqq \int_{\mathbb{R}} \left(q(t_0, \xi_v[t, x](t_0)) \partial_2 \xi_v[t, x](t_0) - o'(x))^+ \right)^2 \, \mathrm{d}x \tag{41}$$

Performing a substitution according to Eq. (40), i.e., $y := \xi_v[t, x](t_0)$, we obtain

$$= \int_{\mathbb{R}} \left(\left(q(t_0, y) - o(\xi_v[t_0, y](t)) \partial_2 \xi_v[t_0, y](t) \right)^+ \right)^2 \, \mathrm{d}y,$$

with $(\cdot)^+ := \max\{\cdot, 0\}$. Thanks to the previously assumed regularity, we can compute the time-derivative and obtain the following by applying the chain rule for $t \in [0, T]$:

$$V'(t) = 2 \int_{\mathbb{R}} \left(q(t_0, y) - o(\xi_v[t_0, y](t)) \partial_2 \xi_v[t_0, y](t) \right)^+ \left(-o(\xi_v[t_0, y](t)) \partial_{23} \xi_v[t_0, y](t) \right) \\ - o'(\xi_v[t_0, y](t)) \partial_3 \xi_v[t_0, y](t) \partial_2 \xi_v[t_0, y](t) \right) \, \mathrm{d}y,$$

and substituting $\partial_3 \xi_v[t_0, y](t) = v(\xi_v[t_0, y](t))$ according to Eq. (40),

$$= -2 \int_{\mathbb{R}} \underbrace{\left(q(t_0, y) - o(\xi_v[t_0, y](t))\partial_2\xi_v[t_0, y](t)\right)^+}_{\geq 0} \\ \cdot \left(o(\xi_v[t_0, y](t))v'(\xi_v[t_0, y](t)) + \partial_y(o(\xi_v[t_0, y](t)))v(\xi_v[t_0, y](t))\right) \, \mathrm{d}y.$$
(42)

This yields, in particular (recalling again the properties of the characteristics in Eq. (40)),

$$V'(t_0) = -2 \int_{\mathbb{R}} \underbrace{\left(q(t_0, y) - o(y)\right)^+}_{\ge 0} \left(o(y)v'(y) + o'(y)v(y)\right) \, \mathrm{d}y. \tag{43}$$

On an abstract level, we have the function V measuring the violation of the obstacle. We also have (the initial datum respects the obstacle) that V(0) = 0. If we can manage to show that

$$V'(t) \le 0 \ \forall t \in [0,T]: \ V(t) > 0,$$
(44)

we know that the obstacle is never violated, i.e. $V \equiv 0$.

Thus, assume there exists $t \in (0, T]$ such that V(t) > 0. Then, there exists $E_>(t) \subset \mathbb{R}$ measurable with Lebesgue-measure greater zero so that

$$q(t, y) - o(y) > 0 \ \forall y \in E_{>0}(t).$$

Recalling Eq. (43), we have

$$V'(t) = -2 \int_{E_{>}(t)} \underbrace{\left(q(t_0, y) - o(y)\right)^{+}}_{>0} \left(o(y)v'(y) + o'(y)v(y)\right) \, \mathrm{d}y,$$

and so, we obtain by postulating Eq. (44) in a weak sense that

$$o(y)v'(y) + o'(y)v(y) \ge 0 \ \forall y \in E_{>}(t).$$

If we keep in mind that we want the velocity at time $t \in [0, T]$ to be maximal in some topology, we obtain the optimization problem (in L^1)

$$\max_{v \in L^{\infty}(\mathbb{R}; [0,1])} \int_{\mathbb{R}} v(x) - 1 \, \mathrm{d}x$$

subject to
$$\frac{\mathrm{d}}{\mathrm{d}y} \left(o(y)v(y) \right) \ge 0 \quad \forall y \in E_{>}(t),$$
(45)

where the inequality constraint is meant distributionally.

For sets $E_{>}(t)$ that are irregular (e.g., Cantor sets), the authors are unccertain on how to characterize solutions to the Eq. (45), but if this could be established, it may even present a more direct way to "solve the obstacle problem." However, assuming

that the set $E_{>}(t)$ is a union of disjoint intervals $(I_j)_{j \in J} \subset \mathbb{R}$ for an index set J, we can solve on each on these intervals Eq. (45).

Theorem 16 (The solution to Eq. (45)). For $t \in [0,T]$, the optimality system in Eq. (45) with the additional assumption that, for an index set J and a set of disjoint intervals $(I_j)_{j \in J}$,

$$E_{>}(t) = \bigcup_{j \in J} I_j, \quad I_j \cap I_k = \emptyset \qquad \forall j \in J, \ \forall k \in J \setminus \{j\}$$

admits a unique solution

$$v_t(x) = \begin{cases} \frac{o(\sup(I_j))}{o(x)}, & x \in I_j, j \in J\\ 1 & else, \end{cases} \quad \forall x \in \mathbb{R}.$$

This characterizes the velocity at time $t \in [0, T]$ so that one can see the solution to the obstacle as the solution of the conservation law

$$\partial_t q(t,x) + \partial_x (v_t(x)q(t,x)) = 0.$$

The choice of considering the solution on characteristics Eq. (41) is due to our requiring the expression to be time-sensitive concerning the obstacle violation and differentiable.

Remarkably, under the assumptions in Thm. 16, the characterization of the velocity coincides with the results in Sec. 5, namely, Eq. (29). This underlines the reasonability of the proposed approach, as in a regular enough setting, all presented approaches (regularization in V_{ε} in Sec. 4, characterization by Rankine–Hugoniot type approach in Sec. 5, and the optimization approach in this Sec. 6) coincide.

7 Visualization by means of a tailored Godunov scheme

In this section, we conduct numerical studies to "validate" the theoretical results about properties of the solution to Eq. (1), i.e.,

$$\partial_t q_{\varepsilon}(t,x) + \partial_x \left(V_{\varepsilon}(o_i(x) - q_{\varepsilon}(t,x)) q_{\varepsilon}(t,x) \right) = 0$$

on $(0,T) \times \mathbb{R}$ with initial condition $q(0,\cdot) = q_0^i$, $i \in \{1,2,3\}$. Here, *if not stated otherwise*, V_{ε} is as in Ex. 1, i.e., $V_{\varepsilon}(s) = 1 - \exp\left(\frac{s}{\varepsilon}\right)$, $s \in \mathbb{R}_{\geq 0}$, and $o_1, o_2, o_3 : \mathbb{R} \to \mathbb{R}$,

•
$$o_1(x) \coloneqq -\exp(-x^2) + \frac{3}{2}$$

• $o_2(x) \coloneqq -\frac{2}{3} \exp\left(-20\left(x + \frac{1}{2}\right)^2\right) - \exp\left(-20x^2\right) + \frac{3}{2}$ (46)
• $o_2(x) \coloneqq \min\left\{\max\left\{|x| - \frac{1}{2}\right\}^2\right\}$

• $o_3(x) \coloneqq \min\left\{\max\left\{|x|, \frac{1}{2}\right\}, \frac{3}{2}\right\}$



Fig. 3 Plots of o_1 and q_{ε} for $\varepsilon = 2^{-10}$ and initial datum $q_0^{(1)}$ specified in Eqs. (46) and (47). The chosen time points are t = 0, t = 0.25, t = 0.5, t = 0.75 (top, from left to right) and t = 1, t = 2, t = 3, t = 4 (bottom, from left to right).

for all $x \in \mathbb{R}$ and the corresponding initial data $q_0^1, q_0^2, q_0^3 : \mathbb{R} \to \mathbb{R}$

$$\bullet \ q_0^1(x) \coloneqq \chi_{[-1.5,-1]}(x) - x\chi_{(-1,0]} \bullet \ q_0^2(x) \coloneqq \max\{0, 1 - 3(x+1)^2\}^2 + \max\{0, 1 - 3(x-1)^2\}^2$$
(47)

$$\bullet \ q_0^3(x) \coloneqq \chi_{[-1.5,-1]}(x)$$

for all $x \in \mathbb{R}$. Of note, o_3 (due to nondifferentiability) and q_0^3 (due to lack of Lipschitz bounds) are *not* in accordance with Asm. 2.

To simulate Eq. (1), we employ a space-dependent Godunov method as laid out in [52, (34)] but use an adaptive step size such that a Courant-Friedrichs-Lévy-type condition (for non-space-dependent flux case, see [53, (13.11)]) holds in every time step.

For the spatial step size, we chose $\Delta x = 10^{-4}$. The adapted temporal step size then leads to several million time steps in every instance.

7.1 Visualization of q_{ε} for different scenarios

In Sec. 7.1, we let $\varepsilon = \frac{1}{1024}$ and observe the behavior of q_{ε} for different obstacles and initial data. Choosing o_1 as the obstacle and q_0^1 as the initial datum, we obtain the result in Fig. 3. Density accumulates "in front of" the obstacle, which can be interpreted as a "traffic jam". Also, density uses almost all the possible space under o_1 . In Fig. 4, we offer some more perspectives that also indicate very clearly that the obstacle is obeyed.

Choosing the data q_0^3 and o_3 , we obtain the results portrayed in Fig. 5.

In fact, the lack of regularity of this data does not prevent the solution from behaving in a physical/intuitive manner. However, at t = 2 and t = 3, some minor diffusion effects can be observed in the sense that the solution seems to "fade out" to its right. This is a numerical problem and would be far greater if one employs a Lax-Friedrichs scheme [52, p.16] instead of a Godunov scheme.

The obstacle in the case of o_2 and q_0^1 (see Fig. 6) is a little more delicate. Still, the solution behaves in the desired manner. Once the first stalactite of the obstacle is



Fig. 4 Plots of q_{ε} (color gradient) and o_1 (see (46)) from several perspectives. The initial datum is q_0^1 (see (47)). The thick red line surrounds the coincidence region $\{\lim_{\varepsilon \searrow 0} q_{\varepsilon} = o\}$.



Fig. 5 Plots of o_3 and q_{ε} for $\varepsilon = 2^{-10}$ and initial datum $q_0^{(3)}$ specified in Eqs. (46) and (47). Times are t = 0, t = 0.25, t = 0.5, t = 0.75 (top) and t = 1, t = 2, t = 3, t = 4 (bottom) from left to right



Fig. 6 Plots of o_2 and q_{ε} for $\varepsilon = 2^{-10}$ and initial datum $q_0^{(1)}$ specified in Eqs. (46) and (47). Times are t = 0, t = 0.25, t = 0.5, t = 0.75 (top) and t = 1, t = 2, t = 3, t = 4 (bottom) from left to right

passed, the density between the two stalactites only starts to increase once the second one is hit. After some time, all the "room" is used by the solution.

7.2 q_{ε} as $\varepsilon \searrow 0$

To observe the behavior of q_{ε} with ε becoming very small, we simulate Eq. (1) with initial datum q_0^1 and obstacle o_1 for several different ε , as shown in Fig. 7.



Fig. 7 Plots of o_1 (see Eq. (46)) and $q_{\varepsilon}(t, \cdot)$ for $\varepsilon \in \{2^{-6}, 2^{-7}, 2^{-8}, 2^{-9}, 2^{-10}\}$, where t = 0.9 in the top row and t = 1.8 in the bottom row. Here, the initial datum is $q_0^{(1)}$ (see Eq. (47)) and V_{ε} is as in Ex. 1 (left column), $V_{\varepsilon}(x) = \frac{1}{2} (\tanh\left(\frac{x}{\varepsilon}\right) + 1)$ (middle column) and $V_{\varepsilon}(x) = \min \{\max\{\frac{x}{\varepsilon}, 0\}, 1\}$ (right column) for all $x \in \mathbb{R}$.

We see that for smaller ε , the solution moves closer to the obstacle. This becomes especially evident around the point x = 0. This behavior can be motivated by the fact that $V_{\varepsilon} \nearrow 1$ as $\varepsilon \searrow 0$ on $[0, \infty)$ (recall Asm. 1). So, the smaller ε , the later q_{ε} slows down.

Thus, as suggested in Thm. 13 1), the convergence of q_{ε} as $\varepsilon \searrow 0$ can be identified numerically.

Moreover, if we choose a different regularization of V_{ε} , namely, $\mathbb{R} \ni x \mapsto \frac{1}{2} \left(\tanh\left(\frac{x}{\varepsilon}\right) + 1 \right)$ or $\mathbb{R} \ni x \mapsto \min \left\{ \max\left\{\frac{x}{\varepsilon}, 0\right\}, 1 \right\}$ (see Fig. 7), we see the same behavior. Note that the latter function (e.g., due to non-differentiability) does not meet the requirements in Asm. 1. In fact, the solutions for larger ε are closer to the solution for $\varepsilon = \frac{1}{1024}$ than in Fig. 7. This is due to the tanh-regularization being more precise.

7.3 Visualization of V_{ε}

Motivated by Thm. 15, we hypothesize that if $t \in [0, T]$ and there exists an interval I(t) such that $\lim_{\varepsilon \searrow 0} q_{\varepsilon} = o$ on $\{t\} \times I(t)$, $\lim_{\varepsilon \searrow 0} V_{\varepsilon}(o(x) - q_{\varepsilon}(t, x)) = \frac{o(b_t)}{o(x)}$ holds for all $x \in I(t)$, where $b_t := \sup I(t)$. As we can not compute $\lim_{\varepsilon \searrow 0} q_{\varepsilon}$, we want to analyze $V_{\varepsilon}(o - q_{\varepsilon})$ for the small value of $\varepsilon = \frac{1}{1024}$. In fact, at least numerically, our examples in Fig. 8 show the desired behavior.



Fig. 8 V^* according to Thm. 15 and $V_{\varepsilon}(o-q_{\varepsilon})$ with $\varepsilon = 2^{-10}$ for times t = 0.25, 1 with q_0^1 , o_1 (top from left to right), t = 0.5, 1 with q_0^2 , o_2 (middle from left to right), t = 0.5, 1 with q_0^3 , o_3 (bottom from left to right). See Eqs. (46) and (47) for the definition of the obstacles and initial data, respectively.

8 Open problems—future research

The most important question not answered in this work is whether the solution to Eq. (28) is unique and if it is independent of the regularization V_{ε} . Moreover, several generalizations are straightforward to some degree:

- The same approach works if we do not assume a constant velocity 1 if the obstacle is not present but rather a space- and time-dependent velocity satisfying required regularity for well-posedness.
- One can also consider nonlinear conservation laws [54], i.e.,

$$\partial_t q + \partial_x f(q) = 0$$

with the same scaling argument, resulting in

$$\partial_t q + \partial_x \big(V_{\varepsilon}(o-q)f(q) \big) = 0$$

and V_{ε} the regularization of the Heaviside function as in Asm. 1.

- The same argument can be made for nonlocal conservation laws [50].
- In the case of systems of conservation laws [54] and multi-d conservation laws, the presented approach may require some refinement. Particularly in the multi-d case, researchers may choose to steer toward the direction of the velocity field, which cannot be represented by a simple rescaling.

Data Availability

There is no associated data.

Acknowledgments

L. Pflug and J. Rodestock have been supported by the DFG – Project-ID 416229255 – SFB 1411.

References

- Berthelin, F., Degond, P., Delitala, M., Rascle, M.: A model for the formation and evolution of traffic jams. Archive for Rational Mechanics and Analysis 187(2), 185–220 (2007) https://doi.org/10.1007/s00205-007-0061-9
- [2] Andreianov, B., Donadello, C., Razafison, U., Rosini, M.D.: Qualitative behaviour and numerical approximation of solutions to conservation laws with non-local point constraints on the flux and modeling of crowd dynamics at the bottlenecks. ESAIM Math. Model. Numer. Anal. 50(5), 1269–1287 (2016) https://doi.org/10. 1051/m2an/2015078
- [3] Andreianov, B., Donadello, C., Rosini, M.D.: A second-order model for vehicular traffics with local point constraints on the flow. Math. Models Methods Appl. Sci. 26(4), 751–802 (2016) https://doi.org/10.1142/S0218202516500172
- [4] Chalons, C., Goatin, P., Seguin, N.: General constrained conservation laws. Application to pedestrian flow modeling. Netw. Heterog. Media 8(2), 433–463 (2013) https://doi.org/10.3934/nhm.2013.8.433
- Berthelin, F.: Existence and weak stability for a pressureless model with unilateral constraint. Math. Models Methods Appl. Sci. 12(2), 249–272 (2002) https://doi. org/10.1142/S0218202502001635
- [6] De Nitti, N., Serre, D., Zuazua, E.: Pointwise constraints for scalar conservation laws with positive wave velocity. cvgmt preprint (2024). http://cvgmt.sns. it/paper/6472/
- [7] Colombo, R.M., Goatin, P.: A well posed conservation law with a variable unilateral constraint. J. Differential Equations 234(2), 654–675 (2007) https: //doi.org/10.1016/j.jde.2006.10.014
- [8] Dymski, N.S., Goatin, P., Rosini, M.D.: Existence of BV solutions for a nonconservative constrained Aw-Rascle-Zhang model for vehicular traffic. J. Math. Anal. Appl. 467(1), 45–66 (2018) https://doi.org/10.1016/j.jmaa.2018.07.025
- [9] Garavello, M., Goatin, P.: The Aw-Rascle traffic model with locally constrained flow. J. Math. Anal. Appl. 378(2), 634–648 (2011) https://doi.org/10.1016/j. jmaa.2011.01.033
- [10] Garavello, M., Villa, S.: The Cauchy problem for the Aw-Rascle-Zhang traffic model with locally constrained flow. J. Hyperbolic Differ. Equ. 14(3), 393–414

(2017) https://doi.org/10.1142/S0219891617500138

- [11] Bayen, A., Friedrich, J., Keimer, A., Pflug, L., Veeravalli, T.: Modeling multilane traffic with moving obstacles by nonlocal balance laws. SIAM Journal on Applied Dynamical Systems 21(2), 1495–1538 (2022) https://doi.org/10.1137/ 20m1366654
- [12] Levi, L.: Obstacle problems for scalar conservation laws. M2AN Math. Model. Numer. Anal. 35(3), 575–593 (2001) https://doi.org/10.1051/m2an:2001127
- [13] Rodrigues, J.F.: On hyperbolic variational inequalities of first order and some applications. Monatsh. Math. 142(1-2), 157–177 (2004) https://doi.org/10.1007/ s00605-004-0238-3
- [14] Rodrigues, J.F.: On the hyperbolic obstacle problem of first order. vol. 23, pp. 253–266 (2002). https://doi.org/10.1142/S0252959902000249 . Dedicated to the memory of Jacques-Louis Lions. https://doi.org/10.1142/S0252959902000249
- [15] Goudon, T., Vasseur, A.: On a model for mixture flows: derivation, dissipation and stability properties. Arch. Ration. Mech. Anal. 220(1), 1–35 (2016) https: //doi.org/10.1007/s00205-015-0925-3
- [16] Gama, R.M., Pedrosa Filho, J.J., Martins-Costa, M.L.: Modeling the saturation process of flows through rigid porous media by the solution of a nonlinear hyperbolic system with one constrained unknown. ZAMM Z. Angew. Math. Mech. 92(11-12), 921–936 (2012) https://doi.org/10.1002/zamm.201100031
- [17] Berthelin, F., Bouchut, F.: Weak solutions for a hyperbolic system with unilateral constraint and mass loss. Ann. Inst. H. Poincaré C Anal. Non Linéaire 20(6), 975–997 (2003) https://doi.org/10.1016/S0294-1449(03)00012-X
- [18] Rossi, E., Weißen, J., Goatin, P., Göttlich, S.: Well-posedness of a non-local model for material flow on conveyor belts. ESAIM: Mathematical Modelling and Numerical Analysis 54(2), 679–704 (2020) https://doi.org/10.1051/m2an/ 2019062
- [19] Fernández-Real, X., Figalli, A.: On the obstacle problem for the 1d wave equation. Mathematics in Engineering 2(4), 584–597 (2020) https://doi.org/10.3934/mine. 2020026
- [20] Bellomo, N., Gibelli, L., Quaini, A., Reali, A.: Towards a mathematical theory of behavioral human crowds. Mathematical Models and Methods in Applied Sciences 32(02), 321–358 (2022)
- [21] Lions, J.-L., Stampacchia, G.: Variational inequalities. Comm. Pure Appl. Math. 20, 493–519 (1967) https://doi.org/10.1002/cpa.3160200302
 - 33

- [22] Kinderlehrer, D., Stampacchia, G.: An Introduction to Variational Inequalities and Their Applications. Pure and Applied Mathematics, vol. 88, p. 313. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London (1980)
- [23] Rodrigues, J.-F.: Obstacle Problems in Mathematical Physics. North-Holland Mathematics Studies, vol. 134, p. 352. North-Holland Publishing Co., Amsterdam (1987). Notas de Matemática, 114. [Mathematical Notes]
- [24] Mignot, F., Puel, J.-P.: Inéquations variationnelles et quasivariationnelles hyperboliques du premier ordre. J. Math. Pures Appl. (9) 55(3), 353–378 (1976)
- [25] Brézis, H.: Problèmes unilatéraux. J. Math. Pures Appl. (9) 51, 1–168 (1972)
- [26] Rudd, M., Schmitt, K.: Variational inequalities of elliptic and parabolic type. Taiwanese J. Math. 6(3), 287–322 (2002) https://doi.org/10.11650/twjm/ 1500558298
- [27] Korte, R., Kuusi, T., Siljander, J.: Obstacle problem for nonlinear parabolic equations. J. Differential Equations 246(9), 3668–3680 (2009) https://doi.org/ 10.1016/j.jde.2009.02.006
- [28] Rodrigues, J.F., Santos, L.: Quasivariational solutions for first order quasilinear equations with gradient constraint. Arch. Ration. Mech. Anal. 205(2), 493–514 (2012) https://doi.org/10.1007/s00205-012-0511-x
- [29] Chalub, F.A.C.C., Rodrigues, J.F.: A class of kinetic models for chemotaxis with threshold to prevent overcrowding. Port. Math. (N.S.) 63(2), 227–250 (2006)
- [30] Bögelein, V., Duzaar, F., Mingione, G.: Degenerate problems with irregular obstacles. J. Reine Angew. Math. 650, 107–160 (2011) https://doi.org/10.1515/ CRELLE.2011.006
- [31] Amorim, P., Neves, W., Rodrigues, J.F.: The obstacle-mass constraint problem for hyperbolic conservation laws. Solvability. Ann. Inst. H. Poincaré C Anal. Non Linéaire 34(1), 221–248 (2017) https://doi.org/10.1016/j.anihpc.2015.11.003
- [32] Kružkov, S.N.: First order quasilinear equations in several independent variables. Mathematics of the USSR-Sbornik 10(2), 217 (1970) https://doi.org/10.1070/ SM1970v010n02ABEH002156
- [33] Alt, H.W.: Linear Functional Analysis. Springer, London (2016). https://doi.org/ 10.1007/978-1-4471-7280-2 . http://dx.doi.org/10.1007/978-1-4471-7280-2
- [34] Evans, L.C., Gariepy, R.F.: Measure Theory and Fine Properties of Functions, Revised Edition. Chapman and Hall/CRC, Boca Raton (2015). https://doi.org/ 10.1201/b18333 . http://dx.doi.org/10.1201/b18333
- [35] Godlewski, E., Raviart, P.A.: Hyperbolic Systems of Conservation Laws.

Mathématiques & applications. Ellipses, Paris (1991). https://books.google.de/books?id=X3qyvAEACAAJ

- [36] Brezis, H.: Functional Analysis, Sobolev Spaces and Partial Differential Equations. Springer, New York (2010). https://doi.org/10.1007/ 978-0-387-70914-7. http://dx.doi.org/10.1007/978-0-387-70914-7
- [37] Simon, J.: Compact sets in the space $L^p(0,T;B)$. Annali di Matematica Pura ed Applicata **146**, 65–96 (1986)
- [38] Milgrom, P., Segal, I.: Envelope theorems for arbitrary choice sets. Econometrica 70(2), 583–601 (2002) https://doi.org/10.1111/1468-0262.00296
- [39] Nocedal, J., Wright, S.: Numerical Optimization. Springer, New York (2006). https://doi.org/10.1007/978-0-387-40065-5 . http://dx.doi.org/10.1007/ 978-0-387-40065-5
- [40] Teschl, G.: Ordinary Differential Equations and Dynamical Systems. American Mathematical Society, Providence (2012). https://doi.org/10.1090/gsm/140 . http://dx.doi.org/10.1090/gsm/140
- [41] Leoni, G.: A First Course in Sobolev Spaces. American Mathematical Society, Providence (2009). https://doi.org/10.1090/gsm/105 . http://dx.doi.org/10. 1090/gsm/105
- [42] BULÍČEK, M., GWIAZDA, P., MÁLEK, J., ŚWIERCZEWSKA-GWIAZDA, A.: On scalar hyperbolic conservation laws with a discontinuous flux. Mathematical Models and Methods in Applied Sciences 21(01), 89–113 (2011) https://doi.org/ 10.1142/s021820251100499x
- [43] Bulíček, M., Gwiazda, P., Świerczewska-Gwiazda, A.: On unified theory for scalar conservation laws with fluxes and sources discontinuous with respect to the unknown. Journal of Differential Equations 262(1), 313–364 (2017) https: //doi.org/10.1016/j.jde.2016.09.020
- [44] Carrillo, J.: Conservation laws with discontinuous flux functions and boundary condition. Journal of Evolution Equations 3(2), 283–301 (2003) https://doi.org/ 10.1007/s00028-003-0095-x
- [45] Martin, S., Vovelle, J.: Convergence of implicit finite volume methods for scalar conservation laws with discontinuous flux function. ESAIM: Mathematical Modelling and Numerical Analysis 42(5), 699–727 (2008) https://doi.org/10.1051/ m2an:2008023
- [46] Dias, J.-P., Figueira, M.: On the approximation of the solutions of the riemann problem for a discontinuous conservation law. Bulletin of the Brazilian Mathematical Society, New Series 36(1), 115–125 (2005) https://doi.org/10.1007/

s00574-005-0031-5

- [47] Dias, J.-P., Figueira, M.: On the riemann problem for some discontinuous systems of conservation laws describing phase transitions. Communications on Pure & Applied Analysis 3(1), 53–58 (2004) https://doi.org/10.3934/cpaa.2004.3.53
- [48] Rankine, W.J.M.: Xv. on the thermodynamic theory of waves of finite longitudinal disturbance. Philosophical Transactions of the Royal Society of London 160, 277– 288 (1870) https://doi.org/10.1098/rstl.1870.0015
- [49] Hugoniot, H.: Memoir on the propagation of movements in bodies, especially perfect gases (first part). J. de l'Ecole Polytechnique 57(3) (1887)
- [50] Keimer, A., Pflug, L.: Existence, uniqueness and regularity results on nonlocal balance laws. Journal of Differential Equations 263(7), 4023–4069 (2017) https: //doi.org/10.1016/j.jde.2017.05.015
- [51] Bouchut, F., James, F.: One-dimensional transport equations with discontinuous coefficients. Nonlinear Analysis: Theory, Methods & Applications 32(7), 891–933 (1998) https://doi.org/10.1016/s0362-546x(97)00536-1
- [52] Zhang, P., Liu, R.-X.: Hyperbolic conservation laws with space-dependent flux: I. characteristics theory and riemann problem. Journal of Computational and Applied Mathematics 156(1), 1–21 (2003) https://doi.org/10.1016/ s0377-0427(02)00880-4
- [53] LeVeque, R.J.: Numerical Methods for Conservation Laws. Birkhäuser, Basel (1992). https://doi.org/10.1007/978-3-0348-8629-1 . http://dx.doi.org/10.1007/ 978-3-0348-8629-1
- [54] Bressan, A.: Hyperbolic Systems of Conservation Laws: The One-Dimensional Cauchy Problem. Oxford University PressOxford, Oxford (2000). https:// doi.org/10.1093/oso/9780198507000.001.0001 . http://dx.doi.org/10.1093/oso/ 9780198507000.001.0001