METRIC LINES IN ENGEL-TYPE GROUPS AND THE NILPOTENT GROUP N_{6.3.1}.

ALEJANDRO BRAVO-DODDOLI

ABSTRACT. Given a sub-Riemannian manifold, which geodesics are "metric lines" (i.e. globally minimizing geodesics)? This article takes the first steps in answering this question for "arbitrary rank" and "nonintegrable" Carnot groups. We classify the metric lines of the Engeltype groups Eng(n) (Theorem B) and give a partial classification for the group of four-by-four nilpotent triangular matrices $N_{6,3,1}$ (Theorem C). The sub-Riamannian structure of the former group is defined on a non-integrable distribution of rank n + 1 and the geodesic flow of the latter group is not algebraically integrable.

1. INTRODUCTION

The Engel-type group and the group of four-by-four nilpotent triangular matrices admit the structure of Carnot groups of step 3, denoted by Eng(n) and $N_{6,3,1}$, respectively. Every Carnot group is a sub-Riemannian manifold, so are Eng(n) and $N_{6,3,1}$. This work provides a complete classification of metric lines Eng(n) and a partial result in $N_{6,3,1}$. To understand the concept of a metric line in the realm of sub-Riemannian geometry, let us provide a formal definition.

Definition 1. Let M be a sub-Riemannian manifold, $dist_M(\cdot, \cdot)$ be the sub-Riemannian distance on M, and $|\cdot| : \mathbb{R} \to [0, \infty)$ be the absolute value. We say that a curve $\gamma : \mathbb{R} \to M$ is a metric line if it is a globally minimizing geodesic, i.e.,

 $|a-b| = dist_M(\gamma(a), \gamma(b))$ for all compact interval $[a, b] \subset \mathbb{R}$.

For the formal definition of a sub-Riemannian geodesic, see [3, sub-sub-Chapter 4.7.2] or [28, sub-Chapter 1.4]. Alternative term for "metric line" is "globally minimizing geodesic", "isometric embedding of the real line" or "infinite-geodesic".

The first main contribution of this study establishes a condition of a sub-Riemannian geodesic in a Carnot group to qualify as a metric line.

Theorem A. Let \mathbb{G} be a Carnot group and $\gamma(t)$ be a sub-Riemannian geodesic in \mathbb{G} parametrized by arc length. A necessary condition for $\gamma(t)$ being

Key words and phrases. Carnot group, non-integrable functions, Global minimizing geodesic, sub-Riemannian geometry.

a metric line is that there exists a unitary vector v in g_1 such that

(1.1)
$$2 = \lim_{t \to \infty} \frac{1}{t} \int_{-t}^{t} \rho(v, (L_{\gamma^{-1}(s)})_* \dot{\gamma}(s)) ds$$

where $(L_{\gamma^{-1}(t)})_*$ is the pushforward of the left translation by $\gamma^{-1}(t)$.

We conjecture that the metric lines in Carnot groups are precisely geodesic satisfying a more robust condition than the one outlined in Theorem A.

Conjecture 2. Let \mathbb{G} be a Carnot group and $\gamma(t)$ be a sub-Riemannian geodesic in \mathbb{G} parametrized by arc length. Sub-Riemannian geodesics satisfying the following condition precisely constitute metric lines in \mathbb{G} ; there exists a unitary vector v in \mathfrak{g}_1 such that

(1.2)
$$v = \lim_{t \to -\infty} (L_{\gamma^{-1}(t)})_* \dot{\gamma}(t) = \lim_{t \to \infty} (L_{\gamma^{-1}(t)})_* \dot{\gamma}(t).$$

We say that a group \mathbb{G} is metabelian if $[\mathbb{G}, \mathbb{G}]$ is abelian. Every metabelian group possesses a maximal normal abelian sub-group \mathbb{A} containing $[\mathbb{G}, \mathbb{G}]$. In a prior work [17], we performed the symplectic reduction of sub-Riemannian geodesic flow on metabelian abelian nilpotent groups (every Carnot group is nilpotent), providing an explicit expression for the reduced Hamiltonian. Both groups, $\operatorname{Eng}(n)$ and $N_{6,3,1}$ serve as examples of metabelian Carnot groups, as indicated by equations (4.1) and (5.1). In this context, we leverage their reduced dynamics to classify sub-Riemannian geodesics and identify a family of candidates to qualify as metric lines.

1.1. The Engel-type group. The Engel-type group Eng(n) is (2n + 2)dimensional sub-Riemannian manifold featuring a rank (n+1) distribution. If n = 1, Eng(n) corresponds to the Engel group. As demonstrated in [17], the sub-Riemannian geodesic flow on Eng(n) is integrable for all natural number n. The sub-Riemannian geodesics on Eng(n) fall into distinct types: line, small oscillations, r-periodic, and r-homoclinic. Sub-sub-Section 4.1.2 provides the formal definitions for each type. Theorem B is the second primary results of this paper, and it offers a comprehensive classification of metric lines in Engel-type group.

Theorem B. The metric lines in Eng(n) are precisely geodesic of line and r-homoclinic types.

Corollary 40 demonstrates that sub-Riemannian geodesics of the type small oscillations and r-periodic do not qualify as metric lines. In addition, Lemma 36 shows that only abnormal geodesics being a metric lines are line geodesics.

A supplementary result concerning the sub-Riemannian geodesic in the Engel-type group is Proposition 44. This proposition states that the projection to \mathbb{R}^{n+1} of sub-Riemannian geodesics with zero angular momentum lies on a plane and satisfies the Euler-Elastica equation. In particular, the projection of a geodesic of the *r*-homoclinic type is the Euler-Solition, as illustrated in Figure 1.1 and 1.2.

METRIC LINES IN ENGEL-TYPE GROUPS AND THE NILPOTENT GROUP $N_{6,3,1}$. 3

1.2. The four-by-four Nilpotent Triangular Matrices Group. The four-by-four nilpotent triangular matrices group $N_{6,3,1}$ is a 6 dimensional sub-Riemannian Manifold featuring a rank three distribution. In [29], Montgomery, R. and Shapiro, M., and Stolin, A. proved that the sub-Riemannian geodesic flow in $N_{6,3,1}$ is not algebraically integrable. The sub-Riemannian geodesics on $N_{6,3,1}$ fall into distinct types: line, small oscillations, periodic, homoclinic, and generic. Theorems B is the third primary results of this paper and offers partial classification of metric lines in four-by-four nilpotent triangular matrices group.

Theorem C. The geodesic of type line and homoclinic are metric lines in $N_{6,3,1}$.

Lemma 57 shows the sub-Riemannian geodesic of the type small oscillations, periodic and unbounded generic do not qualify as metric lines. In addition, Lemma 55 shows that only abnormal geodesics being a metric lines, are line geodesics. The problem remains open for a particular type of bounded generic geodesic whose reduced dynamics approach arbitrarily close to an equilibrium point, refer to Corollary 12 and Lemma 57 for more details.

In [31], R. Perline and S. Tabachnikov say, "We have observed that the presence of Elastica in a geometrical context often indicates there is an integrable system floating around in the background". Proposition 58 offers additional evidence to this observation. Despite the non-algebraic integrability of the sub-Riemannian geodesic flow in N_{6,3,1}, certain regions in the cotangent bundle T^* N_{6,3,1} exhibit zero angular momentum, rendering the sub-Riemannian geodesic flow integrable on these regions. Moreover, Proposition 58 establishes a connection to the Euler-Elastica Problem: the projection to \mathbb{R}^3 of sub-Riemannian geodesics with zero angular momentum lies on a plane and satisfies the Euler-Elastica equation. In particular, the projection of a geodesic of the homoclinic type is the Euler-Solition, as depicted in 1.1 and 1.2.

1.3. History of the conjecture 2 and the Euler-Elastica problem. We summarize the preceding results that lead us to Conjecture 2.

- Metric lines in step two Carnot group precisely correspond to linegeodesics, consult Definition 5.
- In [8, 7, 9, 10], A. Andertov and Y. Sachkov employed optimal synthesis to prove the conjecture for the Heisenberg and the Engel groups.
- In [11], A. Ardentov and E. Hakavuori utilized optimal synthesis to demonstrate the conjecture for the Cartan group.

It's worth noting that the Heisenberg and Engel groups are isomorphic as a Carnot group to the 1-jet space and the 2-jet spaces, respectively.

4



FIGURE 1.1. The images show the space R^3 , with coordinate (x_1, x_2, θ_0) , and the projections by π of the homoclinic-geodesic c(t).

• In [16], we presented a partial outcome for the conjecture in the k-jet space and offered an alternative proof to the result obtained by A. Andertov and Y. Sachkov's result.

Theorems B and C represent generalizations of the result obtained by A. Andertov and Y. Sachkov result in the following manner: the metric line within the Engel group is such that its projection to \mathbb{R}^2 is the Euler-Soliton. To delve into the details of the Euler-Elastica problem, one can refer to [33, sub-Chapter 5.5] for an introduction from the perspective of Calculus of Variations, [3, sub-sub-chapter 7.8.3] for its connections with the rolling sphere, [23, sub-sub-Chapter 11.2.3] for its relationships with Dubis' problem, and consult [8] or [13] for its connections with sub-Riemannian geodesics in the Engel group.

It is noteworthy that the optimal synthesis method necessitates explicit integration of the sub-Riemannian geodesic flow. Consequently, the earlier classifications of metric lines in Carnot Groups were conducted under specific conditions, including the Carnot group being metabelian and of rank two, and the sub-Riemannian geodesic flow being integrable.

Theorems B and C demonstrate that the approach employed in this paper is more versatile than optimal synthesis. Our method involves defining an intermediate sub-Riemannian space \mathbb{R}_F^{n+1} and sub-Riemannian submersion $\pi_F : \mathbb{G} \to \mathbb{R}_F^{n+1}$, detailed in sub-Section 2.2. Proposition 4 establishes that the horizontal lift of a metric line is a metric line. As a result, we streamline the investigation of metric lines on a Carnot group \mathbb{G} by focusing on metric lines in \mathbb{R}_F^{n+1} . Finally, the forth main contribution of this study is Theorem D, offering sufficient conditions for a sub-Riemannian geodesic in \mathbb{R}_F^{n+1} to qualify as a metric line.



FIGURE 1.2. The images show two solutions to the Euler-Elastica problem in the (x, θ_0) -plane. The first panel presents a generic solution, and the second panel displays the Eulersoliton solution.

Outline. Section 2 presents the preliminary concepts and results necessary to prove Theorem B and C. Sub-Section 2.1 briefly describe the Carnot groups as a sub-Riemannian manifolds, introduce the metabelian definition and presents the algebraic assumptions used during the paper, consult equation (2.1). Under these assumptions, sub-Section 2.1 exposes the exponential coordinates of the second kind and the reduced Hamiltonian flow, the main consequence of the symplectic reduction is the **Background Theorem**, which establishes the correspondent between a pair (F, x(t)) and the sub-Riemannian geodesic in a metabelian Carnot group, where F(x) is a polynomial and x(t) is a solution to the reduced dynamics. Sub-Section 2.2 defines the intermediate sub-Riemannian Manifolds called magnetic space. In addition, sub-Section 2.2 defines the period map, the cost function, and some properties of the sub-Riemannian geodesics in the magnetic space. Sub-Section 2.3 presents and proves Theorem D, the method to verify that sub-Riemannian geodesics qualifies as a metric line.

The main goal of sub-Section 4 is to define the Engel group Eng(n) and prove Theorem B. Sub-section 4.1 presents the reduced Hamiltonian for the sub-Riemannian geodesic flow on Engenl type group Eng(n) and the classify the sub-Riemannian geodesics. Sub-Section 4.2 provide the essential properties of the geodesics whose reduced dynamics is given by radial anharmonic oscillator, one of these results is Proposition 44, which makes the relation between sub-Riemannian geodesics with zero angular momentum and the Euler-Elastica problem. Sub-Section 4.3 introduces the magnetic space used to prove Theorem C and some essential results. Sub-Section 4.4 proves Theorem B and has a similar structure than section 4.

Acknowledgment. I acknowledge my advisor for my Ph.D. degree, Richard Montgomery, who introduced me to sub-Riemannian geometry, Carnot groups, and Metric lines, and for his invaluable help during my time at UCSC and after. I acknowledge my mentor, Anthony Block, and the mathematics department of the University of Michigan, where this work was done, for supporting my application and made me part of the department.

Thank you to Enrico Le Donne, Nicola Padua, Gil Bor, Andrei Ardentov, and Yuri Sachkov. Enrico Le Donne for inviting me to the University of Fribourg and introducing me to the metabelian groups; Nicola Paddeu and Gil Bor for e-mail conversations regarding the course of this work; and Andrei Ardentov and Yuri Sachkov, whose works inspired me to create this paper.

2. Preliminary

We will present the fundamental concepts and results necessary to prove the main theorems of the paper.

2.1. Carnot Groups As Sub-Riemannian Manifold. A Carnot group \mathbb{G} is a simple connected Lie group with a graded, nilpotent Lie algebra \mathfrak{g} , and its first layer \mathfrak{g}_1 Lie-generates \mathfrak{g} . Every Carnot group \mathbb{G} has a canonical projection $\pi : \mathbb{G} \to \mathbb{R}^{d_1}$, where $\mathbb{G}/[\mathbb{G}, \mathbb{G}] \simeq \mathfrak{g}_1 \simeq \mathbb{R}^{d_1}$ and d_1 is the dimension of the layer \mathfrak{g}_1 . To endow \mathbb{G} with a left-invariant sub-Riemannian structure, we define the non-integrable distribution at a point g in \mathbb{G} by $\mathcal{D}_g := (L_g)_*\mathfrak{g}_1$. Additionally, we equip \mathcal{D} with an inner product by declaring the basis of left-invariant vector fields in \mathcal{D} to be orthonormal. Alternatively, we can achieve this by equipping \mathbb{R}^{d_1} with the Euclidean inner product, and then declaring the inner product on \mathcal{D} to be the pullback of the Euclidean product by π . In this manner, a curve $\gamma(t)$ on \mathbb{G} and tangent to \mathcal{D} has the same arc length that its projection $\pi(\gamma(t))$. Such a submersion possessing this property is termed a sub-Riemannian submersion. Let us formalize this definition.

Definition 3. Let (M, \mathcal{D}_M, g_M) and (N, \mathcal{D}_N, g_N) be two sub-Riemannian manifolds and $\phi : M \to N$ be a submersion. We say that ϕ is a sub-Riemannian submersion if $\phi_*\mathcal{D}_M = \mathcal{D}_N$ and $\phi^*g_N = g_M$.

Every sub-Riemannian submersion is accompanied by an inverse map at the curve level known as the horizontal lift, and it is unique up to translations. A classic result pertaining to metric lines and their horizontal lift is as follows.

Proposition 4. Let $\phi : M \to N$ be a sub-Riemannian submersion. If $\gamma(t)$ is the horizontal lift of a metric line in N, then $\gamma(t)$ is a metric line in M.

The proof of Proposition 4 is given in [15, p. 154]. In every Carnot group, there exists a family of geodesics called line-geodesic.

Definition 5. Let \mathbb{G} be a Carnot group. We say that a sub-Riemannian geodesic $\gamma(t)$ is a line-geodesic if $\gamma(t)$ is the lift of a line in \mathbb{R}^{d_1} .

An alternative term for "line-geodesic" is "one parameter sub-group". Proposition 4 and Definition 5 lead to the following corollary.

Corollary 6. Every geodesic line in a Carnot group is a metric line.

2.1.1. *Metabelian Carnot groups*. Let us revisit the definition of metabelian groups and discuss some of their properties.

Definition 7. We say that a group \mathbb{G} is metabelian if $[\mathbb{G}, \mathbb{G}]$ is abelian. Let \mathbb{G} be a metabelian Carnot group. We denote by \mathbb{A} the maximal normal abelian subgroup containing $[\mathbb{G}, \mathbb{G}]$, and \mathfrak{a} being the Lie algebra of \mathbb{A} .

See [32] for a more extensive explanation of the algebraic properties of metabelian groups.

Let us consider the left action of \mathbb{A} on \mathbb{G} . As this action is both free and proper, the quotient $\mathcal{H} := \mathbb{G}/\mathbb{A}$ is well defined. Consequently, \mathbb{G} is isomorphic as a manifold to $\mathbb{A} \times \mathcal{H}$. This is a pivotal aspect in the symplectic reduction of the cotangent bundle of \mathbb{G} .

Given the Lie algebra \mathfrak{a} , we can decompose the \mathfrak{g}_1 as the direct sum of two sub-spaces using the sub-Riemannian inner product. Let us introduce these spaces.

Definition 8. Let \mathbb{G} be a metabelian Carnot group, and let \mathfrak{a} be the Lie algebra of a maximal abelian subgroup \mathbb{A} containing $[\mathbb{G}, \mathbb{G}]$. We denote by \mathfrak{v} the intersection of \mathfrak{a} and \mathfrak{g}_1 , and we define \mathfrak{h} as the orthogonal complement of \mathfrak{v} with respect to the sub-Riemannian inner product, then $\mathfrak{g}_1 = \mathfrak{h} \oplus \mathfrak{v}$.

The condition $[\mathfrak{h}, \mathfrak{h}] = 0$ implies we can identify \mathcal{H} with a subgroup and view \mathbb{G} as the direct product $\mathbb{A} \rtimes \mathcal{H}$. For the remainder of this work, our focus will be on metabelian Carnot groups satisfy to have step 3 and the following conditions;

(2.1) $[\mathfrak{h},\mathfrak{h}] = 0$, $\dim \mathfrak{h} = n$, $\dim \mathfrak{v} = 1$, $\dim \mathfrak{g}_2 = n$, and $\dim \mathfrak{g}_3 = 1$.

The conditions from equation (2.1) imply the non-integrable distribution \mathcal{D} has rank n + 1 and $\dim \mathbb{A} = n + 2$. Both the Engel-type group $\operatorname{Eng}(n)$ and the group of four-by-four nilpotent matrices $N_{6,3,1}$ fulfill these conditions. Additionally, we will build a sub-Riemannian manifold \mathbb{R}_F^{n+2} and a sub-Riemannian submersion $\pi_F : \mathbb{G} \to \mathbb{R}_F^{n+2}$ utilizing these properties.

2.1.2. Exponential Coordinates Of Second Type. Given the above conditions from equation (2.1), we will define a basis for \mathfrak{g} : we denote by E_1, \ldots, E_n the basis for \mathfrak{h} and $E_0^{\mathfrak{a}}, \ldots, E_{n+1}^{\mathfrak{a}}$ the basis for \mathfrak{a} , where $E_0^{\mathfrak{a}}$ the basis for \mathfrak{v} . Therefore, $\{E_1, \ldots, E_n, E_0^{\mathfrak{a}}, \ldots, E_{n+1}^{\mathfrak{a}}\}$ constitutes a basis for \mathfrak{g} .

The assumption of \mathfrak{g} being nilpotent implies the exponential map is a global diffeomorphism. Leveraging this fact, we proceed to define exponential coordinates of second type.

Definition 9. We endows \mathbb{G} with a unique chart $(\mathbb{R}^{n+2} \times \mathcal{H}, \Phi)$. A pair (θ, x) in $\mathbb{R}^{n+2} \times \mathcal{H}$, where θ is in \mathbb{R}^{n+2} and x is in \mathcal{H} , defines a point g in \mathbb{G} through the map Φ given by

$$g = \Phi(\theta, x) = \prod_{k=0}^{n+1} \exp(\theta_k E_k^{\mathfrak{a}}) * \prod_{i=1}^n \exp(x_n E_n),$$

where * is multiplication on \mathbb{G} and exp : $\mathfrak{g} \to \mathbb{G}$ is the exponential map. We call the coordinates (θ, x) exponential coordinates of second type.

The establishment of exponential coordinates of the second type results in the subsequent proposition.

Lemma 10. Let \mathbb{G} be a metabelian Carnot group satisfying the conditions from equation (2.1), then the left-invariant vector fields generating the nonintegrable distribution \mathcal{D} are given by

$$X_i(g) := (L_g)_* E_i = \frac{\partial}{\partial x_i} \quad 1 \le i \le n,$$

$$Y_0(g) := (L_g)_* E_0^{\mathfrak{a}} = \frac{\partial}{\partial \theta_0} + \sum_{k=1}^n x_k \frac{\partial}{\partial \theta_k} + P(x) \frac{\partial}{\partial \theta_{n+1}},$$

where $P(x) : \mathcal{H} \to \mathbb{R}$ is a homogeneous polynomial of degree 2.

Lemma 10 is a particular result from a more general statement in the context of a general metabelian Carnot groups, consult [17, Lemma 3.2].

2.1.3. The Reduced Hamiltonian H_{μ} . We will briefly summarize the main ideas and results from [17], adding the conditions from equation (2.1). Let $T^*\mathbb{G}$ be the cotangent bundle of \mathbb{G} endowed with canonical symplectic structure, and let H_{sR} be the Hamiltonian function governing the sub-Riemannian geodesic flow, for the formal definition of the sub-Riemannian geodesic flow see [28, sub-Chapter] or [3, sub-sub-Chapter 4.7.2.]. We consider the Hamiltonian action \mathbb{A} on $T^*\mathbb{G}$, which fixes the Hamiltonian H_{sR} . This action gives rise to a momentum map $J: T^*\mathbb{G} \to \mathfrak{a}^*$. When we consider the standard coordinates $(p,g) = (p_{\theta}, p_x, \theta, x)$ on T^*G , the momentum map is expressed simply as $J(p,g) = p_{\theta}$. For a more detailed explanation of the momentum map, refer to [30] or [2, sub-Chapter 4.2].

Let μ be in \mathfrak{a}^* , and consider its inverse image $J^{-1}(\mu) \subset T^*\mathbb{G}$. Since there exists a symplectomorphism between $T^*\mathbb{G}$ and $T^*\mathbb{A} \times T^*\mathcal{H}$, it follows $J^{-1}(\mu)$ is isomorphic as a manifold to $\mathbb{A} \times T^*\mathcal{H}$. The abelian nature of \mathbb{A} implies the co-isotropic group \mathbb{A}_{μ} is equal to \mathbb{A} . Consequently, we conclude there exists a symplectomorphism between the reduced space $T^*\mathbb{G}//\mathbb{A}_{\mu}$ and $T^*\mathcal{H}$, where $T^*\mathcal{H}$ is endowed with the canonical symplectic structure. In addition, if $\Pi : T^*\mathbb{G} \to T^*\mathcal{H}$ is the projection defined by $\Pi(p,g) = (p_x, x)$, then Π is a symplectic map, and the reduced Hamiltonian H_{μ} is given by

(2.2)
$$H_{\mu}(p_x, x) = \frac{1}{2} \sum_{i=1}^{n} p_{x_i}^2 + \frac{1}{2} F_{\mu}^2(x),$$

where if $\mu = (a_0, \ldots, a_{n+1})$ then $F_{\mu}(x) = a_0 + \sum_{k=1}^n a_k x_k + a_{n+1} P(x)$. For a comprehensive overview of symplectic reduction, please refer to [26] or [30]. The Hamiltonian system is given by

(2.3)
$$\dot{x}_i = p_{x_i}, \text{ and } \dot{p}_{x_i} = -F_\mu(x)\frac{\partial F}{\partial x_i}$$

The condition $H_{\mu}(p_x, x) = \frac{1}{2}$ implies the reduced dynamics occur within a closed set called hill region, and the corresponding sub-Riemannian geodesic is parametrized by arc length.

Definition 11. Let H_{μ} be the reduced Hamiltonian given by equation (2.2). We say that a closed and connected set $\Omega_{\mu} \subset \mathcal{H}$ is a hill region, if $|F_{\mu}(x)| < 1$ for every x in the interior of Ω_{μ} and $|F_{\mu}(x)| = 1$ for every x in the boundary of Ω_{μ} .

Let us outline the procedure for constructing a sub-Riemannian geodesic for a pair $(F_{\mu}, x(t))$: Let us assume that x(t) is the *x*-component of a solution to the reduced dynamic system (2.2) with energy $H_{\mu} = \frac{1}{2}$. Having the solution $(p_x(t), x(t))$, we define a curve $\gamma(t)$ in \mathbb{G} by the following equation

(2.4)
$$\dot{\gamma}(t) = \sum_{i=1}^{n} \dot{x}_i(t) X_i(\gamma(t)) + F_\mu(x(t)) Y_0(\gamma(t)),$$

where we used $\dot{x}_i(t) = p_{x_i}(t)$ from Hamilton equation (2.3).

The **Background Theorem** establishes the correspondent between a solution to the reduced dynamics and the sub-Riemannian geodesic in G.

Background Theorem. Let \mathbb{G} be metabelian Carnot group satisfying the conditions from equation (2.1). The above procedure results in a geodesic in \mathbb{G} parametrized by arc length and momentum μ in \mathfrak{a}^* . Conversely, every arc length parameterized geodesic in \mathbb{G} with momentum μ in \mathfrak{a}^* can be obtained through this procedure using the Hamiltonian H_{μ} .

In [5, 6, 27], Anzaldo-Meneses and Monroy-Perez Monroy proved the **Background Theorem** within the framework of the jet space of real function on one variable, an example of a metabelian Carnot group. Their work served as inspiration for us to extend and generalize this result to metabelian nilpotent groups, for more details refer to [17].

2.1.4. Classification of Normal Geodesic in \mathbb{G} . We classify the sub-Riemannian geodesics according to their reduce dynamics. Let $\gamma(t)$ be a geodesic in \mathbb{G} corresponding to the pair $(F_{\mu}, x(t))$, then x(t) is the x-component of the solution $(p_x(t), x(t))$ of the reduced system (2.3). By the theory of n-degree of freedom system, $(p_x(t), x(t))$ is regular or singular.

(Regular geodesic) We say a geodesic $\gamma(t)$ is regular, if its reduced dynamics is regular. The reduce dynamics is regular if $(p_x(t), x(t))$ is not asymptotic to a equilibrium point.

(Singular geodesic) We say a geodesic $\gamma(t)$ is singular, if its reduced dynamics is singular. The reduce dynamics is singular if $(p_x(t), x(t))$ is asymptotic to a equilibrium point

More specific, a normal geodesic $\gamma(t)$ is only one of the following type:

(Line geodesic) We say a geodesic $\gamma(t)$ is a line, if the projected curve $\pi(\gamma(t))$ is a line in \mathbb{R}^{n+1} . Line geodesic correspond to constant polynomial (uniform rectilinear motion) or trivial solution of the reduced dynamics (equilibrium points).

(Bounded geodesic) We say a geodesic $\gamma(t)$ is regular bounded, if its reduced dynamics is regular and bounded. The reduce dynamics is unbounded if there exists a compact set $K \subset \mathcal{H}$ such that x(t) is in K for all t.

(Unbounded geodesic) We say a geodesic $\gamma(t)$ is regular unbounded, if its reduced dynamics is regular and unbounded. The reduce dynamics is unbounded if there does not exist a compact set $K \subset \mathcal{H}$ such that x(t) is in K for all t.

(Homoclinic geodesic) We say a geodesic $\gamma(t)$ is homoclinic if its reduced dynamics is a homoclinic orbit. The reduced dynamics has a homoclinic orbit if and only if $\lim_{t\to\pm\infty}(p_x(t), x(t)) = (0, x_0)$.

(Heteroclinic geodesic) We say a geodesic $\gamma(t)$ is heteroclinic if its reduced dynamics is a heteroclinic orbit. The reduced dynamics has a heteroclinic orbit if and only if the limits $\lim_{t\to\infty}(p_x(t), x(t)) = (0, x_1)$ and $\lim_{t\to-\infty}(p_x(t), x(t)) = (0, x_2)$ and $x_1 \neq x_2$.

(Turn-back geodesic) We say a heteroclinic geodesic $\gamma(t)$ is turn-back if $F_{\mu}(x_1)F_{\mu}(x_2) = -1$.

(Direct-type geodesic) We say a heteroclinic geodesic $\gamma(t)$ is direct-type if $F_{\mu}(x_1)F_{\mu}(x_2) = 1$.

Theorem A implies that turn-back geodesics do not qualify as a metric lines.

Corollary 12. The following geodesics do not qualify as metric lines.

- Turn back geodesics.
- Bounded generic geodesic whose reduced dynamics lay on a regular region of the potential $F_{\mu}(x)$.

Corollary 12 is a particular result for a more general statement from [17, Corollary 1.5]. A consequence of Theorem A is Lemma 34, which provides an alternative proof for the turn-back case.

2.2. The (n+2)-Dimensional Magnetic Space. In [15], we introduced the 3-dimensional sub-Riemannian manifold, denoted by \mathbb{R}_F^3 and called "magnetic space", whose geometry depends on the choice of a polynomial F(x). Now, we aim to generalize this concept by defining a sub-Riemannian manifold $\mathbb{R}_F^{n+2} = \mathcal{H} \times \mathbb{R}^2$ as follows: we consider the global coordinates (x, y, z), then the Pfaffian equation dz - F(x)dy = 0 defines a non-integrable distribution \mathcal{D}_F , and the sub-Riemannian metric on \mathcal{D}_F is $ds_{\mathbb{R}_F^{n+2}}^2 = (\sum_{i=1}^n dx_i^2 + dy^2)|_{\mathcal{D}_F}$. We provided a sub-Riemannian submersion π_F factoring the sub-Riemannian submersion $\pi : \mathbb{G} \to \mathbb{R}^{n+1}$, that is, $\pi = pr \circ \pi_F$, where the target of π_F is \mathbb{R}_F^{n+2} and the target of pr is \mathbb{R}^{n+1} . If $F(x) = a_0 + \sum_{k=1}^n a_k x_k + a_{n+1} P(x)$, then the projections π_F and pr are given in coordinates by

(2.5)
$$\pi_F(x,\theta) = (x,\theta_0, \sum_{\ell=0}^{n+1} a_\ell \theta_\ell) = (x,y,z), \text{ and } pr(x,y,z) := (x,y).$$

It follows that π_F maps the frame $\{X_1, \ldots, X_n, Y_0\}$, defined in Lemma 10, into the frame $\{\tilde{X}_1, \ldots, \tilde{X}_n, \tilde{Y}\}$, that is,

$$\tilde{X}_i := \frac{\partial}{\partial x_i} = (\pi_F)_* X_i \text{ for all } i = 1, \dots, n,$$
$$\tilde{Y} := \frac{\partial}{\partial y} + F(x) \frac{\partial}{\partial z} = (\pi_F)_* Y_0.$$

We conclude that $\{\tilde{X}_1, \ldots, \tilde{X}_n, \tilde{Y}\}$ is a global framed for \mathcal{D}_F . An explanation of the name "magnetic space" is given in [15, sub-Section 4.1].

For future work we may be interesting in work with metabelian Carnot with same algebraic condition from equation (2.1) but arbitrary step s, then we will state Theorem A in this more general context. The following proposition tells every metabelian Carnot group with the above conditions has a sub-Riemannian submersion π_F to some magnetic space \mathbb{R}_F^{n+2} .

Proposition 13. Let \mathbb{G} be a metabelian Carnot group with the algebraic condition $[\mathfrak{h}, \mathfrak{h}] = 0$, dim $\mathfrak{v} = 1$ and step s, then there exist a magnetic space \mathbb{R}_F^{n+2} and a sub-Riemannian submersion $\pi_F : \mathbb{G} \to \mathbb{R}_F^{n+2}$ for a polynomial F generating geodesic in the spirit of the **Background Theorem**.

Proof. Using coordinates of second type and building the sub-Riemannian submersion in the same way that equation (2.5).

2.2.1. Geodesics In The Magnetic Space. The Hamiltonian function governing the sub-Riemannian geodesic flow in \mathbb{R}^{n+2}_{F} is

(2.6)
$$H_F(p_x, p_y, p_z, x, y, z) = \frac{1}{2} \sum_{i=1}^n p_{x_i}^2 + \frac{1}{2} (p_y + F(x)p_z)^2.$$

We say a curve c(t) = (x(t), y(t), z(t)) is a \mathbb{R}_F^{n+1} -geodesic parametrized by arc length in \mathbb{R}_F^{n+1} , if it is the projection of the sub-Riemannian geodesic flow with the condition $H_F = \frac{1}{2}$. Since H_F does not depend on the coordinates y and z, then the momentum p_y and p_z are constant of motion, and the translation $\varphi_{(y_0,z_0)}(x, y, z) = (x, y + y_0, z + z_0)$ is an isometry.

Definition 14. Let us denote by $dist_{\mathbb{R}_{F}^{n+2}}(,)$ the sub-Riemannian distance and by $Iso(\mathbb{R}_{F}^{n+2})$ the isometry group of \mathbb{R}_{F}^{n+2} .

For a more comprehensive understanding of the sub-Riemannian distance definition and the sub-Riemannian group of isometries, consult [28, Chapter 1.4] or [3, sub-Chapter 3.2].

Setting $p_y = a$ and $p_z = b$ prompted the formulation of the following definition.

Definition 15. We say that the two-dimensional linear space Pen_F is the pencil of F(x), if $Pen_F := \{G(x) = a + bF(x) : (a, b) \in \mathbb{R}^2\}$.

We define the lift of a curve in \mathbb{R}^{n+2}_F to a curve in \mathbb{G} .

Definition 16. Let c(t) be a curve in \mathbb{R}_F^{n+2} . We say that a curve $\gamma(t)$ in \mathbb{G} is the lift of c(t) = (x(t), y(t), z(t)) if $\gamma(t)$ solves

$$\dot{\gamma}(t) = \dot{x}(t)X(\gamma(t)) + G(x(t))Y_0(\gamma(t)).$$

Now, we delineate the \mathbb{R}_{F}^{n+2} -geodesics, their lifts, and their connection with the sub-Riemannian geodesics in G.

Proposition 17. Let c(t) be a \mathbb{R}^{n+2}_F -geodesic for G(x) in Pen_F , then the component x(t) satisfies the one-degree of freedom given by the Hamiltonian function:

(2.7)
$$H_G(p_x, x) := \frac{1}{2}p_x^2 + \frac{1}{2}(a + bF(x))^2 = \frac{1}{2}p_x^2 + \frac{1}{2}G^2(x).$$

And the coordinates y(t) and z(t) satisfy

 $\dot{y} = G(x(t))$ and $\dot{z} = G(x(t))F(x(t))$. (2.8)

Moreover, every \mathbb{R}_F^{n+2} -geodesic is the π_F -projection of a geodesic in \mathbb{G} corresponding to G(x) in Pen_F. Conversely, the lifts of a \mathbb{R}_{F}^{n+2} -geodesic are precisely those geodesics corresponding to polynomials in Pen_F .

The proof is the same as the proposition presented in [15, sub-Section 4.1]in the context of the jet space and the magnetic space \mathbb{R}^3_F .

Corollary 18. Let $\gamma(t)$ be a sub-Riemannian geodesic in \mathbb{G} corresponding to the polynomial F(x) and let c(t) be the curve given by $\pi_F(\gamma(t))$, then c(t)is a \mathbb{R}^{n+2}_{F} -geodesic corresponding to the pencil (a,b) = (0,1).

We classify the sub-Riemannian geodesics in \mathbb{R}^{n+2}_F according to their reduce dynamics defined by the reduced Hamiltonian H_G , equation (2.7), in the same way as we did in sub-sub-Section 2.1.4.

In sub-Riemannian geometry, there are two types of geodesics: normal and abnormal. The sub-Riemannian geodesic flow governs the normal geodesics, while the abnormal geodesics are determined by the endpoint map. Lemma 19 characterizes the abnormal geodesics in \mathbb{R}^{n+2}_{F} . For a more in-depth understanding of the endpoint map and abnormal geodesics, refer to [28, Chapter 3], [3, sub-sub-Chapter 4.3.2], or [18].

Lemma 19. The space \mathbb{R}_F^{n+2} possesses two families of abnormal curves; (Family of vertical lines) We say a curve c(t) is a abnormal vertical line curve, if c(t) is tangent to the vector field \tilde{Y} and $x(t) = x^*$ is a constant point in \mathcal{H} such that $dF|_{x=x^*} = 0$. So c(t) qualify as a metric line.

(Family of horizontal isocontour) We say a curve c(t) is abnormal horizontal isocontour, if c(t) is tangent to space \mathcal{H} and F(x(t)) = const. c(t) qualify as a metric line if and only if c(t) is a line geodesic.

Here we used the expressions horizontal and vertical for curves c(t) in \mathbb{R}_{F}^{n+2} tangent to the sub-bundles $(\pi_{F} \circ L_{q})_{*}\mathfrak{h}$ and $(\pi_{F} \circ L_{g})_{*}\mathfrak{v}$, respectively. The proof of Lemma 19 is in the Appendix A.

METRIC LINES IN ENGEL-TYPE GROUPS AND THE NILPOTENT GROUP $N_{6,3,1}$.13

2.2.2. Cost Map In Magnetic Space. In [15, sub-Section 7.2], we defined the Cost map and utilized it to establish a family of metric lines in the jest space. Here, we introduce Cost, a crucial tool for formulating the statement of Theorem D.

Definition 20. Let (c, \mathcal{T}) be a pair of a \mathbb{R}^3_F -geodesic c(t) parametrized by arc-length, and a time interval $\mathcal{T} := [t_0, t_1]$. For every pair (c, \mathcal{T}) , we denote by $\Delta(c, \mathcal{T})$ the change suffer by the time t, and the coordinates y, and z after the geodesic c(t) travel during the time interval \mathcal{T} , then the function $\Delta : (c, \mathcal{T}) \to [0, \infty) \times \mathbb{R}^2$ is given by

(2.9)
$$\Delta(c, \mathcal{T}) := (\Delta t(c, \mathcal{T}), \Delta y(c, \mathcal{T}), \Delta z(c, \mathcal{T})) \\ := (t_1 - t_0, y(t_1) - y(t_0), z(t_1) - z(t_0))$$

For every pair (c, \mathcal{T}) , we define the function $Cost : (c, \mathcal{T}) \to [0, \infty) \times \mathbb{R}$ by

(2.10)
$$Cost(c, \mathcal{T}) := (Cost_t(c, \mathcal{T}), Cost_y(c, \mathcal{T}))$$

where

$$Cost_t(c, \mathcal{T}) = \Delta t(c, \mathcal{T}) - \Delta y(c, \mathcal{T})$$
$$Cost_y(c, \mathcal{T}) = \Delta y(c, \mathcal{T}) - \Delta z(c, \mathcal{T}).$$

We call $Cost(c, \mathcal{T})$ the cost function of c(t).

We interpret $Cost_t(c, \mathcal{T})$ as the cost that takes to the geodesic c(t) travel in the direction of the *y*-component in the positive direction. To provide further context to this interpretation, we present the following lemma.

Lemma 21. Let c(t) and $\tilde{c}(t)$ be two \mathbb{R}_{F}^{n+2} -geodesics. Let us assume that they travel from a point A to a point B in the time intervals \mathcal{T} and $\tilde{\mathcal{T}}$, respectively. If $Cost_{t}(c,\mathcal{T}) < Cost_{t}(\tilde{c},\tilde{\mathcal{T}})$, then the arc length of c(t) is shorter that the arc length of $\tilde{c}(t)$.

Lemma 21 extends an equivalent result in the context of magnetic space \mathbb{R}^3_F , and the proof remains the same; refer to [16, Lemma 21].

If c(t) is a geodesic for G(x), we can rewrite the maps $\Delta(c, \mathcal{T})$ and $Cost(c, \mathcal{T})$ in terms of G(x).

Definition 22. Let c(t) be a \mathbb{R}_F^{n+2} -geodesic and $\mathcal{T} = [t_0, t_1]$ be a time interval. val. We denote by $s(t) : \mathcal{T} \to \mathbb{R}$ the arc length of the curve x(t) in \mathcal{H} , that is,

$$s(t) = \int_{t_0}^t \sqrt{p_{x_1}^2(t) + \dots + p_{x_n}^2(t)} dt = \int_{t_0}^t \sqrt{1 - G^2(x(t))} dt.$$

Because s(t) is a monotone-increasing function, we can consider the inverse function t(s) and parametrize x in terms of s. We use this fact to express the map $\Delta(c, \mathcal{T})$ and $Cost(c, \mathcal{T})$ in terms of G(x).

Proposition 23. Let (c, \mathcal{T}) be a pair where c(t) is a \mathbb{R}^{n+2}_F -geodesic for G(x) in Pen_F , and \mathcal{T} is a time interval, let x be parametrized by s, then

we can rewrite the map $\Delta(c, \mathcal{T})$ from definition 20 in terms of G(x) and the arc-length $s(\mathcal{T}) := S$ as follows:

$$\Delta(c,\mathcal{T}) = (\int_{\mathcal{S}} \frac{ds}{\sqrt{1 - G^2(x(s))}}, \int_{\mathcal{S}} \frac{G(x(s))ds}{\sqrt{1 - G^2(x(s))}}, \int_{\mathcal{S}} \frac{G(x(s))F(x(s))ds}{\sqrt{1 - G^2(x(s))}})$$

In the same way, we can rewrite the map Cost(c, T) from definition 20 as follows:

$$Cost(c, \mathcal{T}) = \left(\int_{\mathcal{S}} \frac{1 - G(x)}{\sqrt{1 - G^2(x)}} ds, \int_{\mathcal{S}} \frac{G(x)(1 - F(x))}{\sqrt{1 - G^2(x)}} ds\right).$$

Proof. Set the change of variable t = t(s), then $ds = \sqrt{1 - G^2(x(t))}dt$, this last expression yield the formula for $\Delta t(c, \mathcal{T})$. For $\Delta y(c, \mathcal{T})$ and $\Delta z(c, \mathcal{T})$, set up the same change of variable and uses the expression $\dot{y} = G(x)$ and $\dot{z} = G(x)F(x)$. For $Cos_t(c, \mathcal{T})$ and $Cos_y(c, \mathcal{T})$ consider the difference of the previous expressions.

Corollary 24. Let c(t) be a \mathbb{R}_{F}^{n+2} -geodesic and let \mathcal{T} be a time interval then $Cost_{t}(c, \mathcal{T}) = 0$ if and only if G(x(t)) = 1 for all t in \mathcal{T} .

A isometry φ and a time translation $t \to t + t^*$ map metric lines in metric lines, then to classify metric lines it is enough to study the following class of geodesics.

Definition 25. We say that two \mathbb{R}_F^{n+2} -geodesics $c_1(t)$ and $c_2(t)$ are related if and only if there exists an isometry $\varphi(x, y, z)$ and a time translation t^* such that $c_1(t) = \varphi(c_2(t+t^*))$. We denote by [c] the class of geodesics induced by the equivalence relationship.

We notice that if [c] is a class of geodesics, then all the representatives are \mathbb{R}_{F}^{n+2} -geodesics corresponding to the same polynomial G(x) in Pen_{F} . Moreover, the equivalence relationship on the space of \mathbb{R}_{F}^{n+2} -geodesics induce a class in the space of solution x(t) of the reduced system, then every class [c] correspond to a unique pair (G, [x]). So, if c(t) is a homoclinic geodesic, then all the representatives of [c] are homoclinic geodesic with the same homoclinic orbit, same statement holds for heteroclinic geodesics.

Definition 26. Let x_0 and x_1 be two point in \mathcal{H} , then:

(1) We denote by $Homc(x_0)$ the set of all the classes of equivalence [c], whose reduced dynamics has a homoclinic orbit asymptotically to x_0 . $Homc(x_0)$ is the union of two disjoint sets, namely

$$Homc^{\pm}(x_0) := \{ (G, [x]) : G(x_0) = \pm 1 \}.$$

If the reduced Hamiltonian H_G has a unique equilibrium point, we omitted x_0 in the notation.

(2) We denote by $Hetc(x_0, x_1)$ the set of all the classes of equivalence [c], whose reduced dynamics has a heteroclinic orbit asymptotically to x_0 and x_1 ,

when $t \to \pm \infty$ respectively. $Hetc(x_0, x_1)$ is the union of two disjoint sets, namely

$$Hetc^{\pm}(x_0, x_1) := \{ (G, x_{\infty}) : G(x_0) = \pm 1 \text{ and } G(x_1) = \pm 1 \}.$$

To determine a class [c] is enough to provide a polynomial G(x) and an initial condition $(p_x(0), x(0))$. In the case when x(0) is in the boundary of the Hill region of G(x), then $p_x(0) = 0$ and it is enough specify just x(0). In sub-Sections 4.4 and 5.3, we will use this fact to characterize the space $Homc^{\pm}(x_0)$.

Proposition 23 yield the following corollary.

Lemma 27. Let c(t) be a \mathbb{R}_F^{n+2} -geodesic and let \mathcal{T}_n be a time interval such that $\lim_{n\to\infty} \mathcal{T}_n = [-\infty, \infty]$, then $\lim_{n\to\infty} Cos_t(c, \mathcal{T}_n)$ is finite if and only if [c] is in $Homc^+(x_0)$ or $Hetc^+(x_0, x_1)$, for some points x_0 and x_1 in \mathcal{H} .

In addition: $\lim_{n\to\infty} Cos_t(c, \mathcal{T}_n)$ does not depend of the representative, [x] has finite arc-length. And, if $\gamma(t)$ is the horizontal lift of c(t), then $\gamma(t)$ is geodesic in \mathbb{G} that satisfies the condition from Theorem A.

Definition 28. For every class [c] in $Homc^+(x_0)$ or $Hetc^+(x_0, x_1)$, we define the normalized period map $\Theta[c]$ by

$$\Theta[c] := (\Theta_1[c], \Theta_2[c]) = \lim_{n \to \infty} Cost(c, \mathcal{T}_n)$$

(2.11)
$$= \left(\int_{\mathcal{S}_{\infty}} \frac{1 - G(x(s))}{\sqrt{1 - G^2(x(s))}} ds, \int_{\mathcal{S}_{\infty}} \frac{G(x(s))(1 - F(x(s)))}{\sqrt{1 - G^2(x(s))}} ds\right),$$

where if $\lim_{n\to\infty} \mathcal{T}_n = [-\infty,\infty]$, then $\mathcal{S}_{\infty} := \lim_{n\to\infty} s(\mathcal{T}_n)$.

2.3. Metric lines in the Magnetic Space. This sub-section presents Theorem D. To make the statement more general we will relax, only in this sub-section, the hypothesis from equation (2.1) in the same way that the statement of Proposition 13. That is we will work in a magnetic space \mathbb{R}_F^{n+2} , where F(x) is a polynomial of degree s - 1 and π_F is a projection from a metabelian Carnot group with the conditions $[\mathfrak{h}, \mathfrak{h}] = 0$, dim $\mathfrak{v} = 1$ and step s. In this way the Theorem D will be helpful for future works. Before presenting the statement of this work, let us introduce some essential definitions.

Notation 29. Let us introduce some classical topology concepts:

- (1) Let K be a subset of \mathbb{R} . We denote by ∂K the boundary of K.
- (2) Let $|| \cdot ||_{\infty}$ be the maximum norm in \mathbb{R}^2 . We say that a sequence in \mathbb{R}^2 is uniformly bounded if there exist a positive constant C such that the maximum norm of the sequence is bounded by C for all n.

The first definition gives a condition for a sequence of geodesics to be bounded in \mathcal{H} .

Definition 30. Let $B_{\mathcal{H}}(r, x_0)$ be the closed ball on \mathcal{H} with radius r and center at x_0 , then we define the following set;

$$Com(B_{\mathcal{H}}(r, x_0)) := \{ (c, \mathcal{T}) : c(t) \text{ is a } \mathbb{R}_F^{n+2} - geodesic, \ \mathcal{T} \text{ is a compact} \\ time \text{ interval, and } x(t) \in B_{\mathcal{H}} \text{ for all } t \in \partial \mathcal{T} \}.$$

We say that region $B_{\mathcal{H}}(r, x_0) \times \mathbb{R}^2$ is geodesically compact, if for every sequence (c_n, \mathcal{T}_n) in $Com(B_{\mathcal{H}}(r))$ holding the following conditions;

(1)
$$\lim_{n\to\infty} \mathcal{T}_n = [-\infty,\infty],$$

(2) $Cost(c_n, \mathcal{T}_n)$ is uniformly bounded.

Then, there exist a compact subset $K_{\mathcal{H}}$ of \mathcal{H} such that $x_n(\mathcal{T}_n) \subset K_{\mathcal{H}}$ for all n.

The second definition helps to associate a polynomial $G_n(x)$ to a sequence of geodesics $c_n(t)$.

Definition 31. We say a sequence of geodesics $c_n(t)$ is strictly normal if $c_n(t)$ is a normal geodesic for all n and every convergent sub-sequence converges to a normal geodesic.

Theorem D. Let F(x) be a polynomial of degree s - 1 from \mathcal{H} to \mathbb{R} . Let $c_h(t)$ be geodesic in \mathbb{R}_F^{n+2} such that [c] is in $Homc^+(x_0)$. If the following conditions hold:

- (1) There exist a ball $B_{\mathcal{H}}(r, x_0)$ such that the region $B_{\mathcal{H}}(r, x_0) \times \mathbb{R}^2$ is geodesically compact.
- (2) The map Θ_2 : Homc⁺(x_0) $\rightarrow \mathbb{R}$ is one to one.
- (3) The map $Cost(G_h, \mathcal{T})$ is bounded for compact interval \mathcal{T} .
- (4) The sequence of minimizing geodesics $c_n(t)$ joining the points c(-n)and c(n) is strictly normal.

Then $c_h(t)$ is a metric line in \mathbb{R}^{n+2}_F .

The same statement is true if we replace $Homc^+(x_0)$ by $Hetc^+(x_0, x_1)$).

The proof is in appendix B and generalize the one given in [16, Theorem 33 and 43] in the context of \mathbb{R}^3_F .

3. Proof of Theorem A

Before proving Theorem A, let us introduce the Carnot dilation and the blowdown of a curve. They are two essential definitions for the proof of the theorem.

Carnot groups have the property of admitting dilatations. The dilatation is a one-parameter group of automorphism of \mathbb{G} , denote by $\delta_u : \mathbb{G} \to \mathbb{G}$ and with u in $\mathbb{R} \setminus \{0\}$. The dilatation is compatible with the metric, that is $dist_{\mathbb{G}}(\delta_u g_1, \delta_u g_2) = |u| dis_{\mathbb{G}}(g_1, g_2)$. If $u \neq 0$ and $\gamma(t)$ is a sub-Riemannian geodesic parametrized by arc length, so is $\gamma_u(t)$, where

$$\gamma_u(t) := \delta_{\frac{1}{u}} \gamma(ut).$$

METRIC LINES IN ENGEL-TYPE GROUPS AND THE NILPOTENT GROUP $N_{6,3,1.17}$

E. Hakavuouri and E. Le Donne developed a method to prove that a geodesic is not a metric line is the following. The idea relies on the concept of a blowdown.

Definition 32. Let \mathbb{G} be a Carnot group, and $\gamma(t)$ be a rectifiable curve in \mathbb{G} . We say $\bar{\gamma}(t)$ is the blowdown of $\gamma(t)$ if $\bar{\gamma}(t) = \lim_{u_n} \gamma_{u_n}(t)$ where u_n is any sequence of scales tending to infinity with n, and the limit being uniform on compact intervals.

E. Hakavuouri ad E. Le Donne proved the following proposition.

Proposition 33. Let \mathbb{G} be a Carnot group. Let $\gamma : \mathbb{R} \to \mathbb{G}$ be a metric line. Then there exists a sequence u_n , with $\lim_{n\to\infty} u_n = \infty$, for which the blowdown $\bar{\gamma}$ converges uniformly on compact sets to a line geodesic parametrized by arc length.

The proof is in [22, Corollary 1.6].

3.0.1. Proof of Theorem A. We will prove that if $\gamma(t)$ is a metric line, then equation 1.2 holds.

Proof. Let us assume $\gamma(t)$ is a metric line, then Proposition 33 implies there exists a sequence u_n such that $\bar{\gamma}(t) = g \exp(tv)$ for some unitary vector v in \mathfrak{g}_1 . If $\mathcal{T} = [-1, 1]$ and $\mathcal{T}_n := [-u_n, u_n]$, then, we have

$$\int_{\mathcal{T}} \rho(v, (L_{\gamma_{u_n}^{-1}(t)})_* \dot{\gamma}_{u_n}(t)) dt = \int_{\mathcal{T}} \rho(v, (L_{(\delta_{\frac{1}{u_n}}\gamma(u_n t))^{-1}})_* \frac{d}{dt} (\delta_{\frac{1}{u_n}}\gamma(u_n t))) dt$$
$$= \frac{1}{u_n} \int_{\mathcal{T}_n} \rho(v, (L_{\gamma^{-1}(\tilde{t})})_* (\dot{\gamma}(\tilde{t}))) d\tilde{t}$$

The first equality follows by the definition of $\gamma_{u_n}(t)$ and the second by the change of variable $u_n t = \tilde{t}$. Taking the limit $n \to \infty$ and using the uniformly convergence of $\bar{\gamma}$ on the compact set K, we have

$$2 = \int_{\mathcal{T}} \rho(v, v) dt = \lim_{n \to \infty} \frac{1}{u_n} \int_{\mathcal{T}_n} \rho(v, (L_{\gamma^{-1}(\tilde{t})})_*(\dot{\gamma}(\tilde{t}))) d\tilde{t}.$$

The following lemma shows that the condition from Conjecture 2 implies the condition from Theorem A.

Lemma 34. Let \mathbb{G} be a Carnot group, then

- Conjecture 2 implies the condition from Theorem A.
- If $\gamma(t)$ is a geodesic in \mathbb{G} such that the following limits $\lim_{t\to\infty} \dot{\gamma}(t)$ and $\lim_{t\to-\infty} \dot{\gamma}(t)$ exist but differ, then $\gamma(t)$ does not qualify as a metric line. Therefore, the turn-back geodesics do not qualify as a metric lines.

Proof. The L'Hopital rule yields the following result: let f(t) be a continuous function, if $\lim_{t\to\infty} f(t) = c$ then $\lim_{t\to\infty} \frac{1}{t} \int_0^t f(s) ds = c$.

The following lemma tells that dilatation of a metric lines is a metric line.

Lemma 35. If $\gamma(t)$ is a metric line in a Carnot group \mathbb{G} , then $\gamma_u(t)$ is a metric line in \mathbb{G} .

The proof of Lemma 35 is in [16, Lemma 7].

4. The Engel-type group $\operatorname{Eng}(n)$

Let $\operatorname{Eng}(n)$ be the Engel-type group, whose Lie algebra $\operatorname{eng}(n)$ is spanned by (2n+2) vectors $E_1, \ldots, E_n, E_0^{\mathfrak{a}}, \ldots, E_{n+1}^{\mathfrak{a}}$ satisfying the following non trivial relations

(4.1)
$$E_i^{\mathfrak{a}} := [E_i, E_0^{\mathfrak{a}}] \text{ and } E_{n+1}^{\mathfrak{a}} := [E_i, E_i^{\mathfrak{a}}]$$

Equation (4.1) shows $\operatorname{Eng}(n)$ is a metabelian Carnot group, the first layer \mathfrak{g}_1 and the Lie algebra \mathfrak{a} are spanned by $\{E_1, \ldots, E_n, E_0^{\mathfrak{a}}\}$ and $\{E_0^{\mathfrak{a}}, \ldots, E_{n+1}^{\mathfrak{a}}\}$, respectively. In addition, its growth vector is (n+1, 2n+1, 2n+2). To endow $\operatorname{Eng}(n)$ with a left-invariant sub-Riemannian metric, we declare the vector from equation (4.1) orthonormal, then \mathfrak{v} has dimension 1 and \mathfrak{h} is spanned by E_1, \ldots, E_n , so equation (4.1) implies $\operatorname{Eng}(n)$ satisfy the conditions from equation (2.1) and we can consider the sub-Riemannian submersion π_F given in sub-Section 2.2.1. In [20], E. Le Donne and T. Moisala studied the sub-Riemannian structure of the Engel-type group. Moreover, in [21], E. Le Donne and F. Tripaldi classified the Carnot groups in low dimension; in this context, they denoted the group $\operatorname{Eng}(2)$ by $N_{6.3.1a}$.

Let $g = (\theta, x)$ be the exponential coordinates of second defined in subsub-Section 2.1.2, where $x = (x_1, \ldots, x_n)$ and $\theta = (\theta_0, \theta_1, \ldots, \theta_{n+1})$, then the following left-invariant vector fields define a frame for the non-integrable distribution \mathcal{D} :

$$X_i = \frac{\partial}{\partial x_i}$$
 for $i = 1, ..., n$, and $Y_0 = \frac{\partial}{\partial \theta_0} + \sum_{i=1}^n x_i \frac{\partial}{\partial \theta_i} + \frac{x_i^2}{2} \frac{\partial}{\partial \theta_{n+1}}$.

Then, the sub-Riemannian metric on \mathcal{D} in the coordinates (θ, x) is given by the expression $ds_{\text{Eng}(n)}^2 = \sum_{i=1}^n dx_i^2 + d\theta_0^2$.

The following lemma characterize the abnormal curves in the Engel-type group.

Lemma 36. The Engel-type group Eng(n) possesses two families of abnormal curves:

(Family of vertical lines) We say a curve $\gamma(t)$ is an abnormal vertical line, if $\gamma(t)$ is tangent to the vector field Y_0 . So $\gamma(t)$ qualify as a metric line.

(Family of horizontal curves) We say a curve $\gamma(t)$ is an abnormal horizontal curves, if $\gamma(t)$ is tangent to sub-space $\mathcal{H} \subset \operatorname{Eng}(n)$. And in addition $\dot{\gamma}(t)$ is perpendicular to a constant vector or tangent to a sphere on \mathcal{H} . So $\gamma(t)$ qualify as a metric line if and only if $\gamma(t)$ is a horizontal line.

The use of the expressions horizontal and vertical is the same as in Lemma 19. The proof of Lemma 36 is in the Appendix A.

4.1. Reduced Hamiltonian. If $\mu = (a_0, \ldots, a_n)$ is in \mathfrak{a}^* , then equation (2.2) implies that the reduced Hamiltonian H_{μ} is of the form

(4.2)
$$H_{\mu}(p_x, x) = \frac{1}{2} \sum_{i=1}^{n} p_{x_i}^2 + \frac{1}{2} (a_0 + \sum_{i=1}^{n} a_i x_i + \frac{a_{n+1}}{2} \sum_{i=1}^{n} x_i^2)^2.$$

When $a_{n+1} = 0$, the reduced Hamiltonian H_{μ} takes on the structure resembling that of a Hamiltonian for small oscillations. Further elaboration on the theory of small oscillations consult [12, Section 23] or [24, Section 23]. In instances where $a_{n+1} \neq 0$, to find a more suitable expression for the reduce Hamiltonian, we establish the following change of coordinates;

$$(\tilde{x}_1, \dots, \tilde{x}_n) = (\frac{a_1}{a_{n+1}} + x_1, \dots, \frac{a_n}{a_{n+1}} + x_n),$$

and define the constants $(\alpha, \beta) := (a_0 - \frac{1}{2a_2} \sum_{i=1}^n a_i^2, \frac{a_{n+1}}{2})$. Thus, the reduced Hamiltonian is the radial an-harmonic oscillator

(4.3)
$$H_{\mu}(p_{\tilde{x}},\tilde{x}) = \frac{1}{2} \sum_{i=1}^{n} p_{\tilde{x}_{i}}^{2} + \frac{1}{2} F_{\mu}^{2}(r), \text{ where } F_{\mu}(r) = \alpha + \beta r^{2},$$

and if $|| \cdot ||_{\mathcal{H}}$ is the Euclidean norm in \mathcal{H} , then $r = ||x||_{\mathcal{H}}$.

If $(x(t), p_x(t))$ is the solution of an *n*-degree of freedom system with radial potential, then it is well known that curve x(t) lays on a two-plane. Let us formalize this idea in the following proposition.

Lemma 37. Let $T^*\mathbb{R}^2_{T^*\mathcal{H}}$ be the 4-dimensional sub-manifold of $T^*\mathcal{H}$ given by

$$T^* \mathbb{R}^2_{T^* \mathcal{H}} := \{ (p_x, x) \in T^* \mathcal{H} : (p_x, x) = (p_{x_1}, p_{x_2}, 0, \dots, 0, x_1, x_2, 0, \dots, 0) \}.$$

If $(p_x(t), x(t))$ is a solution to the an-harmonic oscillator, given by (4.3), with the initial condition $(p_x(0), x(0))$ in $T^*\mathbb{R}^2_{\mathcal{H}}$, then $(p_x(t), x(t))$ is in $T^*\mathbb{R}^2_{T^*\mathcal{H}}$ for all time.

Moreover, let $\Psi_Q(p_x, x)$ be the cotangent lift action of SO(n) on $T^*\mathcal{H}$, then every solution $(p_x(t), x(t))$ to the radial an-harmonic oscillator, given by (4.3), has the form $(p_x(t), x(t)) = \Psi_Q(\tilde{p}_x(t), \tilde{x}(t))$ where $(\tilde{p}_x(t), \tilde{x}(t))$ is a solution laying in $T^*\mathbb{R}^2_{T^*\mathcal{H}}$.

Proof. Let us first prove that if $(p_x(0), x(0))$ in $T^*\mathbb{R}^2_{\mathcal{H}}$, then $(p_x(t), x(t))$ is in $T^*\mathbb{R}^2_{T^*\mathcal{H}}$ for all time: By construction the Hamiltonian functions is invariant under the action of SO(n) on $T^*\mathcal{H}$, then SO(n) defines a momentum map $J: T^*\mathcal{H} \to \mathfrak{so}^*(n)$. Using the Killing form, we can identify the $\mathfrak{so}^*(n)$ with $\mathfrak{so}(n)$, then $J: T^*\mathcal{H} \to \mathfrak{so}(n)$ and a standard computation shows that $J(p_x, x) = p_x \wedge x$. Therefore, on one side, the skew-symmetric matrix associated with the two-form $p_x \wedge x$ has rank 2; on the second side, the conservation of the momentum map implies $p_x(0) \wedge x(0) = p_x(t) \wedge x(t)$. So the $p_x(t) \neq x(t)$ generates the same plan for all time t.

Since the reduced Hamiltonian function H_{μ} is SO(n)-invariant, the action maps solutions of the Hamiltonian system into solutions, see [19, Section 6],

then $(p_x(t), x(t)) = \Psi_Q(\tilde{p}_x(t), \tilde{x}(t))$ is a solution for the reduced Hamiltonian system, where $(\tilde{p}_x(t), \tilde{x}(t))$ is a solution laying in $T^* \mathbb{R}^2_{T^*\mathcal{H}}$.

Proposition 37 implies it is enough to understand the planar an-harmonic oscillator to describe and classify the dynamics of the reduced Hamiltonian H_{μ} .

4.1.1. The Planar Radial An-Harmonic Oscillator. Let us restrict the reduced Hamiltonian given by equation (4.3) to the sub-manifold $T^*\mathbb{R}^2_{\mathcal{H}}$ and write it in polar coordinates,

(4.4)
$$H_{\mu}(p_x, x)|_{T^*\mathbb{R}^2_{T^*\mathcal{H}}} = H_{\mu}(p_r, p_{\theta}, r, \theta) = \frac{1}{2}(p_r^2 + \frac{p_{\theta}^2}{r^2}) + \frac{1}{2}F_{\mu}^2(r).$$

Being the potential radial implies the coordinate θ is cyclic, then p_{θ} is a constant of motion. Let us fix $p_{\theta} = \ell$, we will denote the effective potential by $\frac{1}{2}V_{ef}(r)$, where $V_{ef}(r) := \frac{\ell^2}{r^2} + F_{\mu}^2(r)$. We reduce Hamiltonian H_{μ} to a one-degree of freedom system

(4.5)
$$H_{(\mu,\ell)}(p_r,r) = \frac{1}{2}p_r^2 + \frac{1}{2}V_{ef}(r).$$

Fixing the energy level $H_{(\mu,\ell)} = \frac{1}{2}$ and using Hamilton equation $p_r = \dot{r}$, we reduce to quadrature the dynamics for the radial coordinate,

(4.6)
$$t - t_0 = \int_{r_0}^r \frac{dr}{\sqrt{1 - V_{ef}(r)}}. \text{ where } r_0 = r(t_0).$$

We notice that the effective potential is non-negative and unbounded, then the conservation of energy $H_{(\mu,\ell)}(p_x(t), x(t)) = \frac{1}{2}$ implies the hill region is compact, then the radial coordinates lay on a closed interval called the radial Hill interval. Let us formalize the definition.

Definition 38. We say an interval $R = [r_{min}, r_{max}]$ is the radial hill interval of the effective potential V_{ef} if $V_{ef}(r_{max}) = V_{ef}(r_{max}) = 1$ and $V_{ef}(r) < 1$ for all r in (r_{min}, r_{max}) .

The following result characterizes the equilibrial points of the planar anharmonic oscillator.

Corollary 39. The reduce Hamiltonian system (4.5) has an equilibrium if and only if $\ell = 0$ and $F_{\mu}(r) = \pm(1 - \beta r^2)$. In addition, this equilibrium point is at $(p_r, r) = (0, 0)$ and has homoclinic orbits.

The potential polynomial $F_{\mu}(r) = \pm (1 - \beta r^2)$ provided by Corollary 39 is the generalization of the polynomial $F_{(\mu)}(x) = \pm (1 - \beta x^2)$ associated to the metric lines in the Engel group, see [16, Theorem B].

4.1.2. Classification of normal sub-Riemannian geodesics in the Engel-type group Eng(n). Let $\gamma(t)$ be a normal sub-Riemannian geodesic different of line type and with momentum μ , then $\gamma(t)$ is only one of the following options:

Small oscillation We say a geodesic $\gamma(t)$ is of the type small oscillation, if $a_{n+1} = 0$.

r-periodic We say a geodesic $\gamma(t)$ is of the type *r*-periodic, if $a_{n+1} \neq 0$ and the radial dynamic is periodic.

r-homoclinic We say a geodesic $\gamma(t)$ is of the type *r*-homoclinic, if $a_{n+1} \neq 0$ and the radial dynamic is a homoclinic orbit.

Corollary 40. The sub-Riemannian geodesic in Eng(n) of the type small oscillations and r-periodic are not metric lines.

Corollary 40 follows by Theorem A.

Proof. By construction the r-periodic geodesics are bounded in a regular region of the potential $F_{\mu}(r)$, then Lemma 12 tells they do not qualify as metric line.

Let $\gamma(t)$ be a geodesic of the oscillatory type, we will show that $\gamma(t)$ does not satisfies the condition form Theorem A. First, we notice that reduced Hamiltonian has a quadratic potential, then we can use the theory of small oscillation. Let *B* be a *n* by *n* symmetric matrix generating the quadratic potential, by construction *B* has rank one. So *B* has (n-1)-eigenvectors with eigenvalue 0 and one eigenvector with positive eigenvalue ω^2 , then after a change of variables the reduced Hamiltonian H_{μ} is the sum of two Hamiltonian functions, namely a (n-1)-degree of freedom system for a free particle $H_{free}(p_{x_1}, \ldots, p_{x_{n-1}})$ and one degree of freedom system for harmonic oscillator $H_{osc}(p_{x_n}, x_n)$ given by

$$H_{free} = \frac{1}{2}(p_{x_1}^2 + \dots + p_{x_{n-1}}^2)$$
 and $H_{osc} = \frac{1}{2}(p_{x_n}^2 + \omega^2 x_n^2).$

If $H_{osc} = 0$, the corresponding geodesic is a line-geodesic. Therefore, we will focus in the case $H_{osc} \neq 0$. Without lose of generality let us assume $\gamma(0) = 0$, then $\dot{\gamma}(t)$ is given by

$$\dot{\gamma}(t) = (p_{x_1}X_1 + \dots + p_{x_{n-1}}X_{n-1} + \sqrt{2H_{osc}}\cos(\omega t)X_n + \frac{\sqrt{2H_{osc}}}{\omega}\sin(\omega t)Y_0)|_{\gamma(t)},$$

we remark $1 = ||\dot{\gamma}|| = 2H_{free} + 2H_{osc}$. Let $v = (v_1, \dots, v_n, v_{n+1})$ be an unitary arbitrary vector in $\mathfrak{eng}(n)$, then

$$\begin{aligned} \lim_{t \to \infty} \frac{1}{t} \int_{-t}^{t} \rho(v, (L_{\gamma^{-1}(s)})_* \dot{\gamma}(s)) ds &|= 2|p_{x_1} v_1 + \dots + p_{x_{n-1}} v_{n-1}| \\ &\leq 4H_{free} < 2. \end{aligned}$$

Therefore, $\gamma(t)$ does not satisfies the condition form Theorem A.

4.2. Radial geodesics.

Proposition 41. The Lie algebra $\mathfrak{eng}(n)$ is invariant under the action of SO(n) given by

$$\tilde{E}_{j} = \sum_{i=1}^{n} Q_{ji} E_{i}, \quad \tilde{E}_{0}^{\mathfrak{a}} = E_{0}^{\mathfrak{a}}, \quad \tilde{E}_{j}^{\mathfrak{a}} = \sum_{i=1}^{n} Q_{ji} E_{i}^{\mathfrak{a}}, \quad \tilde{E}_{n+1}^{\mathfrak{a}} = E_{n+1}^{\mathfrak{a}},$$

where $Q := (Q_{ij})$ is in SO(n). Moreover, the action on $\mathfrak{eng}(n)$ induces an isometric action ψ_Q on $\operatorname{Eng}(n)$. If (θ, x) are the exponential coordinates of the second type, then the action $\psi_Q(\theta, x) = (\tilde{\theta}, \tilde{x})$ given by

$$\tilde{\theta_0} = \theta_0, \ \tilde{\theta}_j = \sum_{i=1}^n Q_{ji}\theta_i, \ \tilde{\theta}_{n+1} = \theta_{n+1}, \ \tilde{x}_j = \sum_{i=1}^n Q_{ji}x_i.$$

The sub-Riemannian geodesic flow in $\operatorname{Eng}(n)$ has n+3 constant of motion in involution given by the Hamiltonian function and the momentum map defined by symplectic action of \mathbb{A} . In addition, it possesses as many constants of motion as the dimension of SO(n). These constants of motion are not in volution. However, it is possible to find n constant more involution using the standard proof for potential with radial symmetry; see [17, Appendix B].

Lemma 42. Let $\operatorname{Eng}_{c}(2)$ be the 6 dimensional sub-manifold of $\operatorname{Eng}(n)$ given by

 $\operatorname{Eng}_{c}(2) := \{(x,\theta) \in \operatorname{Eng}(n) : (x,\theta) = (x_{1}, x_{2}, \dots, \theta_{0}, \theta_{1}, \theta_{2}, c_{1}, \dots, c_{n}, \theta_{n+1})\},\$ where $c = (c_{1}, \dots, c_{n})$. If $\gamma(t)$ is a normal sub-Riemannian geodesic in $\operatorname{Eng}(n)$ with momentum $a_{n+1} \neq 0$ such that $\gamma(0)$ is in $\operatorname{Eng}_{c}(2)$ and $\dot{\gamma}(0) = T_{\gamma(0)} \operatorname{Eng}(n)$, then $\gamma(t)$ lies in $\operatorname{Eng}_{c}(2)$ for all t.

The proof of Lemma 42 is the same as the one for Lemma 37.

Corollary 43. Every normal sub-Riemannian geodesic in Eng(n) with momentum $a_{n+1} \neq 0$ has the form $\gamma(t) = \psi_Q(\gamma_0(t))$, where ψ_Q is given by Proposition and $\gamma_0(t)$ is a normal sub-Riemannian geodesic in Eng_c(2).

Corollary 43 implies it is enough to prove Theorem B for the case Eng(2).

Proposition 44. Let $\gamma(t)$ be a normal sub-Riemannian geodesic in Eng(n) with momentum $a_{n+1} \neq 0$ and whose reduce dynamics has zero angular momentum. If $c(t) := \pi(\gamma(t))$, where $\pi : \text{Eng}(n) \to \mathbb{R}^{n+1}$ is the sub-Riemannian projection, then c(t) lays in a two-plane and solves the Euler-Elastica problem.

The Euler-Elastica problem has many ways to characterize. To prove Proposition 44, we will utilize the following criterion: A plane curve is deemed a solution to the Euler-Elastica problem if, at each point, its curvature is directly proportional to the oriented distance from the curve to a specified line, referred to as the directrix.

Proof. By Corollary 43, it is enough to consider a normal sub-Riemannian geodesic in Eng(2). In addition, without loss of generality, we can assume the initial condition $\gamma(0) = (x_1(0), 0, 0, 0, 0, 0)$ for some $x_1(0)$ in \mathbb{R} . Let $(p_x(t), x(t))$ be the solution to the reduced system associated to $\gamma(0)$. The sub-Riemannian geodesic $\gamma(t)$ has angular momentum zero if and only if $p_x(t)$ is parallel to the radial direction, then the reduced solution has the form $(p_x(t), x(t)) = (p_{x_1}(t), 0, x_1(t), 0)$. If $F_{\mu}(x)$ is the polynomial from equation (4.2), then $F_{\mu}(x(t)) = a_0 + a_1 x_1(t) + \frac{a_2}{2} x_1^2(t)$ for all t. Without lose of generality, after a translation in the axis x_1 , we can consider $a_1 = 0$.

By construction, the projected curve c(t) lays on the two-plane given by $(x_1, 0, \theta_0)$. In addition, the **Background Theorem** tells $||\dot{c}(t)|| = 1$ and $\dot{c}(t) = (p_{x_1}(t), 0, F_{\mu}(x(t)))$. We use Hamilton equations (2.3) to find

$$\ddot{c}(t) = \frac{\partial F_{\mu}(x(t))}{\partial x_{1}} (-F_{\mu}(x(t)), 0, p_{x_{1}}(t)), \text{ where } \frac{\partial F_{\mu}(x(t))}{\partial x_{1}} = a_{2}x_{1}(t).$$

So the curvature of c(t) is equal to linear function $a_2x_1(t)$, that is, the curvature is proportional to the oriented distance from $(x_1(t), 0, \theta_0(t))$ to the directrix defined by $x_1 = 0$.

4.3. The Magnetic Space For the Engel-Type Group. Let π_F be the sub-Riemannian submersion from the Engel-type group $\operatorname{Eng}(n)$ to the magnetic space \mathbb{R}_F^{n+2} . We will assume the polynomial $F_{\mu}(x)$ has the form $\alpha + \beta r^2$ where r = ||x|| and $\beta \neq 0$, then we will study some of the properties of this magnetic space.

Let us consider the isometric action $\varphi_{(Q,y_0,z_0)}$ of $SO(n) \times R^2$ on \mathbb{R}_F^{n+2} . If Q is in SO(n) and (y_0, z_0) is in \mathbb{R}^2 , then $\varphi_{(Q,y_0,z_0)}(x, y, z) = (Qx, y+y_0, z+z_0)$.

Lemma 45. Let $R^4_{\mathbb{R}^{n+2}_F}$ be a four-dimensional sub-manifold of \mathbb{R}^{n+2}_F given by

$$R^4_{\mathbb{R}^{n+2}_F} := \{ (x, y, z) \in \mathbb{R}^{n+2}_F : 0 = x_3 = \dots = x_n \}.$$

If c(t) is a normal sub-Riemannian geodesic in \mathbb{R}_F^{n+2} such that c(0) is in $\mathbb{R}_{\mathbb{R}_F^{n+2}}^4$ and $\dot{c}(0)$ is in $T_{c(0)}\mathbb{R}_{\mathbb{R}_F^{n+2}}^4$, then c(t) lies in $\mathbb{R}_{\mathbb{R}_F^{n+2}}^4$ for all time t.

The proof of Lemma 45 is the same as the one for Lemma 37.

Corollary 46. Every normal sub-Riemannian geodesic in \mathbb{R}_F^{n+2} has the form $\gamma(t) = \varphi_{(Q,y_0,z_0)}c_0(t)$ where $c_0(t)$ is a normal sub-Riemannian geodesic in $\mathbb{R}_{\mathbb{R}^{n+2}}^4$ for all t.

Corollary implies that it is enough to understand the sub-Riemannian geodesics in \mathbb{R}^4_F to describe the sub-Riemannian geodesics in \mathbb{R}^{n+2}_F .

In Section 2.2, we defined the pencil Pen_F as the two-dimensional vector space given by the polynomials with form G(x) = a + bF(x), inspired in this definition, we will consider the pencil of effective potentials.

Definition 47. We say that the three-dimensional space Pen_V is the pencil of F(r), if $Pen_V := \{V_{ef}(r) = \frac{\ell^2}{r^2} + G^2(r) : G(r) \in Pen_F\}.$

We can rewrite $\Delta(c, \mathcal{T})$ as a function of the effective potential.

Lemma 48. Let c(t) be an \mathbb{R}^4_F -geodesic for $V_{ef}(r)$ in Pen_V and let \mathcal{T} be a time interval, then $\Delta(c, \mathcal{T})$ from Definition 20 can be rewritten in terms of the effective potential $V_{ef}(r)$ as follows,

$$\Delta(c,\mathcal{T}) = \left(\int_{r(\mathcal{T})} \frac{dr}{\sqrt{1 - V_{ef}(r)}}, \int_{r(\mathcal{T})} \frac{G(r)dr}{\sqrt{1 - V_{ef}(r)}}, \int_{r(\mathcal{T})} \frac{G(r)F(r)dr}{\sqrt{1 - V_{ef}(r)}}\right).$$

In the same way, the map $Cost(c, \mathcal{T})$ from Definition 20 can be rewritten as follows;

$$Cost(c, \mathcal{T}) = 2(\int_{r(\mathcal{T})} \frac{1 - G(r)}{\sqrt{1 - V_{ef}(r)}} dr, \int_{r(\mathcal{T})} \frac{(1 - F(r))G(r)}{\sqrt{1 - V_{ef}(r)}} dr).$$

The proof of Lemma 48 is similar to the proof of Proposition 23.

Corollary 49. Every to sub-Riemannian geodesic associated to (G, ℓ, R) are related by isometry $\varphi_{(Q,y_0,z_0)}(x, y, z)$.

We can generalized the normalized period map for every sub-Riemannian geodesic in \mathbb{R}^4_F -geodesic using the radial Hill interval.

Definition 50. The period map $\Theta : (G, \ell, R) \to [0, \infty] \times \mathbb{R}$ is given by

$$\begin{split} \Theta(G,\ell,R) &:= (\Theta_1(G,\ell,R), \Theta_2(G,\ell,R)) \\ &:= 2(\int_R \frac{1-G(r)}{\sqrt{1-V_{ef}(r)}} dr, \int_R \frac{G(r)(1-F(r))}{\sqrt{1-V_{ef}(r)}} dr). \end{split}$$

4.4. **Proof of Theorem B.** This Sub-section is devoted to proving Theorem B. Let $\gamma_h(t)$ be an arbitrary *r*-homoclinic geodesic in $\operatorname{Eng}(n)$ for a polynomial $F_h(x)$. We will consider the space R_F^{n+2} and the R_F^{n+2} -geodesic $c_h(t) := \pi_{F_h}(\gamma(t))$, then we will prove the following Theorem.

Theorem 51. Let $\gamma_h(t)$ be an arbitrary r-homoclinic geodesic in $\operatorname{Eng}(n)$ for a polynomial $F_h(x)$. If $c_h(t) := \pi_{F_h}(\gamma(t))$, then $c_h(t)$ is a metric line in R_F^{n+2} .

Lemma 35 and Corollary 46 imply it is enough to consider the space \mathbb{R}_F^4 and the *r*-homoclinic geodesic $\gamma_h(t)$ corresponding to the polynomial $F_h(r) := 1 - 2r^2$ and with initial condition c(0) = (1, 0, 0, 0), then by construction $c(t) = (x_1(t), 0, y(t), z(t))$. In addition, the time reversibility of the reduce Hamiltonian H_μ implies $x_1(n) = x_1(-n)$ for all n, so $\Delta x(c_h, [-n, n]) := x(n) - x(-n) = 0$ for all n.

4.4.1. Set UP The Proof of Theorem 51. Theorem 51 is a consequence of Theorem D and Corollary 40. Let us verify the conditions of Theorem D. The following lemma characterizes the r-homoclinic geodesic in $\mathbb{R}^4_{F_d}$ and proves conditions 2 and 3.

Lemma 52. The set of all r-homoclinic classes of geodesic Homc(0) is given by

$$Homc(0)^{\pm} := \{ (G, \ell, x_0) : G(r) = \pm (1 - \beta r^2), \ 0 < \beta, \ \ell = 0, \ and \ ||x_0|| = \frac{2}{\beta} \}.$$

Moreover, the map $\Theta_2 : Homc(0)^{\pm} \to \mathbb{R}$ is one to one, and $Cost(c_h, \mathcal{T})$ is bounded by $\Theta_h := \Theta_1(F_h, 0, [0, 1])$ for all compact interval \mathcal{T} .

Proof. The first part of the proof is consequence of Corollary 39. Let us proof $\Theta_2 : Homc(0)^{\pm} \to \mathbb{R}$ is one to one: let $[c_{\beta}]$ be a class corresponding to the polynomial $G(r) = (1 - \beta r^2)$, then if we set up the change of variable $r = \sqrt{\frac{2}{\beta}}\tilde{r}$, we have

$$\Theta_2([c_\beta]) = 4 \int_{[0,\sqrt{\frac{2}{\beta}}]} \frac{r^2(1-\beta r^2)dr}{\sqrt{1-(1-\beta r^2)^2}} = 4\left(\sqrt{\frac{2}{\beta}}\right)^3 \int_{[0,1]} \frac{\tilde{r}^2(1-2\tilde{r}^2)d\tilde{r}}{\sqrt{1-(1-2\tilde{r}^2)^2}}.$$

We conclude that $\Theta_2([c_\beta])$ is one-to-one when we vary β . The last part proof is the same as the equivalent result for the space \mathbb{R}^3_F , see [16, Lemma 34].

The following lemma verifies condition 1 from Theorem D.

Lemma 53. The space $\mathbb{R}^4_{F_h}$ with $F(r) = 1 - 2r^2$ is geodesically compact.

We will prove Lemma 53 in the Appendix C. The following corollary tells us $c_n(t)$ is not a sequence of line geodesics in particular show condition 4.

Corollary 54. Let n be layer than T_h^* , where T_h^* is given by Corollary 67, then the sequence of geodesics $c_n(t)$ is neither a sequence of geodesics lines nor convergent to a geodesic line. In particular, $c_n(t)$ is strictly normal.

The proof is similar to an equivalent corollary in the context of the jet space, [16, Corollary 50].

4.4.2. Proof of Theorem 51.

Proof. Let $\gamma(t)$ be a *r*-homoclinic geodesic in Eng(*n*) for a polynomial $F_h(r) = 1 - \beta r^2$, and let $c_h(t)$ be the projection of $\gamma(t)$ by π_{F_h} . By Lemma 35, it is enough to consider the *r*-homoclinic geodesic for the polynomial $F(r) = 1 - 2r^2$. Lemma 53, Lemma 52, Corollaries 49 and 54 imply the Conditions 1, 2, 3, and 4, respectively. So Theorem D shows $c_h(t)$ is a metric line in $\mathbb{R}_{F_h}^{n+2}$.

4.4.3. Proof of Theorem B.

Proof. Corollary 40 implies the *r*-periodic geodesics and small oscillation geodesics are not metric lines in Eng(n). Corollary 6 shows the line geodesics are line. Finally, Theorem 51 verify that if $\gamma(t)$ is a *r*-homolinic geodesic, then $c(t) := \pi(\gamma(t))$ is a metric line in \mathbb{R}_F^{n+2} . So Proposition 4 tells $\gamma(t)$ is a metric line in Eng(n).

5. The four-by-four nilpotent triangular matrices group

Let $N_{6,3,1}$ be the whose Lie algebra is spanned by 6 vectors $E_1, E_2, E_0^{\mathfrak{a}}, E_1^{\mathfrak{a}}, E_2^{\mathfrak{a}}$ and $E_3^{\mathfrak{a}}$ satisfying the following non trivial relations

(5.1)
$$E_i^{\mathfrak{a}} := [E_i, E_0^{\mathfrak{a}}] \text{ and } E_3^{\mathfrak{a}} := [E_1, E_2^{\mathfrak{a}}] = [E_2, E_1^{\mathfrak{a}}].$$

Equation (5.1) shows $N_{6,3,1}$ is a metabelian Carnot group, the first layer \mathfrak{g}_1 and the Lie algebra \mathfrak{a} are spanned by $\{E_1, E_2, E_0^{\mathfrak{a}}\}$ and $\{E_0^{\mathfrak{a}}, \ldots, E_{n+1}^{\mathfrak{a}}\}$, respectively. To endow $N_{6,3,1}$ with a left-invariant sub-Riemannian metric, we declare the vector $\{E_1, E_2, E_0^{\mathfrak{a}}\}$ to be orthonormal, then \mathfrak{v} has dimension 1 and \mathfrak{h} is spanned by $\{E_1, E_2\}$, so equation (4.1) implies that $N_{6,3,1}$ satisfy the condition from equation 2.1, and we can consider the sub-Riemannian submersion π_F given in sub-Section 2.2.1. In [21], E. Le Donne and F. Tripaldi denoted the four-by-four nilpotent triangular matrices group by $N_{6,3,1}$, here we used the same nation. In Appendix D, we show the matrix representation of the group $N_{6,3,1}$.

Let $g = (\theta, x)$ be the exponential coordinates of second defined in subsub-Section 2.1.2, where $x = (x_1, x_2)$ and $\theta = (\theta_0, \theta_1, \theta_2, \theta_3)$. The following set of left-invariant vector fields is a frame for the non-integrable distribution \mathcal{D} :

$$X_1 = \frac{\partial}{\partial x_1}, \ X_2 = \frac{\partial}{\partial x_2} \text{ and } Y_0 = \frac{\partial}{\partial \theta_0} + x_1 \frac{\partial}{\partial \theta_1} + x_2 \frac{\partial}{\partial \theta_2} + x_1 x_2 \frac{\partial}{\partial \theta_3}.$$

Then, the sub-Riemannian metric on \mathcal{D} in the coordinates (θ, x) is given by the expression $ds_{N_{6,3,1}}^2 = dx_1^2 + dx_2^2 + d\theta_0^2$.

The following lemma characterize the abnormal curves in the group $N_{6.3,1}$.

Lemma 55. The Engel-type group Eng(n) possesses two families of abnormal curves;

(Family of vertical lines) We say a curve $\gamma(t)$ is an abnormal line, if $\gamma(t)$ is tangent to the vector field Y_0 . So $\gamma(t)$ qualify as a metric line.

(Family of horizontal curve) We say a curve $\gamma(t)$ is an abnormal horizontal curve, if $\gamma(t)$ is tangent to sub-space $\mathcal{H} \subset N_{6,3,1}$, and in addition $\dot{\gamma}(t)$ is perpendicular to a constant vector or $\gamma(t)$ is tangent to hyperbola. So $\gamma(t)$ qualify as a metric line if and only if $\gamma(t)$ is a horizontal line.

We used the terms horizontal and vertical in the same was as Lemma 19. The proof of Lemma 55 is in the Appendix A.

5.1. Reduced Hamiltonian. If $\mu = (a_0, a_1, a_2, a_3)$ is in \mathfrak{a}^* , then equation (2.2) implies that the reduced Hamiltonian H_{μ} is of the form

(5.2)
$$H_{\mu}(p_x, x) = \frac{1}{2}(p_{x_1}^2 + p_{x_2}^2) + \frac{1}{2}(a_0 + a_1x_1 + a_2x_2 + a_3x_1x_2)^2.$$

In the case $a_{n+1} = 0$, the reduced Hamiltonian H_{μ} has the form of a small oscillation system. In the case $a_{n+1} \neq 0$, if we set up the change of coordinates

$$(\tilde{x}_1, \tilde{x}_2) = (\frac{a_1 - a_2}{\sqrt{2}a_3} + x_1, \frac{a_1 + a_2}{\sqrt{2}a_3} + x_2),$$

and define the constants $(\alpha, \beta) := (a_0 - \frac{a_1 a_2}{a_3}, \frac{a_3}{2})$, then the reduced Hamiltonian is

(5.3)
$$H_{\mu}(p_{\tilde{x}},\tilde{x}) = \frac{1}{2}(p_{\tilde{x}_1}^2 + p_{\tilde{x}_2}^2) + \frac{1}{2}(\alpha + \beta \frac{\tilde{x}_2^2 - \tilde{x}_1^2}{2})^2.$$

The reduced Hamiltonian system is not meromorphic integrable; a simple proof of this is using the Darboux points method, consult [25] for more details about the method. The angular momentum $L(p_{\tilde{x}}, \tilde{x}) := p_{\tilde{x}_1}\tilde{x}_2 - p_{\tilde{2}}\tilde{x}_1$ is not conservative quantitative in general for the reduced Hamiltonian system (5.3). However, this system has two regions where the angular momentum $L(p_x, x) = 0$ is constant of motion. So the sub-Riemannian geodesic flow is integrable in these regions, namely the regions $(p_{\tilde{x}_1}, 0, \tilde{x}_1, 0)$ and $(0, p_{\tilde{x}_2}, 0, \tilde{x}_2)$. For simplicity we will replace $(\tilde{x}_1, \tilde{x}_2)$ by (x_1, x_2) for the rest of the Section 5.

The following result characterizes the equilibrium points of the reduced Hamiltonian system (5.3).

Corollary 56. The reduced Hamiltonian system (5.3) has an equilibrium if and only if and $F_{\mu}(r) = \pm (1 + \beta(x_2^2 - x_1^2))$. In addition, this equilibrium point is at $(p_{x_1}, p_{x_2}, x_1, x_2) = (0, 0, 0, 0)$ and has two homoclinic orbits.

Proof. By Hamilton equations, given by (2.3), a point $(p_1, p_2.x_1, x_2)$ is a equilibrium point if and only if $(p_{x_1}, p_{x_2}) = (0, 0)$ and (x_1, x_2) is such that $F_{\mu}(x_1, x_2) = 1$ and $\nabla F_{\mu}(x_1, x_2) = (0, 0)$. The reduced Hamiltonian system has an equilibrium if and only if $F_{\mu}(r) = \pm (1 - \beta(x_2^2 - x_1^2))$. Since the system is Hamiltonian, studying the linear system associated with the equilibrium point is enough to understand its nature. The central manifold has dimension two, and the stable and untestable manifolds have dimension 1. By constructing the homoclinic orbit, we can see that stable and unstable manifold intersect. For instance, if $F_{\mu}(x) = 1 - \beta(x_2^2 - x_1^2)$, or $F_{\mu}(x) = 1 - \beta(-x_2^2 + x_1^2)$ with $0 < \beta$, then the solutions with initial conditions $(0, 0, \pm \sqrt{\frac{2}{\beta}}, 0)$ or $(0, 0, 0, \pm \sqrt{\frac{2}{\beta}})$ are homoclinic orbits.

5.1.1. Classification of the sub-Riemannian geodesics in N_{6,3,1}. Let $\gamma(t)$ be a normal sub-Riemannian geodesic in N_{6,3,1} different of line type, and with momentum μ , then $\gamma(t)$ is only one of the following options:

(Small oscillations) We say a geodesic $\gamma(t)$ is of the type small oscillations, if $a_3 = 0$.

(**Periodic**) We say a geodesic $\gamma(t)$ is of the type periodic if $a_3 \neq 0$, $L(p_x, x) = 0$, and the reduced dynamic is periodic.

(Homoclinic) We say a geodesic $\gamma(t)$ is of the type homoclinic if $a_3 \neq 0$ and the reduced dynamic is a homoclinic orbit.

(Generic) We say a geodesic $\gamma(t)$ is of the type generic if $a_3 \neq 0$ and $L(p_x, x) \neq 0$.

Lemma 57. The sub-Riemannian geodesic in $N_{6,3,1}$ of the type small oscillations, periodic and generic, are not metric lines.

Proof. Lemma 12 implies bounded-generic, and periodic type geodesics do not qualify as metric lines, the proof for the small oscillation is the same as the one for corollary 40.

Let $\gamma(t)$ be an unbounded-generic geodesic; the goal is to show that $\gamma(t)$ does not satisfy the condition from Theorem A. We will use the original coordinates (x_1, x_2) and the Hamiltonian function H_{μ} given by (5.2). Let us consider first the case for the polynomial $F_{\mu}(x) = \beta x_1 x_2$. Without loss of generality, let us assume that the dynamics for $x_1(t)$ is such that $x_1(t) \to \infty$ when $t \to \infty$, then we can find a time t^* such that $x_1(t) > C > 0$ if $t^* < t$. The intuitive idea of the proof is that by Sturm's comparison theorem, consult [34, Theorem 5.20] or [19, sub-Chapter 27.7], the x_2 component is an oscillatory solution, see equation 5.4 below, and it is well know that the integral of oscillatory solutions is finite. Let us formalize the idea:

We use Hamilton equations to find a second-order equation for the coordinates x_1 and x_2 given by

(5.4)
$$\ddot{x}_1 = -\beta^2 x_2^2(t) x_1(t)$$
, and $\ddot{x}_2 = -\beta^2 x_1^2(t) x_2(t)$.

First, we notice that $\ddot{x}_1(t) < 0$ if $t^* < t$, then $x_1(t)$ is decelerating. If $\dot{x}(t) = 0$, in finite time t_1^* , then $\dot{x}(t)$ change sign and keep decreasing until $x_1(t) = 0$. So we conclude that $\dot{x}_1(t)$ is not zero if $t^* < t$. By the above discussion it is enough to consider the cases where the asymptotic behavior is $x_1(t) = ct^{\ell_1} + O(t^{-\ell_2})$ where $0 < \ell_1 \le 1, 0 < \ell_2 \le 1$, and 0 < c < 1, then for big enough t, the second equation from (5.4) is

(5.5)
$$\ddot{x}_2 = -\beta^2 c^2 t^{2\ell_1} x_2(t),$$

we set up the change of variable $x_2(t) = \sqrt{t}u(\frac{\beta c t^{\ell_1+1}}{\ell_1+1})$, then equation 5.5 become in the following Bessel equation and the properties of Bessel functions implies that $x_2(t)$ has the following asymptotic behavior is given by

$$t^2\ddot{u} + t\dot{u} + (t^2 - \frac{1}{2(\ell_1 + 1)})u = 0$$
, and $x_2(t) = \frac{\sin(\frac{\beta c t^{\ell_1 + 1}}{\ell_1 + 1})}{\sqrt{t^{\ell_1}}} + O(t^{\frac{-3(\ell_1 + 1)}{2}}).$

Reefer to [34, Problem 5.31] for more details about Bessel functions.

Let $v = (v_1, v_2, v_3)$ be an unitary vector in the Lie algebra, then

$$\lim_{t \to \infty} |\frac{1}{t} \int_{t^*}^t \rho(v, (L_{\gamma^{-1}(s)})_* \dot{\gamma}(s)) ds| = |v_1 c| < 1.$$

Since the asymptotic behavior implies

$$0 = \lim_{t \to \infty} \frac{1}{t} \int_{t^*}^t \dot{x}_2(s) ds = \lim_{t \to \infty} \frac{1}{t} \int_{t^*}^t x_1(s) x_2(s) ds.$$

It is enough to consider the lower bound t^* in the above integrals, since the integral from 0 to t^* is finite. The proof is the same for the case $t \to \infty$. Therefore, $\gamma(t)$ does not satisfy the condition from Theorem A.

Let us consider a general polynomial $F_{\mu}(x)$, after a translation we write $F_{\mu}(x) = \alpha + \beta x_1, x_2$. Let us consider the curvilinear coordinates defined by

the equipotential curve $F_{\mu}(x) = 0$ and its perpendicular direction $N(y_1)$. If the equipotential curve is $(x_1, x_2) = (y_1, \frac{-\alpha}{\beta y_1})$, then the change of coordinates is given by $(x_1, x_2) = (y_1, \frac{-\alpha}{\beta y_1}) + y_2 N(y_1)$. We will show that Hamiltonian function $H_{\mu}(p_y, y)$ has the same form as the above case when we neglect the terms $\frac{1}{x_1^2}$. Indeed, limit $x_1(t) \to \infty$ then $\frac{1}{x_1^2} \to \infty$, so we approximate the perpendicular vector $N(y_1) \approx (0, 1)$ and set the change of coordinates $(x_1, x_2) = (y_1, \frac{-\alpha}{\beta y_1} + y_2)$. Computing the reduced Hamiltonian in the coordinates $(p_{y_1}, p_{y_2}, y_1, y_2)$, we find

$$H_{\mu}(p_{y_1}, p_{y_2}, y_1, y_2) := \frac{1}{2}(p_{y_1}^2 + p_{y_2}^2) + \frac{1}{2}\beta^2 y_1^2 y_2^2.$$

Then, we repeat the above proof.

5.2. Sub-Riemannian geodesic in $N_{6,3,1}$.

Proposition 58. Let $\gamma(t)$ be a normal sub-Riemannian geodesic in N_{6,3,1} with momentum $a_3 \neq 0$ and whose reduce dynamics has zero angular momentum. If $c(t) := \pi(\gamma(t))$, where $\pi : N_{6,3,1} \to \mathbb{R}^3$ is the sub-Riemannian projection, then c(t) lays in a two-plane and solves the Euler-Elastica problem.

The proof of proposition 58 is the same as the proof of Proposition 44.

5.3. **Proof of Theorem C.** This sub-section is devoted to proving Theorem C. Let $\gamma_h(t)$ be an arbitrary homoclinic geodesic in N_{6,3,1} for a polynomial $F_h(x)$. We will consider the space \mathbb{R}^4_F and the \mathbb{R}^4_F -geodesic $c_h(t) := \pi_{F_h}(\gamma(t))$, then we will prove the following theorem.

Theorem 59. Let $\gamma_h(t)$ be an arbitrary homoclinic geodesic in N_{6,3,1} for a polynomial $F_h(x)$. If $c_h(t) := \pi_{F_h}(\gamma(t))$, then $c_h(t)$ is a metric line in \mathbb{R}_F^4 .

If $F_h(x) := 1 + \beta(x_2^2 - x_1^2)$, then we notice that \mathbb{R}_F^4 has a discrete group of isometries generating by the reflections $R_1(x_1, x_2, y, z) = (-x_1, x_2, y, z)$ and $R_2(x_1, x_2, y, z) = (x_1, -x_2, y, z)$. Therefore, the Carnot dilation, and the group of isometries $Iso(\mathbb{R}_F^4)$ imply it is enough to consider the space \mathbb{R}_F^4 and the homoclinic geodesic $\gamma_h(t)$ corresponding to the polynomial $F_h(x) := 1 + 2(x_2^2 - x_1^2)$ and with initial condition c(0) = (1, 0, 0, 0), then by construction $c(t) = (x_1(t), 0, y(t), z(t))$.

5.3.1. Set up the proof of Theorem 59. Theorem 59 is a consequence of Theorem D and Lemma 57. Let us verify the conditions of Theorem D. The following lemma characterizes the homoclinic geodesic in $\mathbb{R}^4_{F_d}$ and proves conditions 2 and 3 from Theorem D.

Lemma 60. The set of all homoclinic classes of geodesic Homc(0) is given by

$$Homc^{\pm} := \{ (G, x_0) : G(x) = \pm (1 + \beta (x_2^2 - x_1^2)), \ 0 < \beta, \ x_0 = (\sqrt{\frac{2}{\beta}}, 0) \}$$
$$\cup \{ (G, x_0) : G(x) = \pm (1 + \beta (-x_2^2 + x_1^2)), \ 0 < \beta, \ x_0 = (0, \sqrt{\frac{2}{\beta}}) \}.$$

Moreover, the map $\Theta_2 : Pen_h^+ \to \mathbb{R}$ is one to one, and $Cost(c_h, \mathcal{T})$ is bounded by $\Theta_h := \Theta_1(F_h, [x])$ for all compact interval \mathcal{T} .

Proof. The first part of the proof is consequence of Corollary 56. Let us prove $\Theta_2 : Pen_h^+ \to \mathbb{R}$ is one to one; if $[c_\beta^1]$ is the class corresponding to the polynomial $G(x) = (1 + \beta(x_2^2 - x_1^2))$ and initial condition $x_0 = (\sqrt{\frac{2}{\beta}}, 0)$, then $x_2(t) = 0$ for all t and we can rewrite $\Theta_2([c_\beta^1])$ as an integral of one variable where $ds = dx_1$. We have that

$$\Theta_2([c_\beta^1]) = 4 \int_{[0,\sqrt{\frac{2}{\beta}}]} \frac{x_1^2(1-\beta x_1^2)dx_1}{\sqrt{1-(1-\beta x_1^2)^2}}$$

If $[c_{\beta}^2]$ is the class corresponding to the polynomial $G(x) = (1 + \beta(-x_2^2 + x_1^2))$ and initial condition $x_0 = (0, \sqrt{\frac{2}{\beta}})$, then $x_1(t) = 0$ for all t and we can rewrite $\Theta_2([c_{\beta}^2])$ as an integral of one variable where $ds = dx_2$. We have that

$$\Theta_2([c_\beta^2]) = -4 \int_{[0,\sqrt{\frac{2}{\beta}}]} \frac{x_2^2(1-\beta x_2^2)dx_2}{\sqrt{1-(1-\beta x_2^2)^2}}.$$

Therefore, we conclude that $\Theta_2([c_1]) = -\Theta_2([c_2])$. We repeat the proof from 52 to show that $\Theta_2([c_1])$ is one-to-one when we vary β .

The following lemma verifies condition 1.

Lemma 61. Let $\mathbb{R}^4_{F_h}$ be a magnetic space, where $F(x) = 1 + 2(x_2^2 - x_1^2)$, then the region $B_1(0) \times \mathbb{R}^2$ of the space is geodesically compact.

Proof. To prove Lemma 57, we will show that unbounded solution behaves as a harmonic oscillator in the perpendicular direction to the asymptote, so the polynomial G(x) oscillates between [-1, 1]. If $\lim_{n\to\infty} \mathcal{T}_n = [-\infty, \infty]$, a necessary condition for $Cost_t(c_n, \mathcal{T}_n)$ to be finite is that G(x(t)) = 1, when $t \to \pm \infty$. So $Cost(c_n, \mathcal{T}_n)$ is not bounded. \Box

The following corollary tells us $c_n(t)$ is not a sequence of line geodesics in particular show condition 4.

Corollary 62. Let n be layer than T_h^* , where T_h^* is given by Corollary 67, then the sequence of geodesics $c_n(t)$ is neither a sequence of geodesics lines nor convergent to a geodesic line. In particular, $c_n(t)$ is strictly normal.

The proof of Corollary 62 is the same to an equivalent result in \mathbb{R}^3_F , see [16, Corollary 50], respectively.

5.3.2. Proof of Theorem 59.

Proof. By the Carnot dilatation, it is enough to consider the homoclinic geodesic for the polynomial $F(x) = 1 + 2(x_2^2 - x_1^2)$. Let $\gamma(t)$ be a homoclinic geodesic in N_{6,3,1} for a polynomial $F_h(x)$, and $c_h(t)$ be the projection of $\gamma(t)$ by π_{F_h} . Lemma 61, Lemma 60, Corollary 62 imply the Conditions 1, 2, 3, and 4, respectively. So Theorem D shows $c_h(t)$ is a metric line in $\mathbb{R}^4_{F_h}$. \Box

5.3.3. Proof of Theorem B.

Proof. Lemma 57 implies that the small oscillations, periodic, and generic geodesics are not metric lines in N_{6,3,1}. Corollary 6 shows the line geodesics are line. Finally, Theorem 59 verify that if $\gamma(t)$ is a homolinic geodesic, then $c(t) := \pi(\gamma(t))$ is a metric line in \mathbb{R}_F^4 . So Proposition 4 tells $\gamma(t)$ is a metric line in Eng(n).

6. CONCLUSION

We present a method to prove that a particular sub-Riemannian geodesic is a metric line by considering its projection to the sub-Riemannian magnetic space \mathbb{R}_F^{n+2} and verifying the conditions of Theorem D. Theorem 51 completely classifies the metric lines in Engel-type group, Theorem C provide a partial result in the classification on the group four-by-four nilpotent triangular matrices. Lemma 57 is the first proof that that the generic-unbounded geodesics are not metric lines.

Future work:

- Provide a complete classification of the metric lines in the metabelian Carnot groups of step 3.
- Provide a general proof that the generic-unbounded geodesics do not qualify as a metric line.
- Prove that bounded-generic geodesics arbitrary close to a critical point do not qualify as a metric line.
- Prove the homoclinic geodesics in the jet space $J^k(\mathbb{R}, \mathbb{R})$ are metric lines.

Two interesting question are:

- Are all the metric lines in a Carnot group of step 3 related to Euler-Solition in the same way that Propositions 44 and 58?
- Are all the metric lines in a Carnot group of step 3 homoclinic?

APPENDIX A. ABNORMAL GEODESICS

A.1. Proof of Lemma 19.

Proof. We will compute the abnormal curves using the Pontryagin's maximum principle. Let us look the following optimal control problem:

(A.1)
$$\dot{c}(t) = u_1 X_1 + \dots + u_n X_n + u_{n+1} Y,$$

with the boundary conditions $\gamma(0)$ and $\gamma(T)$, we notice that equation (A.1) implies $\dot{x}_i = u_i$. It follows from Cauchy-Schwarz inequality that minimizing the sub-Riemannian length is equivalent to minimizing the action

(A.2)
$$\int_0^T \frac{u_1^2 + \dots + u_{n+1}^2}{2} dt \to \min,$$

with the condition $u_1^2 + \cdots + u_{n+1}^2$. Thus, the Hamiltonian associated to the optimal control problem (A.1) and (A.2) for the abnormal case is

(A.3)
$$H(p, x, y, z, u) = u_1 p_{x_1} + \dots + u_n p_{x_n} + u_{n+1} (p_y + p_z F(x)).$$

From the Pontrygin's maximum principle, we obtain a Hamiltonian system for the variables $p_{x_1}, \cdots, p_{x_n}, p_y$ and p_z :

(A.4)
$$\dot{p}_{x_i} = u_{n+1} \frac{\partial F}{\partial x_i}$$
 for $i = 1, \dots, n, \ \dot{p}_y = 0$ and $\dot{p}_z = 0$,

the maximum condition

(A.5)
$$\max_{u \in \mathbb{R}^{n+1}} H(p(t), x(t), u) = H(p(t), \hat{x}(t), \hat{y}(t), \hat{z}(t), \hat{u}(t))$$

where $\hat{u}(t), \hat{c}(t) = (\hat{x}(t), \hat{y}(t), \hat{z}(t))$ is the optimal process, and the condition of non-triviality $p(t) \neq 0$. Therefore, the maximum condition imply

(A.6)
$$0 = p_{x_i}$$
, for $i = 1, ..., n$, and $0 = p_y + pzF(x)$.

We notice that the non-trivial condition imply p_z is a non-zero constant. By derivating equation (A.6) with respect time and using equation (A.4), we obtain;

(A.7)
$$0 = u_{n+1} \frac{\partial F}{\partial x_i}$$
 for $i = 1, ..., n$, and $0 = \frac{\partial F}{\partial x_1} u_1 + \dots + \frac{\partial F}{\partial x_n} u_n$

The left equation from (A.7) tells $u_{n+1} = 0$ or $\frac{\partial F}{\partial x_i} = 0$ for $i = 1, \ldots, n$. (Family of vertical lines) Let us assume $\frac{\partial F}{\partial x_i} = 0$ for $i = 1, \ldots, n$, then equation A.7 implies $u_1 = \cdots = u_n = 0$. We conclude that $u_{n+1} = \pm 1$, so c(t) is a vertical line.

(Family of horizontal isocontour) Let us assume $u_{n+1} = 0$, then c(t)is tangent to \mathcal{H} and the second equation from (A.7) implies the curve x(t)belongs to a level set of F(x), a curve in the Euclidean sub-space \mathcal{H} is a metric line if and only if is a line.

A.2. Proof of Lemma 36.

Proof. Following the strategy from the proof of Lemma 19. We have

(A.8)
$$H(p,g,u) = u_1 P_{X_1} + \dots + u_n P_{X_n} + u_{n+1} P_{Y_0},$$

where $P_{X_1}, \dots, P_{X_n}, P_{Y_0}$ are the left-invariant momentum function associated to the left invariant vector fields tangent to \mathcal{D} . From the Pontrygin's maximum principle, we obtain a Hamiltonian system for the left-invariant momentum functions:

(A.9)
$$\dot{P}_{X_i} = -u_{n+1}P_{Y_i} \text{ for } i = 1, \dots, n, \quad \dot{P}_{Y_0} = u_1P_{Y_1} + \dots + u_nP_{Y_n}, \\ \dot{P}_{Y_i} = u_iP_{Y_{n+1}} \text{ for } i = 1, \dots, n, \text{ and } \dot{P}_{Y_{n+1}} = 0.$$

Then, the maximum condition imply

(A.10)
$$0 = -u_{n+1}P_{Y_i}$$
, for $i = 1, ..., n$, and $0 = u_1P_{Y_1} + \dots + u_nP_{Y_n}$.

(Family of vertical lines) Let us consider the case $P_{Y_1} = \cdots = P_{Y_n} = 0$, the no trivial condition implies $P_{Y_{n+1}} \neq 0$. Since P_{Y_1}, \cdots, P_{Y_n} are constant, then equation (A.9) says $0 = u_i P_{Y_{n+1}}$ for $i = 1, \ldots, n$, so $0 = u_i$ for $i = 1, \ldots, n$. We conclude that $u_{n+1} = \pm 1$, so $\gamma(t)$ is a vertical line.

(Family of horizontal curves) Let us consider the case $P_{Y_{n+1}} = 0$, then (A.9) and the non-trivial condition show $(P_{Y_1}, \dots, P_{Y_n})$ is a constant vector different of zero. Therefore, the (A.10) implies $u_{n+1} = 0$ and $(P_{Y_1}, \dots, P_{Y_n})$. $(u_1, \dots, u_n) = 0$, so $\gamma(t)$ is a Horizontal curve tangent to \mathcal{H} and perpendicular to the vector $(P_{Y_1}, \dots, P_{Y_n})$.

Let us consider the case $P_{Y_{n+1}} \neq 0$, then equation (A.9) yields $u_i = \frac{\dot{P}_{Y_i}}{P_{Y_{n+1}}}$, plugin the last expression (A.10) implies

$$0 = \dot{P}_{Y_1} P_{Y_1} + \dots + \dot{P}_{Y_n} P_{Y_n} \text{ then } R^2 = P_{Y_1}^2 + \dots + P_{Y_n}^2.$$

We conclude that $(P_{Y_1}, \dots, P_{Y_n})$ is never the zero vector if $R \neq 0$, so $u_{n+1} = 0$, and $\gamma(t)$ is tangent to a sphere in \mathcal{H} . If R = 0, then $\gamma(t)$ is vertical line.

A.3. Proof of Lemma 55.

Proof. Following the strategy from the proof of Lemma 36. We have

(A.11)
$$H(p, g, u) = u_1 P_{X_1} + u_2 P_{X_2} + u_3 P_{Y_0},$$

where $P_{X_1}, P_{X_2}, P_{Y_0}$ are the left-invariant momentum function associated to the left invariant vector fields tangent to \mathcal{D} . From the Pontrygin's maximum principle, we obtain a Hamiltonian system for the left-invariant momentum functions:

(A.12)
$$\dot{P}_{X_1} = -u_3 P_{Y_1}, \quad \dot{P}_{X_2} = -u_3 P_{Y_2}, \text{ and } \dot{P}_{Y_0} = u_1 P_{Y_1} + u_2 P_{Y_2}, \\ \dot{P}_{Y_1} = u_2 P_{Y_3}, \quad \dot{P}_{Y_2} = u_1 P_{Y_3} \quad \text{and} \quad \dot{P}_{Y_{n+1}} = 0.$$

Then, the maximum condition imply

(A.13)
$$0 = -u_{n+1}P_{Y_i}$$
, for $i = 1, ..., n$, and $0 = u_1P_{Y_1} + \dots + u_nP_{Y_n}$.

(Family of vertical lines) The assumption $P_{Y_1} = P_{Y_2} = 0$ implies $0 = u_1 = u_2$, in same way that in Lemma 36. We conclude that $u_{n+1} = \pm 1$, so $\gamma(t)$ is a vertical line.

(Family of horizontal curves) The assumption $P_{Y_3} = 0$ implies $u_3 = 0$ and $(P_{Y_1}, P_{Y_2}) \cdot (u_1, u_2) = 0$, in same way that in Lemma 36. We conclude that $\gamma(t)$ is a Horizontal curve tangent to \mathcal{H} and perpendicular to the constant vector (P_{Y_1}, P_{Y_2}) .

Let us consider the case $P_{Y_3} \neq 0$, then equation (A.12) yields $u_1 = \frac{P_{Y_2}}{P_{Y_3}}$ and $u_2 = \frac{\dot{P}_{Y_1}}{P_{Y_3}}$, plugin the last expression (A.13) implies

$$0 = \dot{P}_{Y_2} P_{Y_1} + \dot{P}_{Y_1} P_{Y_2} \text{ then } R^2 = P_{Y_1} P_{Y_2}.$$

We conclude that (P_{Y_1}, P_{Y_2}) is never the zero vector if $R \neq 0$, so $u_3 = 0$, and $\gamma(t)$ is tangent to a hyperbola in \mathcal{H} . If R = 0, then $\gamma(t)$ is vertical line. \Box

Appendix B. Proof of Theorem D

The goal is to show that for arbitrary T > 0, the geodesic $c_h(t)$ is a minimizer in the interval [-T, T]. The strategy to prove this goal is the following: for all n > T, we will take a sequence of minimizing geodesics $c_n(t)$ in the interval $[0, T_n]$ and joining the points $c_h(-n) = c_n(0)$ and $c_h(n) = c_n(T_n)$, see figure B.1. Thus, we will show that exist a convergent sub-sequence $c_{n_j}(t)$ converging to a homoclinic geodesic $c_{\infty}(t)$ in $Min(K, \mathcal{T})$, where K is a compact sub-set of \mathbb{R}_F^{n+2} and \mathcal{T} is a compact interval. Conditions 2 implies that $c_{\infty}(t)$ is a homoclinic geodesic for $G_h(x)$, and there ψ in $Iso(\mathbb{R}_F^{n+2})$ such that $c_d([-T,T]) \subset \psi(c_{\infty}(\mathcal{T}))$. By Lemma 64, see below, we will conclude $c_h(t)$ is minimizing in [-T,T]. Since T is arbitrary, $c_h(t)$ is a metric line.

B.0.1. Prelude of Proof. Let us present two classical results on metric spaces.

Proposition 63. Let K be a compact subset of \mathbb{R}_F^{n+2} and let \mathcal{T} be a compact time interval. Let us define the following set of \mathbb{R}_F^{n+2} -geodesics

 $Min(K,\mathcal{T}) := \{\mathbb{R}_F^{n+2} \text{-}geodesics \ c(t) : c(\mathcal{T}) \subset K \text{ and } c(t) \text{ is minimizing in } \mathcal{T}\}.$

Then $Min(K, \mathcal{T})$ is a sequentially compact set with respect to the uniform topology.

The proof of Proposition 63 is the same as the equivalent result in the context of the magnetic space \mathbb{R}^3_F , see [16, Proposition 28]. The following lemma is the last necessary tool to prove Theorem D.

Lemma 64. Let $c_1(t)$ be a \mathbb{R}_F^{n+2} -geodesic in $Min(K, \mathcal{T})$ and let $c_2(t)$ be a \mathbb{R}_F^{n+2} -geodesic. If $\varphi(x, y, z)$ is an isometry such that $c_2(\mathcal{T}') \subset \varphi(c_1(\mathcal{T}))$, then $c_2(t)$ is in $Min(\varphi(K), \mathcal{T}')$.

Definition 65. Let $\mathcal{P}(\mathcal{H}, k)$ be the vector space of polynomial from \mathcal{H} to \mathbb{R} of degree bounded by k, and let $||F||_{\infty} := \sup_{x \in K_{\mathcal{H}}} |F(x)|$ be the uniform norm where $K_{\mathcal{H}}$ is a compact set. We denote by $B(K_{\mathcal{H}}, k)$ the closed ball of radius 1.

The following Proposition is a classical result in metric space.



FIGURE B.1. The images show the (x, y)-plane and the projections by pr of the homoclinic-geodesic $c_h(t)$ and the sequence of minimizing geodesics $c_n(t)$.

Proposition 66. $B(K_{\mathcal{H}}, k)$ is a compact set.

Consult [16, Appendix A] for the proof of Proposition 66 for the case $\dim \mathcal{H} = 1$, the general proof is the same.

B.0.2. Set Up The Proof of Theorem D. Let us consider a homoclinic geodesic $c_h(t) = (x_h(t), y_h(t), z_h(t))$ holding the conditions from Theorem D.

Corollary 67. There exist a positive time T_h^* such that if $T_h^* < t$, then $y_h(t) > 0$ and $y_h(-t) < 0$.

Proof. Since $c_h(t)$ is a homoclinic geodesic for a polynomial $G_h(x)$ in $Pen_h^+(x_0)$, then $\lim_{t\to\pm\infty} y(t) = \pm\infty$.

Let T be arbitrary large and consider the points $c_h(-n)$ and $c_h(n)$ where T < n and n is in N. Let $c_n(t) = (x_n(t), y_n(t), x_n(t))$ be a sequence of minimizers geodesics in the interval $[0, T_n]$ such that:

(B.1)
$$c_n(0) = c_h(-n), \quad c_n(T_n) = c_h(n) \text{ and } T_n \le n$$

We call the equations and inequality from (B.1) the endpoints and shorter conditions, respectively. Since the endpoints holds for all n, the sequence $c_n(t)$ has the asymptotic conditions:

(B.2)
$$\lim_{n \to \infty} c_n(0) = (x_0, -\infty, -\infty) \text{ and } \lim_{n \to \infty} c_n(T_n) = (x_0, \infty, \infty).$$

In addition, the endpoint condition implies difference of the endpoint are equal

$$\Delta y(c_h, [-n, n]) = \Delta y(c_n, [0, T_n]), \text{ and } \Delta y(c_h, [-n, n]) = \Delta y(c_n, [0, T_n]),$$
for all *n*. Therefore;

(B.3)
$$Cost_t(c_h, [-n, n]) \ge Cost_t(c_n, [0, T_n]), \\ Cost_y(c_h, [-n, n]) = Cost_y(c_n, [0, T_n]),$$

for all n. Condition 3 implies $Cost(c_n, [0, T_n])$ is uniformly bounded and Lemma ?? yields the asymptotic period condition;

(B.4)
$$\lim_{n \to \infty} Cost_y(c_n, [0, T_n]) = \Theta_2(G_h, [x]).$$

We remark that a equation (B.4) does not tell that $c_n(n)$ converges to $c_h(t)$ neither $G_n(c)$ converges to $G_h(x)$. Only tells that the sequence $Cost_y(c_n, [0, T_n])$ in \mathbb{R} converges to the value $\Theta_2(G_h, [x])$.

Condition 4 implies $c_n(t)$ is a normal geodesic, so we can assume that $c_n(t)$ is geodesic for a polynomial $G_n(x)$ and with reduced dynamics $x_n(t)$. The construction of the geodesic $c_n(t)$ is such that the initial condition is unbounded. The following proposition provides a bounded initial condition.

Proposition 68. Let n be a natural number layer than T_h^* , where T_h^* is given by corollary 67, then there exist a compact sub-set $K_0 \subset R_{F_h}^{n+2}$ and a time t_n^* in $(0, T_n)$ such that $c_n(t_n^*)$ is in K_0 for all $n > T_h^*$.

Proposition 68 extends equivalent Proposition within the magnetic space \mathbb{R}_F^3 , the proof's strategy remaining consistent except for the final step. Subsequently, we proceed with the proof. Conversely, Corollary 70, Lemmas 71 and 72 generalize analogous results within the magnetic space \mathbb{R}_F^3 , with the proof methodology remaining unchanged. For reference, consult [16, Corollary 40, Lemma 41 and Lemma 42].

Proof. Let n be a natural number larger than T_d^* . By construction, $y_n(0) < 0$ and $y_n(T_n) > 0$, the intermediate value theorem implies that exist a t_n^* in $(0, T_n)$ such that $y_n(t_n^*) = 0$. As we saw above $Cost(c_n, [0, T_n])$ is uniformly bounded, and Condition 1 tells that the region $B(r, x_0) \times \mathbb{R}^2$ is geodesically compact, then there exists a constant C_1 and a compact set K_x such that $||Cost(c_n, [0, T_n])||_{\infty} < C_1$ for all n, and $x_n(t)$ is in K_x for all t in $[0, T_n]$.

Let us prove that $z_n(t_n^*)$ is bounded: as we saw the endpoint condition implies equation B.3, in particular $Cost_y(c_d, [-n, n]) = Cost_y(c_n, [0, T_n])$. The the definition of $Cost_y$ yields

$$z_n(t_n^*) - z_n(0) = \Delta z(c_n, [0, t_n^*]) = \Delta y(c_n, [0, t_n^*]) - Cost_y(c_n, [0, t_n^*]),$$

$$z_d(0) - z_d(-n) = \Delta z(c_d, [-n, 0]) = \Delta y(c_d, [-n, 0]) - Cost_y(c_d, [-n, 0]).$$

By construction;

$$\Delta y(c_n, [0, t_n^*]) = \Delta y(c_d, [-n, 0]), \ z_d(0) = 0 \text{ and } z_n(0) = z_d(-n),$$

then

(B.5)
$$\begin{aligned} |z_n(t_n^*)| &= |\Delta z(c_n, [0, t_n^*]) - \Delta z(c_d, [-n, 0])| \\ &\leq |Cost_y(c_n, [0, t_n^*])| + |Cost_y(c_d, [-n, 0])| \end{aligned}$$

Condition 3 says $Cost_y(c_d, [-n, 0])$ is bounded by some constant C_1 . Lemma below tells $Cost_y(c_n, [0, t_n^*])$ is bounded by some constant C_2 .

Let K_0 be the compact set given by $K_x \times [-1,1] \times [-C_1 - C_2, C_1 + C_2]$. We just proved $c_n(t_n^*)$ is in K_0 . Let us prove that $Cost_y(c_n, [0, t_n^*])$ is bounded by C_2 .

Lemma 69. There exist a constant C_2 such that $Cost_y(c_n, [0, t_n^*])$ is bounded by C_2 .

The strategy is to use the well-know result that a function from sequentially compact set to the real number attain a maximum.

Proof. Let Ω_n be the Hill region containing the point $x_n(t_n^*)$, let us notice that the interior of Ω_n is not empty: Indeed, if Ω is an isolated point then $G_n(x_n(t_n^*)) = \pm 1$ and $x_n(t_n^*)$ is a critical point of F. The by Lemma 19, the geodesic $c_n(t)$ is a abnormal geodesic and this is a contradiction with condition 4. Therefore, we for every n we can find a compact set $K_{\mathcal{H}}^n$ such that $x_n(t_n^*)$ is in $K_{\mathcal{H}}^n$ and $K_{\mathcal{H}}^n$ is inside Ω_n . Let $\tilde{K}_{\mathcal{H}}^n$ be the image of the compact set $K_{\mathcal{H}}^n$ by the translation $x \to x - x_n(t_n^*)$, $\tilde{K}_{\mathcal{H}}^n$ contains 0 for all n and the compact set $\tilde{K}_{\mathcal{H}}$ given by the intersection $\bigcap_{n \in \mathbb{N}} \tilde{K}_{\mathcal{H}}^n$ is not empty. Moreover, the interior is not empty by condition 4, using the above argument. Therefore, the sequence of polynomial $G_n(x - x_n(t_n^*))$ lays in the compact set $B(\tilde{K}_{\mathcal{H}}, s - 1)$.

The family of curves $x_n([0, t_n^*])$ is uniformly bounded and equicotinuous, since Condition 1 implies $x_n([0, t_n^*])$ belongs to $K_{\mathcal{H}}$ for all n and is smooth, then the family of curves $x_n([0, t_n^*])$ is sequentially compact.

Therefore, by Proposition 23 the domain of function $Cost_y(c_n, [0, t_n^*])$ is sequentially compact set. So $Cost_y(c_n, [0, t_n^*])$ it bounded by some constant C_2 .

Let us reparametrize the minimizing geodesic $c_n(t)$ sequence. Let $\tilde{c}_n(t)$ be the minimizing geodesic in the interval $\mathcal{T}_n := [-t_n^*, T_n - t_n^*]$ given by $\tilde{c}_n(t) := c_n(t + t_n^*)$, then $\tilde{c}_n(0)$ is bounded.

Corollary 70. There exist a subsequence \mathcal{T}_{n_i} such that $\mathcal{T}_{n_i} \subset \mathcal{T}_{n_{i+1}}$.

For simplicity, we will use the notation \mathcal{T}_n for the subsequence \mathcal{T}_{n_i} .

Lemma 71. Let N be a natural number larger than T_h^* , Then there exist a compact set $K_N \subset \mathbb{R}_F^{n+2}$ such that $\tilde{c}_n(t)$ is in $Min(K_n, \mathcal{T}_N)$ if n > N.

Therefore, $\tilde{c}_n(t)$ has a convergent subsequence $\tilde{c}_{n_j}(t)$ converging to a geodesic $c_{\infty}(t)$ in $Min((K_N, \mathcal{T}_N))$. Condition 4 implies that $c_{\infty}(t)$ is a normal geodesic, then there exist a polynomial $G_{\infty}(x)$ in Pen_F . The following Lemma tells $G_h(x) = G_{\infty}(x)$.

Lemma 72. The unique polynomial in $Pen_F(x_0)$ satisfying the asymptotic conditions given by (B.2) and (B.4) is $G_h(x)$.

B.0.3. Proof of Theorem D.

Proof. Let $\tilde{c}_n(t)$ be the sequence of geodesic defined by the endpoints condition given by equation (B.1). By Lemma 71, for all $N > T_h^*$ there exist a compact set K_N and compact time interval \mathcal{T}_N such that $\tilde{c}(t)$ is in

 $Min(K_n, \mathcal{T}_n)$ if n > N. By proposition 63, $\tilde{c}(t)$ has a convergent subsequence converging to a geodesic $c_{\infty}(t)$ in $Min(K_n, \mathcal{T}_n)$. Condition 4 implies $c_{\infty}(t)$ is normal geodesic for a polynomial $G_{\infty}(x)$ in $Pen_F(x_0)$. Lemma 72 tells us that $G_h(x) = G_{\infty}(x)$.

Since $c_{\infty}(t)$ and $c_h(t)$ are homolinic geodesics for $G_h(x)$ whose reduce dynamics converge to x_0 , they belong to the same equivalence class, this implies there exist a map ψ in $Iso(\mathbb{R}_F^{n+1})$ sending $c_{\infty}(t)$ to $c_h(t)$. Using N is arbitrary and $c_h([-T,T])$ is bounded, we can find a compact set $K := K_N$ and $\mathcal{T} := \mathcal{T}_N$ such that $c_d([-T,T]) \subset \psi(c_{\infty}(\mathcal{T}))$ and c_{∞} is in $Min(K,\mathcal{T})$. Lemma 64 implies that $c_h(t)$ is minimizing in [-T,T] and T is arbitrary. Therefore, $c_h(t)$ is a metric line in \mathbb{R}_F^{n+2} .

Appendix C. Proof of Lemma 53

C.1. Proof Lemma 53.

Proof. Let $c_n(t) = (x_n(t), y_n(t), z_n(t))$ be a sequence of geodesic in \mathbb{R}_F^{n+2} with $F(r) = 1 - br^2$ traveling during a time interval \mathcal{T}_n and with travel curve $\mathcal{C}(c_n, \mathcal{T}_n)$ such that $x_n(\partial \mathcal{T}_n)$ is in $B_{\mathcal{H}}(r)$ for all n. We will prove that if \mathcal{C}_n is unbounded, then $Cost(c_n, \mathcal{T}_n)$ is unbounded.

The sequence of geodesics $c_n(t)$ induces a sequence of pairs (G_n, ℓ_n) . The first goal is to extract a subsequence of converging pairs (G_n, ℓ_n) . One side, If $r(\mathcal{T}_n) = [(r_{min})_n, (r_{max})_n]$, we consider the affine map $h_n(\tilde{r}) = (r_{min})_n + u_n \tilde{r}$ with $u_n := (r_{max})_n - (r_{min})_n$, which maps [0, 1] to $r(\mathcal{T}_n)$. We define $\tilde{G}_n(\tilde{r}) = G_n(h_n(\tilde{r}))$, so \tilde{G}_n is in $\mathcal{P}(k)$. On the other side, we notice the condition $x_n(\partial \mathcal{T}_n)$ implies $|\ell_n| \leq r$, then ℓ_n is a sequence of bounded real number, then there exist a subsequence $(\tilde{G}_{n_j}, \ell_n)$ converging to (\tilde{G}, ℓ) . Let us proceed by the following cases: case $\tilde{G}(\tilde{r}) \neq 1$ and case $\tilde{G}(\tilde{r}) = 1$.

Case $\tilde{G}(\tilde{r}) \neq 1$: we find $Cost_t(c_n, \mathcal{T}_n) = u_n Cots_t(\tilde{G}_n, \ell_n, [0, 1])$, by setting the change of variable $r = h_n(\tilde{r})$. By Fatou's lemma

$$0 < Cost_t(\tilde{G}, \ell, [0, 1]) \leq \liminf_{n_j \to \infty} Cost_t(\tilde{G}_{n_j}, \ell_{n_j}, [0, 1]).$$

Then, $u_{n_j} \to \infty$ implies $Cost_t(c_{n_j}, \mathcal{T}_{n_j})$.

Case $\tilde{G}(\tilde{r}) = 1$: there exist n^* such that $\tilde{G}_{n_j}(\tilde{r}) > \frac{1}{2}$ for all \tilde{r} in [0, 1] if $n_j > n^*$. We notice $1 - F(r) = br^2$, then

$$|Cost_y(G_{n_j}, \ell_{n_j}, \mathcal{R}_{n_j})| > \frac{1}{\sqrt{3}} \int_{\mathcal{R}_{n_j}} |b| r^2 dr.$$

So $|Cost_y(G_{n_j}, \ell_{n_j}, \mathcal{R}_{n_j})| \to \infty$, when $n_j \to \infty$.

Appendix D. Matrix representation of $N_{6,3,1}$

Let us denote by E_{ij} the square matrix with entry 1 where the *i*th row and *j*th column meet, all the other entries being zero. If $\hat{E}_1, \hat{E}_2, \hat{E}_0^{\mathfrak{a}}, \hat{E}_1^{\mathfrak{a}}, \hat{E}_2^{\mathfrak{a}}$ and $\hat{E}_3^{\mathfrak{a}}$ are the matrix representation of the elements of the Lie algebra $E_1, E_2, E_0^{\mathfrak{a}}, E_1^{\mathfrak{a}}, E_2^{\mathfrak{a}}$ and $E_3^{\mathfrak{a}}$. Then

$$\hat{E}_1 = E_{21}, \ \hat{E}_2 = -E_{43}, \ \hat{E}_0^{\mathfrak{a}} = E_{32}, \ \hat{E}_1^{\mathfrak{a}} = E_{31}, \ \hat{E}_2^{\mathfrak{a}} = E_{43}, \ \hat{E}_3^{\mathfrak{a}} = E_{41}.$$

References

- [1]
- R. Abraham and J.E. Marsden. Foundations of Mechanics. AMS Chelsea publishing. AMS Chelsea Pub./American Mathematical Society, 2008.
- [3] A.A Agrachev, D. Barilari, and U. Boscain. A Comprehensive Introduction to Sub-Riemannian Geometry. Cambridge University Press, 2019.
- [4] A.A. Agrachev and Y.L. Sachkov. Control Theory from the Geometric Viewpoint. Control theory and optimization. Springer, 2004.
- [5] A. Anzaldo-Meneses and F. Monroy-Perez. Goursat distribution and sub-riemannian structures. Journal of Mathematical Physics, 44(12):6101–6111.
- [6] A. Anzaldo-Meneses and F. Monroy-Perez. Integrability of nilpotent sub-riemannian structures.
- [7] A.A. Ardentov and Y.L. Sachkov. Conjugate points in nilpotent sub-riemannian problem on the engel group. *Journal of Mathematical Sciences*, 195(3):369–390.
- [8] A.A. Ardentov and Y.L. Sachkov. Extremal trajectories in a nilpotent sub-riemannian problem on the engel group. *Sbornik: Mathematics*, 202(11):1593–1615.
- [9] A.A. Ardentov and Y.L. Sachkov. Cut time in sub-riemannian problem on engel group. 2014.
- [10] A.A. Ardentov and Y.L. Sachkov. Maxwell strata and cut locus in sub-riemannian problem on engel group. *Regular and Chaotic Dynamics*, 22, 09 2017.
- [11] Andrei Ardentov and Eero Hakavuori. Cut time in the sub-riemannian problem on the cartan group. ESAIM: Control, Optimisation and Calculus of Variations, 28, 01 2022.
- [12] V. I. Arnol'd. Mathematical methods of classical mechanics. Springer Science.
- [13] A. Bravo-Doddoli. Higher elastica: Geodesics in the jet space. European Journal of Mathematics.
- [14] A. Bravo-Doddoli. No periodic geodesic in the jet space. Pacific Journal of Mathematics.
- [15] A. Bravo-Doddoli and R. Montgomery. Geodesics in Jet Space. Regular and Chaotic Dynamics, 2021.
- [16] Alejandro Bravo-Doddoli. Metric lines in jet space, 2023.
- [17] Alejandro Bravo-Doddoli, Enrico Le Donne, and Nicola Paddeu. Sympletic reduction of the sub-riemannian geodesic flow for metabelian nilpotent groups, 2023.
- [18] R. Bryant and L. Hsu. Rigidity of integral curves of rank 2 distributions. *Inventiones mathematicae*, 114(2):435–462, 1993.
- [19] R. Cooke and V.I. Arnold. Ordinary Differential Equations. Springer Textbook. Springer Berlin Heidelberg, 1992.
- [20] Enrico Le Donne and Terhi Moisala. Semigenerated step-3 carnot algebras and applications to sub-riemannian perimeter, 2020.
- [21] Enrico Le Donne and Francesca Tripaldi. A cornucopia of carnot groups in low dimensions, 2020.
- [22] Eero Hakavuori and Enrico Le Donne. Blowups and blowdowns of geodesics in carnot groups. Journal of Differential Geometry, 123, 02 2023.
- [23] Velimir Jurdjevic. Geometric Control Theory. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1996.
- [24] L.D. Landau and E.M. Lifshitz. Mechanics third edition: Volume 1 of course of theoretical physics. 1976.

- [25] Andrzej J. Maciejewski and Maria Przybylska. Darboux points and integrability of Hamiltonian systems with homogeneous polynomial potential. *Journal of Mathematical Physics*, 46(6):062901, 05 2005.
- [26] Jerrold Marsden and Alan Weinstein. Reduction of symplectic manifolds with symmetry. Reports on Mathematical Physics, 5, 1974.
- [27] F. Monroy-Pérez and A. Anzaldo-Meneses. Optimal control on nilpotent lie groups. Journal of Dynamical and Control Systems, 8:487–504, 2002.
- [28] R. Montgomery. A tour of subriemannian geometries, their geodesics and applications. Number 91. American Mathematical Soc.
- [29] R. Montgomery, M. Shapiro, and A. Stolin. Chaotic geodesics in carnot groups. preprint on webpage at https://arxiv.org/abs/dg-ga/9704013, 1997.
- [30] Juan-Pablo Ortega and Tudor Ratiu. Momentum Maps and Hamiltonian Reduction. 01 2004.
- [31] Ron Perline and Sergei Tabachnikov. Geometry of integrable linkages, 2023.
- [32] Derek J.S. Robinson. A Course in the Theory of Groups. Springer, 2 edition, 1995.
 [33] Joel Storch. A First Course in the Calculus of Variations [Bookshelf], volume 36. 10
- [55] Joel Storch. A First Course in the Calculus of Variations [Booksneif], volume 50, 10 2016.
- [34] G. Teschl. Ordinary Differential Equations and Dynamical Systems. Graduate studies in mathematics. American Mathematical Soc.

Alejandro Bravo: Department of Mathematic University of Michigan, Ann Arbor, MI 48109, U.S.

Email address: Abravodo@umich.edu