Is every triangle a trajectory of an elliptical billiard?

Vladimir Dragović^{1,3} and Milena Radnović^{2,3}

¹The University of Texas at Dallas, Department of Mathematical Sciences ²The University of Sydney, School of Mathematics and Statistics ³Mathematical Institute SANU, Belgrade vladimir.dragovic@utdallas.edu, milena.radnovic@sydney.edu.au

Abstract

Using Marden's Theorem from geometric theory of polynomials, we show that for every triangle there is a unique ellipse such that the triangle is a billiard trajectory within that ellipse. Since 3-periodic trajectories of billiards within ellipses are examples of the Poncelet polygons, our considerations provide a new insight into the relationship between Marden's Theorem and the Poncelet Porism, two gems of exceptional classical beauty. We also show that every parallelogram is a billiard trajectory within a unique ellipse. We prove a similar result for the self-intersecting polygonal lines consisting of two pairs of congruent sides, named "Darboux butterflies". In each of three considered cases, we effectively calculate the foci of the boundary ellipses.

Keywords: Elliptical billiards; Marden's Theorem; 3-periodic billiard trajectories; 4-periodic billiard trajectories; Darboux butterflies; conics

AMS subclass: 51N20, 30C15, 37J35, 70H06

Contents

1	Introduction	2
2	Triangles 2.1 Triangular billiards	
3	Conics 3.1 Elliptical billiards	
4	The Siebeck–Marden theorem	8
5	3-periodic trajectories of billiards within ellipses	11
6	Convex 4-periodic trajectories of billiards within ellipses 6.1 Is every parallelogram a trajectory of an elliptical billiard?	11 11 13
7	Nonconvex 4-periodic elliptical billiard trajectories 7.1 How do nonconvex 4-periodic trajectories look like?	

1 Introduction

Recall that mathematical billiard in a planar domain is a dynamical system where a particle moves without constraints within the domain, and obeys the billiard reflection law when it hits the boundary [KT1991, Tab2005]. Thus, billiard trajectories are polygonal lines with vertices at the boundary, such that two consecutive sides form congruent angles with the tangent line to the boundary at their joint vertex, see Figure 1. Mathematical billiards are idealized models in many aspects: a usual billiard




Figure 1: Billiard reflection law: the angle of incidence with the tangent line at the bouncing point on the boundary equals the angle of reflection.

ball is replaced by a material point, the friction and spin are neglected. Such models have natural applications, for example in geometric optics. Here, we assume that the billiard particle is of the unit mass and it moves under the inertia between the impacts, i.e. uniformly along straight lines.

Billiards within ellipses have been intensively studied, see for example [Bol1990, Koz2003, KT1991, DR2011, DR2014, ADSK2016, KS2018, BM2017, BM2022, Glu2021, CZ2023, GRK2021, GKR2023, DGR2022, FV2023] and references therein.

In this note, we will give the affirmative answer to the title questions. The main ingredients in our considerations of triangular trajectories are Marden's theorem from the geometry of polynomials [Mar1966] and the classical Ceva Theorem from elementary geometry. Periodic trajectories of billiards within ellipses can be seen as an instance of so-called Poncelet polygons, which are closed polygonal lines inscribed in one conic an circumscribed about the other conic, see e.g. [DR2011]. The Poncelet Theorem from projective geometry of conics states that if for a given pair of conics there is one Poncelet polygon, then there are infinitely many such polygons and all have the same number of sides. Thus, we are in this work dealing with an interaction between Marden's Theorem and the Poncelet Theorem, which are both recognized by their exceptional classical beauty. Previously, a strong relationship between these two theorems was observed in [Dra2011] in a different context.

This paper is organised as follows. We review basic facts about billiards within triangles and the Ceva Theorem in Section 2, and billiards within ellipses in Section 3.1. We recall Marden's Theorem in Section 4. In Section 5, we prove that each triangle is a billiard trajectory within a unique ellipse. For the proof that all convex 4-periodic elliptical billiard trajectories are parallelograms, see Section 6.1, where we also show the the converse statement, see Theorem 6.3. All nonconvex 4-periodic elliptical billiard trajectories are described in Section 7.1 as so-called Darboux butterflies. The paper is concluded by Section 7.2, where we show that every Darboux butterfly is a billiard trajectory within a unique ellipse, see Theorem 7.12. In each of the three cases, of a triangle, a parallelogram, and a Darboux butterfly, we effectively calculate the foci of the boundary ellipse.

2 Triangles

In this section consists of two parts: in Subsection 2.1, we review main properties of the billiards within a triangle in the Euclidean plane, and in Subsection 2.2 we revisit the Ceva theorem.

2.1 Triangular billiards

A trajectory of such a billiard is a polygonal line, finite or infinite, with vertices on the sides of the triangle, such that consecutive edges of the trajectory satisfy the billiard reflection law. i.e. they form the same angle with the side of the triangle which their common vertex lie on, see Figure 2.




Figure 2: Billiard motion in a triangle.

The reflection is not well defined in the vertices of the triangle, thus we omit from our consideration trajectories falling in a vertex.

Remark 2.1 Note the following minimization property of the billiard reflection: if X, Y are two points on the same side of the a given line ℓ , then the length XL + LY, for $L \in \ell$, will be the smallest when the segments XL and LY satisfy the billiard reflection law off ℓ , see Figure 3.




Figure 3: The minimization property of billiard reflection: the shortest path connecting points X and Y which visits line ℓ is the billiard path XLY. Its length equals the segment XY', where Y' is symmetric to Y with respect to ℓ . For any other point $L' \in \ell$, we have XL' + L'Y = XL' + L'Y' > XY'.

One of the first natural questions for any dynamical system is to establish if its periodic trajectories exist and find them if they do. Thus, next we recall a proof of the existence a periodic trajectory within any acute triangle.

Theorem 2.2 (see e.g. Theorem 2.2 [DR2011]) Let $\triangle ABC$ be an acute triangle, and K, L, M the feet of its altitudes. Then $\triangle KLM$ is the triangle with minimal perimeter inscribed in $\triangle ABC$ and it represents a unique 3-periodic trajectory of the billiard with $\triangle ABC$.

Proof. Let M' be an arbitrary given point on side AB. In order to find points $K' \in BC$, $L' \in AC$ such that the triangle K'L'M' has the minimal perimeter, denote by M_1 , M_2 the points symmetric to M' with respect to sides BC and AC respectively. Then K' and L' are intersection points of M_1M_2 with sides BC and AC respectively, see Figure 4.




Figure 4: For a fixed point M' on AB, we construct points K', L' on the remaining two sides of the triangle, such that $\triangle K'L'M'$ has smallest possible perimeter.

The perimeter of $\triangle K'L'M'$ is equal to the length of segment M_1M_2 . We observe that M_1M_2 is a side of the isosceles triangle CM_1M_2 , whose angle $\angle M_1CM_2 = 2\angle BCA$ does not depend on the choice of point M'. Since $CM_1 \cong CM_2 \cong CM'$, the segment M_1M_2 is shortest when the point M' is such that CM' is the altitude of the triangle ABC from the vertex A, i.e. the minimal perimeter is attained when M' = M. It immediately follows that points K' and L', as constructed above, are then also the feet of the corresponding altitudes, see Figure 5.




Figure 5: The vertices of the triangle with smallest perimeter inscribed in $\triangle ABC$ are the feet K, L, M of the altitudes.

The minimizing property from Remark 2.1 implies that $\triangle KLM$, as the triangle with smallest perimeter inscribed in $\triangle ABC$ is a billiard trajectory within $\triangle ABC$.

Now, let us show uniqueness. Suppose that $K_1L_1M_1$ is another periodic trajectory within the triangle ABC, such that $K_1 \in BC$, $L_1 \in AC$, $M_1 \in AB$. Then, according to the billiard reflection law, we can denote: $k := \angle CK_1L_1 = \angle BK_1M_1$, $l := \angle CL_1K_1 = \angle AL_1M_1$, $m := \angle AM_1L_1 = \angle BM_1K_1$. Denoting by $\angle A$, $\angle B$, $\angle C$ the angles of the triangle ABC, we get that the sums of the angles in the triangles AL_1M_1 , BK_1M_1 , CK_1L_1 are:

$$\angle A + l + m = \angle B + k + m = \angle C + k + l = 180^{\circ}$$

which gives: $k = \angle A$, $l = \angle B$, $m = \angle C$.

From there, the sides of the triangle $K_1L_1M_1$ are parallel to the corresponding sides of the triangle KLM. The assumption that $K_1L_1M_1$ is a periodic billiard trajectory then implies $K_1 = K$, $L_1 = L$, $M_1 = M$, as shown in Figure 6. Namely, if K_1 is between K and C, then M_1 lies in the interior of $\triangle ABC$. This follows from the similarity of triangles KLM and $K_1L_1M_1$, by using the Thales theorem. The same argument shows that M_1 lies outside $\triangle ABC$ if K_1 is between K and $K_1L_1M_2$.

The previous proof essentially relies on the assumption that $\triangle ABC$ is acute, since otherwise some of the feet of the altitudes are not inner points of the sides. Thus, additional discussion in needed for




Figure 6: The trajectories with segments parallel to the segments of KLM are 6-periodic.

the right and obtuse triangles.

It is easy to see that there are periodic billiard trajectories within a right triangle: one of them, the polygonal line KLMNMLK, is shown in Figure 7.




Figure 7: A periodic trajectory of the billiard within a right triangle.

After such elementary considerations for acute and right triangles, one can stay amazed by the fact that it is not known if billiards within general obtuse triangles have any periodic trajectories! There are examples for some special cases and also an intriguing computer-based proof for the existence of periodic billiard trajectories when the obtuse angle does not exceed 100°, see [Sch2006, Sch2009].

2.2 Ceva's theorem

Now we will formulate the classical Ceva Theorem and one of its applications.

Theorem 2.3 (Ceva's Theorem) Let $\triangle ABC$ be a given triangle and O point not belonging to the lines AB, BC, AC. If the lines AO, BO, CO meet the lines BC, AC, AB at K, L, M respectively, then

$$\frac{AM}{MB} \cdot \frac{BK}{KC} \cdot \frac{CL}{LA} = 1. \tag{2.1}$$

In Ceva's Theorem, the signed lengths of segments are used in (2.1). For example, the quantity AM/MB is positive when M is between A and B and negative otherwise.

Notice that Ceva's theorem can be applied to an acute triangle $\triangle ABC$ and its 3-periodic billiard trajectory KLM, since the altitudes intersect at one point, see Figure 5 and Theorem 2.2.

We will use Ceva's theorem to prove the following statement.

Lemma 2.4 Let ABC be a given triangle and K, L, M its feet of the altitudes from A, B, C respectively. Then there exist real numbers m_1 , m_2 , m_3 such that point K divides BC in the ratio $m_2:m_3$, point L divides CA in the ratio $m_3:m_1$, and M divides AB in the ratio $m_1:m_2$. Moreover

$$m_1: m_2: m_3 = (b^2 + c^2 - a^2): (a^2 + c^2 - b^2): (a^2 + b^2 - c^2),$$
 (2.2)

where a, b, c respectively denote the lengths of the sides BC, AB, AC.

Proof. Since the altitudes of a triangle intersect at the orthocenter, Ceva's theorem implies the existence of numbers m_1 , m_2 , m_3 . The similarities of right triangles $\triangle KBA \sim \triangle MBC$, $\triangle LCB \sim \triangle KCA$, $\triangle MAC \sim \triangle LAB$ imply the following relations: KB/AB = MB/BC, CL/CB = CK/AC, AM/AC = LA/AB, yielding

$$m_1 : m_2 : m_3 = (AL \cdot AC) : (BM \cdot BA) : (CK \cdot CB).$$

The Pythagorean theorem gives

$$CK^2 = b^2 - AK^2$$
, $AL^2 = c^2 - BL^2$, $BM^2 = a^2 - CM^2$,

thus

$$m_1: m_2: m_3 = b\sqrt{c^2 - BL^2}: c\sqrt{a^2 - CM^2}: a\sqrt{b^2 - AK^2}$$

Observing that $a \cdot AK = b \cdot BL = c \cdot CM = 2A$, where A is the area of the triangle, we get:

$$m_1: m_2: m_3 = \sqrt{c^2b^2 - 4A^2}: \sqrt{a^2c^2 - 4A^2}: \sqrt{a^2b^2 - 4A^2}.$$

Finally, applying the Heron formula $\mathcal{A} = \sqrt{s(s-a)(s-b)(s-c)}$, with s = (a+b+c)/2, and simplifying, we get (2.2).

We conclude this section with the formulation of the converse Ceva's theorem.

Theorem 2.5 (Converse Ceva's Theorem) If points K, L, and M are chosen on the lines BC, AC and AB respectively such that the relation (2.1) is satisfied then either AK, BL and CM intersect at one point or all three are parallel.

3 Conics

3.1 Elliptical billiards

In this section, we review most important properties of billiards within an ellipse. The most famous one is the focal property of elliptic billiards, which is illustrated in Figure 8.

Proposition 3.1 (First focal property of ellipses) Let \mathcal{E} be an ellipse with foci F_1, F_2 and $A \in \mathcal{E}$ an arbitrary point. Then segments AF_1 , AF_2 satisfy the billiard reflection law off \mathcal{E} .




Figure 8: First focal property of ellipses.

If the billiard particle is launched from one focus of the ellipse, the focal property implies that the segments of the trajectory alternately contain the foci F_1 and F_2 . Next proposition characterizes the trajectories which do not contain the foci.

Proposition 3.2 Suppose that the billiard particle is traveling along segment s which does not contain any focus F_1 , F_2 of the boundary ellipse \mathcal{E} . Then we have:





Figure 9: Two segments satisfying the billiard reflection law of ellipse \mathcal{E} are tangent to the same conic, which is confocal with \mathcal{E} .

- If both foci of \mathcal{E} are on the same side of s, then there is unique ellipse \mathcal{E}' sharing the same foci with \mathcal{E} and touching the segment. Moreover, after reflection off the boundary, the next segment of the billiard trajectory will also be tangent to \mathcal{E}' , see Figure 9;
- If s crosses the segment F_1F_2 , then there is a hyperbola \mathcal{H} sharing the same foci with \mathcal{E} and touching the line containing s. Moreover, after reflection off the boundary, the next segment of the billiard trajectory will also be tangent to \mathcal{H} .

Proposition 3.2 implies the following key property of elliptical billiards: each trajectory of the billiard within ellipse has a caustic – a curve such that each segment of the trajectory lies on a tangent line of that curve, see Figure 10. For elliptical billiards, caustics are ellipses hyperbolas confocal with the boundary, including degenerate ones, which can be identified as horizontal and vertical axes of the ellipse \mathcal{E} .






Figure 10: The caustics of billiard trajectories.

A direct consequence of Propositions 3.1 and 3.2 is the well-known second focal property of ellipses, illustrated in Figure 11.

Proposition 3.3 (Second focal property of ellipses) Let \mathcal{E} be an ellipse with foci F_1, F_2 and A an arbitrary point outside ellipse \mathcal{E} . Denote by t_1 and t_2 the two tangents from A to ellipse \mathcal{E} . Then the angle between t_1 and the segment AF_1 is equal to the angle between AF_2 and t_2 .

In the next lemma, we note that the type of the caustic may be determined by the period of a trajectory.

Lemma 3.4 A periodic trajectory of the elliptic billiard with odd period has an ellipse as the caustic.

Proof. Suppose that the caustic of a given trajectory is a hyperbola. Denote by F_1 , F_2 the focal points of the boundary ellipse. Then every segment of the billiard trajectory intersects the segment F_1F_2 . Thus, for a periodic trajectory there will be an even number of intersections of the trajectory with F_1F_2 . Consequently, the period of a closed trajectory with hyperbola as caustic must be even.




Figure 11: Second focal property of ellipses

3.2 Pascal's theorem

We review here the classical Pascal's theorem.

Theorem 3.5 (Pascal Theorem, see e.g. [DR2011], **Section 4.4)** Let C be a non-degenerate conic. Consider six points $M, N, O, P, Q, R \in C$. Then the points $D = l_{MN} \cap l_{PQ}$, $E = \ell_{NO} \cap \ell_{QR}$ and $F = \ell_{OP} \cap \ell_{RM}$ are collinear.

Corollary 3.6 Let a non-degenerate conic C be given together with its four points M, N, P, Q. The intersection of the tangents to the conic at M and P is collinear with the points of intersection $\ell_{MN} \cap \ell_{PQ}$ and $\ell_{NP} \cap \ell_{MQ}$.

Corollary 3.7 (Simpson's line) Given three points M, N, P on a non-degenerate conic C. Then the intersections of the tangents to the conic at M, N, P with the opposite sides of the triangle MNP are collinear.

The classical Pappus theorem is a version of Pascal's theorem for degenerate conics.

4 The Siebeck-Marden theorem

Marden's theorem is one of the fundamental results in geometric theory of polynomials and rational functions. That theorem has a long history which is well described in the famous Marden's book [Mar1966]. The earliest version of this theorem, up to our best knowledge, goes back to 1864 when Siebeck (see [Sie1864]) formulated and proved it for the case of polynomials with simple roots. Here we consider only the case n = 3.

Theorem 4.1 (Siebeck, [Sie1864], n = 3) Let P(z) be a polynomial of degree 3 with complex coefficients, such that its zeros α_1 , α_2 , α_3 are simple and noncollinear. Then there exists an ellipse \mathcal{E} tangent to every line segment $[\alpha_i, \alpha_j]$ at the midpoint, see Figure 12. Moreover, the foci of the curve \mathcal{E} are zeros of the derivative polynomial P'(z).

We note that the ellipse which touches the midpoints of the triangle sides is the Steiner ellipse, i.e. the ellipse of the maximal area inscribed in that triangle. It is interesting to note that the ratio of areas of any triangle and its Steiner ellipse equals $3\sqrt{3}/(4\pi)$, see e.g. [GSO2020].

The nontrivial and interesting result of Siebeck still attracts a lot of attention, see for example [Kal2008, Dra2011]. In [Pra2004], this theorem is attributed to van den Berg, and two proofs are presented: the original proof of van der Berg from 1888 [vdB1888] and another one from [Sch1982]. We note that the main points of the proof of Siebeck theorem for n=3 from [Kal2008] are based on Propositions 3.1 and 3.2, which indicates a deep relationship with elliptic billiards. We are going to exploit that connection in next Section 5.

Siebeck's theorem can be extended to the cases with non-simple roots, namely to the functions of the form:

$$P(z) = (z - \alpha_1)^{m_1} (z - \alpha_2)^{m_2} (z - \alpha_3)^{m_3}.$$
 (4.1)




Figure 12: Siebeck theorem. If α_1 , α_2 , α_3 are zeros of a cubic polynomial P(z) and β_1 , β_2 zeros of its derivative P'(z), then there is an ellipse with foci β_1 , β_2 inscribed in triangle with vertices α_1 , α_2 , α_3 . Moreover, the ellipse is touching the sides at midpoints.

Each zero of the derivative P'(z) is then either equal to some α_i such that $m_i > 1$ or it is a zero of the logarithmic derivative of P:

$$F(z) = (\log P(z))' = \frac{m_1}{z - \alpha_1} + \frac{m_2}{z - \alpha_2} + \frac{m_3}{z - \alpha_3}.$$
 (4.2)

Notice that F(z) has two zeros. The following statement then holds, see also Figure 13.

Theorem 4.2 ([Mar1966, Theorem 4.2 for n = 3]) Consider a function P(z) of the form (4.1), such that α_1 , α_2 , α_3 are noncollinear complex numbers, and m_1 , m_2 , m_3 non-zero real constants. Let β_1 , β_2 are the zeros of the logarithmic derivative (4.2) of P(z). Then there is a conic with foci β_1 , β_2 touching each line of the segment $[\alpha_i, \alpha_j]$ in a point dividing that segment in the ratio $m_i : m_j$.




Figure 13: Marden theorem. An example for $m_1 = 1$, $m_2 = 2$, $m_3 = \sqrt{3}$.

Now we are going to expand on those results from the geometric theory of polynomials, in order to give complete characterization of ellipses inscribed in triangles.

Lemma 4.3 Suppose that F_1 is a point inside a triangle. Then there exists a unique ellipse inscribed in the triangle, such that one of its foci is point F_1 .

Proof. Denote the triangle by ABC. Denote by F_1' , F_1'' , F_1''' the points symmetric to F_1 with respect to sides BC, AC, AB respectively. Then the second focal point of the ellipse inscribed in $\triangle ABC$ is obtained as the intersection of bisectors of angles $F_1''AF_1'''$ and $F_1''CF_1'$. Namely, if F_2 is a second focal point, then triangle $CF_1''F_2$ is congruent to triangle $CF_1'F_2$. Thus, angle F_2CF_1'' is equal to angle F_2CF_1'' .

Theorem 4.4 Let ABC be a given triangle, and \mathcal{E} an ellipse inscribed in it, touching the sides BC, AC, AB in points K, L, M respectively. Then the lines AK, BL, CM are concurrent. (See Figure 14.)




Figure 14: The lines connecting triangle vertices with the points where the inscribed ellipse touches the opposite sides are concurrent.

Proof. Let φ be an affine transformation that maps the ellipse \mathcal{E} to a circle \mathcal{E}' . That circle is inscribed in triangle $\varphi(ABC)$.

Since segments of each pair $\varphi(A)\varphi(L)$ and $\varphi(A)\varphi(M)$, $\varphi(B)\varphi(K)$ and $\varphi(B)\varphi(M)$, $\varphi(C)\varphi(K)$ and $\varphi(C)\varphi(L)$ are of equal lengths as tangent segments to the circle from the same vertex, the converse Ceva's theorem (Theorem 2.5) gives that lines $\varphi(AK)$, $\varphi(BL)$, $\varphi(CL)$ are concurrent.

The statement follows immediately. \Box

In order to align further this matter with the Marden theorem, we provide an alternative proof of the last theorem.

Proof. (The second proof of Theorem 4.4.) Let F_1 be one of the foci of \mathcal{E} . Denote the complex numbers corresponding to A, B, C, and F_1 by z_1 , z_2 , z_3 , and q respectively. We search for nonzero real numbers m_1 , m_2 , m_3 such that q is one of the zeros of the derivative of $(z-z_1)^{m_1}(z-z_2)^{m_2}(z-z_3)^{m_3}$. This imposes one complex relation on real numbers m_1 , m_2 , m_3 . That complex relation gives two real relations which uniquely determine three real numbers m_1 , m_2 , m_3 up to a nonzero real factor. According to Marden's theorem (Theorem 4.2), there is an ellipse \mathcal{E}_1 inscribed in triangle ABC which has F_1 as one of its foci, with the other focus corresponding to the second zero of the derivative. Since ellipses \mathcal{E} and \mathcal{E}_1 are both inscribed in $\triangle ABC$ and both have F_1 as one of their foci, Lemma 4.3 implies that they coincide.

Theorem 4.5 Every ellipse inscribed in a triangle is a Marden ellipse, i.e. there exist positive real numbers m_1 , m_2 , m_3 such that the foci of the ellipse are the zeros of the logarithmic derivative of the

function $(z - \alpha_1)^{m_1}(z - \alpha_2)^{m_2}(z - \alpha_3)^{m_3}$, with α_1 , α_2 , α_3 being the complex numbers corresponding to the vertices of the triangle. The numbers m_1 , m_2 , m_3 are unique up to a non-zero factor.

Proof. Denote the triangle by ABC, the inscribed ellipse by \mathcal{E} , and by K, L, M the common points of \mathcal{E} with sides BC, AC, AB respectively. According to Theorem 4.4, the lines AK, BL, CM are concurrent, thus Ceva's theorem implies that (2.1) is satisfied. This relation determines positive numbers m_1 , m_2 , m_3 uniquely up to a non-zero factor.

5 3-periodic trajectories of billiards within ellipses

Now, after reviewing billiards and Marden's theory in previous sections, we are equipped and ready to address the title question.

Theorem 5.1 Every triangle is a 3-periodic trajectory of the billiard within an ellipse. That ellipse is uniquely determined.

Proof. Let an arbitrary triangle KLM be given. Through each of its vertices, we construct a line orthogonal to the bisector of the corresponding angle of the triangle. As a result, we obtain a new triangle ABC, formed by these three perpendiculars. Notice that $\triangle ABC$ is acute. By construction, KLM is a 3-periodic billiard trajectory within triangle ABC. According to Theorem 2.2, points K, L, M are the feet of the altitudes of triangle ABC, thus the three lines AK, BL, CM intersect at one point – the orthocenter of $\triangle ABC$. Thus, according to Ceva's Theorem (see Theorem 2.3), there exist nonzero real numbers m_1 , m_2 , m_3 such that point K divides BC in the ratio $m_2 : m_3$, point L divides CA in the ratio $m_3 : m_1$, and M divides AB in the ratio $m_1 : m_2$. The numbers m_1 , m_2 , m_3 can be calculated in terms of the lengths of the sides of triangle ABC, see Lemma 2.4 and relation (2.2).

Now, according to Marden's theorem (Theorem 4.2), there exists a conic \mathcal{E} which touches the sides AB, BC, CA respectively at M, K, L. Since K, L, M are inner points of those sides, \mathcal{E} must be an ellipse inscribed in $\triangle ABC$, see Figure 13. By construction, KLM is a billiard trajectory within ellipse \mathcal{E} .

6 Convex 4-periodic trajectories of billiards within ellipses

6.1 Is every parallelogram a trajectory of an elliptical billiard?

Let us recall that every 4-periodic convex elliptical billiard trajectory is a parallelogram, see, for example, [DR2019, Theorem 4]. In this section, in Theorem 6.3, we will show that being a convex 4-periodic elliptical billiard trajectory represents a complete characterization of parallelograms.

We start by formulating the following useful statement.

Lemma 6.1 Let a 4-periodic convex billiard trajectory within an ellipse be given, see Figure 15. Then we have:

- the tangent lines to the ellipse at the reflection points form a rectangle;
- the parallelogram is also a closed billiard trajectory within the rectangle;
- the diagonals of the rectangle are parallel to the sides of the parallelogram.

Proof. Follows by straightforward calculation of the angles and application of the billiard reflection law. \Box

Lemma 6.2 Any parallelogram is a closed billiard trajectory within a rectangle. Moreover, such a rectangle is uniquely determined for each parallelogram.




Figure 15: A convex 4-periodic trajectory within ellipse and the tangent lines at the points of reflection.

Theorem 6.3 Any parallelogram is a closed billiard trajectory within an ellipse. Moreover, such an ellipse is uniquely determined for a given parallelogram.

We immediately provide a compact, synthetic proof of this theorem, based on the Pascal theorem (see Theorem 3.5). An alternative, analytic proof is given in Section 6.2.

Proof. Let EFGH be a given parallelogram. According to Lemma 6.2, there is a unique rectangle within which EFGH is a closed billiard trajectory, see Figure 16.




Figure 16: A 4-periodic billiard trajectory within a rectangle.

Denote by a the side of the rectangle containing point E and by C the unique conic circumscribed about EFGH such that a is its tangent line. We want to prove that the conic is touching the remaining sides of the rectangle exactly at the points F, G, H.

Applying Pascal's theorem to \mathcal{C} and its points E, E, F, F, G, H, (see Corollary 3.6) we get that the intersection points of the following pairs of lines are collinear: a and FG; EF and GH; b' and AD, where b' is the tangent line to \mathcal{C} at F. Denote $X = a \cap FG$, $Y = b' \cap EH$. Since EF and GH are parallel, this implies that the line XY also must be parallel to them, thus Y is the intersection point of EH with the line which is parallel to EF and contains X.

Parallelogram EFXY is rhombus, since EF = FX, so its diagonals are orthogonal to each other, which implies that $Y \in b$, where b is the side of the rectangle containing point F. Thus b = b'.

Similarly, we can prove that the remaining two sides of the rectangle are also tangent to \mathcal{C} .

It remains to show that C is an ellipse. Each degenerate conic which contains points E, F, G, H is intersecting transversely the sides of the rectangle, thus C must be non-degenerate. No two tangent

lines of a parabola are parallel, thus C is either hyperbola or ellipse. For a hyperbola, we can notice that a pair of parallel lines cannot touch the same branch, and that the touching points of the lines containing sides of the rectangle would be placed on the extensions of those sides, which does not correspond to the geometric arrangement that we got here.

We conclude that C is the unique ellipse within which parallelogram EFGH is a closed billiard trajectory.

6.2 An analytic proof of Theorem 6.3

The following four lemmas and the theorem constitute an analytic proof of Theorem 6.3.

Lemma 6.4 Suppose that a rectangle and a point on one of its sides are given. Then there is a unique 4-periodic billiard trajectory within the rectangle such that the given point is one of its vertices.

Proof. Follows from the fact that the segments of any 4-periodic billiard trajectory within a rectangle are parallel to its diagonals, see Figure 16. \Box

Lemma 6.5 Consider the rectangle determined by A, E, H and denote it by AEO_1H . It is similar to the rectangle ABCD. Let O be the center of the rectangle ABCD. The points A, O_1 and O are collinear.

Lemma 6.6 Let F_1, F_2 be the foci of an ellipse tangent to AB at E and to AD at H. Assume the foci are symmetric with respect to O. Denote by F'_1 the point symmetric to F_1 w.r.t. AB and by F''_1 the point symmetric to F_1 w.r.t. AD. Denote by X and Y the projections of F_1 to AB and AD respectively. Then:

$$F_1'E||OX(=\frac{1}{2}F_1'F_2), \quad F_1''H||OY(=\frac{1}{2}F_1''F_2)$$
 (6.1)

and OX = OY.

Note that $d = F_1E + F_2E = F_1'F_2 = F_1''F_2 = 2OX = 2OY$, where d is the defining "rope length" for the ellipse \mathcal{E} .

Lemma 6.7 Let Y_1 be the intersection of EO_1 and XO and let X_1 be the intersection of HO_1 and YO. Then AEY_1Y and AXX_1H are rectangles.

Proof. The proof follows from the fact that $XF_1'EY_1$ and $F_1''YX_1H$ are parallelograms. Thus, $F_1'X = EY_1$ and $F_1''Y = HX_1$. We get $EY_1 = XF_1$ and $HH_1 = YF_1$.

Theorem 6.8 Given a right angle HAE. Denote by O_1 the fourth vertex of the rectangle HAEO₁. Let O be a point of the ray AO_1 . Then, there exists a unique (up to the symmetry w.r.t. O) point F_1 such that its mirror images F'_1 and F''_1 w.r.t. AE and AH respectively, satisfy the following

$$F_1'E||OX, \quad F_1''H||OY, \quad OX = OY,$$
 (6.2)

where $\{X\} = AE \cap F_1F_1'$ and $\{Y\} = AH \cap F_1F_1''$.

Proof. Denote by O_x and O_y the projections of O at rays AE and AH respectively. Denote the side lengths as follows

$$AX = x, AE = e, AO_x = o_x, AY = y, AH = h, AO_y = o_y.$$

Here we assume the values e, h, o_x, o_y to be known, while x and y need to be determined. The condition OX = OY is equivalent to

$$(o_x - x)^2 - (o_y - y)^2 = o_x^2 - o_y^2, (6.3)$$

by using the Pythagorean theorem. Recall that by X_1 we denoted the intersection of HO_1 and YO. According to Lemma 6.7, AXX_1H is a rectangle. Let $\{X_2\} = XX_1 \cap O_yO$. Thus, AXX_2O_y is a rectangle as well. From the similarity $\triangle OO_yY \sim \triangle OX_2X_1$ we get

$$(o_x - x)(o_y - y) = o_x(o_y - h). (6.4)$$

We use the notation $\mathcal{X} = o_x - x$ and $\mathcal{Y} = o_y - y$. From equations (6.3) and (6.4), we get the following biquadratic equation

$$\mathcal{X}^4 - (o_x^2 - o_y^2)\mathcal{X}^2 - o_x^2(o_y - h)^2 = 0.$$
(6.5)

Let us study the corresponding quadratic equation

$$Q: \hat{X}^2 - (o_x^2 - o_y^2)\hat{X} - o_x^2(o_y - h)^2 = 0.$$
(6.6)

The discriminant of (6.6) is

$$\mathcal{D}_Q = (o_x^2 - o_y^2)^2 + 4o_x^2(o_y - h)^2.$$

The discriminant is positive, except in the following cases:

$$o_x = o_y = h, (6.7)$$

when $\mathcal{D}_Q = 0$. Thus, quadratic equation (6.6) always has two real solutions \hat{X}_1, \hat{X}_2 . Their product

$$\hat{X}_1 \cdot \hat{X}_2 = -o_x^2 (o_y - h)^2 < 0, \quad o_y \neq h$$

is negative, unless $o_y = h$. Thus, for $o_y \neq h$, exactly one of the solutions is positive, say \hat{X}_1 . Thus,

$$(o_x - x)^2 = \hat{X}_1.$$

From the last equation, there are two options for x: $x_1 = o_x - \sqrt{\hat{X}_1}$ and $x_2 = o_x + \sqrt{\hat{X}_1}$. One of them gives F_1 and another one F_2 . Hence, F_1 and F_2 are two points symmetric with respect to O, the only two points that satisfy the conditions of the theorem.

The exceptional case (6.7) corresponds to the situation in which ABCD is a square, with H being the midpoint of AD. In this case the ellipse \mathcal{E} is a circle and $F_1 = F_2 = O$.

In the case $o_y = h$ and $o_x^2 > o_y^2$, we again get a unique positive solution \hat{X}_1 .

Thus, we have concluded the analytic proof of Theorem 6.3. As an outcome, we get the exact formulas for the foci of the ellipse.

7 Nonconvex 4-periodic elliptical billiard trajectories

7.1 How do nonconvex 4-periodic trajectories look like?

Following [BLPT2020, Definition 3.4], we introduce the notion of a Darboux butterfly.

Definition 7.1 Any polygonal self-intersecting line consisting of four sides, such that the first and the third are congruent as well as the second and the fourth side is called a Darboux butterfly (or a bow tie).

Lemma 7.2 Every Darboux butterfly consists of legs and diagonals of an isosceles trapezoid.

Proof. Let GHKL be a Darboux butterfly. Then $\triangle GHK$ is congruent to $\triangle KLG$ and they are placed from the same side of the line GL. Thus, HL is parallel to GL, and GH is congruent to KL.

Lemma 7.3 Every 4-periodic elliptical billiard trajectory, having a hyperbola as a caustic, is a Darboux butterfly.

Proof. We know that every 4-periodic convex elliptical billiard trajectory is a parallelogram, see, for example, [DR2019, Theorem 4]. From [Sta2021], Theorem 1, it follows that for each 4-periodic trajectory of an elliptical billiard, having a hyperbola as the caustic, there exists a 4-periodic elliptical billiard trajectory having an ellipse as the caustic such that the corresponding sides of the two trajectories are congruent.

Thus, we come to the following question.

7.2 Is every Darboux butterfly an elliptical billiard trajectory?

We start by showing that every Darboux butterfly is a billiard trajectory within a kite.

Lemma 7.4 Every Darboux butterfly is a billiard trajectory in a unique kite, consisting of two congruent acute triangles.

Proof. Consider a Darboux butterfly GHKL. Denote by S the intersection of GL and HK. Let the bisector of $\angle HGS$ and the bisector of $\angle SHG$ intersect at O_1 . Construct the lines GB and HB orthogonal to GO_1 at G and to O_1H at H, respectively. The angle $\angle GO_1H$ is obtuse since $\angle O_1GH + \angle O_1HG < \pi/2$. Since in the quadrilateral BHO_1G the angles at G and H are right, we get that $\angle GBH < \pi/2$.

Let the bisector of $\angle SKL$ and the bisector of $\angle SLK$ intersect at O_2 . We construct the lines KD and LD orthogonal to KO_2 at K and to LO_2 at L respectively. Denoting the intersection of BG and DK by A and denoting the intersection of BH and DL by C, we get a quadrilateral ABCD, which is a kite. Polygonal line GHKL is a billiard trajectory within this kite.

Lemma 7.5 Every kite, consisting of two congruent acute triangles, has a unique Darboux butterfly 4-periodic billiard trajectory inscribed in it. The vertices of this trajectory are the footings of the altitudes of each of these two triangles at the sides of the kite. This 4-periodic billiard trajectory is obtained as the concatenation of two 3-periodic billiard trajectories, each of the two being the unique 3-periodic trajectory inscribed in each of the two acute triangles.

Proof. Consider a kite ABCD with acute triangles ABC and ACD. Let GHKL be a Darboux butterfly 4-periodic billiard trajectory in it. Let S be the intersection of GL and HK. Let O_1S be the bisector of $\angle GSH$ and O_2S the bisector of $\angle KSL$. Then O_1, S, O_2 are collinear. Construct the line ℓ orthogonal to O_1O_2 at S. Let ℓ intersect BA and BC at A_1 and C_1 , respectively. Similarly, Let ℓ intersect DA and DC at A_2 and C_2 , respectively. Since $\triangle GSH \cong \triangle KLS$, the triangles BA_1C_1 and DA_2C_2 are congruent and symmetric w.r.t. ℓ . Thus, ℓ coincides with AC and $S \in AC$.

Since $S \in AC$ and O_1O_2 is orthogonal to AC, we see that GSH and KSL are 3-periodic billiard trajectories of $\triangle ABC$ and $\triangle ADC$, respectively. Such 3-periodic trajectories are unique within any given acute triangle and their vertices are the footings of the altitudes of the triangle, see Theorem 2.2. Thus, G and H are the footings of the altitudes from A and C in $\triangle ABC$ and L and K are the footings of the altitudes from A and C in $\triangle ADC$.

As a small digression, we are going to show that not all 4-periodic billiard trajectories within kites are Darboux butterflies.

Lemma 7.6 Consider a kite ABCD such that the congruent triangles $\triangle ABC$ and $\triangle ADC$ are acute. Construct a line parallel to AC. Let A_1, A_2, C_1, C_2 be the intersections of that line with AB, AD, CB, and CD respectively. Construct the footings of the altitudes of $\triangle A_1B_1C_1$ from A_1 and C_1 , denoted as H and G, and the altitudes from C_2 and A_2 of $\triangle DC_2A_2$, denoted K and L. Then, HGKL is a billiard trajectory.

Lemma 7.7 If an isosceles trapezoid GKLH is given, then the orthogonal lines to the bisectors of its angles at the vertices form a kite ABCD consisting of two congruent right triangles ABC and ADC,

with right angles at B and D respectively. The trapezoid is a 4-periodic billiard trajectory within this kite.

Conversely, if a kite ABCD consisting of two congruent right triangles ABC and ADC is given, then it admits (infinitely many) 4-periodic billiard trajectories in the form of an isosceles trapezoid.

Theorem 7.8 Consider an acute triangle ABC. Let G and H be the footings of its altitudes from C and A, respectively. Then, there exists a unique ellipse \mathcal{E} with its foci being symmetric w.r.t. AC, which is tangent to AB and BC at G and H, respectively.

Lemma 7.9 Consider an acute triangle ABC. Let G and H be the footings of its altitudes from C and A respectively. Let \mathcal{E} be an ellipse with the foci being symmetric w.r.t. AC, which is tangent to AB and BC at G and H respectively. Let F_1 be the focal point inside $\triangle ABC$. Denote the projections of F_1 at AB, BC, and BA by O, Y, and X respectively. Denote by \bar{F}_1 " and \bar{F}_1' the projections of F_1 at the altitudes AG and CH respectively. Then, the points O, \bar{F}_1 ", Y are collinear as well as the points O, \bar{F}_1', X . Moreover OX = OY.

Proof. Let F'_1 and F_1 " be the points symmetric to F_1 w.r.t. AB and BC respectively. Then, $F_2F'_1||OX$ and $OX = F_2F'_1/2$. The points F'_1 , H, F_2 are collinear. The segment \bar{F}'_1H is parallel and congruent to XF'_1 , since \bar{F}'_1H and XF'_1 are the opposite sides of a rectangle. Thus, XF'_1 is parallel and congruent to \bar{F}_1H . We conclude that $XF_1H\bar{F}'_1$ is a parallelogram. Thus, XF'_1 is parallel to F'_1H . So, \bar{F}'_1 is collinear with X and X0. A proof that X1, X2, X3 are collinear and X4 follows along the same lines.

Since G and H belong to the same ellipse with the foci F_1, F_2 , we have:

$$OX = \frac{F_2 F_1'}{2} = \frac{F_2 F_1''}{2} = OY.$$

Proposition 7.10 Consider an acute triangle ABC. Let G and H be the footings of its altitudes from C and A, respectively. There exists a unique point F_1 inside $\triangle ABC$ such that its projections $O, X, Y, \bar{F}'_1, \bar{F}_1$ " at AC, AB, BC, CG, AH respectively, satisfy the following:

- (i) O, \bar{F}'_1, X are collinear;
- (ii) O, \bar{F}_1 ", Y are collinear:
- (iii) OX = OY.

Lemma 7.11 Consider a triangle ABC such that neither of the angles at A or C are right. Let G be the footing of its altitudes from C. All the points F such that the projections O, X, \bar{F}'_1 of F at AC, AB, CG respectively are collinear, belong to the circle with a diameter AC. Conversely, any point F which belongs to the circle with a diameter AC has the property that the projections O, X, \bar{F}'_1 of F at AC, AB, CG respectively are collinear.

Proof. We first prove Lemma 7.11. Without loss of generality, we may set A = (0,0), $B = (b_1, b_2)$ and C = (1,0). Denote the coordinates of $F = (f_1, f_2)$. A direct calculation shows that the collinearity condition for points O, X, \bar{F}'_1 is equivalent to:

$$CC1: (b_1-1)b_2(-f_1+f_1^2+f_2^2)=0.$$
 (7.1)

Since neither of the angles at A or C are right, $b_1 \neq 1$ and $b_2 \neq 0$. Thus, the condition CC1, (7.1), is equivalent to:

$$CC1': -f_1 + f_1^2 + f_2^2 = 0.$$
 (7.2)

The last relation (7.2) is equivalent to the condition that the point F belongs to the circle

$$\left(f_1 - \frac{1}{2}\right)^2 + f_2^2 = \frac{1}{4},$$

which is the circle with AC as a diameter.

It is interesting to note that the locus of points F from Lemma 7.11 does not depend on the vertex B.

Proof. Now we turn to a proof of Proposition 7.10. We will use the same notation as in the proof of Lemma 7.11. From Lemma 7.11, we see that conditions (i) and (ii) are equivalent and they reduce to (7.2). A direct calculation for condition (iii) gives

$$b_2^2((1-2b_1)f_2^2 + (1-b_1)f_1^2 + 2(b_1^2 + b_2^2)f_1 - (b_1^2 + b_2^2)) = 0. (7.3)$$

Observing that $b_2 \neq 0$ and substituting $f_2^2 = f_1 - f_1^2$ from (7.2) into (7.3), we get

$$f_1 = \frac{b_1^2 + b_2^2}{(1 - b_1)^2 + b_2^2}. (7.4)$$

Thus, $F_1 = (f_1, f_2)$ which satisfies conditions (i), (ii), and (iii) is uniquely determined through (7.4) and (7.2). This finishes the proof of Proposition 7.10.

This finishes the proof of Theorem 7.8. As a consequence, we get our final

Theorem 7.12 For every Darboux butterfly there exists a unique ellipse such that the Darboux butterfly is a billiard trajectory within that ellipse.

Proof. Given a Darboux butterfly GHKL, according to Lemma 7.4, consider the unique kite consisting of two congruent acute triangles ABC and ADC for which GHKL is a billiard trajectory. According to Lemma 7.5, the vertices of this trajectory are the footings of the altitudes of each of these two triangles at the sides of the kite. This 4-periodic billiard trajectory is obtained as the concatenation of two 3-periodic billiard trajectories, each of the two being the unique 3-periodic trajectory inscribed in each of the two acute triangles. Thus, G and H are the footings of the altitudes of the acute triangle ABC from C and A respectively. Then, according to Theorem 7.8, there exists a unique ellipse \mathcal{E} with foci being symmetric w.r.t. AC, which is tangent to AB and BC at G and G respectively. By symmetry, this ellipse \mathcal{E} is also tangent to G and G and G respectively, since G and G are the footings of the altitudes of the acute triangle G from G and G respectively. Thus, the Darboux butterfly G and G is a billiard trajectory within the ellipse G. The ellipse G is a unique conic containing four points G and G is a unique to G and G is a unique conic containing four points G and G is a unique to G is a unique conic containing

Acknowledgements

The research was supported by the Australian Research Council, Discovery Project 190101838 Billiards within quadrics and beyond, the Science Fund of Serbia grant Integrability and Extremal Problems in Mechanics, Geometry and Combinatorics, MEGIC, Grant No. 7744592 and the Simons Foundation grant no. 854861.

References

[ADSK2016] A. Avila, J. De Simoi, and V. Kaloshin, An integrable deformation of an ellipse of small eccentricity is an ellipse, Ann. of Math. (2) **184** (2016), no. 2, 527–558, DOI 10.4007/annals.2016.184.2.5. MR3548532

[vdB1888] F. J. van der Berg, Nogmaals over afgeleide Wortelpunten, Nieuw Archiev voor Wiskunde 15 (1888), 100–164.

- [BM2017] M. Bialy and A. E. Mironov, Angular billiard and algebraic Birkhoff conjecture, Adv. Math. 313 (2017), 102–126, DOI 10.1016/j.aim.2017.04.001. MR3649222
- [BM2022] ______, The Birkhoff-Poritsky conjecture for centrally-symmetric billiard tables, Ann. of Math. (2) 196 (2022), no. 1, 389–413, DOI 10.4007/annals.2022.196.1.2. MR4429262
- [Bol1990] S. V. Bolotin, Integrable Birkhoff billiards, Vestnik Moskov. Univ. Ser. I Mat. Mekh. 2 (1990), 33-36.
- [BLPT2020] G. Bor, M. Levi, R. Perline, and S. Tabachnikov, Tire Tracks and Integrable Curve Evolution, Int. Math. Res. Not. IMRN 2020 (2020), no. 9, 2698–2768.
 - [CZ2023] P. Corvaja and U. Zannier, Finiteness theorems on elliptical billiards and a variant of the Dynamical Mordell-Lang Conjecture, Proceedings of the London Mathematical Society 127 (2023), no. 5, 1268-1337.
 - [Dra2011] V. Dragović, Poncelet-Darboux curves, their complete decomposition and Marden theorem, Int. Math. Res. Not. IMRN 15 (2011), 3502–3523.
 - [DR2011] V. Dragović and M. Radnović, Poncelet Porisms and Beyond, Springer Birkhauser, Basel, 2011.
 - [DR2014] _____, Pseudo-integrable billiards and arithmetic dynamics, Journal of Modern Dynamics 8 (2014), no. 1, 109–132.
 - [DR2019] V. Dragović and M. Radnović, Caustics of Poncelet polygons and classical extremal polynomials, Regul. Chaotic Dyn. 24 (2019), no. 1, 1–35.
- [DGR2022] V. Dragović, S. Gasiorek, and M. Radnović, Billiard ordered games and books, Regul. Chaotic Dyn. 27 (2022), no. 2, 132–150.
 - [FV2023] A. Fomenko and V. Vedyushkina, Billiards and Integrable Systems, Russian Mathematical Surveys 78 (2023), no. 5, 881–954.
- [GRK2021] R. Garcia, D. Reznik, and J. Koiller, New properties of triangular orbits in elliptic billiards, Amer. Math. Monthly 128 (2021), no. 10, 898–910, DOI 10.1080/00029890.2021.1982360. MR4343349
- [GKR2023] R. Garcia, J. Koiller, and D. Reznik, Loci of 3-periodics in an elliptic billiard: Why so many ellipses?, J. Symbolic Comput. 114 (2023), 336–358, DOI 10.1016/j.jsc.2022.06.001. MR4446194
- [GSO2020] G. Glaeser, H. Stachel, and B. Odehnal, The universe of quadrics, Springer, Berlin, 2020.
- [Glu2021] A. Glutsyuk, On polynomially integrable Birkhoff billiards on surfaces of constant curvature, J. Eur. Math. Soc. (JEMS) 23 (2021), no. 3, 995–1049, DOI 10.4171/jems/1027. MR4210728
- [Kal2008] D. Kalman, An Elementary Proof of Marden's Theorem, The American Mathematical Monthly 115 (2008), 330–337.
- [KS2018] V. Kaloshin and A. Sorrentino, On the local Birkhoff conjecture for convex billiards, Ann. of Math. (2) 188 (2018), no. 1, 315–380, DOI 10.4007/annals.2018.188.1.6. MR3815464
- [KT1991] V. Kozlov and D. Treshchëv, Billiards, Amer. Math. Soc., Providence RI, 1991.
- [Koz2003] V. V. Kozlov, Rationality conditions for the ratio of elliptic integrals and the great Poncelet theorem, Vestnik Moskov. Univ. Ser. I Mat. Mekh. **71** (2003), no. 4, 6–13 (Russian); English transl., Moscow Univ. Math. Bull. **58** (2004), no. 4, 1–7.
- [Mar1966] M. Marden, Geometry of Polynomials, 2nd ed., Math. Surveys, vol. 2, AMS, 1966.
- [Pra2004] V. V. Prasolov, *Polynomials*, Algorithms and Computation in Mathematics, vol. 11, Springer-Verlag, Berlin, 2004. Translated from the 2001 Russian second edition by Dimitry Leites.
- [Sch1982] I. J. Schoenberg, Mathematical time exposures, Mathematical Association of America, Washington, DC, 1982.
- [Sch2006] R. E. Schwartz, Obtuse triangular billiards. I. Near the (2,3,6) triangle, Experiment. Math. 15 (2006), no. 2, 161–182.
- [Sch2009] _____, Obtuse triangular billiards. II. One hundred degrees worth of periodic trajectories, Experiment. Math. 18 (2009), no. 2, 137–171.
- [Sie1864] J. Siebeck, Ueber eine neue analytische Behandlungweise der Brennpunkte, J. Reine Angew. Math. 64 (1864), 175–182.
- [Sta2021] H. Stachel, *Isometric Billiards in Ellipses and Focal Billiards in Ellipsoids*, Journal for Geometry and Graphics **25** (2021), no. 1, 97–118.
- [Tab2005] S. Tabachnikov, Geometry and billiards, Student Mathematical Library, vol. 30, American Mathematical Society, Providence, RI, 2005.