# Tensor Krylov subspace methods via the T-product for large Sylvester tensor equations

F. BOUYGHF<sup>a,c</sup>, M. EL GUIDE<sup>b</sup>, A. EL ICHI<sup>a,c</sup>

 <sup>a</sup> LabMIA-SI, University of Mohammed V Rabat, Morocco.
 <sup>b</sup> Africa Institute for Research in Economics and Social Sciences (AIRESS), FGSES, Mohammed VI Polytechnic University, Rabat, Morocco.
 <sup>c</sup> LMPA, University of the Littoral Opal Coast, Calais, France.

### Abstract

In the present paper, we introduce new tensor krylov subspace methods for solving large Sylvester tensor equations. The proposed method uses the well-known T-product for tensors and tensor subspaces. We introduce some new tensor products and the related algebraic properties. These new products will enable us to develop third-order the tensor FOM (tFOM), GMRES (tGMRES), tubal Block Arnoldi and the tensor tubal Block Arnoldi method to solve large Sylvester tensor equation. We give some properties related to these method and present some numerical experiments.

Keywords: Arnoldi, Krylov subspaces, Sylvester equations, Tensors, T-products.

#### 1. Introduction

The aim of this paper is to present numerical Tensor Krylov subspace methods for solving Sylvester tensor equation

$$\mathcal{M}(\mathfrak{X}) = \mathfrak{C},\tag{1}$$

where  $\mathcal{M}$  is a linear operator that could be described as

$$\mathcal{M}(\mathfrak{X}) = \mathcal{A} \star \mathfrak{X} - \mathfrak{X} \star \mathcal{B},\tag{2}$$

<sup>\*</sup>Corresponding author

*Email addresses:* fatimabouyghf3@gmail.com (F. BOUYGHF), mohamed.elguide@um6p.ma (M. EL GUIDE ), elichi.alaa@gmail.com (A. EL ICHI )

where  $\mathcal{A}$ ,  $\mathfrak{X}$ ,  $\mathcal{B}$  and  $\mathcal{C}$  are three-way tensors, leaving the specific dimensions to be defined later, and  $\star$  is the T-product introduced by Kilmer and Martin [17, 18].

Consider the following Sylvester matrix equation

$$AX + XB = C. (3)$$

In the existing body of literature, various methodologies have been proposed to address the solution of Sylvester matrix equations, as articulated in Equation (3). When confronted with matrices of relatively small dimensions, established direct methods, as advocated in seminal works such as [1] and [9], are often recommended. These direct methods leverage Schur decomposition to transform the original equation into a more amenable form, thereby facilitating resolution through forward substitution.

For larger Sylvester matrix equations, iterative projection techniques have been advanced, as evidenced by studies like [5], [13], and [22]. These methods employ Galerkin projection approaches, including both classical and block Arnoldi techniques. By employing such projection methods iteratively, lower-dimensional Sylvester matrix equations are derived, subsequently tackled through direct methods for efficient resolution. Notably, comprehensive approaches to Krylov subspace methods for solving linear systems are elucidated in [4], providing a unified perspective on these iterative methodologies.

It's essential to consider the nature and size of the matrices involved when choosing an appropriate solution strategy, with direct methods favored for smaller matrices and iterative projection methods preferred for handling larger Sylvester matrix equations. The utilization of Schur decomposition and Galerkin projection techniques underscores the adaptability and scalability of these methods across varying problem sizes.

This paper focuses on the development of efficient and robust iterative Krylov subspace methods using the T-product for solving the Sylvester tensor equation (STE) represented by (1). Specifically, when dealing with small-sized tensors in (1), our aim is to extend the matrixoriented direct methods outlined in [1] and [9] to third-order tensors, employing the T-product formalism. This extension leads to the formulation of the *t-Bartels-Stewart* algorithm.

For larger tensors, we introduce a novel method termed as orthogonal and oblique projection onto a tensor Krylov subspace. Two specific instances of this approach, namely the tensor Full Orthogonalization Method (tFOM) and the tensor Generalized Minimal Residual Method (tGMRES), are examined. Additionally, we present well-known tensor Tubal Block Krylov methods utilizing the T-product to transform the original large Sylvester equation into a lower-dimensional *STE*. In this context, we describe the Tubal Block Arnoldi (TBA) as a generalization of the block Arnoldi matrix.

This paper focuses on the development of efficient and robust iterative Krylov subspace methods using the T-product for solving the Sylvester tensor equation (STE) represented by (1). Specifically, when dealing with small-sized tensors in (1), our aim is to extend the matrixoriented direct methods outlined in [1] and [9] to third-order tensors, employing the T-product formalism. This extension leads to the formulation of the *t-Bartels-Stewart* algorithm.

For larger tensors, we introduce a novel method termed as orthogonal and oblique projection onto a tensor Krylov subspace. Two specific instances of this approach, namely the tensor Full Orthogonalization Method (tFOM) and the tensor Generalized Minimal Residual Method (tGMRES), are examined. Additionally, we present well-known tensor Tubal Block Krylov methods utilizing the T-product to transform the original large Sylvester equation into a lower-dimensional *STE*. In this context, we describe the Tubal Block Arnoldi (TBA) as a generalization of the block Arnoldi matrix.

This work contributes to the expansion of iterative methods for solving Sylvester tensor equations, catering to both small and large-sized tensors through the adaptation of established matrix-based techniques and the introduction of novel tensor projection approaches.

User reformulate :The paper is organized as follows: In Section 2, we give notations and definitions related to the T-product. More details of all notations and definitions given in this section are given in [6, 7, 8]. In Section 3, we develop the (tFOM) and (tGMRES) methods. The *t-Bartels-Stewart* method will be introduced in Section 4. After defining a tubal QR factorisation algorithm in Section 5, we will establish the tensor tubal-Block Arnoldi process (TBA) that allows us to introduce the tubal-Block Arnoldi for solving large Sylvester tensor equation (TBAS) method. Finally, some numerical tests are reported in Section 6.

The paper is structured as follows: In Section 2, we present notations and definitions pertaining to the T-product. Comprehensive details for all the notations and definitions introduced in this section can be found in [6, 7, 8]. Section 3 is dedicated to the development of the tensor Full Orthogonalization Method (tFOM) and the tensor Generalized Minimal Residual Method (tGMRES). In Section 4, we introduce the *t-Bartels-Stewart* method. Following this, Section 5 outlines the tubal QR factorization algorithm before establishing the Tensor Tubal-Block Arnoldi process (TBA). This process enables the introduction of the Tubal-Block Arnoldi for Solving Large Sylvester Tensor Equation (TBAS) method. Finally, Section 6 presents numerical tests to validate and assess the performance of the proposed methods.

#### 2. Notation and background

A tensor is a multidimensional array of data. The number of indices of a tensor is called modes or ways. Notice that a scalar can be regarded as a zero mode tensor, first mode tensors are vectors and matrices are second mode tensor. The order of a tensor is the dimensional of the array needed to represent it, also known as ways or modes. For a given N-mode (or order-N) tensor  $\mathcal{X} \in \mathbb{R}^{n_1 \times n_2 \times n_3 \dots \times n_N}$ , the notation  $x_{i_1,\dots,i_N}$  (with  $1 \leq i_j \leq n_j$  and  $j = 1, \dots, N$ ) stands for the element  $(i_1, \dots, i_N)$  of the tensor  $\mathcal{X}$ . The norm of a tensor  $\mathcal{A} \in \mathbb{R}^{n_1 \times n_2 \times \dots \times n_\ell}$  is specified by

$$\|\mathcal{A}\|_{F}^{2} = \sum_{i_{1}=1}^{n_{1}} \sum_{i_{2}=1}^{n_{2}} \cdots \sum_{i_{\ell}=1}^{n_{\ell}} a_{i_{1}i_{2}\cdots i_{\ell}}^{2}$$

Corresponding to a given tensor  $\mathcal{A} \in \mathbb{R}^{n_1 \times n_2 \times n_3 \dots \times n_N}$ , the notation

$$\underbrace{\mathfrak{X}_{\vdots \ldots \vdots k}}_{(N-1)\text{-times}} k , \text{ for } k = 1, 2, \ldots, n_N$$

denotes a tensor in  $\mathbb{R}^{n_1 \times n_2 \times n_3 \dots \times n_{N-1}}$  which is obtained by fixing the last index and is called frontal slice. Fibers are the higher-order analogue of matrix rows and columns. A fiber is defined by fixing all the indexes except one.

In this paper, a tensor is of third order, i. e., N=3, that will be denoted by the calligraphic script letters, say  $\mathcal{A} = [a_{ijk}]_{i,j,k=1}^{n_1,n_2,n_3}$ . We use capital letters to denote matrices, lower case letters to denote vectors, boldface lower case letters to denote tube fibers (tubal scalars or tubes) and boldface upper-case letters to denote block diagonal matrix. Using MATLAB notation,  $\mathcal{A}(:, j, k)$ ,  $\mathcal{A}(i, :, k)$  and  $\mathcal{A}(i, j, :)$  denote mode-1, mode-2, and mode-3 fibers, respectively. The notations  $\mathcal{A}(i, :, :)$ ,  $\mathcal{A}(:, j, k)$ , and  $\mathcal{A}(:, :, k)$  stand for the i-th horizontal, j-th lateral, and k-th frontal slices of  $\mathcal{A}$ , respectively. The j-th lateral slice is also denoted by  $\overrightarrow{\mathcal{A}}_j$ . Tt is a tensor of

size  $(n_1 \times 1 \times n_3)$  and will be referred to as a tensor column. Moreover, the k-th frontal slices of  $\mathcal{A}$  is a matrix size  $(n_1 \times n_2)$  denoted by  $\mathcal{A}^{(k)}$ .

## 2.1. Definitions and properties of the T-product

In this part, we briefly review some concepts and notations related to the T-product, see [3, 18, 17] for more details. Let  $\mathcal{A} \in \mathbb{R}^{n_1 \times n_2 \times n_3}$  be a third-order tensor, then the operations beirc, unfold and fold are defined by

$$\operatorname{bcirc}(\mathcal{A}) = \begin{pmatrix} \mathcal{A}^{(1)} & \mathcal{A}^{(n_3)} & \mathcal{A}^{(n_3-1)} & \dots & \mathcal{A}^{(2)} \\ \mathcal{A}^{(2)} & \mathcal{A}^{(1)} & \mathcal{A}^{(n_3)} & \dots & \mathcal{A}^{(3)} \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \mathcal{A}^{(n_3)} & \mathcal{A}^{(n_3-1)} & \ddots & \mathcal{A}^{(2)} & \mathcal{A}^{(1)} \end{pmatrix} \in \mathbb{R}^{n_1 n_3 \times n_2 n_3},$$
$$\operatorname{unfold}(\mathcal{A}) = \begin{pmatrix} \mathcal{A}^{(1)} \\ \mathcal{A}^{(2)} \\ \vdots \\ \mathcal{A}^{(n_3)} \end{pmatrix} \in \mathbb{R}^{n_1 n_3 \times n_2}, \quad \operatorname{fold}(\operatorname{unfold}(\mathcal{A})) = \mathcal{A}.$$

Let  $\widetilde{\mathcal{A}}$  be the tensor obtained by applying the discrete Fourier transform DFT matrix  $F_{n_3} \in \mathbb{C}^{n_3 \times n_3}$ ; (for more details about DFT matrix, see [12]) on all the 3-mode tubes of the tensor  $\mathcal{A}$ . With the Matlab command fft, we have

$$\widetilde{\mathcal{A}} = \mathtt{fft}(\mathcal{A}, [], 3), \text{ and } \mathtt{ifft}(\widetilde{\mathcal{A}}, [], 3) = \mathcal{A},$$

where ifft denotes the Inverse Fast Fourier Transform.

Let  ${\bf A}$  be the matrix

$$\mathbf{A} = \operatorname{Diag}(\widetilde{\mathcal{A}}) = \begin{pmatrix} \widetilde{\mathcal{A}}^{(1)} & & \\ & \widetilde{\mathcal{A}}^{(2)} & & \\ & & \ddots & \\ & & & \ddots & \\ & & & & \widetilde{\mathcal{A}}^{(n_3)} \end{pmatrix},$$
(4)

and the matrices  $\widetilde{\mathcal{A}}^{(i)}$ 's are the frontal slices of the tensor  $\widetilde{\mathcal{A}}$ . The block circulant matrix bcirc( $\mathcal{A}$ ) can be block diagonalized by using the DFT matrix and this gives

$$(F_{n_3} \otimes I_{n_1}) \operatorname{bcirc}(\mathcal{A}) (F_{n_3}^* \otimes I_{n_2}) = \mathbf{A}$$
(5)

where  $F_{n_3}^*$  denotes the conjugate transpose of  $F_{n_3}$  and  $\otimes$  is the Kronecker matrix product. Next we recall the definition of the T-product.

**Definition 2.1.** The *T*-product (\*) between two tensors  $\mathcal{A} \in \mathbb{R}^{n_1 \times n_2 \times n_3}$  and  $\mathcal{B} \in \mathbb{R}^{n_2 \times m \times n_3}$  is the  $n_1 \times m \times n_3$  tensor given by:

$$\mathcal{A} \star \mathcal{B} = \text{fold}(\text{bcirc}(\mathcal{A})\text{unfold}(\mathcal{B})).$$

Notice that from the relation (4), we can show that the product  $\mathcal{C} = \mathcal{A} \star \mathcal{B}$  is equivalent to  $\mathbf{C} = \mathbf{AB}$ . So, the efficient way to compute the T-product is to use Fast Fourier Transform (FFT). The following algorithm allows us to compute in an efficient way the T-product of the tensors  $\mathcal{A}$  and  $\mathcal{B}$ .

Algorithm 1 Computing the T-product via FFT Inputs:  $\mathcal{A} \in \mathbb{R}^{n_1 \times n_2 \times n_3}$  and  $\mathcal{B} \in \mathbb{R}^{n_2 \times m \times n_3}$ 

Output:  $\mathcal{C} = \mathcal{A} \star \mathcal{B} \in \mathbb{R}^{n_1 \times m \times n_3}$ .

- 1. Compute  $\widetilde{\mathcal{A}} = \mathtt{fft}(\mathcal{A}, [], 3)$  and  $\widetilde{\mathcal{B}} = \mathtt{fft}(\mathcal{B}, [], 3)$ .
- 2. Compute each frontal slices of  $\widetilde{\mathfrak{C}}$  by

$$C^{(i)} = \begin{cases} A^{(i)}B^{(i)}, & i = 1, \dots, \lfloor \frac{n_3 + 1}{2} \rfloor \\ conj(C^{(n_3 - i + 2)}), & i = \lfloor \frac{n_3 + 1}{2} \rfloor + 1, \dots, n_3 \end{cases}$$

3. Compute  $\mathcal{C} = ifft(\widetilde{C}, [], 3)$ .

For the T-product, we have the following definitions

### **Definition 2.2.** [17]

- 1. The identity tensor  $\mathcal{J}_{n_1n_1n_3}$  is the tensor whose first frontal slice is the identity matrix  $I_{n_1n_1}$  and the other frontal slices are all zeros.
- Let z ∈ ℝ<sup>1×1×n<sub>3</sub></sup>, the tubal rank of z is the number of its non-zero Fourier coefficients. If the tubal-rank of z is equal to n<sub>3</sub>, we say that it is invertible and we denote by (z)<sup>-1</sup> the inverse of z if and only if: z ★ (z)<sup>-1</sup> = (z)<sup>-1</sup> ★ z = e. where unfold(e) = (1,0,0...,0)<sup>T</sup>.

3. An  $n_1 \times n_1 \times n_3$  tensor  $\mathcal{A}$  is invertible, if there exists a tensor  $\mathcal{B}$  of order  $n_1 \times n_1 \times n_3$ such that

$$\mathcal{A} \star \mathcal{B} = \mathcal{I}_{n_1 n_1 n_3}$$
 and  $\mathcal{B} \star \mathcal{A} = \mathcal{I}_{n_1 n_1 n_3}$ .

In that case, we set  $\mathbb{B} = \mathcal{A}^{-1}$ . It is clear that  $\mathcal{A}$  is invertible if and only if  $\operatorname{bcirc}(\mathcal{A})$  is invertible.

- 4. The transpose of A is obtained by transposing each of the frontal slices and then reversing the order of transposed frontal slices 2 through  $n_3$ .
- 5. If A, B and C are tensors of appropriate order, then

$$(\mathcal{A} \star \mathcal{B}) \star \mathcal{C} = \mathcal{A} \star (\mathcal{B} \star \mathcal{C}).$$

6. Suppose A and B are two tensors such  $A \star B$  and  $B^T \star A^T$  are defined. Then

$$(\mathcal{A} \star \mathcal{B})^T = \mathcal{B}^T \star \mathcal{A}^T.$$

**Definition 2.3.** Let A and B two tensors in  $\mathbb{R}^{n_1 \times n_2 \times n_3}$ . Then

1. The scalar inner product is defined by

$$\langle \mathcal{A}, \mathcal{B} \rangle = \sum_{i_1=1}^{n_1} \sum_{i_2=1}^{n_2} \sum_{i_3=1}^{n_3} a_{i_1 i_2 i_3} b_{i_1 i_2 i_3}.$$

2. The norm of A is defined by

$$\|\mathcal{A}\|_F = \sqrt{\langle \mathcal{A}, \mathcal{A} \rangle}.$$

# **Definition 2.4.** [17]

1. An  $n_1 \times n_1 \times n_3$  tensor Q is orthogonal if

$$Q^T \star Q = Q \star Q^T = \mathcal{I}_{n_1 n_1 n_3}.$$

2. A tensor is called f-diagonal if its frontal slices are orthogonal matrices. It is called upper triangular if all its frontal slices are upper triangular.

**Definition 2.5.** [21](Block tensor based on T-product) Suppose  $\mathcal{A} \in \mathbb{R}^{n_1 \times m_1 \times n_3}$ ,  $\mathcal{B} \in \mathbb{R}^{n_1 \times m_2 \times n_3}$ ,  $\mathcal{C} \in \mathbb{R}^{n_2 \times m_1 \times n_3}$  and  $\mathcal{D} \in \mathbb{R}^{n_2 \times m_2 \times n_3}$  are four tensors. The block tensor

$$\begin{bmatrix} \mathcal{A} & \mathcal{B} \\ \mathcal{C} & \mathcal{D} \end{bmatrix} \in \mathbb{R}^{(n_1+n_2)\times(m_1+m_2)\times n_3}$$

is defined by compositing the frontal slices of the four tensors.

**Proposition 2.1.** Let  $\mathcal{A}, \mathcal{A}_1 \in \mathbb{R}^{n \times s \times n_3}$ ,  $\mathcal{B}, \mathcal{B}_1 \in \mathbb{R}^{n \times p \times n_3}$ ,  $\mathcal{A}_2 \in \mathbb{R}^{\ell \times s \times n_3}$ ,  $\mathcal{B}_2 \in \mathbb{R}^{\ell \times p \times n_3}$ ,  $\mathcal{C} \in \mathbb{R}^{s \times n \times n_3}$ ,  $\mathcal{D} \in \mathbb{R}^{p \times n \times n_3}$  and  $\mathcal{F} \in \mathbb{R}^{n \times n \times n_3}$ . Then

1. 
$$\mathcal{F} \star [\mathcal{A} \ \mathcal{B}] = [\mathcal{F} \star \mathcal{A} \ \mathcal{F} \star \mathcal{B}] \in \mathbb{R}^{n \times (s+p) \times n_3}.$$
  
2.  $\begin{bmatrix} \mathcal{C} \\ \mathcal{D} \end{bmatrix} \star \mathcal{F} = \begin{bmatrix} \mathcal{C} \star \mathcal{F} \\ \mathcal{D} \star \mathcal{F} \end{bmatrix} \in \mathbb{R}^{(s+p) \times n \times n_3}.$   
3.  $[\mathcal{A} \ \mathcal{B}] \star \begin{bmatrix} \mathcal{C} \\ \mathcal{D} \end{bmatrix} = \mathcal{A} \star \mathcal{C} + \mathcal{B} \star \mathcal{D} \in \mathbb{R}^{n \times n \times n_3}.$   
4.  $\begin{bmatrix} \mathcal{A}_1 & \mathcal{B}_1 \\ \mathcal{A}_2 & \mathcal{B}_2 \end{bmatrix} \star \begin{bmatrix} \mathcal{C} \\ \mathcal{D} \end{bmatrix} = \begin{bmatrix} \mathcal{A}_1 \star \mathcal{C} + \mathcal{B}_1 \star \mathcal{D} \\ \mathcal{A}_2 \star \mathcal{C} + \mathcal{B}_2 \star \mathcal{D} \end{bmatrix} \in \mathbb{R}^{(\ell+n) \times n \times n_3}.$ 

Now we introduce the T-diamond tensor product.

**Definition 2.6.** Let  $\mathcal{A} = [\mathcal{A}_1, \dots, \mathcal{A}_p] \in \mathbb{R}^{n_1 \times p \times n_3}$ , where  $\mathcal{A}_i \in \mathbb{R}^{n_1 \times s \times n_3}$ ,  $i = 1, \dots, p$  and let  $\mathcal{B} = [\mathcal{B}_1, \dots, \mathcal{B}_\ell] \in \mathbb{R}^{n_1 \times \ell \times n_3}$  with  $\mathcal{B}_j \in \mathbb{R}^{n_1 \times s \times n_3}$ ,  $j = 1, \dots \ell$ . Then, the product  $\mathcal{A}^T \diamond \mathcal{B}$  is the  $p \times \ell$  matrix given by :

$$(\mathcal{A}^T \diamondsuit \mathcal{B})_{i,j} = \langle \mathcal{A}_i, \mathcal{B}_j \rangle$$
.

We consider the following Sylvester tensor equation

$$\mathcal{A} \star \mathfrak{X} + \mathfrak{X} \star \mathcal{B} = \mathfrak{C},\tag{6}$$

where  $\mathcal{A} \in \mathbb{R}^{n \times n \times n_3}$ ,  $\mathcal{B} \in \mathbb{R}^{q \times q \times n_3}$  and  $\mathfrak{X}, \mathcal{C} \in \mathbb{R}^{n \times q \times n_3}$  respectively.  $\mathcal{A}, \mathcal{B}$  and  $\mathcal{C}$  are given,  $\mathfrak{X}$  the unknown tensor to be determined.

**Theorem 2.1.** Sylvester tensor equation (6) has a unique solution if and only if  $\Gamma(\mathbf{A}) \cap \Gamma(\mathbf{B}) = \emptyset$ , where the block diagonal matrix  $\mathbf{A}, \mathbf{B}$  are defined by (4), and  $\Gamma(\mathbf{A}), \Gamma(\mathbf{B})$  denotes the set of eigenvalues of the matrix  $\mathbf{A}$  and  $\mathbf{B}$  respectively.

**Proof** It is easy to show that the product  $\mathcal{A} \star \mathcal{X} + \mathcal{X} \star \mathcal{B} = \mathcal{C}$  is equivalent to  $\mathbf{C} = \mathbf{A}\mathbf{X} + \mathbf{X}\mathbf{B}$  (Sylvester matrix equation). which has a unique solution if and only  $\Gamma(\mathbf{A}) \cap \Gamma(\mathbf{B}) = \emptyset$ .  $\Box$ 

# 3. Tensor tFOM and tGMRES algorithms

## 3.1. The tArnoldi method

Consider the following tensor linear system of equations

$$\mathcal{M}(\mathfrak{X}) = \mathfrak{C} \tag{7}$$

where  $\mathcal{M}$  an linear operator,  $\mathcal{C}$  and  $\mathfrak{X} \in \mathbb{R}^{n \times s \times p}$ . We introduce the tensor Krylov subspace  $\mathfrak{TK}_m(\mathcal{M}, \mathcal{V})$  associated to the T-product, defined for the pair  $(\mathcal{M}, \mathcal{V})$  as follows

$$\mathcal{TK}_m(\mathcal{M}, \mathcal{V}) = \mathrm{Tspan}\{\mathcal{M}, \mathcal{M}(\mathcal{V}), \dots, \mathcal{M}^{m-1}(\mathcal{V})\}$$
(8)

where  $\mathcal{M}^{i-1}(\mathcal{V}) = \mathcal{M}^{i-2}(\mathcal{M}(\mathcal{V}))$ , for i = 2, ..., m and  $\mathcal{A}^0$  is the identity tensor. In the following algorithm, we define the Tensor tArnoldi algorithm.

Algorithm 2 Tensor tArnoldi algorithm
1. Input. $\mathcal{A} \in \mathbb{R}^{n \times n \times p}$ , $\mathcal{V} \in \mathbb{R}^{n \times s \times p}$ and the positive integer $m$ .
2. Set $\beta = \ \mathcal{V}\ _F, \ \mathcal{V}_1 = \frac{\mathcal{V}}{\beta}$
3. For $j = 1,, m$
(a) $\mathcal{W} = \mathcal{M}(\mathcal{V}_j)$
(b) for $i = 1,, j$
i. $h_{i,j} = \langle \mathcal{V}_i, \mathcal{W} \rangle$
ii. $\mathcal{W} = \mathcal{W} - h_{i,j} \mathcal{V}_i$
(c) End for
(d) $h_{j+1,j} = \ \mathcal{W}\ _F$ . If $h_{j+1,j} = 0$ , stop; else
(e) $\mathcal{V}_{j+1} = \mathcal{W}/h_{j+1,j}$ .
4. End and return $\mathbb{V}_m$

It is not difficult to show that after m steps of Algorithm 2, the tensors  $\mathcal{V}_1, \ldots, \mathcal{V}_m$ , form an orthonormal basis of the tensor global Krylov subspace  $\mathcal{TK}_m(\mathcal{M}, \mathcal{V})$ . Let  $\mathbb{V}_m$  be the  $(n \times$   $(sm) \times p$ ) tensor with frontal slices  $\mathcal{V}_1, \ldots, \mathcal{V}_m$  and let  $\widetilde{H}_m$  be the  $(m+1) \times m$  upper Hesenberg matrix whose elements are the  $h_{i,j}$ 's defined by Algorithm 2. Let  $H_m$  be the matrix obtained from  $\widetilde{H}_m$  by deleting its last row;  $H_{.,j}$  will denote the *j*-th column of the matrix  $H_m$  and  $\mathcal{A} \star \mathbb{V}_m$ is the  $(n \times (sm) \times p)$  tensor with frontal slices  $\mathcal{A} \star \mathcal{V}_1, \ldots, \mathcal{A} \star \mathcal{V}_m$ :

$$\mathbb{V}_m := [\mathcal{V}_1, \dots, \mathcal{V}_m] \quad \text{and} \quad \mathcal{W}_m := [\mathcal{M}(\mathcal{V}_1), \dots, \mathcal{M}(\mathcal{V}_m)].$$
(9)

We introduce the product  $\circledast$  defined by

$$\mathbb{V}_m \circledast y = \sum_{j=1}^m y_j \mathcal{V}_j, \ y = (y_1, \dots, y_m)^T \in \mathbb{R}^m,$$

and we set

$$\mathbb{V}_m \circledast H_m = [\mathbb{V}_m \circledast H_{.,1}, \dots, \mathcal{V}_m \circledast H_{.,m}]$$

Then, it is easy to see that for all vectors u and v in  $\mathbb{R}^m$ , we have

$$\mathbb{V}_m \circledast (u+v) = \mathbb{V}_m \circledast u + \mathbb{V}_m \circledast v \quad \text{and} \quad (\mathbb{V}_m \circledast H_m) \circledast u = \mathbb{V}_m \circledast (H_m \ u).$$
(10)

With these notations, we can show the following result that will be useful later on.

**Proposition 3.1.** Let  $\mathbb{V}_m$  be the tensor defined by  $[\mathcal{V}_1, \ldots, \mathcal{V}_m]$  where  $\mathcal{V}_i \in \mathbb{R}^{n \times s \times p}$  are defined by the Tensor tArnoldi algorithm. Then, we have

$$\|\mathbb{V}_m \circledast y\|_F = \|y\|_2, \ \forall y = (y_1, \dots, y_m)^T \in \mathbb{R}^m.$$
 (11)

**Proof** From the definition of the product  $\circledast$ , we have  $\sum_{j=1}^{m} y_j \mathcal{V}_j = \mathbb{V}_m \circledast y$ . Therefore,

$$\|\mathbb{V}_m \circledast y\|_F^2 = \left\langle \sum_{j=1}^m y_j \mathcal{V}_j, \sum_{j=1}^m y_j \mathcal{V}_j \right\rangle_F.$$

But, since the tensors  $\mathcal{V}_i$ 's are orthonormal, it follows that

$$\|\mathbb{V}_m \circledast y\|_F^2 = \sum_{j=1}^m y_j^2 = \|y\|_2^2,$$

which shows the result.

With the above notations, we can easily prove the results of the following proposition.

**Proposition 3.2.** Suppose that m steps of Algorithm 2 have been run. Then, the following statements hold:

$$\mathcal{W}_m = \mathcal{V}_m \circledast H_m + h_{m+1,m} \left[ \mathcal{O}_{n \times s \times p}, \dots, \mathcal{O}_{n \times s \times p}, \mathcal{V}_{m+1} \right], \tag{12}$$

$$\mathcal{W}_m = \mathbb{V}_{m+1} \circledast \widetilde{H}_m, \tag{13}$$

$$\mathbb{V}_m^T \Diamond \mathbb{W}_m = H_m, \tag{14}$$

$$\mathbb{V}_{m+1}^T \diamondsuit \mathbb{W}_m = \widetilde{H}_m, \tag{15}$$

$$\mathbb{V}_m^T \Diamond \mathbb{V}_m = I_m, \tag{16}$$

where  $I_m$  the identity matrix and O is the tensor having all its entries equal to zero.

**Proof** From Algorithm 2, we have 
$$\mathcal{M}(\mathcal{V}_j) = \sum_{i=1}^{j+1} h_{i,j} \mathcal{V}_i$$
. Using the fact that  $\mathcal{W}_m := [\mathcal{M}(\mathcal{V}_1), \dots, \mathcal{M}(\mathcal{V}_m)],$ 

the *j*-th frontal slice of  $\mathcal{W}_m$  is given by

$$(\mathcal{W}_m)_j = \mathcal{M}(\mathcal{V}_j) = \sum_{i=1}^{j+1} h_{i,j} \mathcal{V}_i.$$

Furthermore, from the definition of the  $\circledast$  product, we have

$$(\mathbb{V}_{m+1} \circledast \widetilde{H}_m)_j = \mathbb{V}_{m+1} \circledast H_{.,j}$$
$$= \sum_{i=1}^{j+1} h_{i,j} \mathcal{V}_i,$$

which proves the first two relations. The other relations follow from the definition of T-diamond product  $\Box$ 

#### 3.2. The tFOM method

In the following, we examined the tensor full orthogonalization (tFOM) method. It could be considered as generalization of the global FOM algorithm [14]. Let  $\mathfrak{X}_0 \in \mathbb{R}^{n \times s \times p}$  be an arbitrary initial guess with the corresponding residual  $\mathcal{R}_0 = \mathfrak{C} - \mathcal{M}(\mathfrak{X}_0)$ . The aim of tensor T-global GMRES method is to find and approximate solution  $\mathfrak{X}_m$  approximating the exact solution  $\mathfrak{X}^*$  of (7) such that

$$\mathfrak{X}_m - \mathfrak{X}_0 = \mathfrak{P}_m \in \mathfrak{TK}_m(\mathcal{M}, \mathfrak{R}_0), \tag{17}$$

and

$$\mathfrak{R}_m \perp \mathfrak{T}\mathfrak{K}_m(\mathcal{M}, \mathfrak{R}_0) \tag{18}$$

 $\mathcal{P}_m \in \mathfrak{TK}_m(\mathcal{M}, \mathcal{R}_0)$  can be expressed as  $\mathcal{P}_m = \mathcal{V}_m \circledast y$  with  $y = (y_1, \ldots, y_m)^T \in \mathbb{R}^m$ . From where the residual  $\mathcal{R}_m$  is given by

$$\mathfrak{R}_m = \mathfrak{R}_0 - \mathfrak{W}_m \circledast y \tag{19}$$

Using (19), the relation (18) can be expressed as

$$\langle \mathcal{V}_i, \mathcal{R}_m \rangle = \langle \mathcal{V}_i, \mathcal{W}_m \circledast y \rangle, \quad i = 1, \dots, m$$
 (20)

where the tensors  $\mathcal{V}_1, \ldots, \mathcal{V}_m$ , form an orthonormal basis of the tensor global Krylov subspace  $\mathcal{TK}_m(\mathcal{M}, \mathcal{R}_0)$ . Using (10) and (12), equation (20) can be expressed as:

$$H_m \ y = \|\mathcal{R}_0\|_F \ e_1^{(m)},\tag{21}$$

where  $e_1^{(m)}$  is the first canonical basis vector in  $\mathbb{R}^m$  and  $H_m$  the Hessenberg matrix of size  $(m \times m)$  obtained from Algorithm 2.

**Proposition 3.3.** At step m, the norm of the residual  $\Re_m = \mathcal{C} - \mathcal{M}(\mathfrak{X}_m) = \Re_0 - \mathfrak{W}_m \circledast y_m$ produced by the tFOM method for tensor equation (1) has the following expression

$$\|\mathcal{R}_m\|_F = h_{m+1,m} \left| y_m^{(m)} \right|.$$
 (22)

where  $y_m^{(m)}$  is the last component of the vector  $y_m$ 

**Proof** At step m, using the relations (19) and (18) the norm of the residual  $\mathcal{R}_m$  can be expressed as

 $\|\mathcal{R}_m\|_F = \|\mathcal{R}_0 - \mathbb{V}_m \circledast (H_m y_m) + h_{m+1,m} [\mathcal{O}_{n \times s \times p}, \dots, \mathcal{O}_{n \times s \times p}, \mathcal{V}_{m+1}] \circledast y_m\|_F.$ since  $H_m y_m = \|\mathcal{R}_0\|_F e_1^{(m)}$  and  $\mathcal{R}_0\|_F (\mathbb{V}_m \circledast e_1^{(m)})$ , we get

$$\|\mathcal{R}_{m}\|_{F} = h_{m+1,m} \|[\mathcal{O}_{n \times s \times p}, \dots, \mathcal{O}_{n \times s \times p}, \mathcal{V}_{m+1}] \circledast y_{m}\|_{F} = h_{m+1,m} |y_{m}^{(m)}|$$

which shows the results.  $\Box$ 

### 3.3. The tGMRES method

In the sequel, we develop the tensor tGMRES algorithm for solving the problem (7). It could be considered as generalization of the global GMERS algorithm [15]. Let  $\mathcal{X}_0 \in \mathbb{R}^{n \times s \times p}$  be an arbitrary initial guess with the corresponding residual  $\mathcal{R}_0 = \mathcal{C} - \mathcal{M}(\mathcal{X}_0)$ . The purpose of tensor tGMRES method is to find and approximate solution  $\mathcal{X}_m$  approximating the exact solution  $\mathcal{X}^*$ of (7) such that

$$\mathfrak{X}_m - \mathfrak{X}_0 \in \mathfrak{TK}_m(\mathcal{M}, \mathfrak{R}_0), \tag{23}$$

with the classical minimization property

$$\|\mathcal{R}_m\|_F = \min_{\mathfrak{X} \in \mathfrak{X}_0 + \mathfrak{TK}_m(\mathcal{M}, \mathcal{R}_0)} \|\mathcal{C} - \mathcal{M}(\mathfrak{X})\|_F.$$
(24)

Let  $\mathfrak{X}_m = \mathfrak{X}_0 + \mathbb{V}_m \circledast y$  with  $y \in \mathbb{R}^m$ , be the approximate solution satisfying (23). Then,

$$\begin{aligned} \mathcal{R}_m = & \mathcal{C} - \mathcal{M}(\mathcal{X}_m) \\ = & \mathcal{C} - \mathcal{M} \left( \mathcal{X}_0 + \mathbb{V}_m \circledast y \right) \\ = & \mathcal{R}_0 - \mathbb{W}_m \circledast y. \end{aligned}$$

It follows then that

$$\|\mathfrak{R}_m\|_F = \min_{y \in \mathbb{R}^m} \|\mathfrak{R}_0 - \mathbb{W}_m \circledast y\|_F,$$

where  $\mathcal{W}_m := [\mathcal{M}(\mathcal{V}_1), \dots, \mathcal{M}(\mathcal{V}_m)]$  is the  $(n \times sm \times p)$  tensor defined earlier. Using Proposition 3.1 and the fact that  $\mathcal{R}_0 = \|\mathcal{R}_0\|_F \mathcal{V}_1$  with  $\mathcal{V}_1 = \mathcal{V}_{m+1} \circledast e_1$ , where  $e_1$  the first canonical basis vector in  $\mathbb{R}^{m+1}$ , we get

$$\begin{split} \|\mathcal{R}_{0} - (\mathcal{A} \star \mathbb{V}_{m}) \circledast y\|_{F} &= \|\mathcal{R}_{0} - (\mathbb{V}_{m+1} \circledast H_{m}) \circledast y\|_{F} \\ &= \|\|\mathcal{R}_{0}\|_{F} (\mathbb{V}_{m+1} \circledast e_{1}) - (\mathbb{V}_{m+1} \circledast \widetilde{H}_{m}) \circledast y\|_{F} \\ &= \|\mathbb{V}_{m+1} \circledast (\|\mathcal{R}_{0}\|_{F} e_{1} - \widetilde{H}_{m} y)\|_{F} \\ &= \|\|\mathcal{R}_{0}\|_{F} e_{1} - \widetilde{H}_{m} y\|_{2}. \end{split}$$

Finally, we obtain

$$\mathfrak{X}_m = \mathfrak{X}_0 + \mathbb{V}_m \circledast y, \tag{25}$$

where,

$$y = \arg \min_{y \in \mathbb{R}^m} || \ ||\mathcal{R}_0||_F \ e_1 - \hat{H}_m y)||_2.$$
(26)

**Proposition 3.4.** At step m, the residual  $\Re_m = \mathcal{C} - \mathcal{M}(\mathfrak{X}_m)$  produced by the tensor Global GMRES method for tensor equation (1) has the following expression

$$\mathfrak{R}_m = \mathbb{V}_{m+1} \circledast \left(\gamma_{m+1} Q_m e_{m+1}\right),\tag{27}$$

where  $Q_m$  is the unitary matrix obtained from the QR decomposition of the upper Hessenberg matrix  $\widetilde{H}_m$  and  $\gamma_{m+1}$  is the last component of the vector  $\|\mathcal{R}_0\|_F Q_m^{\mathrm{T}} e_1$  and  $e_{m+1} = (0, 0, \dots, 1)^{\mathrm{T}} \in \mathbb{R}^{m+1}$ .

Furthermore,

$$\left\|\mathcal{R}_{m}\right\|_{F} = \left|\gamma_{m+1}\right|. \tag{28}$$

**Proof** At step m, the residual  $\mathcal{R}_m$  can be expressed as

$$\mathcal{R}_m = \mathbb{V}_{m+1} \circledast \left(\beta e_1 - \widetilde{H}_m y_m\right)$$

by considering the QR decomposition  $\widetilde{H}_m = Q_m \widetilde{U}_m$  of the  $(m+1) \times m$  matrix  $\widetilde{H}_m$ , we get

$$\mathfrak{R}_m = (\mathbb{V}_{m+1} \circledast Q_m) \circledast \left(\beta Q_m^T e_1 - \widetilde{U}_m y_m\right).$$

Since y solves problem (26), it follows that

$$\mathfrak{R}_m = \mathbb{V}_{m+1} \circledast (\gamma_{m+1} Q_m e_{m+1})$$

where  $\gamma_{m+1}$  is the last component of the vector  $\beta Q_m^T e_1$ . Therefore,

$$\begin{aligned} \|\mathcal{R}_{m}\|_{F} &= \|\mathbb{V}_{m+1} \circledast (\gamma_{m+1}Q_{m}e_{m+1})\|_{F} \\ &= \|\gamma_{m+1}Q_{m}e_{m+1}\|_{2} \\ &= |\gamma_{m+1}|, \end{aligned}$$

which shows the results.  $\Box$ 

### 4. t-Bartels-Stewart method for Sylvester tensor equations of small size

In this section, we introduce the *t-Bartels-Stewart* method for Sylvester tensor equations of small size based on T-product formalism, as a generalisation of the well known Bartels and Stewart algorithm proposed in [1]. Motivated by the matrix case, the *t-Bartels-Stewart* method

- Algorithm 3 tFOM and tGMRES Algorithms 1. Input.  $\mathcal{A} \in \mathbb{R}^{n \times n \times n_3}$ ,  $\mathcal{B} \in \mathbb{R}^{s \times s \times n_3}$ ,  $\mathfrak{X}_0, \mathfrak{C} \in \mathbb{R}^{n \times s \times n_3}$ , the maximum number of iteration Iter<sub>max</sub> an integer m and a tolerance tol.
  - 2. **Output.**  $\mathfrak{X}_m \in \mathbb{R}^{n \times s \times n_3}$  the approximate solution of (7).
  - 3. Compute  $\mathcal{R}_0 = \mathcal{C} \mathcal{M}(\mathfrak{X}_0)$ .
  - 4. For  $k = 1, \ldots$ , Iter<sub>max</sub>

    - (b) Solve the problem :  $\begin{cases} H_m \ y_m = \|\mathcal{R}_0\|_F \ e_1^{(m)} \ (\mathbf{tFOM method}) \\ y_m = \arg \ \min_{y \in \mathbb{R}^m} || \ ||\mathcal{R}_0||_F \ e_1^{(m+1)} \widetilde{H}_m y_m ||_2. \ (\mathbf{tGMRES method} \ ) \end{cases}$ (c) Compute  $\mathfrak{X}_m = \mathfrak{X}_0 + \mathbb{V}_m \circledast y_m$
  - 5. If  $||\mathcal{R}_m||_F < tol$  then

return  $\mathfrak{X}_m$ ;

- 6. else  $\mathfrak{X}_0 = \mathfrak{X}_m$  and go to Step 2.
- 7. **Output.**  $\mathfrak{X}_m \in \mathbb{R}^{n \times s \times n_3}$  the approximate solution of (7).

is based on transforming the tensors  $\mathcal{A}$  and  $\mathcal{B}$  into *t*-real Schur form that will be defined later. This gives an new triangular tensor equation that will be solved by *t*-back-substitution method. First, we introduce the *t*-real schur decomposition.

**Theorem 4.1.** Let  $\mathcal{A} \in \mathbb{R}^{n \times n \times n_3}$ . Then  $\mathcal{A}$  can be factored as

$$\mathcal{A} = \mathcal{U} \star \mathcal{R} \star \mathcal{U}^T, \tag{29}$$

where  $\mathfrak{R} \in \mathbb{R}^{n \times n \times n_3}$  quasi upper triangular tensor (each frontal slice of  $\mathfrak{R}$  is quasi upper triangular) and  $\mathcal{U} \in \mathbb{R}^{n \times n \times n_3}$  orthogonal tensor.

**Proof** We have

$$(F_{n_3} \otimes I_{n_1}) \operatorname{bcirc}(\mathcal{A}) (F_{n_3}^* \otimes I_{n_1}) = \mathbf{A} = \begin{pmatrix} \widetilde{\mathcal{A}}^{(1)} & & \\ & \widetilde{\mathcal{A}}^{(2)} & & \\ & & \ddots & \\ & & & \widetilde{\mathcal{A}}^{(n_3)} \end{pmatrix}.$$

Next, we compute the schur matrix decomposition of each frontal slice  $\widetilde{\mathcal{A}}^{(i)}$ ,  $i = 1, \ldots, n_3$ . Then

$$\begin{pmatrix} \widetilde{\mathcal{A}}^{(1)} & & \\ & \ddots & \\ & \widetilde{\mathcal{A}}^{(n_3)} \end{pmatrix} = \begin{pmatrix} \widetilde{\mathcal{U}}^{(1)} & & \\ & \ddots & \\ & & \widetilde{\mathcal{U}}^{(n_3)} \end{pmatrix} \begin{pmatrix} \widetilde{\mathcal{R}}^{(1)} & & \\ & \ddots & \\ & & \widetilde{\mathcal{R}}^{(n_3)} \end{pmatrix} \begin{pmatrix} \widetilde{\mathcal{U}}^{(1)T} & & \\ & & \ddots & \\ & & & \widetilde{\mathcal{U}}^{(n_3)T} \end{pmatrix}$$
(30)  
Since  $(F_{n_3}^* \otimes I_{n_1}) \begin{pmatrix} \widetilde{\mathcal{U}}^{(1)} & & \\ & \ddots & \\ & & & \widetilde{\mathcal{U}}^{(n_3)} \end{pmatrix} (F_{n_3} \otimes I_{n_1}), (F_{n_3}^* \otimes I_{n_1}) \begin{pmatrix} \widetilde{\mathcal{R}}^{(1)} & & \\ & \ddots & \\ & & & \widetilde{\mathcal{R}}^{(n_3)} \end{pmatrix} (F_{n_3} \otimes I_{n_3}) \end{pmatrix} (F_{n_3} \otimes I_{n_1})$ are block circulant matrices, by apply-  
 $\widetilde{\mathcal{U}}^{(n_3)T}$ 

ing the appropriate matrices  $(F_{n_3}^* \otimes I_{n_1}), (F_{n_3} \otimes I_{n_1})$  to the left and right of each matrix in (30) respectively, and folding up the result. This give the result.  $\Box$ 

The *t*-real schur decomposition can be computed by the Algorithm 4.

Next, we transform the tensors  $\mathcal{A}$  and  $\mathcal{B}$  into *t-real schur* forms  $\mathcal{A} = \mathcal{U}_{\mathcal{A}} \star \mathcal{R}_{\mathcal{A}} \star \mathcal{U}_{\mathcal{A}}^{T}$  and

# Algorithm 4 t-real schur decomposition

1. Input:  $\mathcal{A} \in \mathbb{R}^{n \times n \times n_3}$ .

- 2. **Output:**  $\mathcal{R} \in \mathbb{R}^{n \times n \times n_3}$  upper triangular tensor.  $\mathcal{U} \in \mathbb{R}^{n \times n \times n_3}$  orthogonal tensor.
- 3. Set \$\tilde{\mathcal{A}}\$ = fft(\$\mathcal{A}\$, [], 3)
  (a) for \$i = 1, ..., n\_3\$
  i. [\$\tilde{\mathcal{Q}}^{(i)}\$\$\tilde{\mathcal{R}}^{(i)}\$] = schur(\$\tilde{\mathcal{A}}^{(i)}\$) (real schur matrix decomposition)
  (b) End
  4. \$\mathcal{Q}\$ = ifft(\$\tilde{\mathcal{Q}}\$, [], 3), \$\mathcal{R}\$ = ifft(\$\tilde{\mathcal{R}}\$, [], 3)
  5. End

 $\mathcal{B} = \mathcal{U}_{\mathcal{B}} \star \mathcal{R}_{\mathcal{B}} \star \mathcal{U}_{\mathcal{B}}^{T}$ . Then the Sylvester tensor equation (6) become

$$(\mathcal{U}_{\mathcal{A}} \star \mathcal{R}_{\mathcal{A}} \star \mathcal{U}_{\mathcal{A}}^{T}) \star \mathcal{X} + \mathcal{X} \star (\mathcal{U}_{\mathcal{B}} \star \mathcal{R}_{\mathcal{B}} \star \mathcal{U}_{\mathcal{B}}^{T}) = \mathcal{C}$$

which equivalent to

$$\mathcal{U}_{\mathcal{A}}^{T} \star \left[ (\mathcal{U}_{\mathcal{A}} \star \mathcal{R}_{\mathcal{A}} \star \mathcal{U}_{\mathcal{A}}^{T}) \star \mathcal{X} + \mathcal{X} \star (\mathcal{U}_{\mathcal{B}} \star \mathcal{R}_{\mathcal{B}} \star \mathcal{U}_{\mathcal{B}}^{T}) \right] \star \mathcal{U}_{\mathcal{B}} = \mathcal{U}_{\mathcal{A}}^{T} \star \mathcal{C} \star \mathcal{U}_{\mathcal{B}}$$

Since  $\mathcal{U}_{\mathcal{A}}$  and  $\mathcal{U}_{\mathcal{B}}$  are orthogonal tensors, we get

$$\mathcal{R}_{\mathcal{A}} \star \mathcal{Y} + \mathcal{Y} \star \mathcal{R}_{\mathcal{B}} = \mathcal{C}_1 \tag{31}$$

where  $\mathcal{Y} = \mathcal{U}_{\mathcal{A}}^T \star \mathcal{X} \star \mathcal{U}_{\mathcal{B}}$  and  $\mathcal{C}_1 = \mathcal{U}_{\mathcal{A}}^T \star \mathcal{C} \star \mathcal{U}_{\mathcal{B}}$ .

**Remark 4.1.** The equation (31) can be solved by the t-back-substitution method. This method follow the same steps of matrix back substitution, where the tube fibers (3-mode fibers), lateral slices and T-product play the role of scalars, vectors and matrix product respectively.

Finally, the *t-Bartels-Stewart* method is implemented by Algorithm 5.

Algorithm 5 t-Bartels-Stewart method	
1. Input: $\mathcal{A} \in \mathbb{R}^{n \times n \times n_3}, \mathcal{B} \in \mathbb{R}^{q \times q \times n_3}$ and $\mathcal{C} \in \mathbb{R}^{n \times q \times n_3}$	

- 2. **Output:**  $\mathcal{X}_S \in \mathbb{R}^{n \times q \times n_3}$  solution of Sylvester tensor equation (6) of small size.
- 3. Compute *t*-real schur forms

$$\mathcal{A} = \mathcal{U}_{\mathcal{A}} \star \mathcal{R}_{\mathcal{A}} \star \mathcal{U}_{\mathcal{A}}^{T}, \quad \mathcal{B} = \mathcal{U}_{\mathcal{B}} \star \mathcal{R}_{\mathcal{B}} \star \mathcal{U}_{\mathcal{B}}^{T}$$

using Algorithm 4.

- 4. Compute  $\mathcal{Y}_S$  solution of tensor equation (31) using *t*-back-substitution method.
- 5. Compute  $\mathfrak{X}_S = \mathfrak{U}_A \star \mathfrak{Y}_S \star \mathfrak{U}_B^T$ .

# 5. Tensor Krylov methods via T-product for solving large Sylvester tensor equations

In this section, we consider the case when the Sylvester tensor equation (1) and (2) is of large size. As in matrix case, iterative projection methods have been developed; see [5, 13, 22]. These methods use Galerkin projection methods, such the classical and the block Arnoldi techniques,

to produce low-dimensional Sylvester matrix equations that are solved by using direct methods. In tensor case, the main idea is to transform the large Sylvester tensor equations using the wellknow Tubal-Block-Arnoldi that will be introduced later into low dimensional equations.

#### 5.1. Tubal Block Arnoldi method

In this subsection, we introduce the Tubal-Block-Arnoldi method via T-product as a generalization of the well-know block Arnoldi method, see [5] for more details.

Let  $\mathcal{A} \in \mathbb{R}^{n_1 \times n_1 \times n_3}$  be a square tensor and  $\mathcal{V} \in \mathbb{R}^{n_1 \times s \times I_3}$ ,  $s \ll n_1$ . The tensor block Krylov subspace is defined by

$$\mathcal{K}_{m}^{Block}(\mathcal{A},\mathcal{V}) = Range\left(\left[\mathcal{V},\mathcal{A}\star\mathcal{V},\ldots,\mathcal{A}^{m-1}\star\mathcal{V}\right]\right) \subset \mathbb{R}^{n_{1}\times s\times n_{3}}$$
(32)

where  $\mathcal{A}^{i} = \mathcal{A} \star \mathcal{A}^{i-1}, i = 1, \dots, m-1, \mathcal{A}^{0} = \mathcal{I}_{n_{1}n_{1}n_{3}}$  and

$$Range(\mathcal{Z}) = \left\{ \overrightarrow{\mathcal{Y}} \in \mathbb{R}^{n_1 \times 1 \times n_3} \mid \overrightarrow{\mathcal{Y}} = \mathcal{Z} \star \overrightarrow{\mathcal{X}}, \ \overrightarrow{\mathcal{X}} \in \mathbb{R}^{s \times 1 \times n_3} \right\}$$

for  $\mathcal{I} \in \mathbb{R}^{n_1 \times s \times \times n_3}$ .

Before describing the Tubal-Block-Arnoldi process, let us first introduce the Tubal-QR Factorization.

**Theorem 5.1.** (Tubal-QR Factorization) Let  $\mathcal{A} = \begin{bmatrix} \overrightarrow{\mathcal{A}}_1, \dots, \overrightarrow{\mathcal{A}}_m \end{bmatrix} \in \mathbb{R}^{n_1 \times m \times n_3}, \overrightarrow{\mathcal{A}}_i \in \mathbb{R}^{n_1 \times 1 \times n_3}$ for  $i = 1, \dots, m$ , there exist orthogonal tensors  $\mathcal{U} \in \mathbb{R}^{n_1 \times n_2 \times n_3}$  and  $\mathcal{R} \in \mathbb{R}^{n_2 \times n_2 \times n_3}$  triangular tensor such that

$$\mathcal{A} = \mathcal{Q} \star \mathcal{R} \tag{33}$$

We call (33) the tubal QR factorization of the tensor  $\mathcal{A}$ .

The Tubal QR factorization is summarized in Algorithm 6.

The function Normalization1( $\vec{\mathcal{A}}$ ) described in [18] allow us to write a given tensor  $\vec{\mathcal{A}} \in \mathbb{R}^{n \times 1 \times n_3}$  as follow:  $\vec{\mathcal{A}} = \vec{\mathcal{U}} \star \mathbf{a}$  where  $\mathbf{a} \in \mathbb{R}^{1 \times 1 \times n_3}$  is invertible and  $\vec{\mathcal{U}}^T \star \vec{\mathcal{U}} = \mathbf{e}$ .

Tubal-Block-Arnoldi method have to build an orthonormal basis of  $\mathbb{V}_m^b$  of the Krylov subspace  $\mathcal{K}_m^{Block}(\mathcal{A}, \mathcal{V}).$ 

Algorithm 6 Tubal-QR Factorization (Tubal-QR)

- 1. Input.  $\mathcal{A} \in \mathbb{R}^{n \times m \times n_3}$ 2. Set  $[\mathcal{V}_1, R_{1,1,:}] = Normalization1(\overrightarrow{\mathcal{A}}_1)$ 3. For  $j = 1, \dots, m$ (a)  $\mathcal{W} = \overrightarrow{\mathcal{A}}_j$ , (b) for  $i = 2, \dots, j - 1$ i.  $\mathcal{R}_{i,j,:} = \mathcal{V}_i^T \star \mathcal{W}$ ii.  $\mathcal{W} = \mathcal{W} - \mathcal{V}_i \star \mathcal{R}_{i,j,:}$ (c) End for (d)  $[\mathcal{Q}_j, \mathcal{R}_{j,j,:}] = Normalization1(\overrightarrow{\mathcal{W}})$ 4. End
- 5. **Output.**  $Q \in \mathbb{R}^{n \times m \times n_3}$  orthogonal and  $\mathcal{R} \in \mathbb{R}^{m \times m \times n_3}$  such that :  $\mathcal{A} = Q \star \mathcal{R}$ .

# Algorithm 7 The Tubal Block Arnoldi Algorithm (TBA)

1. Input.  $\mathcal{A} \in \mathbb{R}^{n_1 \times n_1 \times n_3}$  be a square tensor,  $\mathcal{V} \in \mathbb{R}^{n_1 \times s \times I_3}$ ,  $s \ll n_1$  and an integer m.

- 2. Set  $[\mathcal{V}_1^b, \mathcal{H}_0] = \text{Tubal-QR}(\mathcal{V}).$
- 3. For j = 1,...,m
  - (a)  $\mathcal{W} = \mathcal{A} \star \mathcal{V}_j$
  - (b) For i = 1,...,j
    - i.  $\mathcal{H}_{i,j} = V_i^{bT} \star \mathcal{W}$
    - ii.  $\mathcal{W} = \mathcal{W} V_i^b \star \mathcal{H}_{i,j},$
  - (c) End.
  - (d)  $[V_{j+1}, \mathcal{H}_{j+1,j}] = \text{Tubal-QR}(\mathcal{W}).$
- 4. End return  $\mathbb{V}_m^b$

Using Definition 2.5, we can put away the tensors  $\mathcal{H}_{i,j} \in \mathbb{R}^{s \times s \times n_3}$  into a block tensors  $\mathbb{H}_m$  and

 $\mathbb{H}_{m+1}$  defined as follow

$$\mathbb{H}_{m+1} = \begin{bmatrix} \mathcal{H}_{1,1} & \mathcal{H}_{1,2} & \cdots & \mathcal{H}_{1,m} \\ \mathcal{H}_{2,1} & \mathcal{H}_{2,2} & \cdots & \mathcal{H}_{2,m} \\ & \ddots & \ddots & \vdots \\ & & \mathcal{H}_{m,m-1} & \mathcal{H}_{m,m} \\ & & & \mathcal{H}_{m+1,m} \end{bmatrix} \in \mathbb{R}^{(m+1)s \times ms \times n_3}$$
$$\mathbb{H}_m = \begin{bmatrix} \mathcal{H}_{1,1} & \mathcal{H}_{1,2} & \cdots & \mathcal{H}_{1,m} \\ \mathcal{H}_{2,1} & \mathcal{H}_{2,2} & \cdots & \mathcal{H}_{2,m} \\ & \ddots & \ddots & \vdots \\ & & \mathcal{H}_{m,m-1} & \mathcal{H}_{m,m} \end{bmatrix} \in \mathbb{R}^{ms \times ms \times n_3}.$$

,

It is not difficult to show that after m steps of Algorithm 7, the tensor  $\mathbb{V}_m^b := [\mathcal{V}_1^b, \dots, \mathcal{V}_m^b] \in \mathbb{R}^{n_1 \times ms \times n_3}$ , where  $\mathcal{V}_i^b \in \mathbb{R}^{n_1 \times s \times n_3}$  form an orthonormal basis of the tensor Block Krylov subspace  $\mathcal{K}_m^{Block}(\mathcal{A}, \mathcal{V})$ .

It is easy to see that  $\mathbb{H}_m$  can be obtained from  $\mathbb{H}_{m+1}$  by deleting the last block row

$$[\mathcal{O}_{ssn_3},\ldots,\mathcal{O}_{ssn_3},\mathcal{H}_{m+1,m}] = \mathcal{H}_{m+1,m} \star \mathbb{E}_m \in \mathbb{R}^{s \times ms \times n_3}$$

where  $\mathcal{O}_{ssn_3}$  denote the zeros tensors of size  $(s \times s \times n_3)$  which all its entries are equal to zeros, and  $\mathbb{E}_m = [\mathcal{O}_{ssn_3}, \dots, \mathcal{O}_{ssn_3}, \mathcal{I}_{ssn_3}] \in \mathbb{R}^{s \times ms \times n_3}$ Let

$$\mathcal{A} \star \mathbb{V}_{m}^{b} := [\mathcal{A} \star \mathcal{V}_{1}, \dots, \mathcal{A} \star \mathcal{V}_{m}] \in \mathbb{R}^{n_{1} \times ms \times n_{3}},$$
$$\mathbb{V}_{m+1}^{b} := [\mathbb{V}_{m}^{b}, \mathcal{V}_{m+1}^{b}] \in \mathbb{R}^{n_{1} \times (m+1)s \times n_{3}},$$
$$\mathbb{H}_{m} = (\mathcal{H}_{i,j})_{1 \leq i,j \leq m} \in \mathbb{R}^{ms \times \times ms \times n_{3}},$$
$$\mathbb{H}_{m+1} = \begin{bmatrix} \mathbb{H}_{m} \\ \mathcal{H}_{m+1,m} \star \mathbb{E}_{m} \end{bmatrix} \in \mathbb{R}^{(m+1)s \times ms \times n_{3}}.$$

With the above notations, we can easily prove the results of the following proposition.

Proposition 5.1. From Algorithm 7, we have

$$\mathcal{A} \star \mathbb{V}_{m}^{b} = \mathbb{V}_{m}^{b} \star \mathbb{H}_{m} + \mathcal{V}_{m+1}^{b} \star (\mathcal{H}_{m+1,m} \star \mathbb{E}_{m}), \tag{34}$$

$$\mathbb{V}_m^{bT} \star \mathcal{A} \star \mathbb{V}_m^b = \mathbb{H}_m,\tag{35}$$

$$\mathbb{V}_{m+1}^{bT} \star \mathcal{A} \star \mathbb{V}_{m}^{b} = \mathbb{H}_{m+1}, \tag{36}$$

$$\mathbb{V}_m^{bT} \star \mathbb{V}_m^b = \mathfrak{I}_{ms} \tag{37}$$

where  $\mathbb{J}_{ms} \in \mathbb{R}^{ms \times ms \times n_3}$ , denote the identity tensor.

Notice that in the case  $n_3 = 1$ , Algorithm 7 reduces to the well known block Arnoldi process. **Proof** The proof come directly from steps of Algorithm 7 and Proposition 2.1.  $\Box$ 

### 5.2. Tubal-Block-Arnoldi method for solving large Sylvester tensor equations (TBAS)

This subsection discusses the computation of an approximate solution of the tensor equations :

$$\mathcal{M}(\mathfrak{X}) = \mathfrak{C},\tag{38}$$

where  $\mathcal{M}$  is a linear operator that could be described as

$$\mathcal{M}(\mathcal{X}) = \mathcal{A} \star \mathcal{X} - \mathcal{X} \star \mathcal{B} \tag{39}$$

where  $\mathcal{A} \in \mathbb{R}^{n \times n \times n_3}$ ,  $\mathcal{B} \in \mathbb{R}^{q \times q \times n_3}$  and  $\mathcal{X}, \mathcal{C} \in \mathbb{R}^{n \times q \times n_3}$  respectively.  $\mathcal{A}, \mathcal{B}$  and  $\mathcal{C}$  are given,  $\mathcal{X}$  the unknown tensor to be determined.

Let  $\mathfrak{X}_0 \in \mathbb{R}^{n \times s \times p}$  be an arbitrary initial guess with the corresponding residual  $\mathfrak{R}_0 = \mathfrak{C} - \mathcal{M}(\mathfrak{X}_0)$ . The purpose of tensor Tubal Block Arnoldi method for solving large Sylvester tensor equations (TBAS) method is to find and approximate solution  $\mathfrak{X}_m$  of the exact solution  $\mathfrak{X}^*$  of (38) such that

$$\mathfrak{X}_m - \mathfrak{X}_0 \in \mathcal{K}_m^{Block}(\mathcal{M}, \mathcal{R}_0), \tag{40}$$

with the classical orthogonality property

$$\mathfrak{R}_m = \mathfrak{C} - \mathcal{M}(\mathfrak{X}_m) \perp \mathcal{K}_m^{Block}(\mathcal{M}, \mathfrak{R}_0).$$
(41)

Using the fact that  $\mathcal{K}_m^{Block}(\mathcal{M}, \mathcal{R}_0) = \mathcal{K}_m^{Block}(\mathcal{A}, \mathcal{R}_0)$  (which not difficult to prove), the relations

(40) and (41) can be expressed as

$$\mathfrak{X}_m - \mathfrak{X}_0 \in \mathcal{K}_m^{Block}(\mathcal{A}, \mathfrak{R}_0), \tag{42}$$

with the classical orthogonality property

$$\mathfrak{R}_m = \mathfrak{C} - \mathcal{M}(\mathfrak{X}_m) \perp \mathcal{K}_m^{Block}(\mathcal{A}, \mathfrak{R}_0).$$
(43)

Since  $\mathbb{V}_m^b$  (defined earlier) form an orthonormal basis of the Krylov subspace  $\mathcal{K}_m^{Block}(\mathcal{A}, \mathcal{R}_0)$ . The relations (42) and (43) can be written as

$$\mathfrak{X}_m = \mathfrak{X}_0 + \mathbb{V}_m^b \star \mathfrak{Y}_m, \quad \text{with } \mathfrak{Y}_m \in \mathbb{R}^{ms \times s \times n_3},$$
(44)

and

$$\mathbb{V}_m^{bT} \star \mathfrak{R}_m = 0. \tag{45}$$

Using the fact that  $\mathcal{R}_m = \mathcal{C} - \mathcal{M}(\mathcal{X}_m) = \mathcal{C} - \mathcal{A} \star \mathcal{X}_m + \mathcal{X}_m \star \mathcal{B}$ , the relations (44) and (45), we get the low dimensional equation

$$\mathbb{H}_m \star \mathcal{Y}_m - \mathcal{Y}_m \star \mathcal{B} = \mathcal{C}_1 \tag{46}$$

where  $\mathcal{C}_1 = \mathbb{V}_m^{bT} \star \mathcal{R}_0$ .

The tensor equation (46) will be solved by using the *t-Bartels-Stewart* method.

**Proposition 5.2.** At step m, the residual norm of  $\Re_m = \mathbb{C} - \mathcal{M}(\mathfrak{X}_m) = \mathbb{C} - \mathcal{A} \star \mathfrak{X}_m + \mathfrak{X}_m \star \mathfrak{B}$ with  $\mathfrak{X}_m = \mathfrak{X}_0 + \mathbb{V}_m^b \star \mathfrak{Y}_m$  can be expressed as follows :

$$||\mathcal{R}_m||_F = ||\mathcal{H}_{m+1,m} \star \mathbb{E}_m \star \mathcal{Y}_m||_F \tag{47}$$

where  $\mathbb{E}_m = [\mathcal{O}_{ssn_3}, \dots, \mathcal{O}_{ssn_3}, \mathcal{I}_{ssn_3}] \in \mathbb{R}^{s \times ms \times n_3}$  and  $\mathcal{Y}_m$  solution of equation (46).

**Proof** At step m, the residual  $\mathcal{R}_m = \mathcal{C} - \mathcal{M}(\mathcal{X}_m) = \mathcal{C} - \mathcal{A} \star \mathcal{X}_m + \mathcal{X}_m \star \mathcal{B}$  with  $\mathcal{X}_m = \mathcal{X}_0 + \mathbb{V}_m^b \star \mathcal{Y}_m$  can be written as :

$$\mathfrak{R}_m = \mathfrak{R}_0 - \mathcal{A} \star \mathfrak{V}_m \star \mathfrak{Y}_m + \mathfrak{V}_m \star \mathfrak{Y}_m \star \mathfrak{B}.$$

Then, using the fact that  $\mathcal{A} \star \mathbb{V}_m^b = \mathbb{V}_m^b \star \mathbb{H}_m + \mathcal{V}_{m+1}^b \star (\mathcal{H}_{m+1,m} \star \mathbb{E}_m)$ , the relation (46) and the fact that the tensor  $\mathcal{V}_{m+1}^b$  is orthogonal, we get

$$||\mathcal{R}_m||_F = ||\mathcal{H}_{m+1,m} \star \mathbb{E}_m \star \mathcal{Y}_m||_F$$

To save memory and CPU-time requirements, the Tubal Block Arnoldi method for solving large Sylvester tensor equation (TBAS) will be used in a restarted mode. This means that we have to restart the algorithm every m inner iterations, where m is a fixed integer. The restarted Tubal Block Arnoldi algorithm for solving (38), denoted by TBAS(m), is summarized as follows: During the computation of the approximate solution  $\mathcal{X}_m$  of (38), we assume that  $\Gamma(\mathbf{H})_m \cap$ 

Algorithm 8 The Tubal-Block-Arnoldi for solving large Sylvester tensor equation TBAS(m)

1. Input.  $\mathcal{A} \in \mathbb{R}^{n \times n \times n_3}$ ,  $\mathcal{V}, \mathcal{B} \in \mathbb{R}^{q \times q \times n_3}$ ,  $\mathcal{X}_0, \mathcal{C} \in \mathbb{R}^{n \times q \times n_3}$ , the maximum number of iteration Iter<sub>max</sub> an integer *m* and a tolerance *tol*.

- 2. Compute  $\mathcal{R}_0 = \mathcal{C} \mathcal{M}(\mathfrak{X}_0)$ .
- 3. For  $k = 1, \ldots, \text{Iter}_{\max}$ 
  - (a) Apply Algorithm 7 to compute  $\mathbb{V}_m^b$  and  $\mathbb{H}_m$ .
  - (b) Apply Algorithm 5 to solve the problem :  $\mathbb{H}_m \star \mathcal{Y}_m \mathcal{Y}_m \star \mathcal{B} = \mathbb{V}_m^{bT} \star \mathcal{R}_0$
  - (c) Compute  $\mathfrak{X}_m = \mathfrak{X}_0 + \mathbb{V}_m^b \star \mathfrak{Y}_m$
- 4. If  $||\mathcal{R}_m||_F < tol$ , stop
- 5. else  $\mathfrak{X}_0 = \mathfrak{X}_m$  and go to Step 2.
- 6. **Output.**  $\mathfrak{X}_m \in \mathbb{R}^{n \times q \times n_3}$  the approximate solution of (38).

 $\Gamma(-\mathbf{B}) = \emptyset$ , where the block diagonal matrix  $\mathbf{H}_{\mathbf{m}}$  and  $-\mathbf{B}$  are defined by (4), and  $\Gamma(\mathbf{H}_m), \Gamma(-\mathbf{B})$  denotes the set of eigenvalues of the matrix  $\mathbf{H}_m$  and  $-\mathbf{B}$  respectively.

#### 6. Numerical experiments

This section performs some numerical tests for the Tensor Tubal-Global GMRES and Tensor Tubal-Global Golub Kahan methods to solve the linear tensor problem (1). All computations were carried out using the MATLAB R2018b environment with an Intel(R) Core i7-8550U CPU @1.80 GHz and processor 8 GB. The stopping criterion was

$$||\mathcal{R}_k||_F < \epsilon,$$

where  $\epsilon = 10^{-6}$  is a chosen tolerance and  $\mathcal{R}_k$  the m-th residual associated to the approximate solution  $\mathcal{X}_k$ . In all the presented tables, we reported the obtained residual norms to achieve the desired convergence, the iteration number and the corresponding cpu-time.

We will compare the results in our method to the results obtained by solving the equivalent problem  $\mathbf{C} = \mathbf{A}\mathbf{X} - \mathbf{X}\mathbf{B}$  where  $\mathbf{A}$ ,  $\mathbf{B}$  and  $\mathbf{C}$  are the matrices

$$\begin{split} \mathbf{A} &= \mathrm{Diag}(\widetilde{\mathcal{A}}) = \begin{pmatrix} \widetilde{\mathcal{A}}^{(1)} & & \\ & \widetilde{\mathcal{A}}^{(2)} & \\ & & \ddots & \\ & & \widetilde{\mathcal{A}}^{(n_3)} \end{pmatrix} = \mathrm{Block}\mathrm{Diag}(\widetilde{\mathcal{A}}^{(1)}, \dots, \widetilde{\mathcal{A}}^{(n_3)}), \\ \mathbf{B} &= \mathrm{Diag}(\widetilde{\mathfrak{B}}) = \mathrm{Block}\mathrm{Diag}(\widetilde{\mathfrak{B}}^{(1)}, \dots, \widetilde{\mathfrak{B}}^{(n_3)}) \\ \mathbf{C} &= \mathrm{Diag}(\widetilde{\mathfrak{C}}) = \mathrm{Block}\mathrm{Diag}(\widetilde{\mathfrak{C}}^{(1)}, \dots, \widetilde{\mathfrak{C}}^{(n_3)}) \end{split}$$

and the matrices  $\widetilde{\mathcal{A}}^{(i)}, \widetilde{\mathcal{B}}^{(i)}$  and  $\widetilde{\mathcal{C}}^{(i)}$ 's are the frontal slices of the tensor  $\widetilde{\mathcal{A}}, \widetilde{\mathcal{B}}$  and  $\widetilde{\mathcal{C}}$  respectively. Notice that **A** is a matrix of size  $(n_0 \times n_0), n_0 = n \times n_3$ , **B** is a matrix of size  $(s_0 \times s_0), s_0 = s \times n_3$ and **C** is a matrix of size  $(n_0 \times s_0)$ .

We compared the required CPU-times (in seconds) to achieve the convergence for the two methods:

- 1. **TBAS**: The tubal block Arnoldi–Sylvester method.
- 2. **BAS** : Resolution of Sylvester matrix equation :  $\mathbf{C} = \mathbf{AX} \mathbf{XB}$  by using the block Arnoldi-Sylvester method introduced in [5].
- 3. **tFOM** : The tensor full orthogonalization method.
- 4. tGMRES : The tensor generalized minimal residual method.

In Table 1, we reported the obtained relative residual norms, the total number of required iterations to achieve the convergence and the corresponding cpu-times for restarted Tubal Block Arnoldi(m). Consider the convection-diffusion equation:

$$\begin{cases} -\mu \,\Delta u + c^T \nabla \, u = f & \text{ in } [0,1]^N \\ u = 0 & \text{ in } \partial \Omega \end{cases}$$
(48)

The tensor  $\mathcal{A}, \mathcal{B}$  are obtained using  $n_3$  frontal slices (which are obtained from a standard finite difference discretization of (48)). In this example, the frontal slices are of size  $n \times n$  and  $s \times s$ , given as follows

$$\mathcal{A}^{(i)} = \frac{\mu}{h_1^2} \operatorname{tridiag}(-1, 2, -1) + \frac{a_i}{4h_1} \begin{bmatrix} 3 & -5 & 1 & & \\ 1 & 3 & -5 & \ddots & \\ & \ddots & \ddots & \ddots & 1 \\ & & 1 & 3 & -5 \\ & & & & 1 & 3 \end{bmatrix}$$

and

$$\mathcal{B}^{(i)} = \frac{\mu}{h_2^2} \text{tridiag}(-1, 2, -1) + \frac{b_i}{4h_2} \begin{bmatrix} 3 & -5 & 1 & & \\ 1 & 3 & -5 & \ddots & \\ & \ddots & \ddots & \ddots & 1 \\ & & 1 & 3 & -5 \\ & & & & 1 & 3 \end{bmatrix}$$

For  $i = 1, ..., n_3$ , we set  $a_i = i, b_i = n_3 + i, h_1 = \frac{1}{n+1}$  and  $h_2 = \frac{1}{s+1}$ . In this example, the right-hand side tensor C is constructed using the Matlab command C =

In this example, the right-hand side tensor C is constructed using the Matlab command  $C = rand(n, s, n_3)$ .

Table 1 reports on the obtained relative residual norms and the corresponding cpu-times to obtain the desired convergence. As can be seen from this table, the restarted Tensor Tubal block Arnoldi method (TBAS)(m) gives good results with a small cpu-times.

#### Conlusion

In this paper, we introduced new tensor krylov subspace methods for solving large Sylvester tensor equations. The proposed method uses the well-known T-product for tensors and tensor subspaces. We developed some new tensor products and the related algebraic properties. These new products will lead us to develop third-order the tensor FOM (tFOM), tensor GMRES (tGMRES), tubal Block Arnoldi and the tensor tubal Block Arnoldi method to solve large Sylvester tensor equation . We give some properties related to these method. The numerical

$n \backslash s \backslash k$	Method	# its.	$  \mathcal{R}_k  _F$	cpu-time in seconds
	TBAS	11	$5.55\times 10^{-7}$	3.03
$1000 \langle 3 \rangle 10$	$\mathbf{BAS}[5]$	65	$7.18\times 10^{-7}$	7.22
	tFOM	66	$7.57\times 10^{-7}$	57.69
	tGMRES	39	$9.50\times 10^{-7}$	25.34
	TBAS	12	$6.86\times 10^{-8}$	12.02
$2000 \backslash 3 \backslash 6$	BAS[5]	69	$3.25\times 10^{-7}$	17.61
	tFOM	69	$6.91\times 10^{-7}$	300.16
	tGMRES	41	$7.38\times 10^{-7}$	108.05

Table 1: Results for Example 2.  $\epsilon = 10^{-6}$ , and  $n_3 = 2$ 

experiments show that the restarted Tensor Tubal block Arnoldi method TBAS(m) gives the best results with a small cpu-times.

Author contribution All three authors have contributed in the same way.

Funding None.

**Data availability** Data sharing not applicable to this article as no data sets were generated or analyzed during the current study. We just generalized existing programs.

## Declarations

Ethical approval Not applicable. External Review board.

Competing interests The authors declare no competing interests.

Conflict of interest The authors declare that they have no conflict of interest.

## References

- R.H. Bartels, G.W. Stewart, Algorithm 432: solution of the matrix equation AX + XB = C, Circ. Syst. Signal Proc. 13 (1994) 820-826.
- [2] R. Bouyouli, K. Jbilou, R. Sadaka, H. Sadok, Convergence properties of some block Krylov subspace methods for multiple linear systems, J. Comput. Appl. Math., 196 (2006) 498-511.
- [3] K. Braman, Third-order tensors as linear operators on a space of matrices, Linear Algebra Appl., 433 (2010) 1241-1253.
- [4] F. BOUYGHF, A. MESSAOUDI, H. SADOK, An unified approach to Krylov subspace methods for solving linear systems, Numerical Algorithms Journal, https://doi.org/10.1007/s11075-023-01648-0.
- [5] A. El Guennouni, K. Jbilou, A.J. Riquet, Block Krylov subspace methods for solving large Sylvester equations, Numer. Algebra 29 (2002) 75-96.
- [6] M. El Guide, A. El Ichi, K. Jbilou, and R. Sadaka, "On tensor GMRES and Golub-Kahan methods via the T-product for color image processing", Electronic Journal of Linear Algebra, 37(2021) 524-543.
- [7] M. El Guide, A. El Ichi, K. Jbilou and F.P.A. Beik, "Tensor Krylov subspace methods via the Einstein product with applications to image and video processing", Applied Numerical Mathematics 181 (2022) 347–363.
- [8] A. El Ichi, K. Jbilou, and R. Sadaka, "On tensor tubal-Krylov subspace methods", Linear and Multilinear Algebra, 70 (2022) 7575-7598.
- [9] G.H. Golub, S. Nash, C. Van Loan, A Hessenberg–Schur method for the problem AX + XB = C, IEEC Trans. Automat. Contr. AC24 (1979) 909–913
- [10] A. El Ichi, Calcul tensoriel et Applications, Ph.D. Thesis, Univérsité du Littoral Cote d'Opale, Calais, 2021.

- [11] K. Glover, D. J. N. Limebeer, J. C. Doyle, E. M. Kasenally, and M. G. Safonov, A characterisation of all solutions to the four block general distance problem, SIAM J. Control Optim., 29 (1991) 283-324.
- [12] G. H. Golub and C. F. Van Loan, Matrix Computations, 3rd ed., Johns Hopkins University Press, Baltimore, 1996.
- [13] D.Y. Hu, L. Reichel, Krylov subspace methods for the Sylvester equation, Linear Algebra Appl. 174 (1992) 283-314
- [14] K. Jbilou A. Messaoudi H. Sadok Global FOM and GMRES algorithms for matrix equations, Appl. Num. Math., 31(1999), 49–63.
- [15] K. Jbilou, H. Sadok, and A. Tinzefte, Oblique projection methods for linear systems with multiple right-hand sides, Electron. Trans. Numer. Anal., 20 (2005) 119-138.
- [16] T. G. Kolda and B. W. Bader, Tensor Decompositions and Applications. SIAM Rev. 51 (2009) 455-500.
- [17] M. E. Kilmer and C. D. Martin, Factorization strategies for third-order tensors, Linear Algebra Appl., 435 (2011) 641-658.
- [18] M. E. Kilmer, K. Braman, N. Hao and R. C. Hoover, Third-order tensors as operators on matrices: a theoretical and computational framework with applications in imaging, SIAM J. Matrix Anal. Appl., 34 (2013) 148-172.
- [19] J. Lasalle and S. Lefschetz, Stability of Lyapunov Direct Methods, Academic Press, New York, 1961.
- [20] B. C. MOORE, Principal component analysis in linear systems: controllability, observability and model reduction, IEEE Trans. Auto. Control, AC-26 (1981) 17-31.
- [21] Y. Miao, L. Qi and Y. Wei, Generalized Tensor Function via the Tensor Singular Value Decomposition based on the T-Product, Lin. Alg. Appl., 590 (2020) 258-303.
- [22] Y. Saad, Numerical solution of large Lyapunov equations, in: M.A. Kaashoek, J.H. van Schuppen, A.C. Ran (Eds.), Proceedings of the International Symposium MTNS-89,

Boston, Signal Processing, Scattering, Operator Theory and Numerical Methods, 3 (1990) 503-511