# SOLUTIONS OF AN ELLIPTIC SYSTEM WITH A NEARLY CRITICAL EXPONENT

#### I.A. GUERRA

ABSTRACT. Consider the problem

$$\begin{aligned} -\Delta u_{\epsilon} &= v_{\epsilon}^{p} \quad v_{\epsilon} > 0 \quad \text{in} \quad \Omega, \\ -\Delta v_{\epsilon} &= u_{\epsilon}^{q_{\epsilon}} \quad u_{\epsilon} > 0 \quad \text{in} \quad \Omega, \\ u_{\epsilon} &= v_{\epsilon} = 0 \quad \text{on} \quad \partial\Omega, \end{aligned}$$

where  $\Omega$  is a bounded convex domain in  $\mathbb{R}^N$ , N > 2, with smooth boundary  $\partial \Omega$ . Here  $p, q_{\epsilon} > 0$ , and

$$\epsilon := \frac{N}{p+1} + \frac{N}{q_{\epsilon}+1} - (N-2).$$

This problem has positive solutions for  $\epsilon > 0$  (with  $pq_{\epsilon} > 1$ ) and no non-trivial solution for  $\epsilon \leq 0$ . We study the asymptotic behaviour of *least energy* solutions as  $\epsilon \to 0^+$ . These solutions are shown to blow-up at exactly one point, and the location of this point is characterized. In addition, the shape and exact rates for blowing up are given.

Résumé. Considéré le problème

$$\begin{array}{rcl} -\Delta u_{\epsilon} &=& v_{\epsilon}^{p} & v_{\epsilon} > 0 & \mathrm{en} & \Omega, \\ -\Delta v_{\epsilon} &=& u_{\epsilon}^{q_{\epsilon}} & u_{\epsilon} > 0 & \mathrm{en} & \Omega, \\ u_{\epsilon} &=& v_{\epsilon} &=& 0 & \mathrm{sur} & \partial\Omega, \end{array}$$

où  $\Omega$  est un domaine convexe et borné de  $\mathbb{R}^N$ , N > 2, avec la frontière régulière  $\partial \Omega$ . Ici  $p, q_{\epsilon} > 0$ , et

$$\epsilon := \frac{N}{p+1} + \frac{N}{q_{\epsilon}+1} - (N-2).$$

Ce problème a les solutions positives pour  $\epsilon > 0$  (avec  $pq_{\epsilon} > 1$ ) et non pas de solution non-trivial pour  $\epsilon \leq 0$ . Nous étudions le comportement asymptotique de solutions d'énergie minimale quand  $\epsilon \to 0^+$ . Ces solutions explosent en un seul point, et la localisation de ce point est characterisé. De plus, la forme et le rythme d'explosion sont donnés.

### 1. INTRODUCTION

We consider the elliptic system

$$-\Delta u_{\epsilon} = v_{\epsilon}^{p} \quad v_{\epsilon} > 0 \quad \text{in} \quad \Omega, \tag{1.1}$$

 $-\Delta v_{\epsilon} = u_{\epsilon}^{q_{\epsilon}} \quad u_{\epsilon} > 0 \quad \text{in} \quad \Omega,$ (1.2)

$$u_{\epsilon} = v_{\epsilon} = 0 \quad \text{on} \quad \partial\Omega, \tag{1.3}$$

<sup>1991</sup> Mathematics Subject Classification. Primary,—, Secondary,—.

*Key words and phrases.* Semilinear elliptic system, asymptotic behaviour, critical exponent. This research was supported by FONDECYT 3040059.

where  $\Omega$  is a bounded convex domain in  $\mathbb{R}^N$ , N > 2, with smooth boundary  $\partial \Omega$ . Here  $p, q_{\epsilon} > 0$ , and

$$\epsilon := \frac{N}{p+1} + \frac{N}{q_{\epsilon}+1} - (N-2).$$
(1.4)

When  $\epsilon \leq 0$ , there is no solution for (1.1)-(1.3), see [18] and [22]. On the other hand when  $\epsilon > 0$ , we can prove existence of solutions obtained by the variational method. In fact, for  $\epsilon > 0$ , the embedding  $W^{2,\frac{p+1}{p}}(\Omega) \hookrightarrow L^{q_{\epsilon}+1}(\Omega)$  is compact for any  $q_{\epsilon} + 1 > (p+1)/p$ , that is  $pq_{\epsilon} > 1$ . Using this, it is not difficult to show that there exists a function  $\bar{u}_{\epsilon}$  positive solution of the variational problem

$$S_{\epsilon}(\Omega) = \inf\left\{ \left\| \Delta u \right\|_{L^{\frac{p+1}{p}}(\Omega)} \mid u \in W^{2,\frac{p+1}{p}}(\Omega), \quad \left\| u \right\|_{L^{q_{\epsilon}+1}(\Omega)} = 1 \right\}, \tag{1.5}$$

see for example [23]. This solution satisfies  $-\Delta \bar{u}_{\epsilon} = \bar{v}_{\epsilon}^{p}$ ,  $-\Delta \bar{v}_{\epsilon} = S_{\epsilon}(\Omega)\bar{u}_{\epsilon}^{q_{\epsilon}}$ , in  $\Omega$ and  $\bar{u}_{\epsilon} = \bar{v}_{\epsilon} = 0$  on  $\partial\Omega$ . After a suitable multiples of  $\bar{u}_{\epsilon}$  and  $\bar{v}_{\epsilon}$ , we obtain  $u_{\epsilon}$  and  $v_{\epsilon}$ solving (1.1)-(1.3). We call  $(u_{\epsilon}, v_{\epsilon})$  the *least energy solution* to (1.1)-(1.3). For others existence results, we refer to [4], [7], [9], [15], and [19].

Note that by setting  $v_{\epsilon} = (-\Delta u_{\epsilon})^{1/p}$ , we can write the system (1.1)-(1.3) only in terms of  $u_{\epsilon}$ , that is

$$-\Delta(-\Delta u_{\epsilon})^{1/p} = u_{\epsilon}^{q_{\epsilon}} \quad u_{\epsilon} > 0 \quad \text{in} \quad \Omega$$
(1.6)

$$u_{\epsilon} = \Delta u_{\epsilon} = 0 \quad \text{on} \quad \partial \Omega.$$
 (1.7)

Concerning the least energy solutions, in [23] it was proved that  $S_{\epsilon}(\Omega) \to S$  as  $\epsilon \downarrow 0$ , where S is independent of  $\Omega$  and moreover is the best Sobolev constant for the inequality

$$\|u\|_{L^{q+1}(\mathbb{R}^N)} \le S^{-\frac{p}{p+1}} \|\Delta u\|_{L^{\frac{p+1}{p}}(\mathbb{R}^N)}$$
(1.8)

with p, q, N satisfying

$$\frac{N}{p+1} + \frac{N}{q+1} - (N-2) = 0.$$
(1.9)

This shows that the sequence  $\{u_{\epsilon}\}_{\epsilon>0}$  of least energy solutions of (1.6)-(1.7) satisfy

$$S_{\epsilon}(\Omega) = \frac{\int_{\Omega} |\Delta u_{\epsilon}|^{\frac{p+1}{p}} dx}{\|u_{\epsilon}\|_{L^{q_{\epsilon}+1}(\Omega)}^{\frac{p+1}{p}}} = S + o(1) \quad \text{as} \quad \epsilon \to 0.$$
(1.10)

Relation (1.9) defines a curve in  $\mathbb{R}^2_+$ , for the variables p and q. This curve is the so-called *Sobolev Critical Hyperbola*. By symmetry, we assume without restriction that

$$2/(N-2) 
(1.11)$$

For each fixed value of p, the strict inequality gives a lower bound for the dimension, i.e.  $N > \max\{2, 2(p+1)/p\}$ .

In this article, we shall study in detail the asymptotic behaviour of the variational solution  $u_{\epsilon}$ , of (1.6)–(1.7) as  $\epsilon \downarrow 0$ , that is, as  $q_{\epsilon}$  approaches from <u>below</u> to q given by the Sobolev Critical Hyperbola (1.9).

The asymptotic behaviour of the equation (1.6)-(1.7) as  $\epsilon \downarrow 0$  has already been studied for  $p = p^*$  and p = 1. Next we recall some of these results to introduce ours. The case  $p = p^*$  is equivalent to consider the single equation

The case p = p is equivalent to consider the single equation

$$-\Delta u_{\epsilon} = u_{\epsilon}^{p^{*}-\epsilon}$$
 in  $\Omega$ , and  $u_{\epsilon} = 0$  on  $\partial \Omega$ .

This problem was studied in [1, 10, 13, 20]. There, exact rates of blow-up were given and the location of blow-up points were characterized. One key ingredient was the Pohozaev identity and the observation that the solution  $u_{\epsilon}$ , scaled in the form  $||u_{\epsilon}||_{L^{\infty}(\Omega)}^{-1}u_{\epsilon}$  converges to U solution of

$$-\Delta U = U^{p^*}, \quad U(y) > 0 \quad \text{for} \quad y \in \mathbb{R}^N$$
(1.12)

$$U(0) = 1, \quad U \to 0, \quad \text{as} \quad |y| \to \infty,$$
 (1.13)

which is unique, explicit, and radially symmetric. For the location of blow-up and the shape of the solution away of the singularity, it was proved that a scaled  $u_{\epsilon}$ , given by  $\|u_{\epsilon}\|_{L^{\infty}(\Omega)}u_{\epsilon}$ , converges to the Green's function G, solution of  $-\Delta G(x, \cdot) = \delta_x$  in  $\Omega, G(x, \cdot) = 0$  on  $\partial\Omega$ . The location of blowing-up points are the critical points of  $\phi(x) := g(x, x)$  (in fact their minima, see [10]), where g(x, y) is the regular part of G(x, y), i.e

$$g(x,y) = G(x,y) - \frac{1}{(N-2)\sigma_N |x-y|^{N-2}}.$$

In [6], a similar result was proven in the case p = 1, (N > 4), where the problem is reduced to study (1.12)–(1.13) with the operator  $\Delta^2$  instead of  $-\Delta$ . Both cases give the blow-up rate

$$\epsilon \| u_{\epsilon} \|_{L^{\infty}(\Omega)}^2 \to C \text{ as } \epsilon \to 0.$$

for some explicit  $C := C(p, N, \Omega) > 0$ . We can ask ourselves if this behaviour is universal. We will see later that this is only a coincide.

Mimicking the above argument, we will study the asymptotic behaviour of the solution  $u_{\epsilon}$  of (1.6)–(1.7) as  $\epsilon \downarrow 0$ . We shall show that  $||u_{\epsilon}||_{L^{\infty}(\Omega)}^{-1} u_{\epsilon}$  converges, as  $\epsilon \downarrow 0$ , to the solution U of the problem

$$-\Delta U = V^p, \quad V(y) > 0 \quad \text{for} \quad y \in \mathbb{R}^N$$
 (1.14)

$$-\Delta V = U^{q}, \quad U(y) > 0 \quad \text{for} \quad y \in \mathbb{R}^{N}$$

$$(1.15)$$

$$U(0) = 1, \quad U \to 0, \quad V \to 0 \quad \text{as} \quad |y| \to \infty.$$
 (1.16)

In [5], it was proved that U and V are radially symmetric, if  $p \ge 1$  and  $U \in L^{q+1}(\mathbb{R}^N)$ and  $V \in L^{p+1}(\mathbb{R}^N)$ . This is the case when considering least energy solutions, see details in section 2. Thus U(r) := U(y) and V(r) := V(y) with r = |y|, moreover U and V are unique, and decreasing in r, see [16, 23]. There exist no explicit form of (U, V) for all  $p \ge 1$ , however to carry out the analysis it is sufficient to know the

asymptotic behaviour of (U, V) as  $r \to \infty$ , which was studied in [16]. They found

$$\lim_{r \to \infty} r^{N-2} V(r) = a \quad \text{and} \quad \begin{cases} \lim_{r \to \infty} r^{N-2} U(r) = b & \text{if} \quad p > \frac{N}{N-2} \\ \lim_{r \to \infty} \frac{r^{N-2}}{\log r} U(r) = b & \text{if} \quad p = \frac{N}{N-2} \\ \lim_{r \to \infty} r^{p(N-2)-2} U(r) = b & \text{if} \quad \frac{2}{N-2} (1.17)$$

The aim of this paper is to show the following results.

**Theorem 1.1.** Let  $u_{\epsilon}$  be a least energy solution of (1.6)–(1.7) and  $p \ge 1$ . Then a) there exists  $x_0 \in \Omega$  such that, after passing to a subsequence, we have

i) 
$$u_{\epsilon} \to 0 \in C^1(\Omega \setminus \{x_0\}),$$
 ii)  $v_{\epsilon} = |\Delta u_{\epsilon}|^{\frac{1}{p}} \to 0 \in C^1(\Omega \setminus \{x_0\})$ 

as  $\epsilon \to 0$  and

iii) 
$$|\Delta u_{\epsilon}|^{\frac{p+1}{p}} \to ||V||_{L^{p+1}(\mathbb{R}^{N})}^{p+1} \delta_{x_{0}} \quad as \quad \epsilon \to 0$$

in the sense of distributions.

b)  $x_0$  is a critical point of

$$\phi(x) := g(x,x) \quad if \quad p \in [N/(N-2), (N+2)/(N-2)) \quad and$$
 (1.18)

$$\tilde{\phi}(x) := \tilde{g}(x,x) \quad if \quad p \in (2/(N-2), N/(N-2))$$
(1.19)

for  $x \in \Omega$ . The function  $\tilde{g}(x,y)$  is defined for  $p \in (2/(N-2), N/(N-2))$  by

$$\tilde{g}(x,y) = \tilde{G}(x,y) - \frac{1}{(p(N-2)-2)(N-p(N-2))(N-2)^p \sigma_N^p |x-y|^{p(N-2)-2}}$$

where  $-\Delta \tilde{G}(x, \cdot) = G^p(x, \cdot)$  in  $\Omega$ ,  $\tilde{G}(x, \cdot) = 0$  on  $\partial \Omega$ .

We observe that regularity of  $\tilde{\phi}$  is needed to compute its critical points in b). We show next that  $\tilde{\phi}$  is regular. By definition of  $\tilde{G}$ , we have

$$\lim_{y \to x} |x - y|^{(p-1)(N-2)} \Delta \tilde{g}(x, y) = -\frac{pg(x, x)}{((N-2)\sigma_N)^{p-1}}$$
(1.20)

for  $x \in \Omega$ . Thus  $-\Delta \tilde{g}(x, \cdot) \in L^q(\Omega)$  for any  $q \in (N/2, N/(p(N-2) - N + 2))$ . This implies, by regularity, that  $\tilde{g}(x, \cdot) \in L^{\infty}(\Omega)$  and therefore  $\tilde{\phi}(x) = \tilde{g}(x, x), x \in \Omega$  is bounded. In addition, we define

$$\hat{g}(x,y) = \tilde{g}(x,y) + \frac{pg(x,x)|x-y|^{N-p(N-2)}}{(N-p(N-2))(2N-p(N-2)-2)((N-2)\sigma_N)^{p-1}} \quad (1.21)$$

and we have for any  $x \in \Omega$  that

$$\lim_{y \to x} |x - y|^{(p-2)(N-2)} \Delta \hat{g}(x, y) = -\frac{p(p-1)g(x, x)}{((N-2)\sigma_N)^{p-2}}.$$
(1.22)

Thus  $\hat{g}(x,y)$  is regular in y for x fixed. Since N > p(N-2), we take first y = x in (1.21) and then the gradient and we find  $\nabla_x \tilde{g}(x,x) = \nabla_x \hat{g}(x,x)$ . Hence  $\tilde{\phi}(x)$  is regular.

To state the next theorems we denote

$$\alpha = \frac{N}{q+1}$$
 and  $\beta = \frac{N}{p+1}$ ,

so the critical hyperbola (1.9) takes the form  $\alpha + \beta = N - 2$ .

Theorem 1.2. Let the assumptions of Theorem 1.1 be satisfied. Then

$$\begin{split} \lim_{\epsilon \to 0^+} \epsilon \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{(N-2)}{\alpha}} &= S^{\frac{1-pq}{p(q+1)}} \|U\|_{L^{q}(\mathbb{R}^{N})}^{q} \|V\|_{L^{p}(\mathbb{R}^{N})}^{p} |\phi(x_{0})| \quad if \quad p > \frac{N}{N-2} \\ \lim_{\epsilon \to 0^+} \epsilon^{\frac{\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{(N-2)}{\alpha}}}{\log(\|u_{\epsilon}\|_{L^{\infty}(\Omega)})} &= \frac{1}{\alpha} a^{\frac{N}{N-2}} S^{\frac{1-pq}{p(q+1)}} \|U\|_{L^{q}(\mathbb{R}^{N})}^{q} |\phi(x_{0})| \quad if \quad p = \frac{N}{N-2} \\ \lim_{\epsilon \to 0^+} \epsilon \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{p(N-2)-2}{\alpha}} &= S^{\frac{1-pq}{p(q+1)}} \|U\|_{L^{q}(\mathbb{R}^{N})}^{q(p+1)} |\tilde{\phi}(x_{0})| \quad if \quad p < \frac{N}{N-2}. \end{split}$$

In particular taking  $p = p^*$ , and using (1.9) we find that  $q = p^*$ . We recover the results in [13, 20], that is

$$\epsilon \|u_{\epsilon}\|^2_{L^{\infty}(\Omega)} \to C \quad \text{as} \quad \epsilon \to 0,$$
 (1.23)

for some explicitly given C > 0. See also [1] for the case  $\Omega = B_R$ .

When N > 4, we can take p = 1, and use (1.11) to find that q = (N+4)/(N-4). Here we recover the result in [2, 6], where they prove that (1.23) holds for some C > 0.

**Theorem 1.3.** Let the assumptions of Theorem 1.1 be satisfied. Then

$$\lim_{\epsilon \to 0} \|u_{\epsilon}\|_{L^{\infty}(\Omega)} v_{\epsilon}(x) = \|U\|_{L^{q}(\mathbb{R}^{N})}^{q} G(x, x_{0}), \quad and \tag{1.24}$$

$$\begin{cases}
\lim_{\epsilon \to 0} \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}} u_{\epsilon}(x) = \|V\|_{L^{p}(\mathbb{R}^{N})}^{p} G(x, x_{0}) \quad if \quad p > \frac{N}{N-2} \\
\lim_{\epsilon \to 0} \frac{\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}} u_{\epsilon}(x) = \frac{1}{\alpha} a^{\frac{N}{N-2}} G(x, x_{0}) \quad if \quad p = \frac{N}{N-2} \\
\lim_{\epsilon \to 0} \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{1}{\alpha}(\beta+p(N-2)-N)} u_{\epsilon}(x) = \|U\|_{L^{q}(\mathbb{R}^{N})}^{pq} \tilde{G}(x, x_{0}) \quad if \quad p < \frac{N}{N-2}
\end{cases}$$
(1.25)

where all the convergences in  $C^{1,\alpha}(w)$  with w any neighborhood of  $\partial\Omega$  not containing  $x_0$ .

**Remark 1.4.** For  $p < \frac{N}{N-2}$ , the convergence in (1.25) can be improved to  $C^{3,\alpha}(\omega)$ . See the proof of the theorem.

**Remark 1.5.** By (2.13), we find that  $\lim_{\epsilon \to 0} \|v_{\epsilon}\|_{L^{\infty}(\Omega)} = V(0) \lim_{\epsilon \to 0} \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}}$ . So, in addition, when p = 1 we have that

$$\epsilon \|v_{\epsilon}\|_{L^{\infty}(\Omega)}^{2(N-4)/N} \to CV(0)^{2(N-4)/N} \quad as \quad \epsilon \to 0.$$

We can extend these results to the problem

$$-\Delta(-\Delta u_{\epsilon})^{1/p} = u_{\epsilon}^{q} + \epsilon u_{\epsilon} \quad u_{\epsilon} > 0 \quad \text{in} \quad \Omega$$
(1.26)

$$u_{\epsilon} = \Delta u_{\epsilon} = 0 \quad \text{on} \quad \partial\Omega \tag{1.27}$$

with  $\epsilon \to 0$ . The existence of positive solutions for this problem can be found in [15] and [19] in the case of a ball. See [14] for related results for p = 1.

**Theorem 1.6.** Let the assumptions of Theorem 1.1 be satisfied. Then

$$\lim_{\epsilon \to 0^+} \epsilon \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{2-\frac{2}{\alpha}} = \|U\|_{L^{2}(\mathbb{R}^{N})}^{-2} \|U\|_{L^{q}(\mathbb{R}^{N})}^{q} \|V\|_{L^{p}(\mathbb{R}^{N})}^{p} |\phi(x_{0})| \quad if \quad p > \frac{N}{N-2}$$

$$and \quad \alpha > 1, \ N > 4 \qquad (1.28)$$

$$\lim_{\epsilon \to 0^+} \epsilon \frac{\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{2-\overline{\alpha}}}{\log(\|u_{\epsilon}\|_{L^{\infty}(\Omega)})} = \frac{1}{\alpha} a^{\frac{N}{N-2}} \|U\|_{L^{2}(\mathbb{R}^{N})}^{-2} \|U\|_{L^{q}(\mathbb{R}^{N})}^{q} |\phi(x_{0})| \quad if \quad p = \frac{N}{N-2}$$

$$and \quad 2 - \frac{2}{\alpha} = 3 - (\frac{N}{N-2})^{2} > 0, \qquad (1.29)$$

$$\lim_{\epsilon \to 0^+} \epsilon \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{2-\frac{2+N-p(N-2)}{\alpha}} = \|U\|_{L^{2}(\mathbb{R}^{N})}^{-2} \|U\|_{L^{q}(\mathbb{R}^{N})}^{q(p+1)} |\tilde{\phi}(x_{0})| \quad if \quad \frac{N+4}{2(N-2)} 
$$and \quad \alpha > \frac{2+N-p(N-2)}{2} \tag{1.30}$$$$

$$\lim_{\epsilon \to 0^+} \epsilon \log(\|u_{\epsilon}\|_{L^{\infty}(\Omega)}) = \frac{\|U\|_{L^{q}(\mathbb{R}^{N})}^{q} \|V\|_{L^{p}(\mathbb{R}^{N})}^{p}}{b^{2}} |\phi(x_{0})| \quad if \quad N = 4, \ p = q = 3, \ (1.31)$$

$$\lim_{\epsilon \to 0^+} \epsilon \log(\|u_{\epsilon}\|_{L^{\infty}(\Omega)}) \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{3-q}{2}} = \frac{\|U\|_{L^{q}(\mathbb{R}^{N})}^{q(p+1)}}{b^{2}} |\tilde{\phi}(x_{0})| \quad if \quad p = \frac{N+4}{2(N-2)} < \frac{N}{N-2},$$
  
and  $q \leq 3$  (1.32)

Note that N > 4 (integer) is equivalent to  $3 - (N/(N-2))^2 > 0$  and also to (N+4)/(2(N-2)) < N/(N-2). This implies that (1.29) holds for p = N/(N-2) and N > 4, and (1.32) holds for  $p = \frac{N+4}{2(N-2)}$ ,  $q \leq 3$  and N > 4.

For example, p = 1 gives q + 1 = 2N/(N - 4) and provided that N > 8, we get

$$\lim_{\epsilon \to 0^+} \epsilon \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{2(N-8)}{N-4}} = C_1 |\tilde{\phi}(x_0)|.$$

For N = 8 and p = 1, we have

$$\lim_{\epsilon \to 0^+} \epsilon \log(\|u_{\epsilon}\|_{L^{\infty}(\Omega)}) = C_1 |\tilde{\phi}(x_0)|.$$

## 2. Preliminaries

Before proving the main theorem, we need some properties of  $u_{\epsilon}$ . Using that  $u_{\epsilon}$  is a minimizing sequence, we have

$$\int_{\Omega} (\Delta u_{\epsilon})^{\frac{p+1}{p}} dx = \int_{\Omega} v_{\epsilon} \Delta u_{\epsilon} dx = \int_{\Omega} u_{\epsilon} \Delta v_{\epsilon} dx = \int_{\Omega} u_{\epsilon}^{q_{\epsilon}+1} dx.$$

Then  $[S + o(1)] \|u_{\epsilon}\|_{L^{q_{\epsilon}+1}(\Omega)}^{\frac{p+1}{p}} = \|u_{\epsilon}\|_{L^{q_{\epsilon}+1}(\Omega)}^{q_{\epsilon}+1}$  implies

$$\lim_{\epsilon \to 0} \int_{\Omega} u_{\epsilon}^{q_{\epsilon}+1} dx = S^{\frac{p_q-1}{p(q+1)}}.$$
(2.1)

**Lemma 2.1.** The minimizing sequence  $u_{\epsilon}$  of (1.10) is such that

$$||u_{\epsilon}||_{L^{\infty}(\Omega)} \to \infty$$

moreover  $\|(-\Delta u_{\epsilon})^{1/p}\|_{L^{\infty}(\Omega)} = \|v_{\epsilon}\|_{L^{\infty}(\Omega)} \to \infty \quad as \quad \epsilon \to 0.$ 

Proof. If  $||u_{\epsilon}||_{L^{\infty}(\Omega)} \to \infty$  then by regularity, we find  $||v_{\epsilon}||_{L^{\infty}(\Omega)} \to \infty$ , see [12, Theorem 3.7]. Now, assume that  $||u_{\epsilon}||_{L^{\infty}(\Omega)} \leq M$  and  $||v_{\epsilon}||_{L^{\infty}(\Omega)} \leq M$ , by elliptic regularity, we have that

$$\|v_{\epsilon}\|_{C^{2+\alpha}(\bar{\Omega})} \le M$$
 and  $\|u_{\epsilon}\|_{C^{2+\alpha}(\bar{\Omega})} \le M$ 

with  $\alpha \in (0,1)$  and some constant M. This implies that there exists  $u^*, v^* \in C^2(\bar{\Omega})$ , such that

$$u_{\epsilon} \to u^*$$
 in  $C^2(\bar{\Omega})$ ,  $v_{\epsilon} \to v^*$  in  $C^2(\bar{\Omega})$  as  $\epsilon \to 0$ .

Hence  $u^*$  satisfies

$$0 \neq \int_{\Omega} (\Delta u^*)^{\frac{p+1}{p}} dx = S \left[ \int_{\Omega} (u^*)^{q+1} dx \right]^{\frac{(p+1)}{p(q+1)}}$$

which contradicts that S can be achieved by a minimizer in a bounded domain, see [23]. In other words there exists no non trivial solution for

$$-\Delta u^* = (v^*)^p, \quad v > 0 \quad \text{in} \quad \Omega \tag{2.2}$$

$$-\Delta v^* = (u^*)^q, \quad u > 0 \quad \text{in} \quad \Omega \tag{2.3}$$

$$u^* = v^* = 0 \quad \text{on} \quad \partial\Omega \tag{2.4}$$

in a convex bounded domain, with p, q satisfying (1.9), see [18], [22].

For any  $\epsilon > 0$ , let  $(u_{\epsilon}, v_{\epsilon})$  be a solution of (1.1–1.3). By the Pohozaev inequality, see [18] or [22], we have for any  $\tilde{\alpha}, \tilde{\beta} \in \mathbb{R}$  that

$$\left(\frac{N}{q_{\epsilon}+1}-\tilde{\alpha}\right)\int_{\Omega} u_{\epsilon}^{q_{\epsilon}+1} dx + \left(\frac{N}{p+1}-\tilde{\beta}\right)\int_{\Omega} v_{\epsilon}^{p+1} dx \qquad (2.5)$$

$$+(N-2-\tilde{\alpha}-\tilde{\beta})\int_{\Omega} (\nabla u_{\epsilon}, \nabla v_{\epsilon}) \, dx = -\int_{\partial\Omega} (\nabla u_{\epsilon}, n) (\nabla v_{\epsilon}, x-y) \, ds.$$
(2.6)

We choose  $\tilde{\alpha} + \tilde{\beta} = N - 2$ ,  $\tilde{\alpha} = \alpha$  and so  $\tilde{\beta} = \beta$ . This implies that

$$\epsilon \int_{\Omega} u_{\epsilon}^{q_{\epsilon}+1} dx = -\int_{\partial\Omega} \frac{\partial u_{\epsilon}}{\partial n} \frac{\partial v_{\epsilon}}{\partial n} (n, x - y) ds.$$
(2.7)

Since  $u_{\epsilon}$  becomes unbounded as  $\epsilon \to 0$  we choose  $\mu = \mu(\epsilon)$  and  $x_{\epsilon} \in \Omega$  such that

$$\mu^{\alpha_{\epsilon}} u_{\epsilon}(x_{\epsilon}) = 1$$

where  $\alpha_{\epsilon} = N/(q_{\epsilon} + 1)$ . Note that  $\mu \to 0$  as  $\epsilon \to 0$ .

First we claim that  $x_{\epsilon}$  stays away from the boundary. This is consequence of moving plane method and interior estimates [8], [11]. Let  $\phi_1$  the positive eigenvalue

of  $(-\Delta, H_0^1(\Omega))$ , normalized to  $\max_{x \in \Omega} \phi_1(x) = 1$ . Since  $p \ge 1$ , multiplying by  $\phi_1$  we obtain

$$\lambda_1 \int_{\Omega} u_{\epsilon} \phi_1 = \int_{\Omega} v_{\epsilon}^p \phi_1 \ge 2\lambda_1 \int_{\Omega} v_{\epsilon} \phi_1 - C \int_{\Omega} \phi_1$$
$$\lambda_1 \int_{\Omega} v_{\epsilon} \phi_1 = \int_{\Omega} u_{\epsilon}^{q_{\epsilon}} \phi_1 \ge 2\lambda_1 \int_{\Omega} u_{\epsilon} \phi_1 - C \int_{\Omega} \phi_1$$

for some  $C = C(p, q, \lambda_1) > 0$ . Hence  $\int_{\Omega} u_{\epsilon} \phi_1 \leq (C/\lambda_1) \int_{\Omega} \phi_1$  which implies  $\int_{\Omega'} u_{\epsilon} \leq C(\Omega')$  with  $\Omega' \subset \Omega$  and  $\int_{\Omega'} v_{\epsilon} \leq C(\Omega')$ . Using the moving planes method [11], we find that there exist  $t_0 \alpha > 0$  such that

$$u_{\epsilon}(x-t\nu)$$
 and  $v_{\epsilon}(x-t\nu)$  are nondecreasing for  $t \in [0, t_0]$ ,

 $\nu \in \mathbb{R}^N$  with  $|\nu| = 1$ , and  $(\nu, n(x)) \ge \alpha$  and  $x \in \partial\Omega$ . Therefore we can find  $\gamma, \delta$  such that for any  $x \in \{z \in \overline{\Omega} : d(z, \partial\Omega) < \delta\} = \Omega_{\delta}$  there exists a measurable set  $\Gamma_x$  with (i)  $meas(\Gamma_x) \ge \gamma$ , (ii)  $\Gamma_x \subset \Omega \setminus \overline{\Omega}_{\delta/2}$ , and (iii)  $u_{\epsilon}(y) \ge u_{\epsilon}(x)$  and  $v_{\epsilon}(y) \ge v_{\epsilon}(x)$  for any  $y \in \Gamma_x$ . Then for any  $x \in \Omega_{\delta}$ , we have

$$u_{\epsilon}(x) \leq \frac{1}{meas(\Gamma_{x})} \int_{\Gamma_{x}} u_{\epsilon}(y) dy \leq \frac{1}{\gamma} \int_{\Omega_{\delta}} u_{\epsilon} \leq C(\Omega_{\delta}), \text{ and}$$
$$v_{\epsilon}(x) \leq \frac{1}{meas(\Gamma_{x})} \int_{\Gamma_{x}} v_{\epsilon}(y) dy \leq \frac{1}{\gamma} \int_{\Omega_{\delta}} v_{\epsilon} \leq C(\Omega_{\delta}).$$

Hence if  $u_{\epsilon}(x_{\epsilon}) \to \infty$ , this implies that  $x_{\epsilon}$  will stay out of  $\Omega_{\delta}$  a neighbordhood of the boundary. This proves the claim.

Let  $x_{\epsilon} \to x_0 \in \Omega$ . We define a family of rescaled functions

$$u_{\epsilon,\mu}(y) = \mu^{\alpha_{\epsilon}} u_{\epsilon}(\mu^{1-\epsilon/2}y + x_{\epsilon})$$
(2.8)

$$v_{\epsilon,\mu}(y) = \mu^{\beta} v_{\epsilon}(\mu^{1-\epsilon/2}y + x_{\epsilon})$$
(2.9)

and find using the definitions of  $\epsilon$ ,  $\alpha_{\epsilon}$  and  $\beta$ , that

$$-\Delta u_{\epsilon,\mu} = v^p_{\epsilon,\mu} \mu^{\alpha_{\epsilon}+2-\epsilon-p\beta} = v^p_{\epsilon,\mu} \quad \text{in} \quad \Omega_{\epsilon}$$
(2.10)

$$-\Delta v_{\epsilon,\mu} = u_{\epsilon,\mu}^{q_{\epsilon}} \mu^{\beta+2-\epsilon-q_{\epsilon}\alpha_{\epsilon}} = u_{\epsilon,\mu}^{q_{\epsilon}} \quad \text{in} \quad \Omega_{\epsilon}$$
(2.11)

 $u_{\epsilon,\mu} = v_{\epsilon,\mu} = 0 \quad \text{on} \quad \partial\Omega_{\epsilon}.$  (2.12)

By equicontinuity and using Arzela-Ascoli, we have that

$$u_{\epsilon,\mu} \to U \quad \text{and} \quad v_{\epsilon,\mu} \to V \quad \text{as} \quad \epsilon \to 0.$$
 (2.13)

in  $C^2(K)$  for any K compact in  $\mathbb{R}^N$ , where (U, V) satisfies (1.14)-(1.16). Now extending  $u_{\epsilon,\mu}$  and  $v_{\epsilon,\mu}$  by zero outside  $\Omega_{\epsilon}$  and using (2.1), by the argument in [21] or [23], we have that  $u_{\epsilon,\mu} \to \overline{U}$  strongly (up to a subsequence) in  $W^{2,\frac{p+1}{p}}(\mathbb{R}^N)$ . In the limit  $\overline{U} \in L^{q+1}(\mathbb{R}^N)$  and  $\overline{V} := (-\Delta \overline{U})^{\frac{1}{p}} \in L^{p+1}(\mathbb{R}^N)$ , and they satisfy (1.14)-(1.16). Since  $p \ge 1$ , the solution  $(\overline{U}, \overline{V})$  is unique and radially symmetric, see [5]. In addition the radial solutions are unique [16, 23], so  $\overline{U} \equiv U$  and  $\overline{V} \equiv V$ , consequently

$$\int_{\mathbb{R}^N} [u_{\epsilon,\mu} - U]^{q+1}(y) \, dy \to 0 \quad \int_{\mathbb{R}^N} [v_{\epsilon,\mu} - V]^{p+1}(y) \, dy \to 0.$$
(2.14)

**Lemma 2.2.** There exists  $\delta > 0$  such that

 $\delta \leq \mu^\epsilon \leq 1.$ 

*Proof.* Since  $\mu \to 0$ , we have  $\mu^{\epsilon} \leq 1$ . By (2.14), we get  $\int_{B_1} u_{\epsilon,\mu}^{q_{\epsilon}+1} dx \geq M$ , but

$$M \leq \int_{B_1} u_{\epsilon,\mu}^{q_{\epsilon}+1} dx = \mu^{\epsilon N/2} \int_{|y-x_{\epsilon}| \leq \mu^{1-\epsilon/2}} u_{\epsilon}^{q_{\epsilon}+1}(y) dy \leq \mu^{\epsilon N/2} \int_{\Omega} u_{\epsilon}^{q_{\epsilon}+1}(y) dy \qquad (2.15)$$

Using the convergence (2.1), we obtain the result.

**Lemma 2.3.** There exists K > 0 such that the solution  $(u_{\epsilon,\mu}, v_{\epsilon,\mu})$  satisfies

$$u_{\epsilon,\mu}(y) \le KU(y) \quad v_{\epsilon,\mu}(y) \le KV(y) \quad \forall y \in \mathbb{R}^N.$$
 (2.16)

We prove this lemma in section 2.3.

**Lemma 2.4.** There exists a constant C > 0 such that

$$\epsilon \leq C\mu^{N-2}h(\mu) \quad with \quad h(\mu) = \begin{cases} 1 & \text{for } p > N/(N-2) \\ |\log(\mu)| & \text{for } p = N/(N-2) \\ \mu^{(p(N-2)-N)} & \text{for } p < N/(N-2). \end{cases}$$
(2.17)

*Proof.* We will establish the following

$$\int_{\partial\Omega} \frac{\partial u_{\epsilon}}{\partial n} \frac{\partial v_{\epsilon}}{\partial n}(n,x) \, dx \le C \mu^{N-2} h(\mu)$$

and from here the result follows applying (2.7). We claim that

$$\left|\frac{\partial u_{\epsilon}}{\partial n}\right| \le C\mu^{\alpha_{\epsilon}} \quad \left|\frac{\partial v_{\epsilon}}{\partial n}\right| \le C\mu^{\beta}h(\mu)$$

In the following M is a positive constant that can vary from line to line and we shall use systematically Lemma 2.2.

For p > N/(N-2), using that  $-p\beta + N = \beta$ , we have

$$\int\limits_{\Omega} v_{\epsilon}^{p}(x) \, dx \leq M \mu^{-p\beta + N(1-\epsilon/2)} \int\limits_{\mathbb{R}^{N}} V^{p}(y) \, dy \leq M \mu^{\beta}$$

and by (2.16) there exists M > 0 such that

$$v_{\epsilon}^{p}(x) \le M \frac{\mu^{\beta+p(N-2)-N-p(N-2)\epsilon/2}}{|x-x_{0}|^{p(N-2)}}.$$
(2.18)

for  $x \neq x_0$ . Using that  $\beta < \beta + p(N-2) - N$ , by Lemma 5.1 we find  $\left|\frac{\partial v_{\epsilon}}{\partial n}\right| \leq C\mu^{\beta}$ . For  $u_{\epsilon}$ , using that  $-q_{\epsilon}\alpha_{\epsilon} + N = \alpha_{\epsilon}$ ,

$$\int_{\Omega} u_{\epsilon}^{q_{\epsilon}} dx \le M \mu^{-q_{\epsilon}\alpha_{\epsilon} + N(1-\epsilon/2)} \int_{\mathbb{R}^{N}} U^{q}(y) \, dy \le M \mu^{\alpha_{\epsilon}}$$

and by (2.16) there exist M > 0 such

$$u_{\epsilon}^{q_{e}}(x) \le M \frac{\mu^{-q_{\epsilon}\alpha_{\epsilon}+q_{\epsilon}(N-2)-q_{\epsilon}(N-2)\epsilon/2}}{|x-x_{0}|^{q_{\epsilon}(N-2)}}$$
(2.19)

for  $x \neq x_0$ . Using that  $\alpha_{\epsilon} < \alpha_{\epsilon} - N + q_{\epsilon}(N-2)$ , by Lemma 5.1, we obtain  $\left|\frac{\partial u_{\epsilon}}{\partial n}\right| \leq C \mu^{\alpha_{\epsilon}}$ . For p < N/(N-2), we have

$$\int_{\Omega} v_{\epsilon}^{p} dx \leq M \mu^{-p\beta + p(N-2)(1-\epsilon/2)} \lim_{\mu \to 0} \frac{1}{\mu^{(p(N-2)-N)(1-\epsilon/2)}} \int_{B_{\frac{1}{\mu^{1-\epsilon/2}}}(x_{\epsilon})} V^{p}(y) dy \quad (2.20)$$
$$\leq M \mu^{\beta + (p(N-2)-N)} \quad (2.21)$$

and for  $x \neq x_0$ , we find (2.18) for  $v_{\epsilon}$  and for  $u_{\epsilon}$  we have

$$\int_{\Omega} u_{\epsilon}^{q_{\epsilon}} \leq M \mu^{-q_{\epsilon}\alpha_{\epsilon} + N(1-\epsilon/2)} \int_{\mathbb{R}^{N}} U^{q}(y) \, dy \leq M \mu^{\alpha_{\epsilon}}$$

and by (2.16) there exist M > 0 such that

$$u_{\epsilon}^{q_{\epsilon}}(x) \le M \frac{\mu^{-q_{\epsilon}\alpha_{\epsilon}+q_{\epsilon}(p(N-2)-2)-q_{\epsilon}(p(N-2)-2)\epsilon/2}}{|x-x_{0}|^{q_{\epsilon}(p(N-2)-2)}}$$
(2.22)

for  $x \neq x_0$ . From these estimates we prove the claim applying Lemma 5.1 and noting that  $\alpha_{\epsilon} < \alpha_{\epsilon} - N + q_{\epsilon}(p(N-2)-2) + (p+1)\epsilon/\alpha_{\epsilon}$ . For the case p = N/(N-2), we proceed as before noting that

$$\int_{\Omega} v_{\epsilon}^{p} dx \le M \mu^{-p\beta + N(1 - \epsilon/2)} |\log(\mu)| \lim_{\mu \to 0} \frac{1}{|\log(\mu)|} \int_{B_{\frac{1}{\mu^{1 - \epsilon/2}}}(x_{\epsilon})} V^{p}(y) dy \le M |\log(\mu)| \mu^{\beta}$$

and for  $x \neq x_0$  we have (2.18). Similarly to (2.22), we obtain that for  $x \neq x_0$ , there exist M > 0 such that

$$u_{\epsilon}^{q_{\epsilon}}(x) \le M \frac{\mu^{-q_{\epsilon}\alpha_{\epsilon}+q_{\epsilon}(N-2)-q_{\epsilon}(N-2)\epsilon/2}}{|x-x_{0}|^{q_{\epsilon}(N-2)}} \log(|x-x_{0}|^{-1+\epsilon/2})^{q_{\epsilon}}.$$
(2.23)

Using this and proceeding and before we prove the claim and the lemma follows.  $\Box$ 

## Lemma 2.5.

$$|\mu^{\epsilon} - 1| = O(\mu^{N-2}h(\mu)\log\mu)$$

*Proof.* By the theorem of the mean  $|\mu^{\epsilon} - 1| = |\mu^{s\epsilon} \epsilon \log \mu|$  for some  $s \in (0, 1)$  and therefore (2.17) gives the result.

## 3. Proof of Theorem 1.2 and 1.3

*Proof of Theorem 1.3.* We start by proving the case  $p > \frac{N}{N-2}$ . We have

$$-\Delta(\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}}u_{\epsilon}) = \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}}v_{\epsilon}^{p} \quad \text{in} \quad \Omega,$$
(3.1)

$$-\Delta(\|u_{\epsilon}\|_{L^{\infty}(\Omega)}v_{\epsilon}) = \|u_{\epsilon}\|_{L^{\infty}(\Omega)}u_{\epsilon}^{q_{\epsilon}} \quad \text{in} \quad \Omega,$$
(3.2)

$$u_{\epsilon} = v_{\epsilon} = 0 \quad \text{on} \quad \partial\Omega. \tag{3.3}$$

We integrate the right hand side of (3.1)

$$\int_{\Omega} \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}} v_{\epsilon}^{p} dx = \mu^{-(p+1)\beta+N+N\epsilon/2} \int_{\Omega_{\epsilon}} v_{\epsilon,\mu}^{p}(y) dy$$

But  $N - (p+1)\beta = 0$ , so using (2.16) by dominated convergence and Lemma 2.5, we get

$$\lim_{\epsilon \to 0} \int_{\Omega} \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}} v_{\epsilon}^{p} dx = \int_{\mathbb{R}^{N}} V^{p}(y) dy = \|V\|_{L^{p}(\mathbb{R}^{N})} < \infty.$$

Similarly, now using

$$\int_{\Omega} \|u_{\epsilon}\|_{L^{\infty}(\Omega)} u_{\epsilon}^{q_{\epsilon}} dx = \mu^{-(q_{\epsilon}+1)\alpha_{\epsilon}+N+N\epsilon/2} \int_{\Omega_{\epsilon}} u_{\epsilon,\mu}^{q_{\epsilon}} dx \to \|U\|_{L^{q}(\mathbb{R}^{N})} < \infty$$
(3.4)

as  $\epsilon \to 0$ . Also using the bound (2.16), we find

$$\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}} v_{\epsilon}^{p}(x) \leq \frac{M\mu^{-(p+1)\beta+p(N-2)-p(N-2)\epsilon/2}}{|x-x_{0}|^{p(N-2)}}$$

for  $x \neq x_0$  and some M > 0. But  $-(p+1)\beta + p(N-2) > 0$  and Lemma 2.2 then  $\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}} v_{\epsilon}^{p}(x) \to 0$  for  $x \neq x_0$ . Also we have

$$||u_{\epsilon}||_{L^{\infty}(\Omega)}u_{\epsilon}^{q_{\epsilon}}(x) \leq \frac{M\mu^{-(q_{\epsilon}+1)\alpha_{\epsilon}+q_{\epsilon}(N-2)-q_{\epsilon}(N-2)\epsilon/2}}{|x-x_{0}|^{q_{\epsilon}(N-2)}}.$$

for  $x \neq x_0$  and some M > 0. But  $-(q_{\epsilon} + 1)\alpha_{\epsilon} + q_{\epsilon}(N-2) > 0$  and Lemma 2.2 then  $\|u_{\epsilon}\|_{L^{\infty}(\Omega)} u_{\epsilon}^{q_{\epsilon}}(x) \to 0$  for  $x \neq x_0$ .

From here we have

$$-\Delta(\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{p}{\alpha}}u_{\epsilon}) \to \|V\|_{L^{p}(\mathbb{R}^{N})}^{p}\delta_{x=x_{0}} \quad \text{and} \quad -\Delta(\|u_{\epsilon}\|_{L^{\infty}(\Omega)}v_{\epsilon}) \to \|U\|_{L^{q}(\mathbb{R}^{N})}^{q}\delta_{x=x_{0}}$$

in the sense of distributions in  $\Omega$ , as  $\epsilon \to 0$ . Let  $\omega$  be any neighborhood of  $\partial \Omega$  not containing  $x_0$ . By regularity theory, see Lemma 5.1, we find

$$\|\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}}u_{\epsilon}\|_{C^{1,\alpha}(w)} \leq C\left[\|\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}}v_{\epsilon}^{p}\|_{L^{1}(\Omega)} + \|\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}}v_{\epsilon}^{p}\|_{L^{\infty}(w)}\right]$$

and a similar bound for  $||||u_{\epsilon}||_{L^{\infty}(\Omega)}v_{\epsilon}||_{C^{1,\alpha}(w)}$ . Consequently

$$\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}} u_{\epsilon} \to \|V\|_{L^{p}(\mathbb{R}^{N})}^{p} G \quad \text{in} \quad C^{1,\alpha}(w) \quad \text{as} \quad \epsilon \to 0.$$
(3.5)

and

$$\|u_{\epsilon}\|_{L^{\infty}(\Omega)}v_{\epsilon} \to \|U\|^{q}_{L^{q}(\mathbb{R}^{N})}G \quad \text{in} \quad C^{1,\alpha}(w) \quad \text{as} \quad \epsilon \to 0.$$
(3.6)

For the case p < N/(N-2), we proceed as before and we have (3.4) and the bound

$$\|u_{\epsilon}\|_{L^{\infty}(\Omega)}u_{\epsilon}^{q_{\epsilon}}(x) \leq \frac{M\mu^{-(q_{\epsilon}+1)\alpha_{\epsilon}+q_{\epsilon}(p(N-2)-2)-q_{\epsilon}(p(N-2)-2)\epsilon/2}}{|x-x_{0}|^{q_{\epsilon}(p(N-2)-2)}}$$

for  $x \neq x_0$  and some M > 0. Using that  $-(q_{\epsilon}+1)\alpha_{\epsilon} + q(p(N-2)-2) = 2(p+1) > 0$ and Lemma 2.2, we get  $\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}} u_{\epsilon}^{q_{\epsilon}}(x) \to 0$  for  $x \neq x_0$  and hence

$$\|u_{\epsilon}\|_{L^{\infty}(\Omega)}v_{\epsilon} \to \|U\|_{L^{q}(\mathbb{R}^{N})}^{q}G \quad \text{in} \quad C^{1,\alpha}(w) \quad \text{as} \quad \epsilon \to 0.$$
(3.7)

Now we claim that

$$\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{1}{\alpha}(\beta+p(N-2)-N)}u_{\epsilon} \to \|U\|_{L^{q}(\mathbb{R}^{N})}^{pq}\tilde{G} \quad \text{in} \quad C^{1,\alpha}(w) \quad \text{as} \quad \epsilon \to 0.$$
(3.8)

We have

$$-\Delta(\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{1}{\alpha}(\beta+p(N-2)-N)}u_{\epsilon}) = \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{1}{\alpha}(\beta+p(N-2)-N)}v_{\epsilon}^{p} = \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{p}v_{\epsilon}^{p}.$$

Since the last term converges to  $(||U||^q_{L^q(\mathbb{R}^N)}G)^p$  in  $C^{1,\alpha}(\omega)$  as  $\epsilon \to 0$  and  $p \ge 1$ , we have

$$\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{1}{\alpha}(\beta+p(N-2)-N)}u_{\epsilon} \to \|U\|_{L^{q}(\mathbb{R}^{N})}^{pq}\tilde{G} \quad \text{in} \quad C^{3,\alpha}(w) \quad \text{as} \quad \epsilon \to 0.$$

For the remaining case p = N/(N-2), we have as  $\epsilon \to 0$ , the convergence

$$\int_{\Omega} \frac{\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{p}{\alpha}}}{|\log(\|u_{\epsilon}\|_{L^{\infty}(\Omega)})|} v_{\epsilon}^{p} dx = \frac{\mu^{-(p+1)\beta+N+N\epsilon/2}}{\alpha_{\epsilon}|\log(\mu)|} \int_{\Omega_{\epsilon}} v_{\epsilon,\mu}^{p} dy \to \frac{1}{\alpha} \lim_{r \to \infty} V(r)^{\frac{N}{N-2}} r^{N} = \frac{a^{\frac{N}{N-2}}}{\alpha}.$$

and the pointwise bound for  $x \neq x_0$ 

$$\frac{\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}}}{|\log(\|u_{\epsilon}\|^{\frac{\beta}{\alpha}})|}v_{\epsilon}^{p}(x) \leq \frac{M\mu^{-p(N-2)\epsilon/2}}{\log(\mu)|x-x_{0}|^{p(N-2)}}$$

By Lemma 2.2,  $\frac{\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}}}{|\log(\|u_{\epsilon}\|^{\frac{\beta}{\alpha}})|} v_{\epsilon}^{p}(x) \to 0 \text{ for } x \neq x_{0}.$  Writing

$$-\Delta\left(\frac{\|u_{\epsilon}\|_{L^{\infty}}^{\frac{\beta}{\alpha}}}{|\log(\|u_{\epsilon}\|^{\frac{\beta}{\alpha}})|}u_{\epsilon}\right) = \frac{\|u_{\epsilon}\|_{L^{\infty}}^{\frac{\beta}{\alpha}}}{|\log(\|u_{\epsilon}\|^{\frac{\beta}{\alpha}})|}v_{\epsilon}^{p},$$

we observe that the last term converges to  $\delta_{x=x_0}$ . By Lemma 5.1, we have

$$\frac{\|u_{\epsilon}\|_{L^{\infty}}^{\frac{\beta}{\alpha}}}{|\log(\|u_{\epsilon}\|^{\frac{\beta}{\alpha}})|}u_{\epsilon} \to \frac{a^{\frac{N}{N-2}}}{\alpha}G \quad \text{in} \quad C^{1,\alpha}(w) \quad \text{as} \quad \epsilon \to 0,$$

and clearly we have (3.6) using (2.23). This completes the proof of the theorem.  $\Box$ 

Proof of Theorem 1.2. For p > N/(N-2) we have

$$\epsilon \|u_{\epsilon}\|^{\frac{N-2}{\alpha}} \int_{\Omega} u_{\epsilon}^{q_{\epsilon}+1} dx = \int_{\partial\Omega} (\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{\beta}{\alpha}} \nabla u_{\epsilon}, n) (\|u_{\epsilon}\|_{L^{\infty}(\Omega)} \nabla v_{\epsilon}, n) (n, x-y) ds$$

By (3.5) and (3.6),

$$\lim_{\epsilon \to 0} \epsilon \|u_{\epsilon}\|_{\Omega}^{\frac{N-2}{\alpha}} \int_{\Omega} u_{\epsilon}^{q_{\epsilon}+1} dx = \|V\|_{L^{p}(\mathbb{R}^{N})}^{p} \|U\|_{L^{q}(\mathbb{R}^{N})}^{q} \int_{\partial\Omega} \frac{\partial G(x, x_{0})}{\partial n} \frac{\partial G(x, x_{0})}{\partial n} (n, x - x_{0}) ds.$$

Also for the case p < N/(N-2), using

$$\epsilon \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{1}{\alpha}(p(N-2)-2)} \int_{\Omega} u_{\epsilon}^{q_{\epsilon}+1} dx$$
  
= 
$$\int_{\partial\Omega} (\|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{1}{\alpha}(\beta+p(N-2)-N)} \nabla u_{\epsilon}, n) (\|u_{\epsilon}\|_{L^{\infty}(\Omega)} \nabla v_{\epsilon}, n) (n, x-y) ds$$

and (3.8) and (3.7), we get

$$\begin{split} \lim_{\epsilon \to 0} \epsilon \|u_{\epsilon}\|_{L^{\infty}(\Omega)}^{\frac{1}{\alpha}(p(N-2)-2)} \int_{\Omega} u_{\epsilon}^{q_{\epsilon}+1} dx \\ &= \|U\|_{L^{q}(\mathbb{R}^{N})}^{q(p+1)} \int_{\partial\Omega} \frac{\partial \tilde{G}(x,x_{0})}{\partial n} \frac{\partial G(x,x_{0})}{\partial n} (n,x-x_{0}) ds. \end{split}$$

The case p = N/(N-2) is analogous.

The proof of the theorems follows from the next lemma.

Lemma 3.1. We have the following identities

$$i) \int_{\partial\Omega} \frac{\partial G(x, x_0)}{\partial n} \frac{\partial G(x, x_0)}{\partial n} (n, x - x_0) \, ds = -(N - 2)g(x_0, x_0)$$

and

$$ii) \int_{\partial\Omega} \frac{\partial \tilde{G}(x, x_0)}{\partial n} \frac{\partial G(x, x_0)}{\partial n} (n, x - x_0) \, ds = -\frac{N}{q+1} \tilde{g}(x_0, x_0)$$

*Proof.* i) was proven in [3], see also [13]. To prove ii) we follow a similar procedure. From [18, 22], for any  $y \in \mathbb{R}^N$ , we have the following identity

$$\int_{\Omega'} \Delta u(x-y,\nabla v) + \Delta v(x-y,\nabla u) - (N-2)(\nabla u,\nabla v)dx = \int_{\partial\Omega'} \frac{\partial u}{\partial n}(x-y,\nabla v) + \frac{\partial v}{\partial n}(x-y,\nabla u) - (\nabla u,\nabla v)(x-y,n)\,ds$$

where  $\Omega' = \Omega \setminus B_r$  with r > 0. For a system  $-\Delta v = 0$  and  $-\Delta u = v^p$ , in  $\Omega'$ , the identity takes the form

$$\int_{\Omega'} \frac{N}{p+1} v^{p+1} - \bar{a} v^{p+1} dx = \int_{\partial\Omega'} \frac{1}{p+1} v^{p+1} (x-y,n) ds$$
$$+ \int_{\partial\Omega'} \frac{\partial u}{\partial n} \left[ (x-y,\nabla v) + \bar{a} v \right] + \frac{\partial v}{\partial n} \left[ (x-y,\nabla u) + \bar{b} u \right] - (\nabla u,\nabla v) (x-y,n) ds \quad (3.9)$$

with  $\bar{a} + \bar{b} = N - 2$ . Let y = 0, choose  $\bar{a} = N/(p+1)$  and take v = G(x,0) and  $u = \tilde{G}(x,0)$ . Using that u = v = 0 on  $\partial\Omega$ , and so  $\nabla u = (\nabla u, n)n$  and  $\nabla v = (\nabla v, n)n$  on  $\partial\Omega$ , we obtain

$$\int_{\partial\Omega} \frac{\partial \tilde{G}}{\partial n} \frac{\partial \tilde{G}}{\partial n}(x,n) ds = \int_{\partial B_r} \frac{1}{p+1} G^{p+1}(x,n) + \frac{\partial \tilde{G}}{\partial n}[(x,\nabla G) + \frac{N}{p+1}G] ds + \int_{\partial B_r} \frac{\partial G}{\partial n}[(x,\nabla \tilde{G}) + \frac{N}{q+1}\tilde{G}] - (\nabla \tilde{G},\nabla G)(x,n) ds.$$

Let k = p(N-2) and  $\Gamma = \sigma_N(N-2)$ . For |x| = r, we have

$$\nabla \tilde{G} = -\frac{1}{\Gamma^p(N-k)} |x|^{-k} x + \nabla \tilde{g}, \quad \nabla G = -\frac{1}{\sigma_N} |x|^{-N} x + \nabla g,$$

$$\frac{\partial \tilde{G}}{\partial n} = -\frac{1}{\Gamma^p (N-k)} |x|^{1-k} + (\nabla \tilde{g}, n), \quad \frac{\partial G}{\partial n} = -\frac{1}{\sigma_N} |x|^{1-N} + (\nabla g, n)$$

$$\begin{aligned} (x,\nabla \tilde{G}) &+ \frac{N}{q+1}\tilde{G} &= (\frac{N}{(q+1)(k-2)} - 1)\frac{1}{\Gamma^p(N-k)}|x|^{2-k} + (x,\nabla \tilde{g}) + \frac{N}{q+1}\tilde{g} \\ (x,\nabla G) &+ \frac{N}{p+1}G &= (\frac{N}{p+1} - (N-2))\frac{1}{\Gamma}|x|^{2-N} + (x,\nabla g) + \frac{N}{p+1}g \end{aligned}$$

$$(\nabla \tilde{G}, \nabla G) = \frac{|x|^{-k-N+2}}{\sigma_N \Gamma^p(N-k)} - \frac{(\nabla g, x)}{\Gamma^p(N-k)} |x|^{(2-N)p} - \frac{(\nabla \tilde{g}, x)}{\sigma_N} |x|^{-N} + (\nabla \tilde{g}, \nabla g)$$

and

$$\frac{1}{p+1}G^{p+1} = \frac{1}{p+1} \left[ \frac{1}{\Gamma^p} |x|^{-k} - \Delta \tilde{g} \right] \left[ \frac{1}{\Gamma} |x|^{2-N} + g \right]$$

From here, we check that terms with  $|x|^{3-N-k}$  cancel out others integral tends to 0 since the integrand are  $o(|x|^{1-N})$  and only remain one term of order  $|x|^{1-N}$ , giving

$$\int_{\partial\Omega} \frac{\partial \tilde{G}}{\partial n} \frac{\partial G}{\partial n}(x,n) \, ds = -\lim_{r \to 0} \frac{1}{\sigma_N r^{N-1}} \int_{\partial B_r} \frac{N}{q+1} \tilde{g} ds = -\frac{N}{q+1} \tilde{g}(0,0).$$

Proof of Theorem 1.1. a) The part ii) follows from Theorem 5.1,

$$\||\Delta u_{\epsilon}|^{\frac{1}{p}}\|_{C^{1,\alpha}(\omega)} \le \|u_{\epsilon}^{q_{\epsilon}}\|_{L^{1}(\Omega)} + \|u_{\epsilon}^{q_{\epsilon}}\|_{L^{\infty}(\omega)}$$

and estimates (2.19), (2.22), and (2.23). Part i) follows from

$$||u_{\epsilon}||_{C^{1,\alpha}(\omega)} \le ||v_{\epsilon}^{p}||_{L^{1}(\Omega)} + ||v_{\epsilon}^{p}||_{L^{\infty}(\omega)}.$$

and estimate (2.18). Finally iii) follows combining ii) with the convergence

$$\int_{\mathbb{R}^N} |\Delta u_\epsilon|^{\frac{p+1}{p}} dx = \int_{\mathbb{R}^N} v_\epsilon^{p+1} dx \to \|V\|_{L^{p+1}(\mathbb{R}^N)}^{p+1}.$$

as  $\epsilon \to 0$ . This completes part a).

For part b), note that from (2.7), we have the vectorial equality  $\int_{\partial\Omega} (\nabla u_{\epsilon}, \nabla v_{\epsilon}) n \, ds =$ 

0. In the limit for  $p \ge N/(N-2)$ , we get

$$\int_{\partial\Omega} (\nabla G(x, x_0), \nabla G(x, x_0)) n \, ds = 0 \tag{3.10}$$

and similarly for p < N/(N-2), we obtain

$$\int_{\partial\Omega} (\nabla \tilde{G}(x, x_0), \nabla G(x, x_0)) n \, ds = 0 \tag{3.11}$$

But we have the following result.

**Lemma 3.2.** For every  $x_0 \in \Omega$ 

$$\int_{\partial\Omega} (\nabla G(x, x_0), n) (\nabla G(x, x_0), n) n \, ds = -\nabla \phi(x_0) \tag{3.12}$$

and

$$\int_{\partial\Omega} (\nabla \tilde{G}(x, x_0), n) (\nabla (\Delta \tilde{G}(x, x_0))^{1/p}, n) n \, ds = -\nabla \tilde{\phi}(x_0). \tag{3.13}$$

Hence combining (3.10) with (3.12), and (3.11) with (3.13), we complete the proof of part b) and the theorem is proven.

*Proof of the Lemma.* Equality (3.12) was proved in [3] and [13]. To prove (3.13), by (3.9) we have

$$\int_{\partial\Omega} \frac{\partial \tilde{G}}{\partial n} \frac{\partial G}{\partial n} n \, ds = \int_{\partial B_r} \left\{ \frac{1}{p+1} G^{p+1} n + \frac{\partial \tilde{G}}{\partial n} \nabla G + \frac{\partial G}{\partial n} \nabla \tilde{G} - (\nabla \tilde{G}, \nabla G) n \right\} ds.$$

Using  $\int_{\partial B_r} n = 0$ , we get  $\int_{\partial \Omega} \frac{\partial \tilde{G}}{\partial n} \frac{\partial G}{\partial n} n \, ds = \frac{1}{(p+1)r^{N-1}} \int_{\partial B_r} \left\{ \frac{1}{\Gamma^p} r^{N-k-1}g - \Delta \tilde{g} \frac{1}{\Gamma}r - \Delta \tilde{g}gr^{N-1} \right\} n \, ds$   $+ \frac{1}{r^{N-1}} \int_{\partial B_r} \left\{ (\nabla \tilde{g}, n) \nabla g + (\nabla g, n) \nabla \tilde{g} - (\nabla \tilde{g}, \nabla g)n \right\} r^{N-1} \, ds$   $- \frac{1}{r^{N-1}} \int_{\partial B_r} \left\{ \frac{1}{\sigma_N} \nabla \tilde{g} + \frac{r^{N-k}}{\Gamma^p(N-k)} \nabla g \right\} \, ds. \quad (3.14)$ 

We use the regular  $\hat{g}(x,0)$  instead of  $\tilde{g}(x,0)$ . Thus

$$\nabla \hat{g}(x,0) = \nabla \tilde{g}(x,0) + \frac{pg(0,0)}{\Gamma^{p-1}(2N-k-2)} |x|^{N-k-2} x, \qquad (3.15)$$

$$\Delta \hat{g}(x,0) = \Delta \tilde{g}(x,0) + \frac{pg(0,0)}{\Gamma^{p-1}} |x|^{N-k-2}.$$
(3.16)

But  $g(x,0) = g(0,0) + (\nabla g(0,0), x) + o(|x|^2)$  and

$$\int_{\partial B_r} r^{-k} g(x,0) n \, ds = \int_{\partial B_1} r^{N-k-1} g(0,0) n \, ds + \int_{\partial B_1} r^{N-k} (\nabla g(0,0), y) n \, ds + o(r^{N-k+1})$$

where y = x/r. Clearly the first integral in the r.h.s is zero and the other terms tends to zero as  $r \to 0$ . Hence

$$\lim_{r \to 0} \frac{1}{r^{N-1}} \int_{\partial B_r} r^{N-k-1} g(x,0) n \, ds = 0.$$
(3.17)

We replace (3.15) and (3.16) in (3.14), to obtain an identity without  $\tilde{g}$ . Using the limit (3.17) and that  $\hat{g}$  and g are regular, we obtain

$$\int_{\partial\Omega} \frac{\partial \tilde{G}}{\partial n} \frac{\partial G}{\partial n} n \, ds = \lim_{r \to 0} \frac{1}{r^{N-1}} \int_{\partial B_r} \frac{1}{\sigma_N} \nabla \hat{g} \, ds = \nabla \hat{g}(0,0) = \nabla \tilde{\phi}(0),$$

where the last equality follows by observation after Theorem 1.1.

## 4. Proof of Lemma 2.3

Let us recall the problem (2.10)-(2.12),

$$-\Delta u_{\epsilon,\mu} = v^p_{\epsilon,\mu} \quad \text{in} \quad \Omega_{\epsilon} \tag{4.1}$$

$$-\Delta v_{\epsilon,\mu} = u^{q_{\epsilon}}_{\epsilon,\mu} \quad \text{in} \quad \Omega_{\epsilon} \tag{4.2}$$

$$u_{\epsilon,\mu} = v_{\epsilon,\mu} = 0 \quad \text{on} \quad \partial\Omega_{\epsilon}$$

$$\tag{4.3}$$

where  $\Omega_{\epsilon} = (\Omega - x_{\epsilon})/\mu^{1-\epsilon/2}$ . Let  $\bar{R} > 0$ . We define  $\sigma(p) := 2 + N - p(N-2)$ , and the scalar function 1  $\cdot c$  ()

$$J(|y|) := \begin{cases} 1 & \text{if } \sigma(p) < 2, \\ |\log(|y|/\bar{R})| & \text{if } \sigma(p) = 2, \\ |y|^{2-\sigma(p)} & \text{if } \sigma(p) > 2. \end{cases}$$

Note that  $\sigma(p) \in [0, N)$  for  $p \in (2/(N-2), (N+2)/(N-2)]$  and  $\sigma(q) \leq 0$ . We consider the transformations

$$z_{\epsilon}(y) = |y|^{2-N} v_{\epsilon,\mu} \left(\frac{y}{|y|^2}\right) \quad \text{and} \quad w_{\epsilon}(y) = \frac{|y|^{2-N}}{J(|y|)} u_{\epsilon,\mu} \left(\frac{y}{|y|^2}\right)$$

in  $\Omega_{\epsilon}^*$ , the image of  $\Omega_{\epsilon}$  under  $x \mapsto x/|x|^2$ .

The next lemma is equivalent to Lemma 2.3, using the asymptotic behaviour (1.17).

## **Lemma 4.1.** Let $(w_{\epsilon}, z_{\epsilon})$ solving

$$-\Delta J(|y|)w_{\epsilon} = |y|^{-\sigma(p)}z_{\epsilon}^{p} \quad in \quad \Omega_{\epsilon}^{*}$$

$$(4.4)$$

$$-\Delta z_{\epsilon} = |y|^{-\sigma(q) + (q_{\epsilon} - q)(N-2)} [J(|y|)w_{\epsilon}]^{q_{\epsilon}} in \quad \Omega_{\epsilon}^{*}$$

$$(4.5)$$

$$w_{\epsilon} = z_{\epsilon} = 0 \quad on \quad \partial \Omega_{\epsilon}^*. \tag{4.6}$$

Then for any fixed  $R \in (0, \overline{R})$ , we have

$$\|w_{\epsilon}\|_{L^{\infty}(\Omega^{R}_{\epsilon})} + \|z_{\epsilon}\|_{L^{\infty}(\Omega^{R}_{\epsilon})} \le C$$

where  $\Omega_{\epsilon}^{R} = \Omega_{\epsilon}^{*} \cap B_{R}$ , and C = C(R) independent of  $\epsilon > 0$  provided  $\epsilon$  is sufficiently small.

*Proof.* Given R > 0, let  $w_0$  and  $z_0$  be solutions of

$$\Delta J(|y|)w_0 = 0$$
 in  $\Omega_{\epsilon}^R$  and  $w_0 = 0$ , on  $\partial \Omega_{\epsilon}^*$   $w_0 = w_{\epsilon}$  on  $\partial B_R$ 

and

$$\Delta z_0 = 0$$
 in  $\Omega_{\epsilon}^R$  and  $z_0 = 0$ , on  $\partial \Omega_{\epsilon}^*$   $z_0 = z_{\epsilon}$  on  $\partial B_R$ .

By the convergence in compact sets of  $w_{\epsilon}$  and  $z_{\epsilon}$ , see (2.13), we have  $|z_{\epsilon}| + |\nabla z_{\epsilon}| + |\nabla z_{\epsilon}|$  $|w_{\epsilon}| + |\nabla w_{\epsilon}| \leq C$  in |y| = R for C independent of  $\epsilon$ . Therefore by the maximum principle, we get

$$|Jw_0| + |\nabla(Jw_0)| + |z_0| + |\nabla z_0| \le C$$
 in  $\Omega^R_{\epsilon}$ .

Define  $\tilde{w} = w_{\epsilon} - w_0$  and  $\tilde{z} = z_{\epsilon} - z_0$ . We now write

$$-\Delta J(|y|)\tilde{w} = a(y)z_{\epsilon} \quad \text{in} \quad \Omega_{\epsilon}^{R}$$

$$(4.7)$$

$$-\Delta \tilde{z} = b(y)J(|y|)w_{\epsilon} \quad \text{in} \quad \Omega_{\epsilon}^{R}$$

$$(4.8)$$

$$\tilde{w} = \tilde{z} = 0 \quad \text{on} \quad \partial \Omega^R_{\epsilon}$$

$$\tag{4.9}$$

where  $a(y) = |y|^{-\sigma(p)} z_{\epsilon}^{p-1}$  and  $b(y) = |y|^{-\sigma(q) + (q_{\epsilon}-q)(N-2)} [J(|y|)w_{\epsilon}]^{q_{\epsilon}-1}$ . Clearly by the maximum principle  $\tilde{w} \geq 0$  and  $\tilde{z} \geq 0$ .

Let P(y) = a(y) and

$$Q(y) = \begin{cases} \frac{1}{M} b(y) & \text{for } y \in B_R \setminus \bar{B}_r \\ b(y) & \text{for } B_r \end{cases}$$

where  $r \in (0, R)$  and M > 1 both independent of  $\epsilon$  and to be determined later. Then

$$b(y)J(|y|)w_{\epsilon} = Q(y)J(|y|)w_{\epsilon} + f(y)$$

where

$$f(y) = (b(y) - Q(y))J(|y|)w_{\epsilon} = \begin{cases} 0 & \text{for } y \in \Omega_{\epsilon} \cap B_{r} \\ (1 - \frac{1}{M})b(y)J(|y|)w_{\epsilon} & \text{for } y \in B_{R} \setminus \bar{B}_{r} \end{cases}$$

It is clear that  $f \in L^{\infty}(\Omega_{\epsilon}^{R})$ , in fact  $||f||_{L^{\infty}(\Omega_{\epsilon}^{R})} \leq (1 - 1/M)r^{-(2+N)}$  by using that  $w_{\epsilon}(y) \leq Cr^{\sigma(p)-N}$  for  $|y| \geq r$ , when p < N/(N-2), and  $w_{\epsilon}(y) \leq Cr^{2-N}$  for  $|y| \geq r$  when p > N/(N-2). A similar bound is obtained for p = N/(N-2). Then we transform (4.7)–(4.8) in the system

$$\begin{aligned} -\Delta J \tilde{w} &= P z_{\epsilon} \quad \text{in} \quad \Omega_{\epsilon}^{R} \\ -\Delta \tilde{z} &= Q J w_{\epsilon} + f \quad \text{in} \quad \Omega_{\epsilon}^{R} \end{aligned}$$

We define  $\eta_2(y) = \chi_{w_{\epsilon} \leq 2\tilde{w}}(y)$  and  $\eta_1(y) = \chi_{z_{\epsilon} \leq 2\tilde{z}}(y)$  for  $y \in \Omega_{\epsilon}^R$ , we find

$$\begin{aligned} -\Delta J\tilde{w} &\leq 2\eta_1 P\tilde{z} + f_1 \quad \text{in} \quad \Omega_{\epsilon}^R \\ -\Delta \tilde{z} &\leq 2\eta_2 Q J\tilde{w} + f_2 \quad \text{in} \quad \Omega_{\epsilon}^R \end{aligned}$$

Where  $f_1 = (1 - \eta_1)Pz_{\epsilon} = \chi_{z_{\epsilon} \leq 2z_0}Pz_{\epsilon} \leq 2Pz_0$  and  $f_2 = f + (1 - \eta_2)QJw_{\epsilon}$  where  $(1 - \eta_2)QJw_{\epsilon} \leq 2QJw_0$ . We write the system in the form

$$-\Delta J\tilde{w} \leq 2\eta_1 P |y|^{\gamma} |y|^{-\gamma} \tilde{z} + f_1 \quad \text{in} \quad \Omega_{\epsilon}^R, \tag{4.10}$$

$$-|y|^{-\gamma}\Delta\tilde{z} \leq 2\eta_2 Q|y|^{-\gamma} J\tilde{w} + f_2|y|^{-\gamma} \quad \text{in} \quad \Omega^R_{\epsilon}, \tag{4.11}$$

$$\tilde{w} = \tilde{z} = 0 \quad \text{on} \quad \partial \Omega_{\epsilon}^{R}.$$
 (4.12)

Let  $u(y) \mapsto 2\eta_2 Q|y|^{-\gamma}u(y)$  and  $u(y) \mapsto 2\eta_1 P|y|^{\gamma}u(y)$  be the multiplication operators  $\mathcal{P}$  and  $\mathcal{Q}$  respectively. Note that a multiplication operator  $\mathcal{C}$  with corresponding function  $c(y) \in L^s(\Omega_{\epsilon}^R)$  is bounded from  $L^{s_1}(\Omega_{\epsilon}^R)$  to  $L^{s_2}(\Omega_{\epsilon}^R)$  with  $1/s_2 = 1/s_1 + 1/s$ .

Formally we define -L the operator as  $u(y) \mapsto -|y|^{-\gamma} \Delta(|y|^{\gamma} u(y))$ . More precisely, in the appendix, we define  $(-\Delta)^{-1}$  and  $(-L)^{-1}$ , which by the Hardy-Littlewood-Sobolev inequality are bounded, independently of  $\epsilon$ , from  $L^{m_1}(\Omega_{\epsilon}^R)$  to  $L^{m_2}(\Omega_{\epsilon}^R)$  with  $1/m_1 = 1/m_2 + 2/N$ . Note that the image of these operators is a function with zero-Dirichlet boundary condition, so they are positive. Then we can write

$$J\tilde{w} \le (-\Delta)^{-1} \mathcal{P}(-L)^{-1} (\mathcal{Q}(J\tilde{w}) + |y|^{-\gamma} f_2) + (-\Delta)^{-1} f_1$$

Denoting by  $K = (-\Delta)^{-1} \mathcal{P}(-L)^{-1} \mathcal{Q}$  and  $h = K|y|^{-\gamma} f_2 + (-\Delta)^{-1} f_1$  we have

$$(I - K)J\tilde{w} \le h$$

The proof is complete finding m large enough such that  $h \in L^m(\Omega_{\epsilon}^R)$  and (I - K) is invertible from  $L^m(\Omega_{\epsilon}^R)$  to  $L^m(\Omega_{\epsilon}^R)$ .

We can estimate  $Q(y)|y|^{-\gamma}$  in  $L^{\frac{q+1}{q-1}}(\Omega^R_{\epsilon})$ , for  $\gamma = 2\sigma(p)/(p+1) \ge 0$ , and note that  $\gamma = -\sigma(q)/(q+1)$  using the Sobolev Hyperbola. Since  $v_{\epsilon,\mu} \to V$  in  $L^{q+1}(\mathbb{R}^N)$ , we

have

$$\int_{\Omega_{\epsilon}^{*}} [J(|y|)w_{\epsilon}(y) - V(y/|y|^{2})|y|^{2-N}]^{q+1}|y|^{-\sigma(q)} \, dy \to 0 \quad \text{as} \quad \epsilon \to 0$$

Therefore for any  $\lambda$ , we can take r small such that

$$\int_{\Omega_{\epsilon}^{r}} [Jw_{\epsilon}]^{(q+1)\frac{q\epsilon-1}{q-1}}(y)|y|^{-\sigma(q)} dy \leq \int_{\Omega_{\epsilon}^{r}} [Jw_{\epsilon}]^{(q+1)}(y)|y|^{-\sigma(q)} dy \leq \frac{\lambda}{2C(\delta)}$$

and M large such that for all  $\epsilon \leq \epsilon_0$  we have

$$\int_{\Omega_{\epsilon}^{R}} [Q(s)|y|^{-\gamma}]^{\frac{q+1}{q-1}} dy \leq C(\delta) \int_{\Omega_{\epsilon}^{r}} [Jw_{\epsilon}]^{(q+1)\frac{q\epsilon-1}{q-1}} |y|^{-\sigma(q)} dy + \frac{C(\delta)}{M^{\frac{q+1}{q-1}}} \int_{B_{R}\setminus B_{r}} [Jw_{\epsilon}]^{(q+1)\frac{q\epsilon-1}{q-1}} |y|^{-\sigma(q)} dy \leq \lambda \quad (4.13)$$

where we have used  $b(y) \leq C(\delta)[Jw_{\epsilon}]^{q_{\epsilon}-1}$  with  $\delta$  given by Lemma 2.2. Now we show that K is bounded from  $L^{m}(\Omega^{R}_{\epsilon})$  to  $L^{m}(\Omega^{R}_{\epsilon})$ .

$$\begin{split} \|KJ\tilde{w}\|_{L^{m}(\Omega_{\epsilon}^{R})} &\leq C_{1}\|\mathcal{P}(-L)^{-1}\mathcal{Q}J\tilde{w}\|_{L^{r}(\Omega_{\epsilon}^{R})} \\ &\leq C_{1}\||y|^{\gamma}2\eta_{1}P\|_{L^{\frac{p+1}{p-1}}(\Omega_{\epsilon}^{R})}\|(-L)^{-1}\mathcal{Q}J\tilde{w}\|_{L^{r'}(\Omega_{\epsilon}^{R})} \\ &\leq C_{1}\||y|^{\gamma}2\eta_{1}P\|_{L^{\frac{p+1}{p-1}}(\Omega_{\epsilon}^{R})}C_{2}\|\mathcal{Q}J\tilde{w}\|_{L^{s'}(\Omega_{\epsilon}^{R})} \\ &\leq C_{1}C_{2}\||y|^{\gamma}2\eta_{1}P\|_{L^{\frac{p+1}{p-1}}(\Omega_{\epsilon}^{R})}\||y|^{-\gamma}2\eta_{2}Q\|_{L^{\frac{q+1}{q-1}}(\Omega_{\epsilon}^{R})}\|J\tilde{w}\|_{L^{m'}(\Omega_{\epsilon}^{R})} \\ &\leq \overline{C}\||y|^{\gamma}P\|_{L^{\frac{p+1}{p-1}}(\Omega_{\epsilon}^{R})}\||y|^{-\gamma}Q\|_{L^{\frac{q+1}{q-1}}(\Omega_{\epsilon}^{R})}\|J\tilde{w}\|_{L^{m'}(\Omega_{\epsilon}^{R})} \end{split}$$

with  $\frac{1}{r} = \frac{1}{m} + \frac{2}{N}$ , so r' > 1 implies m > N/(N-2).  $\frac{1}{r} = \frac{p-1}{p+1} + \frac{1}{r'}$  and  $\frac{1}{s'} = \frac{1}{r'} + \frac{2}{N}$ , so condition b) in (5.1) implies N - 2 + N/m > 2N/(p+1) and s' > 1 implies m > (q+1)/2 so a) in (5.1) holds since  $\gamma > 0$  and  $\frac{1}{s'} = \frac{q-1}{q+1} + \frac{1}{m'}$ . Since

$$\frac{q-1}{q+1} + \frac{p-1}{p+1} = \frac{4}{N}$$
, we have  $m' = m$ 

By

$$\int_{a_{\epsilon}^{*}} [z_{\epsilon}(y) - U(y/|y|^{2})|y|^{2-N}]^{p+1}|y|^{-\sigma(p)} \, dy \to 0 \quad \text{as} \quad \epsilon \to 0,$$

we deduce that  $|||y|^{\gamma-\sigma(p)}z_{\epsilon}^{p-1}||_{L^{\frac{p+1}{p-1}}(\Omega_{\epsilon}^{R})} = |||y|^{\gamma}P||_{L^{\frac{p+1}{p-1}}(\Omega_{\epsilon}^{R})} \leq C(\epsilon_{0})$  with  $C(\epsilon_{0}) > 0$ and for all  $\epsilon \in (0, \epsilon_{0})$ . Since  $\lambda$  in (4.13) can be arbitrarely small then the norm of K is small and so  $I - K \colon L^{m}(\Omega_{\epsilon}^{R}) \to L^{m}(\Omega_{\epsilon}^{R})$  invertible for m large. We have that

$$|||y|^{-\gamma} f_2||_{L^m(\Omega_{\epsilon}^R)} \le r^{-\gamma} ||f_2||_{L^{\infty}(\Omega_r^R)} (\operatorname{meas}(\Omega_r^R))^{1/m}$$

is bounded, since  $f_2$  is zero outside  $\Omega_r^R$  and

$$\|\Delta^{-1}f_1\|_{L^m(\Omega^R_{\epsilon})} \le C_1 \|f_2\|_{L^r(\Omega^R_{\epsilon})} \le \|f_1\|_{L^{\infty}(\Omega^R_{r})} (\operatorname{meas}(\Omega^R_{\mu}))^{1/r} \le C(z_0) (\operatorname{meas}(B_R))^{1/r}$$

This implies  $||Jw||_{L^m(\Omega^R)} \leq M$  for every m large, and consequently for every  $m \geq 1$ . (Use the  $w_0$  to get that  $||Jw_{\epsilon}||_{L^m(\Omega^R)} \leq M$ ). Now we have that

$$-\Delta \tilde{z} = b(y)Jw_{\epsilon} = |y|^{-\sigma(q) + (q-q_{\epsilon})(N-2)}[Jw_{\epsilon}]^{q_{\epsilon}}.$$

Since  $\sigma(q) \leq 0$ , if we take *m* large such that  $mq_{\epsilon} > N/2$  then

$$\|\tilde{z}\|_{L^{\infty}(\Omega_{\epsilon}^{R})} \leq \tilde{M}$$
 and therefore  $\|z_{\epsilon}\|_{L^{\infty}(\Omega_{\epsilon}^{R})} \leq M$  (4.14)

for some M independent of  $\epsilon$ . We study now each case of J separately. We have

$$-\Delta J w_{\epsilon} = |y|^{-\sigma(p)} z_{\epsilon}^{p} \quad \text{in} \quad \Omega_{\epsilon}^{*}.$$
(4.15)

a) In the case J = 1, since  $\sigma(p) < 2$ , using (4.14), we have  $-\Delta \tilde{w}_{\epsilon} \in L^q(\Omega)$  for any  $q \in (N/2, N/\sigma(p))$ . By regularity, we get

$$\|w_{\epsilon}\|_{L^{\infty}(\Omega^{R}_{\epsilon})} \leq M.$$

b) For  $J(|y|) = -\log(|y|/\bar{R}) > \log(\bar{R}/R)$ , we have

$$-\Delta \tilde{w} - \frac{\nabla J}{J} \nabla \tilde{w} - \frac{\Delta J}{J} \tilde{w} = \frac{1}{J|y|^2} z_{\epsilon}^p \quad \text{in} \quad \Omega_{\epsilon}^R$$

or equivalently

$$-\Delta \tilde{w} + \frac{1}{J|y|^2} (y, \nabla \tilde{w}) + \frac{1}{J|y|^2} (N-2)\tilde{w} = \frac{1}{J|y|^2} z_{\epsilon}^p \quad \text{in} \quad \Omega_{\epsilon}^R.$$

Using (4.14), we can take  $u = \tilde{w} - M$  with  $M = \sup_{\epsilon > 0} \sup_{y \in \Omega_{\epsilon}^{R}} z_{\epsilon}^{p}(y)/(N-2)$ , and we get

$$-J|y|^2\Delta u + (y,\nabla u) + (N-2)u \le 0 \quad \text{in} \quad \Omega^R_\epsilon.$$

Since u = -M < 0 on the boundary,  $u \leq 0$  in  $\Omega_{\epsilon}^{R}$ . This gives  $w_{\epsilon} \leq M$  in  $\Omega_{\epsilon}^{R}$ . For the remaining case, p < N/(N-2) we have

$$-\Delta \tilde{w} - \frac{\nabla J}{J} \nabla \tilde{w} - \frac{\Delta J}{J} \tilde{w} = \frac{1}{|y|^2} z_{\epsilon}^p \quad \text{in} \quad \Omega_{\epsilon}^R.$$

As before, defining  $u = \tilde{w} - M$  with  $M = \sup_{\epsilon > 0} \sup_{y \in \Omega_{\epsilon}^{R}} z_{\epsilon}^{p} / [(\sigma(p) - 2)(N - \sigma(p))]$  then

$$-|y|^2 \Delta u - (2 - \sigma(p))(y, \nabla u) - (2 - \sigma(p))(N - \sigma(p))u \le 0 \quad \text{in} \quad \Omega_{\epsilon}^R$$

Since u = -M < 0 on the boundary,  $u \leq 0$  in  $\Omega_{\epsilon}^{R}$ . This implies  $w_{\epsilon} \leq M$  in  $\Omega_{\epsilon}^{R}$ .

### 5. Appendix

Let N > 2. Let h and v be a function in  $L^{s'}(\Omega^R_{\epsilon})$ . Given the Green's function G solution of  $-\Delta G(x, \cdot) = \delta_x$  in  $\Omega^R_{\epsilon}$ ,  $G(x, \cdot) = 0$  on  $\partial \Omega^R_{\epsilon}$ , we define

$$(-\Delta)^{-1}h(\xi) = \int_{\Omega_{\epsilon}^{R}} G(x,\xi)h(x) \, dx \quad \xi \in \Omega_{\epsilon}^{R}.$$

and

$$(-L)^{-1}v(\xi) = |\xi|^{-\gamma} \int_{\Omega_{\epsilon}^{R}} G(x,\xi) |x|^{\gamma} v(x) \, dx \quad \xi \in \Omega_{\epsilon}^{R}.$$

Note that G is positive, so both operators are positive. We know that  $(-\Delta)^{-1}$  is bounded, independently of  $\epsilon$ , from  $L^{s'}(\Omega_{\epsilon}^R)$  to  $L^{r'}(\Omega_{\epsilon}^R)$  with 1/r' = 1/s' - 2/N. Next we prove the same result for  $(-L)^{-1}$ . By the weighted Hardy-Littlewood-Sobolev inequality [5, 17], for  $|\xi|^{-\gamma} f \in L^{s'}(\Omega_{\epsilon}^R)$ , we have that

$$\|\xi^{-\gamma}(-\Delta)^{-1}f\|_{L^{r'}(\Omega_{\epsilon}^{R})} \le 2\||\xi|^{-\gamma} \int_{\Omega_{\epsilon}^{R}} \frac{C}{|x-\xi|^{N-2}} f(x) \, dx\|_{L^{r'}(\Omega_{\epsilon}^{R})} \le C\||\xi|^{-\gamma} f\|_{L^{s'}(\Omega_{\epsilon}^{R})}$$

for  $1 < s' < r' < \infty$ , with 1/r' = 1/s' - 2/N and

a) 
$$-\gamma < N(1 - 1/s') = N - 2 - N/r'$$
 and b)  $\gamma < N/r'$ . (5.1)

In other words, for any  $v \in L^{s'}(\Omega^R_{\epsilon})$ , we have

$$\begin{aligned} \|(-L)^{-1}v\|_{L^{r'}(\Omega_{\epsilon}^{R})} &= \||\xi|^{-\gamma}(-\Delta)^{-1}|x|^{\gamma}v\|_{L^{r'}(\Omega_{\epsilon}^{R})} \\ &\leq 2\||\xi|^{-\gamma} \int_{\Omega_{\epsilon}^{R}} \frac{C}{|x-\xi|^{N-2}} |x|^{\gamma}v(x) \, dx\|_{L^{r'}(\Omega_{\epsilon}^{R})} \\ &\leq C\|v\|_{L^{s'}(\Omega_{\epsilon}^{R})}. \end{aligned}$$
(5.2)

Lemma 5.1. Let u solve

$$\begin{cases} -\Delta u = f & in \quad \Omega \subset \mathbb{R}^N \\ u = 0 & on \quad \partial \Omega \end{cases}$$

Let  $\omega$  be a neighborhood of  $\partial\Omega$ . Then

$$\|u\|_{W^{1,q}(\Omega)} + \|\nabla u\|_{C^{0,\alpha}(\omega')} \leq C(\|f\|_{L^1(\Omega)} + \|f\|_{L^{\infty}(\omega)})$$
  
for  $q < N/(N-1)$ ,  $\alpha \in (0,1)$  and  $\omega' \subset \omega$  is a strict subdomain of  $\omega$ .

#### Acknowledgements

The author thank Prof. J. Dávila for many useful discussions.

#### References

- F.V. ATKINSON AND L.A. PELETIER, *Elliptic equations with nearly critical growth*, J. Diff. Eq. 70 (1987), 349–365.
- [2] M. BEN AYED AND K. EL MEHDI, On a biharmonic equation involving nearly critical exponent, preprint
- [3] H. BREZIS AND L.A. PELETIER Asymptotics for elliptic equations involving critical growth, Partial Differential Equations and the Calculus of Variations, Vol. I, Birkhuser Boston, Massachusetts, (1989), 149-192.
- [4] PH. CLÉMENT, D.G. DE FIGUEIREDO, AND E. MITIDIERI, Positive solutions of semilinear elliptic systems, Comm. PDE 17 (1992) 923–940.
- [5] X. CHEN, C. LI AND B. OU Classification of solutions for a system of integral equations Comm. in PDE, 30 2005, 59–65.

- [6] K.-S. CHOU AND D. GENG, Asymptotics of positive solutions for a biharmonic equation involving critical exponent Diff. and Int. Eq. 13(7-9) (2000) 921-940.
- [7] D.G. DE FIGUEIREDO AND P. FELMER On superquadratic elliptic systems, Trans. Amer. Math. Soc. 343 (1994), 99-116.
- [8] D.G. DE FIGUEIREDO, P.L. LIONS AND R.D. NUSSBAUM, A priori estimates and existence of positive solutions of semilinear elliptic equations J. Math. Pures. Appl. 61 1982, 41-63.
- [9] D.G. DE FIGUEIREDO AND B. RUF *Elliptic Systems with Nonlinearities of Arbitrary Growth*, Trans. Amer. Math. Soc. 343 (1994), 99-116.
- [10] M. FLUCHER AND J. WEI, Semilinear Dirichlet problem with nearly critical exponent, asymptotic location of hot spots Manuscripta Matematica 94 1997, 337–346.
- [11] B. GIDAS, W. NI, AND L. NIRENBERG, Symmetry and related properties via the maximum principle Comm. Math. Phys. 68 1979, 209-243.
- [12] D. GILBARG AND N.S. TRUDINGER, Elliptic Partial Differential Equations of Second Order, second edition. Springer-Verlag, Berlin-New York, 1983.
- [13] Z-C. HAN, Asymptotic approach to singular solutions for nonlinear elliptic equations involving critical Sobolev exponent Ann. Inst. Henri Poincaré, 8(2) 1991, 159-174.
- [14] M. HAMMAMI, Concentration phenomena for fourth-order elliptic equations with critical exponent Electron. J. Differential Equations 121 2004, 22p.
- [15] J. HULSHOF AND R.C.A.M VAN DER VORST, Differential systems with strongly indefinite variational structure, J. Funct. Anal. 114 (1993), 32–58.
- [16] J. HULSHOF AND R.C.A.M VAN DER VORST, Asymptotic Behaviour of ground states Proc. Amer. Math. Soc. 124(8) 1996, 2423–2431.
- [17] E. H. LIEB, Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities, Annals of Math. 118 1983, 349–374
- [18] E. MITIDIERI, A Rellich type identity and applications, Comm. Partial Diff. Equations 18(1-2) 1993, 125-151.
- [19] L.A. PELETIER AND R.C.A.M VAN DER VORST, Existence and nonexistence of positive solutions of non-linear elliptic systems and the biharmonic equation Diff. Int. Eq. 5 no. 4 1992, 747-767.
- [20] O. REY Proof of two conjectures of H. Brézis and L.A. Peletier. Manuscripta Matematica 65 1 1989, 19–37.
- [21] M. STRUWE, Global compactness result for elliptic boundary value problems involving limiting nonlinearities, Math. Z. vol 187 1984, 511–517
- [22] R.C.A.M VAN DER VORST, Variational identities and applications to differential systems, Arch. Rational. Mech. Anal. 116, (1991), 375-398.
- [23] X.J. WANG, Sharp constant in a Sobolev inequality, Nonlinear Anal. T.M.A. 20 (1993), 261-268

Departamento de Matematica y C. C., Universidad de Santiago, Casilla 307, Correo 2, Santiago, Chile

*E-mail address*: iguerra@usach.cl