# New constructions for covering designs

Daniel M. Gordon, 1 Greg Kuperberg, 2 and Oren Patashnik 1

<sup>1</sup>Center for Communications Research, 4320 Westerra Ct., San Diego, CA 92121 <sup>2</sup>Department of Mathematics, University of California, One Shields Avenue, Davis, CA 95616 (Dated: 16 February 1995)

A (v, k, t) covering design, or covering, is a family of k-subsets, called blocks, chosen from a v-set, such that each t-subset is contained in at least one of the blocks. The number of blocks is the covering's size, and the minimum size of such a covering is denoted by C(v, k, t). This paper gives three new methods for constructing good coverings: a greedy algorithm similar to Conway and Sloane's algorithm for lexicographic codes [6], and two methods that synthesize new coverings from preexisting ones. Using these new methods, together with results in the literature, we build tables of upper bounds on C(v, k, t) for  $v \le 32$ ,  $k \le 16$ , and  $t \le 8$ .

# 1. INTRODUCTION

Let the covering number C(v,k,t) denote the smallest number of k-subsets of a v-set that cover all t-subsets. These numbers have been studied extensively. Mills and Mullin [19] give known results and many references. Hundreds of papers have been written for particular values of v, k, and t. The best general lower bound on C(v,k,t), due to Schönheim [27], comes from the following inequality:

# Theorem 1

$$C(v,k,t) \geq \left\lceil \frac{v}{k}C(v-1,k-1,t-1) \right\rceil.$$

Iterating this gives the Schönheim bound  $C(v,k,t) \ge L(v,k,t)$ , where

$$L(v,k,t) = \left\lceil \frac{v}{k} \left\lceil \frac{v-1}{k-1} \dots \left\lceil \frac{v-t+1}{k-t+1} \right\rceil \dots \right\rceil \right\rceil.$$

Sometimes a lower bound of de Caen [7] is slightly better than the Schönheim bound when k and t are not too small:

$$C(v,k,t) \ \geq \ \frac{(t+1)(v-t)}{(k+1)(v-k)} \binom{v}{t} \bigg/ \binom{k}{t} \, .$$

The best general upper bound on C(v,k,t) is due to Rödl [26]: Define the *density* of a covering to be the average number of blocks containing a t-set. The minimum density of a (v,k,t) covering is  $C(v,k,t)\binom{k}{t}/\binom{v}{t}$  and is obviously at least 1. Rödl shows that for k and t fixed there exist coverings with density approaching 1 as v gets large. Erdős and Spencer [11] give the bound

$$C(v,k,t) \binom{k}{t} / \binom{v}{t} \le 1 + \ln \binom{k}{t},$$

which is weaker but applies to all v, k, and t. Furthermore it can be improved by at most a factor of  $4 \ln 2 \approx 2.77$  asymptotically, because a  $(v,v-1,\lfloor v/2 \rfloor)$  covering that achieves the Schönheim lower bound has density asymptotic to v/4, while the Erdős-Spencer upper bound in that case corresponds to a density asymptotic to  $v \ln 2$ .

This paper presents new constructions for coverings. The greedy method of Section 2 produces reasonably good coverings and it is completely general—it applies to all possible

values of v, k, and t, and it doesn't rely on the existence of other good coverings. The finite geometries of Section 3 produce very good (often optimal) coverings, but they apply only to certain sets of v, k, and t values. The induced-covering method of Section 4, which constructs coverings from larger ones, and the dynamic programming method of Section 5, which constructs coverings from smaller ones, both apply to all parameter values, but they rely on preexisting coverings. (We show in a paper with Spencer [12] that the greedy construction, as well as the induced-covering method applied to certain finite geometry coverings, both produce coverings that match Rödl's bound.) Finally, the previously known methods of Section 6, when combined with the methods of earlier sections, yield the tables of upper bounds in Section 7.

# 2. GREEDY COVERINGS

Our greedy algorithm for generating coverings is analogous to the surprisingly good greedy algorithm of Conway and Sloane [6] for generating codes. That algorithm may be stated very concisely: To construct a code of length n and minimum distance d, arrange the binary n-tuples in lexicographic order, and repeatedly choose the first one in the list that is distance d or more from all n-tuples chosen earlier; the n-tuples chosen are the codewords. The resulting code is called a lexicographic code, or lexicode.

This simple method has several nice features: Lexicodes tend to be fairly good (at packing codewords into the space), they are linear, and they include some well-known codes such as Hamming codes and the binary Golay codes. Brouwer, Shearer, Sloane, and Smith [3, page 1349] use the same method to make constant weight codes, by choosing only *n*-tuples of a given weight.

The greedy algorithm does not require lexicographic order. Brualdi and Pless [4] show that a large family of orders lead to linear codes. And sometimes Gray code orders, for example, lead to better codes.

Constructing good codes and good constant weight codes are packing problems. But a similar method applies to covering problems. A greedy (v,k,t) covering is one generated by the following algorithm:

1. Arrange the *k*-subsets of a *v*-set in a list.

- Choose from the list the k-subset that contains the maximum number of t-sets that are still uncovered. In case of ties, choose the k-subset occurring earliest in the list.
- 3. Repeat Step 2 until all *t*-sets are covered.

The list of k-sets can be in any order. Some natural orders are lexicographic, colex (which is similar to lexicographic but the subsets are read from right to left rather than left to right), and a generalized Gray code order (where successive sets differ only by one deletion and one addition). The resulting lists, when k = 3 and v = 5, are

```
123 124 125 134 135 145 234 235 245 345 (lexicographic);
123 124 134 234 125 135 235 145 245 345 (colex);
123 134 234 124 145 245 345 135 235 125 (gray).
```

Nijenhuis and Wilf [22] give algorithms to generate lexicographic and Gray code orders. Stanton and White [30] discuss colex algorithms.

It is natural to investigate the greedy algorithm with random order, too, since we know [12] that random order does well asymptotically. To keep with the constructive spirit of this paper, we used an easily reproduced "random" permutation of the k-sets. To generate the permutation, start with the k-sets lexicographically ordered in positions 1 through  $\binom{v}{k}$ , then successively swap the k-sets in positions i and i+j, for  $i=1,2,\ldots,\binom{v}{k}$ , where j is  $X_i$  mod  $\binom{v}{k}-i+1$  and where the sequence of pseudo-random X's comes from the linear congruential generator  $X_{i+1}=(41X_i+7) \mod 2^{30}$ . The seed  $X_0$  is 1, and when there are multiple random-order runs on the same set of (v,k,t) parameters, the subsequent seeds are 2, 3, .... Knuth [15] discusses the linear congruential method.

Greedy coverings are not in general optimal, but as happens with codes (Brouwer, Shearer, Sloane, and Smith [3], Brualdi and Pless [4], Conway and Sloane [6]) they are often quite good—about 42% of the table entries come from greedy coverings. Interestingly, the Steiner system S(24,8,5), which Conway and Sloane [6, page 347] showed is a constant-weight lexicographic code, also arises as a greedy covering.

The problem with greedy coverings is that they are expensive to compute. Our implementation of the algorithm above uses two arrays: one with  $\binom{\nu}{k}$  locations corresponding to the k-subsets, and one with  $\binom{\nu}{t}$  locations corresponding to the t-subsets. Each k-set array location contains the number of uncovered t-sets contained in that k-set, and is initialized to  $\binom{k}{t}$ . Each t-set array location contains a 0 or 1, indicating whether that t-set has been covered. Each time through Step 2, each t-set contained in the selected k-set must be checked. If the t-set is uncovered, it is marked as covered, and each k-set containing it must have its array location decremented. For fixed k and t, the algorithm asymptotically takes time and space  $O(\nu^k)$ .

We ran a program to generate greedy coverings for all entries in our tables, for all four orders described above. For random order, we used  $10^e$  runs, where  $e = 3[v \le 20] + [v \le 15] + [v \le 10] + [k \le 10] + [k \le 5] + 2[U]$  and where U is the predicate 't = 2 and C(v, k, 2) is unknown' (the symbol [P] is 1 if the predicate P is true, 0 otherwise).

For the range of parameters of our tables, the four orders produced coverings of roughly the same size, but lexicographic order performed slightly better on average than colex order, which performed better than Gray code order, which performed better than a single run of random order.

#### 3. FINITE GEOMETRY COVERINGS

Finite geometries may be used to construct very good coverings for certain sets of parameters. Anderson [2] has a nice discussion of finite geometries.

Let  $\operatorname{PG}(m,q)$  denote the projective geometry of dimension m over  $\operatorname{GF}(q)$ , where q is a prime power. The points of  $\operatorname{PG}(m,q)$  are the equivalence classes of nonzero vectors  $u=(u_0,u_1,\ldots,u_m)$ , where two vectors u and v are equivalent if  $u=\lambda v$  for some nonzero  $\lambda\in\operatorname{GF}(q)$ . There are  $(q^{m+1}-1)/(q-1)$  such points.

A k-flat is a k-dimensional subspace of PG(m,q), for  $1 \le k \le m$ , determined by m-k independent homogeneous linear equations. A k-flat has  $(q^{k+1}-1)/(q-1)$  points, and there are  $\begin{bmatrix} m+1 \\ k+1 \end{bmatrix}_q$  different k-flats in PG(m,q), where

$$\begin{bmatrix} n \\ k \end{bmatrix}_{q} = \frac{(q^{n} - 1)(q^{n-1} - 1)\dots(q^{n-k+1} - 1)}{(q^{k} - 1)(q^{k-1} - 1)\dots(q - 1)}$$

is the *q*-binomial coefficient.

By removing all points with  $u_0 = 0$  we obtain the affine (or Euclidean) geometry AG(m,q). It has  $q^m$  points and  $q^{m-k} {m \brack k}_q$  different k-flats, each of which contains  $q^k$  points.

For either geometry, any k+1 independent points determine a k-flat, and k+1 dependent points are contained in multiple k-flats, so the k-flats cover every set of k+1 points. Thus, taking the points of the geometry as the v-set of the covering, and taking the points of a k-flat as a block of the covering, we get the following two theorems.

# Theorem 2

$$C\left(\frac{q^{m+1}-1}{q-1}, \frac{q^{k+1}-1}{q-1}, k+1\right) \le {m+1 \brack k+1}_q.$$

# Theorem 3

$$C(q^m, q^k, k+1) \leq q^{m-k} \begin{bmatrix} m \\ k \end{bmatrix}_q$$

Equality holds for both theorems when k = m - 1 or k = 1. Theorem 2 is due to Ray-Chaudhuri [25], and Theorem 3 follows easily from results of Abraham, Ghosh, and Ray-Chaudhuri [1], although the idea of using finite geometries to construct coverings dates back at least to Veblen and Bussey [38] in 1906.

### 4. INDUCED COVERINGS

The main drawback of the finite geometry coverings is that they exist only for certain families of parameters. But they are such good coverings that they can be used to construct pretty good coverings for other parameters.

Suppose we have a good (v,k,t) covering, say from a geometry, and we want to construct a (v',k',t) covering, where v' < v and k' < k. Consider the family of sets obtained from the (k-element) blocks by randomly choosing v' elements of the v-set, deleting all other elements from the blocks, and throwing out any blocks with fewer than t elements (since those blocks cover no t-sets).

The remaining blocks cover all t-subsets of the v' elements, but have different sizes. Suppose some block has  $\ell$  elements. If  $\ell = k'$  its size is correct as is, and it becomes a block of our new covering. If  $\ell < k'$ , add any  $k' - \ell$  elements to the block. And if  $\ell > k'$ , replace the block by an  $(\ell, k', t)$  covering, which covers all t-sets the original block covered.

The new blocks each have k' elements, and together they cover all t-sets, so the new family forms a (v',k',t) induced covering.

In small cases, the method tends to do best when k'/k is about v'/v. In large cases, the method does well if for every  $\ell$  near v'k/v, a good  $(\ell,k',t)$  covering is available. Also, it need not start with a finite geometry covering—any (v,k,t) covering will do. But generally the better the covering it starts with, the better the result.

The induced coverings in our tables come either from using the simple special cases of Section 6.1 or from finite geometries. We constructed each finite geometry covering based on PG(m,p) and AG(m,p) with  $p \le 11$  prime and with at most  $10^4$  points and  $10^6$  flats. For each such covering, and for each v and k in the relevant table, we used a random set of v points to construct an induced covering as described above, trying 100 random sets in each case.

# 5. COMBINING SMALLER COVERINGS

Suppose we want to form a  $(v_1+v_2,k,t)$  covering. Let the  $(v_1+v_2)$ -set be the disjoint union of a  $v_1$ -set and a  $v_2$ -set. Given an s with  $0 \le s \le t$ , choose a  $(v_1,\ell,s)$  covering and a  $(v_2,k-\ell,t-s)$  covering for some  $\ell$ , which must be in the range  $s \le \ell \le k-t+s$ . For each possible arrangement of t elements as an s-subset of the  $v_1$ -set and a (t-s)-subset of the  $v_2$ -set, there is an  $\ell$ -set from the first covering and a  $(k-\ell)$ -set from the second covering whose union is a k-set that covers the t-set. Thus the number of blocks that cover all such t-sets is at most the product of the sizes of the two coverings. Choosing an optimal  $\ell$  for each s gives us our  $(v_1+v_2,k,t)$  covering built up from smaller coverings. This construction gives the bound

$$C(v_1 + v_2, k, t) \ \leq \ \sum_{s=0}^t \min_{\ell} C(v_1, \ell, s) \cdot C(v_2, k - \ell, t - s) \, .$$

Furthermore we can try all choices of  $v_1$  and  $v_2$  summing to the v of interest.

The coverings produced by this method tend to have some redundancy. To remove redundancy when  $v_1 = 2$ , for example, we can try combining a (v,k,t) covering and a (2,0,0)

covering (which has one block, the empty set), along with a (v,k-2,t-1) covering and a (2,2,2) covering. This forms a (v+2,k,t) covering, and is sometimes an improvement over the basic construction above:

$$C(v+2,k,t) \leq C(v,k,t) + C(v,k-2,t-1).$$

This example has replaced the s and s+1 terms of the basic construction's bound, when s=1, with the single term

$$\min_{\ell} C(v_1, \ell, s+1) \cdot C(v_2, k-\ell, t-s).$$

The new term corresponds to covering any t-subset having either s or s+1 elements in the  $v_1$ -set, by using one product of coverings, rather than two. If changing  $C(v_1, \ell, s)$  to  $C(v_1, \ell, s+1)$  does not cost too much, the bound will improve.

To generalize this combining of terms, define  $c_{i,j}$  for  $0 \le i \le j \le t$  to be the number of blocks required to cover any t-subset that has between i and j elements in the  $v_1$ -set, and between t-j and t-i elements in the  $v_2$ -set. Since  $c_{i,j} \le c_{i,r} + c_{r+1,j}$  for any  $i \le r < j$ , we have

$$\begin{split} c_{i,j} \; \leq \; \min \bigl( \min_{\ell} C(\mathbf{v}_1,\ell,j) \cdot C(\mathbf{v}_2,k-\ell,t-i), \\ & \quad \min_{i \leq r < j} \bigl( c_{i,r} + c_{r+1,j} \bigr) \bigr) \,. \end{split}$$

Using dynamic programming, we may efficiently compute a bound for  $c_{0,t}$ , which is an upper bound for  $C(v_1+v_2,k,t)$ .

This general construction produces about 30% of the entries in our tables. It includes as special cases several of the simple constructions of Section 6.1, as well as the direct-product construction of Morley and van Rees [21], which yields the bound

$$C(2v+v, v+k+v, t+s+1) < C(v,k,t) + C(v+v, k+v, s)$$
.

#### 6. OTHER CONSTRUCTIONS

### **6.1.** Simple Constructions

There are several simple and well-known methods for building coverings from other coverings. All but the last of these methods are special cases of the methods in the previous two sections.

Adding a random element to each block of a (v,k,t) covering gives a (v,k+1,t) covering of the same size. Thus

$$C(v,k+1,t) \leq C(v,k,t)$$
.

Adding a new element to a v-set, and including it in every block in a (v,k,t) covering, forms a (v+1,k+1,t) covering of the same size, hence

$$C(v+1,k+1,t) \leq C(v,k,t)$$
.

Combining a (v,k,t) covering and a (v,k-1,t-1) covering over the same v-set, by adding a new v+1st element to all of the blocks of the (v,k-1,t-1) covering but to none of the

blocks of the (v,k,t) covering, forms a (v+1,k,t) covering, of size the sum of the other two sizes, thus

$$C(v+1,k,t) < C(v,k,t) + C(v,k-1,t-1).$$

Those constructions are special cases of the method of Section 5.

Deleting one element from a v-set, and adding a random element to any block of a (v,k,t) covering that contains the deleted element, creates a (v-1,k,t) covering of the same size. Thus

$$C(v-1,k,t) \leq C(v,k,t)$$
.

Choosing the element of a covering that occurs in the fewest blocks, throwing away all other blocks, and then throwing away the chosen element, results in a (v-1,k-1,t-1) covering. This method, due to Schönheim, is a reformulation of Theorem 1; the corresponding upper bound is

$$C(v-1,k-1,t-1) \leq \left\lfloor \frac{k}{v}C(v,k,t) \right\rfloor.$$

Those two constructions are special cases of the induced-covering method of Section 4.

Replacing each element of the *v*-set in a (v,k,t) covering by m different elements gives an (mv,mk,t) covering of the same size, thus

$$C(mv, mk, t) \leq C(v, k, t)$$
.

### 6.2. Steiner Systems

A *Steiner system* is a covering in which the covering density is 1—every t-set is covered exactly once. Clearly a Steiner system is an optimal covering, as well as an optimal packing, and C(v,k,t) = L(v,k,t). The projective and affine coverings by lines (1-flats), for example, are Steiner systems. Brouwer, Shearer, Sloane, and Smith [3, page 1342] and Chee, Colbourn, and Kreher [5] give tables of small Steiner systems.

If a (v,k,t) Steiner system exists then C(v+1,k,t) = L(v+1,k,t). This result is due to Schönheim [27, Theorem II]; the proof also appears in Mills and Mullin [19, Theorem 1.3].

### 6.3. Turán Theory

The *Turán number*  $T(n, \ell, r)$  is the minimum number of r-subsets of an n-set such that every  $\ell$ -subset contains at least one of the r-subsets. It is easy to see that

$$C(v,k,t) = T(v,v-t,v-k)$$
,

so covering numbers are just Turán numbers reordered. The two sets of numbers, however, have been studied for different parameter ranges (de Caen's lower bound in the introduction, for instance, is useful primarily for Turán theory ranges). Most papers on coverings have  $\nu$  large compared with k and t, while most papers on Turán numbers have

n large compared with  $\ell$  and r, often focusing on the quantity  $\lim_{n\to\infty} T(n,\ell,r)/\binom{n}{r}$  for fixed  $\ell$  and r. Thus Turán theory usually studies C(v,k,t) for k and t not too far from v.

Fifty years ago Turán [37] determined  $T(n,\ell,2)$  exactly, showing that C(v,v-2,t)=L(v,v-2,t), the Schönheim lower bound. He also gave upper bounds and conjectures for T(n,4,3) and T(n,5,3), which stimulated much of the research. The results labeled 'Turán theory' in our tables either are described in recent survey papers by de Caen [8] and Sidorenko [29], or follow from constructions due to de Caen, Kreher, and Wiseman [10] or to Sidorenko [28].

Sidorenko [28] also recently told us of a Turán theory construction, similar in spirit to the combining constructions of Section 5, that improves many bounds in the table. In terms of covering theory, let x be an element occurring in the most blocks of a (v,k,t) covering, and replace x by x' and x'': If a block b did not contain x, replace it by two blocks,  $b \cup \{x'\}$  and  $b \cup \{x''\}$ ; if b did contain x, replace it by the single block  $b - \{x\} \cup \{x',x''\}$ . Finally, add a (v-1,k+1,t+1) covering on the same elements minus x' and x''. It is not hard to see that this is a (v+1,k+1,t+1) covering, and that it gives the bound

$$C(v+1,k+1,t+1)$$
  
  $\leq |(2v-k)C(v,k,t)/v| + C(v-1,k+1,t+1).$ 

### 6.4. Cyclic Coverings

Another well-known method that is often successful when applicable—when the size of a prospective covering is v—is to construct a cyclic covering: Choose some k-subset as the first block, and choose the v-1 cyclic shifts of that block as the remaining blocks. Trying this for all possible k-sets is fairly cheap, and frequently it produces a covering. The entries  $C(19,9,3) \leq 19$  and C(24,10,3) = 24 in our tables, for example, are generated by the k-sets  $1\ 2\ 3\ 4\ 6\ 8\ 13\ 14\ 17$  and  $1\ 2\ 3\ 5\ 6\ 8\ 12\ 13\ 15\ 21$ , and are unmatched by any other method.

Incidentally, if the size of a prospective covering is a multiple of v, say 2v, the same method applies by taking the cyclic shifts of two starting blocks; the few cases we tried for this variation produced no improvements in the tables.

# 6.5. Hill-Climbing

For cases of interest—with v not too large—random coverings are not very good, but hill-climbing sometimes finds good coverings: Start with a fixed number of random k-sets, say  $L(v,k,t)+\varepsilon$  for some small integer  $\varepsilon$ . Rank the k-sets by the number of t-sets they cover that no other k-set covers, and replace one with lowest rank by another random k-set. Repeat until all t-sets are covered or until time runs out.

We found a few good coverings with this method, but Nurmela and Östergård [23] went much further, using simulated annealing—a more sophisticated hill-climbing—to find many good coverings. In fact many of the bounds in the tables could be improved, by starting with a covering produced

by one of the other methods and then hill-climbing; but generally the improvements would be small.

# 7. TABLES OF UPPER BOUNDS ON C(v,k,t)

We constructed Tables 2 through 8 using the methods described above, together with results from the literature. Each table entry indicates the upper bound, the method of construction, and whether the covering is known to be optimal. We have tried to provide constructions for as many sets of parameters as possible, so we list a method of construction from this paper even when a result in the literature achieves the same bound. When two different methods produce the same size covering, we've given precedence to the method listed earlier in the Key to the tables.

About 93% of the 1631 nontrivial (v>k>t) upper bounds in the tables come from one of the constructions described in this paper. For each of the remaining upper bounds, there is a source in our reference list that describes the result, although to keep our reference list reasonably short we have often given a secondary source rather than the original. (Mills and Mullin [19] give an extensive list of previous results and references.) Sources for Steiner systems, Turán number bounds, and simulated annealing coverings appear in Sections 6.2, 6.3, and 6.5; the Todorov constructions come from papers by Todorov [31, 33, 34] and Todorov and Tonchev [36]; and the remaining upper bounds appear in Table 1. The covering number C(24, 18, 17) is listed in Table 1, even though it doesn't occur in the other tables, because it yields a (15,9,8) simple induced covering (of Section 6.1).

Gordon et al. [13] construct an optimal (12,6,3) covering, using a block-array construction. That method directly extends to the (18,9,4) covering given in Table 1, and a similar construction gives four other coverings listed in the table.

Most of the lower bounds used to establish optimality follow from the Schönheim inequality (Theorem 1); and a few others are listed as equalities in Table 1. For the rest: If t = 2, the lower bound is explained by Mills and Mullin [19] when it is less than 14 or has  $v \le 5$ , or explained by Todorov [34] otherwise; if t = 3, it's either Mills and Mullin or Todorov and Tonchev [36]; and if  $4 \le t \le 8$ , it's either Mills [18, Theorem 2.3], Todorov [32, Theorem 4], or Sidorenko's Turán theory survey [29].

How good are our bounds? For t = 2, very good—most of the entries are known to be optimal, and the largest gap between an entry's lower and upper bound is currently only a factor of 1.12. That largest gap rises with t, though, to 1.89 for t = 4, to 2.98 for t = 6, and to 3.72 for t = 8. We believe that our lower bounds tend to be closer to the truth than our upper bounds; it's quite possible that all the upper bounds are within a factor of 3, but probably not a factor of 2, of optimal.

Most of the entries in the tables for t>2 are not optimal, and we would appreciate knowing of any better coverings. Please send communications to the first author, at gordon@ccrwest.org.

	1
bound	reference
$C(29,5,2) \le 44$	Lamken [16]
C(31,7,2) = 26	Todorov [34] techniques (lower bound)
C(12,6,3) = 15	Gordon et al. [13]
$C(14,6,3) \le 25$	Lotto covering [17]
$C(15,6,3) \le 31$	Lotto covering [17]
$C(16,6,3) \le 38$	Hoehn [14]
C(18,6,3) = 48	Lotto covering [17]
$C(30,6,3) \le 237$	Lotto covering [17]
C(11,7,4) = 17	Sidorenko [28]
$C(14,6,4) \le 87$	Hoehn [14]
$C(18,6,4) \le 258$	Lotto covering [17]
$C(18,9,4) \le 43$	Gordon et al. [13]
$C(20, 10, 4) \le 43$	block-array construction
$C(24, 12, 5) \le 86$	block-array construction
$C(30,15,5) \le 120$	block-array construction
$C(12,8,6) \le 51$	Morley [20]
$C(32, 16, 6) \le 286$	block-array construction
C(15, 12, 8) = 30	Radziszowski and Sidorenko [24]
C(24, 18, 17) = 21252	de Caen [8]

Table 1: Miscellaneous results

# Key to Tables 2 through 8

l — greedy covering, lexicographic order

c — greedy covering, colex order

g — greedy covering, Gray code order

r — greedy covering, random order

p — projective geometry covering

a — affine geometry covering

o — cyclic covering

m — multiple of smaller covering

e — simple dynamic programming (Section 6.1)

i — simple induced covering (Section 6.1)

d — dynamic programming method (Section 5)

*i* — induced covering

u — Sidorenko Turán construction (Section 6.3)

s — Steiner system

t — Turán theory

x — covering with small k and t; see Mills and Mullin [19, §3]

y — covering with fixed size; see Mills and Mullin [19, §4]

v — Todorov construction

w — was known previously; see Table 1

n — Nurmela-Östergård simulated annealing covering

*h* — hill-climbing

\* — optimal covering

$v \setminus k$	3 4 5 6 7 8 9 10 11 12 13 14 15 16
3	1*
4	$3^{l*} 1^*$
5	$4^{l*} 3^{l*} 1^*$
6	$6^{o*}3^{l*}3^{l*}1^{*}$
7	$7^{l*} 5^{l*} 3^{l*} 3^{l*} 1^{*}$
8	$11^{l*} 6^{l*} 4^{l*} 3^{l*} 3^{l*} 1^{*}$
9	$12^{r*} 8^{l*} 5^{l*} 3^{l*} 3^{l*} 3^{l*} 1^{*}$
10	$17^{r*} 9^{l*} 6^{j*} 4^{m*} 3^{l*} 3^{l*} 3^{l*} 1^{*}$
11	$19^{r_1}1^{0*}7^{r*}6^{l*}4^{l*}3^{l*}3^{l*}3^{l*}1^{*}$
12	$24^{r}12^{o*}9^{r*}6^{l*}5^{l*}3^{l*}3^{l*}3^{l*}3^{l*}1^{*}$
13	$26^{r}13^{l}10^{l}7^{l}6^{l}4^{d}3^{l}3^{l}3^{l}3^{l}1^{l}$
14	$33^{r}18^{l}12^{l}7^{m}6^{v}5^{l}4^{l}3^{l}3^{l}3^{l}3^{l}1^{s}$
15	$35^{l}19^{r}13^{r}10^{l}7^{l}6^{l}4^{m}3^{l}3^{l}3^{l}3^{l}3^{l}1^{*}$
16	43 <sup>1</sup> *20 <sup>a</sup> *15 <sup>r</sup> *10 <sup>1</sup> * 8 <sup>y</sup> * 6 <sup>1</sup> * 5 <sup>1</sup> * 4 <sup>m</sup> *3 <sup>1</sup> * 3 <sup>1</sup> * 3 <sup>1</sup> * 3 <sup>1</sup> * 3 <sup>1</sup> * 1 <sup>*</sup>
17	46 <sup>r</sup> 26 <sup>c</sup> 16 <sup>r</sup> 12 <sup>l*</sup> 9 <sup>r*</sup> 7 <sup>l*</sup> 6 <sup>l*</sup> 5 <sup>l*</sup> 4 <sup>l*</sup> 3 <sup>l*</sup> 3 <sup>l*</sup> 3 <sup>l*</sup> 3 <sup>l*</sup> 3 <sup>l*</sup>
18	54*27*18°12**********************************
19	57, 31x19°15°111* 91* 71* 61* 51* 4e* 31* 31* 31* 31*
20	$67^{r}35^{r}21^{c}16^{v}12^{l*}9^{r*}7^{j*}6^{l*}6^{l*}4^{m*}4^{l*}3^{l*}3^{l*}3^{l*}$
21	$70^{j}37^{x}21^{l}17^{v}13^{l}11^{l}7^{m}7^{l}6^{l}5^{l}4^{e}3^{l}3^{l}3^{l}$
22	81 <sup>r</sup> 39 <sup>x</sup> 27 <sup>l</sup> *19 <sup>m</sup> f3 <sup>y</sup> 11 <sup>l</sup> * 9 <sup>y</sup> * 7 <sup>m</sup> *6 <sup>y</sup> * 6 <sup>l</sup> * 5 <sup>l</sup> * 4 <sup>m</sup> *3 <sup>l</sup> * 3 <sup>l</sup> *
23	$85^{j}46^{v}28^{l}21^{v}16^{v}12^{l}10^{l}*8^{j}*7^{l}*6^{l}*5^{l}*4^{d}*4^{l}*3^{l}*$
24	$96^{j}48^{x}30^{j}22^{v}17^{v}12^{m}1^{l*}8^{v*}7^{j*}6^{l*}6^{l*}5^{l*}4^{m*}3^{l*}$
25	$100^{j}50^{j}30^{a}23^{v}18^{v}13^{j}11^{l}10^{l}7^{v}7^{l}6^{l}5^{v}4^{m}4^{e}$
26	$113^{e}59^{e}37^{e}24^{v}20^{j}13^{m}12^{l}10^{m}8^{v}7^{m}6^{v}6^{l}5^{l}4^{m}$
27	117 <sup>a</sup> 61 <sup>x</sup> 38 <sup>x</sup> 27 <sup>o</sup> 20 <sup>v</sup> 17 <sup>l</sup> 12 <sup>m</sup> 11 <sup>e</sup> 9 <sup>v</sup> 7 <sup>j</sup> 7 <sup>l</sup> 6 <sup>l</sup> 5 <sup>m</sup> 5 <sup>l</sup>
28	$131^{e}63^{s}43^{d}28^{o}22^{v}18^{r}14^{j}11^{l}10^{l}*7^{m*7}e^{*}6^{l}*6^{l}*5^{l}*$
29	$136^{j}73^{e}44^{w}31^{j}24^{v}18^{l}4^{v}12^{l}10^{v}9^{j}7^{e}7^{l}6^{l}6^{l}$
30	$150^{j}75^{x}48^{x}31^{j}25^{v}19^{m}15^{v}13^{m}1^{y*}9^{m*}8^{j*}7^{m*}6^{m*}6^{l*}$
31	$155^{l}78^{x}50^{x}31^{p}26^{v}20^{j}18^{l}13^{v}12^{l}10^{l}8^{v}7^{j}7^{l}6^{l}$
32	171 <sup>l</sup> *88 <sup>x</sup> *54 <sup>j</sup> 38 <sup>e</sup> *31 <sup>l</sup> 20 <sup>m</sup> *19 <sup>r</sup> 15 <sup>m</sup> 12 <sup>l</sup> *10 <sup>m</sup> *9 <sup>y</sup> * 7 <sup>y</sup> * 7 <sup>e</sup> * 6 <sup>l</sup> *

Table 2: t = 2

$v \backslash k$	4	5	6	7	8	9	10	11	12	13	14	15 1	16
4	1*												
5	$4^{l*}$	1*											
6	6°*	$4^{l*}$	1*										
7	12 <sup>r*</sup>	$5^{l*}$	$4^{l*}$	1*									
8	$14^{l*}$	80*	$4^{l*}$	$4^{l*}$	1*								
9	$25^{l*}$	$12^{l*}$	$7^{j*}$		•	1*							
10	30 <sup>r*</sup>	17 <sup>r*</sup>	$10^{l*}$	$6^{l*}$	$4^{l*}$	$4^{l}$	* 1*						
11	47 <sup>r</sup> *	$20^{j*}$	11 <sup>o*</sup>	$8^{r*}$	$5^{l*}$	$4^{l}$	* 4 <sup>l</sup>	* 1*					
12	57 <sup>x*</sup>												
13	78 <sup>x</sup> *												
14	91 <sup>s*</sup>	47 <sup>e</sup>	$25^{w}$	$14^{i*}$	11 <sup>h</sup>	* 8 <sup>d</sup>	* 5 <sup>n</sup>	*4 <sup>l</sup>	* $4^{l}$	$^*4^{l}$	* 1*		
15	124 <sup>e*</sup>												
16	$140^{l*}$												
17	183 <sup>l*</sup>												
18	207 <sup>x</sup> *												
19	261 <sup>e</sup> 1												
20	285 <sup>s</sup> *1	45 <sup>e</sup>	75 <sup>d</sup>										
21	352 <sup>e</sup> *1				42 <sup>e</sup>								
22	385 <sup>j</sup> *2					-					-	-	-
23	466 <sup>e</sup> *2												
24	510 <sup>x</sup> *2												
25	600 <sup>x</sup> *2												
26	650 <sup>s</sup> *2												
27	763 <sup>e</sup> *3												-
28	819s*3												
29	950 <sup>e</sup> *4												
30	1020 <sup>x</sup> *5												
31	1170 <sup>e</sup> 5												
32	1240 <sup>l</sup> *6	19 <sup>c</sup> 3	12 <sup>d</sup> 1	86 <sup>e</sup> 1	40 <sup>m</sup>	90 <sup>d</sup>	67ª	52 <sup>i</sup>	38 <sup>n</sup>	$32^{o}$	$24^{d}$	22 <sup>e</sup> 1	14 <sup>m*</sup>

Table 3: t = 3

$v \backslash k$	5	6	7	8	9	10	11	12	13	14 1	5 16	ó
5	1*											_
6	$5^{l*}$	1*										
7	$9^{l*}$	$5^{l*}$	1*									
8	20 <sup>r*</sup>	$7^{j*}$	$5^{l*}$	1*								
9	30 <sup>r*</sup>	$12^{l*}$	$6^{l*}$	$5^{l*}$	1*							
10	51 <sup>r</sup>	$20^{r*}$	10°*		$5^{l*}$	1*						
11	66 <sup>j*</sup>	32 <sup>n*</sup>	$17^{j*}$	9 <sup>j</sup> *		$5^{l*}$						
12	113 <sup>e</sup> *	$41^{n}$	$24^{n}$	12 <sup>d*</sup>								
13	157 <sup>n</sup>	66 <sup>n</sup>	$30^{n}$	19 <sup>r</sup>								
14	235 <sup>e</sup>	87 <sup>w</sup>		27 <sup>r</sup>	16 <sup>d</sup>	9 <sup>m</sup>				* 1*		
15		134 <sup>e</sup>	59 <sup>j</sup>				$8^{d*}$			* 5 <sup>l</sup> *		
16		178 <sup>d</sup>	_	30 <sup>a</sup> *		19 <sup>d</sup>	$12^{j}$	$7^{m}$			$5^{l*}$ 1	
17			119 <sup>d</sup>				16 <sup>j</sup>				$5^{l*}$ 5	
18			157 <sup>r</sup>		43 <sup>w</sup>			$12^l$		-	5 <sup>l</sup> * 5	
19	926 <sup>u</sup>		187 <sup>d</sup>								$6^{l*}$ 5	
20			246 <sup>l</sup> 1					$20^{o}$			*8 <sup>d</sup> * 5	
21	1431 <sup>g</sup>										$9^{m*7}$	
22	1746 <sup>g</sup>		253 <sup>j</sup> 1					$31^j$			$2^{d}$ 9	
23	1771 <sup>j</sup> *										$5^{d}11$	
24	2237 <sup>e</sup> *1		357 <sup>l</sup> *2								$9^d 12$	
25	2706 <sup>u</sup> 1 3306 <sup>e</sup> 1										23 <sup>a</sup> 17 27 <sup>d</sup> 19	
26 27	3906 <sup>u</sup> 1											
28	4669 <sup>e</sup> 1											
29	5427 <sup>u</sup> 1											
30	6239 <sup>l</sup> 2											
31	$6852^{j}2$											
-	7843 <sup>l</sup> 3											
32	1045 3	∠00° I	/12:0	2000	נייסטני	172"	240°	133	120"	70"3	94° 30	,

Table 4: t = 4

$v \setminus k$	6	7	8	9	10	11	12	13	14	15	16
6	1*										
7	$6^{l*}$	1*									
8	$12^{l*}$	$6^{l*}$	1*								
9	30 <sup>r</sup> *	90*	$6^{l*}$	1*							
10	50 <sup>r</sup> *	$20^{d*}$	$8^{j*}$	$6^{l*}$	1*						
11	$100^{n}$	$34^{j}$	$16^{j*}$	$7^{l*}$	$6^{l*}$	1*					
12	132 <sup>s*</sup>	59 <sup>n</sup>	$26^{t*}$	12 <sup>o*</sup>	$6^{l*}$	$6^{l*}$					
13	245 <sup>e</sup> *	88 <sup>n</sup>	$43^{n}$	$19^{d}$	$11^{j*}$			1*			
14	385 <sup>u</sup>	$154^e$	66 <sup>r</sup>	36 <sup>r</sup>	14 <sup>o*</sup>	$10^{j*}$	$6^{l*}$		1*		
15	$620^{e}$	$224^u$	$108^{r}$	$49^d$	$30^{r}$	$13^{d*}$	9 <i>j</i> *	$6^{l*}$	$6^{l*}$	1*	
16	$840^{l}$	$358^e$	$118^{l}$	79 <sup>e</sup>	$41^d$	$22^d$	$12^m$	$8^{l*}$	$6^{l*}$		* 1*
17	1277 <sup>e</sup>	506 <sup>r</sup>	$208^{e}$	94 <sup>u</sup>	$58^{d}$	$36^{j}$	$17^d$	11 <sup>j*</sup>	$7^{l*}$	$6^l$	* 6 <sup>l</sup> *
18	1791 <sup>u</sup>	$696^{l}$	$296^{d}$	149 <sup>e</sup>	$71^{d}$		$24^{d}$		9 <sup>m</sup>	* 6 <sup>l</sup>	* 6 <sup>l</sup> *
19	$2501^{l}$	$930^{l}$	419 <sup>g</sup>	199 <sup>u</sup>	$113^{d}$		$39^d$		$14^{j}$	9 <sup>e</sup>	* 6 <sup>l</sup> *
20	3297 <sup>g</sup>	$1239^{l}$	541 <sup>c</sup>	$267^{d}$	$130^i$		$42^d$		$18^d$		* 8 <sup>m</sup> *
21	4322 <sup>g</sup>	$1617^{l}$	677 <sup>g</sup>	$369^{d}$	199 <sup>d</sup>	$110^d$			$28^d$	$16^a$	$12^e$
22	5558 <sup>g</sup>	$2088^{l}$	746 <sup>c</sup>		$241^{i}$	$150^{d}$		58 <sup>d</sup>	$34^m$	$22^o$	$14^d$
23	7064 <sup>g</sup>		759 <sup>c</sup>	$622^{d}$		194 <sup>d</sup>			52 <sup>d</sup>	-	
24	7084 <sup>s*</sup>			$748^{d}$		$266^d$			59 <sup>m</sup>		
25		$4121^{l}$ 1			$494^{d}$	$335^{i}$ 1			67 <sup>j</sup>	-	$37^d$
26	11954 <sup>u</sup>	$4680^{j}1$							$67^{i}$	62 <sup>a</sup>	$43^{m}$
27	15260 <sup>e</sup>	$4680^{j}2$								67 <sup>e</sup>	$50^d$
28	19042 <sup>u</sup>	4680 <sup>s</sup> *2								, ,	55 <sup>d</sup>
29	23711 <sup>e</sup>	6169 <sup>e</sup> *3									$62^{j}$
30	28960 <sup>u</sup>	7991 <sup>u</sup> 4									
31	33715 <sup>j</sup>	9966 <sup>d</sup> 5									
32	36544 <sup>l</sup> 1	2660 <sup>d</sup> 6	6430 <sup>d</sup> 3	3465 <sup>u</sup> 2	2069 <sup>d</sup> 1	263 <sup>l</sup> 7	744 <sup>i</sup> 4	196 <sup>i</sup> 2	293 <sup>d</sup> 1	191 <sup>a</sup>	62 <sup>a*</sup>

Table 5: t = 5

$v \backslash k$	7	8	9	10	11	12	13	14	15	16	$v \backslash k$	8	9	10	11	12	13	14	15	16
7	1*										8	1*								
8	$7^{l*}$	1*									9	$8^{l*}$	1*							
9	$16^{l*}$	$7^{l*}$	1*								10	$20^{j*}$	$8^{l*}$	1*						
10	45 <sup>r*</sup>	$12^{j*}$	$7^{l*}$	1*							11	63 <sup>j</sup> *	$15^{l*}$	$8^{l*}$	1*					
11	$84^{j}$	$29^{t*}$	$10^{j*}$	$7^{l*}$	1*						12	$126^{t}$	$40^{r*}$	12°*	$8^{l*}$	1*				
12	$177^{n}$	51 <sup>w</sup>	$22^{d*}$	-	,	1*					13	$297^{n}$	$79^{n}$	$30^{d}$	$11^{j*}$	$8^{l*}$	1*			
13	$264^{n}$	$104^{n}$	$40^{t}$	16 <sup>d</sup> *	$8^{l*}$	•	1*				14	$474^{j}$	$183^{e}$	$58^{t}$	$22^{d*}$	$10^{j*}$	$8^{l*}$	1*		
14	509 <sup>e</sup>	$179^{u}$	81 <sup>r</sup>	$29^{d}$	14 <sup>0*</sup>	$7^{l*}$	$7^{l*}$	1*			15	983 <sup>e</sup>	$325^{d}$	$132^{r}$	$45^d$	$18^{d*}$	$9^{l*}$	$8^{l*}$	1*	
15	869 <sup>u</sup>	$333^e$	$128^{d}$	59 <sup>d</sup>	$21^{j}$	$13^{j*}$	$7^{l*}$	$7^{l*}$	1*		16	1806 <sup>u</sup>	$636^d$	$232^{d}$	99 <sup>r</sup>	$28^d$	16 <sup>0*</sup>	$8^{l*}$	$8^{l*}$	1*
16	1489 <sup>e</sup>	$522^{u}$	$219^{r}$	$95^{d}$	$46^d$	$19^{j}$	$12^{j*}$		,	1*	17	3295 <sup>e</sup>	$1093^{r}$	$407^{r}$	$163^{d}$	$72^{d}$	$26^{j}$	$15^{j*}$	$8^{l*}$	$8^{l*}$
17	$2234^{u}$	$829^{r}$	$305^{u}$	156 <sup>r</sup>	$70^d$	$36^{j}$	$17^{d}$	$11^{j*}$	$7^{l*}$	$7^{l*}$	18	5354 <sup>u</sup>	$1775^{c}$	659 <sup>c</sup>	$283^{r}$	$122^{d}$	$50^d$	$24^{j}$	$14^{j*}$	$8^{l*}$
18	3511 <sup>e</sup>	$1240^{r}$	506 <sup>r</sup>	$213^{d}$		55 <sup>r</sup>	$28^d$	15 <sup>j</sup> *	$10^{j*}$	$7^{l*}$	19	8865 <sup>e</sup>	$2800^{l}$	$1048^{r}$	$448^{r}$	$210^{d}$	$90^{d}$	$42^{j}$	$19^{d*}$	
19	$5219^{u}$	$1802^{l}$	$737^{r}$	$345^{r}$	164 <sup>d</sup>	$93^d$	$42^{j}$	$22^{d}$	13 <sup>j*</sup>	$9^{l*}$	20	$13838^{l}$	$4277^{c}$	1607 <sup>r</sup>	693 <sup>r</sup>	$327^{r}$		$60^d$	$34^{d}$	$17^{j*}$
20	$7522^{g}$	$2550^{l}$	1049 <sup>r</sup>	$492^{r}$	254 <sup>r</sup>	$126^{d}$	$71^{d}$	$32^{d}$	19 <sup>d</sup>	12 <sup>m*</sup>	21	20664 <sup>g</sup>	$6388^{l}$	$2407^{c}$	$1042^{g}$			131 <sup>e</sup>	$50^d$	$28^d$
21	$10453^{g}$	$3543^{l}$	$1466^{c}$	691 <sup>g</sup>		-, -	$94^d$	$58^d$	$27^d$	$17^{j}$	22	$30045^{g}$		3509 <sup>c</sup>			$372^{c}$		$94^d$	$40^d$
22	$14290^{g}$			947 <sup>g</sup>				, .	$46^d$		23	42944 <sup>g</sup>							$144^{d}$	$76^d$
23	19200 <sup>g</sup>										24	60164 <sup>g</sup>								
24	$25481^{g}$										25	$83017^{l}$								
25	31597 <sup>u</sup> 1	11317 <sup>c</sup>	3951 <sup>u</sup>	$2035^{d}$ 1	$160^{l}$	$647^{g}$	$329^{u}$	$203^{d}$	19 <sup>d</sup>	$82^{d}$	26	$112252^{l}$	$35103^{l}$ 1	$2896^{l}$	$5834^{l}$	$2803^{l}$ 1	1449 <sup>c</sup>	$743^{i}$	454 <sup>r</sup>	$243^{d}$
26	40918 <sup>e</sup> 1	14635 <sup>1</sup>	5067 <sup>e</sup>	$2452^{d}1$	$422^{d}$	$792^{i}$	$482^{e}$	$232^{d}$	147 <sup>d</sup>	$97^d$	27	150647 <sup>l</sup>	47150 <sup>c</sup> 1	.7597 <sup>l</sup>	$7856^{l}$	3784 <sup>c</sup> 1	1955 <sup>c</sup> 1	$073^{l}$	618 <sup>c</sup>	367 <sup>g</sup>
27	52746 <sup>u</sup> 1										-	197976 <sup>l</sup>								
28	68006 <sup>e</sup> 2											$259931^{l}$								
29	86749 <sup>u</sup> 2	26893 <sup>u</sup> 1	$0866^{d}$	5241 <sup>d</sup> 2	2857 <sup>d</sup> 1	1726 <sup>c</sup>	$965^{d}$	572 <sup>d</sup> 3	$325^{u}2$	$214^{e}$	30	337223 <sup>l</sup> 1	.06616 <sup>l</sup> 4	10540 <sup>l</sup> 1	$7879^{l}$	8641 <sup>l</sup> 4	1495 <sup>1</sup> 2	473 <sup>c</sup> 1	$427^{l}$	$741^{i}$
30	$109220^{l}$ 3	33062 <sup>e</sup> 1	$3149^{d}$	$6622^d$ 3	$3732^{d}$	2159 <sup>c</sup> 1	$155^{d}$	657 <sup>d</sup>	134 <sup>d</sup> 2	$234^{d}$		430492 <sup>j</sup> 1								
	133062 <sup>j</sup>										32	532248 <sup>l</sup> 1	.74784 <sup>l</sup> 6	66824 <sup>l</sup> 2	9423 <sup>c</sup> 1	4252 <sup>c</sup> 7	7418 <sup>g</sup> 4	097 <sup>c</sup> 2	$342^{i} 1$	$190^{i}$
32	154130 <sup>l</sup> 5	50743 <sup>u</sup> 2	21140 <sup>d</sup> 1	10556 <sup>d</sup> 5	862 <sup>c</sup> 3	3285 <sup>c</sup> 1	944 <sup>c</sup> 1	$087^{i}$	$709^{d}$	286 <sup>w</sup>				Tah	le 7: <i>t</i> =	= 7				

Table 7: t = 7

Table 6: t = 6

$v \backslash k$	9	10	11	12	13	14	15	16
9	1*							
10	$9^{l*}$	1*						
11	25 <sup>r*</sup>	$9^{l*}$	1*					
12	$84^{t*}$	$18^{l*}$	$9^{l*}$	1*				
13	$185^{t}$	$52^{t*}$	$15^{d*}$	9 <sup>l</sup> *	1*			
14	482 <sup>e</sup>	$121^{u}$	$40^{d}$	$13^{d*}$	$9^{l*}$	1*		
15	$790^{j}$	$300^{u}$	$81^{t}$	$30^{d*}$	$12^{d*}$	$9^{l*}$	1*	
16	1773 <sup>e</sup>	$553^{d}$	$209^{r}$	$65^d$	$24^{d*}$	$11^{d*}$	$9^{l*}$	1*
17	3499 <sup>u</sup>	$1160^{r}$	$393^{d}$	153 <sup>r</sup>	$44^d$	$20^{d*}$	$10^{l*}$	$9^{l*}$
18	6794 <sup>e</sup>	$2083^{c}$	$717^{r}$	$280^{r}$	$107^{d}$	$34^{d}$	18 <sup>0</sup> *	
19	$11827^{u}$	$3579^{r}$	$1227^{l}$	$487^{r}$	$192^{d}$	$76^d$	$31^d$	$17^{d*}$
20	20692 <sup>e</sup>	5934 <sup>c</sup>	$2055^{l}$	814 <sup>r</sup>	$355^{r}$	$150^{d}$	57 <sup>d</sup>	$26^{d}$
21	$33718^{g}$	$9499^{l}$	$3313^{g}$	$1321^c$	$582^{g}$	$274^{c}$	$96^d$	$49^{d}$
22	52674 <sup>g</sup>	$14900^{l}$	5186 <sup>g</sup>	$2072^{c}$	$915^{g}$	$437^{l}$	$219^{l}$	$71^{d}$
23	80027g	$22699^{g}$	$7917^{l}$	$3182^{l}$	$1410^g$	$674^{l}$	$316^d$	$160^{d}$
24	$119064^{l}$	$33830^{c}$	$11828^c$	$4765^{l}$	$2118^{l}$	$1013^{l}$	$517^{l}$	$254^{d}$
25	$172071^{l}$	$49556^{l}$	$17331^{c}$	$7000^{g}$	$3118^{l}$	$1498^{l}$	$765^{l}$	$409^{l}$
26	$246965^{l}$	$71206^{c}$	24924 <sup>c</sup> 1	10079 <sup>c</sup>	$4504^{c}$	2166 <sup>c</sup> 1	$110^{l}$	597 <sup>l</sup>
27	$347268^{l}$	100709 <sup>l</sup>	34976 <sup>l</sup> 1	$14320^{l}$	$6400^{c}$	3086 <sup>l</sup> 1	1583 <sup>c</sup>	$853^{l}$
28	$480708^{l}$ 1	140394 <sup>c</sup>	$49017^{l}$			4329 <sup>c</sup> 2	2221 <sup>c</sup> 1	$202^{g}$
29	$650404^{l}$ 1		$67625^{l}$ 2			5992°3		
30	879517 <sup>l</sup> 2		92034 <sup>l</sup> 3					
31	1174351 <sup>l</sup> 3							
32	1530641 <sup>l</sup>	467414 <sup>l</sup> 1	164722 <sup>l</sup> 6	57117 <sup>c</sup> 3	30228 <sup>c</sup> 1	.4697 <sup>l</sup> 7	7601 <sup>l</sup> 4	130 <sup>i</sup>

Table 8: t = 8

### Acknowledgments

We thank W. H. Mills, Nick Patterson, Alexander Sidorenko, D. T. Todorov, and an anonymous referee for some

constructive suggestions and for pointing out literature results we were unaware of. We are particularly grateful to Alexander Sidorenko for allowing us to publish his Turán construction in Section 6.3.

- C. T. Abraham, S. P. Ghosh, and D. K. Ray-Chaudhuri. File organization schemes based on finite geometries. *Information and Control*, 12:143–163, 1968.
- [2] Ian Anderson. Combinatorial Designs: Construction Methods, chapter 4. Halsted Press (a division of Wiley), 1990. Originally published by Ellis Horwood Limited, Chichester, England.
- [3] A. E. Brouwer, James B. Shearer, N. J. A. Sloane, and Warren D. Smith. A new table of constant weight codes. *IEEE Transactions on Information Theory*, 36(6):1334–1380, 1990.
- [4] Richard A. Brualdi and Vera S. Pless. Greedy codes. *Journal of Combinatorial Theory*, series A, 64(1):10–30, 1993.
- [5] Yeow Meng Chee, Charles J. Colbourn, and Donald L. Kreher. Simple *t*-designs with  $v \le 30$ . *Ars Combinatoria*, 29:193–258, 1990
- [6] John H. Conway and N. J. A. Sloane. Lexicographic codes: Error-correcting codes from game theory. *IEEE Transactions on Information Theory*, 32(3):337–348, 1986.
- [7] D. de Caen. Extension of a theorem of Moon and Moser on complete subgraphs. *Ars Combinatoria*, 16:5–10, 1983. The lower bound is stated in terms of Turán numbers.
- [8] D. de Caen. The current status of Turán's problem on hypergraphs. In P. Frankl, Z. Füredi, G. Katona, and D. Miklós, editors, *Extremal Problems for Finite Sets*, pages 187–197. János Bolyai Mathematical Society, Budapest, 1994. Proceedings of a conference held in Visegrád, Hungary, 1991.
- [9] D. de Caen, D. L. Kreher, S. P. Radziszowski, and W. H. Mills. On the covering of *t*-sets with *t*+1-sets: *C*(9,5,4) and *C*(10,6,5). *Discrete Mathematics*, 92:65–77, 1991.
- [10] D. de Caen, D. L. Kreher, and J. Wiseman. On constructive upper bounds for the Turán numbers T(n,2r+1,2r). Congressus Numerantium, 65:277–280, 1988. Proceedings of the Nineteenth Southeastern Conference on Combinatorics, Graph Theory, and Computing. Specific bounds appear in de Caen et al. [9].
- [11] Paul Erdős and Joel Spencer. *Probabilistic Methods in Combinatorics*, pages 74–75. Academic Press, 1974.
- [12] Daniel M. Gordon, Greg Kuperberg, Oren Patashnik, and Joel H. Spencer. Asymptotically optimal covering designs. Preprint, July 1994.
- [13] Daniel M. Gordon, Oren Patashnik, John Petro, and Herbert Taylor. Minimum (12,6,3) covers. Submitted for publication, March 1994.
- [14] Gerald Hoehn. Private email, 9 August 1994.
- [15] Donald E. Knuth. Seminumerical Algorithms, volume 2 of The Art of Computer Programming, section 3.2.1. Addison-Wesley, second edition, 1981.
- [16] E. R. Lamken, W. H. Mills, R. C. Mullin, and S. A. Vanstone. Coverings of pairs by quintuples. *Journal of Combinatorial Theory*, series A, 44(1):49–68, 1987. The result appeared originally in Lamken's PhD thesis *Coverings, Orthogonally Resolvable Designs and Related Combinatorial Configurations*, University of Michigan, 1983.
- [17] Tables for Lotto systems. In *Medicine and Sports*. Sofia, Bulgaria, 1983. Cited by D. T. Todorov, private communication.
- [18] W. H. Mills. Covering designs I: Coverings by a small number

- of subsets. Ars Combinatoria, 8:199-315, 1979.
- [19] W. H. Mills and R. C. Mullin. Coverings and packings. In Jeffrey H. Dinitz and Douglas R. Stinson, editors, *Contemporary Design Theory: A Collection of Surveys*, pages 371–399. Wiley, 1992.
- [20] M. E. Morley. The complements of some covering designs and questions on the overall significance. *Australasian Journal of Combinatorics*, 4:285–289, 1991.
- [21] M. Morley and G. H. J. van Rees. Lottery schemes and covers. *Utilitas Mathematica*, 37:159–165, 1990.
- [22] Albert Nijenhuis and Herbert S. Wilf. Combinatorial Algorithms: For Computers and Calculators, chapter 3. Academic Press, second edition, 1978.
- [23] Kari J. Nurmela and Patric R. J. Östergård. Upper bounds for covering designs by simulated annealing. *Congressus Numer*antium, 96:93–111, 1993.
- [24] S. P. Radziszowski and A. Sidorenko. Computer result. Cited by an anonymous referee.
- [25] D. K. Ray-Chaudhuri. Combinatorial information retrieval systems for files. SIAM Journal on Applied Mathematics, 16(5):973–992, 1968. Corollary 1 is the result cited.
- [26] Vojtěch Rödl. On a packing and covering problem. *European Journal of Combinatorics*, 5(1):69–78, 1985.
- [27] J. Schönheim. On coverings. *Pacific Journal of Mathematics*, 14:1405–1411, 1964.
- [28] Alexander Sidorenko. Private email: T(n,7,4) construction and T(11,7,4) lower bound, 24–25 October 1994; general Turán construction, 28 December 1994.
- [29] Alexander Sidorenko. What we know and what we do not know about Turán numbers. *Graphs and Combinatorics*, to appear.
- [30] Dennis Stanton and Dennis White. *Constructive Combinatorics*, section 1.2. Springer-Verlag, 1986.
- [31] D. T. Todorov. Combinatorial Coverings. PhD thesis, University of Sofia, 1985. Cited in another of his papers [34].
- [32] D. T. Todorov. On some covering designs. *Journal of Combinatorial Theory*, series A, 39(1):83–101, 1985.
- [33] D. T. Todorov. On the covering of pairs by 13 blocks. Comptes Rendus de l'Académie Bulgare des Sciences, 38(6):691–694, 1985.
- [34] D. T. Todorov. A table for the coverings of pairs. In *Proceedings* of the 15th Conference of the Union of Bulgarian Mathematicians, pages 472–481, 1986. The lower bound techniques also appear in a more available source [35].
- [35] D. T. Todorov. Lower bounds for coverings of pairs by large blocks. *Combinatorica*, 9(2):217–225, 1989.
- [36] D. T. Todorov and V. D. Tonchev. On some coverings of triples. Comptes Rendus de l'Académie Bulgare des Sciences, 35(9):1209–1211, 1982.
- [37] Paul Turán. Eine Extremalaufgabe aus der Graphentheorie. Mat. Fiz. Lapok, 48:436–452, 1941. In Hungarian, German summary. Cited in Mathematical Reviews, volume 8, page 284, 1947
- [38] Oswald Veblen and W. H. Bussey. Finite projective geometries. *Transactions of the AMS*, 7:241–259, 1906.