ON WHAT I DO NOT UNDERSTAND (AND HAVE SOMETHING TO SAY): PART I

SAHARON SHELAH

ABSTRACT. This is a non-standard paper, containing some problems in set theory I have in various degrees been interested in. Sometimes with a discussion on what I have to say; sometimes, of what makes them interesting to me, sometimes the problems are presented with a discussion of how I have tried to solve them, and sometimes with failed tries, anecdote and opinion. So the discussion is quite personal, in other words, egocentric and somewhat accidental. As we discuss many problems, history and side references are erratic, usually kept at a minimum ("see . . ." means: see the references there and possibly the paper itself).

The base were lectures in Rutgers Fall '97 and reflect my knowledge then. The other half, concentrating on model theory, will subsequently appear. I thank Andreas Blass and Andrzej Rosłanowski for many helpful comments.

Contents

1.	Cardinal problems and pcf	2
2.	The quest for the test: on the theory of Iterated Forcing for the	
	continuum	11
3.	Case studies for iterated forcing for the reals	18
4.	Nicely defined forcing notions	27
5.	To prove or to force, this is the question	37
6.	Boolean Algebras and iterated forcing	43
7.	A taste of Algebra	47
8.	Partitions and colourings	50
Ref	References	

¹⁹⁹¹ Mathematics Subject Classification. Primary 03Exx; Secondary 05C55, 03E99, 04-00. Key words and phrases. Set theory, cardinal arithmetic, pcf theory, forcing, iterated forcing, large continuum, nep, nicely definable forcing, combinatorial set theory, Boolean algebras, set theoretic algebra, partition calculus, Ramsey theory.

I would like to thank Alice Leonhardt for the beautiful typing. Publication 666 Based on lectures in the Rutgers Seminar, Fall 1997 are: §1, §2, §5, §7, §8.

1. Cardinal Problems and PCF

Here, we deal with cardinal arithmetic as I understand it (see [150] or [154]), maybe better called cofinality arithmetic (see definitions below). What should be our questions? Wrong questions usually have no interesting answers or none at all. Probably the most popular question is:

Problem 1.1. Is $pp(\aleph_{\omega}) < \aleph_{\omega_1}$?

Recall:

Definition 1.2. Let \mathfrak{a} be a set of regular cardinals (usually $|\mathfrak{a}| < \min(\mathfrak{a})$). We define

- 1. $pcf(\mathfrak{a}) = \{cf(\prod \mathfrak{a}/D) : D \text{ is an ultrafilter on } \mathfrak{a} \}.$
- 2. $\operatorname{cf}(\prod \mathfrak{a}) = \min\{|F| : F \subseteq \prod \mathfrak{a} \text{ and } (\forall g \in \prod \mathfrak{a})(\exists f \in F)(g \leq f)\}.$
- 3. For a filter D on \mathfrak{a} , $\operatorname{tcf}(\prod \mathfrak{a}/D) = \lambda$ means that in $\prod \mathfrak{a}/D$ there is an increasing cofinal sequence of length λ .
- 4. For a singular cardinal μ and a cardinal θ such that $cf(\mu) \leq \theta < \mu$ let

$$\operatorname{pp}_{\theta}(\mu) = \sup \big\{ \operatorname{tcf}(\prod \mathfrak{a}/I) : \quad \mathfrak{a} \subseteq \operatorname{Reg} \cap \mu, \ |\mathfrak{a}| < \min(\mathfrak{a}), \ \sup(\mathfrak{a}) = \mu, \\ I \text{ an ideal on } \mathfrak{a} \text{ such that } J_{\mathfrak{a}}^{\operatorname{bd}} \subseteq I, \text{ and } |\mathfrak{a}| \leq \theta \big\},$$

were for a set A of ordinals with no last element, J_A^{bd} is the ideal of bounded subsets of A.

- 5. Let $pp(\mu) = pp_{cf(\mu)}(\mu)$.
- 6. We define similarly $pp_{\Gamma}(\mu)$ for a family (equivalently: a property) Γ of ideals; e.g., $\Gamma(\theta,\tau)$ = the family of $(<\tau)$ -complete ideals on a cardinal $<\theta$, $\Gamma(\theta)$ = $\Gamma(\theta^+,\theta)$.

Definition 1.3. 1. For a partial order P,

$$\operatorname{cf}(P) = \min\{|Q| : Q \subseteq P \text{ and } (\forall p \in P)(\exists q \in Q)(p \leq q)\}.$$

2. For cardinals $\lambda, \mu, \theta, \sigma$,

$$\operatorname{cov}(\lambda, \mu, \theta, \sigma) = \min\{|\mathcal{A}| : \quad \mathcal{A} \subseteq [\lambda]^{<\mu} \text{ and any } a \in [\lambda]^{<\theta} \text{ is included}$$
in the union of $<\sigma$ members of \mathcal{A} }.

Problem 1.1 is for me the right form of

Question 1.4. 1. Assume \aleph_{ω} is strong limit. Is $2^{\aleph_{\omega}} < \aleph_{\omega_1}$? 2. Assume $2^{\aleph_0} < \aleph_{\omega}$. Is $(\aleph_{\omega})^{\aleph_0} < \aleph_{\omega_1}$?

2. Assume
$$2^{\aleph_0} < \aleph_{\omega}$$
. Is $(\aleph_{\omega})^{\aleph_0} < \aleph_{\omega}$?

Why do I think 1.1 is a better form? Because we know that:

- $(*)_1$ If \aleph_{ω} is strong limit, then $2^{\aleph_{\omega}} = (\aleph_{\omega})^{\aleph_0}$ (classical cardinal arithmetic).
- $(*)_2 \operatorname{pp}(\aleph_{\omega}) = \operatorname{cf}([\aleph_{\omega}]^{\aleph_0}, \subseteq) \text{ (see [95])},$
- $(*)_3 \aleph_{\omega}^{\aleph_0} = 2^{\aleph_0} + \operatorname{cf}([\aleph_{\omega}]^{\aleph_0}, \subseteq) \text{ (trivial)}.$

So the three versions are equivalent and say the same thing when they say something at all, but Problem 1.1 is always meaningful.

To present what I think are central problems, we can start from what I called the solution of the "Hilbert's first problem", see [119] (though without being seconded).

Theorem 1.5. For $\lambda \geq \beth_{\omega}$, there are $\kappa < \beth_{\omega}$ and $\mathcal{P} \subseteq [\lambda]^{\triangleleft \beth_{\omega}}$, $|\mathcal{P}| = \lambda$ such that every $A \in [\lambda]^{\leq \beth_{\omega}}$ is equal to the union of $< \kappa$ members of \mathcal{P} .

¹that is, most people who are aware of this direction, will mention it, and probably many have tried it to some extent

So \mathcal{P} is "very dense". E.g., if $c : [\lambda]^n \longrightarrow \beth_n$ then for some $B_m \in \mathcal{P}$ (for $m < \omega$), the restrictions $c \upharpoonright [B_m]^n$ are constant and $|B_m| = \beth_m$. We can replace \beth_ω by any strong limit cardinal $> \aleph_0$.

In [98, $\S 8$] the following application of 1.5 to the theory of Boolean Algebras is proved:

Theorem 1.6. If B is a c.c.c. Boolean algebra and $\mu = \mu^{\beth_{\omega}} \leq |B| \leq 2^{\mu}$, then B is μ -linked, i.e., $B \setminus \{0\}$ is the union of μ sets of pairwise compatible elements.

(See also [129], [131] and Hajnal, Juhász and Szentmiklossy [53].) We also have the following application:

Theorem 1.7 (See [155]). If X is a topological space (not necessarily T_2) with λ points, $\mu \leq \lambda < 2^{\mu}$ and $> \lambda$ open sets and μ is strong limit of cofinality \aleph_0 , then X has $\geq 2^{\mu}$ open sets.

Another connection to the general topology is the following

Definition 1.8. For topological spaces X, Y and a cardinal θ , write $X \to (Y)^1_{\theta}$ iff for every partition $\langle X_i : i < \theta \rangle$ of X into θ parts, X has a closed subspace Y' homeomorphic to Y which is included in one part of the partition.

Arhangel'skii asked whether for every compact Hausdorff space X,

$$X \rightarrow (Cantor discontinuum)_2^1$$
.

Arhangel'skii's problem $+\neg CH$ is sandwiched between two pcf statements of which we really do not know whether they are true. If, for simplicity $2^{\aleph_0} \ge \aleph_3$, then e.g.:

- $(*)_1$ if for no \mathfrak{a} , $\mathfrak{a} \subseteq \text{Reg}$, where Reg is the class of regular cardinals, $|\mathfrak{a}| \geq \aleph_2$, $\prod \mathfrak{a}/[\mathfrak{a}]^{\leq \aleph_0}$ is $\sup(\mathfrak{a})$ -directed, then the answer is: for every Hausdorff space X, we have $X \nrightarrow (\text{Cantor discontinuum})_2^1$ and more.
- (*)₂ If for some $\mathfrak{a} \subseteq \operatorname{Reg} \setminus 2^{<\kappa}$, $|\mathfrak{a}| = 2^{\aleph_0} \le \kappa$ and $\prod \mathfrak{a}/[\mathfrak{a}]^{\le \aleph_0}$ is $\sup(\mathfrak{a})$ -directed, then in some forcing extension there exists a zero-dimensional Hausdorff space X such that $X \to (\operatorname{Cantor\ discontinuum})_2^1$.

The Stone-Čech compactification of this space gives a negative answer to Arhangel'skii's question.

(On the problem, see [119] and more in [112].)

However, we can start from inside pcf theory.

Problem 1.9. Is $pcf(\mathfrak{a})$ countable for each countable set of cardinals?

This seems to me more basic than 1.1, but yet 1.1 is weaker. I think it is better to look at the battlefield between independence by forcing from large cardinals and proofs in ZFC (I would tend to say between the armies of Satan and God but the armies are not disjoint).

The advances in pcf theory show us ZFC is more powerful than expected before. I will try to give a line of statements on which both known methods fail – so far.

Conjecture 1.10. If \mathfrak{a} is a set of regular cardinals $> |\mathfrak{a}|$, then for no inaccessible λ the intersection $\lambda \cap \operatorname{pcf}(\mathfrak{a})$ is unbounded in λ .

Conjecture 1.11. For every $\mu \geq \aleph_{\omega}$, for every $\aleph_n < \aleph_{\omega}$ large enough there is no $\lambda < \mu$ of cofinality \aleph_n such that $\operatorname{pp}_{\Gamma(\aleph_n)}(\lambda) > \mu$ (or replace $\aleph_n < \aleph_{\omega}$ by $\aleph_{\alpha} < \aleph_{\omega^2}$ or even $\aleph_{\alpha} < \aleph_{\omega_1}$, or whatever).

Conjecture 1.12. (A) It is consistent, for any uncountable θ (e.g., \aleph_1), that for some λ

$$\theta \le |\{\mu < \lambda : \operatorname{cf}(\mu) = \aleph_0, \operatorname{pp}(\mu) > \lambda\}|.$$

(B) It is consistent that for some λ , the set

$$\{\mu < \lambda : \operatorname{cf}(\mu) > \aleph_0, \ \operatorname{pp}_{\aleph_1 - \operatorname{complete}}(\mu) > \lambda\}$$

is infinite.

Those three conjectures seem to be fundamental. Note that having ZFC-provable answer in 1.10, 1.11, but independent answer for 1.12 are conscious choices. For all of those problems, present methods of independence fail, and in addition they are known to require higher consistency strength. Of course, we can concentrate on other variants; e.g., in 1.12(B) use θ instead of \aleph_0 .

Other problems tend to be sandwiched between those, or at least those more basic problems are embedded into them. E.g., 1.11 implies that in 1.5 we can replace \beth_{ω} by \aleph_{ω} if we replace equal by included (or demand $\lambda \geq \sum_{n < \omega} 2^{\aleph_n}$) and

this implies $|\mathfrak{a}| \leq \aleph_0 \Rightarrow |\operatorname{pcf}(\mathfrak{a})| \leq \aleph_{\omega}$, while e.g. $|\mathfrak{a}| \leq \aleph_{\omega n} \Rightarrow |\operatorname{pcf}(\mathfrak{a})| \leq \aleph_{\omega n+\omega}$ implies the analog of 1.5 for \aleph_{ω^2} , see [119], [114]. See [114] for more on the ZFC side; it is very helpful in preventing futile attempts to force.

Note that $\operatorname{pp}(\aleph_{\omega}) > \aleph_{\omega_1}$, implies that for some countable \mathfrak{a} , $\operatorname{pcf}(\mathfrak{a})$ is uncountable, which implies that clause (A) from Conjecture 1.12 holds. Also $\operatorname{pp}(\aleph_{\omega}) > \aleph_{\omega_2}$ implies that for some countable \mathfrak{a} , $|\operatorname{pcf}(\mathfrak{a})| \geq \aleph_2$ which implies that clause (B) of Conjecture 1.12 fails).

So there is no point to try to prove $CON(pp(\aleph_{\omega}) > \aleph_{\omega_1})$ before having the consistency of 1.12(A) and, thus, $CON(pp(\aleph_{\omega}) > \aleph_{\omega_1})$ is a more specialized case. (Also if we look at the earlier history of consistency proofs – clearly there is no point to start with Problem 1.1).

In Conjecture 1.10 the situation (which we say is impossible) may look bizarre, as $pcf(\mathfrak{a})$ is extremely large. Of course, much better is $|pcf(\mathfrak{a})| <$ "first inaccessible $> |\mathfrak{a}|$ " and even $|pcf(\mathfrak{a})| \le |\mathfrak{a}|^{+\omega}$, which follows from Conjecture 1.11. Of course, replacing in 1.10, " λ inaccessible" by " λ -Mahlo" is still a very important conjecture while getting $pcf(\mathfrak{a}) <$ "the first fixed point $> |\mathfrak{a}|$ " is much better, so why from all variants of 1.10, those we have just mentioned and others, "the accumulation inaccessible" was chosen? The point is that it implies

(*)
$$cf(\prod pcf(\mathfrak{a})) = cf(\prod \mathfrak{a}),$$

if \mathfrak{a} is a set of regular cardinals $> |\mathfrak{a}|$

(see [154, Ch.VIII,§3], [94]; note that in the notation of [94], conjecture 1.10 says that $pcf(\mathfrak{a}) \in J_{\mathfrak{a}}^*$.) If there is a failure of Conjecture 1.10 then consistently (*) fails. We can force by (< λ)-complete forcing iterating adding $f \in \prod(pcf(\mathfrak{a}) \cap \lambda)$ dominating the old product (or for any μ , just adding μ many λ -Cohen functions, i.e., forcing with

$$\{f: f \text{ is a partial function from } \mu \text{ to } \lambda, \text{ Dom}(f) | < \lambda\}\}$$
.

So 1.10 denotes the significant dividing line between chaos and order.

Concerning the last conjecture 1.12, maybe the proofs in Gitik and Shelah [43] are relevant. There we force for hypermeasurable cardinals $\kappa_0 < \kappa_1 < \ldots < \kappa_n$ with a forcing which makes each κ_i hypermeasurable indestructible under reasonable forcing notions, including those which may add new Prikry sequences of ordinals

 $> \kappa_{\ell}$ of length $< \kappa_{\ell}$. (So in this case supercompact cannot serve, unlike in many proofs which do with hypermeasurable cardinals what is relatively easy to do with supercompact cardinals.) Let $\lambda = \lambda^{<\lambda} > \kappa_n$, $\theta_{\ell} < \lambda$. Then we blow up 2^{κ_n} to λ , change cf (κ_n) to θ_n ; blow up $2^{\kappa_{n-1}}$ to λ , change cf (κ_{n-1}) to σ_{n-1} , etc.

The point is that when we arrive to κ_i the forcing so far is fairly " κ_i -complete for pure extensions", etc, so does not destroy " κ_i is λ -hypermeasurable". So for Conjecture 1.12 we (fix the desired cofinality θ and we) need to do it not n times but θ^* times (θ^* = inverse order of θ) so we need "anti-well-founded iteration". In other words, we have $\langle \kappa_i : i < \theta \rangle$ increasing; κ_i is λ -hypermeasurable indestructible (necessarily in a strong way), and $\lambda > \sum_{i < \theta} \kappa_i$.

FIRST TRY:

We may try to define by induction on $i < \theta$, \triangleleft -decreasing sequences $\bar{\mathbb{P}}^i = \langle \mathbb{P}^i_j : j \leq i \rangle$ of forcing notions such that $|\mathbb{P}^i_0| = \lambda_i$,

$$\Vdash_{\mathbb{P}_0^i}$$
 " $\operatorname{cf}(\kappa_j) = \aleph_0$ for $j < i$ "

(or whatever fixed value, but \aleph_0 is surely easier), \mathbb{P}_j^i is κ_j^+ -c.c., purely κ_j -complete, \mathbb{P}_0^i makes $\operatorname{pp}(\kappa_j) = \lambda$ for every j < i.

In successor stage - no problem: i = j + 1 and

$$\begin{array}{rcl} \mathbb{P}^{i+1}_{i+1} & = & \text{(blowing up } 2^{\kappa} \text{ to } \lambda \text{ changing } \text{cf}(\kappa_{j}) \text{ to } \aleph_{0}) \\ \mathbb{P}^{i+1}_{j} & = & \mathbb{P}^{i+1}_{i+1} * \mathbb{P}^{i}_{j}. \end{array}$$

Not good enough: \mathbb{P}_i^{i+1} changes the definition of: "blowing up 2^{κ_j} to λ " as there are more ω -sequences. So we should correct ourselves to $|\mathbb{P}_j^i| = \sum_{\zeta \in i} \kappa_{\zeta}$:

$$\mathbb{P}_j^i$$
 blows up 2^{κ_j} to essentially $\sum_{\zeta < i} \kappa_{\zeta}$.

So we have to prove the forcing notions extend as they should. If $\bar{\mathbb{P}}^i$ is defined, there is no problem to choose an appropriate \mathbb{P}^{i+1}_{i+1} . Now for each $j \leq i$ separately we would like to choose \mathbb{P}^{i+1}_j to be a \lessdot -extension of \mathbb{P}^i_j and of \mathbb{P}^{i+1}_{i+1} , but we have to do it for all $j \leq i$ together. The limit case seems harder.

* * *

Why, in 1.12(A), do we have $\theta \geq \aleph_1$? Moti Gitik shows consistency for $\theta = \aleph_0$ by known methods.

AUDIENCE QUESTION: How dare you conjecture ZFC can show 1.10, 1.11?

For Conjecture $1.12\,\mathrm{I}$ have a scenario for an independence proof (outlined above). For 1.10 and 1.11 the statements imply there is quite a complicated pcf structure you necessarily drag with you. So it is reasonable to assume that if we shall know enough theorems on the pcf structure we shall get a contradiction. Of course, those arguments are not decisive.

* * *

Traditionally, remnants of GCH have strongly influenced the research on cardinal arithmetic, so e.g. people concentrate on the strong limit case, see [154, AG], [95]; probably also it was clear what to do and easier. On the other hand, [154] aims to get "exponentiation-free theorems", so we put forward:

Thesis 1.13. "Everything" is expressibly by cases of pp_J (and 2^{κ} for κ regular).

E.g. in [96, §2] this is done to the tree power of λ ,

$$\lambda^{\langle \kappa, \operatorname{tr} \rangle} = \sup\{|\lim_{\kappa} (T)| : T \text{ a tree with } \leq \lambda \text{ nodes and } \kappa \text{ levels}\},$$

where $\lim_{\kappa}(T)$ is the set of κ -branches of T (well, using $\kappa^{\langle \kappa, \text{tr} \rangle}$ for regular κ , which is malleable by forcing, a relative of 2^{κ} for κ regular).

But maybe there are also forcing proofs by which we can get interesting situations say below the continuum, whose strong limit counterparts are false, or have bigger consistency strength, or at least are harder to prove. The known forcing proofs may be open to such variations, e.g., when we add many Prikry sequences to one κ we may have the order between them such that every condition decides little about it. The following problem may be relevant to 1.13, and anyhow is a central one.

Problem 1.14. For a singular cardinal $\mu > \theta = cf(\mu)$, is

$$cov(\mu, \mu, \theta^+, \theta) = pp_{\Gamma(\theta^+, \theta)}(\mu)$$
?

Note that other cases of cov can be reduced to those above. Now, this is almost proved: it holds when $\theta = \operatorname{cf}(\mu) > \aleph_0$. Furthermore, if μ is strong limit, $\aleph_0 = \operatorname{cf}(\mu)$ and the two expressions in 1.14 are not equal, both are quite large above μ as in Conjecture 1.12. Also, e.g., for a club of $\delta < \omega_1$

$$cov(\beth_{\delta}, \beth_{\delta}, \aleph_1, \aleph_0) = pp(\beth_{\delta}),$$

(see [95], the " \beth_{ω_1} " can be weakened to strong limit in cov sense). But

Question 1.15. Can we force that there is $\mu < 2^{\aleph_0}$ such that $cf(\mu) = \aleph_0$ and $cov(\mu, \mu, \aleph_1, \aleph_0) > pp(\mu)$?

[Why $< 2^{\aleph_0}$? As blowing up the continuum does not change the situation, proving the consistency for $\mu < 2^{\aleph_0}$ can be only easier. But for $\mu < 2^{\aleph_0}$ maybe it is even consistent that

$$cov(\mu, \mu, \aleph_1, \aleph_0) > \mu^+ = pp(\mu),$$

that is, by our present ignorance, it is even possible that the behaviour below the continuum is different than above it.]

Note that all cases of λ^{κ} can be reduced to cases of 2^{θ} , θ regular, and $\operatorname{cf}([\mu]^{\leq \theta}, \subseteq)$ where $\mu > \operatorname{cf}(\theta) = \theta \geq \operatorname{cf}(\mu)$.

Why? If κ is regular, $\lambda \leq 2^{\kappa}$ then $\lambda^{\kappa} = 2^{\kappa}$. If κ is regular and $\lambda > 2^{\kappa}$ then $\lambda^{\kappa} = \operatorname{cf}([\lambda]^{\leq \kappa}, \subseteq)$. So assume κ is singular and let $\sigma = \operatorname{cf}(\kappa)$ and $\kappa = \sum_{i < \sigma} \kappa_i$, where

each κ_i is regular and $\sigma < \kappa_i < \kappa$, so $\lambda^{\kappa} = \lambda^{\sum \{\kappa_i : i < \sigma\}} = \prod_{i < \sigma} \lambda^{\kappa_i}$. Thus, if $\lambda \leq 2^{\kappa}$ then

$$\lambda^{\kappa} = 2^{\kappa} = \prod_{i < \sigma} 2^{\kappa_i} = (\sum_{i < \sigma} 2^{\kappa_i})^{\sigma} = \operatorname{cf}([\sum_{i < \sigma} 2^{\kappa_i}]^{\sigma}, \subseteq).$$

Lastly, if $\lambda > 2^{\kappa}$

$$\lambda^{\kappa} = \prod_{i < \sigma} \lambda^{\kappa_i} = (\sum_{i < \sigma} \lambda^{\kappa_i})^{\sigma} = (\max_{i < \sigma} \ \lambda^{\kappa_i})^{\sigma} = \max_{i < \sigma} \ \lambda^{\kappa_i} = \max_{i < \sigma} \ \mathrm{cf}([\lambda]^{\kappa_i}, \subseteq)$$

(on the third equality see Hajnal and Hamburger [52], [140, 2.11(4),p. 164]).

If the answer to 1.14 is yes, then we can reduce all cases of λ^{κ} and of cov to statements on cases of pp.

Problem 1.16. If $cf(\mu) = \aleph_0$, is $pp(\mu)$ equal to $pp_{J^{bd}}^{cr}(\mu)$, where

 $\mathrm{pp}^{\mathrm{cr}}_{J^{\mathrm{bd}}_{\omega}}(\mu) = \sup\{\lambda: \quad \text{for some increasing sequence } \langle \lambda_n: n < \omega \rangle \text{ of regular } \\ \quad \text{cardinals converging to } \mu \text{ we have } \lambda = \mathrm{tcf}(\prod_{n < \omega} \lambda_n/J^{\mathrm{bd}}_{\omega})\} ?$

A variant is: except when $\operatorname{pp}^{\operatorname{cr}}_{J^{\operatorname{bd}}_{-}}(\mu)$ has cofinality \aleph_0 and $\operatorname{pp}(\mu)$ is its successor.

By pcf calculus, if $pp(\mu) < \mu^{+\omega_1}$ then this is true. Similarly, if $\theta < \mu_0 < \mu$ and

$$(\forall \mu')([\operatorname{cf}(\mu') \leq \theta \& \mu' \in (\mu_0, \mu) \Rightarrow \operatorname{pp}(\mu') < \mu^{+\theta^+}]$$

then $pp(\mu) = pp_{\theta}(\mu)$ and see [158, 6.5]. Also, by [95, Part C], e.g., for a club of $\delta < \omega_1, \ \mu = \beth_{\delta}$ satisfies the conclusion.

* * *

On pcf for set theories with weak versions of Choice (say DC_{κ} , the dependent choice of length κ) see [163].

Problem 1.17. Develop combinatorial set theory generally and, in particular, pcf theory using only little choice (say DC_{κ}).

Inner model theory and descriptive set theory are not hampered by lack of choice, and much was done on variants of the axiom of choice. [163] may be a beginning of combinatorial set theory, and pcf in particular; i.e., it is enough to show that there are interesting theorems. In particular

Question 1.18. (a) Does DC_{κ} for κ large enough imply the existence of a proper class of regular cardinals?

(b) Does DC_{κ} for κ large enough imply that for a class of λ , $\mathcal{P}(\lambda)$ is not the union of $<\lambda$ sets, each of cardinality $\leq \lambda$?

See more in [163]. Gitik [40] had proved

$$CON((\forall \delta)(cf(\delta) \leq \aleph_0))$$

relative to suitable large cardinals. Woodin asked if

$$CON (DC_{\aleph_0} + (\forall \delta)(cf(\delta) \leq \aleph_1))$$
.

Specker asked if, consistently, for every λ , for some $\langle A_n : n < \omega \rangle$ we have $\mathcal{P}(\lambda) = \bigcup_{n = \infty} A_n$, $|A_n| \leq \lambda$.

* * *

On how the problem of the existence of universal objects is connected to pcf see Kojman and Shelah [67], and [161], [109]. The following conjecture will simplify the answers:

Conjecture 1.19. For every limit of limit cardinals μ , for arbitrarily large regular $\lambda < \mu$, we have

$$(\forall^* \mu_1 < \mu)[\operatorname{cf}(\mu_1) = \lambda \Rightarrow \operatorname{pp}_{\Gamma(\lambda)}(\mu_1) < \mu],$$

where \forall^* means "for every large enough".

After we learned that, on the one hand, $2^{\aleph_{\omega}}$ (\aleph_{ω} strong limit) has a bound (in fact, every $2^{\aleph_{\delta}}$, if \aleph_{δ} is strong limit > $|\delta|$, in [132, Ch.XIII]), and on the other hand there are bounds for 2^{μ} , μ the ω_1 -th fix point (when μ is strong limit or less), it becomes natural to ask:

Conjecture 1.20. If \aleph_{δ} is the first fix point (i.e., the first such that $\aleph_{\delta} = \delta$, so it has cofinality \aleph_0), then $\operatorname{pp}(\aleph_{\delta}) < (2^{\aleph_0})^+$ -th fix point.

(Even assuming GCH below \aleph_{δ} and proving just $\operatorname{pp}(\aleph_{\delta})$ < "the first inaccessible" is good, but " $<\omega_4$ -th fix point" is better, and " $<\omega_1$ -th fix point" is best, but seems pointless to ask as long as 1.1 is open).

Note that we almost know: if \aleph_{δ} is the ω_1 -th fix point (strong limit), then $pp(\aleph_{\delta}) < \omega_4$ -th fix point, we know it if the answer to 1.10 is yes, see [152] and see [154, Ch.V].

* * *

Traditionally we have asked: "can we find all the laws of cardinal arithmetic?" This had been accomplished for regular cardinals, and we prefer

Problem 1.21. Find all the rules of the pcf calculus or at least find more (or show that the set of rules is inherently too complicated).

Note: if for simplicity $|\operatorname{pcf}(\mathfrak{a})| < \min(\mathfrak{a})$, then on $\operatorname{pcf}(\mathfrak{a})$ the pcf structure is naturally a compact topology: \mathfrak{b} is closed iff $\mathfrak{b} = \operatorname{pcf}(\mathfrak{b})$, and the theorem on existence of generators $\langle \mathfrak{b}_{\lambda} : \lambda \in \operatorname{pcf}(\mathfrak{a}) \rangle$ says that the topology is a particularly nice one. If 1.10 holds this is true whenever $|\mathfrak{a}| < \min(\mathfrak{a})$ (see [94]).

There may well be some "global phenomena". Also there may be special behaviour near

 $\min\{\lambda : \text{ for some } A \subseteq \lambda, \text{ there is no indiscernible class for } K[A]\},$

as above it the covering theorem (Dodd and Jensen [26]) shows that cardinal arithmetic is trivial. On the other hand, on the behaviour below it, see [154, Ch.V].

An extreme case of our non-understanding concerning global behavior is:

Question 1.22. Is it possible that:

if \mathfrak{a} is a set of odd [even] regular cardinals $> |\mathfrak{a}|$, then every $\theta \in \operatorname{pcf}(\mathfrak{a})$ is odd [even]?

(where $\aleph_{2\alpha}$ is even and $\aleph_{2\alpha+1}$ is odd).

Instead of looking more on $pp(\aleph_{\omega})$ we may ask if the best result was derived from the known laws of cardinal arithmetic.

Question 1.23. Let $\ell < 4$. Can there be $\delta \in [\omega_{\ell}, \omega_{\ell+1})$ and a closure operation $c\ell$ on $\mathcal{P}(\delta+1)$ such that all the rules used in the proof of $pp(\aleph_{\omega}) < \aleph_{\omega_4}$ hold? (see Jech and Shelah [57]).

Question 1.24. 1. Characterize the possible sequences

$$\langle J_{\leq \theta}[\{\aleph_n : n \in [1, \omega)\}] : \theta \in \operatorname{pcf}\{\aleph_n : n \in [1, \omega)\}\rangle.$$

2. For every ordinal γ characterize the possible $\langle J_{<\theta}[\mathfrak{a}] : \theta \in \operatorname{pcf}(\mathfrak{a}) \rangle$ up to isomorphism when $\operatorname{otp}(\mathfrak{a}) = \gamma$.

[For \mathfrak{a}' , \mathfrak{a}'' we have an isomorphism if there there is a one–to–one order preserving $f:\mathfrak{a}'\longrightarrow\mathfrak{a}''$ such that $\{J_{<\theta}[\mathfrak{a}'']:\theta\in\operatorname{pcf}(\mathfrak{a}'')\}=\{\{f[\mathfrak{b}]:\mathfrak{b}\in J_{<\theta}[\mathfrak{a}']\}:\theta\in\operatorname{pcf}(\mathfrak{a}')\}.$

* * *

I feel that

Thesis 1.25. Proving a theorem from ZFC + "cardinal arithmetic assumptions" is a "semi ZFC result".

This view makes proofs from cases of the failure of the SCH related to the thesis below more interesting.

Thesis 1.26. Assumptions on the failure of GCH (and even more so, of SCH) are good assumptions, practical ones, in the sense that from them you can deduce theorems.

Traditionally this is how instances of GCH were treated (with large supporting evidence). Clearly 1.26 may be supported by positive evidence (though hard to refute), whereas 1.25 remains a matter of taste. So Magidor would stress looking at "existence of a large cardinal" as semi-ZFC axioms (unlike some randomly chosen consistent theorems), which seems to mean in our terminology that we will look at consequences of it as semi-ZFC theorems. Jensen stresses that showing ψ holds in a universe with structure is much better than mere consistency (so the fine structure in **L** was the only one we know of at one time, but e.g. K is no less good than **L**; the statement in [102] was inaccurate).

I agree with both, just to a lesser degree. Kojman criticized 1.25 saying cases of failure of SCH are large cardinal assumption in disguise; and I agree that $2^{\lambda} > \aleph_{\lambda^{+4}}$ is a weaker assumption than $2^{\beth_{\omega}} > \beth_{\omega}^+$, but I still stick to 1.25. We may hope to really resolve problems by partitioning to cases according to what the cardinal arithmetic is.

* * *

DISCUSSION: The following should be obvious, but I have found that mentioning them explicitly is helpful. Assume, e.g., that $cf(\mu) = \aleph_0$, $pp(\mu) > \mu^{+\omega_n}$, n > 0 (ω_n for simplicity) and let

(*)_{μ ,n} for stationarily many $\delta < \omega_n$ of cofinality \aleph_0 , $\operatorname{pp}_{\aleph_n}(\mu^{+\delta}) < \mu^{+\omega_n}$ (a "soft" assumption, see [154, IX,§4]).

Then we can find pairwise disjoint countable $\mathfrak{a}_i \subseteq \text{Reg} \cap \mu$ unbounded in μ and $\alpha_i < \omega_n$ successor, strictly increasing and such that

$$\mu^{+\alpha_i} = \max \operatorname{pcf}(\mathfrak{a}_i), \quad \mu^{+\alpha_i} \notin \operatorname{pcf}(\bigcup_{j \neq i} \mathfrak{a}_j),$$

moreover $\mu' < \mu \implies \mu^{\alpha_i} = \max \operatorname{pcf}(\mathfrak{a}_i \setminus \mu').$

[Why? We can find, by the assumption and Fodor Lemma, $\alpha^* < \omega_n$ such that

$$\alpha \in [\alpha^*, \omega_n) \implies \max \operatorname{pcf}\{\mu^{+\beta+1} : \beta \in (\alpha_0, \alpha)\} < \mu^{+\omega_n}.$$

By the assumption $pp(\mu) \ge \mu^{+\omega_n}$, there is $\mathfrak{a} \subseteq \mu \setminus \omega_n$, $|\mathfrak{a}| = \aleph_n$ such that $\alpha < \omega_n \Rightarrow \mu^{+\alpha+1} \in pcf(\mathfrak{a})$. First assume $2^{\aleph_n} < \mu$, so without loss of generality $min(\mathfrak{a}) > 2^{\aleph_n}$,

and we have a smooth closed generating sequence $\langle \mathfrak{b}_{\lambda} : \lambda \in \operatorname{pcf}(\mathfrak{a}) \rangle$ for $\operatorname{pcf}(\mathfrak{a})$ (so $\mathfrak{b}_{\lambda} \subseteq \operatorname{pcf}(\mathfrak{a})$, etc.). Now choose by induction on $i < \omega_n$ pairs $(\alpha_i, \mathfrak{b}'_i)$ as follows. If $\langle \alpha_j : j < i \rangle$ has been defined, we know that

$$\max \ \mathrm{pcf}\{\mu^{+\beta}: \beta \ \mathrm{successor}, \, \alpha^* \leq \beta \leq (\alpha^*+2) \cup \bigcup_{j < i} \alpha_j\} < \mu^{+\omega_n},$$

and hence we can find $m_i < \omega$ and successor ordinals $\gamma_\ell^i \in [\alpha^*, (\alpha^* + 2) \cup \bigcup_{j < i} \alpha_j]$ (for $\ell < m_i$) such that

$$\{\mu^{+\beta} : \beta \text{ a successor}, \alpha^* \leq \beta \leq (\alpha^* + 1) \cup \bigcup_{j < i} \alpha_j\} \subseteq \bigcup_{\ell < m_i} \mathfrak{b}_{\gamma_\ell^i}.$$

Let $\alpha_i < \omega_n$ be the minimal successor such that

$$\mu^{+\alpha_i} > \max \operatorname{pcf}\{\mu^{+\beta} : \beta \text{ a successor, } \alpha^* \leq \beta \leq (\alpha^* + 1) \cup \bigcup_{i \leq i} \alpha_i\},$$

and let $\mathfrak{a}_i = \mathfrak{b}_{\alpha_i} \setminus \bigcup_{\ell < m_i} \mathfrak{b}_{\gamma_\ell}$. If $\neg (2^{\aleph_\omega} < \mu)$ use the end of [158, §6].]

If, weakening $(*)_{\mu,n}$, we assume that for some $\alpha^* < \omega_n$ we have

$$\delta > \alpha^* \& \delta < \omega_n \text{ is limit } \Rightarrow \operatorname{pp}(\mu^{+\delta}) < \mu^{+\omega_n},$$

then we can get the same conclusion. Of course, omitting $(*)_{\mu,n}$ if $2^{\aleph_0} < \omega_n$, by the Δ -system lemma, we can get $\langle (\mathfrak{a}_i, \alpha_i) : i < \omega_n \rangle$ as above but demanding only $i \neq j \Rightarrow \mu^{+\alpha_i} \notin \operatorname{pcf}(\mathfrak{a}_j)$. Of course, we cannot let $\alpha_i = i+1$, as e.g. for some infinite $A \subseteq \omega$, $\mu^{+\omega+1} = \operatorname{tcf}(\prod_{n \in A} \mu^{+n}/J_A^{\operatorname{bd}})$, and hence $\mu^{+\omega+1} \in \operatorname{pcf}(\bigcup_{n \in A} \mathfrak{a}_n)$.

Another remark: Even if $\operatorname{pcf}(\mathfrak{a})$ is large and \mathfrak{a} is countable, we can find a c.c.c. forcing notion \mathbb{Q} such that in $\mathbf{V}^{\mathbb{Q}}$ we can find $\langle \mathfrak{b}_{\lambda} : \lambda \in \operatorname{pcf}(\mathfrak{a}) \setminus \mathfrak{a} \rangle$ satisfying: $\mathfrak{b}_{\lambda} \subseteq \mathfrak{a}$ has order type ω and $\prod \mathfrak{b}_{\lambda}/J_{\mathfrak{b}_{\lambda}}^{\operatorname{bd}}$ has true cofinality λ . [Why? If $\langle \mathfrak{b}_{\lambda} : \lambda \in \operatorname{pcf}(\mathfrak{a}) \rangle$ is a generating sequence, let \mathbb{Q} force for each λ an ω -sequence $\subseteq \mathfrak{b}_{\lambda}$, almost disjoint to \mathfrak{b}_{λ_1} for $\lambda_1 < \lambda$.] Such forcing does not change the pcf structure (in fact, if $\langle \mathfrak{b}_{\lambda} : \lambda \in \operatorname{pcf}(\mathfrak{a}) \rangle$ is a generating sequence for \mathfrak{a} , \mathbb{Q} is a $\min(\mathfrak{a})$ -c.c. forcing notion, then $\langle \mathfrak{b}_{\lambda} : \lambda \in \operatorname{pcf}(\mathfrak{a}) \rangle$ is still a generating sequence for \mathfrak{a} , witnessed by the same $\langle f_{\alpha}^* : \alpha < \lambda \rangle$).

Question 1.27. For a regular cardinal θ , can we find an increasing sequence $\langle \lambda_i : i < \theta \rangle$ of regular cardinals such that for some successor λ and $f_{\alpha} \in \prod_{i < \theta} \lambda_i$ for $\alpha < \lambda$ we have:

(*) if C_i is a club of λ_i for $i < \theta$, then for every large enough $\alpha < \lambda$ for every large enough $i < \theta$ we have $f_{\alpha}(i) \in C_i$.

By [116, §6] an approximation to this holds: if μ is a strong limit singular cardinal, pp(μ) =⁺ 2 $^{\mu}$ and $\lambda = 2^{\mu} = \text{cf}(2^{\mu})$ then the answer is yes, i.e. (*) holds true, but 2^{μ} may be a limit cardinal (if 2^{μ} is singular, a related statement holds).

Question 1.28. Assume $\kappa = \operatorname{cf}(\kappa)$, $\langle \mu_i : i \leq \kappa \rangle$ is an increasing continuous sequence of strong limit cardinals, for nonlimit i, $\operatorname{cf}(\mu_i) = \aleph_0$ and $\prod_{i \leq \kappa} \mu_i^{+n}/J_\kappa^{\operatorname{bd}}$ has

true cofinality μ^{+n} . Can we find an interesting colouring theorem on μ^{+n} ? (The point is that for $n \geq 2$, we can have both a colouring as μ^{+n} is a successor of regulars (as in [147], [160]) and using a witness to $\operatorname{tcf}(\mu_i^{+n}/J_\kappa^{\operatorname{bd}}) = \mu^{+n}$ as in [98], [116].) The question is whether combining we shall get something startling.

- Question 1.29. 1. Are there non-metrizable first countable Hausdorff topological spaces which are \aleph_2 -metrizable (i.e., the induced topology on any $\leq \aleph_1$ points is metrizable)?
 - 2. Are there non-collectionwise Hausdorff, first countable Hausdorff topological spaces which are \aleph_1 -collectionwise Hausdorff?

See [93]. Concerning hopes to answer yes note that if SCH fails (or just $cf(\mu) = \aleph_0$, $pp(\mu) > \mu^+$) then there are examples (see [93, §1]), so we are allowed to assume $2^{\beth_{\omega}} = \beth_{\omega}^+$, etc.

Question 1.30. Let D be an ultrafilter on κ and $\operatorname{Spc}(D) = \{\prod_{i < \kappa} \lambda_i / D, \ \lambda_i \ge 2^{\kappa} \}$

for $i < \kappa$ }. Is $\operatorname{Spc}(D)$ equal to $\{\mu : 2^{\kappa} \le \mu = \mu^{< \operatorname{reg}(D)}\}$? (Where $\operatorname{reg}(D) = \sup\{\theta : \text{ for some } A_i \in D, i < \theta, \text{ for every } \alpha < \kappa, \text{ the number of } i < \theta \text{ such that } \alpha \in A_i \text{ is finite } \}$.)

See on this [96] where some information is gained.

Question 1.31. For which $\lambda \geq \mu$ we can find an almost disjoint family $\mathcal{A} \subseteq [\lambda]^{\aleph_0}$ such that

$$(\forall X \in [\lambda]^{\mu})(\exists A \in \mathcal{A})(A \subseteq^* X)$$
?

At least when $\lambda \geq \mu = \beth_{\omega}$? (See [119], [112]).

Question 1.32. Is it consistent that for some strong limit singular cardinal μ , for no regular $\lambda \in [\mu, 2^{\mu}]$ do we have a c.c.c. Boolean Algebra which is not λ -Knaster?

On related ZFC constructions see [98], [116]; see also §6 here.

Question 1.33. Are all the assumptions in the result of [117] (see below) necessary? In particular, are assumptions (a), (b), (c) below sufficient?

Theorem 1.34 (See [117]). Assume that

- (a) V is our universe of sets, W is another model of ZFC (i.e., a transitive class of V containing all the ordinals),
- (b) κ is a regular cardinal in \mathbf{V} ,
- (c) (\mathbf{W}, \mathbf{V}) has κ -convering (that is, every set of $< \kappa$ ordinals from \mathbf{V} is included in a set of $< \kappa$ ordinals from \mathbf{W}),
- (d) the successor of κ in \mathbf{V} is the same as its succesor in \mathbf{W} , call it κ^+ ,
- (e) (\mathbf{W}, \mathbf{V}) has κ^+ -convering.

Then (\mathbf{W}, \mathbf{V}) has the strong κ -covering (that is, for every structure M with universe an ordinal α and a countable vocabulary, and a set X from \mathbf{V} of cardinality $< \kappa$, there is a set Y from \mathbf{W} of cardinality $< \kappa$ including X which is the universe of an elementary submodel of M).

2. The quest for the test: on the theory of Iterated Forcing for the continuum

On the subject see [164], and recent papers, too, but this section is hampered by some works in progress.

The issue is:

- **Problem 2.1.** (a) Assuming we know something about each iterand \mathbb{Q}_i , what can we say about \mathbb{P}_{α} , where $\langle \mathbb{P}_i, \mathbb{Q}_j : i \leq \alpha, j < \alpha \rangle$ is an iteration (which may be FS (finite support), or CS (countable support) or RSC (revised countable support) and more)?
 - (b) Find more useful ways to iterate (say, new "supports").

So "c.c.c. is preserved by FS iteration", "properness is preserved by CS iteration" can be seen as prototypes. But also many times: "adding no Cohen real over \mathbf{V} ", "adding no dominating real over \mathbf{V} ", etc., and, very natural, "adding no new real".

Note that this is not the same as having forcing axioms, e.g., having (the very important) MA does not discard the interest in FS iterations of c.c.c. forcing. The point is that in many questions you want to add reals for some purpose (which appear as generic sets for some forcing notions), but not another (e.g., a well-ordering of ω of order type ω_1). Also considering an axiom speaking on forcing notions with some property, when considering a candidate, a forcing notion \mathbb{P} , during an iteration we may force that it will not satisfy the property, discard it instead "honestly" forcing with it.

What we get by iterations as above can be phrased as having some axioms, but we have many combinations of adding reals of kinds A, B, and C while preserving properties Pr₁, Pr₂, in other words practically one preservation theorem may be used in many such contexts.

In fact, some of the most intriguing problems are fine distinctions: adding solution to one kind, but not to a close variant, e.g., the old problem:

```
Question 2.2. CON(\mathfrak{p} < \mathfrak{t})? (Note that if \mathfrak{p} < \mathfrak{t} then 2^{\aleph_0} \ge \aleph_3. See 3.7).
```

With FS iteration, all values of the continuum were similar, except \aleph_1 (well, also there is a distinction between regular and singular).

In fact, the advances in proper forcing make us "rich in forcing" for $2^{\aleph_1} = \aleph_2$, making the higher values more mysterious. (So in [164, Ch.VII,VIII] we separate according to the size of the \mathbb{Q}_i 's and whether we add reals, but we concentrate on the length ω_2). So, because we know much more how to force to get $2^{\aleph_0} = \aleph_2$, the independence results on the problems of the interrelation of cardinal invariants of the continuum have, mostly dealt with relationships of two cardinals, as their values are $\{\aleph_1, 2^{\aleph_0}\}$. Thus, having only two possible values $\{\aleph_1, \aleph_2\}$ among any three, two are equal; the Pigeonhole Principle acts against us. As we are rich in our knowledge to force for $2^{\aleph_0} = \aleph_2$, naturally we are quite poor concerning ZFC results. If we try for cardinal invariants c_1, c_2 to prove they consistently are \aleph_1, \aleph_2 , respectively, much of our way exists (quoting existing preservation theorems) and we can look at the peculiarities of those invariants which may be still intractable. We are not poor concerning forcing for $2^{\aleph_0} = \aleph_1$ (and are rich in ZFC). But for $2^{\aleph_0} \geq \aleph_3$ we are totally lost: very poor in both directions. We would like to have iteration theory for length $\geq \omega_3$. I tend to think good test problems will be important in developing such iterations.

In some senses, most suitable is

Problem 2.3. Investigate cardinal invariants of the continuum showing ≥ 3 may have prescribed order.

Of course, the lack of forcing ability does not stop you from proving hopeful ZFC theorems about them, if true. Now I think there are some, but:

Thesis 2.4. They are camouflaged by the independent statements.

[Yes, I really believe there are interesting restrictions.] However, once we prove 90 percent of the problems are independent we will know where to look (as in hindsight occurs in cardinal arithmetic). So cardinal invariants from this perspective are excellent excuses to find iteration theorems. Mainly for $2^{\aleph_0} \geq \aleph_3$, but, of course, there is more to be said on $2^{\aleph_0} = \aleph_1$ (though not for 2.3), and even $2^{\aleph_0} = \aleph_2$.

Without good test problems you are in danger of imitating the king who painted the target after shooting the arrow. Let us consider some additional well known problems:

Question 2.5 (See Just, Mathias, Prikry and Simon [63]). Is there a filter D on ω such that:

- (a) every co-finite subset of ω belongs to D,
- (b) D is a P-filter (i.e., if $A_n \in D$ for $n < \omega$, then for some $A \in D$, $n < \omega \Rightarrow A \subseteq^* A_n$),
- (c) D is not feeble, i.e., if $0 = n_0 < n_1 < \dots$, then for some $A \in D$ for infinitely many $i < \omega$ we have $[n_i, n_{i+1}) \cap A = \emptyset$.

Question 2.6 (See Garcia–Ferreira and Just [39]). Is there an almost disjoint family $\mathcal{A} \subseteq [\omega]^{\aleph_0}$ (i.e., $(\forall A \neq B \in \mathcal{A})[|A \cap B| < \aleph_0]$) of cardinality \mathfrak{b} satisfying the following condition:

```
if A_n \in \mathcal{A} are pairwise distinct and h : \omega \longrightarrow \omega
then for some B \in \mathcal{A} we have (\exists^{\infty} n)(A_n \cap B \nsubseteq h(n))?
```

If not, then $2^{\aleph_0} > \aleph_{\omega}$; on both questions see the discussions after 2.13.

 $\textbf{Question 2.7} \ (\text{See van Mill [179, Problem 4, p.563], Miller [79, Problem 9.1]}).$

```
CON(no P-point and no Q-point)?
```

If so, $2^{\aleph_0} \geq \aleph_3$. [Why? Mathias [74] showed that if \mathfrak{d} (the minimal size of a dominating family) is \aleph_1 , then there is a Q-point. Ketonen [66] showed that $\mathfrak{d} = 2^{\aleph_0}$ implies the existence of P-points.]

Question 2.8. $CON(\omega(\omega+1))$ with box product topology is not paracompact)?

```
If so, 2^{\aleph_0} \geq \aleph_3. See on this Williams [181].
```

Question 2.9 (See Miller [79, Problem 16.3]).

CON(Borel Conjecture and Dual Borel Conjecture)?

(See 3.3).

Question 2.10. CON(cf(cov(meagre)) < additivity(meagre))? (See before 2.14).

Problem 2.11. 1. CON(every function $f: {}^{\omega}2 \longrightarrow {}^{\omega}2$ is continuous when restricted to some non-null set)?

[Here "null" means of Lebesgue measure zero.]

2. Similarly for other natural ideals. This in particular means if \mathbb{Q} is a nicely defined forcing notion (see §5 below, e.g., Souslin c.c.c.), η a \mathbb{Q} -name of a real, $A \subseteq {}^{\omega}2$ is called (\mathbb{Q}, η) -positive if for every countable $\tilde{N} \prec (\mathcal{H}(\chi), \in, <^*_{\chi})$ to

which \mathbb{Q} , $\underline{\eta}$ belong, some $\eta \in A$ is $\underline{\eta}[G]^N$ for some $G \subseteq \mathbb{Q}^N$ generic over N; so the question for such \mathbb{Q} is "CON(every $f : {}^{\omega}2 \longrightarrow {}^{\omega}2$ has a continuous restriction to some (\mathbb{Q}, η) -positive set A)?"

- 3. Is the following consistent: if $A \subseteq {}^{\omega}2$ is non-null, $f: A \longrightarrow {}^{\omega}2$ then for some positive $B \subseteq A$, $f \upharpoonright B$ is continuous. Similarly for general ideals as in part (2).
- 4. If $A \subseteq {}^{\omega}2 \times {}^{\omega}2$ is not equivalent to a Borel set modulo one ideal I_1 (as described in part (2) above), then for some continuous $f: {}^{\omega}2 \longrightarrow {}^{\omega}2$, the set $\{\eta \in {}^{\omega}2 : (\eta, f(\eta)) \in A\}$ is not equivalent to a Borel set modulo another ideal I_0 for suitable pairs (I_0, I_1) .

See Fremlin [36], Ciesielski [19, Theorem 3.13, Problem 5]; [157] shows "yes" for (2) for non meagre, Ciesielski and Shelah [20] prove "yes" for (4) for non meagre, on work in progress see Rosłanowski and Shelah [87, §2]. With Juris Steprāns we have had some discussions on trying to use the oracle cc to the case of non-megre ideal in (2). See 3.8.

NOTE: Mathematicians who are not set theorists generally consider "null" as senior to "meagre", that is as a more important case; set theorists inversely, as set-theoretically Cohen reals are much more manageable than random reals and have generalizations, relatives, etc. Particularly, in FS iterations, we get Cohen reals "for free" (in the limit), which kills our chances for many things and until now we have nothing parallel for random reals (but see [87]).

Judah suggests:

Question 2.12 (V = L). Find a forcing making $\mathfrak{d} = \aleph_3$ but not adding Cohen reals.

I am skeptical whether this is a good test question, as you may make $\mathfrak{d} = \aleph_3 = \mathfrak{b}$ by c.c.c. forcing, then add \aleph_1 random reals $\langle \nu_i : i < \omega_1 \rangle$ by a measure algebra; so over $\mathbf{L}[\langle \nu_i : i < \omega_1 \rangle]$ we have such a forcing. But certainly "not adding Cohen" is important, as many problems are resolved if $\text{cov}(\text{meagre}) = 2^{\aleph_0}$.

There is a basic question for us:

Problem 2.13. Is there an iteration theorem solving all the problems described above or at least for all cases involving large continuum not adding Cohen reals?

I suspect not, and the answers will be ramified.

Let us review some problems. Now, Problems 2.5, 2.6 are for $2^{\aleph_0} > \aleph_{\omega}$, as: in 2.5, if $\operatorname{cf}([\mathfrak{d}]^{\aleph_0}, \subseteq) = \mathfrak{d}$ then there is such a filter (see [63]), and also in 2.6, if $\operatorname{cf}([\mathfrak{b}]^{\aleph_0}, \subseteq) = \mathfrak{b}$ there is a solution (see Just, Mathias, Prikry and Simon [63]).

It may well be that the solution will look like: let μ be a strong limit singular cardinal with $2^{\mu} \geq \mu^{++} > \mu^{+}$ and we use FS iteration of length μ^{++} . This will be great, but probably does not increase our knowledge of iterations. If on the other hand along the way we will add new ω -sequences say to μ (say cf(μ) = \aleph_0) and necessarily we use more complicated iteration, then it will involve better understanding of iterations, probably new ones.

We can ([164, Ch.XIV]) iterate up to "large" κ , and for many $\alpha < \kappa$, α strongly inaccessible, we have \mathbb{Q}_{α} change its cofinality to \aleph_0 . Sounds nice, but no target yet.

* * *

We may note that "FS iterations of c.c.c. forcing notions" is not dead. Concerning 2.3 and 2.10, there are recent indications that FS iteration of c.c.c. still can be exploited even in cases for which for a long time we thought new supports are needed. We can iterate with FS, $\langle \mathbb{P}_i, \mathbb{Q}_i : i < \alpha \rangle$, where \mathbb{Q}_i is (partially random or is Cohen) adding a generic real r_i , \mathbb{Q}_i is Cohen forcing or random forcing over $\mathbf{V}[\langle r_j : j \in A_i \rangle]$, where $A_i \subseteq i$ and each \mathbb{Q}_i is reasonably understood; but we do not require $j \in A_i \Rightarrow A_j \subseteq A_i$ (so called transitive memory). It is not so immediate to understand this sort of iterations, e.g., can the iteration add a dominating real?

It appears that if the A_i 's are sufficiently closed, it will not, see [100], more generally look at [120]. There we prove:

CON(\exists non-null A such that the null ideal restricted to A is \aleph_1 -saturated).

Clearly we should use a measurable cardinal κ , a normal ultrafilter \mathcal{D} on κ in \mathbf{V} and we add κ random reals $\langle r_{\zeta} : \zeta < \kappa \rangle$, but how do we make

$$A \in \mathcal{D}^{\mathbf{V}} \Rightarrow \{r_{\zeta} : \zeta \in \kappa \setminus A\} \text{ is null } ?$$

Cohen forcing does the job, but unfortunately too strongly. (In the case with non meagre, by Komjath [69], there are no problems in this respect). The solution is that we use FS iteration, but first we add 2^{κ} Cohens, that is \mathbb{Q}_i is Cohen forcing for $i < 2^{\kappa}$; only, then we add the (somewhat) random reals:

$$\mathbb{Q}_{\lambda+\zeta}$$
 for $\zeta < \kappa$ is random forcing over $\langle r_{\lambda+\xi} : \xi < \zeta \rangle$, $\langle r_i : i \in A_\zeta \subseteq \lambda \rangle$.

We use $\{r_i : i < \lambda\}$ such that: r_i makes $\{r_{\lambda+\zeta} : \zeta \text{ satisfies } i \in A_\zeta\}$ null for $i < 2^{\aleph_0}$. So we need: for $A \in \mathcal{D}^{\mathbf{V}}$ for some $i < \lambda$ for every $\xi < \kappa, \xi \in \kappa \setminus A \iff i \notin A_\xi$. This works for specially chosen A_ζ 's.

Problem 2.14. 1. Can you make this into a general method?

2. Can you deal with n or even κ kinds of "reals" (getting interesting results)?

What does this mean? This means that we use FS iteration $\langle \mathbb{P}_i, \mathbb{Q}_j : i \leq \delta, j < \delta \rangle$ and $h: \alpha \longrightarrow \beta$, for $\zeta < \beta$, \mathbb{R}_{ζ} is a nep c.c.c. forcing notion in $\tilde{\mathbf{V}}$, (on nep see §5 and more see [107]; e.g., \mathbb{R}_{ζ} is Cohen, random, or as in [86], [88] or whatever), and $\underline{s}_{\zeta} \in {}^{\omega}2$ is a generic real for \mathbb{R}_{ζ} , and \mathbb{Q}_i is $\mathbb{R}_{h(\zeta)}$ as interpreted in $\mathbf{V}[\langle \underline{r}_j : j \in A_i \rangle]$ and \underline{r}_j is $\underline{s}_{h(j)}$ there, and $A_i \subseteq i$. So the idea is that $0 = \delta_0 < \delta_i < \ldots < \delta_n = \delta$ and $j \in [\delta_{\ell}, \delta_{\ell+1}) \Rightarrow h(j) = \ell$.

* * *

In [103] we use \aleph_{ε} -support. This is less than ($< \aleph_1$)-support (i.e., countable support). This looks quite special, but

Problem 2.15. Can we make a general (interesting) theorem?

We can note that long FS iterations not only add Cohen reals, they also add, e.g., \aleph_2 —Cohens, i.e. generics for $\{f: f \text{ a finite function from } \omega_2 \text{ to } \{0,1\} \}$. So we may like to iterate, allowing to add Cohen reals but not \aleph_2 —Cohens in the sense above. This is done in [103], but the family of allowable iterands can be probably widened.

* * *

If we agree that preservation theorems are worthwhile, then after not collapsing \aleph_1 , probably the most natural case is adding no reals. Now, whereas properness seems to me both naturally clear and covers considerable ground for not collapsing \aleph_1 and there are reasonable preservation theorems for "proper +X" for many natural properties X (e.g., adding no dominating reals, see [164, Ch.VI, $\S1$, $\S2$, $\S3$], [164, Ch.XVIII, §2]), the situation with NNR (no new real) is inherently more complicated. In the early seventies when I heard on Jensen's CON(GCH+SH), I thought it would be easy to derive an axiom; some years later this materialized as Abraham, Devlin and Shelah [5], but reality is not as nice as dreams. One obstacle is the weak diamond, see Devlin and Shelah [25], more in [132, Ch.XIV,§1], [164, AP,§1], [105]. For a time the iteration theorem in [132, Ch.V,§5,§7,Ch.VIII,§4] seemed satisfactory to me. [There we use two demands. The first was D-completeness (this is a "medicine" against the weak diamond, and $\mathbb D$ is a completeness system, \aleph_1 -complete in [132, Ch V, §5,7], that is any countably many demands are compatible, and 2complete in [132, Ch VIII, §4], that is any two demands are compatible). The second demand was α -properness for each countable ordinal α (or relativized version, see [132, Ch VIII, §4], [164, Ch VIII, §4]).] But [138, §1] (better [164, Ch.XVIII, §1]) gives on the one hand very nice and easy forcing notions not adding reals (running away from club guessing sequences) which are not covered as they fail ($<\omega_1$)properness and on the other hand, shows by a not so nice example that generally you cannot just omit the $(<\omega_1)$ -properness demand and promise an iteration theorem covering them. The problem concerning that forcing was resolved (promised in [138], carried out in a different way in lectures in MSRI '89 = [164, Ch.XVIII,§2]), but resulted in a dichotomy: we can get by forcing CON(ZFC+CH+SH) and we can get by forcing CON(ZFC + CH + no club guessing), but can we have both? More generally, can we have two other such contradictory statements (more generally for such results see Shelah and Zapletal [171]).

Question 2.16. Can we have two statements of the form²

$$(\forall x \in \mathcal{H}(\aleph_2))(\exists y \in \mathcal{H}(\aleph_2))\varphi$$

each consistent with

```
CH + [Axiom(\mathbb{Q} is (<\omega_1)-proper and \mathbb{D}-complete for some simple 2-completeness system \mathbb{D})]
```

but not simultaneously?

(We may change the axiom used, we may speak directly on the iteration; we may deal with CS and proper or with RCS and semi-proper, etc.)

Note: possible failure of iteration does not prove a ZFC consequence, we may have freedom in the iteration only in some stages (like c.c.c. productive under MA).

This leaves me in bad shape: the iteration theorems seem not good enough, but the test problem (of getting both) does not seem so good. Now, [113] deals with NNR solving the specific dichotomy (and really satisfies the [138] promise circumvented in [164, Ch.XVIII, §2]) but left 2.15 open.

Eisworth suggested to me (motived by Abraham and Todorčević [4])

Question 2.17. Is the following consistent with ZFC+CH

 $^{^{2}\}mathcal{H}(\lambda)$ is the family of sets with transitive closure of cardinality $<\lambda$

(*) if $A_{\alpha} \in [\omega_1]^{\aleph_0}$ and $\alpha < \beta \Rightarrow A_{\alpha} \subseteq A_{\beta}$ mod finite, and for every stationary $S \subseteq \omega_1$ the set $\bigcup \{ [A_{\alpha}]^{<\aleph_0} : \alpha \in S \}$ contains $[E]^{<\aleph_0}$ for some club E of ω_1 , then for some club C of ω_1 we have

$$(\forall \alpha < \omega_1)(\exists \beta < \omega_1)(C \cap \alpha \subseteq A_\beta).$$

For long, an exciting problem for me has been

- **Problem 2.18.** 1. Can we find a consequence of ZFC + CH which "stands behind" the "club objection to NNR", e.g. it implies the failure of CH + $Axiom(\mathbb{Q} \text{ proper } \mathbb{D}\text{--complete}$ for some single 2-completeness system)?
 - 2. Similarly for other limitations on iteration theorems?

Question 2.19. Is "CH + \mathcal{D}_{ω_1} is \aleph_2 -saturated" consistent, where \mathcal{D}_{ω_1} is the club filter on ω_1 ?

Recall that a filter D on a set A is λ -saturated if there are no $A_i \in D^+$ for $i < \lambda$ such that $i < j \implies A_i \cap A_j = \emptyset \mod D$.

See [164, Ch.XVI]. Woodin proved that if there is a measurable cardinal then no. So we may look at L[A], $A \subseteq \kappa$ codes $\mathcal{H}(\chi)$, κ large and try to collapse it to ω_2 .

Note that by [105], if CH + \mathcal{D}_{ω_1} is \aleph_2 -saturated, then essentially we have the weak diamond for three colours (or any finite number).

* * *

Baumgartner [11] asked

Question 2.20. Is it consistent that $2^{\aleph_0} > \aleph_2$ and any two \aleph_2 -dense subsets of \mathbb{R} of cardinality \aleph_2 (that is, any interval has \aleph_2 points) are isomorphic (as linear orders).

I think it is more reasonable to try

Question 2.21. Is it consistent that: $2^{\aleph_0} > \lambda \geq \aleph_2$ and there are no two far subsets $A \in [\mathbb{R}]^{\lambda}$, where

Definition 2.22. The (linear orders) I, J are θ -far if there is no linear order of cardinality θ embedded into both. If θ is omitted, we mean $\min\{|I|, |J|\}$.

On OCA' (i.e., OCA_{\aleph_1,ω}, see the definition below) see Abraham, Rubin and Shelah [2], continued for OCA" in Todorčević [178], Veličković [180]; on a parallel for subsets of the plain which follows from MA, see Steprāns and Watson [176].

Question 2.23. 1. Is OCA'_{\aleph_2} consistent? Is OCA''_{\aleph_2} consistent?

- 2. The parallel problems for κ^2 and λ , even for $\lambda = \kappa^+$, $\kappa > \aleph_0$, where
- **Definition 2.24.** 1. OCA'_{λ,κ} means $\lambda \leq 2^{\kappa}$ and: for any $A \in [\kappa^2]^{\lambda}$ and an open symmetric set $\mathcal{U} \subseteq \kappa^2 \times \kappa^2$ there is $B \subseteq A$ of cardinality λ such that $\{(a,b): a \neq b \text{ are from } B\}$ is included in \mathcal{U} or is disjoint to \mathcal{U} (we use the space κ^2 for simplicity).
 - 2. OCA_{λ,κ} is defined similarly only we have $B_i \subseteq A$ for $i < \kappa$, $A = \bigcup_{i < \kappa} B_i$, each B_i as in part (1).
 - 3. If we omit λ we mean $\lambda = \kappa^+$, if in addition we omit κ , we mean $\kappa = \aleph_0$.

3. Case studies for iterated forcing for the reals

The following was suggested during the lecture on $\S 2$ by Juhász who was in the audience:

Question 3.1. Does CH imply that there is an S-space of cardinality \aleph_2 , where S-space is defined as being regular, hereditarily separable, not Lindelöf?

Eisworth prefers the variant:

Does CH imply the existence of a locally compact S-space?

This problem looks important, but it is not clear to me if it is relevant to developing iteration theorems, though an existence proof may be related to the weak diamond, consistency to NNR iterations.

The same goes for the well known:

Question 3.2. $CON(\mathfrak{d} < \mathfrak{a})$?

This definitely seems not to be connected to the iteration problem. It seems to me that a good test problem for our purpose in §2 should have one step clear but the iteration problematic, whereas for those two problems the situation is the inverse.

Note: by existing iteration theorems to get the consistency of $\mathfrak{d} < \mathfrak{a} + 2^{\aleph_0} = \aleph_2$ it is enough to show

(*) for any MAD family $\{A_i : i < i^*\} \subseteq [\omega]^{\aleph_0}$, there is an ${}^{\omega}\omega$ -bounding proper forcing notion $\mathbb Q$ of cardinality \aleph_1 adding $A \in [\omega]^{\aleph_0}$ almost disjoint to each A_i .

You are allowed to assume CH (start with $\mathbf{V} \models 2^{\aleph_0} = \aleph_1 + 2^{\aleph_1} = \aleph_2$ and use CS iteration of such forcing notion); even \diamondsuit_{\aleph_1} (if $\mathbf{V} \models \diamondsuit_{\aleph_1}$). We can weaken $|\mathbb{Q}| = \aleph_1$ to " \mathbb{Q} satisfies \aleph_2 -pic" (this is a strong form of \aleph_2 -c.c. good for iterating proper forcing, see [164, Ch.VIII,2.1,p.409]). If you agree to use large cardinals, it is okay to assume in (*) that an appropriate forcing axiom holds and not restrict $|\mathbb{Q}|$, and as we can first collapse 2^{\aleph_0} to \aleph_1 , we can get \diamondsuit_{\aleph_1} for "free". I idly thought to use free forcing for the problem, ([164, Ch.IX]), but no illumination resulted.

We can try in another way: start with a universe with a forcing axiom (say MA) and force by some \mathbb{P} , which makes $\mathfrak{d} = \aleph_1$, but \mathbb{P} is understood well enough and we can show that \mathfrak{a} is still large (just as adding a Cohen real to a model of MA preserves some consequences of MA (see Roitman [81], Judah and Shelah [58]]). So clearly FS iteration will not do.

I think that a more interesting way is to consider, assuming CH,

$$K_{\omega_{1}} = \{(\bar{\mathbb{P}}, \bar{\underline{r}}) : \quad \bar{\mathbb{P}} = \langle \mathbb{P}_{i} : i < \omega_{1} \rangle \text{ is } \Leftrightarrow \text{-increasing, } |\mathbb{P}_{i}| \leq \aleph_{1}, \\ \bar{\underline{r}} = \langle \underline{r}_{i} : i < \omega_{1} \rangle, \ \underline{r}_{i} \text{ is a } \mathbb{P}_{i+1}\text{-name,} \\ \Vdash_{\mathbb{P}_{i+1}} \text{"} \ \underline{r}_{i} \in {}^{\omega}\omega \text{ dominates } ({}^{\omega}\omega)^{\mathbf{V}^{\mathbb{P}_{i}}}] \text{"} \}$$

ordered naturally, and for a generic enough ω_2 -limit $\langle (\bar{\mathbb{P}}^{\zeta}, \bar{r}^{\zeta}) : \zeta < \omega_2 \rangle$ we may use $\bigcup_{\substack{i < \omega_1, \\ \zeta < \omega_2}} \mathbb{P}_i^{\zeta}$. Another way to try is the non-Cohen Oracle [108]. The difference is

small. Also the " $\omega_2 + \omega_1$ -length mix finite/countable pure support iteration" seems similar.

I have just heard about $\mathrm{CON}(\mathfrak{u} < \mathfrak{a})$ being an old problem, clearly related to $\mathrm{CON}(\mathfrak{d} < \mathfrak{a})$. I do not see much difference at present.

Another direction is to develop the historic \aleph_{ε} -support iteration from [103].

Discussion 3.3. Concerning 2.9, I had not really considered it (except when Judah spoke to me about it) but just before the lecture, Bartoszyński reminded me of it (see [6]). Now, "the" proof of CON(Borel conjecture) is by CS iteration of Laver forcing (see Laver [73]), whereas the consistency proof of the dual is adding many Cohen reals (see Carlson [18]). So in a (hopeful) iteration proving consistency we have two kinds of assignments. We are given, say in stage α , in $\mathbf{V}^{\mathbb{P}_{\alpha}}$ a set $A = \{\eta_i : i < \omega_1\} \subseteq {}^{\omega}2$, and we should make it not of strong measure zero, so we should add an increasing sequence $\bar{n} = \langle n_\ell : \ell < \omega \rangle$ of natural numbers such that for no $\bar{\nu} = \langle \nu_\ell : \ell < \omega \rangle \in \prod_{\ell < \omega} {}^{(n_\ell)}2$ do we have $(\forall i < \omega_1)(\exists^\infty \ell)(\nu_\ell \lhd \eta_i)$. Now, even if we define \mathbb{Q}_{α} to add such \bar{n} , we have to preserve it later, so it is easier to preserve, for some family $F \subseteq \prod_{\ell < \omega} 2^{n_\ell}$, the demand

$$(\forall f \in F) \neg (\exists \langle \nu_{\ell,k} : \ell < \omega, k < f(\ell) \rangle) (\forall i < \omega_1) (\exists^{\infty} \ell) (\exists k < f(\ell)) (\nu_{\ell,k} \neg \eta_i).$$

The second kind of assignment which we have in stage α is the following. In $\mathbf{V}^{\mathbb{P}_{\alpha}}$, we are given $A = \{\eta_i : i < \omega_1\} \subseteq {}^{\omega}2$ and we should make it non-strongly meagre, so we should add, by \mathbb{Q}_{α} , a subtree $T_{\alpha} \subseteq {}^{\omega}>2$ (i.e., $\langle \rangle \in T_{\alpha}$, $\eta \in T_{\alpha} \& \nu \lhd \eta \Rightarrow \nu \in T_{\alpha}$, $\eta \in T_{\alpha} \Rightarrow (\exists \ell < 2)(\eta \widehat{\ } \langle \ell \rangle \in T_{\alpha})$) of positive measure (i.e., $0 < \inf\{|T_{\alpha} \cap {}^{n}2|/2^{n} : n < \omega\}$) such that $(\forall \eta \in {}^{\omega}2)(\exists i < \omega_1)[\eta \oplus \eta_i \notin \bigcup_{r} (T_{\alpha})^{[r]}]$, where

$$(T_{\alpha})^{[n]} = \{ \nu : \text{ for some } \rho \in T_{\alpha} \text{ we have } \ell g(\nu) = \ell g(\rho) \text{ and } (\forall \ell) (n \leq \ell < \ell g(\nu) \Rightarrow \rho(\ell) = \nu(\ell)) \}.$$

Again we have to preserve this.

A way to deal with such preservation problems is to generalize "oracle c.c.c." (see [164, Ch.IV]) replacing Cohen by other things. To explain this, it seems reasonable to look at the "oracle for random" (or even sequence of c.c.c. Souslin forcing, from [108]). This evolves to: for iterations of length $\leq \omega_2$ of forcing notions of cardinality \aleph_1 , prove that we can preserve the following condition on $\mathbb{P} = \mathbb{P}_{\alpha}$ for some $\langle M_{\delta}, M_{\delta}^+, r_{\delta} : \delta \in S \rangle$, $S \subseteq \omega_1$ stationary such that $\langle M_{\delta} : \delta \in S \rangle$ is an oracle, i.e., a \diamond^* -sequence and $M_{\delta} \models \delta = \omega_1$, $M_{\delta} \models \mathrm{ZFC}_*^-$, $M_{\delta}^+ \models$ " $\mathrm{ZFC}_*^- + M_{\delta}$ is countable" and r_{δ} is random over M_{δ}^+ . Now without loss of generality, $\mathbb{P} \subseteq \omega_1$ and

```
 \{\delta \in S: \ \mathbb{P} \upharpoonright \delta \in M_{\delta}, \text{ and for every } p \in \mathbb{P} \cap \delta, \text{ for some } q \text{ we have } \\ p \leq q \in \mathbb{P} \text{ and } q \Vdash \text{``} r_{\delta} \text{ is random over } M_{\delta}^{+}[\mathcal{G}_{\mathbb{P}} \cap \delta] \text{'`} \} \ \in \mathcal{D}_{\omega_{1}} \upharpoonright S
```

(so this is like the oracle c.c.c. ([164, Ch.IV]), but the support is not countable so on other stationary $S_1 \subseteq \omega_1 \setminus S$ we may have different behaviour). Of course, we use "small" S so that we have "space" for more demands, see [108]. But trying to explain it (to Rosłanowski) it seemed the proof is too simple, so we can go back to good old CS and just preserving an appropriate property, a watered-down relative in the nep family ([107]).

We mainly try to combine the two iterations (of Cohen and of Laver forcing notions):

Definition 3.4. A forcing notion \mathbb{Q} is 1–e.l.c. if the following condition is satisfied:

```
whenever \chi is large enough, M_0 \prec M_1 \prec (\mathcal{H}(\chi), \in), \mathbb{Q} \in M_0, M_0 \in M_1 and M_0, M_1 are countable and p \in \mathbb{Q} \cap M_0, then for some condition q \in \mathbb{Q} stronger than p we have
```

 $q \Vdash$ "for every $\mathcal{I} \in M_1$ such that $\mathcal{I} \cap M_0$ is predense in \mathbb{Q}^{M_0} we have $\mathcal{G}_{\mathbb{Q}} \cap \mathcal{I} \neq \emptyset$ ".

(Note that $q \Vdash$ " $M_0[G_{\mathbb{Q}} \cap M_0]$ is a generic extension of M_0 for a forcing notion which M_1 thinks is countable".)

Note: e.l.c. stands for elementary locally Cohen. This is, of course, close to Cohen, or more accurately is another way to present strongly proper. But we also seem to need Laver forcing (or a close relative of it), but it is far from being strongly proper. Still it satisfies the parallel if we demand " $\mathcal{I} \subseteq Q^{M_1}$ is predense under pure extensions", i.e., with the same trunk. This approach seems to me promising but it is not clear what it delivers.

We may consider a more general definition (and natural preservation):

Definition 3.5. Let Pr be a property. A forcing notion \mathbb{Q} with generic $X \subseteq \alpha_{\mathbb{Q}}$ (i.e. $V[\mathcal{G}_{\mathbb{Q}}] = V[X[\mathcal{G}_{\mathbb{Q}}]]$, $\alpha_{\mathbb{Q}}$ an ordinal) is called e.l.—Pr forcing if:

for χ large enough, if $\mathbb{Q}, \tilde{\chi} \in M_1 \prec M_2 \prec (\mathcal{H}(\chi), \in), M_1, M_2$ countable, $M_1 \in M_2, p \in \mathbb{Q} \cap M_1$, then we can find q, \mathbb{Q}' such that

- (a) $p \leq q \in \mathbb{Q}$,
- (b) $\mathbb{Q}' \in M_2$ is a forcing notion with $X' \subseteq \alpha_{\mathbb{Q}}$ generic,
- (c) $M_2 \models \Pr(\mathbb{Q}', M_1, p),$
- (d) $q \Vdash X \upharpoonright M_2$ is a \mathbb{Q}' -generic over p and for some set $G' \subseteq (\mathbb{Q}')^{M_2}$, generic over M_2 we have $X \upharpoonright M_2 = X'[G']$.

This seems to me interesting but though Laver forcing satisfies some relatives of those properties it does not seem to be enough.

Note: this definition tells us that generically for many countable models $M \prec (\mathcal{H}(\chi), \in)$, we have some $q \in G_{\mathbb{Q}}$ which is almost (M, \mathbb{Q}) -generic, but not quite. The "almost" is because this holds for another forcing \mathbb{Q}' . So when the whole universe is extended generically for \mathbb{Q} , M "fakes" and is instead extended generically for \mathbb{Q}' . So for preservation in iteration it is not natural to demand $M_2 \prec (\mathcal{H}(\chi), \in)$, but rather to proceed as in [107], this will be n.e.l.-Pr.

We may wonder (considering 2.9) whether we can replace Laver forcing in the proof of the consistency of the Borel conjecture, by a forcing notion not adding a dominating real. So a related question to 2.9 is

Question 3.6.

$$CON(\mathfrak{b} = \aleph_1 + Borel Conjecture)$$
?

It is most natural to iterate, one basic step will be \mathbb{Q} , adding an increasing sequence $\langle p_i : i < \omega \rangle$ such that on the one hand:

(a) no old non-dominated family $\subseteq {}^{\omega}\omega$ is dominated (or at least some particular old family remains undominated),

while on the other hand

(b) for any uncountable $A \subseteq {}^{\omega}2$, from **V**, we have:

$$\Vdash_{\mathbb{Q}}$$
 " for no $\eta_i \in {}^{n_i}2$, $(i < \omega)$ do we have $(\forall \nu \in A)(\exists^{\infty}i)(\eta_i \triangleleft \nu)$ ",

or at least

(b)' like (b) for one A given by bookkeeping.

(To preserve we need to strengthen the statement, replacing $\langle \tilde{\eta}_i : i < \omega \rangle$ by a thin enough tree.) The $\tilde{\eta}_i$ should "grow" fast enough, so naturally we think of forcing notions as in Rosłanowski and Shelah [85], [87], which proved easily checked

sufficient conditions for what we desire (so in the "neighborhood" of Blass and Shelah [14]). But what should be the norm?

* * *

Discussion 3.7. Concerning $\mathfrak{p} < \mathfrak{t}$, I have made quite a few failed tries. Some try to use long iterations ($\geq \aleph_{\omega+1}$) or a new support. But also I thought that Blass and Shelah [15] would be a reasonable starting point, the point is how to extend \aleph_1 -generated filters to a good enough P-point.

That is, trying to force $\mathfrak{p} = \aleph_2$, $\mathfrak{t} = \aleph_3 = \mathfrak{c}$ start, say, with $\mathbf{V} = \mathbf{L}$ and use a FS iteration $\langle \mathbb{P}_i, \mathbb{Q}_j : i \leq \omega_3, j < \omega_3 \rangle$, where \mathbb{Q}_i is a Cohen forcing adding $r_i \in {}^{\omega}2$ for some i's, and \mathbb{Q}_i is shooting an ω -sequence through a P-point filter (or ultrafilter) on ω for some $i \geq \omega_2$. The point is that when we have to find a \leq^* -lower bound to the downward directed $\mathcal{A} \in [\mathcal{P}(\omega)]^{\aleph_1}$, we extend it to a P-point, possibly also for the ω_2 -towers we have to do this. It is natural to try to preserve, for $\alpha \in [\omega_2, \omega_3)$, the statement:

```
in \mathbf{V}^{\mathbb{P}_{\alpha}}, noting that \mathcal{H}(\aleph_1) has cardinality \aleph_2, if \mathcal{H}(\aleph_1) = \bigcup_{\alpha < \omega_2} M_{\alpha}, M_{\alpha} increasing continuous, ||M_{\alpha}|| < \aleph_2, then the following set is = \emptyset \mod \mathcal{D}_{\omega_2} + S_1^2:
```

 $\left\{\delta: \text{ if some } a\in M_\delta\cap[\omega]^{\aleph_0} \text{ is almost included in } r_i^{-1}(\{1\}) \text{ for many } i<\delta, \\ \text{ then } a \text{ is almost disjoint to } r_\delta^{-1}(\{1\}) \right\}.$

* * *

Discussion 3.8. Concerning 2.11 consider the problem "every $f: \mathbb{R} \longrightarrow \mathbb{R}$ is continuous on a non-null set".

We can try to use a forcing notion which looks locally random (like the forcing for "non meager set" of [157] looked locally like the Cohen forcing notion) or a mixture of random and quite bounding ones. Such forcing notions are considered in [87], do they help for "every function $f: {}^{\omega}2 \longrightarrow {}^{\omega}2$ is continuous on a non-null set"?

How can we try to prove the consistency of "for every non-meagre $A \subseteq {}^{\omega}2$ and $f:{}^{\omega}2 \longrightarrow {}^{\omega}2$ for some non-meagre $B \subseteq A$, $f \upharpoonright B$ is continuous"? We may use CS or even FS iteration of length ω_2 , (with $\mathbf{V} \models \mathrm{GCH} + \diamondsuit_{\{\delta < \aleph_2 : \mathrm{cf}(\delta) = \aleph_1\}} + S_{\alpha} \subseteq \omega_1$ $(\alpha < \omega_2)$ increasing $\mathrm{mod} \ \mathcal{D}_{\omega_1}$).

In Stage α we have $\bar{r}^{\alpha} = \langle r_i^{\alpha} : i \in S_{\alpha} \rangle$ such that $\bar{r}^{\alpha} \in N \prec (\mathcal{H}(\chi), \in) \Rightarrow r_{N \cap \omega_1}^{\alpha}$ is forced to be Cohen over N and $\beta < \alpha \Rightarrow \{i \in S_{\beta} : r_i^{\beta} \neq r_i^{\alpha}\}$ is not stationary.

Sometimes in Stage α , bookkeeping gives us a \mathbb{P}_{α} -name A_{α} of a non-meagre subset of $^{\omega}2$ and we choose $\bar{r}^{\alpha+1}$ such that $\bar{r}^{\alpha+1} \upharpoonright S_{\alpha} = \bar{r}^{\alpha}$ and $\{r_i^{\alpha+1} : i \in S_{\alpha+1} \setminus S_{\alpha}\}$ is a non-meagre subset of A_{α} .

Sometimes in Stage α , bookkeeping gives us a stationary subset S^{α} of S_{α} (from \mathbf{V}) and a \mathbb{P}_{α} -name f_{α} of a function from $\{r_{i}^{\alpha}:i\in S^{\alpha}\}$ to ${}^{\omega}2$ and we try to choose $\bar{r}^{\alpha+1}$ such that: $\bar{r}^{\alpha+1}\upharpoonright S_{\alpha}=\bar{r}^{\alpha},\ \{r_{i}^{\alpha+1}:i\in S_{\alpha+1}\setminus S_{\alpha}\}\subseteq \{r_{i}^{\alpha}:i\in S^{\alpha}\}$ and $f_{\alpha}\upharpoonright \{r_{i}^{\alpha+1}:i\in S_{\alpha+1}\setminus S_{\alpha}\}$ is continuous. So the aim is that in $\mathbf{V}^{\mathbb{P}_{\omega_{2}}}$, every non-meagre $A\subseteq {}^{\omega}2$ contains a subset of the form $\{r_{i}^{\alpha}:i\in S_{\alpha+1}\setminus S_{\alpha}\}$ and

 $S' \subseteq S_{\alpha} \& \alpha < \omega_1 \& S'$ stationary $\Rightarrow \{r_i^{\alpha} : i \in S'\}$ is non-meagre.

We may try to define iterations for forcing related to measure: we can use CS, or try to imitate the measure algebra, there are various ways to interpret it. If each \mathbb{Q}_i is as in [85], so each condition has possible pair: a norm $\in \omega$ and real $r \in (0,1)$ and using those we define what is a condition in the iteration. See [87].

More on preservation (for not necessarily c.c.c. ones) commutativity, associativity, generic sets and countable for pure/finite for a pure support iterations see [108]. Remember that an automorphism F of $\mathcal{P}(\omega)$ /finite is called trivial if it is induced by a permutation f of \mathbb{Z} (the integers, where $\omega = \{n \in \mathbb{Z} : n \geq 0\}$) such that $\{n \in \mathbb{Z} : n < 0 \iff f(n) \geq 0\}$ is finite.

Question 3.9. What can AUT($\mathcal{P}(\omega)$ /finite) be? Is it consistent that AUT($\mathcal{P}(\omega)$ /finite) is not the group of trivial automorphisms of $\mathcal{P}(\omega)$ /finite, but is of cardinality continuum (or even is generated by adding one automorphism to the subgroup of the trivial auto-

morphisms of $\mathcal{P}(\omega)/\text{finite}$)?

It is reasonable try to combine Shelah and Steprāns [169] and the later part of the proof in [164, Ch.IV,§6] (from being "locally trivial" to being trivial).

* * *

Discussion 3.10. Next we deal with the variants of OCA and isomorphisms or farness of sets of \aleph_2 reals, i.e. 2.21, 2.24.

Concerning 2.21 (on "no far $A, B \in [^{\omega}2]^{\lambda}$ "), assume that for $\ell = 1, 2$ we have $A_{\ell} = \{\eta_{\alpha}^{\ell} : \alpha < \lambda\} \subseteq {}^{\omega}2$ with no repetitions. Considering Baumgartner [10] and Abraham, Rubin and Shelah [2], it is natural to try to find $\bar{f} = \langle f_{\alpha} : \alpha < \lambda \rangle$ such that:

- (a) f_{α} is a partial, countable, non-empty function from ω_2 to ω_2 , (for the present aim, $\text{Dom}(f_{\alpha})$ a singleton in (a) and $\gamma = 1$ in (b) are fine, so we assume so),
- (b) for some $\gamma = \gamma^* \leq \omega$, the sequence

$$\langle \bigcup_{n < \gamma} (\text{Dom}(f_{\gamma \alpha + n}) \cup \text{Rang}(f_{\gamma \alpha + n})) : \alpha < \lambda \rangle$$

is a sequence of pairwise disjoint sets.

We let $\hat{f}_{\alpha} = \{(\eta_i^1, \eta_{f_{\alpha}(i)}^2) : i \in \text{Dom}(f_{\alpha})\}$, so $\hat{f}_{\alpha}(\eta)$ is well defined iff $i \in \text{Dom}(f_{\alpha})$, $\eta = \eta_i^1$, and then $\hat{f}_{\alpha}(\eta_i^1) = \eta_{f_{\alpha}(i)}^2$.

It is natural to try the following forcing notion

 $\mathbb{Q}_{\bar{f}} = \big\{g: \quad g \text{ is a finite 1-to-1 order preserving function from } A_1 \text{ to } A_2,$ which has the form $\bigcup_{\ell=1}^n f_{\alpha_\ell}, \ \alpha_\ell < \omega_2, \text{ and such that}$ $\ell_1 \neq \ell_2 \quad \Rightarrow \quad \neg (\exists \alpha) (\{\alpha_{\ell_1}, \alpha_{\ell_2}\} \subseteq [\gamma^*\alpha, \gamma^*\alpha + \gamma^*]) \big\}.$

The order is the inclusion.

It is enough to have " $\mathbb{Q}_{\bar{f}}$ satisfies the c.c.c." (for some \bar{f} as above): clearly $\bigcup \{g: g \in G_{\mathbb{Q}_{\bar{f}}}\}$ is an order preserving function from some $A'_1 \subseteq A_1$ into A_2 ; but does it have cardinality λ ? Essentially yes, as e.g. if $\mathrm{cf}(\lambda) > \aleph_0$ then some $p \in \mathbb{Q}_{\bar{f}}$ forces this. For this it is enough to have: if $u \in [\lambda]^{\aleph_1}$ then $\mathbb{Q}_{\bar{f} \upharpoonright u}$ satisfies the c.c.c. and $\lambda = \aleph_2$, it is enough to check for $u = \alpha \in [\omega_1, \omega_2)$. So as in [2], it is enough that

(c) if $n < \omega$ and $C \subseteq {}^{2n}({}^{\omega}2)$ is closed and

$$(\eta_0, \eta_1, \dots, \eta_{2n-1}) \in C \quad \Rightarrow \quad (\forall \ell < m < n)(\eta_{2\ell} <_{\ell ex} \eta_{2m} \Leftrightarrow \eta_{2\ell+1} <_{\ell ex} \eta_{2m+1})$$
and there are $p_{\zeta} = \{(\eta_{\zeta,\ell}, \nu_{\zeta,\ell}) : \ell \leq n\} \in \mathbb{Q}_{\bar{f}} \text{ for } \zeta < \omega_1, \ \ell < n \text{ with}$

$$(\zeta_1, \ell_1) \neq (\zeta_2, \ell_2) \quad \Rightarrow \quad \eta_{\zeta_1, \ell_1} \neq \eta_{\zeta_2, \ell_2} \text{ and } \nu_{\zeta_1, \ell_1} \neq \nu_{\zeta_2, \ell_2},$$

hence

$$(\forall \zeta < \omega_1)(\forall \ell < m < n)(\eta_{\zeta,\ell} <_{\ell ex} \eta_{\zeta,m} \Leftrightarrow \nu_{\zeta,\ell} <_{\ell ex} \nu_{\zeta,m})$$

(if $\gamma^* = 1$, and each $\text{Dom}(f_{\alpha})$ is a singleton, $p_{\zeta} \in \mathbb{Q}_{\bar{f}}$ means, in addition only $\{(\eta_{\zeta,\ell}, \nu_{\zeta,\ell})\} = f_{\alpha_{\zeta,\ell}}$ for some $\alpha_{\zeta,\ell}$), $(p_{\alpha}'', p_{\alpha}'', \dots, p_{\alpha-1}'') \in C$ such that $p_{\alpha}' \neq p_{\alpha}''$

then there are $(\eta_0', \eta_1', \dots, \eta_{2n-1}'), (\eta_0'', \eta_1'', \dots, \eta_{2n-1}'') \in C$ such that $\eta_\ell' \neq \eta_\ell''$ and $\eta_{2\ell}' <_{\ell ex} \eta_{2\ell}'' \Leftrightarrow \eta_{2\ell+1}' <_{\ell ex} \eta_{2\ell}''$.

Any counterexample to clause (c) induces a continuous partial function for which we get dependencies. If $\mathbf{V} \models \mathrm{CH}$, and \mathbb{P} is adding Cohen reals, then in $\mathbf{V}^{\mathbb{P}}$ this holds, so it is natural to try to retain during the iteration similarly to this case. For 2.20 we can use similar, but somewhat more involved forcing notion as in [2].

There are similar considerations on OCA. We may consider trying to get negative ZFC results, so Kojman and Shelah [67] seems to me a reasonable starting point (of course, the problem there is different).

* * *

Baumgartner [12] defines

Definition 3.11. 1. For a (non-principal) ultrafilter D on ω , and a countable ordinal δ , we say D is a δ -ultrafilter if:

- (*) for every function f from ω to ω_1 , for some $A \in D$ we have $\operatorname{otp}(f(A)) < \delta$.
- 2. We say that D is a weak δ -ultrafilter if:
- (*) for every function f from ω into δ , for some $A \in D$ we have $\operatorname{otp}(f(A)) < \delta$.
- 3. We say that D is a NWD if for every function f from ω to \mathbb{R} , for some $A \in D$, f(A) is a nowhere dense subset of \mathbb{R} .
- 4. For an ideal I, D is an I-filter if for any $f : \text{Dom}(D) \longrightarrow \text{Dom}(I)$ and $A \in D^+$ there is $B \subseteq A, B \in D^+$ such that $f(B) \in I$.

Then he asked whether such ultrafilters exist (if CH yes, so):

Question 3.12. Prove the consistency of "there is no δ -ultrafilter on ω ".

It seemed the solution of the related "CON(there is no NWD-ultrafilter)" in [165] should give this, but it did not, and it is not clear if the question 3.12 is harder (the NWD eluded me several times, but when solved, the solution seems a straightforward generalization of CON(no P-point), which was also a priori the natural starting point).

A nice feature of P-points is that "D generates a P-point ultrafilter on ω " is preserved in limit for CS iterations, so P-points generated by $\aleph_1 < 2^{\aleph_0}$ sets are gotten naturally. Are they the only ones? Of course, by c.c.c. forcing P you may have ultrafilters on ω generated by $< 2^{\aleph_0}$ sets, and forcing by a subforcing $\mathbb{Q} \triangleleft \mathbb{P}$, in $\mathbf{V}^{\mathbb{Q}}$ we get an ultrafilter preserved (see more in Brendle and Shelah [17]); but we have no understanding, though the suggestion in 3.2 may help.

Question 3.13. Are ultrafilters D as defined below in 3.21 preserved in limit stages of CS iterations? This means that:

if $\bar{\mathbb{Q}} = \langle \mathbb{P}_i, \mathbb{Q}_j : i \leq \delta, j < \delta \rangle$ is a CS iteration of proper forcing notions and \tilde{D} is an ultrafilter as above in \mathbf{V} , then $(\forall i < \delta)(\boxtimes_i) \Rightarrow (\boxtimes_{\delta})$, where

 $(\boxtimes_{\beta}) \Vdash_{\mathbb{P}_{\beta}}$ "in $\mathbf{V}^{\mathbb{P}_{\beta}}$ the filter on ω that the family D generates in $\mathbf{V}^{\mathbb{P}_{\beta}}$ is an ultrafilter".

Less nice, but still good, is to prove the preservation of "D generates an ultrafilter +Pr" (where Pr is some additional property like: $({}^{\omega}\omega)^{\mathbf{V}}$ is dominating).

The following 3.14 - 3.22 suggest an approach to question 3.13.

Definition 3.14. 1. Let

 $\mathcal{T} = \{ t : t \subseteq {}^{\omega}{}^{>}\omega, \ t \text{ has a } \lhd -\text{minimal element rt}(t), \\ t \text{ is closed under initial segments of length } \ge \ell g(\text{rt}(t)), \\ \text{for } \eta \in t, \ \text{Suc}_t(\eta) = \{ \eta \widehat{\ \ } \langle \ell \rangle : \eta \widehat{\ \ } \langle \ell \rangle \in t \} \text{ is empty or infinite} \}.$

For $t \in \mathcal{T}$ let $h_t : t \longrightarrow \omega_1 \cup \{\infty\}$ be defined by

$$h_t(\eta) = \bigcup \{h_t(\nu) + 1 : \nu \in \operatorname{Suc}_t(\eta)\}.$$

(So $h_t(\eta) = \infty$ iff there is an ω -branch through η).

We say that t is standard if

$$\eta \in t \& \beta < h_t(\eta) \implies (\forall^{\infty} \nu \in \operatorname{Suc}_t(\eta))(\beta \leq h_t(\nu)).$$

If not said otherwise, every t is standard. Note:

$$s \in \text{sub}(t) \& t \text{ is standard} \Rightarrow s \text{ is standard},$$

where on sub(t) see part (8) below.

2. For an ordinal $\alpha < \omega_1$, let

$$\mathcal{T}_{\alpha} = \{t \in \mathcal{T} : \operatorname{Rang}(h_t) \subseteq \omega_1 \text{ and } h_t(\operatorname{rt}(t)) = \alpha\},\ \mathcal{T}_{<\alpha} = \bigcup_{\beta < \alpha} \mathcal{T}_{\beta}.$$

3. For $t \in \mathcal{T}$ let

 $\mathcal{A}_t = \big\{ \bar{A}: \quad \bar{A} = \langle A_\eta : \eta \in t \rangle, \ A_\eta \in [\omega]^{\aleph_0}, \ \text{and if} \ \eta \in t \ \text{is not maximal}, \\ \text{then} \ \langle A_\nu : \nu \in \mathrm{Suc}_T(\eta) \rangle \ \text{is a sequence of pairwise disjoint} \\ \text{subsets of} \ A_\eta \big\},$

$$\begin{array}{rcl}
\mathcal{A}_{\alpha} & = & \bigcup_{\alpha} \left\{ \mathcal{A}_{t} : t \in \mathcal{T}_{\alpha} \right\}, \\
\mathcal{A}_{<\alpha} & = & \bigcup_{\beta < \alpha} \mathcal{A}_{\beta}.
\end{array}$$

- 4. For $t \in \mathcal{T}$, $\eta \in t$ let $t^{[\eta]} = \{\nu : \eta \leq \nu \in t\} \in \mathcal{T}$.
- 5. For $t \in \mathcal{T}$ let $\lim_{t \to \infty} \{ \eta \in \omega : (\forall \ell < \omega) (\ell \geq \ell g(\operatorname{rt}(t)) \Rightarrow \eta \upharpoonright \ell \in t) \}$.
- 6. $\max(t) = \{ \eta \in t : \operatorname{Suc}_t(\eta) = \emptyset \}.$
- 7. We say that y is a front of $t \in \mathcal{T}$ if: $y \subseteq t$, $(\forall \eta, \nu \in y)(\neg \eta \triangleleft \nu)$ and

$$(\forall \eta)[(\eta \in \lim(t) \vee \eta \in \max(t)) \ \Rightarrow \ (\exists \ell \leq \ell g(\eta))(\eta \upharpoonright \ell \in y)].$$

We let $fr(t) = \{y : y \text{ is a front of } t\}.$

8. For $t \in \mathcal{T}$ let $\mathrm{sub}(t) = \{s \in \mathcal{T} : s \subseteq t \text{ and } \max(s) = \max(t) \cap s\}$. Clearly for a standard t and $s \in \mathrm{sub}(t)$ we have $h_s = h_t \upharpoonright s$. Let

$$\operatorname{sub}^-(t) = \{ s \in \mathcal{T} : s \subseteq t, \ s \in \operatorname{sub}(t) \text{ and } \operatorname{rt}(s) = \operatorname{rt}(t) \}.$$

- 9. For $s, t \in \mathcal{T}$ and a set \mathcal{Y} of fronts of t let $\mathrm{FT}^0_{\mathcal{Y}}(t,s)$ be the set of embeddings $f: s \longrightarrow t$ (i.e., f is one-to-one, $\mathrm{Dom}(f) = s$, $\mathrm{Rang}(f) \subseteq t$, $f(\mathrm{rt}(s)) = \mathrm{rt}(t)$ and $(\forall \eta, \nu \in t)(\eta \lhd \nu \Rightarrow f(\eta) \lhd f(\nu)))$ which respect each $y \in \mathcal{Y}$, i.e., $\{\eta: f(\eta) \in y\}$ is a front of s for every $y \in \mathcal{Y}$.
- 10. If \mathcal{Y} is the set of all fronts of t we may omit it.
- 11. $FT^1(t,s)$ is the set of $f \in FT^0(t,s)$ such that

$$\eta \in \operatorname{Suc}_s(\operatorname{rt}(s)) \quad \Rightarrow \quad h_s(\eta) = h_t(f(\eta)).$$

12. $FT^2(t,s)$ is the set of all $f:s \longrightarrow t$ such that:

$$\eta \lhd \nu \Leftrightarrow f(\eta) \lhd f(\nu),$$

 $\{f(\eta) : \eta \in \operatorname{Suc}_s(\operatorname{rt}(s))\}\$ is a front of t , and for $\eta \in \operatorname{Suc}_s(\operatorname{rt}(s)), \quad f \upharpoonright s^{[\eta]}$ is one-to-one onto $t^{[f(\eta)]}.$

13. For an ideal I let

$$\mathcal{ST} = \mathcal{ST}_I = \{(t,g): t \in \mathcal{T} \text{ and } g: t \setminus \{\operatorname{rt}(t)\} \longrightarrow \operatorname{Dom}(I) \text{ are such that}$$

 $\eta \in t \setminus \max(t) \Rightarrow \{g(\nu): \nu \in \operatorname{Suc}_T(\eta)\} \in I^+\}.$

We usually omit I if it is clear from the context (here it is fixed).

- 14. For (t^1, g^1) , $(t^2, g^2) \in \mathcal{ST}$ for $\ell = 0, 1, 2$ let $\mathcal{FT}^{\ell}((t^1, g^1), (t^2, g^2))$ be the set of $f \in FT^{\ell}(t^1, t^2)$ such that: $g^2 = g^1 \circ f$.
- 15. Let

$$\mathcal{T}\mathcal{T}_{t} = \{(t, \bar{A}, g) : \bar{A} \in \mathcal{A}_{t} \text{ and } (t, g) \in \mathcal{S}\mathcal{T}\}, \\
\mathcal{T}\mathcal{T} = \bigcup_{t} \mathcal{T}\mathcal{T}_{t}, \\
\mathcal{T}\mathcal{T}_{\alpha} = \{(t, \bar{A}, g) : t \in \mathcal{T}_{\alpha} \text{ and } (t, \bar{A}, g) \in \mathcal{T}\mathcal{T}_{t}\}, \\
\mathcal{T}\mathcal{T}_{<\alpha} = \bigcup_{\beta < \alpha} \mathcal{T}\mathcal{T}_{\beta}.$$

- 16. Let $\mathcal{FT}^{\ell}((t^1, \bar{A}^1, g^1), (t^2, \bar{A}^2, g^2))$ be the set of $f \in \mathcal{FT}^{\ell}((t^1, g^1), (t^2, g^2))$ such that $\eta \in t^2 \implies A^1_{f(\eta)} \subseteq A^2_{\eta}$.
- 17. If we omit ℓ (in FT^{ℓ} , \mathcal{FT}^{ℓ}) we mean $\ell = 0$.

Definition 3.15. We define a partial order on $TT_{\leq \alpha}$:

$$(t^1, \bar{A}^1, g^1) \leq^\ell (t^2, \bar{A}^2, g^2) \quad \text{ if and only if } \quad \mathcal{FT}^\ell((t^1, \bar{A}^1, g^1), (t^2, \bar{A}^2, g^2)) \neq \emptyset.$$

Observation 3.16. \leq^{ℓ} really is a partially order of TT.

Definition 3.17. 1. For $t \in \mathcal{T}$, let

fsub
$$(t) = \{ s \in \mathcal{T} : \text{ for some finite } w \subseteq \operatorname{Suc}_t(\operatorname{rt}(t)) \text{ we have } s = \{ \eta \in t : \neg (\exists \nu \in w) (\nu \leq \eta) \} \}.$$

2. For $(t^1, \bar{A}^1, g^1), (t^2, \bar{A}^2, g^2) \in \mathcal{TT}$ we define $\mathcal{FT}^\ell_*((t^1, \bar{A}^1, g^1), (t^2, \bar{A}^2, g^2))$ as the set of all f such that for some $t^3 \in \mathrm{fsub}(t^2)$ we have

$$f \in \mathcal{FT}^{\ell}((t^1, \bar{A}^1, g^1), (t^3, \bar{A}^2 \upharpoonright t^3, g^2 \upharpoonright t^3)).$$

- 3. $(t^1, \bar{A}^1, g^1) \leq_*^{\ell} (t^2, \bar{A}^2, g^2)$ if and only if $\mathcal{FT}_*^{\ell}((t^1, \bar{A}^1, g^1), (t^2, \bar{A}^2, g^2))$ is not empty.
- **Fact 3.18.** 1. $<^{\ell}_*$ is a partial order of TT such that: $<^{\ell}$ is a subset of $<^{\ell}_*$.
 - 2. Any $<_*^{\ell}$ -increasing chain of length ω in $TT_{<\alpha}$ has an upper bound in $TT_{<\alpha}$.
 - 3. $TT_{\leq \omega_1} = TT_{<\omega_1}$.

Proposition 3.19. 1. If $(t, \bar{A}, g) \in \mathcal{TT}_{\leq \alpha}$, and $B \subseteq \omega$,

then for some $(t', \bar{A}', g') \in \mathcal{TT}_{\leq \alpha}$ we have:

- (α) $(t, \bar{A}, g) \leq^1 (t', \bar{A}', g')$, in fact $t' \in \text{sub}(t)$,
- (β) $\bigcup \{A_η : η \in t \setminus \{\text{rt}(t)\}\}\$ is a subset of B or is disjoint to B.
- 2. Similarly for TT.

Proof. (1) By induction on α .

(2) Similarly.

Proposition 3.20. If $(t, \bar{A}, g) \in \mathcal{TT}_{\leq \alpha}$ and E is an equivalence relation on ω , then for some (t, \bar{A}', g') and front y of t' we have:

- $(\alpha) (t, \bar{A}, g) \leq^1 (t', \bar{A}', g'),$
- (β) for η ∈ y, $A'_η$ is included in one E-equivalence class,
- (γ) for $\eta \neq \nu$ from y, the E-equivalence classes in which A'_{η} , A'_{ν} are included, are distinct (hence disjoint),
- (δ) on $A'_{\mathrm{rt}(t')}$ we have: E is either trivial or refines $\langle A'_{\eta} : \eta \in \mathrm{Suc}_{t'}(\mathrm{rt}(t')) \rangle$.

Proof. By induction on α .

Proposition 3.21 (CH). Let I be an ideal such that $|Dom(I)| \leq \aleph_1$ and

$$(\forall X \in I^+)(\exists Y \in I^+)(Y \subseteq X \& |Y| = \aleph_0).$$

There is a \leq^1_* -increasing sequence $(\langle t^{\zeta}, \bar{A}^1, g^{\zeta} \rangle : \zeta < \omega_1)$ of members of $\mathcal{TT}_{\leq \omega_1}$ such that $\{\bigcup_{\eta \in t^{\zeta}} A^1_{\eta} : \zeta < \omega_1\}$ generates a non-principal ultrafilter D on ω which

is a Q-point, and for every equivalence relation E on ω , for some ζ , $\langle A_{\eta}^{\zeta} : \eta \in \operatorname{Suc}_{t^{\zeta}}(rt(t^{\zeta})) \rangle$ refines $E \upharpoonright A_{\operatorname{rt}(t^{\zeta})}^{\zeta}$.

Remark: This construction gives an ultrafilter D on ω , a Q-point such that

$$D' \leq_{RK} D \Rightarrow D'$$
 is not NWD.

Proposition 3.22. In 3.21, if in addition an ideal I' satisfies

(*) $|\mathrm{Dom}(I')| \leq \aleph_1$ and if $(t, \bar{A}, g) \in \mathcal{TT}_{<\omega_1}$ and $g' : \omega \longrightarrow \mathrm{Dom}(I')$ then for some $(t', \bar{A}', g') \geq (t, \bar{A}, g)$ we have $g'[A'_{\langle \rangle}] \in I'$,

then we can demand that D is an I'-ultrafilter (see 3.11(4)).

* * *

There are many problems on the σ -versions of cardinal invariants, and I think for some the method of [100], [120] is relevant, e.g.

Question 3.23 (See Brendle and Shelah [17]). Does $\chi_{\sigma}(D) = \chi(D)$ for all ultrafilters D on ω ? Recall that

 $\chi(D) = \min\{|\mathcal{A}|: \mathcal{A} \subseteq D \text{ and for every } A \in D \text{ for some } B \in \mathcal{A} \text{ we have } B \subseteq A\}$ $\chi_{\sigma}(D) = \min\{|\mathcal{A}|: \mathcal{A} \subseteq {}^{\omega}D \text{ is such that for every } \bar{A} \in {}^{\omega}D, \text{ for some }$ $\bar{B} \in \mathcal{A} \text{ we have } (\forall n < \omega)(\exists m < \omega)(B_m \subseteq {}^*A_n)\}.$

So a reasonable scenario to prove the consistency of a negative answer runs as follows: let, e.g., $\mu = \aleph_{\omega}$. We use FS iteration of c.c.c. forcing notions, $\langle \mathbb{P}_i, \mathbb{Q}_j : i \leq \delta^*, j < \delta^* \rangle$. We have \mathbb{P}_i -names \mathcal{D}_u^i , $\mathcal{A}_{u,\gamma}$ (for $\gamma < \gamma_u^i$ and $u \in [\mu]^{<\aleph_0}$) such that:

- \tilde{D}_u^i is the filter on ω generated by $\{\tilde{A}_{u,\gamma}: \gamma < \gamma_u^i\}$ and the co-bounded sets, γ_u^i are increasing with i, $\tilde{D}_u^i \subseteq \tilde{D}_v^i$ if $u \subseteq v$, and $\tilde{D}_u^j \subseteq \tilde{D}_u^i$ for j < i.

To simplify we decide:

(*) if
$$j < i$$
, $\Vdash_{\mathbb{P}_j}$ " $\underline{A} \subseteq \omega$ and $u \in [\mu]^{<\aleph_0}$ and $\underline{A} \subseteq^* \underline{A}_{u,\gamma}$ for every $\gamma < \gamma_u^j$ " then $\Vdash_{\mathbb{P}_i}$ "for every $v \in [\mu]^{<\aleph_0}$, $\underline{A} \notin \underline{\mathcal{D}}_v^i$ ".

Also for the following it seems reasonable to try to be influenced by [100], [120].

Question 3.24 (See Brendle and Shelah [17]). Can $\pi\chi(D)$ be singular, where

$$\pi\chi(D) = \min\{|\mathcal{A}|: \mathcal{A} \subseteq [\omega]^{\aleph_0}, \text{ and for every } B \in D \text{ for some } A \in \mathcal{A} \text{ we have } A \subseteq^* B\}.$$

4. NICELY DEFINED FORCING NOTIONS

Rosłanowski and Shelah [85], and [107], [164] relate as algebraic three dimensional varieties relate to manifolds in \mathbb{R}^n and these, in turn, relate to general topology. In [107] (on nep and snep) and in Judah and Shelah [56] (on Souslin forcings) we deal with forcing notions defined in an absolute enough way; in [85] (more in [87], [88]) with forcing notions defined in an explicit way (say as tress and generally by creatures), in [164] we deal with forcing notions related to the continuum.

Our problem with speaking about [85], [86], [87] and [88] is that much work is in progress, still orthogonal to it is the question whether in the main theorems of [85], all the assumptions are needed. That is, within the framework of condition trees or ω -sequences of creatures, are the demands on the norms necessary? This is dealt with for the conditions for properness in [85], showing necessity but there are still gaps remaining.

Question 4.1. Are the sufficient conditions for properness in [85, §2] necessary? The test case (chosen in [85]) is

$$\mathbb{Q} = \{ \langle w_n : n < \omega \rangle : w_n \subseteq 2^n, w_n \neq \emptyset \text{ and } \lim_{n \to \omega} |w_n| \} = \infty$$

ordered by $\bar{w} \leq \bar{w}' \iff (\forall n \in \omega)(w'_n \subseteq w_n)$.

Though properness is the main thing and there we look for counterexamples only for properness, it is interesting to know:

Question 4.2. Concerning other theorems of [85], are they sharp?

There are more specialized problems, probably solvable in this context.

Question 4.3. Is there an ω -bounding forcing notion adding a perfect set of random reals?

It seems this should not be hard if true.

The following problems (raised by Komjath and Steprāns respectively) seem to me a matter of choosing the right variant of [85] or [87] and having the right finite combinatorics.

1. Can each $A \in [{}^{\omega}2]^{\aleph_1}$ be null while the union of some \aleph_1 lines Question 4.4. in $\mathbb{R} \times \mathbb{R}$ is not null?

2. For reals $0 < a_0 < a_1 \le 1$, is it consistent with ZFC that: for $\ell < 2$,

 $\ell = 0$ iff some $A \in [\mathbb{R}]^{\aleph_1}$ has positive Hausdorff capacity for a_{ℓ} ?

* * *

I suppose that the feeling that the Cohen forcing notion and the random real forcing notion occupy a special place is old; probably more in the version speaking on the ideal of null sets and the ideal of meagre sets. I feel the former version is more interesting. For me this translates to

Problem 4.5. Among Souslin c.c.c. forcing notions, are Cohen forcing and random forcing special?

Some progress was made in [156].

Theorem 4.6. If a Souslin c.c.c. forcing notion \mathbb{Q} adds $\tilde{\eta} \in {}^{\omega}\omega$ not dominated by any old $\nu \in {}^{\omega}\omega$, then forcing with \mathbb{Q} adds a Cohen real. (The "Souslin" is needed for enough absoluteness, so with the existence of large cardinals we can allow a larger family).

So the Cohen forcing notion is the minimal one among Souslin c.c.c. forcing notions adding an undominated real, so it is natural to conjecture:

Problem 4.7. Show that any Souslin c.c.c. forcing notion adding a real adds a Cohen real or adds a random real.

This really will show that Cohen and random are special.

In a sense the realm of Souslin c.c.c. forcing notion can be looked as being divided between the ω -bounding (with Random forcing as prototype) and those forcing notions adding an undominated real (with Cohen forcing as prototype); we can further distinguish those adding a dominating real.

However, the situation is very unbalanced: among Souslin c.c.c. forcing notions adding an undominated real we have many examples and a <*-minimal one, Cohen, (see Definition 4.8 below).

On the other side we have no idea what occurs among the ${}^{\omega}\omega$ -bounding ones: probably random real is the unique one, but it is not out of the question that there is a plethora (adding one or many randoms is an irrelevant distinction; we can even order $(\mathbb{Q}, \underline{r})$, \underline{r} a \mathbb{Q} -name of a real such that the order depends only on the subforcing \underline{r} generates).

Definition 4.8. Let $\mathbb{Q}_1, \mathbb{Q}_2$ be definitions of forcing notions (absolute enough) say as in [107], or Souslin.

- 1. $\mathbb{Q}_1 \leq_0^* \mathbb{Q}_2$ if forcing with \mathbb{Q}_2 adds a generic for \mathbb{Q}_1 and we let $<^*$ mean $<_0^*$.
- 2. $\mathbb{Q}_1 \leq_1^* \mathbb{Q}_2$ means: for some n, if we force by iteration n times of \mathbb{Q}_2 , we add a generic for \mathbb{Q}_1 .
- 3. $\mathbb{Q}_1 \leq_{2,fs}^* \mathbb{Q}_2$ is defined similarly using FS iteration of length $< \omega_1$.
- 4. $\mathbb{Q}_1 \leq_{2,cs}^* \mathbb{Q}_2$ is defined similarly using CS of length $< \omega_1$.

Note: 4.7 is on the interval between the control measure problem (see Fremlin [36]) and von Neumann question which says: is any complete c.c.c. Boolean Algebra which as a forcing is ω —bounding, a measure algebra. Another way to express the thought that Cohen and random are special was Kunen's conjecture, see Kunen [71], Kechris and Solecki [65], Solecki [174], [173] and Rosłanowski and Shelah [86].

It is natural to investigate the partial orders from 4.8. So,

Problem 4.9. Investigate the quasi order \leq^* (and its variants) for \mathbb{Q} which are nep (see [107]) or which are c.c.c. \aleph_0 -nep or which are c.c.c. \aleph_0 -snep.

[We may concentrate on those with a generic real (those are the main interest for 4.13(1), (2) below).]

An example is (and probably not hard):

Question 4.10. Prove that dominating real forcing (i.e., the Hechler forcing notion) is \leq^* -minimal among Souslin c.c.c. forcing notions adding a dominating real. [For \leq^*_1 this is easy.]

Looking more serious are

Question 4.11. Can you characterize the $<^*$ -minimal \mathbb{Q} , which add a Cohen real but are not equivalent to the Cohen forcing (hopefully there is one or at least there are only few).

Question 4.12. Can you characterize the $<^*$ -minimal \mathbb{Q} among the non-minimal \mathbb{Q} which add a dominating real but are stronger than the Hechler forcing notion?

A positive solution of 4.7 would also show that the only symmetric Souslin c.c.c. forcing notions are the Cohen forcing and the random forcing (by [107, §9]).

* * *

Why should we be interested in Souslin proper or in nep forcing or better yet, why am I? The reason has been iteration theorems; when you are interested in iterating some very special forcing notions, the proof of their properness gives more, e.g., the existence of generic conditions over models occurs also for countable models of versions of ZFC which are not necessarily $\prec (\mathcal{H}(\chi), \in)$. Moreover, some things are preserved by iterations and this is helpful for specific problems which is the point of Judah and Shelah [56]. [In [56] this was phrased using descriptive set theory, getting Souslin proper. However, this does not cover the Sacks forcing notion, the Laver forcing notion, etc., which was accomplished by nep.]

Needless to say, I think iteration theorems for forcing are important and interesting (otherwise, normally I would not have written a book on the subject - see §2).

Another basic reason is that the family of nep forcing notions forms a natural class. Now, while I feel that general sets are much more basic and interesting then families of definable ones, and so prefer $\mathcal{P}(\mathcal{P}(\omega))$ to the family of projective sets, certainly they are interesting and natural.

Another reason is "large" ideals. Let I be a κ -complete ideal on λ . Gitik and Shelah [42] start by proving that $\mathcal{P}(\kappa)/I$ cannot be (the Boolean algebra which up to isomorphism is equivalent to) the Cohen forcing or random real forcing, an old question which Fremlin promoted (see [37]), which comes from asking: can the classical result of Solovay [175] (saying that consistently 2^{\aleph_0} is real valued measurable, now the Maharam type there was large) be improved to get small Maharam type.

But then [42] turns to:

Problem 4.13. 1. Prove that $\mathcal{P}(\kappa)/I$ cannot be a Souslin c.c.c. forcing generated by the name of one real $\tilde{\eta}$ (where I is a κ -complete ideal on κ or at least \aleph_1 -complete).

- 2. Similarly for Souslin proper (or weaken the definability demand natural as the existence of the ideal implies more absoluteness).
- 3. Even reasonable subclasses or cases are interesting.

Problem 4.14. Similarly, we can ask about a σ^+ -complete ideal I on κ such that $\mathcal{P}(\kappa)/I$ has a dense subset isomorphic to a partial order defined in $(\mathcal{H}_{<\sigma}(\kappa'), \in)$ with parameters.

In Gitik and Shelah [42], [44], [41], in addition to information on adding not too many random or Cohen reals, and (toward 4.13) to general criteria for impossibility, we consider more specific cases (see then [172]). The problems lead us to properties of definable forcing notions like symmetry. The theorems on Cohen and random reals use the symmetry (i.e, the Fubini theorem), but other properties pop up naturally, e.g., for Souslin c.c.c. forcing \mathbb{Q} with a dominating real η as generic, to show impossibility it suffices to show: $\Vdash_{\mathbb{Q}}$ " $\mathfrak{b} = \aleph_1$ " (by [42]). Maybe the work on the ideals is done and we just need to verify that always at least one of the criteria applies (at least for large subclasses). Now [156], [107, §8,§9] comes to my mind.

Considerations like this lead to questions like

Question 4.15. Find sufficient conditions on \mathbb{Q} for " $\mathbb{Q} * \text{Random}/\text{Random}$ adds no random real".

(This question is chosen since it is also interesting because if the condition is reasonable enough, it suffices for proving CON(cov*(null) < non(meagre)), see Bartoszyński, Rosłanowski and Shelah [8], [7].)

Question 4.16. Investigate commuting pairs (see [107]).

For such considerations I had felt that a peculiar property of Cohen forcing and random forcing is their satisfying: "being a maximal antichain is a Borel property"; this leads to

Definition 4.17. A forcing notion \mathbb{Q} is very Souslin c.c.c. if it is Souslin c.c.c. and also the notion of " $\{r_n : n < \omega\}$ is a maximal antichain" is Σ_1^1 .

We hope this will turn out to be a good dividing line of the Souslin c.c.c. forcing (so helping to prove theorems). This is because I suspect the answer to the following is yes.

Question 4.18. Prove: If \mathbb{Q} is a Souslin c.c.c. forcing notion, say with generic real \underline{r} and it is not very Souslin c.c.c. above any $p \in \mathbb{Q}$ then $\Vdash_{\mathbb{Q}}$ " $\mathfrak{b} = \aleph_1$ ". (This should help 4.13 by [42, §4]). See on this [108].

As in [107] we can define (restricting κ to be \aleph_0 for simplicity)

Definition 4.19. 1. A forcing notion \mathbb{Q} is ω -nw-nep if there is a sequence $\bar{\varphi} = \langle \varphi_0, \varphi_1, \varphi_2 \rangle$ of Σ_1^1 definitions such that:

- (a) the set of members of \mathbb{Q} and $\leq^{\mathbb{Q}}$ are Σ_1^1 sets (of reals) defined by φ_0, φ_1 , respectively,
- (b) if N is an $(\bar{\varphi}, \omega)$ -nw-candidate (that is, a model of ZFC, suitable version of ZFC, not necessarily well-founded but with standard ω), and with the real parameters of the φ_{ℓ} 's, and $p \in \mathbb{Q}^N = \{x : N \models \varphi_0(x)\}$, then for some $q \in \mathbb{Q}$, we have
 - $(\alpha) p \leq q,$

- (β) q is (N,\mathbb{Q}) -generic, which means that for every $\mathcal{I} \in \mathrm{pd}(N,\mathbb{Q}) =$ $\{\mathcal{J} \in N : N \models "\mathcal{J} \text{ is predense in } \mathbb{Q}"\}, \text{ for some list } \langle r_n : n < \omega \rangle \text{ of }$ $\{x: N \models \text{``} x \in \mathcal{I} \text{''}\}\$ we have $\{r_n: n < \omega\}$ is predense above q,
- (γ) moreover $\varphi_2(q, \langle r_n : n < \omega \rangle)$ holds,
- (c) if $\varphi_2(q', \langle r'_n : n < \omega \rangle)$ then the set $\{r'_n : n < \omega\}$ is predense above q'. 2. Omitting the "nw" or writing just "w" means we allow only well-founded candidates.
- 3. [On Ur see below]. We say $r \subseteq Ur$ (or $r \subseteq \mathcal{H}_{\langle \aleph_1}(Ur)$ in the w-case) is generic for \mathbb{Q} if: r is a \mathbb{Q} -name and " $a \in r$ " is determined by the truth value $\varphi_{r,a}[\mathcal{G}_{\mathbb{Q}}]$ and the sequence $\langle \varphi_{r,a} : a \rangle$ is definable in $\mathfrak{B}_{\mathbb{Q}}$ (see [108]).

Now, there are more examples of ω -nw-nep forcing notions in addition to Cohen and random: say the Sacks forcing notion. But about the Laver forcing notion we should beware; note that we can guarantee that for every $\mathcal{I} \in \mathrm{pd}(\mathbb{Q}, N)$, for some front X of q we have $\eta \in X \implies q^{[\eta]}$ is above a member of \mathcal{I} , but being a front is not absolute from $(\bar{\varphi}, \omega)$ -nw-candidates, as they are not necessarily well-founded. In fact, we can easily craft a counterexample: assume $N \models "\alpha$ a countable ordinal", but from the outside not well ordered. There are $f, T \in N$ (so actually $f = f_{\alpha}$, $t = T_{\alpha}$) such that

```
N\models \text{``} \quad T\subseteq {}^{\omega>}\omega \text{ is closed under initial segments}, \quad \langle\rangle\in T,
                         f: T \longrightarrow \alpha + 1, \qquad f(\langle \rangle) = \alpha,
                         \begin{array}{lll} f(\eta) > 0 & \Rightarrow & (\forall n)(\eta ^\frown \langle n \rangle \in T), \\ f(\eta) = 0 & \Rightarrow & (\forall n)(\eta ^\frown \langle n \rangle \notin T), \\ f(\eta) = \beta + 1 & \Rightarrow & (\forall n < \omega)[\eta ^\frown \langle n \rangle \in T \ \& \ f(\eta ^\frown \langle n \rangle) = \beta], \end{array}
                         if f(\eta) = \delta is limit
                          then \langle f(\eta \hat{\ } \langle n \rangle) : n < \omega \rangle is strictly increasing with limit \delta ".
```

Let

$$\begin{array}{lcl} \mathcal{I}_0 & = & \{(^{\omega>}\omega)^{[\eta]}: \eta \in T, \ f(\eta) = 0\}, \\ \mathcal{I}_{n+1} & = & \{(^{\omega>}\omega)^{[\eta^\frown \rho]}: \eta \in \mathcal{I}_n, \ \rho \in T, \ f(\rho) = 0\}. \end{array}$$

Clearly, above no q is every \mathcal{I}_n predense. Still,

Theorem 4.20. The ω_{ω} -bounding and almost ω_{ω} -bounding forcing notions covered by [85] and [87] are all ω -nw-nep.

What about iterations? Now, for unifying the treatment of finite support and countable support we revise our definition to have two quasi-orders $\leq^{\mathbb{Q}}, \leq^{\mathbb{Q}}_{\mathrm{pr}}$ such that $p \leq_{\text{pr}} q \Rightarrow p \leq q$. Hence in Definition 4.19 we add $\varphi_{1,2}$ (absolute just like φ_1) serving as a definition of $p \leq_{\text{pr}} q$. The support is countable but finite for the apure cases, i.e., only for finitely many α , $\neg(\emptyset_{\alpha} \leq_{\text{pr}} p(\alpha))$. First assume the length is $\alpha^* < \omega_1$, so we can use a parameter coding a well ordering on ω with this order type. We should repeat the proof in [107] in order to prove preservation in this case, but we better not use the $L_{\omega_1,\omega}$ -completeness as there, as we have problem with well-foundedness. So we just demand: elements in \mathbb{P}_{α} have depth $<\omega\alpha$ (or so).

What about long iterations? It seems to me, at least now, better (and fit to [107], too) to use a set of urelements \mathbf{Ur} ; let \mathfrak{C} [and \mathfrak{B}] be models with universes \mathbf{Ur} [or $\subseteq \mathbf{Ur}$ and $\mathcal{S} \subseteq [\mathbf{Ur}]^{\leq \aleph_0}$ be unbounded (usually stationary, if $a \in \mathcal{S}$ then $\mathfrak{B} \upharpoonright a \subseteq \mathfrak{B}$ and $\mathfrak{C} \upharpoonright a \subseteq \mathfrak{C}$), and anyhow the family of nw-candidates should be $\langle (\aleph_1)$ -directed and uord $\subseteq \mathbf{Ur}$ is a well ordered set (which will serve as the length of the iteration). Now, a candidate will be a countable model $N \subseteq (\mathcal{H}(\chi), \in)$ of ZFC^-_* (where $\mathcal{H}(\chi)$ includes the urelements), $N \cap \mathbf{Ur} \in S$, where \mathbf{Ur} and the relations of \mathfrak{B} and \mathfrak{C} are the considered relations. We define an nw-candidate similarly but now \in^N is a relation on N and N is not necessarily well-founded (but the order type of the well ordered ordinals of N is $> otp(N \cap uord)$. In Definition 4.19 we demand, of course, that φ_{ℓ} are upward absolute from the nw-candidates. Now we can use uord as the index set for iteration (instead of the true ordinals) and there are no problems.

This set-up looks like a nice army with no enemy yet, but this seems to me a natural dividing line among the nep forcing notions and therefore reasonable for our interest in those forcing notions per se. I hope it will help, particularly with 4.13 (and even more so by the c.c.c. version see 4.22 below).

More on preservation (for not necessarily c.c.c. forcing notions), commutativity, associativity of generic sets, and countable for pure/finite for apure support iterations see [108].

A restricted version of the large continuum is

Problem 4.21. Can we have long $(> \omega_2)$ iterations of ${}^{\omega}\omega$ -bounding forcing notions (or at least nw-nep ones) not collapsing cardinals and not adding Cohen reals?

What about the c.c.c. (nw-nep) ones (or even very Souslin c.c.c.) ones?

Definition 4.22. \mathbb{Q} is c.c.c.-nw-nep if it is a pair $\langle \varphi_0, \varphi_1 \rangle$ of formulas such that

- $(\alpha) \ \mathbb{Q} = \bigcup \{\mathbb{Q}^N : N \text{ is an nw-candidate}\} \subseteq {}^{\omega}\text{uord, similarly}$ $\leq_Q = \bigcup \{\leq_{\mathbb{Q}}^N : N \text{ is an nw-candidate}\},$ recall that in N, \mathbb{Q}^N , $\leq_{\mathbb{Q}}^N$ are defined by the appropriate φ_{ℓ} , (β) all φ_{ℓ} are upward absolute among nw-candidates,
- (γ) if $N \models$ " $\mathcal{I} \subseteq \mathbb{Q}$ is predense", then $\mathcal{I}^{\mathbb{Q}}$ is really predense.

Proposition 4.23. 1. The Cohen and random forcing notions are c.c.c.-nwnep.

- 2. The class of c.c.c.-nw-nep is closed under FS iterations.
- 3. This class is also closed under subforcings.

And I am curious to know:

Problem 4.24. Does 4.23 exhaust all c.c.c-nw-nep forcing notions (at least those with a generic real)?

Being interested in classifying nep c.c.c. forcing notions, we may consider sweetness; the discussion below is in fact an introduction to [88]. Sweetness phenomenons are when we can build homogeneous forcing notions (as in [136, §7,§8]), sour phenomenons are strong negations (as in [136, §6]).

Problem 4.25. 1. For which (\mathbb{Q}, η) , nep c.c.c. forcing notions, is it consistent that:

(a) there is a (\mathbb{Q}, η) -generic real over $\mathbf{L}[A]$ for every $A \subseteq \omega_1$, and

(b) for every subset $B \in \mathbf{L}[\mathbb{R}]$ of ${}^{\omega}2$ for some $A \subseteq \omega$, for a dense set of $p \in (\mathbb{Q}, \underline{\eta})$: for some truth value \mathbf{t} , if η is a $(\mathbb{Q}^{\mathbf{L}[A]}_{\geq p}, \underline{\eta})$ -generic real then $\eta \in B \Leftrightarrow \mathbf{t}$.

So for this question, random reals are complicated (see [136, §6]), whereas Cohen real and universal-meagre one (and dominating = Hechler reals) are low, see Judah and Shelah [59].

More generally,

Problem 4.26. Let $(\mathbb{Q}, \underline{\eta})$ be a nep c.c.c. forcing notion, and κ be a cardinal number. Let $I_{(\mathbb{Q}, \underline{\eta})}^{\kappa}$ be the κ -complete ideal generated by sets of the form

$$A_N = \{ \eta : \eta \text{ is not } (\mathbb{Q}, \eta) \text{-generic over } N \}$$

for countable models $N \prec (\mathcal{H}(\chi), \in)$ to which (\mathbb{Q}, η) belongs.

What is the consistency strength of "every projective set, or even every set from $\mathbf{L}[\mathbb{R}]$, is equal to a Borel set modulo $I_{(\mathbb{O},\eta)}^{\kappa}$ "?

We hope for a strong dichotomy phenomena, i.e., if the answer above is negative for (\mathbb{Q}, η) (so the sweetness fails), then a strong negation holds, so we call such phenomena sourness.

Definition 4.27. Let $\mathbb{Q}_1, \mathbb{Q}_2$ be nep c.c.c. forcing notions definable in \mathbf{L} (or $\mathbf{L}[r]$). We say that $\mathbb{Q}_1, \mathbb{Q}_2$ are explicitly sour over Cohen *if* we can find \mathbb{Q}_{ℓ} -names $\tilde{\eta}_{\ell}$ of Cohen reals (for $\ell = 1, 2$) such that

```
if G_{\ell} \subseteq \mathbb{Q}_{\ell} is generic over L (or L[r]) for \ell = 1, 2 then \eta_1[G_1] \neq \eta_2[G_2].
```

We should note that there may be homogeneity for wrong reasons, i.e., maybe when we force, very few $(\mathbb{Q}, \underline{\eta})$ -generic reals over \mathbf{L} are added and then homogeneity holds for "degenerated" reasons; we may call such cases sacharin. For more on this direction see [87].

Speaking about the class of sweet forcing notions we should mention the following problem.

Problem 4.28. For any cardinal κ and a large cardinal property (or consistency strength) we may ask the following.

- 1. Is there a widest class K of absolute enough forcing notions such that for some forcing notion \mathbb{P} we have
 - (a) $\Vdash_{\mathbb{P}} \kappa = \aleph_1$,
 - (b) \mathbb{P} is homogeneous for complete subforcings from \mathcal{K} , moreover
 - (c) if $\mathbb{P}^* \triangleleft \mathbb{P}$, \mathbb{P}^* has a generic real then $\mathbb{P}/\mathcal{G}_{\mathbb{P}^*}$ satisfies (b) in $\mathbf{V}[\mathcal{G}_{\mathbb{P}^*}]$.
- 2. If not, at least give a wide enough such class.
- 3. Are there two classes $\mathcal{K}_1, \mathcal{K}_2$ as above, such that there is no class with the respective property including $\mathcal{K}_1 \cup \mathcal{K}_2$? Or even that the consistency strength of the (now) obvious conclusion is higher than the given one?

Now, the variants of sweetness try to deal with the case of $\kappa = \aleph_1$ and the consistency strength ZFC (see Rosłanowski and Shelah [88, §3]); the theory of determinacy is applicable to the case $\kappa = \aleph_1$ and maximal consistency strength (see Woodin [182]), and [115] intends to deal with the case of "ZFC + κ is strongly inaccessible" (and no further consistency strength assumptions).

Question 4.29. Is there a sweet forcing notion (see [136, §7]), preferably natural, such that it cannot be completely embedded into the forcing notion constructed in [136, §7] (it was gotten by composing \mathcal{UM} , amalgamating and direct limits), or at least not "below \mathcal{UM} " (in the sense of \leq_1^* , see 4.8)?

* * *

Let me now mention free iterations:

In nep (also above) we can replace CS by free limit as in [132, Ch.IX]. (This was my third proof for preservation of properness (the first was like [108, §2], the second was like the one in [132, Ch.III]; this third proof looks very natural but no real reason for replacing CS iteration by it has appeared).

In particular

Proposition 4.30. Definition of $L_{\omega_1,\omega}$ -free iteration is absolute enough, so we have our φ_2 (we ignore the nw).

Preservation (of reasonable properties) by CS iterations of proper forcing (or variants) seems to me a worthwhile subject. For ${}^{\omega}\omega$ -bounding the situation is nice, "proper $+{}^{\omega}\omega$ -bounding" is preserved, and analogous results hold for a large family of properties even, e.g., "D generates a P-point ultrafilter on ω ". But some properties do not fit, though still are preserved in limits (see [164, Ch.XVIII,§3]). For example, we shall consider below the case of " $A \subseteq {}^{\omega}2$ is non-null". The simple preservation is: in the existence of generic conditions we can preserve "a given old $\eta \in {}^{\omega}2$ is random over the model $N[\mathcal{G}_{\mathbb{P}}]$ ".

Question 4.31. Assume that

- (a) \mathbb{Q} is a CS iteration, or a $L_{\omega_1,\omega}$ -free iteration,
- (b) each \mathbb{Q}_{α} is proper, or a nep forcing notion,
- (c) each \mathbb{Q}_{α} is "non-null for (\mathbb{S}, r) -preserving" (if each \mathbb{Q}_{α} is nep and r is, e.g., random, then by [107] this is equivalent to not making old "2 null).

Does \mathbb{P}_{α} have the property from clause (c)? (so we have four versions of the question, as for clause (a) and for clause (b) we can choose the first or the second possibilities).

Assume:

(*) each \mathbb{Q}_i has a generic $r_i \subseteq \alpha_i$.

(Hence \mathbb{P}_{α} have this property as it is preserved).

We assume knowledge of free iterations (see [164, IX,§1,§2]); in short, if $\mathbb{P}_n \Leftrightarrow \mathbb{P}_{n+1}$ for $n < \omega$, let \mathbb{P}_{ω} be

- $\{\psi: \ \psi \text{ is a sentence in the } L_{\omega_1,\omega} \text{ propositional calculus with the set of propositional variables } \bigcup \{\mathbb{P}_n: n < \omega\} \text{ such that in some forcing extension of } \mathbf{V} \text{ there is } G \subseteq \bigcup \{\mathbb{P}_n: n < \omega\} \text{ satisfying}$
 - (a) for each $n < \omega$, $G \cap \mathbb{P}_n$ is generic over \mathbf{V} ,
 - (b) looking at G as assigning truth values to members of $\bigcup \{\mathbb{P}_n : n < \omega\}$, it assigns the value truth to $\psi\}$.

The order of \mathbb{P}_{ω} is the natural one.

 $^{{}^3\}mathbb{S}$ is a nep forcing notion, \underline{r} is a hereditarily countable \mathbb{S} -name, so $(\mathbb{S},\underline{r})$ induces an ideal on ${}^{\omega}2$, \mathbb{S} may be random and then the ideal is the ideal of null sets.

We deal with an $L_{\omega_1,\omega}$ -free iteration $\bar{\mathbb{Q}}$ such that

$$\Vdash_{\mathbb{P}_i}$$
 " \mathbb{Q}_i is nep and $(\omega_2)^{\mathbf{V}^{\mathbb{P}_i}}$ not null in $(\omega_2)^{\mathbf{V}^{\mathbb{P}_i * \mathbb{Q}_i}}$ ".

This is quite a wide case. What does it mean for a successor $\alpha = \ell g(\bar{\mathbb{Q}})$? Say, $\alpha = 2$ so we know that in $\mathbf{V}^{\mathbb{Q}_0}$, $(^{\omega}2)^{\mathbf{V}}$ is not null, and in $\mathbf{V}^{\mathbb{Q}_0*\mathbb{Q}_1}$, $(^{\omega}2)^{\mathbf{V}^{\mathbb{Q}_0}}$ is not null. But for nep forcing notions preserving the non-nullity of $^{\omega}2$ implies preserving the non-nullity of any old non-null set by [107, §7]. There is no problem with successor stages. So now assume $\delta = \ell g(\bar{\mathbb{Q}})$ is a limit ordinal of countable cofinality.

As \mathbb{P}_{δ} is nep it is enough to assume

(**) $p_0 \Vdash_{P_\delta}$ " $\tilde{T} \subseteq {}^{\omega}>2$ is a subtree and $\operatorname{Leb}\left(\operatorname{lim}(\tilde{T})\right) > 0$ ", and $\delta = \bigcup_{n < \omega} i_n$, $i_n < i_{n+1}$, and without loss of generality $\tilde{T} \cap {}^{n \geq 2}$ is a hereditarily countable \mathbb{P}_{i_n} -name.

and to find an old $\eta \in {}^{\omega}2$ such that $p_0 \nVdash_{\mathbb{P}_{\delta}}$ " $\eta \notin \lim(T)$ ". Now, let N be a \mathbb{Q} -candidate to which $\{T, p_0, \langle i_n : n < \omega \rangle\}$ belongs (without loss of generality $N \prec (\mathcal{H}(\chi), \in)$, with χ large enough). Let $G_0 \subseteq \operatorname{Levy}(\aleph_0, (2^{\mathbb{P}_{\delta}})^N)$ be generic over N. Let $p_1 \in N[G_0]$ be such that $p_1 \in (\mathbb{P}_{\delta})^{N[G_0]}$, $N[G_0] \models$ " $p_0 \leq p_1$ and p_1 is explicitly (N, \mathbb{P}_{δ}) -generic". Let $G_1 = G_{1,\delta} \subseteq \mathbb{P}_{\delta}^{N[G_0]}$ be generic over $N[G_0]$ such that $p_1 \in G_{1,\delta}$ and let $G_{1,i_n} = G_1 \cap \mathbb{P}_{i_n}^{N[G_0]}$. Let s be a random real over $N[G_0][G_1]$ (if we replace random by other nep-explicitly demand even more models), hence over N too. Clearly we can choose such $G_0, G_{1,\delta}$ in \mathbb{V} . So N[s] is a class (= definable) of $N[G_0][G_1][s]$, and clearly N[s] is a \mathbb{Q} -candidate. Also, there is $s' \in \lim(T[G_1])$, $s' \equiv s$ (i.e., $s' \in {}^{\omega}2$ and the set $\{\ell < \omega : s'(\ell) \neq s(\ell)\}$ is finite). Let us define ψ as

$$\psi = p \ \& \ \bigwedge_{n < \omega} \left[s' \upharpoonright n \in (\underline{T} \cap {}^n 2)[\underline{\mathcal{G}}_{\mathbb{P}_{i_n}}] \right]$$

(note that, by the assumption, there is $\langle \psi_{n,\eta} : n < \omega, \eta \in {}^{n}2 \rangle$, $\psi_{n,\eta}$ an $L_{\omega_{1},\omega}$ -sentence for $\mathbb{P}_{i_{n}}$, i.e. using (countably many) variables $q \in \mathbb{P}_{i_{n}}$, such that

$$p_0 \Vdash$$
 " $\eta \in T \cap {}^n 2$ iff $\psi_{n,\eta} \in G_{P_{i_n}}$ "

for $n < \omega$, $\eta \in {}^{n}2$, so, up to equivalence,

$$\psi = p \& \bigwedge \psi_{n,s' \upharpoonright n}$$

(recall that $T \cap n^2$ is a \mathbb{P}_{i_n} -name). The problem is whether $\psi \in \mathbb{P}_{\delta}$. Now, $\psi \in N[s]$, so by absoluteness to show $\psi \in \mathbb{P}_{\delta}$ it suffices to show that this holds in N[s].

So we need

$$N[s] \models \text{``} \psi \ \& \ \bigwedge_n \Psi_{\mathbb{P}_{i_n}} \text{ has a Boolean-valued model "},$$

where $\Psi_{\mathbb{P}_{i_n}} = (\bigwedge \{\bigvee_{q \in \mathcal{I}} q : \mathcal{I} \subseteq \mathbb{P}_{i_n} \text{ is predense } \})^{N[s]}$. But $N[G_1][G_2][s]$ is a generic extension of N[s], so it is enough to prove it there; so there is no problem. In fact for the case of "non-null" the answer is yes (for CS iterations use [164, Ch.XVIII,3.8, pp 912–916]).

On the other hand, in full generality the answer to 4.31 is no; note that life is harder if we want to preserve positiveness for $I_{(\mathbb{Q},\underline{\eta})}$, where \mathbb{Q} is (nep but) not c.c.c., on this and more see [108].

Another possible direction is

Problem 4.32. 1. Is there an interesting theory of nicely definable forcing notions in $\mathcal{H}(\theta)$ or $\mathcal{H}_{<\theta}(\sigma)$?

- 2. Similarly for the theory of iterations (see later).
- 3. Generalize [85], replacing \aleph_0 by κ . Probably require $\kappa = \kappa^{<\kappa} > \aleph_0$ or even that κ is strongly inaccessible.

E.g., let for simplicity κ be strongly inaccessible, D^* a normal filter on κ . For a cardinal $\theta < \kappa$ let a θ -creature \mathfrak{c} consist of $(R^{\mathfrak{c}}, \operatorname{pos}^{\mathfrak{c}}, \operatorname{val}^{\mathfrak{c}})$, where: $R^{\mathfrak{c}}$ is a θ^+ -complete forcing notion, $\operatorname{pos}^{\mathfrak{c}}$ is a non-empty set, $\operatorname{val}^{\mathfrak{c}}$ is a function from $R^{\mathfrak{c}}$ to $\mathcal{P}(\operatorname{pos}^{\mathfrak{c}}) \setminus \{\emptyset\}$ such that $R^{\mathfrak{c}} \models \text{``} x \leq y\text{'`} \Rightarrow \operatorname{val}^{\mathfrak{c}}(y) \subseteq \operatorname{val}^{\mathfrak{c}}(x)$.

A κ -normed tree is $(T^*, \bar{\mathfrak{c}}, \bar{\theta})$, where $T^* \subseteq \kappa$ -Ord is a subtree, usually closed under increasing sequences of length $\langle \kappa, \bar{\mathfrak{c}} = \langle \mathfrak{c}_{\eta} : \eta \in T^* \rangle$, \mathfrak{c}_{η} is a θ_{η} -creature, $\bar{\theta} = \langle \theta_{\eta} : \eta \in T^* \rangle$ and $\operatorname{pos}^{\mathfrak{c}_{\eta}} = \operatorname{Suc}_{T^*}(\eta)$.

We can consider

```
\begin{array}{l} \mathbb{Q}^1_{(T^*,\bar{\mathfrak{c}}^*,\bar{\theta}),D^*} = \\ \mathbb{Q}^1 = \left\{ (T,\bar{r}) : (\mathbf{a}) \quad T \text{ is a subtree of } T^*, \ (<\kappa)\text{-closed}, \\ (\mathbf{b}) \quad \bar{r} = \langle r_\eta : \eta \in \operatorname{sp}(T)) \rangle, \ r_\eta \in R^{\mathfrak{c}_\eta} \text{ and } \operatorname{val}^{\mathfrak{c}_\eta}(r_\eta) = \operatorname{Suc}_T(\eta), \\ (\mathbf{c}) \quad \text{for every } \eta \in \lim_{\kappa} (T), \text{ we have } \{\zeta < \kappa \colon \eta \upharpoonright \zeta \in \operatorname{sp}(T)\} \in D^* \} \end{array}
```

with the natural order, where $\operatorname{sp}(T)=\{\eta\in T: (\exists^{\geq 2}x)[\eta^\frown\langle x\rangle\in T]\}$. Trivially, forcing with \mathbb{Q}^1 adds neither bounded subsets of κ nor sequences of ordinals of length $<\kappa$. For this forcing notion, if $T^*_\alpha=:\{\eta\in T^*:\ell g(\eta)=\alpha\}$ has cardinality $<\kappa$ for each $\alpha<\kappa$, and $(\forall\eta\in\lim_\kappa(T^*))[\{\zeta<\kappa:\theta_{\eta}\mid\zeta>\zeta\}\in D^*]$ (or there is $A\in D^*$ such that $\zeta\in A$ & $\eta\in T$ & $\ell g(\eta)=\zeta$ \Rightarrow $\theta_\eta>\zeta$), then forcing with $\mathbb{Q}^1_{(T^*,\bar{\mathfrak{c}}^*,\bar{\theta}),D^*}$ does not collapse κ^+ . Also, this forcing is $(<\kappa)$ —complete κ -bounding and \mathbb{Q}^1 —names of $\tau:\kappa$ — Ord can be read continuously on T for a dense set of $(T,\bar{r})\in\mathbb{Q}^1$. Moreover, this property is preserved by $(\leq\kappa)$ —support iterations (by [104, §1], or directly).

We can allow gluing (i.e., putting together with η many nodes above, creating a new forcing notion, i.e. creatures, see [85, 3.3(2),§6.3]).

We may consider $\bar{\mathfrak{c}} = \langle \bar{\mathfrak{c}}_{\zeta} : \zeta < \kappa \rangle$, \mathfrak{c}_{ζ} a θ_{ζ} -creature and for some club E of κ , $\zeta \in E \& \xi \geq \zeta \quad \Rightarrow \quad \theta_{\xi} > \zeta$, and consider

$$\mathbb{Q}^2_{\bar{\mathfrak{c}}} = \left\{ (w, \bar{r}) : \text{for some } \zeta, \ w \in \prod_{\varepsilon < \zeta} \mathrm{pos}^{\mathfrak{c}_{\varepsilon}}, \ \bar{r} = \langle r_{\varepsilon} : \varepsilon \in [\zeta, \kappa] \rangle, \ r_{\varepsilon} \in R^{\mathfrak{c}_{\varepsilon}} \right\}$$

with the natural order. If $\zeta < \kappa \implies |R^{\mathfrak{c}_{\zeta}}| + |\operatorname{pos}^{\mathfrak{c}_{\zeta}}| < \kappa$ and

$$(\alpha) \ \kappa = \sup\{\zeta : \text{for } \xi \in [\zeta,\kappa), \ R^{\mathfrak{e}_{\xi}} \text{ is } | \prod_{\varepsilon < \zeta} \text{pos}^{\mathfrak{e}_{\varepsilon}}|^{+} - \text{complete}\}, \text{ or }$$

$$(\beta) \diamondsuit_{\kappa}$$

then forcing does not collapse κ^+ and is $(<\kappa)$ -complete but the "read continuously" is problematic for case (β) .

Again, we may allow the forcing notion to be omittory (see [85, 2.1.1]), i.e., allow

$$\mathbb{Q}^3_{\bar{\mathfrak{c}}} = \{(w,\bar{r}): \text{ for some bounded } u \subseteq \kappa \text{ and } A \in D, \\ w \in \prod_{\varepsilon \in u} \mathrm{pos}^{\mathfrak{c}_\varepsilon}, \text{ and } \bar{r} = \langle r_\alpha : \alpha \in A \rangle \}$$

and/or a combination of creatures (i.e., the function Σ) and/or we may allow memory (=the object which \mathfrak{c}_{α} produces depends on the earlier \mathfrak{c}_{β} 's) and/or gluing.

Note that natural nice enough \mathfrak{c} 's make us regain "read continuously" and its parallels.

All those generalize naturally. But if in $\mathbb{Q}^1_{(T^*,\bar{\varsigma},\theta),D^*}$ we allow D^* to be the co-bounded filter, the proofs fail. There is more to be said and done.

Of course, we can now carry out generalizations of various independence results on cardinal invariants, e.g. between variants of (f,g)-bounding and slaloms (= corsets), see [151], [164, Ch.VI,§2], Goldstern and Shelah [46], Rosłanowski and Shelah [87]. We may consider in the tree version for every limit $\delta \in S$ ($S \subseteq \kappa$ is a stationary set) to omit tops for one or just $\leq |\delta|$ branches in a condition $p = (T, \bar{r})$, provided for each $\eta \in T$ we have $\langle |\{\rho : \eta \lhd \rho \in T, \ \ell g(\rho) = \ell g(\eta) + \alpha\}| : \alpha < \kappa \rangle$ goes to κ .

Recall that whereas Cohen forcing and many others have κ -parallels, not so with random real forcing.

Problem 4.33. 1. Prove that there is no reasonable parallel, say no κ^+ -c.c. forcing notion such that any new member of κ is bounded by some old one.

2. Similarly as far as the parallel of [156] is concerned.

Concerning nep forcing notions, we may define when " \mathbb{Q} is semi-nep" as follows: for some nep forcing notion \mathbb{Q}^* we have

- (a) $\mathbb{Q} \subseteq \mathbb{Q}^*$ (that is the set of elements of \mathbb{Q} is a subset of the set of elements of \mathbb{Q}^* and the order of \mathbb{Q} is the order of \mathbb{Q}^* restricted to \mathbb{Q}),
- (b) if N is a Q-candidate, $\mathbb{S} \in N$, $\mathbb{S}^N = \mathbb{Q}^* \cap N$, $p \in \mathbb{S}$, $p \in N$ then for some q, $p \leq q$, q is (N, \mathbb{S}) -generic (in an explicit way).

5. To prove or to force, this is the question

On many things we confidently "know" that they are independent, "just" a proof is needed (for many others we know that forcing will not help by absoluteness). The rest we may think are decidable, but actually we are not sure. More fall in the middle; our intuitions do not give an answer or worse: they give an answer which oscillates in time. In the mid-seventies, I was interested in (see Abraham and Shelah [3]):

Question 5.1. Is there an Aronszajn tree T and a function $c: T \longrightarrow \{\text{red}, \text{ green}\}$ (or more colours) such that for any uncountable set $A \subseteq T$, all colors appear on the set $\{\eta \cap \nu : \eta, \nu \in A \text{ are incomparable}\}$, where $\eta \cap \nu$ is the maximal common lower bound?

Why? Baumgartner [11] proved that among uncountable real linear orders (e.g., with density \aleph_0) there may be a minimal one under embeddability. This follows from

CON(if
$$A, B \in [\mathbb{R}]^{\aleph_1}$$
 are \aleph_1 -dense, then they are isomorphic).

So we may ask: among uncountable linear orders, can there be finitely many such that any other embeds one of them? (call such a family a base).

A base should contain a real order and ω_1 and ω_1^* . Any linear order into which none of them embeds is necessarily a Specker order (= take an Aronszajn tree, order it lexicographically).

You may ask: Can there be a "minimal" order among those? But there cannot. It is known ([124], answering a question of Countryman) that there is a Specker order L such that the product $L \times L$ (with the product order) is the union of countably many chains (comes from a very special Aronszajn tree). Hence L and L^* (its inverse) embed no common uncountable chain.

Now, consistently any two candidates for L are isomorphic or anti-isomorphic. So, if we can also put the additional forcing together, then we will have a candidate for a basis which seems extremely likely. We just need to start with a Specker order, i.e., an Aronszajn tree with a lexicographic order, under the circumstances (okay to force a little), without loss of generality with any node having two immediate successors. Look at it as an Aronszajn tree, and remember Abraham and Shelah [3]. So without loss of generality, on a club the tree is isomorphic to the one for L, and let Dom(c) = T,

$$c(\eta) = \begin{cases} & \text{green} & \text{if the two linear orders make the same} \\ & & \text{decision about the two immediate successors of } \eta, \\ & & \text{red} & & \text{otherwise}, \end{cases}$$

and so the problem 5.1 arises.

A common property of some of the problems discussed below (5.2, 5.3, 5.4) is a difference between asking about $S_{\theta}^{\lambda} = \{\delta < \lambda : \mathrm{cf}(\delta) = \theta\}$ and asking about a stationary $S \subseteq S_{\theta}^{\lambda}$ such that $S_{\theta}^{\lambda} \setminus S$ is stationary too; a difference of which I became aware in [126] (e.g. \diamondsuit_{S_1} & $\neg \mathring{\diamondsuit}_{S_2}$ is possible for disjoint stationary subsets of ω_1) after much agony.

Question 5.2 (GCH). If μ is singular, do we have $\diamondsuit_{S_{\mathfrak{g}}^{\mu^+}}$? (Those are the only cases left.)

Similarly for inaccessibles, see [110].

If we try to force consistency of the negation, note that (for μ strong limit singular)

$$2^{\mu} = \mu^{+} + \square_{\mu^{+}} \quad \Rightarrow \quad \diamondsuit_{S_{\mathrm{cf}(\mu)}^{\mu^{+}}} \qquad \text{when } \mathrm{cf}(\mu) > \aleph_{0}$$

(see [137, 3.2,p.1030]). So we need large cardinals (hardly surprising for successors of singular cardinals). See more in Džamonja and Shelah [30]).

Probably it is wiser to try to force this for "large" μ . Changing the cofinality of a supercompact cardinal μ to \aleph_0 , where " μ is prepared" is not helpful as after the forcing some old $S_{\aleph_0}^{\mu^+}$ is added to the old $S_{\mu}^{\mu^+}$ to make the new $S_{\aleph_0}^{\mu^+}$.

If $\mathbf{V} \models \neg \diamondsuit_{S_{\mu}^{\mu^+}}$ (which may be forced but you need enough indestructibility of

measure), you get $\neg \diamondsuit_S$ for some non-reflecting stationary $S \subseteq S_{\aleph_0}^{\mu^+}$, but we have it "cheaply" by forcing (say, starting with **L**; see [137], better [111]). So maybe it is wiser to start with μ of cofinality \aleph_0 .

Let μ be a limit of large cardinals and try to add enough subsets to μ to "kill" $\Diamond_{\mathrm{cf}(\mu)}^{\mu^{\top}}$. Our knowledge of such forcing for such cases is limited at the present. But ZFC + GCH still give an approximation (see more [111]):

- (*) if $S \subseteq S_{\mathrm{cf}(\mu)}^{\mu^+}$ is stationary then for some $\langle \langle \alpha_{\delta,i} : i < \mathrm{cf}(\mu) \rangle : \delta \in S \rangle$ we have: (a) $\alpha_{\delta,i}$ is increasing with limit δ ,

 - (b) if $\theta < \mu$, $f: \mu^+ \longrightarrow \theta$, then

 $(\exists \text{ stationarily many } \delta \in S)(\forall i < \text{cf}(\mu))(f(\alpha_{\delta,2i}) = f(\alpha_{\delta,2i+1})).$

Note: Having two equal values inside a group calls for division (or subtraction) so that we get a known value. So, if we are trying to guess homomorphisms from a group G with $|G| = \mu^+$ to H, $|H| = \theta < \mu$, $G = \bigcup_{i < \mu^+} G_i$, G_i strictly increasing

continuous, $|G_i| < \mu^+$, and $S \subseteq S_{\mathrm{cf}(\mu)}^{\mu^+}$ is stationary, then we can find $\bar{\eta} = \langle \langle g_{\delta,i} : i < \mathrm{cf}(\mu) \rangle : \delta \in S \rangle$ such that $g_{\delta,i} \in G_{\alpha_{\delta,2i+2}} \setminus G_{\alpha_{\delta,2i}}$ and for every homomorphism $h: G \longrightarrow H$ there are stationarily many δ such that $(\bigwedge_i h(g_{\delta,i}) = e_H)$ (relevant

to Whitehead groups). Without loss of generality, $|G_{i+1} \setminus G_i| = \mu$, $|G_0| = \mu$, the universe of G_i is $\mu \times (1+i)$ (or $g_i \in G_{i+1} \setminus G_i$ uses the question on $f: f(i) = f(g_i)$). See [111].

QUESTION: Can we have something similar for any sequence $\bar{\eta}$?

Answer: No. We have quite a bit of freedom (e.g., demand $\alpha_{\gamma,i} \in A^*$, $A^* \in [\mu^+]^{\mu^+}$ fixed) but certainly not for any.

In fact, for any stationary non-reflecting set $S \subseteq S_{\mathrm{cf}(\mu)}^{\mu^+}$ and any sequence $\bar{\eta} = \langle \langle \alpha_{\delta,i} : i < \mathrm{cf}(\mu) \rangle : \delta \in S \rangle$ with $\alpha_{\delta,i}$ increasing with limit δ we can force:

for every
$$\langle h_{\delta} \in {}^{\mathrm{cf}(\mu)}\theta : \delta \in S \rangle$$
, $\theta < \mu$, there is $h \in {}^{(\mu^+)}\theta$ such that $(\forall \delta \in S)(\forall^{\infty}i)(h(\alpha_{\delta,i}) = h_{\delta}(i))$.

This is a strong negation of the earlier statements (see [137], [111]). A related ZFC result is that for a singular cardinal μ , the restriction of the club filter $\mathcal{D}_{\mu^+} \upharpoonright S_{\mathrm{cf}(\mu)}^{\mu^+}$ is not μ^{++} -saturated (see Gitik and Shelah [45]).

A well known problem is

Question 5.3. 1. For a regular cardinal $\lambda > \aleph_2$, can $\mathcal{D}_{\lambda^+} \upharpoonright S_{\lambda}^{\lambda^+}$ be λ^{++} saturated?

2. Similarly just adding the assumption GCH.

* * *

Many "club guessings" are true (see [95]), but I have looked in vain several times on:

Question 5.4. Let λ be a regular uncountable cardinal. Can we find a sequence $\langle \langle \alpha_{\delta,i} : i < \lambda \rangle : \delta \in S_{\lambda}^{\lambda^+} \rangle$ such that $\alpha_{\delta,i}$ are increasing continuous in i with limit δ , and for every club E of λ^+

 $(\exists \text{ stationarily many } \delta)(\exists \text{ stationarily many } i)(\alpha_{\delta,i+1}, \alpha_{\delta,i+2} \in E),$

or other variants (just $\alpha_{\delta,i+1} \in E$ is provable, $\alpha_{\delta,i} \in E$ is trivial under the circumstances).

This is interesting even under GCH, particularly as by Kojman and Shelah [68] (essentially) we get from it that there is a λ^+ -Souslin tree.

We may think instead of trying to prove for $S, S_1 \subseteq S_{\lambda}^{\lambda^+}$ being stationary disjoint, that we can force the failure for S (with GCH). This works (see [110]) but $S = S_{\lambda}^{\lambda^+}$ is harder. The present forcing proofs fail, but also using "first counterexample" fails. We may consider proving: GCH $\Rightarrow \neg$ GSH (where GSH is Generalized Souslin Hypothesis). Let us look at two successor cases λ^+, λ^{++} (λ regular). How can this help? Assume that there is no λ^{++} -Souslin tree and GCH holds. It follows

that every stationary $S \subseteq S_{\leq \lambda}^{\lambda^{++}}$ reflects in $S_{\lambda^{+}}^{\lambda^{++}}$ (see Gregory [48]), moreover it is enough to assume just that there is no $(<\lambda^{+})$ -complete Souslin tree, by Kojman and Shelah [68]. Hence

- (\otimes) there is $S^* \subseteq \lambda^+$ such that:
 - (a) (square on S^*) $C = \langle C_{\alpha} : \alpha \in S^* \rangle, C_{\alpha} \subseteq \alpha \text{ closed, otp}(C_{\alpha}) \leq \lambda,$ if α is limit then $\sup(C_{\alpha}) = \alpha$, and $\beta \in C_{\alpha} \Rightarrow C_{\beta} = \beta \cap C_{\alpha},$ (b) $S^* \cap S_{\lambda}^{\lambda^+}$ is stationary.

(Why? This follows by [148, §4] which says that, e.g., $S_{\lambda^{+}}^{\lambda^{++}}$ is the union of λ^{+} sets with squares.) This looks as if it should help, but I did not find yet how.

Džamonja and Shelah [30] introduced

Definition 5.5. We say that λ strongly reflects at θ if $\theta < \lambda$ are regular uncountable cardinals and for some $F: \lambda \longrightarrow \theta$ for every $\delta \in S^{\lambda}_{\theta}$ for some club C of $\delta, F \upharpoonright \delta$ is strictly increasing (equivalently, is one-to-one).

This helps to prove variants of \clubsuit on the critical stationary subset of μ^+ when μ is singular, i.e., on $S_{\mathrm{cf}(\mu)}^{\mu^+}$, see Džamonja and Shelah [30], [29], and on independence results Cummings, Džamonja and Shelah [21] and Džamonja and Shelah [28].

Question 5.6. Can we get something parallel when $cf(\mu) = \aleph_0$?

Question 5.7. Can we prove that for some strong limit singular cardinal μ and a regular cardinal $\theta < \mu$ we have \clubsuit_S , where $S = S_{\theta}^{\mu^+}$?

On $I[\lambda]$ see [128], [141], [152]. We know that e.g.

$$\{\delta < \aleph_{\omega+1} : \operatorname{cf}(\delta) = \aleph_1\} \notin I[\aleph_{\omega+1}]$$

is consistent with GCH, but

Problem 5.8. 1. Can
$$\{\delta < \aleph_{\omega+1} : \text{cf}(\delta) = \aleph_2\} \notin I[\aleph_{\omega+1}]$$
?
2. Can $\{\delta < (2^{\aleph_0})^{+\omega+1} : \text{cf}(\delta) = (2^{\aleph_0})^+\} \notin I[(2^{\aleph_0})^{+\omega+1}]$?

Now [148, §4], dealing with successors of regulars, raises the question

- **stion 5.9.** 1. Let λ be inaccessible $> \aleph_0$. Is I_{λ}^{sq} (see Definition 5.10 below) non-trivial, i.e., does it include stationary sets of cofinality $\sigma \in (\aleph_0, \lambda) \cap \text{Reg}$? Does it include such S which is large in some sense (e.g., for every such σ)?
 - 2. Similarly for successor of singular.

Definition 5.10. For a regular cardinal $\lambda > \aleph_0$ let

$$I_{\lambda}^{\mathrm{sq}} = \big\{ A \subseteq \lambda : \quad \text{for some partial square } \bar{C} = \langle C_{\delta} : \delta \in S^1 \rangle, \\ S^1 \subseteq \lambda \text{ and the set } A \setminus S^1 \text{ is not stationary in } \lambda \big\}.$$

Definition 5.11. A linear order I is μ -entangled if: for any pairwise distinct $t_{i,\ell} \in I$, $i < \mu$, $\ell < n$, for any $w \subseteq \{0, \ldots, n-1\}$ there are i < j such that

$$t_{i,\ell} < t_{j,\ell} \quad \Leftrightarrow \quad \ell \in w.$$

If $|I| = \mu$ then we omit μ .

Question 5.12. Is there an entangled linear order of cardinality λ^+ , where $\lambda = \lambda^{\aleph_0}$?

A "yes" answer will solve a problem of Monk [80] on the spread of ultraproducts of Boolean Algebras; see [162].

With the help of pcf we can build entangled linear orders in λ^+ for many λ which means: provably for a proper class of λ 's.

The interesting phenomenon is: from instances of GCH, we can give a positive answer, and if $\mu^{\aleph_0} = \mu$, $2^{\mu} > \aleph_{\mu^{+4}}$ we have: for many $\delta < \mu^{+4}$ we get a positive answer to 5.12 with $\lambda = \aleph_{\delta}$. On the other hand, from $\mu = \mu^{\aleph_0}$, $2^{\mu} < \mu^{+\omega}$ we can also prove a positive answer.

In fact, in the remaining case there are quite heavy restrictions on pcf. A typical universe with negative answer to 5.12 (we think that it) will satisfy: for a strong limit cardinal μ , $2^{\mu} = \mu^{+\omega+1}$, and for $\mathfrak{a} \subseteq \operatorname{Reg} \cap \mu$, $\mu = \sup(\mathfrak{a})$, $\operatorname{pcf}(\mathfrak{a}) \setminus \mu$ essentially concentrates on $\mu^{+\omega+1}$ (say $\langle \mu_i : i < \operatorname{cf}(\mu) \rangle$ increasing continuous, if \mathfrak{a} is disjoint from $\{\mu_i^{+n} : i < \operatorname{cf}(\mu), \ 0 < n < \omega\}$ then $\emptyset = \operatorname{pcf}(\mathfrak{a}) \cap (\mu, \mu^{+\omega})$.) See [162].

Maybe our knowledge of forcing will advance. Note that we need not only to have pcf structure as indicated, but also to take care of the non-pcf phenomena as well for constructing entangled linear order as in 5.12.

Considering a ZFC proof of existence, it seems most reasonable to assume toward a contradiction that the answer is no and consider strong limit singular μ of uncountable cofinality. So we know $2^{\mu} < \aleph_{\mu^{+4}}$ and being more careful even $2^{\mu} < \aleph_{\mu^{+}}$. Let $\gamma(*) = \min\{\gamma: 2^{(\mu^{+\gamma})} > 2^{\mu}\}$, so necessarily $\gamma(*)$ is a successor ordinal, say $\gamma(*) = \beta(*) + 1$. Let $\lambda =: \mu^{+\beta(*)}$. We may consider trying to construct an entangled linear order of cardinality $(2^{\mu})^{+}$, using the weak diamond on $\lambda^{+} = \mu^{+\beta(*)+1}$. Moreover, we know that there are trees T with λ^{+} levels and $\leq \lambda^{+}$ nodes and at least $(2^{\mu})^{+}$ many λ^{+} -branches (even $\lambda^{+}2 = \bigcup_{\zeta < 2^{\mu}} \lim_{\lambda^{+}} (T_{\zeta})$ for some subtrees T_{ζ} of

$$^{\lambda^{+}>}2$$
, $|T_{\zeta}| = \lambda^{+}$ above).

Moreover, a relative of $\diamondsuit_{\lambda^+}^*$ holds. All this seems reasonably promising, but has failed so far to solve the problem.

I have also considered to repeat the proof of the weak diamond for λ^+ to try to show that a tree with infinite splitting in the above representation is necessary.

Problem 5.13. Can we prove that a stronger version of the weak diamond holds for some λ ? E.g., a version with more than two colours and/or fixing the cofinality. We shall be glad to get even just the definable weak diamond. See [139], [97, §3], [105] and [101].

* * *

Our ignorance about such problems may well come from our gaps in forcing theory.

A major problem (more exactly a series of problems) is

Problem 5.14. 1. Can we have a reasonable theory of iterations (and/or forcing axioms) for $(\langle \lambda \rangle)$ -complete forcing notions $(\lambda = \lambda^{\langle \lambda \rangle})$?

- 2. Similarly for forcing notions not changing cofinalities of cardinals $< \lambda$?
- 3. Similarly for forcing notions preserving μ^+ and not adding bounded subsets to μ , μ a strong limit singular cardinal?

See some recent information on the first in [110], [104], [111], and even much less on the second [164, Ch.XIV], and on the third Mekler and Shelah [76], Džamonja and Shelah [27].

Though much was done on forcing for the function 2^{λ} and some specific problems, our flexibility is not as good as for 2^{\aleph_0} in forcing theory.

Particularly intriguing are solutions where we know some λ exists but do not know which. The dual problem is iterated forcing of length Ord (class forcing); now for such iteration it is particularly hard to control in the neighborhood of singulars.

Problem 5.15. Prove the consistency of: for every λ (or regular λ) a suitable forcing axiom holds.

Relevant is "GCH fails everywhere" (see Foreman and Woodin [35]). Now Cummings and Shelah [23], [24] is a modest try and 1.22 is relevant.

Specific well known targets are

Problem 5.16. Is GSH consistent? (GSH is the generalized Souslin hypothesis: for every regular uncountable λ there is no λ -Souslin tree.)

Problem 5.17. Is it consistent that for no regular $\lambda > \aleph_1$ do we have a λ -Aronszajn tree (see Abraham [1], Cummings and Foreman [22]).

A relevant problem is 6.4.

I have found partition theorems on trees with ω levels very useful and interesting (see Rubin and Shelah [89], and [134], [132], [164, X, XI, XV,2.6]). In [145, 13,p.1453] and [127, Ch.VIII,§1] trying to prove a theorem on the number of nonisomorphic models of a pseudo elementary class we arrived at the following problem [without loss of generality, try with $2^{\lambda} = \lambda^{+}$ and see [116], by absoluteness]:

Question 5.18. Assume $m(*) < \omega$, $2^{\lambda_n} < \lambda_{n+1}$, M is a model with vocabulary of cardinality θ , $\theta + \mu < \lambda_0$, $a_\eta^i \in M$ for $i < \mu$, $\eta \in T = \bigcup_n \prod_{\ell < n} \lambda_\ell$. Can we find a strictly increasing function $h:\omega\longrightarrow\omega$ and one-to-one functions $f_n^i:\prod_{k< n}\lambda_k\longrightarrow\prod_{k< h(n)}\lambda_k$ such that

- (a) for $n < m, \eta \in \prod_{\ell < m} \lambda_{\ell}$ we have $f_n^i(\eta \upharpoonright \ell) = (f_m^i(\eta)) \upharpoonright h(\ell)$, (b) for $n < \omega, m(*) < \omega, i_0 < \ldots < i_{m(*)-1} < \mu \text{ and } \eta_{\ell}, \nu_{\ell} \in \prod_{k < n} \lambda_k \text{ for } \ell < m(*), \text{ the tuples } \langle a_{\eta_0}^{i_0}, a_{\eta_1}^{i_1}, \ldots, a_{\eta_{m(*)-1}}^{i_{m(*)-1}} \rangle, \langle a_{\nu_0}^{i_0}, a_{\nu_1}^{i_1}, \ldots, a_{\nu_{m(*)-1}}^{i_{m(*)-1}} \rangle \text{ realize the same type in } M^2$ the same type in M?

(If on λ_n there is a "large ideal" (see [159]) life is easier, see [145].)

6. Boolean Algebras and Iterated Forcing

We turn to Boolean Algebras. Monk has made extensive lists of problems about Boolean Algebras (which inspired more than few works of mine). His problems mostly go systematically over all possible relations; our perspective is somewhat different.

Among my results on Boolean Algebras I like 6.1 stated below (see [140]), but the result did not draw much attention though the paper was noticed (see Bonnet and Monk [16], Juhász [61]).

Theorem 6.1. If B is a Boolean Algebra of cardinality $\geq \beth_{\omega}$ and $\lambda = \operatorname{Id}(B)$ (the number of ideals of the Boolean Algebra) then $\lambda = \lambda^{\beth_{\omega}}$.

(We can instead of ideals of Boolean Algebras speak about open subsets of a compact Hausdorff topology, and we can replace \beth_{ω} by any singular strong limit).

So we are left with

Question 6.2. Is it true that for any large enough Boolean Algebra B we have $\mathrm{Id}(B)=\mathrm{Id}(B)^{<\theta}$ when, e.g., $\theta=\log_2(|B|)$, or at least for some constant $n,\ \theta=\min\{\mu: \beth_n(\mu)\geq |B|\}$?

(Similarly for compact spaces).

By [140], for every B there is such n. Of course, in non-specially constructed universes the answer is yes. If you like to try consistency, you have to use the phenomena proved consistent in Gitik and Shelah [43].

On the other hand, a ZFC proof may go in a different way than [140]. Related (see Juhász [60]) is

Question 6.3. What can be the number of open sets of a T_2 topology? T_3 topology? One with clopen basis?

It seems interesting to consider the following

Problem 6.4. Is there a class of cardinals λ (or just two) such that there is a (λ^+, λ) -thin tall superatomic Boolean Algebra B (i.e., $|B| = \lambda^+$, B is superatomic and for every $\alpha < \lambda^+$, B has $\leq \lambda$ atoms of order α), provably in ZFC?

It is well known that if $\lambda = \lambda^{<\lambda}$ then there is a (λ^+, λ) -thin tall superatomic Boolean Algebra, so for $\lambda = \aleph_0$ there is one, so for negative consistency we need "GCH fails everywhere or at least for every large enough λ ". Also note that trivially if there is a λ^+ -tree (i.e., one with λ^+ levels each of cardinality $\leq \lambda$), then there is such (λ^+, λ) -thin tall superatomic Boolean Algebra.

The point is that for several problems in Monk [80]: Problems 72, 74, 75 and ZFC versions of Problems 73, 77, 78, 79 (all solved in Roslanowski and Shelah [84] in the original version, i.e. showing consistency) there is no point to try to get positive answers as long as we do not know it for 6.4.

Also for several problems of [80] (Problems 49, 57, 58, 61, 63, 87) there is no point to try to get consistency of non-existence as long as we have not proved the consistency of the GSH (generalized Souslin hypothesis) which says there are no λ^+ -Souslin trees or there is no λ -Souslin tree for $\lambda = cf(\lambda) > \aleph_0$. For some others this is not provable, but it still seems very advisable to wait for the resolution of GSH

Problem 6.5. Usually the question on cardinal invariants \mathbf{inv}_1 , \mathbf{inv}_2 is "do we always have $\mathbf{inv}_1(B) \leq \mathbf{inv}_2(B)$?", or "do we always have $2^{\mathbf{inv}_1(B)} \geq \mathbf{inv}_2(B)$?"

But maybe there are relations like $\mathbf{inv}_1(B) \leq (\mathbf{inv}_2(B))^{+n}$ for some fixed $n < \omega$ (or for ω). A particularly suspicious case is $|B| \leq (\mathbf{irr}_n(B))^+$, where $n \in [2, \omega]$ and

 $\operatorname{irr}_n(B) = \sup \{|X|: X \subseteq B \text{ is } n\text{-irredundant which means:}$ if m < 1+n and $a_0, \ldots, a_m \in X$ are pairwise distinct then a_0 is not in the subalgebra generated by $\{a_1, \ldots, a_m\}$.

(If $n = \omega$ we may omit it.)

I think that for $n = \omega$ the case $|B| = \aleph_2$ had appeared in an old list of Monk, but not in [80]. We may also ask $|B| \leq (\operatorname{irr}_n(B))^{+m}$ for $n, m \in [2, \omega]$.

Of course, the open case is when, say, $|B| = \lambda^{++}$, $2^{\lambda} \ge \lambda^{++}$. Thinking about this problem, I was sure the answer is consistently no (consistently yes is easy, even in the Easton model). Moreover, I feel I know how to do it: let $\lambda = \lambda^{<\lambda}$, $\lambda^+ < \theta < \mu$, \mathfrak{B} be a suitable algebra on μ with $\le \lambda$ functions. A member p of the forcing notion $\mathbb P$ consists of $w^p \in [\mu]^{<\lambda}$ and a Boolean algebra B^p generated by $\{x_i : i \in w^p\}$ but such that

if $B^p \models$ " $x_{i_0} = \sigma(x_{i_1}, \dots, x_{i_n})$ " for ordinals $i_0, \dots, i_n \in w^p$ and a Boolean term σ

then $i_0 \in c\ell_{\mathfrak{B}}(\{i_1,\ldots,i_n\})$ and possibly more

(the order of \mathbb{P} is the natural one). This is to reconcile the demand "the Boolean algebra has cardinality μ , so without loss of generality we have to ask $i < j \Rightarrow x_i \neq x_j$ " and the λ^+ -c.c. Of course, $\mathcal{B} = \bigcup \{B^p : p \in \mathcal{G}_{\mathbb{P}}\}$.

Note: if $\lambda = \aleph_0$ we have more freedom. (The expected proof goes: if

 $p \Vdash$ " $X \subseteq B$, $X = \{y_i : i < \theta\}$ with no repetition exemplifies $\operatorname{irr}(B)^+ \ge |B|$ "

then we can find $p_i, p \leq p_i \in \mathbb{P}, p_i \Vdash "y_i = \sigma_i(x_{\alpha(i,0)}, \dots, x_{\alpha(i,n_i)})$ ". Hence, if $(\forall \alpha < \theta)(|\alpha|^{\leq \lambda} < \theta = \operatorname{cf}(\theta))$, without loss of generality, $\sigma_i = \sigma, n_i = n^*$ and $\langle (p_i, \langle \alpha(i, \ell) : \ell < n^* \rangle) : i < \theta \rangle$ forms a Δ -system (moreover indiscernible as a sequence of sequences of ordinals of length $< \lambda$). Let $\{\alpha(i, \ell) : \ell < n^* \} \subseteq w^p = w^* \cup \{\gamma_{i,\zeta} : \zeta < \zeta^* \}, \zeta^* < \lambda$, etc, and we have to find $n < \omega_1, i_0 < \ldots < i_n < \theta$ and q above p_{i_0}, \ldots, p_{i_n} such that $B^q \models "y_{i_0} = \sigma^*(y_{i_1}, \ldots, y_{i_n})$ ". So it is natural to demand $\zeta < \zeta^* \Rightarrow \gamma_{i_0,\zeta} \in c\ell_{\mathfrak{B}}\{\gamma_{i_1,\zeta}, \ldots, \gamma_{i_n,\zeta}\}$. So if $\lambda = \aleph_0$ we may use $n > |w^{p_i}|$. But this approach has not converged to a proof.)

So (see Monk [80, Problem 28])

- **Question 6.6.** 1. Is there a class of (or just one) λ such that for some Boolean Algebra B of cardinality λ^+ we have $\operatorname{irr}(B) = \lambda$?
 - 2. Similarly for $irr_n(B)$.

Colouring theorems (e.g. [160]) are not enough for a construction.

Question 6.7. 1. For which pairs (λ, θ) of cardinals $\lambda \geq \theta$ is there a superatomic Boolean Algebra with $> \lambda$ elements, λ atoms and every $f \in \operatorname{Aut}(B)$ moves $< \theta$ atoms?

(That is $|\{x : B \models \text{``} x \text{ an atom and } f(x) \neq x \text{'`}\}| < \theta$).

- 2. In particular, is it true that for some θ , for a proper class of λ 's there is such Boolean Algebra?
- 3. Replace "automorphism" by "one-to-one endomorphism".

See some results in [99, $\S1,\S2$] for θ strong limit singular. (It may be interesting to try: with n depending on the arity of the term as in [99].)

Concerning attainment in ZFC:

- Question 6.8. 1. Can we show the distinction made between the attainments of variants of hL (and hd), in a semi-ZFC way? That is, in Rosłanowski and Shelah [83] such examples are forced. "Semi ZFC" means can we prove such examples exist after adding to ZFC only restrictions on cardinal arithmetic?
 - 2. Similarly for other consistency results. (Well, preferably of low consistency strength).

In view of [98], [116] it is reasonable to consider

Problem 6.9. In 6.5, 6.6 replace $\operatorname{irr}_n(B)$ by $\operatorname{irr}_n(\bar{a}, B)$ (this is true for other cardinal invariants as well), see Definition 6.10 below.

Definition 6.10. Let B be a Boolean Algebra and $\bar{a} = \langle a_i : i < \lambda \rangle$ be a sequence of elements of B.

- 1. $\operatorname{irr}_n(\bar{a}, B) = \sup\{|X| : X \subseteq \lambda \text{ and } \langle a_i : i \in X \rangle \text{ is } n\text{-irredundant}\}.$
- 2. Similarly for other invariants of the "universal family" from Rosłanowski and Shelah [82] (see Definition 1.1 there).

Question 6.11. Is there (at least consistently) a Boolean Algebra B^* , such that if B is a Boolean Algebra extending B^* then for some ultrafilters D_1, D_2 on B we have: $(B, D_1), (B, D_2)$ are not isomorphic, i.e., no automorphism of B maps D_1 onto D_2 .

A close topological relative is: "is there a homogeneous compact Hausdorff space of cellularity $> 2^{\aleph_0}$ " (van Douwen, see Kunen [72]).

* * *

There are some lemmas in [140] which help to prove 6.1, and I would like to know whether the bounds used there are the best possible. Those lemmas also show that for some cardinal invariants (for Boolean Algebras or topologies), defined by supremum, if the supremum is not attained, then the value is "almost" regular (the classical result of Erdös and Tarski on the cellularity on Boolean Algebra (or topology) says it is regular, whereas we get in [140] that, e.g., the spread satisfies $2^{\text{cf}(s^+(B))} > s^+(B)$, for a Boolean Algebra B, $2^{2^{\text{cf}(s^+(X))}} > s^+(X)$ for a Hausdorff space, $2^{\text{cf}(s^+(X))} > s^+(X)$ for a T_3 space T_3 .

Question 6.12. Can we find more applications of the theorems (and proofs) in [140] implying (or saying) that if the supremum in some cardinal invariants for a space \mathcal{U} (or a Boolean Algebra B, or whatever) is not attained, then it has large cofinality?

A recent application is in Rosłanowski and Shelah [84, $\S 6$]. This is a converse to 6.8.

For the spread (and the hereditarily Lindelof degree and the hereditarily density) the results are best possible (see Juhász and Shelah [62]), and for regular spaces we have better results ([140, 5.1] also best possible) but are the bounds in the claims below best possible?

Definition 6.13. 1. φ is nice for X if φ is a function from the family of subsets of the topological space X to cardinals satisfying

$$\varphi(A) \leq \varphi(A \cup B) \leq \varphi(A) + \varphi(B) + \aleph_0$$
 (i.e., monotonicity and subadditivity).

2. We say φ is (χ, μ) -complete provided that if $A_i \subseteq X, \varphi(A_i) < \chi$ for $i < \mu$ then $\varphi(\bigcup_{i < \mu} A_i) < \chi$.

Let $C(\varphi, \mu) = \{\chi : \varphi \text{ is } (\chi, \mu)\text{-complete}\}.$

- 3. We say φ is $(<\lambda,\mu)$ -complete, if for arbitrarily large $\chi<\lambda,\ \varphi$ is (χ,μ) -complete.
- 4. Let Ch_{φ} be the following function from X to cardinals:

$$Ch_{\varphi}(y) = \min\{\varphi(u) : y \in u \in \tau(X)\},\$$

where $\tau(X)$ is the topology of X, that is the family of open sets.

Remark 6.14. 1. We can replace μ by $<\mu$ and $i<\mu$ by $i<\alpha<\mu$, and make suitable changes later.

- 2. In our applications we can restrict the domain of φ to the Boolean Algebra generated by $\tau(X)$ and even more, e.g., in 6.15, 6.17 below to simple combinations of the $u_{i,\xi,\zeta}$.
- 3. We can change the definition of (< λ, μ)–complete to

if
$$A_i \subseteq X$$
 (for $i < \mu$) and $\sup_{i < \mu} \varphi(A_i) < \lambda$ then $\varphi(\bigcup_{i < \mu} A_i) < \lambda$,

without changing our subsequent use. [We then will use: if $\varphi(A_{\alpha}) < \chi_i$ for $\alpha < \mu$ then $\varphi(\bigcup_{\alpha < \mu} A_{\alpha}) < \chi_{i+1}$].

Lemma 6.15. Suppose λ is a singular cardinal of cofinality θ , $\lambda = \sum_{i < \theta} \chi_i$, $\chi_i < \lambda$,

 $\theta < \lambda$ and $\mu = \beth_5(\theta)^+$, or even just $\mu = \beth_2(\beth_2(\theta)^+)^+$. Assume that

- (i) φ is nice for X,
- (ii) $X_{\chi_i} = \{ y \in X : \operatorname{Ch}_{\varphi}(y) \ge \chi_i \}$ has cardinality $\ge \mu$ for $i < \theta$,
- (iii) φ is $(< \lambda, \mu)$ -complete.

Then there are open sets $u_i \subseteq X$ (for $i < \theta$) such that

$$\varphi(u_i \setminus \bigcup_{j \neq i} u_j) \ge \chi_i.$$

Remark 6.16. If $|\{y \in X : \operatorname{Ch}_{\varphi}(y) \geq \chi_i\}| < \mu$, it essentially follows from (χ_i, μ) -completeness that $\varphi(X_{\chi_i}) \geq \lambda$, where $X_{\chi} = \bigcup \{u \in \tau(X) : \varphi(u) < \chi\}$. Otherwise, $\varphi(X \setminus X_{\chi_i}) \geq \lambda$ by subadditivity, but $\varphi(X \setminus X_{\chi_i}) \leq \prod \{\varphi(\{y\}) : y \in X \setminus X_{\chi_i}\}$, so by (χ_i, μ) -completeness for some $y \in X$, $\varphi(\{y\}) \geq \chi_i$, which is impossible for the instances which interest us.

Lemma 6.17. Suppose that X is a Hausdorff space, λ is a singular cardinal, $\theta = \operatorname{cf}(\lambda)$, $\lambda = \sum_{i < \theta} \chi_i$, $\chi_i < \lambda$, $\mu < \lambda$ and clauses (i), (ii), (iii) of 6.15 hold (for φ).

- 1. If $\mu = \beth_2(\theta)^+$ (or even $\sum_{\sigma < \theta} \beth_2(\sigma)^+$), then there are open sets u_i (for $i < \theta$) such that $\varphi(u_i \setminus \bigcup_{j>i} u_j) \ge \chi_i$.
- 2. If $X = \bigcup \{u : \varphi(u) < \lambda\}$, and μ is as in part (1), then there are open sets u_i (for $i < \theta$) such that $\varphi(u_i \setminus \bigcup_{j \neq i} u_j) \ge \chi_i$.

3. If $\mu \geq \beth_3(2^{<\theta})^+$, φ is $(<\chi_0,\mu)$ -complete, then there are u_i (for $i < \theta$) such that $\varphi(u_i \setminus \bigcup_{j \neq i} u_j) \geq \chi_0$ (so λ , χ_i (0 < $i < \theta$) are irrelevant).

Remark 6.18. Part (1) of the lemma is suitable to deal with Boolean Algebras, part (2) with the existence of $\{x_{\alpha} : \alpha < \lambda\}$ such that for every $\alpha < \lambda$ for some u, $x_{\alpha} \in u \cap \{x_{\beta} : \beta < \lambda\} \subseteq \{x_{\beta} : \beta \leq \alpha\}$.

Now.

Question 6.19. Are the cardinal bounds in 6.14 — 6.17 best possible?

7. A TASTE OF ALGEBRA

I have much interest in Abelian groups, but better see Eklof and Mekler [31].

Thomas prefers to deal just with short elegant proofs of short elegant problems (for me the second demand suffices). So he was rightly happy when proving that for any infinite group G with no center, $\gamma(G) < (2^{|G|})^+$, where $G^{[0]} = G$, $G^{[1]}$ is the automorphism group of G considered as an extension of G, $G^{[i+1]} = (G^{[i]})^{[1]}$, $G^{[\delta]} = \bigcup_{i < \delta} G^{[i]}$, so $G^{[i]}$ is an increasing sequence of groups with no center, and $\gamma(G) = \min\{\gamma: G^{[\gamma]} = G^{[\gamma+1]}\}$.

But is there a better cardinal bound? No, for |G| regular $> \aleph_0$, see Just, Shelah and Thomas [64], but we are left with:

Question 7.1. If G is a countable group with a trivial center, then do we have $\gamma(G) < \omega_1$? What about singular |G|?

I heard about the following problem (see Hamkins [54]).

Problem 7.2. If G is a group possibly with center, $G^{[i]}$ is defined as above but we have just a homomorphism $h_{i+1,i}:G^{[i]}\longrightarrow G^{[i+1]}$ with the center of $G^{[i]}$ being the kernel (and in limit stages take the direct limit), is there a bound to $\gamma(G)$ really better than the first strongly inaccessible > |G| (gotten by Hamkins [54])?

Thomas also had started investigating cofinalities of some natural groups, (see Sharp and Thomas [91], [92], Thomas [177]). He drew me to it and I was particularly glad to see that pcf pops in naturally; e.g., (see Shelah and Thomas [170]) if $\lambda_n \in \mathrm{CF}(\mathrm{Sym}(\omega))$ and $\lambda \in \mathrm{pcf}\{\lambda_n : n < \omega\}$ then $\lambda \in \mathrm{CF}(\mathrm{Sym}(\omega))$, where

Definition 7.3. 1. $CF(G) = \{\theta : \theta = cf(\theta) \text{ and there is an increasing sequence of proper subgroups of } G \text{ of length } \theta \text{ with union } G\}.$

2.
$$\operatorname{cf}(G) = \min[\operatorname{CF}(G) \setminus \{\aleph_0\}].$$

Though we found some information about $\operatorname{cf}(\prod_n \operatorname{alt}(n))$ (see Saxl, Shelah and Thomas [166], where $\operatorname{alt}(n)$ is the group of even permutations of $\{0,\ldots,n-1\}$), we remained baffled by

Question 7.4. Is it consistent that $\aleph_2 \leq \operatorname{cf}(\prod_{n < \omega} \operatorname{alt}(n))$?

It is natural to try to use iterations of length ω_2 , where each iterand consists of trees with norms (see Rosłanowski and Shelah [85]). Naturally, a norm on $\mathcal{P}(\operatorname{alt}(n))$ will be such that if $\operatorname{nor}(A) \geq m+1$ and σ is a group term, then we can have for "many" $f_1, \ldots, f_k, g \in \operatorname{alt}(n)$ that $A' = \{h \in A : \sigma(h, f_1, \ldots, f_{k^*}) = g\}$ has

 $\mathbf{nor}(A') \geq m$ toward destroying a guess on an approximation to a lower subgroup exemplifying $\omega_1 \in \mathrm{CF}(\prod_{n < \omega} \mathrm{alt}(n))$. This helps for $\aleph_0 \notin \mathrm{CF}(\prod_{n < \omega} \mathrm{alt}(n))$, but fails for the purpose of 7.4.

* * *

My interest in lifting for the measure algebra started when Talagrand promised me "flowers on your grave from every measure theorist" (a little hard to check), and Fremlin said essentially the same, conventionally (see [135], [164, Ch.IV]). But this does not solve some other problems from Fremlin's list, from which I particularly like

Question 7.5. Assume CH (or even GCH or just prove consistency).

Do we have lifting for every measure algebra?

Which means: let $\mathcal{B}(I)$ be the algebra of subsets of I2 generated from clopen ones by countable unions and intersections, $\mu_{\mathcal{B}}$ the Lebesgue measure on \mathcal{B} (so we get the so-called Maharam algebra), $\mathbf{I} = \{A \in \mathcal{B} : \mu_{\mathcal{B}}(A) = 0\}$ (so \mathbf{I} is the ideal of null sets). A lifting is a homomorphism from \mathcal{B}/\mathbf{I} into \mathcal{B} such that

$$X \in \mathcal{B} \implies f(X/\mathbf{I}) = X \mod \mathbf{I}.$$

Naturally, we think $I_0 \subseteq I_1 \Rightarrow \mathcal{B}(I_0) \subseteq \mathcal{B}(I_1)$ (by identifying) and for an increasing sequence $\bar{I} = \langle I_\alpha : \alpha < \alpha^* \rangle$ we let $\mathcal{B}(\bar{I}) = \bigcup_{\alpha < \alpha^*} \mathcal{B}(I_\alpha)$. In the positive direction we may try to prove by induction on λ ; then we will be naturally drawn to proving: for any $\mathcal{P}^-(n)$ -diagram $\langle \mathcal{B}(\bar{I}_s) : s \in \mathcal{P}^-(n) \rangle$, where $\mathcal{P}^-(A) = \{u : u \subseteq A, u \neq A\}$, and a sequence of liftings $\bar{f} = \langle f_s : s \in \mathcal{P}^-(n) \rangle$ satisfying a reasonable induction hypothesis, $\bigcup_s f_s$ can be extended to a lifting of $\bigcup_s I_{s,\alpha}$ (as e.g. in [133], Sageev and Shelah [90]). For the negative direction we may think of using a partition theorem.

* * *

For a long time I have been interested in compactness in singular cardinals; i.e., whether if something occurs for "many" subsets of a singular λ of cardinality $< \lambda$, it occurs for λ . For the positive side (on the filters see 7.7 below)

Theorem 7.6. Let λ be a singular cardinal, $\chi^* < \lambda$. Assume that \mathbf{F} is a set of pairs (A, B) (written usually as B/A; \mathbf{F} stands for free) $A, B \subseteq \mathcal{U}$ satisfying the axioms II, III, IV, VI, VII below. Let $A^*, B^* \subseteq \mathcal{U}$, $|A^*| = \lambda$, then $B^*/A^* \in \mathbf{F}$ if B^*/A^* is λ -free in a weak sense which means (see Definition 7.7 below):

- (*)₀ for the $\mathcal{D}_{\chi^*}(B^*)$ -majority of $B \in [B^*]^{<\lambda}$ we have $B/A^* \in \mathbf{F}$, or just
- (*)₁ the set $\{\mu < \lambda : \{B \in [B^*]^{\mu} : B/A^* \in \mathbf{F}\} \in \mathcal{E}_{\mu}^{\mu^+}(B^*)\}$ contains a club of λ , or at least
- (*)₂ for some set C of cardinals $< \lambda$, unbounded in λ and closed (meaningful only if $cf(\lambda) > \aleph_0$), for every $\mu \in C$, for an $\mathcal{E}^{\mu^+}_{\mu}(B^*)$ -positive set of $B \in [B^*]^{\mu}$ we have $B/A^* \in \mathbf{F}$.

The axioms are

Ax II: $B/A \in \mathbf{F} \quad \Leftrightarrow \quad A \cup B/A \in \mathbf{F}$,

Ax III: if $A \subseteq B \subseteq C$, $B/A \in \mathbf{F}$ and $C/B \in \mathbf{F}$ then $C/A \in \mathbf{F}$,

Ax IV: if $\langle A_i : i \leq \theta \rangle$ is increasing continuous, $\theta = \operatorname{cf}(\theta)$, $A_{i+1}/A_i \in \mathbf{F}$ then $A_{\theta}/A_0 \in \mathbf{F}$,

Ax VI: if $A/B \in \mathbf{F}$ then for the \mathcal{D}_{χ^*} -majority of $A' \subseteq A$, $A'/B \in \mathbf{F}$ (see below),

Ax VII: if $A/B \in \mathbf{F}$ then for the \mathcal{D}_{χ^*} -majority of $A' \subseteq A$, $A/B \cup A' \in \mathbf{F}$.

(Of course we can get variants by putting more or less into the statement.)

- **Definition 7.7.** 1. Let \mathcal{D} be a function giving for any set B^* a filter $\mathcal{D}(B^*)$ on $\mathcal{P}(B^*)$ (or on $[B^*]^{\mu}$). Then to say "for the \mathcal{D} -majority of $B \subseteq B^*$ (or $B \in [B^*]^{\mu}$) we have $\varphi(B)$ " means $\{B \subseteq B^* : \varphi(B)\} \in \mathcal{D}(B^*)$ (or $\{B \in [B^*]^{\mu} : \neg \varphi(B)\} = \emptyset \mod \mathcal{D}$).
 - 2. Let $\mathcal{D}_{\mu}(B^*)$ be the family of $Y \subseteq \mathcal{P}(B^*)$ such that for some algebra M with universe B^* and $\leq \mu$ functions,

 $Y\supseteq\{B\subseteq B^*:B\neq\emptyset\text{ is closed under the functions of }M\}.$

3. $\mathcal{E}_{\mu}^{\mu^+}(B^*)$ is the collection of all $Y \subseteq [B^*]^{\mu}$ such that: for some χ, x such that $\{B^*, x\} \in \mathcal{H}(\chi)$, if $\bar{M} = \langle M_i : i < \mu^+ \rangle$ is an increasing continuous sequence of elementary submodels of $(\mathcal{H}(\chi), \in)$ such that $x \in M_0$ and $\bar{M} \upharpoonright (i+1) \in M_{i+1}$, then for some club C of μ^+ , $i \in C \implies M_i \cap B^* \in Y$.

On the filters see Kueker [70], and [123, §3]. The theorem was proved in [123] but with two extra axioms, however it included the full case of varieties (i.e., including the non-Schreier ones). Later, the author eliminated those two extra axioms: Ax V in Ben David [13], and Ax I in [125] (answering a question of Fleissner on providing a combinatorial proof of the compactness). Hodges [55] contains also presentation of variants of this result.

There are some cases of incompactness (see Fleissner and Shelah [34], and [146]).

Problem 7.8. 1. Are there general theorems covering the incompactness phenomena?

2. Are there significantly better compactness theorems (for uncountable cofinality, of course)?

Related is

Question 7.9. What can be

 $\{\lambda : \text{there is a } \lambda \text{-free for } \mathcal{V} \text{ algebra } M \text{ of cardinality } \lambda \text{ which is not free}\},$

for a variety \mathcal{V} (at least with countable vocabulary)? (See Eklof and Mekler [31], Mekler and Shelah [77], Mekler, Shelah and Spinas [78].)

* * *

There are cases of strong dichotomy: if $\geq \lambda$ then $\geq 2^{\lambda}$, related to groups (see [142], Grossberg and Shelah [49], [50], and [118]; on Abelian groups see Fuchs [38]).

Question 7.10. [V = L] If $\lambda > \operatorname{cf}(\lambda) > \aleph_0$, G is a torsion free Abelian group of cardinality λ , can $\lambda = \nu_p(\operatorname{Ext}(G, \mathbb{Z}))$?

The cardinal $\nu_p(\operatorname{Ext}(G,\mathbb{Z}))$ is the dimension of $\{x \in \operatorname{Ext}(G,\mathbb{Z}) : px = 0\}$ as a vector space over $\mathbb{Z}/p\mathbb{Z}$. To avoid Ext note that (see Fuchs [38]) this group can be represented as $\operatorname{Hom}(G,\mathbb{Z}/p\mathbb{Z})/\{h/p : h \in \operatorname{Hom}(G,\mathbb{Z})\}$. If G is torsion free, then the group $\operatorname{Ext}(G,\mathbb{Z})$ is divisible and hence the ranks $\nu_p(\operatorname{Ext}(G,\mathbb{Z}))$ for p prime, and $\nu_0(\operatorname{Ext}(G,\mathbb{Z}))$, the rank of $\{x \in \operatorname{Ext}(G,\mathbb{Z}) : x \text{ torsion}\}$, determine $\operatorname{Ext}(G,\mathbb{Z})$ up to an isomorphism. If we assume $(\mathbf{V} = \mathbf{L} \text{ and})$ there is no weakly compact cardinal, this question is the only piece left for characterizing the possible such $\operatorname{Ext}(G,\mathbb{Z})$, see Mekler, Rosłanowski and Shelah [75].

Question 7.11. What is the first cardinal $\lambda = \lambda_{\kappa}$ such that: for every ring R of cardinality $\leq \kappa$, if there is endorigid (or rigid, or 1-to-1 rigid) R-module of size $\geq \lambda$, then there are such R-modules in arbitrarily large cardinals? (I.e. Hanf numbers).

8. Partitions and colourings

Remember (see Erdős, Hajnal, Maté and Rado [33])

Definition 8.1. 1. $\lambda \to (\alpha)^n_{\kappa}$ means: for every colouring $c : [\lambda]^n \longrightarrow \kappa$ there is a set $X \subseteq \lambda$ of order type α such that $c \upharpoonright [X]^n$ is constant.

- 2. $\lambda \to [\alpha]_{\kappa}^n$ means: for every colouring $c : [\lambda]^n \longrightarrow \kappa$ there is a set $X \subseteq \lambda$ of order type α such that $\operatorname{Rang}(c \upharpoonright [X]^n) \neq \kappa$.
- 3. $\lambda \to [\alpha]_{\kappa,\sigma}^n$ means: for every colouring $c : [\lambda]^n \longrightarrow \kappa$ there is a set $X \subseteq \lambda$ of order type α such that $\operatorname{Rang}(c \upharpoonright [X]^n)$ has cardinality $< \sigma$.

Definition 8.2. M is a Jonsson algebra if it is an algebra with countably many functions with no proper subalgebra of the same cardinality. (See [154], [95].)

Definition 8.3. $\Pr_1(\lambda, \mu, \theta, \sigma)$ means: there is $c : [\lambda]^2 \longrightarrow \theta$ such that if $u_i \in [\lambda]^{<\sigma}$ (for $i < \mu$) are pairwise disjoint and $\gamma < \theta$ then for some $i < j < \mu$ we have $c \upharpoonright (u_i \times u_j)$ is constantly γ . (See [154], [95].)

There are many more variants.

It irritates me that after many approximations, I still do not know (better for me "consistently no", better for set theory "yes") the answer to the following.

Question 8.4. If μ is singular, is there a Jonsson algebra on μ^+ ? (and even better $\Pr_1(\mu^+, \mu^+, \mu^+, \operatorname{cf}(\mu))$?)

Also the requirements on an inaccessible to get colouring theorems may well be an artifact of our inability, so let us state minimal open cases.

Question 8.5. 1. Let λ be the first ω -Mahlo cardinal. Does $\lambda \rightarrow [\lambda]^2_{\lambda}$ or at least $\lambda \rightarrow [\lambda]^2_{\theta}$ for $\theta < \lambda$?

2. Let λ be the first inaccessible cardinal which is $(\lambda \cdot \omega)$ -Mahlo. Is there a Jonsson algebra on λ ? (Even better $\lambda \rightarrow [\lambda]_{\lambda}^{2}$?)

In both parts it is better to have $\Pr_1(\lambda, \lambda, \lambda, \aleph_0)$, etc; it is interesting even assuming GCH.

Whereas under GCH the relation \rightarrow for cardinals is essentially understood (see Erdős, Hajnal, Maté and Rado [33]), the case of ordinals is not. As by [122] (GCH

for simplicity) $\alpha < \lambda = \mathrm{cf}(\lambda) \Rightarrow \lambda^{++} \to (\lambda^{+} + \alpha)_{2}^{2}$ (on the problem see [33]), and by Baumgartner, Hajnal and Todorčević [9] for $k < \omega$,

$$\alpha < \lambda = \operatorname{cf}(\lambda) \quad \Rightarrow \quad \lambda^{++} \to (\lambda^{+} + \alpha)_{k}^{2},$$

it remains open:

Question 8.6 (GCH). When does $\lambda^{++} \to (\lambda^{+} + \alpha)_{\aleph_0}^2$, where $2 \le \alpha < \lambda$? (The GCH assumption is for simplicity only.)

By [153] we know that if $\lambda = \lambda^{<\lambda}$ (e.g., $\lambda = \aleph_1 = 2^{\aleph_0}$) then possibly 2^{λ} is very large (in particular $> \lambda^{+\omega}$), but $2^{\lambda} \nrightarrow (\lambda \times \omega)_2^2$. However, $\lambda^{+2k} \to (\lambda \times n)_k^2$ (by [153, bottom of p.288]), so

Question 8.7. For $\lambda = \lambda^{<\lambda}$, $\lambda > \aleph_0$, and $k < \omega$ and $n < \omega$, what is the minimal m such that $\lambda^{+m} \to (\lambda \times n)_k^2$?

(Baumgartner, Hajnal and Todorčević [9, p.2, end of $\S 0$] prefer to ask whether $\lambda^{++} \to (\lambda + \omega)_3^2$ for $\lambda = \aleph_1$, so $\lambda = \lambda^{<\lambda}$ means CH as they choose another extreme case of the unknown).

Now, for me a try at consistency of negative answers for 8.6 calls for using historic forcing (see Shelah and Stanley [168], Rosłanowski and Shelah [83, §3]; it is explained below). On the other hand, large cardinals may make some positive results easier.

Question 8.8. Assume that $\lambda > \kappa > |\zeta| + \sigma$, and κ is a compact cardinal, and $\lambda = \lambda^{<\lambda}$. Does it follow that $\lambda^+ \to (\lambda + \zeta)^2_{\sigma}$?

I tend to think the answer is yes. If so, then we cannot expect, when $\lambda = \lambda^{<\lambda}$, that a λ -complete λ^+ -c.c. forcing notion will not add a counterexample c to $\lambda^+ \to (\lambda + \alpha^*)^2_{\aleph_0}$. Let for simplicity $\alpha^* = 2$. Also we should expect that for such c, for every $u \in [\lambda^+]^{<\lambda}$ we can find $f_n : u \times u \longrightarrow \omega$ for $n < \omega$ such that $f_n(\alpha, \alpha) = n$, and for no distinct $\alpha, \beta, \gamma \in u$ do we have:

$$c(\{\alpha,\beta\}) = n = f_n(\alpha,\beta)$$
 or $f_n(\alpha,\beta) = f_n(\alpha,\gamma) = c(\{\beta,\gamma\}).$

The point of historic forcing is that we know what kind of object our forcing notion has to add. In our case, we assume $\lambda = \lambda^{<\lambda}$ and a condition p has to give $u^p \in [\lambda^+]^{<\lambda}$ and $c^p : [u^p]^2 \longrightarrow \omega$ and by the above considerations also $f_n : u^p \times u^p \longrightarrow \omega$ (for $n < \omega$), and for having some leeway we let $f_n : u^p \times u^p \longrightarrow [\omega]^{\aleph_0}$ such that $n \in f_n(\alpha, \alpha)$, and we demand

(*) for no distinct $\alpha, \beta \in u^p$ and m do we have

$$m = c(\{\alpha, \beta\}) \in f_n(\alpha, \beta) \cap f_n(\alpha, \alpha),$$

(**) for no distinct α, β, γ from u^p do we have

$$c(\{\beta,\gamma\}) \in f_n(\alpha,\beta) \cap f_n(\alpha,\gamma).$$

Moreover, we choose $\langle A_{\eta} : \eta \in {}^{\omega >} \omega \rangle$, $A_{\eta} \in [\omega]^{\aleph_0}$, $\langle A_{\eta \frown \langle \ell \rangle} : \ell < \omega \rangle$ are pairwise disjoint subsets of A_{η} and demand $f_n(\alpha, \beta) \in \{A_{\eta} : \eta \in {}^{\omega >} \omega \}$.

Considering that our forcing will be strategically ($<\lambda$)-complete (as λ -complete seems too much both if the answer to 8.8 is yes and because of the properties of historic forcing in general), in order that there will be no $A \subseteq \lambda^+$, $\operatorname{otp}(A) = \lambda + \alpha^*$ such that $\underline{c} \upharpoonright [A]^2$ is constant, we have a set \mathcal{T} of cardinality $<\lambda$ such that (for simplicity $\lambda = \mu^+$, maybe also μ regular):

(a) each $x \in \mathcal{T}^p$ has the form $x = (a, \delta, b) = (a^x, \delta^x, b^x)$, where

- (i) $\delta < \lambda^+$, $cf(\delta) = \lambda$,
- (ii) $a \subseteq u^p \setminus \delta$, $otp(a) = \zeta$,
- (iii) $b \subseteq u^p \cap \delta$, $otp(b) = \mu$,
- (b) if $(a', \delta', b'), (a'', \delta'', b'') \in \mathcal{T}^p$ then $|b' \cap b''| < \mu$ (probably not necessary),
- (c) $c^p \upharpoonright [a \cup b]^2$ is constantly n(x),
- (d) for no $\alpha \in u^p \cap \delta \setminus \sup(b)$ do we have

$$(\forall \beta \in a \cup b)[c^p(\{\alpha, \beta\}) = n(x)].$$

A p like above will be a precondition. Now the "history" enters and the proof should be clear just as the definition of the set of the conditions should roll itself.

So atomic conditions will have u^p a singleton, each condition will have a history telling how it was created: each step in the history corresponds to one of the reasons for creating a condition in the proof. Naturally, a major reason is the proof of the λ^+ -c.c. by the Δ -system lemma. So assume for $\ell=1,2$ that $p_\ell\in\mathbb{P},\ \alpha_1<\alpha_2,\ u^{p_1}\cap\alpha_1=u^{p_2}\cap\alpha_2,\ u^{p_1}\subseteq\alpha_2,\ {\rm otp}(u^{p_1})={\rm otp}(u^{p_2})$ and the order preserving mapping ${\rm OP}_{u^{p_2},u^{p_1}}$ from u^{p_1} onto u^{p_2} maps p_1 to p_2 . We have to amalgamate p_1 and p_2 getting q, so we have to determine f_n^q on $(u^{p_1}\setminus u^{p_2})\times(u^{p_2}\setminus u^{p_1})$, and $c^q(\{\alpha,\beta\})$ for $\alpha\in u^{p_1}\setminus u^{p_2},\ \beta\in u^{p_2}\setminus u^{p_1}$.

But our vision is that there is one line of history, so should the history of q continue the history of p_1 with p_2 joining or should the history of q continue the history of p_2 with p_1 joining? Both are O.K., but we get two distinct conditions q', q'' which, however, are equivalent; i.e. $q' \leq q'' \leq q'$.

Generally going back and changing the history as above we get an equivalent condition if this is done finitely many times (this also explains why we get strategical ($< \lambda$)-completeness and not λ -completeness). That is, we define $p \leq_{\rm pr} q$ iff p appears in the history of q, $p \sim q$ if p is gotten from q by finitely many changes as above in the history, and lastly $p \leq q$ iff $(\exists p')(p' \sim p \& p' \leq_{\rm pr} q)$.

What is left? "Only" carrying out the amalgamation (using and guaranteeing the conditions, and probably changing them so that they will fit).

Question 8.9. What are the best cardinals needed for the canonization theorems in [130]?

An old well-known problem is

Question 8.10. Is $\aleph_1 \to [\aleph_1; \aleph_1]_2^2$ consistent? (And variants, connected to the *L*-space problem; this seems related to 5.1).

Another problem of Erdős is (the answer is consistently yes, see [144], even colouring also no edges, but provability in ZFC is not clear):

Question 8.11. Is there a graph G with no K_4 (complete graph on 4 vertices) such that $G \to (K_3)^2_{\aleph_0}$, that is for any colouring of the edges by \aleph_0 colours there is a monochromatic triangle?

We can ask it for K_k, K_{k+1} instead of k=3 and colouring r-tuples instead of pairs; the answer still is consistently yes (see [144]), so the problem is in ZFC. See [106, Ch.III,§1] on a connection to model theory.

* * *

A very nice theorem of Hajnal [51] says that, e.g., for any finite graph G and κ for some graph $H, H \to (G)^2_{\kappa}$, but leaves as a mystery:

Question 8.12. Let G be a countable graph, is there a graph H such that

$$H \to (G)^2_{\aleph_0}$$
?

Starting from a problem of Erdös and Hajnal [32], I have been very interested in consistency results, e.g., of the form $\lambda \to [\mu]_3^2$, $\kappa < \mu < \lambda \le 2^{\kappa}$ (see [121]). Usually those are really canonization theorems, for a fixed natural coloring, any other is, on a large set, computable from it. Those results help sometimes in consistency results (just as, e.g., the Erdös-Rado theorem helps in ZFC results). Still it seems to me worthwhile to know.

- Question 8.13. 1. Can we put together the results of, e.g., [121], Shelah and Stanley [167] and [149]? Assume that $\kappa = \kappa^{<\kappa} < \lambda$, λ is, e.g., strongly inaccessible large enough. Can we find a $(<\kappa)$ -complete, κ^+ -c.c. forcing notion $\mathbb P$ such that in $\mathbf V^{\mathbb P}$:
 - (a) for $\sigma < \kappa$ and $\mu < \lambda$ we can find μ', λ' such that $\mu < \mu' < \lambda' < \lambda$ and $\lambda' \to [\mu']_{\sigma,2}^2$;
 - (b) if κ is a measurable indestructibly by adding many Cohens, then also the parallel results for colouring *n*-tuples (see [149]);
 - (c) if $\kappa = \aleph_0$, we also have results on colouring n-tuples simultaneously for all n?
 - 2. Add the hopeful consistency answer for 8.14, 8.15.

Probably easier are, e.g.

Question 8.14. Is it consistent that for some n, $2^{\aleph_0} = \aleph_n = \lambda \to [\aleph_1]_3^2$? (The exact n is less exciting for me, the main division line seems to me \aleph_ω , of course best to know the exact λ .)

Question 8.15. Is it consistent that $2^{\aleph_0} > \lambda \to [\aleph_1]_3^2$?

* * *

On Finite Combinatorics. Spencer, Szemeredi and Alon told me that finding $\lim(\log(r_2^2(n)/n))$ is a major problem (see Definition 8.16 below), but the difference between lower and upper bounds seems to me negligible.

Definition 8.16. $r_k^m(n)$ is the minimal r such that $r \to (n)_k^m$.

Erdös and Hajnal ask, and I find more convincing, the following.

Question 8.17. What is the order of magnitude of $r_2^3(n)$?

We expect it should be 2^{2^n} , or e.g. $2^{2^{(n^{\varepsilon})}}$ for some $\varepsilon > 0$. But we cannot rule out its being 2^n or e.g. $2^{n^{1/\varepsilon}}$ for some $\varepsilon > 0$.

Here the difference is large.

Note that for four colours the problem (what is $r_4^3(n)$) is settled; but I think the true question is:

Question 8.18. Determine (order of magnitude is OK) $f_k(n,r), f_k^+(n,c)$ where:

(a) $f_3(n,c)$ is the minimal m such that for every $\mathbf{d}: [m]^3 \longrightarrow \{0,\ldots,c-1\}$, there are $A \in [m]^n$ and a strictly increasing function $h:A \longrightarrow \{0,\ldots,2^n-1\}$ such that for $\ell_0 < \ell_1 < \ell_2$

in A, the value of $\mathbf{d}(\{\ell_0, \ell_1, \ell_2\})$ is determined by the quantifier–free type $\langle h(\ell_0), h(\ell_1), h(\ell_2) \rangle$ in B_n^3 ,

where B_n^3 has the universe $\{0, \dots, 2^n - 1\}$ and two relations: (viewed as n^2) the lexicographic order and

$$\{(\eta_0, \eta_1, \eta_2) : \eta_0 <_{\ell x} \eta_1 <_{\ell x} \eta_2 \text{ and } \ell g(\eta_0 \cap \eta_1) < \ell g(\eta_1 \cap \eta_2)\}.$$

- (b) $f_3^+(n,c)$ is defined similarly but for every pregiven Rang(h) we can find such h.
- (c) B_n^k is defined below by induction of k, and then $f_k(n,c)$, $f_k^+(n,c)$ are defined analogously to f_3 , f_3^+ .
- (d) Define canonization numbers $g_k(n), g_k^+(n)$, which is the first m such that: if $d: [n]^k \to C$, with no restriction on the cardinality of C, then we can find A, h as above, and a quantifier free formula ϕ in the vocabulary of B_n^k such that for any $u_1, u_2 \in [A]^k$ we have

$$d(u_1) = d(u_2) \quad \text{iff} \quad \phi(\dots, h(\ell_1), \dots; \dots, h(\ell_2), \dots)_{\ell_1 \in u_1, \ell_2 \in u_2} \text{ is satisfied in } B_n^k.$$

The explicit way to describe B_n^k , by induction on k is: it has a linear order $<_k : B_n^2$ is the structure (n, <),

 B_n^{k+1} has universe $B_n^k\{0,1\}$ and the relation:

 $\eta <_{k+1} \nu$ if and only if

for some $y = y(\eta, \nu) \in B_n^k$ we have $\eta(y) = 0$, $\nu(y) = 1$ and

$$\eta \upharpoonright \{x \in B_n^k : x <_k y\} = \nu \upharpoonright \{x \in B_n^k : x <_k y\}.$$

For an m-place relation $R^{B_n^k}$ of B_n^k , $R^{B_n^{k+1}}$ is a 2m-place relation on B_n^{k+1} , namely

$$\{\langle \eta_0, \dots, \eta_{2m-1} \rangle : \eta_\ell \in B_n^{k+1}, \ \eta_\ell <_{k+1} \eta_{\ell+1} \text{ and } \langle y(\eta_0, \eta_1), y(\eta_2, \eta_3), \dots \rangle \in R^{B_n^k} \}.$$

Now it is not clear how fast the number in 8.18 grows, e.g., we cannot exclude $2^{2^{n+c+k}}$. The main question is whether it grows like h(k)-iterated exponentiation in n (say c fixed) with h going to infinity, or with h constant. Of course, enriching somewhat the structure is not a great loss to me.

Question 8.19. Let $f^*(n,c)$ be the first m such that if $\langle A_\ell : \ell < m \rangle$ are pairwise disjoint, $|A_\ell| = m$ for $\ell < n$ and

$$F: \{w \subseteq \bigcup_{\ell} A_{\ell}: |w \cap A_{\ell}| \in \{1, 2\}, \ (\exists!\ell)(|w| = 1)\} \longrightarrow C,$$

where |C| = c then for some $x_{\ell} \neq y_{\ell}$ from A_{ℓ} for $\ell < n$ we have

$$\ell^* < n \implies F(\{x_\ell, y_\ell : \ell \neq \ell^*\} \cup \{x_{\ell^*}\}) = F(\{x_\ell, y_\ell : \ell \neq \ell^*\} \cup \{y_{\ell^*}\}).$$

Again the main question for me is: Does $f^*(n,c)$ grow as a fixed iterated exponentiation?

(This is connected to the van der Waerden theorem, see [143]).

On the Ramsey Theory see Graham, Rothschild and Spencer [47].

Definition 8.20. 1. For a group G and a subset A of G, and a group H let $H \to (G)^A_\sigma$ mean:

if **d** is a function with domain H and range of cardinality $\leq \sigma$,

then for some embedding h of G into H the function \mathbf{d} restricted to h(A) is constant.

- 2. For a group G and a subset A of G, and group H let $H \to [G]_{\sigma,\tau}^A$ mean: if \mathbf{d} is a function with domain H and range of cardinality $< \sigma$,
 - then for some embedding h of G into H the range of the function \mathbf{d} restricted to h(A) has cardinality $< \tau$.

If we omit τ , we mean just that the range is not equal to σ .

- 3. For a group G and an equivalence relation E on G, and group H let $H \to (G)^E_\sigma$ mean:
 - if **d** is a function with domain H and range of cardinality $< \sigma$.

then for some embedding h of G into H, for any $x, y \in G$ which are E-equivalent we have $\mathbf{d}(h(x)) = \mathbf{d}(h(y))$.

- 4. In part (1) (or (2), or (3)) we can replace A (or the domain of E) by the family of subgroups of G isomorphic to a fixed group K, and then \mathbf{d} is a function with domain being the set of subgroups of G isomorphic to K.
- 5. Like part (4), but we replace "subgroups isomorphic to K" by "embedding of K", and then replace " \rightarrow " by " \rightarrow *".

Discussion 8.21. There is a connection between the two last definitions: the first implies a special case of the second one, when we restrict ourselves to permutation groups of some finite set, and A is the set of conjugates of the permutation just interchanging two elements.

Problem 8.22. 1. Investigate the arrows from Definition 8.20.

2. In particular, consider the case when A is a set of pairwise conjugate members of G each of order two.

RECENT ADVANCES:

During the winter of 1999, Gitik told me that he can start with

$$\mathbf{V} \models \text{``} \kappa_n \text{ hypermeasurable of order } \lambda_n, \\ \lambda_n \text{ first (strongly) inaccessible } > \kappa_n, \ \lambda_n < \kappa_{n+1}, \ \lambda > \kappa = \sum_n \kappa_n \text{''},$$

and find a forcing notion \mathbb{P} , not adding bounded subsets of $\kappa = \sum_{n < \omega} \kappa_n$, satisfying the κ^{++} -c.c., and making $2^{\kappa} \geq \lambda$. I have conjectured that combining this proof with earlier proof, you can demand \mathbb{P} makes λ_n be the n-th inaccessible cardinal, $\kappa = \sum_n \lambda_n$, GCH holds below κ and $2^{\kappa} \geq \lambda$, so $\operatorname{pp}(\kappa) \geq \lambda$. Gitik has confirmed this conjecture with κ_n Mahlo. This proves that though 1.20 is open, other theorem which holds for λ of cofinality \aleph_1 cannot be generalized to cofinality \aleph_0 .

References

- [1] Uri Abraham. Aronszajn trees on \aleph_2 and \aleph_3 . Ann. Pure Appl. Logic, 24:213–230, 1983.
- [2] Uri Abraham, Matatyahu Rubin, and Saharon Shelah. On the consistency of some partition theorems for continuous colorings, and the structure of ℵ₁-dense real order types. Annals of Pure and Applied Logic, 29:123–206, 1985.
- [3] Uri Abraham and Saharon Shelah. Isomorphism types of Aronszajn trees. Israel Journal of Mathematics, 50:75–113, 1985.
- [4] Uri Abraham and Stevo Todorčević. Partition properties of ω₁ compatible with CH. Fundamenta Mathematicae, 152:165–181, 1997.

- [5] Uri Avraham (Abraham), Keith J. Devlin, and Saharon Shelah. The consistency with CH of some consequences of Martin's axiom plus 2^{ℵ0} > ℵ1. Israel Journal of Mathematics, 31:19–33, 1978.
- [6] Tomek Bartoszyński and Haim Judah. Set Theory: On the Structure of the Real Line. A K Peters, Wellesley, Massachusetts, 1995.
- [7] Tomek Bartoszyński, Andrzej Rosłanowski, and Saharon Shelah. After all, there are some inequalities which are provable in ZFC. *Journal of Symbolic Logic*, to appear, math.LO/9711222⁴.
- [8] Tomek Bartoszyński, Andrzej Rosłanowski, and Saharon Shelah. Adding one random real. Journal of Symbolic Logic, 61:80–90, 1996, math.LO/9406229.
- [9] James Baumgartner, Andras Hajnal, and Stevo Todorčević. Extensions of the Erdos-Rado Theorems. In Finite and Infinite Combinatorics in Set Theory and Logic, pages 1–18. Kluwer Academic Publishers, 1993. N.W. Sauer et. al. eds.
- [10] James E. Baumgartner. Decomposition of embedding of trees. Notices Amer. Math. Soc., 17:967, 1970.
- [11] James E. Baumgartner. All ℵ₁-dense sets of reals can be isomorphic. Fund. Math., 79:101–106, 1973.
- 12] James E. Baumgartner. Ultrafilters on ω. The Journal of Symbolic Logic, 60:624–639, 1995.
- [13] Shai Ben David. On Shelah's compactness of cardinals. Israel J. of Math., 31:34–56 and 394, 1978
- [14] Andreas Blass and Saharon Shelah. There may be simple P_{\aleph_1} and P_{\aleph_2} -points and the Rudin-Keisler ordering may be downward directed. Annals of Pure and Applied Logic, 33:213–243, 1987.
- [15] Andreas Blass and Saharon Shelah. Ultrafilters with small generating sets. Israel Journal of Mathematics, 65:259–271, 1989.
- [16] Robert Bonnet and Donald Monk. Handbook of Boolean Algebras, vol. 1–3. North Holland, 1989.
- [17] Joerg Brendle and Saharon Shelah. Ultrafilters on ω their ideals and their cardinal characteristics. Transactions of the American Mathematical Society, accepted, math.LO/9710217.
- [18] Timothy J. Carlson. Strong measure zero and strongly meager sets. Proceedings of the American Mathematical Society, 118:577–586, 1993.
- [19] Krzysztof Ciesielski. Set theoretic real analysis. J. of Applied Analysis, 3:143–190, 1997.
- [20] Krzysztof Ciesielski and Saharon Shelah. Category analog of sub-measurability problem. Fundamenta Mathematicae, submitted, math.LO/9905147.
- [21] James Cummings, Mirna Džamonja, and Saharon Shelah. A consistency result on weak reflection. Fundamenta Mathematicae, 148:91–100, 1995, math.LO/9504221.
- [22] James Cummings and Matthew Foreman. The tree property. Adv. Math., 133:1–32, 1998.
- [23] James Cummings and Saharon Shelah. A model in which every infinite Boolean algebra has many subalgebras. *Journal of Symbolic Logic*, 60:992–1004, 1995, math.LO/9509227.
- [24] James Cummings and Saharon Shelah. Cardinal invariants above the continuum. Annals of Pure and Applied Logic, 75:251–268, 1995, math.LO/9509228.
- [25] Keith J. Devlin and Saharon Shelah. A weak version of \diamondsuit which follows from $2^{\aleph_0} < 2^{\aleph_1}$. Israel Journal of Mathematics, 29:239–247, 1978.
- [26] Tony Dodd and Ronald B. Jensen. The covering lemma for K. Ann. of Math Logic, 22:1–30, 1982.
- [27] Mirna Džamonja and Saharon Shelah. Universal graphs at successors of singular strong limits. prevrint.
- [28] Mirna Džamonja and Saharon Shelah. Weak reflection can first happen at the successor of singular. Preprint.
- [29] Mirna Džamonja and Saharon Shelah. On squares, outside guessing of clubs and $I_{< f}[\lambda]$. Fundamenta Mathematicae, 148:165–198, 1995, math.LO/9510216.
- [30] Mirna Džamonja and Saharon Shelah. Saturated filters at successors of singulars, weak reflection and yet another weak club principle. Annals of Pure and Applied Logic, 79:289– 316, 1996, math.LO/9601219.
- [31] Paul C. Eklof and Alan Mekler. Almost free modules; Set theoretic methods. North Holland Library, 1990.

⁴Papers with numbers of the form math.XX/xxxxxxx are electronically available from the XXX Mathematics Archive at http://front.math.ucdavis.edu/

- [32] Paul Erdős and Andras Hajnal. Unsolved Problems in set theory. In Axiomatic Set Theory, volume XIII Part I of Proc. of Symp. in Pure Math., pages 17–48, Providence, R.I., 1971. AMS
- [33] Paul Erdős, Andras Hajnal, A. Maté, and Richard Rado. Combinatorial set theory: Partition Relations for Cardinals, volume 106 of Studies in Logic and the Foundation of Math. North Holland Publ. Co, Amsterdam, 1984.
- [34] William G. Fleissner and Saharon Shelah. Collectionwise Hausdorff: incompactness at singulars. Topology and its Applications, 31:101–107, 1989.
- [35] Matthew Foreman and Hugh Woodin. The generalized continuum hypothesis can fail everywhere. *Annals Math.*, 133:1–36, 1991.
- [36] David Fremlin. Problem list. circulated notes (1994).
- [37] David Fremlin. Real-valued-measurable cardinals. In Set Theory of the Reals, volume 6 of Israel Mathematical Conference Proceedings, pages 151–304, 1993.
- [38] Laszlo Fuchs. Infinite Abelian Groups, volume I, II. Academic Press, New York, 1970, 1973.
- [39] Salvador Garcia-Ferreira and Winfried Just. Two examples of relatively pseudocompact spaces. Questions and Answers in General Topology, 17:35–45, 1999.
- [40] Moti Gitik. All uncountable cardinals can be singular. Israel Journal of Mathematics, 35:61–88, 1980.
- [41] Moti Gitik and Saharon Shelah. More on real-valued measurable cardinals and forcing with ideals. Israel Journal of Mathematics, submitted, math.LO/9507208.
- [42] Moti Gitik and Saharon Shelah. Forcings with ideals and simple forcing notions. Israel Journal of Mathematics, 68:129–160, 1989.
- [43] Moti Gitik and Saharon Shelah. On certain indestructibility of strong cardinals and a question of Hajnal. Archive for Mathematical Logic, 28:35–42, 1989.
- [44] Moti Gitik and Saharon Shelah. More on simple forcing notions and forcings with ideals. Annals of Pure and Applied Logic, 59:219–238, 1993.
- [45] Moti Gitik and Saharon Shelah. Less saturated ideals. Proceedings of the American Mathematical Society, 125:1523–1520, 1997, math.LO/9503203.
- [46] Martin Goldstern and Saharon Shelah. Many simple cardinal invariants. Archive for Mathematical Logic, 32:203–221, 1993, math.LO/9205208.
- [47] Ronald Graham, Bruce L. Rothschild, and Joel Spencer. Ramsey Theory. Willey Interscience Series in Discrete Mathematics. Willey, New York, 1980.
- [48] John Gregory. Higher Souslin trees and the generalized continuum hypothesis. *Journal of Symbolic Logic*, 41(3):663–671, 1976.
- [49] Rami Grossberg and Saharon Shelah. On the structure of $\operatorname{Ext}_p(G, \mathbf{Z})$. Journal of Algebra, 121:117–128, 1989. See also [GrSh:302a] below.
- [50] Rami Grossberg and Saharon Shelah. On cardinalities in quotients of inverse limits of groups. Mathematica Japonica, 47(2), 1998.
- [51] Andras Hajnal. True embedding partition relations. In Finite and Infinite Combinatorics in Sets and Logic, pages 135–152. Kluwer Academic Publishers, 1993.
- [52] Andras Hajnal and Peter Hamburger. Halmazelmélet (Set Theory). Tankönyvkiaó Vállalat, Budapest, Hungary.
- [53] Andras Hajnal, Istvan Juhász, and Z. Szentmiklossy. On the structure of CCC partial orders. *Algebra Universalis*, to appear.
- [54] Joel D. Hamkins. Every group has a terminating transfinite automorphism tower. Proc. Amer. Math. Soc., 126:3223–3226, 1998.
- [55] Wilfrid Hodges. For singular λ , λ -free, implies free. Algebra Universalis, 12:205–220, 1981.
- [56] Jaime Ihoda (Haim Judah) and Saharon Shelah. Souslin forcing. The Journal of Symbolic Logic, 53:1188–1207, 1988.
- [57] Thomas Jech and Saharon Shelah. Possible pcf algebras. Journal of Symbolic Logic, 61:313–317, 1996, math.LO/9412208.
- [58] Haim Judah and Saharon Shelah. MA(σ-centered): Cohen reals, strong measure zero sets and strongly meager sets. Israel Journal of Mathematics, 68:1–17, 1989.
- [59] Haim Judah and Saharon Shelah. Baire Property and Axiom of Choice. Israel Journal of Mathematics, 84:435–450, 1993, math.LO/9211213.
- [60] Istvan Juhász. Cardinal functions. II. In Handbook of set-theoretic topology, pages 63–109. North Holland Publ. Co, 1984.

- [61] Istvan Juhász. Cardinal functions. In Miroslav Hušek and Jan van Mill, editors, Recent progress in general topology (Prague, 1991), pages 417–441. North-Holland Publishing Co., Amsterdam, 1992.
- [62] Istvan Juhász and Saharon Shelah. How large can a hereditarily separable or hereditarily Lindelof space be? Israel Journal of Mathematics, 53:355–364, 1986.
- [63] Winfried Just, A. R. D. Mathias, Karel Prikry, and Petr Simon. On the existence of large p-ideals. *Journal of Symbolic Logic*, 55(2):457–465, 1990.
- [64] Winfried Just, Saharon Shelah, and Simon Thomas. The Automorphism Tower Problem III: Closed Groups of Uncountable Degree. Advances in Mathematics, accepted.
- [65] Alexander S. Kechris and Sławomir Solecki. Approximation of analytic by Borel sets and definable countable chain conditions. *Israel Journal of Mathematics*, 89:343–356, 1995.
- [66] Jussi Ketonen. On the existence of P-points in the Stone-Čech compactification of integers. Fund. Math., 92:91–94, 1976.
- [67] Menachem Kojman and Saharon Shelah. Non-existence of Universal Orders in Many Cardinals. Journal of Symbolic Logic, 57:875–891, 1992, math.LO/9209201.
- [68] Menachem Kojman and Saharon Shelah. The universality spectrum of stable unsuperstable theories. Annals of Pure and Applied Logic, 58:57–72, 1992, math.LO/9201253.
- [69] Peter Komjath. On second-category sets. Proc. Amer. Math. Soc., 107:653-654, 1989.
- [70] David W. Kueker. Countable approximations and Lowenheim-Skolem theorems. Ann. Math. Logic, 11:57–103, 1977.
- [71] Kenneth Kunen. Random and Cohen Reals. In K. Kunen and J. E. Vaughan, editors, Handbook of Set-Theoretic Topology, pages 887-911. Elsevier Science Publishers B.V., 1984.
- [72] Kenneth Kunen. Large homogeneous compact spaces. In J. van Mill and G.M. Reed, editors, Open Problems in Topology. Elsevier Science Publishers B.V, 1990.
- [73] Richard Laver. On the consistency of Borel's conjecture. Acta Math., 137:151–169, 1976.
- [74] A.R.D. Mathias. 0# and the p-point problem. In Higher Set Theory (Proc. Conf., Math. Forschungsinst., Oberwolfach, 1977), volume 669 of Lecture Notes in Math., pages 375–383. Springer, Berlin, 1978.
- [75] Alan H. Mekler, Andrzej Rosłanowski, and Saharon Shelah. On the p-rank of Ext. Israel Journal of Mathematics, to appear, math.LO/9806165.
- [76] Alan H. Mekler and Saharon Shelah. Uniformization principles. The Journal of Symbolic Logic, 54:441–459, 1989.
- [77] Alan H. Mekler and Saharon Shelah. Almost free algebras . Israel Journal of Mathematics, 89:237–259, 1995, math.LO/9408213.
- [78] Alan H. Mekler, Saharon Shelah, and Otmar Spinas. The essentially free spectrum of a variety. Israel Journal of Mathematics, 93:1–8, 1996, math.LO/9411234.
- [79] Arnold W. Miller. Arnie Miller's problem list. In Haim Judah, editor, Set Theory of the Reals, volume 6 of Israel Mathematical Conference Proceedings, pages 645–654. Proceedings of the Winter Institute held at Bar–Ilan University, Ramat Gan, January 1991.
- [80] Donald Monk. Cardinal Invariants of Boolean Algebras, volume 142 of Progress in Mathematics. Birkhäuser Verlag, Basel–Boston–Berlin, 1996.
- [81] Judy Roitman. Adding a random or Cohen real: topological consequences and the effect on Martin's axiom. Fundamenta Mathematicae, 103:47–60, 1979.
- [82] Andrzej Rosłanowski and Saharon Shelah. Cardinal invariants of ultrapoducts of Boolean algebras. Fundamenta Mathematicae, 155:101–151, 1998, math.LO/9703218.
- [83] Andrzej Rosłanowski and Saharon Shelah. Forcing for hL, hd and Depth. Colloquium Mathematicum, accepted, math.LO/9808104.
- [84] Andrzej Rosłanowski and Saharon Shelah. More on cardinal functions on Boolean algebras. Annals of Pure and Applied Logic, accepted, math.LO/9808056.
- [85] Andrzej Rosłanowski and Saharon Shelah. Norms on possibilities I: forcing with trees and creatures. Memoirs of the AMS, to appear, math.LO/9807172.
- [86] Andrzej Rosłanowski and Saharon Shelah. Norms on possibilities II: more ccc ideals on 2^{ω} . Journal of Applied Analysis, 3:103–127, 1997, math.LO/9703222.
- [87] Andrzej Rosłanowski and Saharon Shelah. Norms on possibilities III: strange subsets of the real line. in preparation.
- [88] Andrzej Rosłanowski and Saharon Shelah. Norms on possibilities IV: ccc forcing notions. Preprint.

- [89] Matatyahu Rubin and Saharon Shelah. Combinatorial problems on trees: partitions, Δ-systems and large free subtrees. Annals of Pure and Applied Logic, 33:43–81, 1987.
- [90] Gershon Sageev and Saharon Shelah. Noetherian ring with free additive groups. Abstracts of the American Mathematical Society, 7:369, 1986.
- [91] James D. Sharp and Simon Thomas. Uniformization problems and the cofinality of the infinite symmetric group. Notre Dame J. Formal Logic, 35:328–345, 1994.
- [92] James D. Sharp and Simon Thomas. Unbounded families and the cofinality of the infinite symmetric group. Arch. Math. Logic, 34:33–45, 1995.
- [93] Saharon Shelah. Remarks on \aleph_1 -CWH not CWH first countable spaces. In Set Theory, Boise ID, 1992–1994, volume 192 of Contemporary Mathematics, pages 103–145.
- [94] Saharon Shelah. Also quite large $\mathfrak{b} \subseteq \mathrm{pcf}(\mathfrak{a})$ behave nicely. math.LO/9906018.
- [95] Saharon Shelah. Analytical Guide and Corrections to [154]. math.LO/9906022.
- [96] Saharon Shelah. Applications of PCF theory. Journal of Symbolic Logic, submitted, math.LO/9804155.
- [97] Saharon Shelah. Categoricity of an abstract elementary class in two successive cardinals. Israel Journal of Mathematics, submitted, math.LO/9805146.
- [98] Saharon Shelah. Cellularity of free products of Boolean algebras (or topologies). Fundamenta Mathematica, accepted, math.LO/9508221.
- [99] Saharon Shelah. Constructing Boolean algebras for cardinal invariants. Algebra Universalis, submitted, math.LO/9712286.
- [100] Saharon Shelah. Covering of the null ideal may have countable cofinality. Fundamenta Mathematicae, submitted, math.LO/9810181.
- [101] Saharon Shelah. Few non-minimal types and non-structure. in preparation math.LO/9906023.
- [102] Saharon Shelah. The future of set theory. In Haim Judah, editor, Set Theory of the Reals, volume 6 of Israel Mathematical Conference Proceedings. Proceedings of the Winter Institute held at Bar–Ilan University, Ramat Gan, January 1991.
- [103] Saharon Shelah. Historic iteration with ℵ_ε-support. Archive for Mathematical Logic, accepted, math.LO/9607227.
- [104] Saharon Shelah. Iteration of λ -complete forcing notions not collapsing λ^+ . in preparation, math.LO/9906024.
- [105] Saharon Shelah. More on Weak Diamond. Preprint, math.LO/9807180.
- [106] Saharon Shelah. Non-structure theory, volume accepted. Oxford University Press.
- [107] Saharon Shelah. Non-elementary proper forcing notions. Journal of Applied Analysis, submitted, math.LO/9712283.
- [108] Saharon Shelah. Non-elementary proper forcing notions II. In preparation.
- [109] Saharon Shelah. Non-existence of universal members in classes of Abelian groups. *Journal of Group Theory*, submitted, math.LO/9808139.
- [110] Saharon Shelah. Not collapsing cardinals $\leq \kappa$ in $(<\kappa)$ -support iterations. Israel Journal of Mathematics, accepted, math.LO/9707225.
- [111] Saharon Shelah. Not collapsing cardinals $\leq \kappa$ in $(< \kappa)$ -support iterations II. Israel Journal of Mathematics, submitted, math.LO/9808140.
- [112] Saharon Shelah. On Arhangelskii's Problem. in preparation, math.LO/9906025.
- [113] Saharon Shelah. On CS iterations not adding reals. in preparation.
- [114] Saharon Shelah. PCF and infinite free subsets. Archive for Mathematical Logic, accepted, math.LO/9807177.
- [115] Saharon Shelah. Projective measurability does not imply projective Baire. in preparation.
- [116] Saharon Shelah. Special Subsets of $^{\mathrm{cf}(\mu)}\mu$, Boolean Algebras and Maharam measure Algebras. General Topology and its Applications Proc. of Prague Topological Symposium 1996, accepted, math.LO/9804156.
- [117] Saharon Shelah. Strong covering without squares. Fundamenta Mathematicae, accepted, math.LO/9604243.
- [118] Saharon Shelah. Strong dichotomy of cardinality. Results in Mathematics, submitted, math.LO/9807183.
- [119] Saharon Shelah. The Generalized Continuum Hypothesis revisited. Israel Journal of Mathematics, accepted, math.LO/9809200.
- [120] Saharon Shelah. The null ideal restricted to a non-null set may be saturated. Asian Journal of Mathematics, submitted, math.LO/9705213.

- [121] Saharon Shelah. Was Sierpiński right? IV. Journal of Symbolic Logic, accepted, math.LO/9712282.
- [122] Saharon Shelah. Notes on combinatorial set theory. Israel Journal of Mathematics, 14:262–277, 1973.
- [123] Saharon Shelah. A compactness theorem for singular cardinals, free algebras, Whitehead problem and transversals. Israel Journal of Mathematics, 21:319–349, 1975.
- [124] Saharon Shelah. Decomposing uncountable squares to countably many chains. *Journal of Combinatorial Theory. Ser. A*, 21:110–114, 1976.
- [125] Saharon Shelah. A combinatorial proof of the singular compactness theorem. Mineograph notes and lecture in a mini-conference, Berlin, August'78, 1977.
- [126] Saharon Shelah. Whitehead groups may be not free, even assuming CH. I. Israel Journal of Mathematics, 28:193–204, 1977.
- [127] Saharon Shelah. Classification theory and the number of nonisomorphic models, volume 92 of Studies in Logic and the Foundations of Mathematics. North-Holland Publishing Co., Amsterdam-New York, xvi+544 pp, \$62.25, 1978.
- [128] Saharon Shelah. On successors of singular cardinals. In Logic Colloquium '78 (Mons, 1978), volume 97 of Stud. Logic Foundations Math, pages 357–380. North-Holland, Amsterdam-New York, 1979.
- [129] Saharon Shelah. Remarks on Boolean algebras. Algebra Universalis, 11:77–89, 1980.
- [130] Saharon Shelah. Canonization theorems and applications. The Journal of Symbolic Logic, 46:345–353, 1981.
- [131] Saharon Shelah. On saturation for a predicate. Notre Dame Journal of Formal Logic, 22:239–248, 1981.
- [132] Saharon Shelah. Proper forcing, volume 940 of Lecture Notes in Mathematics. Springer-Verlag, Berlin-New York, xxix+496 pp, 1982.
- [133] Saharon Shelah. Classification theory for nonelementary classes, I. The number of uncountable models of $\psi \in L_{\omega_1,\omega}$. Part B. Israel Journal of Mathematics, 46:241–273, 1983.
- [134] Saharon Shelah. Constructions of many complicated uncountable structures and Boolean algebras. *Israel Journal of Mathematics*, 45:100–146, 1983.
- [135] Saharon Shelah. Lifting problem of the measure algebra. Israel Journal of Mathematics, 45:90–96, 1983.
- [136] Saharon Shelah. Can you take Solovay's inaccessible away? Israel Journal of Mathematics, 48:1–47, 1984.
- [137] Saharon Shelah. Diamonds, uniformization. The Journal of Symbolic Logic, 49:1022–1033,
- [138] Saharon Shelah. More on proper forcing. The Journal of Symbolic Logic, 49:1034–1038, 1984.
- [139] Saharon Shelah. More on the weak diamond. Annals of Pure and Applied Logic, 28:315–318, 1985
- [140] Saharon Shelah. Remarks on the numbers of ideals of Boolean algebra and open sets of a topology. In Around classification theory of models, volume 1182 of Lecture Notes in Mathematics, pages 151–187. Springer, Berlin, 1986.
- [141] Saharon Shelah. Appendix: on stationary sets (in "Classification of nonelementary classes. II. Abstract elementary classes"). In Classification theory (Chicago, IL, 1985), volume 1292 of Lecture Notes in Mathematics, pages 483–495. Springer, Berlin, 1987. Proceedings of the USA-Israel Conference on Classification Theory, Chicago, December 1985; ed. Baldwin, J.T.
- [142] Saharon Shelah. Can the fundamental (homotopy) group of a space be the rationals? Proceedings of the American Mathematical Society, 103:627–632, 1988.
- [143] Saharon Shelah. Primitive recursive bounds for van der Waerden numbers. Journal of the American Mathematical Society, 1:683–697, 1988.
- [144] Saharon Shelah. Consistency of positive partition theorems for graphs and models. In Set theory and its applications (Toronto, ON, 1987), volume 1401 of Lecture Notes in Mathematics, pages 167–193. Springer, Berlin-New York, 1989. ed. Steprans, J. and Watson, S.
- [145] Saharon Shelah. The number of pairwise non-elementarily-embeddable models. The Journal of Symbolic Logic, 54:1431–1455, 1989.
- [146] Saharon Shelah. Incompactness for chromatic numbers of graphs. In A tribute to Paul Erdős, pages 361–371. Cambridge Univ. Press, Cambridge, 1990.

- [147] Saharon Shelah. Strong negative partition above the continuum. The Journal of Symbolic Logic, 55:21–31, 1990.
- [148] Saharon Shelah. Reflecting stationary sets and successors of singular cardinals. Archive for Mathematical Logic, 31:25–53, 1991.
- [149] Saharon Shelah. Strong Partition Relations Below the Power Set: Consistency, Was Sierpiński Right, II? In Proceedings of the Conference on Set Theory and its Applications in honor of A. Hajnal and V.T.Sos, Budapest, 1/91, volume 60 of Colloquia Mathematica Societatis Janos Bolyai. Sets, Graphs, and Numbers, pages 637–638. 1991, math.LO/9201244.
- [150] Saharon Shelah. Cardinal arithmetic for skeptics. American Mathematical Society. Bulletin. New Series, 26:197–210, 1992, math.LO/9201251.
- [151] Saharon Shelah. Viva la difference I: Nonisomorphism of ultrapowers of countable models. In Set Theory of the Continuum, volume 26 of Mathematical Sciences Research Institute Publications, pages 357–405. Springer Verlag, 1992, math.LO/9201245.
- [152] Saharon Shelah. Advances in Cardinal Arithmetic. In Finite and Infinite Combinatorics in Sets and Logic, pages 355–383. Kluwer Academic Publishers, 1993. N.W. Sauer et al (eds.).
- [153] Saharon Shelah. On $CH + 2^{\aleph_1} \to (\alpha)_2^2$ for $\alpha < \omega_2$. In Logic Colloquium'90. ASL Summer Meeting in Helsinki, volume 2 of Lecture Notes in Logic, pages 281–289. Springer Verlag, 1993, math.LO/9308212.
- [154] Saharon Shelah. Cardinal Arithmetic, volume 29 of Oxford Logic Guides. Oxford University Press, 1994.
- [155] Saharon Shelah. Cardinalities of topologies with small base. Annals of Pure and Applied Logic, 68:95–113, 1994, math.LO/9403219.
- [156] Saharon Shelah. How special are Cohen and random forcings i.e. Boolean algebras of the family of subsets of reals modulo meagre or null. *Israel Journal of Mathematics*, 88:159–174, 1994, math.LO/9303208.
- [157] Saharon Shelah. Possibly every real function is continuous on a non-meagre set. Publications de L'Institute Mathématique Beograd, Nouvelle Série, 57(71):47-60, 1995, math.LO/9511220.
- [158] Saharon Shelah. Further cardinal arithmetic. Israel Journal of Mathematics, 95:61–114, 1996, math.LO/9610226.
- [159] Saharon Shelah. Large Normal Ideals Concentrating on a Fixed Small Cardinality. Archive for Mathematical Logic, 35:341–347, 1996, math.LO/9406219.
- [160] Saharon Shelah. Colouring and non-productivity of N₂-cc. Annals of Pure and Applied Logic, 84:153–174, 1997, math.LO/9609218.
- [161] Saharon Shelah. Non existence of universals for classes like reduced torsion free abelian groups under non neccessarily pure embeddings. In Advances in Algebra and Model Theory. Editors: Manfred Droste und Ruediger Goebel, volume 9, pages 229–286. Gordon and Breach, 1997, math.LO/9609217.
- [162] Saharon Shelah. On σ -entangled linear orders. Fundamenta Mathematicae, 153:199–275, 1997, math.LO/9609216.
- [163] Saharon Shelah. Set Theory without choice: not everything on cofinality is possible. Archive for Mathematical Logic, 36:81–125, 1997, math.LO/9512227. A special volume dedicated to Prof. Azriel Levy.
- [164] Saharon Shelah. Proper and improper forcing. Perspectives in Mathematical Logic. Springer, 1998.
- [165] Saharon Shelah. There may be no nowhere dense ultrafilter. In Makowsky, J.A. and Ravve, E.V., editor, Proceedings of the Logic Colloquium Haifa'95, volume 11 of Lecture Notes in Logic, pages 305–324. Springer, 1998, math.LO/9611221.
- [166] Saharon Shelah, Jan Saxl, and Simon Thomas. Infinite products of finite simple groups. Transactions of the American Mathematical Society, 348:4611–4641, 1996, math.IG/9605202.
- [167] Saharon Shelah and Lee Stanley. Consistency of partition relation for cardinal in $(\lambda, 2^{\lambda})$. Journal of Symbolic Logic, submitted, math.LO/9710216.
- [168] Saharon Shelah and Lee Stanley. A theorem and some consistency results in partition calculus. Annals of Pure and Applied Logic, 36:119–152, 1987.
- [169] Saharon Shelah and Juris Steprans. Somewhere trivial automorphisms. Journal of the London Mathematical Society, 49:569–580, 1994, math.LO/9308214.

- [170] Saharon Shelah and Simon Thomas. The Cofinality Spectrum of The Infinite Symmetric Group. Journal of Symbolic Logic, 62:902–916, 1997, math.LO/9412230.
- [171] Saharon Shelah and Jindra Zapletal. Canonical models for ℵ₁ combinatorics. Transactions of the American Mathematical Society, accepted, math.LO/9806166.
- [172] Shelah, Saharon. On "I is a κ -complete ideal on κ such that $\mathcal{P}(\kappa)/I$ is the universal meager forcing".
- [173] Sławomir Solecki. Analytic ideals and their applications. Annals of Pure and Applied Logic, to appear.
- [174] Sławomir Solecki. Analytic ideals. Bulletin of Symbolic Logic, 2:339–348, 1996.
- [175] Robert M. Solovay. Real valued measurable cardinals. In Axiomatic set theory, volume XIII, Part 1 of Proc. of Symposia in Pure Math., pages 397–428, Providence R.I, 1971. A.M.S.
- [176] Juris Steprāns and Stephen W. Watson. Homeomorphisms of manifolds with prescribed behaviour on large dense sets. Bull. London Math. Soc., 19:305–310, 1987.
- [177] Simon Thomas. The cofinalities of the infinite-dimensional classical groups. J. Algebra, 179:704–719, 1996.
- [178] Stevo Todorčević. Partition problems in Topology, volume 84 of Contemporary Mathematics. American Mathematical Society, Providence, RI, 1989.
- [179] Jan van Mill. An Introduction to $\beta\omega$. In K. Kunen and J. E. Vaughan, editors, *Handbook of Set-Theoretic Topology*, pages 503–567. Elsevier Science Publishers, Amsterdam, 1984.
- [180] Boban Veličković. Applications of the open coloring axiom. In H. Judah, W. Just, and H. Woodin, editors, Set Theory of the Continuum (Berkeley, CA, 1989), volume 26 of Mathematical Sciences Research Institute Publications, pages 137–154. Springer Verlag, 1992.
- [181] Scott W. Williams. Box Products. In K. Kunen and J. E. Vaughan, editors, Handbook of Set-Theoretic Topology, pages 169–200. Elsevier Science Publishers, Amsterdam, 1984.
- [182] Hugh Woodin. The Axiom of Determinacy, Forcing Axioms and the Nonstationary Ideal, volume 1 of de Gruyter Series in Logic and its Applications. De Gruyter, in press.

Institute of Mathematics, The Hebrew University of Jerusalem, 91904 Jerusalem, Israel, and Department of Mathematics, Rutgers University, New Brunswick, NJ 08854, USA

E-mail address: shelah@math.huji.ac.il URL: http://www.math.rutgers.edu/~shelah