On elliptic solutions of the quintic complex one-dimensional Ginzburg–Landau equation

S. Yu. Vernov

Skobeltsyn Institute of Nuclear Physics, Moscow State University, Vorob'evy Gory, Moscow, 119992, Russia

Abstract

The Conte-Musette method has been modified for the search of only elliptic solutions to systems of differential equations. A key idea of this a priory restriction is to simplify calculations by means of the use of a few Laurent series solutions instead of one and the use of the residue theorem. The application of our approach to the quintic complex one-dimensional Ginzburg-Landau equation (CGLE5) allows us to find elliptic solutions in the wave form. We also find restrictions on coefficients, which are necessary conditions for the existence of elliptic solutions for the CGLE5. Using the investigation of the CGLE5 as an example, we demonstrate that to find elliptic solutions the analysis of a system of differential equations is more preferable than the analysis of the equivalent single differential equation.

Key words: Standing wave, the elliptic function, the Laurent series, the residue theorem, the quintic complex one-dimensional Ginzburg–Landau equation *PACS*: 05.04.-a, 02.30.-f, 02.70.Wz, 47.27.-i

1 Introduction

At present methods for construction of special solutions of nonintegrable systems in terms of elementary (more precisely, degenerated elliptic) and elliptic functions are actively developed [2,7,8,9,16,17,21,25,26,27,29,31,32,33,37,40,41,43,49] (see also [30] and references therein). Some of these methods are intended for the search for elliptic solutions only [26], others allow to find either solutions in terms of elementary functions only [7,8,31,49] or both types of solutions [2,9,16,17,21,25,27,29,32,33,37,40,41,43]. Note that the methods [40,29] allow to find multivalued solutions as well.

Email address: svernov@theory.sinp.msu.ru (S. Yu. Vernov).

Elliptic and degenerate elliptic functions are single-valued functions, therefore, such solutions of a nonintegrable system exist only if there exist the Laurent series solutions of it. Such local solutions can be constructed by means of the Ablowitz–Ramani–Segur algorithm of the Painlevé test [1] (see also [24,30,39]). Moreover for a wide class of dynamical systems using this method one can find all possible Laurent series expansions of solutions. In this way one obtains solutions only as formal series, that is sufficient, because really only a finite number of coefficients of these series is used. Examples of construction of such solutions are given in [31,42]. The Laurent series solutions give the information about the global behavior of differential systems and assist to look for both exact solutions [32] and the first integrals [28]. The Laurent series solutions can be used to prove the nonexistence of elliptic solutions [23,47] as well.

In [32] R. Conte and M. Musette have proposed a new method for construction of single-valued special solutions of nonintegrable differential equations. A key idea of this method is the use of the Laurent series solutions to transform the initial differential equation into a nonlinear system of algebraic equations. Using this method one can in principal find all elliptic and degenerate elliptic solutions. Unfortunately if the initial differential equation includes the large number of numeric parameters, then it is difficult to solve the obtained nonlinear system of algebraic equations.

The goal of this paper is to propose a modification of the Conte–Musette method, which allows to seek elliptic solutions only. We show that in this case it is possible to fix some parameters of the initial differential system and therefore simplify the resulting system of algebraic equations. To do this we use the Hone's method, which has been proposed to prove the non-existence of elliptic solutions [23]. Note that using our approach one can find in principal all elliptic solutions.

In [32] the authors have transformed the initial system of two coupled ordinary differential equations into the equivalent single differential equation and only after this have constructed the Laurent series solutions. In our paper we demonstrate that the analysis of the system of differential equations may be more useful than the analysis of the equivalent differential equation. Moreover in this paper we show that if a system of differential equations includes a few functions it is possible to find the analytic form of a function which satisfies this system even without knowledge of other functions in the analytic form and without elimination of them from the system.

2 The Conte–Musette method for the system of differential equations

In [32] R. Conte and M. Musette have proposed a way for searching of elliptic and degenerate elliptic solutions to a polynomial autonomous differential equation. In this section we reformulate this method for a system of such equations:

$$F_i(\vec{\mathbf{y}}_{;t}^{(n)}, \vec{\mathbf{y}}_{;t}^{(n-1)}, \dots, \vec{\mathbf{y}}_{;t}, \vec{\mathbf{y}}) = 0, \qquad i = 1, \dots, N,$$
(1)

where
$$\vec{\mathbf{y}} = \{y_1(t), y_2(t), \dots, y_L(t)\}$$
 and $y_{j;t}^{(k)} = \frac{d^k y_j}{dt^k}$.

It is known that any elliptic function (including degenerate one) is a solution of some first order polynomial autonomous differential equation. The classical results of P. Painlevé, L. von Fuchs, C.A.A. Briot and J.-C. Bouquet allow one to construct the suitable form of such an equation, whose general solution is a meromorphic function with poles of order p (see details in [32])¹:

$$\sum_{k=0}^{m} \sum_{j=0}^{(p+1)(m-k)/p} h_{j,k} y^{j} y_{t}^{k} = 0, \qquad h_{0,m} = 1,$$
(2)

in which m is a positive integer number and $h_{j,k}$ are constants to be determined. The general solution of (2) is either an elliptic function, or a rational function of $e^{\gamma x}$, γ being some constant, or a rational function of x. Note that the third case is a degeneracy of the second one, which in turn is a degenerate case of the first one.

The Conte–Musette algorithm is the following [32] (see also [9]):

- (1) Choose a positive integer number m, define the form of Eq. (2) and calculate the number of unknown coefficients $h_{j,k}$.
- (2) Construct solutions of system (1) in the form of the Laurent series. If such solutions do not exist or they correspond to known exact solutions, then no unknown single-valued solutions exist. Note that, since system (1) is autonomous, the coefficients of the Laurent series do not depend on the position of the singular point. They may depend on values of the numerical parameters of (1). In addition, some of these coefficients (the number of which is less than the order of system (1)) may take arbitrary values and have to be considered as new numerical parameters. One should compute more coefficients of the Laurent series than the

¹ The summation in (2) runs over nonnegative integer j that are less than or equal to (p+1)(m-k)/p

number of numerical parameters in the Laurent series plus the number of $h_{j,k}$.

- (3) Choose a Laurent series expansion for some function y_k and substitute the obtained Laurent series coefficients into Eq. (2). This substitution transforms (2) into a linear and overdetermined system in $h_{j,k}$ with coefficients depending on numerical parameters.
- (4) Eliminate coefficients $h_{j,k}$ and get a nonlinear system in parameters.
- (5) Solve the obtained nonlinear system.

R. Conte and M. Musette note that a computer algebra package is highly recommended for using of their method [9]. Steps of this algorithm can be implemented in computer algebra systems separately.

For the given system it is easy to calculate the Laurent series solutions to any accuracy. These computations base on the Painlevé test, which has been implemented in the most popular computer algebra systems [4,34,38,50]. Note that when one has computed a sufficient number of the Laurent series coefficients he can forget about the system of differential equations and work only with coefficients of the obtained series. The first package of computer algebra procedures, which realize the third and the fourth steps of the algorithm, has been written in AMP [13] by R. Conte. One can also use our Maple and RE-DUCE packages of procedures, which are accessible in Internet [44] and are described in [45,46]. So, one passes the first four steps of algorithm without any difficulties.

At the fifth (last) step one should solve an overdetermined system of nonlinear algebraic equations. The standard method for solving of such systems is the construction of a lexicographically ordered Gröbner basis [11]. The Buchberger algorithm [5,6,11] guarantees a construction of a Gröbner basis in a finite number of steps². However, appropriate estimates of the number of steps required for constructing of this basis do not exist. The required computer memory depends, in the general case, exponentially on the number of the unknowns. Therefore, this number should be made as small as possible.

The purpose of this paper is to show that we can essentially simplify the algebraic system of equations, which we have to solve on the last step of the Conte–Musette method, if we search the elliptic solutions only.

In [32] R. Conte and M. Musette have used their method to find wave solutions of the complex cubic Ginzburg–Landau equation (CGLE3). The nonexistence of elliptic travelling and standing wave solutions of the CGLE3 has been proved in [23] and [47] respectively. In Section 4 we seek the elliptic solutions of the quintic complex Ginzburg–Landau equation (CGLE5) using our modification

 $^{^{2}}$ An alternative method for constructing Gröbner bases is the involutive algorithm [18,19].

of the Conte–Musette method. Note that both the CGLE3 and the CGLE5 have only one-parameter Laurent-series solutions in the wave form and there exist only a finite number of such solutions.

Our approach can be effectively used in investigation of any system (1) or a single differential equation, for which only a finite number of different Laurentseries solutions exist. Note that for wide class of such differential equations it has been proved that all their meromorphic solutions are elliptic (maybe degenerated) functions [15].

3 Properties of the elliptic functions

Let us recall some definitions and theorems. The function $\rho(z)$ of the complex variable z is a doubly-periodic function if there exist two numbers ω_1 and ω_2 with $\omega_1/\omega_2 \notin \mathbb{R}$, such that for all $z \in \mathbb{C}$

$$\varrho(z) = \varrho(z + \omega_1) = \varrho(z + \omega_2). \tag{3}$$

By definition a double-periodic meromorphic function is called an elliptic function [14,22]. These periods define the period parallelograms with vertices z_0 , $z_0 + N_1\omega_1$, $z_0 + N_2\omega_2$ and $z_0 + N_1\omega_1 + N_2\omega_2$, where N_1 and N_2 are arbitrary natural numbers and z_0 is an arbitrary complex number. The fundamental parallelogram of periods is called a parallelogram of period, which does not include other parallelogram of periods, that corresponds to $N_1 = N_2 = 1$.

The classical theorems for elliptic functions [14,22] prove that

- If an elliptic function has no poles then it is a constant.
- The number of elliptic function poles within any finite period parallelogram is finite.
- The sum of residues within any finite period parallelogram is equal to zero (the residue theorem).
- If $\rho(z)$ is an elliptic function then any rational function of $\rho(z)$ and its derivatives is an elliptic function as well.
- For each elliptic function $\rho(z)$ there exist such $m \ (m \ge 2)$ and such coefficients $h_{i,j}$ that $\rho(z)$ is a solution of Eq. (2).

Lemma 1 An elliptic function can not have two poles with the same Laurent series expansions in its fundamental parallelogram of periods.

Proof.

Let some elliptic function $\varrho(\xi)$ has two poles in points ξ_0 and ξ_1 , which belong

to the fundamental parallelogram of periods. The corresponding Laurent series are the same and have the convergence radius R. Then the function $v(\xi) = \rho(\xi - \xi_0) - \rho(\xi - \xi_1)$ is an elliptic function as a difference between two elliptic functions with the same periods. At the same time for all ξ such that $|\xi| < R$ $v(\xi) = 0$, therefore, $v(\xi) \equiv 0$ and $\rho(\xi - \xi_0) \equiv \rho(\xi - \xi_1)$ and $\xi_1 - \xi_0$ is a period of $\rho(\xi)$. It contradicts to our assumption that both points ξ_0 and ξ_1 belong to the fundamental parallelogram of periods.

4 Construction of elliptic solutions

4.1 The quintic complex Ginzburg–Landau equation

The one-dimensional quintic complex Ginzburg–Landau equation (CGLE5) is a generalization of the one-dimensional cubic complex Ginzburg–Landau equation [20] (CGLE3), which is one of the most-studied nonlinear equations (see [3] and references therein). Moreover, the CGLE5 is a generic equation which describes many physical phenomena, for example, the behaviour of travelling patterns in binary fluid convection [36] and the large-scale behavior of many nonequilibrium pattern-forming systems [10].

The CGLE5 is as follows

$$i\mathcal{A}_t + p\mathcal{A}_{xx} + q|\mathcal{A}|^2\mathcal{A} + r|\mathcal{A}|^4\mathcal{A} - i\gamma\mathcal{A} = 0,$$
(4)

where subscribes denote partial derivatives: $\mathcal{A}_t \equiv \frac{\partial \mathcal{A}}{\partial t}, \ \mathcal{A}_{xx} \equiv \frac{\partial^2 \mathcal{A}}{\partial x^2}, \ p, q, r \in \mathbb{C}$ and $\gamma \in \mathbb{R}$.

One of the most important directions in the study of the CGLE5 is the consideration of its travelling wave reduction [2,9,12,31,35,36]:

$$\mathcal{A}(x,t) = \sqrt{M(\xi)} e^{i(\varphi(\xi) - \omega t)}, \quad \xi = x - ct, \qquad c \in \mathbb{R}, \quad \omega \in \mathbb{R}.$$
(5)

Substituting (5) in (4) and multiplying both sides of this equation on $4M^2/A$ we obtain

$$2pM''M - pM'^{2} + 4ip\psi MM' + 2(2\omega - ic - 2i\gamma + + 2c\psi - 2p\psi^{2} + 2ip\psi')M^{2} + 4qM^{3} + 4rM^{4} = 0,$$
(6)

where $\psi \equiv \varphi' \equiv \frac{d\varphi}{d\xi}$, $M' \equiv \frac{dM}{d\xi}$. Equation (6) is a system of two equations: both real and imaginary parts of its left-hand side have to be equal to zero.

Dividing (6) by p and separating real terms from imaginary ones, we obtain the following system

$$\begin{cases} 2MM'' - M'^2 - 4M^2\tilde{\psi}^2 - 2\tilde{c}MM' + 4g_iM^2 + 4d_rM^3 + 4u_rM^4 = 0, \\ M\tilde{\psi}' + \tilde{\psi}\left(M' - \tilde{c}M\right) - g_rM + d_iM^2 + u_iM^3 = 0, \end{cases}$$
(7)

where new real variables are as follows

$$u_r + iu_i = \frac{r}{p}, \qquad d_r + id_i = \frac{q}{p}, \qquad s_r - is_i = \frac{1}{p}, \tag{8}$$

$$g_r + ig_i = (\gamma + i\omega)(s_r - is_i) + \frac{1}{2}c^2s_is_r + \frac{i}{4}c^2s_r^2,$$
(9)

and

$$\tilde{\psi} \equiv \psi - \frac{cs_r}{2}, \qquad \tilde{c} \equiv cs_i.$$
(10)

System (7) includes seven numerical parameters: g_r , g_i , d_r , d_i , u_r , u_i and \tilde{c} . Note that to obtain (7) from (6) we assume that the functions $M(\xi)$ and $\psi(\xi)$ are real.

The standard way to construct exact solutions for system (7) is to transform it into the equivalent third order differential equation for M. We rewrite the first equation of system (7) as

$$\tilde{\psi}^2 = \frac{G}{M^2},\tag{11}$$

where

$$G \equiv \frac{1}{2}MM'' - \frac{1}{4}M'^2 - \frac{\tilde{c}}{2}MM' + g_iM^2 + d_rM^3 + u_rM^4.$$
 (12)

From (11) it follows that

$$\tilde{\psi}'\tilde{\psi} = \frac{G'M - 2GM'}{2M^3},\tag{13}$$

Multiplying the second equation of (7) on $\tilde{\psi}$ and substituting (11) and (13) in it, we express $\tilde{\psi}$ in terms of M and its derivatives:

$$\tilde{\psi} = \frac{G' - 2\tilde{c}G}{2M^2 \left(g_r - d_i M - u_i M^2\right)},$$
(14)

and obtain the third order equation for M:

$$(G' - 2\tilde{c}G)^2 + 4GM^2(g_r - d_iM - u_iM^2)^2 = 0.$$
(15)

4.2 The Laurent series solutions

Below we consider the case

$$\frac{p}{r} \notin \mathbb{R},\tag{16}$$

which corresponds to the condition $u_i \neq 0$. In this case Eq. (15) is not integrable [31] and its general solution (which should depend on three arbitrary integration constants) is not known. Using the Painlevé analysis [31] it has been shown that single-valued solutions of (7) can depend on only one arbitrary parameter. System (7) is autonomous, so this parameter is ξ_0 : if $M = f(\xi)$ is a solution, then $M = f(\xi - \xi_0)$, where $\xi_0 \in \mathbb{C}$ has to be a solution. Special solutions in terms of elementary functions have been found in [2,9,31,35]. All known exact solutions of (7) are elementary (rational or hyperbolic) functions. The full list of these solutions is presented in [9]. The purpose of this section is to find at least one elliptic solution of (7).

System (7) is invariant under the transformation:

$$\tilde{\psi} \to -\tilde{\psi}, \quad g_r \to -g_r, \quad d_i \to -d_i, \quad u_i \to -u_i,$$
(17)

therefore we can assume that $u_i > 0$ without loss of generality. Moreover, using scale transformations:

$$M \to \lambda M, \quad d_r \to \frac{d_r}{\lambda}, \quad d_i \to \frac{d_i}{\lambda}, \quad u_r \to \frac{u_r}{\lambda^2}, \quad u_i \to \frac{u_i}{\lambda^2},$$
 (18)

we can always put $u_i = 1$.

Let us construct the Laurent series solutions to system (7). We assume that in a sufficiently small neighborhood of the singularity point ξ_0 the functions $\tilde{\psi}$ and M tend to infinity as some powers of $\xi - \xi_0$:

$$\widetilde{\psi} = A(\xi - \xi_0)^{\alpha} \quad \text{and} \quad M = B(\xi - \xi_0)^{\beta},$$
(19)

where α and β are negative integer numbers and, of course, $A \neq 0$ and $B \neq 0$. Substituting (19) into (7) we obtain that two or more terms in the equations of system (7) balance if and only if $\alpha = -1$ and $\beta = -1$. In other words in this case these terms have equal powers and the other terms can be ignored as $t \longrightarrow t_0$. We obtain values of A and B from the following algebraic system:

$$\begin{cases} B^2 \left(3 - 4A^2 + 4u_r B^2 \right) = 0, \\ 2A - B^2 = 0. \end{cases}$$
(20)

System (20) has four nonzero solutions:

$$A_1 = u_r + \frac{1}{2}\sqrt{4u_r^2 + 3}, \qquad B_1 = \sqrt{2u_r + \sqrt{4u_r^2 + 3}},$$
 (21)

$$A_2 = u_r + \frac{1}{2}\sqrt{4u_r^2 + 3}, \qquad B_2 = -\sqrt{2u_r + \sqrt{4u_r^2 + 3}}, \tag{22}$$

$$A_3 = u_r - \frac{1}{2}\sqrt{4u_r^2 + 3}, \qquad B_3 = \sqrt{2u_r - \sqrt{4u_r^2 + 3}}$$
 (23)

and

$$A_4 = u_r - \frac{1}{2}\sqrt{4u_r^2 + 3}, \qquad B_4 = -\sqrt{2u_r - \sqrt{4u_r^2 + 3}}.$$
 (24)

Therefore, system (7) has four types of the Laurent series solutions. Denote them as follows:

$$\tilde{\psi}_k = \frac{A_k}{\xi} + a_{k,0} + a_{k,1}\xi + \dots, \qquad M_k = \frac{B_k}{\xi} + b_{k,0} + b_{k,1}\xi + \dots, \tag{25}$$

where k = 1..4.

Let $M(\xi)$ is a nontrivial elliptic function. Note that if $\tilde{\psi}$ is a constant, then from the second equation of system (7) it follows that M can not be a nontrivial elliptic function. Therefore, using (14), we conclude that $\tilde{\psi}(\xi)$ has to be a nontrivial elliptic function as well.

Let us consider the fundamental parallelogram of periods for the function $M(\xi)$ and define a number of its poles in this domain. Let M has a pole of type M_1 , hence, according to the residue theorem, it should has a pole of type M_2 (it can not have a pole of type M_4 because u_r is a real parameter). So $\tilde{\psi}$ has poles with the Laurent series $\tilde{\psi}_1$ and $\tilde{\psi}_2$. As an elliptic function it should have a pole of type $\tilde{\psi}_3$ or $\tilde{\psi}_4$ as well. It means that the function $M(\xi)$ should have a pole of type M_3 and, hence, a pole of type M_4 . So $M(\xi)$ should have at least four different poles in its the function $M(\xi)$ can not have the same poles in

the fundamental parallelogram of periods. Therefore, $M(\xi)$ has exactly four poles in its fundamental parallelogram of periods. In this case by means of the residue theorem for $\tilde{\psi}$ we obtain

$$u_r = 0. (26)$$

We obtain that the CGLE5 with $u_r \neq 0$ has no elliptic solution in the wave form. In the case $u_r = 0$ possible elliptic solutions should have four simple poles in the fundamental parallelogram of periods, and, therefore, has the following form [22]:

$$M(\xi - \xi_0) = C + \sum_{k=1}^{4} B_k \zeta(\xi - \xi_k),$$
(27)

where the function $\zeta(\xi)$ is an integral of the Weierstrass elliptic function multiplied by -1:

$$\zeta'(\xi) = -\wp(\xi),\tag{28}$$

C and ξ_k are constants to be defined. We also should define periods of the Weierstrass elliptic function.

To obtain restrictions on other parameters, we use the Hone's method [23] and apply the residue theorem to the functions $\tilde{\psi}^2$, $\tilde{\psi}^3$, and so on. The residue theorem for the function $\tilde{\psi}^2$ gives the equation:

$$\sum_{k=1}^{4} A_k a_{k,0} = 0.$$
⁽²⁹⁾

The values of $a_{k,0}$ are as follows $(u_r = 0)$:

$$a_{1,0} = \frac{\sqrt{3}}{48} \left(6\tilde{c} - \sqrt[4]{27}d_i - 15\sqrt[4]{3}d_r \right), \tag{30}$$

$$a_{2,0} = \frac{\sqrt{3}}{48} \left(6\tilde{c} + \sqrt[4]{27}d_i + 15\sqrt[4]{3}d_r \right), \tag{31}$$

$$a_{3,0} = -\frac{\sqrt{3}}{48} \left(6\tilde{c} + i \left(\sqrt[4]{27}d_i - 15\sqrt[4]{3}d_r \right) \right),$$
(32)

$$a_{4,0} = -\frac{\sqrt{3}}{48} \left(6\tilde{c} - i \left(\sqrt[4]{27}d_i - 15\sqrt[4]{3}d_r \right) \right).$$
(33)

Substituting A_k and $a_{k,0}$ in (29), we obtain

$$\sum_{k=1}^{4} A_k a_{k,0} = \frac{3}{4} \tilde{c} = 0, \tag{34}$$

therefore $\tilde{c} = 0$.

For the function $\tilde{\psi}^3$ the residue theorem gives:

$$\sum_{k=1}^{4} A_k \left(A_k a_{k,1} + a_{k,0}^2 \right) = 0.$$
(35)

The values $a_{k,1}$ are as follows (we put $\tilde{c} = 0$):

$$a_{1,1} = \frac{1}{384} \left(3d_i^2 - 63d_r^2 - 66\sqrt{3}d_i d_r + 128\sqrt{3}g_i \right), \qquad a_{2,1} = a_{1,1}, \quad (36)$$

$$a_{3,1} = \frac{1}{384} \left(3d_i^2 - 63d_r^2 + 66\sqrt{3}d_i d_r - 128\sqrt{3}g_i \right), \qquad a_{4,1} = a_{3,1}.$$
(37)

Equation (35) is equivalent to

$$d_i^2 + 27d_r^2 = 0 \quad \to \quad d_i = \pm i\sqrt{27}d_r.$$
 (38)

The parameters d_r and d_i should be real, therefore, $d_r = 0$ and $d_i = 0$. So, consideration of $\tilde{\psi}^2$ and $\tilde{\psi}^3$ gives three restrictions:

$$\tilde{c} = 0, \qquad d_r = 0 \qquad \text{and} \qquad d_i = 0. \tag{39}$$

The residue theorem for $\tilde{\psi}^4$ gives the restriction

$$g_i g_r = 0. (40)$$

Considering $\tilde{\psi}^5$ and $\tilde{\psi}^6$, we do not obtain new restrictions on coefficients. Taking into account (26) and (39) we obtain system (7) in the following form:

$$\begin{cases} 2MM'' - {M'}^2 - 4M^2 \tilde{\psi}^2 + 4g_i M^2 = 0, \\ \tilde{\psi}'M + \tilde{\psi}M' - g_r M + M^3 = 0. \end{cases}$$
(41)

To find elliptic solutions to system (41) we use the Conte–Musette method. The function $\tilde{\psi}(\xi)$ can have not four but two different Laurent series expansions, whereas the functions $M(\xi)$ should have four different Laurent series expansions, so it is easier to find $\tilde{\psi}(\xi)$ than $M(\xi)$. Equation (2) with m = 1 has no elliptic solution. Let $\tilde{\psi}(\xi)$ satisfies Eq. (2) with m = 2:

$$\tilde{\psi'}^{2} + \left(\tilde{h}_{2,1}\tilde{\psi}^{2} + \tilde{h}_{1,1}\tilde{\psi} + \tilde{h}_{0,1}\right)\tilde{\psi'} + \\ + \tilde{h}_{4,0}\tilde{\psi}^{4} + \tilde{h}_{3,0}\tilde{\psi}^{3} + \tilde{h}_{2,0}\tilde{\psi}^{2} + \tilde{h}_{1,0}\tilde{\psi} + \tilde{h}_{0,0} = 0.$$

$$\tag{42}$$

Substituting in (42) the Laurent series of $\tilde{\psi}$, which begins from A_1 (more exactly we use the first ten coefficients), we obtain the following solution $\tilde{h}_{k,j}$ for an arbitrary value of the parameter $g_r \neq 0$ and $g_i = 0$:

$$\tilde{h}_{4,0} = -\frac{4}{3}, \ \tilde{h}_{0,0} = -\frac{g_r^2}{9}, \ \tilde{h}_{3,0} = \tilde{h}_{2,0} = \tilde{h}_{1,0} = \tilde{h}_{0,1} = \tilde{h}_{1,1} = \tilde{h}_{2,1} = 0, (43)$$

a few solutions with $g_i = 0$ and $g_r = 0$ and no solution for $g_i \neq 0$.

The straightforward substitution of the functions

$$\breve{\psi} = \frac{\sqrt{3}}{2t}, \qquad \breve{M} = \pm \frac{\sqrt[4]{3}}{t}, \tag{44}$$

or

$$\hat{\psi} = -\frac{\sqrt{3}}{2t}, \qquad \hat{M} = \pm i\frac{\sqrt[4]{3}}{t}$$
(45)

in (41) with $g_r = 0$ and $g_i = 0$ proves that they are exact solutions. The coefficients of the Laurent-series solutions do not include arbitrary parameters, so the obtained solutions are unique single-valued solutions and system (41) has no elliptic solution for these values of parameters.

In the case of solutions (43) the function $\tilde{\psi}(\xi)$ satisfies the equation

$$\tilde{\psi'}^2 = \frac{4}{3}\tilde{\psi}^4 + \frac{g_r^2}{9}.$$
(46)

The polynomial in the right hand side of (46) has four different roots, therefore $\tilde{\psi}$ is a non-degenerate elliptic function [22].

Surely we do not rigorously prove the existence of elliptic solutions to the CGLE5. More precisely, we calculate fifty coefficients of the Laurent series of the function $\tilde{\psi}$, which satisfies (41) with $g_i = 0$ and find that they coincide with corresponding coefficients of the Laurent series of the exact solution to Eq. (46).

For rigorous proof we should find the function $M(\xi)$ and check that this function is a solution of (15). The function $M(\xi)$ in a parallelogram of periods has four different Laurent series expansions, so we should choose the parameter msuch that solutions of Eq. (2) have four poles in its fundamental parallelogram of periods. Minimal possible value of m is equal to 4. The general form of (2) for m = 4 and p = 1 is the following:

$$M^{\prime 4} + (h_{2,3}M^2 + h_{1,3}M + h_{0,3})M^{\prime 3} + + (h_{4,2}M^4 + h_{3,2}M^3 + h_{2,2}M^2 + h_{1,2}M + h_{0,2})M^{\prime 2} + + (h_{6,1}M^6 + h_{5,1}M^5 + h_{4,1}M^4 + h_{3,1}M^3 + h_{2,1}M^2 + h_{1,1}M + + h_{0,1})M^\prime + h_{8,0}M^8 + h_{7,0}M^7 + h_{6,0}M^6 + h_{5,0}M^5 + + h_{4,0}M^4 + h_{3,0}M^3 + h_{2,0}M^2 + h_{1,0}M + h_{0,0} = 0.$$
(47)

Substituting the Laurent series M_k from (25), we transform the left hand side of (47) into the Laurent series, which has to be equal to zero. Therefore, we obtain the algebraic system in $h_{i,j}$ and g_r . The first algebraic equation, which corresponds to $1/\xi^8$ is

$$B_k^4 \left(h_{8,0} B_k^4 - h_{6,1} B_k^3 + h_{4,2} B_k^2 - h_{2,3} B_k + 1 \right) = 0, \tag{48}$$

where B_k is defined by (21)–(24). If we would attempt to find all elliptic and degenerate elliptic solutions then we should use only one of B_k and can express, for example, $h_{8,0}$ via $h_{6,1}$, $h_{4,2}$ and $h_{2,3}$. We seek only elliptic solutions, so we know that all B_k have to satisfy (48) and can consider Eq. (48) as the following system:

$$\begin{cases} h_{8,0}B_1^4 - h_{6,1}B_1^3 + h_{4,2}B_1^2 - h_{2,3}B_1 + 1 = 0, \\ h_{8,0}B_2^4 - h_{6,1}B_2^3 + h_{4,2}B_2^2 - h_{2,3}B_2 + 1 = 0, \\ h_{8,0}B_3^4 - h_{6,1}B_3^3 + h_{4,2}B_3^2 - h_{2,3}B_3 + 1 = 0, \\ h_{8,0}B_4^4 - h_{6,1}B_4^3 + h_{4,2}B_4^2 - h_{2,3}B_4 + 1 = 0. \end{cases}$$

$$(49)$$

Using the explicit values of B_k from (21)–(24), we obtain that

$$h_{8,0} = -\frac{1}{3}, \quad h_{4,2} = 0, \quad h_{6,1} = 0, \quad h_{2,3} = 0.$$
 (50)

Taking into account (50), from other equations of the algebraic system we obtain

$$h_{6,0} = \frac{4}{3}g_r, \quad h_{4,0} = -\frac{16}{9}g_r^2, \quad h_{2,0} = \frac{64}{81}g_r^3,$$
 (51)

all other $h_{i,j}$ are equal to zero. So, the equation for M has the form

$$M'^{4} = \frac{1}{81}M^{2} \left(3M^{2} - 4g_{r}\right)^{3}.$$
(52)

Equation (15) at $u_i = 1$, $u_r = 0$, $\tilde{c} = 0$, $d_r = 0$, $d_i = 0$ and $g_i = 0$ has the form:

$$\frac{1}{4} \left(M''' \right)^2 - \left(2MM'' - M'^2 \right) \left(M^2 - g_r \right)^2 = 0.$$
(53)

We multiply Eq. (53) on M'^2 and use Eq. (52) to express all derivatives of $M(\xi)$ in terms of the function $M(\xi)$. The straightforward calculation shows that any solution of (52) satisfies (53). So, we obtain elliptic wave solutions of the CGLE5. If $s_i \neq 0$ these solutions are the standing wave solutions, in the opposite case $(s_i = 0)$ the solutions can have an arbitrary speed c.

Note that we obtain (15) from (6) using the condition that $M(\xi)$ is a real function. For $g_r < 0$ and any initial value of M we obtain real solutions. In the case $g_r > 0$ there exists the minimal possible initial value of M for which real solutions exist and only particular solutions of (52) are suitable elliptic solutions to the CGLE5. The function $M(\xi)$ has the form (27), the values of constants can be determined from (52).

Summing up we can conclude that our modification of the Conte–Musette method allows us to get two results: we obtain new elliptic wave solutions of the CGLE5, and we prove that these solutions are unique elliptic solutions for the CGLE5 with $g_r \neq 0$.

From (40) it follows that elliptic solutions can exist if g_r or g_i is equal to zero. For all nonzero values of g_r and zero g_i we have found elliptic solutions. In the case $g_r = g_i = 0$ there is no elliptic solution. In the case of zero g_r and nonzero g_i we substitute the obtained Laurent series solutions M_k into Eq. (2) with $m = 1, \ldots, 4$ and do not obtain neither elliptic functions nor degenerate elliptic solutions. We hope that the more detail analysis of this case allows us to find all elliptic solutions for the CGLE5.

5 Construct of elliptic solutions for nonintegrable systems

The approach, which we have considered in the previous section, can be applicable to many nonintegrable systems. The best applicability is nonintegrable systems of autonomous nonlinear ordinary differential equations with so-called finiteness property [15]: there is only a finite number of formal Laurent series that satisfy the system. For such systems we can propose the following way to the search for elliptic solutions:

- (1) Calculate a few first terms of all solutions of system (1) in the form of the Laurent series.
- (2) Choose the function y_k , which should be elliptic. Check should other functions be elliptic or not.
- (3) Using the residue theorem define values of numeric parameters at which the solution y_k can be an elliptic function.
- (4) Define a minimal number of poles for candidates to elliptic solutions. Using this number, choose a positive integer m, define the form of Eq. (2) and calculate the number of unknown coefficients $h_{j,k}$.
- (5) Calculate the sufficient number of coefficients for all Laurent series of y_k and substitute the obtained coefficients into Eq. (2). This substitution transforms (2) into a linear and overdetermined system in $h_{j,k}$ with coefficients depending on numerical parameters.
- (6) Eliminate coefficients $h_{i,k}$ and get a nonlinear system in parameters.
- (7) Solve the obtained nonlinear system.

We restrict ourselves to the search of elliptic solutions only. This makes possible to consider not one Laurent series, but as much as possible. This is a key idea of our modification of the Conte–Musette method.

Using the Conte–Musette method in the original form, one in principal would be able to find the elliptic solutions of the CGLE5, but calculations would be cumbersome, because he should reobtain all known solutions in terms of elementary functions to obtain new elliptic solutions. One has to construct and to solve an algebraic system, which includes at least 32 equations, which are linear in 24 unknowns $h_{i,j}$ and nonlinear in 7 parameters. In our approach, we seek only elliptic solutions and, using the Hone's method, reduce the number of arbitrary numerical parameters to 1. Consideration of a few Laurent series, instead of one also simplifies the obtained system of algebraic equations. We conclude that the use of the Conte–Musette method in combination with the Hone's method is very effective.

In contrast to [23,32] we consider a system of differential equations instead of the equivalent single differential equation and demonstrate that the system is more convenient for analysis than a single equation for M. If we analyse only the Laurent series expansions of the function M, then we should also consider the case when M has not four, but only two different Laurent series expansions, beginning from B_1 and B_2 or B_3 and B_4 . The consideration of the Laurent series expansions of two functions: $\tilde{\psi}$ and M, allows us to reject this possibility. Note that in our opinion if one seek degenerate elliptic solutions using the Laurent series then a system of equations is more convenient for analysis than the equivalent single equation as well.

Moreover, in contrast to traditional methods when we use the Conte–Musette method we can choose a function, which analytic form should be found. Instead of the functions $\tilde{\psi}$ and M we can consider some combination, for example, a polynomial, of this functions and seek this combination in the analytic form. Note that we have no need of any differential equation for this combination. So, we can conclude that the use of the Laurent series solutions gives new insight on problem of construction of exact solutions for nonintegrable systems.

6 Comparison with standard methods

Let us compare our approach with other methods for construction of special solutions of nonintegrable systems (see [9] as a review of these methods).

Without the use of the Laurent series solutions it would be very difficult to find the elliptic solutions of the CGLE5, because the form of Eq. (47) is very complex. For example, the Fan's technology [16,17] allows to find solutions of nonintegrable systems, which are polynomials of the function ρ , that satisfies the following equation:

$${\varrho'}^2 = \sum_{j=0}^N c_j \varrho^j,\tag{54}$$

where N and c_j are constants to be determined. The function $\tilde{\psi}$ is a solution of a similar equation with N = 4, but, using the Fan's approach, it is impossible to find $\tilde{\psi}(\xi)$ without knowledge of $M(\xi)$, which satisfies the more complex equation (47). The Kudryashov's method [26] as well as the methods [9,16,17,21,25,27,32,33,37,40,41,43], proposed to search both elliptic and elementary solutions, attempt to find solutions for the initial differential system at all values of the numeric parameters such that single-valued solutions can exist. The use of the residue theorem and the Hone's method allows us to fix some of these parameters and to simplify calculations without loss of elliptic solutions. Note that we not only find elliptic solutions for $g_r \neq 0$. Using the standard methods one can say nothing about uniqueness of the obtained

special elliptic solutions.

To find solutions for the CGLE5 the authors of papers [2,31,35] put some restrictions on dependence between $\tilde{\psi}$ and M. The use of Laurent solutions allows us to search $\tilde{\psi}$ without any restrictions and without eliminating M from system (7).

7 Conclusions

In this paper we propose a new approach for the search of elliptic solutions to systems of differential equations. The proposed algorithm is a modification of the Conte–Musette method [32]. We restrict ourselves to the search of elliptic solutions only. A key idea of this restriction is to simplify calculations by means of the use of a few Laurent series solutions instead of one and the use of the residue theorem.

The application of our approach to the quintic complex one-dimensional Ginzburg–Landau equation (CGLE5) allows to find elliptic solutions in the wave form. Let us point out that the obtained solutions are the first elliptic solutions for the CGLE5. Using the investigation of the CGLE5 as an example, we demonstrate that to find elliptic solutions the analysis of a system of differential equations is more preferable than the analysis of the equivalent single differential equation.

We also find restrictions on coefficients, which are necessary conditions for the existence of elliptic solutions for the CGLE5. To do this we develop the Hone's method [23]. We show that this method is useful not only to prove the nonexistence of elliptic solutions, but also to find new elliptic solutions. Note that the Hone's method and, therefore, our approach, are so effective in the case of the CGLE5, because coefficients of the Laurent series solutions depend only on parameters of equations, i.e. they do not include additional arbitrary parameters (have no resonances). It is an important problem to generalize the Hone's method on the Laurent series solutions with resonances.

Another way for future investigations is the generalization of the Conte-Musette method on the case of multivalued solutions. Some results in this direction have been obtained in [43,48].

Acknowledgements

This work has been supported in part by Russian Federation President's Grant NSh-1685.2003.2.

References

- M.J. Ablowitz, A. Ramani, H. Segur, A Connection between Nonlinear Evolution Equations and Ordinary Differential Equations of P-type. I & II, J. Math. Phys. 21 (1980) 715–721, & 1006–1015.
- [2] N.N. Akhmediev, V.V. Afanasjev, J.M. Soto-Crespo, Singularities and spesial soliton solutions of the cubic-quintic complex Ginzburg–Landau equation, *Rev. Phys. E* 53 (1996) 1190–1201.
- [3] I. Aranson, L. Kramer, The World of the Complex Ginzburg–Landau Equation, Rev. Mod. Phys. 74 (2002) 99–143; cond-mat/0106115.
- [4] D. Baldwin, W. Hereman, Symbolic Software for the Painleve Test of Nonlinear Ordinary and Partial Differential Equations, J. Nonlinear Math. Phys. 13 (2006) No. 1, 90–110; nlin.SI/0505004.
- [5] B. Buchberger, Theoretical Basis for the Reduction of Polynomials to Canonical Forms, SIGSAM Bulletin 39 (1976) 19–29.
- B. Buchberger, Some Properties of Gröbner–Bases for Polynomial Ideals, SIGSAM Bulletin 40 (1976) 19–24.
- [7] R. Conte, M. Musette, Link Between Solitary Waves and Projective Riccati Equations, J. Phys. A 25 (1992) 5609–5623.
- [8] R. Conte, M. Musette, Linearity inside nonlinearity: exact solutions to the complex Ginzburg–Landau equation, *Physica D* **69** (1993) 1–17.
- [9] R. Conte, M. Musette, Solitary waves of nonlinear nonintegrable equations, chapter in the book "Dissipative solitons" (Lect. Notes Phys., Vol. 661), eds. N.N. Akhmediev and A. Ankiewicz, Springer, Berlin, 2005; nlin.PS/0407026
- [10] M.C. Cross, P.C. Hohenberg, Pattern formation outside of equilibrium, Rev. Mod. Phys. 65 (1993) 851–1112.
- [11] J. Davenport, Y. Siret, E. Tournier, Calcul Formel, Systemes et Algorithmes de Manipulations Algebriques, Masson, Paris, New York, 1987.
- [12] A. Doelman, Breaking the hidden symmetry in the Ginzburg–Landau equation, *Physica D* 97 (1996) 398–428.
- [13] J.-M. Drouffe, Simplex AMP reference manual, version 1.0 (1996), SPhT, CEA Sacley, F-91191 Gif-sur-Yvette Cedex, 1996.

- [14] A. Erdélyi et al. (eds.), Higher Transcendental Functions (based, in part, on notes left by H. Bateman), Vol. 3, MC Graw-Hill Book Company, New York, Toronto, London, 1955.
- [15] A.E. Eremenko, Meromorphic traveling wave solutions of the Kuramoto-Sivashinsky equation, nlin.SI/0504053.
- [16] E. Fan, Multiple travelling wave solutions of nonlinear evolutions equations using a unified algebraic methods, J. of Phys. A: Math. Gen. 35 (2002) 6853– 6872.
- [17] E. Fan, An algebraic method for finding a series of exact solutions to integrable and nonintegrable nonlinear evolutions equations, J. of Phys. A: Math. Gen. 36 (2003) 7009–7026.
- [18] V.P. Gerdt, Yu.A. Blinkov, Involutive Bases of Polynomial Ideals, Math. Comp. Simul. 45 (1998) 519–542; math.AC/9912027; Minimal Involutive Bases, Math. Comp. Simul. 45 (1998) 543–560; math.AC/9912029.
- [19] V.P. Gerdt, Involutive Algorithms for Computing Gröbner Basis, In "Computational Commutative and Non-commutative Algebraic Geometry", S. Cojocaru, G. Pfister, V. Ufnarovski (eds.), NATO Science Series, IOP Press, 2005, pp. 199–225; math.AC/0501111.
- [20] V.L. Ginzburg, L.D. Landau, On the theory of superconductors, Zh. Eksp. Teor. Fiz. (Sov. Phys. JETP) 20 (1950) 1064–1082; English translation in L.D. Landau, Collected Papers, Oxford, Pergamon Press, 1965, p. 546.
- [21] R. Grimshaw, M. Pavlov, Exact periodic steady solutions for nonlinear wave equations: A new approach, *Phys. Lett. A* 251 (1999) 25–30.
- [22] von A. Hurwitz, Allgemeine Funktionentheorie und Elliptische Funktionen, von R. Courant, Geometrische Funktionentheorie, Springer-Verlag, Berlin, New York, 1964.
- [23] A.N.W. Hone, Non-existence of elliptic travelling wave solutions of the complex Ginzburg–Landau equation, *Physica D* 205 (2005) 292–306.
- [24] A.N.W. Hone, Painlevé test, Singularity Structure, and Integrability, nlin.SI/0502017.
- [25] N.A. Kudryashov, Exact solutions of a generalized equation of Ginzburg– Landau, Matematicheskoye modelirovanie 1 (1989) 151–158 {in Russian}.
- [26] N.A. Kudryashov, Nonlinear differential equations with exact solutions expressed via the Weierstrass function, nlin.CD/0312035.
- [27] N.A. Kudryashov, Simplest equation method to look for exact solutions of nonlinear differential equations, nlin.SI/0406007.
- [28] N.A. Kudryashov, Painlevé property and the first integrals of nonlinear differential equations, nlin.SI/0408041.

- [29] N.A. Kudryashov, M.V. Demina, Polygons for finding exact solutions of nonlinear differential equations, nlin.SI/0601035.
- [30] N.A. Kudryashov, Analytic Theory of Nonlinear Differential Equations, Moscow–Izhevsk, RCD, 2004, {in Russian}.
- [31] Ph. Marcq, H. Chaté, R. Conte, Exact Solutions of the One-Dimensional Quintic Complex Ginzburg–Landau Equation, *Physica D* 73 (1994) 305–317; patt-sol/9310004.
- [32] M. Musette, R. Conte, Analytic solitary waves of nonintegrable equations, *Physica D* 181 (2003) 70–79; nlin.PS/0302051.
- [33] A.G. Nikitin, T.A. Barannyk, Solitary wave and other solutions for nonlinear heat equations, math-ph/0303004.
- [34] F. Renner, A Constructive REDUCE Package Based Upon the Painlevé Analysis of Nonlinear Evolutions Equations in Hamiltonian and/or Normal Form, Comp. Phys. Comm. 70 (1992) 409–416.
- [35] W. van Saarloos, P.C. Hohenberg, Fronts and pulses in the complex Ginzburg– Landau equation near a subcritical bifurcations, *Phys. Rev. Lett.* 64 (1990) 749–752
- [36] W. van Saarloos, P.C. Hohenberg, Fronts, pulses, sources and sinks in generalized complex Ginzburg–Landau equations, *Physica D* 56 (1992) 303– 367, Erratum 69 (1993) 209
- [37] G.S. Santos, Application of Finite Expansion in Elliptic Functions to Solve Differential Educations, J. of the Physical Society of Japan 58 (1989) 4301–4310.
- [38] C. Scheen, Implementation of the Painlevé Test for Ordinary Differential Equations, *Theor. Comp. Sci.* 187 (1997) 87–104.
- [39] M. Tabor, Chaos and Integrability in Nonlinear Dynamics, Wiles, New York, 1989.
- [40] E.I. Timoshkova, A New Class of Trajectories of Motion in the Hénon–Heiles Potential Field, *Rus. Astron. J.* 76 (1999) 470–475, {in Russian}, *Astron. Rep.* 43 (1999) 406–411, {in English}.
- [41] S.Yu. Vernov, E.I. Timoshkova, On two nonintegrable cases of the generalized Hénon–Heiles system, *Physics of Atomic Nuclei (Yadernaya Fizica)* 68 (2005) 1947–1955 {in English}; 2008–2016 {in Russian}; math-ph/0402049.
- [42] S.Yu. Vernov, Constructing solutions for the generalized Hénon–Heiles system through the Painlevé test, *Theor. Math. Phys. (TMF)* 135 (2003) 409–419 {in Russian}, 792–801 {in English}; math-ph/0209063.
- [43] S.Yu. Vernov, Construction of Special Solutions for Nonintegrable Systems, J. of Nonlin. Math. Phys. 13 (2006) No. 1, pp. 50–63.
- [44] S.Yu. Vernov, Packages of Procedures for Maple 9 ('ellipso') and REDUCE 3.7, 2005, http://theory.sinp.msu.ru/~svernov/programs/

- [45] S.Yu. Vernov, Construction of Single-valued Solutions for Nonintegrable Systems with the Help of the Painlevé Test, in: *Proceedings of the International Conference "Computer Algebra in Scientific Computing"* (St. Petersburg, Russia, 2004), V.G. Ganzha, E.W. Mayr, E.V. Vorozhtsov (Eds.), Technische Universitat, Munchen, Garching, 2004, pp. 457–465; nlin.SI/0407062.
- [46] S.Yu. Vernov, Interdependence between the Laurent-Series and Elliptic Solutions of Nonintegrable Systems, in: the Lecture Notes of Computer Science (Springer) LNCS, V. 3718 (the Proceedings of International Conference "Computer Algebra in Scientific Computing" (CASC 2005, Kalamata, Greece)), Editors: V.G. Ganzha, E.W. Mayr and E.V. Vorozhtsov, Springer-Verlag, Berlin, Germany, 2005, pp. 457–468.
- [47] S.Yu. Vernov, Proof of the Absence of Elliptic Solutions of the Cubic Complex Ginzburg–Landau Equation, *Theor. Math. Phys. (Teor. Mat. Fiz.)* 146 (2006) 131–139 {in English}, 161–171 {in Russian}; Preprint version: On elliptic solutions of the cubic complex one-dimensional Ginzburg–Landau equation, nlin.PS/0503009.
- [48] S.Yu. Vernov, Construction of Exact Particular Solutions of Nonintegrable Systems by Means of Formal Laurent and Puiseux Series, *Programming and Computer Software*, **32** (2006) No. 2, pp. 77–83.
- [49] V.A. Vladimirov, E.V. Kutafina, Exact Travelling Wave Solutions of Some Nonlinear Evolutions Equations, *Rep. Math. Phys.* 54 (2004) 261–271.
- [50] G.Q. Xu, Z.B. Li, Symbolic Computation of the Painlevé Test for Nonlinear Partial Differential Equations Using Maple, *Comp. Phys. Comm.* 161 (2004) No. (1-2), pp. 65–75.